3.1

4

Let $Q(n)$ be the predicate "$n^2 \leq 30$."

a

Write $Q(2)$, $Q(-2)$, $Q(7)$, and $Q(-7)$, and indicate which of these statements are true and which are false.

> ✓ **Answer** ⌄
>
> $Q(2) : 4 \leq 30$ True
> $Q(-2) : 4 \leq 30$ True
> $Q(7) : 49 \leq 30$ False
> $Q(-7) : 49 \leq 30$ False

b

Find the truth set of $Q(n)$ if the domain of n is $\mathbf{Z}$, the set of all integers.

> ✓ **Answer**
>
> $n \leq \sqrt{30}$
> $n \leq 5$
> $n \in (-\infty, 5]$

7

Find the truth set of each predicate.

c

Predicate: $1 \leq x^2 \leq 4$, domain: $\mathbf{R}$

> ✓ **Answer**

$(1 \leq x \leq 2) \vee (1 \leq -x \leq 2)$
$x \in [-2, -1] \cup [1, 2]$

d

Predicate: $1 \leq x^2 \leq 4$, domain: $\mathbf{Z}$

✓ Answer

$(1 \leq x \leq 2) \vee (1 \leq -x \leq 2)$
$x \in [-2, -1] \cup [1, 2]$

12

Find counterexamples to show that the statement is false

$\forall$ real numbers x and y, $\sqrt{x + y} = \sqrt{x} + \sqrt{y}$.

✓ Answer

Let $x = 1$ and $y = 1$
$\sqrt{1 + 1} \neq \sqrt{1} + \sqrt{1}$
$\sqrt{2} \neq 2$

17

Rewrite the following in the form "$\exists$ _____ x such that _____."

b

Some real numbers are rational.

✓ Answer

$\exists$ some $x \in \mathbf{R}$ such that x is rational

20

Rewrite the following statement informally in at least two different ways without using variables or the symbol $\forall$ or the words "for all."

$\forall$ real numbers x, if x is positive then the square root of x is positive.

> ✓ **Answer**
>
> - If x is real and positive, then its square root is positive
> - If x is positive and its square root is not positive, then x is not a real number

22

Rewrite the following in the form "$\forall$ _____ x, if _____ then _____."

a

All Java programs have at least 5 lines.

> ✓ **Answer**
>
> $\forall$ Java programs x, x has at least 5 lines.

28

Rewrite each statement without using quantifiers or variables. Indicate which are true and which are false, and justify your answers as best as you can.

Let the domain of x be the set D of objects discussed in mathematics courses, and let $\mathrm{Real}(x)$ be "x is a real number," $\mathrm{Pos}(x)$ be "x is a positive real number," $\mathrm{Neg}(x)$ be "x is a negative real number," and $\mathrm{Int}(x)$ be "x is an integer."

a

$\mathrm{Pos}(0)$

> ✓ **Answer**
>
> 0 is a positive real number.

b

$$\forall x, \text{Real}(x) \land \text{Neg}(x) \rightarrow \text{Pos}(-x)$$

> ✓ **Answer**
>
> If the inverse of an object discussed in mathematics courses is not positive, then the object is either not a real number, or the object is not negative.

3.2

1

Which of the following is a negation for "All discrete mathematics students are athletic"? More than one answer may be correct.

1. There is a discrete mathematics student who is nonathletic.
2. All discrete mathematics students are nonathletic.
3. There is an athletic person who is not a discrete mathematics student.
4. No discrete mathematics students are athletic.
5. Some discrete mathematics students are nonathletic.
6. No athletic people are discrete mathematics students.

> ✓ **Answer**
>
> 1, 5

5

Write a negation for each of the following statements.

a

Every valid argument has a true conclusion.

> ✓ **Answer**
>
> There is an argument with a false conclusion

b

All real numbers are positive, negative, or zero.

12

Determine whether the proposed negation is correct. If it is not, write a correct negation.

Statement: The product of any irrational number and any rational number is irrational.

Proposed negation: The product of any irrational number and any rational number is rational.

17

Write a negation for each statement

$\forall$ integers d, if $\frac{6}{d}$ is an integer, then $d = 3$

29

Write the contrapositive, converse, and inverse. Indicate as best as you can which of these statements are true and which are false. Give a counterexample for each that is false.

$\forall n \in Z$, if n is prime then n is odd or $n = 2$.

48

Use the facts that the negation of a $\forall$ statement is a $\exists$ statement and that the negation of an if-then statement is an and statement to rewrite the statement without using the word necessary or sufficient.

Being a polynomial is not a sufficient condition for a function to have a real root.

> ✓ **Answer**
>
> $x \in$ set of all functions
> $P : x$ is a polynomial
> $Q : x$ has a real root
> $P \wedge \neg Q$

3.3

2

Let $G(x, y)$ be "$x^2 > y$" Indicate which of the following statements are true and which are false.

a

$G(2, 3)$

> ✓ **Answer**
>
> $4 > 3$ True

b

$G(1, 1)$

> ✓ **Answer**
>
> $1 > 1$ False

c

$G\left(\frac{1}{2}, \frac{1}{2}\right)$

> ✓ **Answer**
>
> $\frac{1}{4} > \frac{1}{2}$ False

d

$G(-2, 2)$

> ✓ **Answer**
>
> $4 > 2$ True

10

Determine whether each of the following statements is true or false.

a

$\forall$ student S, $\exists$ a dessert D such that S chose D.

> ✓ **Answer**
>
> True, Yuen chose a Pie, Tim chose a Pie and a Cake, and Uta chose a Pie

b

$\forall$ student S, $\exists$ a salad T such that S chose T.

19

Rewrite the statement in English without using the symbol ∀ or ∃ or variables and expressing your answer as simply as possible. Also, write a negation for the statement.

$\exists x \in \mathbf{R}$ such that for every real number y,
$x + y = 0$

✓ **Answer**

For every real number, there is another real number such that their sum is 0. There is a real number that does not have another a real number where their sum is 0.

23

Rewrite the statement in English without using the symbol ∀ or ∃ or variables. Also, indicate whether the statement is true or false.

a

∀ nonzero real number r, ∃ a real number s such that $rs = 1$.

✓ **Answer**

Every nonzero real number has another real number where their product is 1. True.

b

∃ a real number r such that ∀ nonzero real number s, $rs = 1$.

✓ **Answer**

There is a real number where multiplying it with any other real number results in 1. False, literally any number except 1.

3.4

2

Use universal instantiation or universal modus ponens to
fill in valid conclusions for the argument

If an integer n equals $2k$ and k is an integer, then n is even.
0 equals 2×0 and 0 is an integer.

> ✓ **Answer**
>
> 0 is even.

14

The argument may be valid by universal modus ponens or universal modus tollens;
others are invalid and exhibit the converse or the inverse error. State whether the
argument is valid or invalid. Justify your answer.

If compilation of a computer program produces error messages, then the program is
not correct.
Compilation of this program does not produce error messages.
∴ This program is correct.

> ✓ **Answer**
>
> This has an inverse error, this assumes the inverse.

22-24

Indicate whether the arguments in 22–24 are valid or
invalid. Support your answers by drawing diagrams.

22

All discrete mathematics students can tell a valid argument from an invalid one.
All thoughtful people can tell a valid argument from an invalid one.
∴ All discrete mathematics students are thoughtful.

> ✓ **Answer**

$A \rightarrow C$

$B \rightarrow C$

$A \not\rightarrow B$

False, assumes the converse of the second statement.

24

No vegetarians eat meat.

All vegans are vegetarian.

∴ No vegans eat meat.

> ✓ **Answer**
>
> $A \rightarrow \neg B$
>
> $C \rightarrow A$
>
> $C \rightarrow \neg B$
>
> True, by chain rule