

# Discrete Mathematics 1

## Chapter 5: Counting

Department of Mathematics  
The FPT university

# Chapter 5: Introduction

# Chapter 5: Introduction

**Topics covered:**

# Chapter 5: Introduction

## Topics covered:

### 5.1 The Basics of Counting

## Topics covered:

5.1 The Basics of Counting

5.2 The Pigeonhole Principle

# Chapter 5: Introduction

## Topics covered:

5.1 The Basics of Counting

5.2 The Pigeonhole Principle

5.3 Permutations and Combinations

## Topics covered:

5.1 The Basics of Counting

5.2 The Pigeonhole Principle

5.3 Permutations and Combinations

5.6 Generating Permutations and Combinations

## Topics covered:

5.1 The Basics of Counting

5.2 The Pigeonhole Principle

5.3 Permutations and Combinations

5.6 Generating Permutations and Combinations



# 5.1 The Basics of Counting

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

### Example 1.

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

**Example 1.** A class has 24 male students and 2 female students.



## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick some one from this class to be a representative to University's Student council is

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick some one from this class to be a representative to University's Student council is  $24+2=26$ .

## 5.1 The Basics of Counting

### The sum rule - The Inclusion-exclusion Principle

- Let  $A$  and  $B$  be two disjoint sets. Then  $|A \cup B| = |A| + |B|$
- Let  $A$  and  $B$  be two arbitrary sets. Then:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick some one from this class to be a representative to University's Student council is  $24+2=26$ .



## Example 2. (Counting numbers)

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.**



**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.** Count:

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.** Count:

- (a) The number of positive integers not exceeding 1000 and divisible by 12

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.** Count:

- (a) The number of positive integers not exceeding 1000 and divisible by 12
- (b) The number of positive integers less than 1000, greater than 100 and divisible by 12

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.** Count:

- (a) The number of positive integers not exceeding 1000 and divisible by 12
- (b) The number of positive integers less than 1000, greater than 100 and divisible by 12
- (c) The number of integers not exceeding 1000 and divisible by 12 or 8

**Example 2.** (Counting numbers) The number of positive integers not exceeding  $n$  and divisible by  $k$  is  $\lfloor n/k \rfloor$ .

**Example 3.** Count:

- (a) The number of positive integers not exceeding 1000 and divisible by 12
- (b) The number of positive integers less than 1000, greater than 100 and divisible by 12
- (c) The number of integers not exceeding 1000 and divisible by 12 or 8
- (d) The number of positive integers less than 1000, divisible by 12 but not divisible by 8.





## Product rule

Let  $A$  and  $B$  be two sets. Then



## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.**

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students.

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.**

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is



## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.**

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions)

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions) Count:

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions) Count:

- (a) The number of functions from a set of 5 elements to a set of 3 elements

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions) Count:

- (a) The number of functions from a set of 5 elements to a set of 3 elements
- (b) The number of one-to-one functions from a set of 5 elements to a set of 3 elements

## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions) Count:

- (a) The number of functions from a set of 5 elements to a set of 3 elements
- (b) The number of one-to-one functions from a set of 5 elements to a set of 3 elements
- (c) The number of one-to-one functions from a set of 3 elements to a set of 5 elements



## Product rule

Let  $A$  and  $B$  be two sets. Then  $|A \times B| = |A| \cdot |B|$

**Example 1.** A class has 24 male students and 2 female students. The number of choices to pick a pair of one male and one female student for a dancing contest is  $24 \times 2 = 48$

**Example 2.** The number of bit strings of length  $n$  is  $2^n$ .

**Example 3.** Count the number of bit strings of length 7 and either starts with a 0 or ends with the two bits 10.

**Example 4.** (Counting functions) Count:

- (a) The number of functions from a set of 5 elements to a set of 3 elements
- (b) The number of one-to-one functions from a set of 5 elements to a set of 3 elements
- (c) The number of one-to-one functions from a set of 3 elements to a set of 5 elements

## 5.2 Pigeonhole Principle

## 5.2 Pigeonhole Principle



## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

**Example.**

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

(1) Among 100 people there are at least

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

- (1) Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  people born in the same month.

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

- (1) Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  people born in the same month.
- (2) How many people must be selected to guarantee that there are at least 10 people born in the same month?



## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

- (1) Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  people born in the same month.
- (2) How many people must be selected to guarantee that there are at least 10 people born in the same month?
- (3) How many cards must be selected from a standard deck of 52 cards to guarantee that there are at least 3 cards of the same suit?

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

- (1) Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  people born in the same month.
- (2) How many people must be selected to guarantee that there are at least 10 people born in the same month?
- (3) How many cards must be selected from a standard deck of 52 cards to guarantee that there are at least 3 cards of the same suit? at least 3 hearts?

## 5.2 Pigeonhole Principle

If  $N$  objects are placed into  $k$  boxes, then there is at least one box that contains at least  $\lceil N/k \rceil$  objects.

### Example.

- (1) Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  people born in the same month.
- (2) How many people must be selected to guarantee that there are at least 10 people born in the same month?
- (3) How many cards must be selected from a standard deck of 52 cards to guarantee that there are at least 3 cards of the same suit? at least 3 hearts?


# The Ramsey number $R(m, n)$

# The Ramsey number $R(m, n)$

A graph of 2 colors (blue, red) is a graph so that between two distinct vertices there is one edge, and this edge is either blue or red.

# The Ramsey number $R(m, n)$

A graph of 2 colors (blue, red) is a graph so that between two distinct vertices there is one edge, and this edge is either blue or red.





Let  $m, n$  be positive integers greater than or equal to 2.



Let  $m, n$  be positive integers greater than or equal to 2.

The Ramsey number  $R(m, n)$  is the least number of vertices of a graph of 2 colors such that we can always find a subgraph of  $m$  vertices whose edges are all blue, or a subgraph of  $n$  vertices whose edges are all red.

Let  $m, n$  be positive integers greater than or equal to 2.

The Ramsey number  $R(m, n)$  is the least number of vertices of a graph of 2 colors such that we can always find a subgraph of  $m$  vertices whose edges are all blue, or a subgraph of  $n$  vertices whose edges are all red.

**Problem.**

Let  $m, n$  be positive integers greater than or equal to 2.

The Ramsey number  $R(m, n)$  is the least number of vertices of a graph of 2 colors such that we can always find a subgraph of  $m$  vertices whose edges are all blue, or a subgraph of  $n$  vertices whose edges are all red.

**Problem.** Find  $R(2, 3)$ ,

Let  $m, n$  be positive integers greater than or equal to 2.

The Ramsey number  $R(m, n)$  is the least number of vertices of a graph of 2 colors such that we can always find a subgraph of  $m$  vertices whose edges are all blue, or a subgraph of  $n$  vertices whose edges are all red.

**Problem.** Find  $R(2, 3)$ ,  $R(2, n)$ ,

Let  $m, n$  be positive integers greater than or equal to 2.

The Ramsey number  $R(m, n)$  is the least number of vertices of a graph of 2 colors such that we can always find a subgraph of  $m$  vertices whose edges are all blue, or a subgraph of  $n$  vertices whose edges are all red.

**Problem.** Find  $R(2, 3)$ ,  $R(2, n)$ ,  $R(3, 3)$

## 4.3 Permutations and Combinations

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set.



## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set. The number of permutations of  $S$  is  $n!$ .

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set. The number of permutations of  $S$  is  $n!$ .
- An  $r$ -permutation of  $S$  is an ordered arrangement of  $r$  elements of  $S$ .

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set. The number of permutations of  $S$  is  $n!$ .
- An  $r$ -permutation of  $S$  is an ordered arrangement of  $r$  elements of  $S$ . The number of  $r$ -permutations is

$$P(n, r) := n(n-1) \cdots (n-r+1) = \frac{n!}{(n-r)!}$$

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set. The number of permutations of  $S$  is  $n!$ .
- An  $r$ -permutation of  $S$  is an ordered arrangement of  $r$  elements of  $S$ . The number of  $r$ -permutations is

$$P(n, r) := n(n-1) \cdots (n-r+1) = \frac{n!}{(n-r)!}$$

- An  $r$ -combination of  $S$  is an unordered selection of  $r$  elements from  $S$ .

## 4.3 Permutations and Combinations

Cho  $S = \{1, 2, \dots, n\}$ .

- A **permutation** of  $S$  is an ordered arrangement of the elements of the set. The number of permutations of  $S$  is  $n!$ .
- An  $r$ -permutation of  $S$  is an ordered arrangement of  $r$  elements of  $S$ . The number of  $r$ -permutations is

$$P(n, r) := n(n-1) \cdots (n-r+1) = \frac{n!}{(n-r)!}$$

- An  $r$ -combination of  $S$  is an unordered selection of  $r$  elements from  $S$ . The number of  $r$ -combinations of  $S$  is

$$C(n, r) := \frac{n!}{r!(n-r)!}$$



## Example 1.

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?



## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students.

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students. How many choices are there to pick 3 students to participate in a Calculus competition?

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students. How many choices are there to pick 3 students to participate in a Calculus competition?  
How many choices to pick 3 students, one will participate in a Calculus competition, one will participate in a Algebra competition, and one will participate in a Discrete Math competition?

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students. How many choices are there to pick 3 students to participate in a Calculus competition?  
How many choices to pick 3 students, one will participate in a Calculus competition, one will participate in a Algebra competition, and one will participate in a Discrete Math competition?
- (c) A class has 9 female and 20 male students. How many ways to pick 11 students for a soccer team if

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students. How many choices are there to pick 3 students to participate in a Calculus competition?  
How many choices to pick 3 students, one will participate in a Calculus competition, one will participate in a Algebra competition, and one will participate in a Discrete Math competition?
- (c) A class has 9 female and 20 male students. How many ways to pick 11 students for a soccer team if
  - (i) the team must have exactly 3 female students?

## Example 1.

- (a) How many permutations of the characters ABC12DE that contain the string BC1?
- (b) A class has 25 students. How many choices are there to pick 3 students to participate in a Calculus competition?  
How many choices to pick 3 students, one will participate in a Calculus competition, one will participate in a Algebra competition, and one will participate in a Discrete Math competition?
- (c) A class has 9 female and 20 male students. How many ways to pick 11 students for a soccer team if
  - (i) the team must have exactly 3 female students?
  - (ii) the team must have at least one female student?



## Example 2.



## Example 2.

(a) How many bit strings of length 8 containing exactly two 0s?

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s?

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s? At least two 0s?

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s? At least two 0s?
- (b) Flip a coin, there are two possible outcomes: head or tail.

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s? At least two 0s?
- (b) Flip a coin, there are two possible outcomes: head or tail. Flip a coin 10 times.

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s? At least two 0s?
- (b) Flip a coin, there are two possible outcomes: head or tail. Flip a coin 10 times. How many possible outcomes are there?

## Example 2.

- (a) How many bit strings of length 8 containing exactly two 0s? At most two 0s? At least two 0s?
- (b) Flip a coin, there are two possible outcomes: head or tail. Flip a coin 10 times. How many possible outcomes are there? How many outcomes with an equal number of heads and tails?

## 5.6 Generating Permutations and Combinations



## 5.6 Generating Permutations and Combinations

Let  $S = \{1, 2, \dots, n\}$ .

## 5.6 Generating Permutations and Combinations

Let  $S = \{1, 2, \dots, n\}$ .

**Problem.** Write an algorithm to generate all permutations or all  $r$ -combinations of  $S$ .

## 5.6 Generating Permutations and Combinations

Let  $S = \{1, 2, \dots, n\}$ .

**Problem.** Write an algorithm to generate all permutations or all  $r$ -combinations of  $S$ .

The following algorithms will be based on lexicographic order (or dictionary order) of permutations or  $r$ -combinations. For example:

$$123 < 132 < 312$$

## 5.6 Generating Permutations and Combinations

Let  $S = \{1, 2, \dots, n\}$ .

**Problem.** Write an algorithm to generate all permutations or all  $r$ -combinations of  $S$ .

The following algorithms will be based on lexicographic order (or dictionary order) of permutations or  $r$ -combinations. For example:

$$123 < 132 < 312$$

Problem:

## 5.6 Generating Permutations and Combinations

Let  $S = \{1, 2, \dots, n\}$ .

**Problem.** Write an algorithm to generate all permutations or all  $r$ -combinations of  $S$ .

The following algorithms will be based on lexicographic order (or dictionary order) of permutations or  $r$ -combinations. For example:

$$123 < 132 < 312$$

Problem: Given a permutation or an  $r$ -combination, find the next permutation or  $r$ -combination in the lexicographic order.

# Generating the Next Permutation

# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

- Work from right to left, find the first two consecutive numbers  $(a_i, a_{i+1})$  with the property that  $a_i < a_{i+1}$ .



# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

- Work from right to left, find the first two consecutive numbers  $(a_i, a_{i+1})$  with the property that  $a_i < a_{i+1}$ .

$$a_1, a_2, \dots, a_i, a_{i+1}, \dots, a_n$$

# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

- Work from right to left, find the first two consecutive numbers  $(a_i, a_{i+1})$  with the property that  $a_i < a_{i+1}$ .

$$a_1, a_2, \dots, a_i, a_{i+1}, \dots, a_n$$

- Among the numbers  $a_{i+1}, a_{i+2}, \dots, a_n$  choose the smallest element  $a_k$  with the property that  $a_k > a_i$ .

$$a_1 a_2 \dots a_i \dots a_k \dots a_n$$

# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

- Work from right to left, find the first two consecutive numbers  $(a_i, a_{i+1})$  with the property that  $a_i < a_{i+1}$ .

$$a_1, a_2, \dots, a_i, a_{i+1}, \dots, a_n$$

- Among the numbers  $a_{i+1}, a_{i+2}, \dots, a_n$  choose the smallest element  $a_k$  with the property that  $a_k > a_i$ .

$$a_1 a_2 \dots a_i \dots a_k \dots a_n$$

- Swap  $a_i$  and  $a_k$

# Generating the Next Permutation

Given a permutation  $a_1 a_2 \dots a_n$  of the set  $S = \{1, 2, \dots, n\}$ .

- Work from right to left, find the first two consecutive numbers  $(a_i, a_{i+1})$  with the property that  $a_i < a_{i+1}$ .

$$a_1, a_2, \dots, a_i, a_{i+1}, \dots, a_n$$

- Among the numbers  $a_{i+1}, a_{i+2}, \dots, a_n$  choose the smallest element  $a_k$  with the property that  $a_k > a_i$ .

$$a_1 a_2 \dots a_i \dots a_k \dots a_n$$

- Swap  $a_i$  and  $a_k$
- Sort the list  $a_{i+1}, a_{i+2}, \dots, a_n$  in the increasing order.

# Generating the Next $r$ -Combination

# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

Given an  $r$ -combination  $a_1 a_2 \dots a_r$ .

# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

Given an  $r$ -combination  $a_1 a_2 \dots a_r$ .

- Work from right to left, find the first number  $a_i$  with the property that  $a_i \neq n - r + i$ .



# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

Given an  $r$ -combination  $a_1 a_2 \dots a_r$ .

- Work from right to left, find the first number  $a_i$  with the property that  $a_i \neq n - r + i$ .

$$a_1 a_2 \dots a_i \dots a_n$$

# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

Given an  $r$ -combination  $a_1 a_2 \dots a_r$ .

- Work from right to left, find the first number  $a_i$  with the property that  $a_i \neq n - r + i$ .

$$a_1 a_2 \dots a_i \dots a_n$$

- Replace the sequence  $a_i, a_{i+1}, \dots, a_n$  by the sequence of consecutive integers, starting from  $a_i + 1$ .

# Generating the Next $r$ -Combination

**Convention:** In the following algorithm, the elements in an  $r$ -combination of  $S = \{1, 2, \dots, n\}$  are listed in the increasing order.

Given an  $r$ -combination  $a_1 a_2 \dots a_r$ .

- Work from right to left, find the first number  $a_i$  with the property that  $a_i \neq n - r + i$ .

$$a_1 a_2 \dots a_i \dots a_n$$

- Replace the sequence  $a_i, a_{i+1}, \dots, a_n$  by the sequence of consecutive integers, starting from  $a_i + 1$ .