

Contents

1	Mean Polytopes and Maximum Entropy Distributions	1
1.1	Maximum entropy problem	1
1.2	The mean polytope	2
1.2.1	Faces	2
1.2.2	Representation in the CP format	5
1.2.3	Examples of elementarily representable faces	6
1.2.4	Partition into relative interiors of faces	7
1.3	Exponential families	7
1.3.1	Representation as elementary Computation-Activation Networks	8
1.3.2	Maximum entropy distributions	9
1.4	Generic maximum entropy distributions	10
1.4.1	Main result	11
1.4.2	Family of maximum entropy distributions	12
1.5	Boolean statistics	12
1.5.1	Characterization of elementary representable faces	13
1.5.2	Interpretation by propositional formulas	15
1.5.3	Faces activated in the TT format	17
1.5.4	Construction of Boolean statistics to polytopes	19
1.5.5	Examples in statistical physics and coding theory	20
1.6	Mean as a Statistic	21
1.7	Face Computation	22
1.7.1	Examples	22
1.7.2	Generic Construction	23
1.7.3	Relation with Facet Complexity	23
	References	24

Chapter 1

Mean Polytopes and Maximum Entropy Distributions

Abstract Motivated by the search for maximum entropy distributions we investigate the convex polytope of mean parameters to a statistic function. We then show that exponential families are maximum entropy distributions to mean parameters in the relative interior of the mean polytope and generalize the result to the faces of the mean polytope. In the case of Boolean statistics we provide a characterization in terms of propositional logic. Further, we investigate the mean parameter as a sufficient statistic and investigate schemes to compute the activation tensor to faces.

1.1 Maximum entropy problem

Given a non-negative and non-vanishing base measure $\nu [X_{[d]}]$, a probability distribution is a non-negative tensor $\mathbb{P} [X_{[d]}]$ such that

$$\langle \mathbb{P} [X_{[d]}], \nu [X_{[d]}] \rangle_{[\emptyset]} = 1.$$

We denote the set of such distributions by

$$\begin{aligned} \Lambda^\nu &= \{ \mathbb{P} [X_{[d]}] : \mathbb{P} [X_{[d]}] \geq 0 [X_{[d]}], \langle \mathbb{P} [X_{[d]}], \nu [X_{[d]}] \rangle_{[\emptyset]} = 1, \\ &\quad \forall_{x_{[d]} \in \times_{k \in [d]} [m_k]} \nu [X_{[d]} = x_{[d]}] = 0 \Rightarrow \mathbb{P} [X_{[d]} = x_{[d]}] = 0 \}. \end{aligned}$$

Note that we here restrict the support of the tensor $\mathbb{P} [X_{[d]}]$ to be included in the support of the base measure, to avoid ambiguity in the representation.

The entropy of a distribution relative to the base measure ν is

$$\mathbb{H}_\nu [\mathbb{P} [X_{[d]}]] = - \langle \mathbb{P} [X_{[d]}], \ln [\mathbb{P} [X_{[d]}]] , \nu [X_{[d]}] \rangle_{[\emptyset]}.$$

Note that this quantity depends on the chosen base measure.

A statistic is a function $t : \times_{k \in [d]} [m_k] \rightarrow \mathbb{R}^p$ assigning a real vector of dimension p to each state $x_{[d]} \in \times_{k \in [d]} [m_k]$. We denote by t_ℓ the restriction to the ℓ -th coordinate of t . The mean parameter of a distribution $\mathbb{P} [X_{[d]}]$ to a statistic t is the vector $\mu [L] \in \mathbb{R}^p$ with the coordinates

$$\mu [L = \ell] = \mathbb{E} [f_\ell] = \langle t_\ell [X_{[d]}], \mathbb{P} [X_{[d]}], \nu [X_{[d]}] \rangle_{[\emptyset]}.$$

To express the computation of the mean parameter as a single contraction, we introduce a coordinate selection variable L of dimension p and the selection encoding $\sigma^t [X_{[d]}, L]$ with coordinates by

$$\sigma^t [X_{[d]} = x_{[d]}, L = \ell] = t_\ell [X_{[d]} = x_{[d]}].$$

The mean parameter is then

$$\mu[L] = \langle \mathbb{P}[X_{[d]}], \sigma^t[X_{[d]}, L] \rangle_{[L]}.$$

The maximum entropy problem given a mean parameter $\mu^*[L]$ is stated by

$$\operatorname{argmax}_{\mathbb{P}[X_{[d]}] \in \Lambda^\nu} \mathbb{H}_\nu[\mathbb{P}[X_{[d]}]] \quad \text{subject to} \quad \langle \mathbb{P}[X_{[d]}], \sigma^t[X_{[d]}, L], \nu[X_{[d]}] \rangle_{[L]} = \mu^*[L] \quad (\mathbf{P}_{t,\mu,\nu})$$

1.2 The mean polytope

Given a statistic function

$$t : \bigtimes_{k \in [d]} [m_k] \rightarrow \mathbb{R}^p$$

on the states $\bigtimes_{k \in [d]} [m_k]$ and a base measure $\nu[X_{[d]}]$ the mean polytope is the set

$$\mathcal{M}_{t,\nu} := \left\{ \langle \mathbb{P}[X_{[d]}], \sigma^t[X_{[d]}, L], \nu[X_{[d]}] \rangle_{[L]} : \mathbb{P}[X_{[d]}] \in \Lambda^\nu \right\}$$

containing all mean parameters for the distributions in Λ^ν . By $\sigma^t[X_{[d]}, L]$ we denote the selection encoding of the statistic (see introduction). Since Λ^ν is the convex hull of the vertices $\frac{1}{\nu[X_{[d]} = x_{[d]}]} \cdot \epsilon_{x_{[d]}}[X_{[d]}]$, and any mean parameter is a linear transformation of the distribution, the mean polytope is the convex hull of the vectors

$$\mathcal{N}_{t,\nu} := \{ \sigma^t[X_{[d]} = x_{[d]}, L] : x_{[d]} \in \bigtimes_{k \in [d]} [m_k], \nu[X_{[d]} = x_{[d]}] \neq 0 \}.$$

1.2.1 Faces

When studying maximum entropy distributions, we will pay special attention to the faces of a polytope, which form the boundary. Here, we define them based on convex hulls and introduce tensor representations by refined base measures.

Definition 1.1 We say that the convex hull of a subset $\mathcal{N}_{t,\nu}^\mathcal{F} \subset \mathcal{N}_{t,\nu}$ is a face of a polytope $\mathcal{M}_{t,\nu}$, if and only if there is a vector $\theta_\mathcal{F}[L] \in \mathbb{R}^p$ such that

$$\mathcal{N}_{t,\nu}^\mathcal{F} = \operatorname{argmax}_{\mu[L] \in \mathcal{N}_{t,\nu}} \langle \mu[L], \theta_\mathcal{F}[L] \rangle_{[\emptyset]}.$$

We denote the face as $\mathcal{F} = \operatorname{conv}(\mathcal{N}_{t,\nu}^\mathcal{F})$. The set of all faces of the mean polytope $\mathcal{M}_{t,\nu}$ is denoted by $L(\mathcal{M}_{t,\nu})$.

The set $L(\mathcal{M}_{t,\nu})$, partially ordered by the inclusion operation, is a lattice (see [10]). We now show that each face itself is a convex polytope corresponding to a refined base measure.

Definition 1.2 The refined base measure with respect to a face $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ is the tensor $\nu^{t,\mathcal{F}}[X_{[d]}]$ with coordinates to $x_{[d]} \in \bigtimes_{k \in [d]} [m_k]$ by

$$\nu^{t,\mathcal{F}}[X_{[d]}] = \begin{cases} \nu[X_{[d]} = x_{[d]}] & \text{if } t(x_{[d]}) \in \mathcal{N}_{t,\nu}^\mathcal{F} \\ 0 & \text{else} \end{cases}.$$

The refined base measure has by definition a smaller support than the base measure. We now show that restricting the distributions in Λ^ν to those in $\Lambda^{\nu^{t,\mathcal{F}}}$ amounts to a restriction of the polytope $\mathcal{M}_{t,\nu}$ to the corresponding face.

Lemma 1.1 For any face $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ we have

$$\mathcal{F} = \mathcal{M}_{t,\nu^{t,\mathcal{F}}}.$$

Proof. For any $x_{[d]} \in \times_{k \in [d]} [m_k]$ we have $\nu^{t, \mathcal{F}} [X_{[d]} = x_{[d]}] \neq 0$ if and only if $t(x_{[d]}) \in \mathcal{N}_{t, \nu}^{\mathcal{F}}$ and $\nu [X_{[d]} = x_{[d]}] \neq 0$. Thus we have

$$\begin{aligned} \mathcal{F} &= \text{conv} \left(\mathcal{N}_{t, \nu}^{\mathcal{F}} \right) \\ &= \text{conv} \left(\sigma^t [X_{[d]} = x_{[d]}, L] : t(x_{[d]}) \in \mathcal{N}_{t, \nu}^{\mathcal{F}}, \nu [X_{[d]} = x_{[d]}] \neq 0 \right) \\ &= \text{conv} \left(\sigma^t [X_{[d]} = x_{[d]}, L] : \nu^{t, \mathcal{F}} [X_{[d]} = x_{[d]}] \neq 0 \right) \\ &= \mathcal{M}_{t, \nu^{t, \mathcal{F}}}. \square \end{aligned}$$

Representability of a distribution with respect to face measures is an equivalent condition for the mean parameter of a distribution to be on a face, as we show next.

Lemma 1.2 *If and only if for a distribution $\mathbb{P} [X_{[d]}] \in \Lambda^\nu$ and a face $\mathcal{F} \in L(\mathcal{M}_{t, \nu})$ we have*

$$\langle \sigma^t [X_{[d]}, L], \mathbb{P} [X_{[d]}], \nu [X_{[d]}] \rangle_{[L]} \in \mathcal{F},$$

then

$$\mathbb{P} [X_{[d]}] \in \Lambda^{\nu^{t, \mathcal{F}}}.$$

Proof. For a distribution $\mathbb{P} [X_{[d]}] \in \Lambda^\nu$ to be in $\Lambda^{\nu^{t, \mathcal{F}}}$, it is enough to show that $\mathbb{P} [X_{[d]}]$ is only supported at $\nu^{t, \mathcal{F}} [X_{[d]}]$. We have

$$\mu [L] = \sum_{x_{[d]}} \mathbb{P} [X_{[d]} = x_{[d]}] \cdot \nu [X_{[d]} = x_{[d]}] \cdot \gamma^t [X_{[d]} = x_{[d]}, L].$$

Now, let $\theta_{\mathcal{F}} [L]$ be a face normal to the face $\mathcal{F}$. We then have

$$\langle \mu [L], \theta_{\mathcal{F}} [L] \rangle_{[\emptyset]} = \sum_{x_{[d]} \in \times_{k \in [d]} [m_k]} \mathbb{P} [X_{[d]} = x_{[d]}] \cdot \nu [X_{[d]} = x_{[d]}] \cdot \langle \gamma^t [X_{[d]} = x_{[d]}, L], \theta_{\mathcal{F}} [L] \rangle_{[\emptyset]}.$$

Now if and only if $\mathbb{P} [X_{[d]} = x_{[d]}] \cdot \nu [X_{[d]} = x_{[d]}]$ is supported only for $x_{[d]}$ with $t(x_{[d]}) \in \mathcal{N}_{t, \nu}^{\mathcal{F}}$ we have that

$$\langle \mu [L], \theta_{\mathcal{F}} [L] \rangle_{[\emptyset]} = \max_{\mu [L] \in \mathcal{M}_{t, \nu}} \langle \mu [L], \theta_{\mathcal{F}} [L] \rangle_{[\emptyset]}$$

which is equal to $\mu [L] \in \mathcal{F}$. Thus, if and only if $\mu [L] \in \mathcal{F}$ then $\mathbb{P} [X_{[d]}]$ is only supported at $x_{[d]}$ in the support of $\nu^{t, \mathcal{F}} [X_{[d]}]$. $\square$

Let us now investigate tensor network representations of refined base measures, based on the basis encoding β^t of a statistic.

Theorem 1.1 *For any face $\mathcal{F}$ of $\mathcal{M}$ we have*

$$\nu^{t, \mathcal{F}} [X_{[d]}] = \langle \beta^t [Y_{[p]}, X_{[d]}], \kappa^{\mathcal{F}} [Y_{[p]}], \nu [X_{[d]}] \rangle_{[X_{[d]}]}$$

where

$$\kappa^{\mathcal{F}} [Y_{[p]}] = \sum_{v[L] \in \mathcal{N}_{t, \nu}^{\mathcal{F}}} \epsilon_{v[L]} [Y_{[p]}].$$

Proof. For any $v[L] \in \mathcal{N}_{t, \nu}^{\mathcal{F}}$ the tensor

$$\tau^v [X_{[d]}] = \langle \beta^t [Y_{[p]}, X_{[d]}], \epsilon_{v[L]} [Y_{[p]}] \rangle_{[X_{[d]}]}$$

is the indicator of the preimage of μ under σ^t . Since preimages of the elements in $\mathcal{N}_{t,\nu}^{\mathcal{F}}$ are disjoint, the support of $\tau^v [X_{[d]}]$ is disjoint and their sum

$$\sum_{v[L] \in \mathcal{N}_{t,\nu}^{\mathcal{F}}} \tau^v [X_{[d]}]$$

is the indicator of the preimage of $\mathcal{F}$ under σ^t . The face measure obeys thus

$$\begin{aligned} \nu^{t,\mathcal{F}} [X_{[d]}] &= \left\langle \left(\sum_{v[L] \in \mathcal{N}_{t,\nu}^{\mathcal{F}}} \tau^v [X_{[d]}] \right), \nu [X_{[d]}] \right\rangle_{[X_{[d]}]} \\ &= \sum_{v[L] \in \mathcal{N}_{t,\nu}^{\mathcal{F}}} \langle \beta^t [Y_{[p]}, X_{[d]}], \epsilon_{v[L]} [Y_{[p]}], \nu [X_{[d]}] \rangle_{[X_{[d]}]} \\ &= \langle \beta^t [Y_{[p]}, X_{[d]}], \kappa^{\mathcal{F}} [Y_{[p]}], \nu [X_{[d]}] \rangle_{[X_{[d]}]} \square \end{aligned}$$

We now investigate the representation of refined base measures by Computation-Activation Networks.

Definition 1.3 Let $\mathcal{G}$ be a hypergraph whose nodes include $[p]$. We say that a face $\mathcal{F}$ is representable by $\mathcal{G}$ if and only if there is a set $\mathcal{U}$ of basis vectors with $\mathcal{U} \cap \mathcal{N}_{t,\nu} = \mathcal{N}_{t,\nu}^{\mathcal{F}}$ and there is a tensor network $\kappa^{\mathcal{G}}$ with respect to the hypergraph $\mathcal{G}$ such that

$$\langle \kappa^{\mathcal{G}} \rangle_{[Y_{[p]}]} = \sum_{v \in \mathcal{U}} \epsilon_v [Y_{[p]}] .$$

We call any such tensor network a face activating tensor network.

Lemma 1.3 If and only if a face is representable by a hypergraph $\mathcal{G}$, we have

$$\frac{1}{\langle \nu^{t,\mathcal{F}} [X_{[d]}] \rangle_{[\emptyset]}} \cdot \nu^{t,\mathcal{F}} [X_{[d]}] \in \Lambda^{t,\mathcal{G},\nu} .$$

Proof. For any $\mathcal{U}$ with $\mathcal{U} : \mathcal{U} \cap \mathcal{N}_{t,\nu} = \mathcal{N}_{t,\nu}^{\mathcal{F}}$ and tensor network $\kappa^{\mathcal{G}}$ respecting the assumptions of Def. 1.3 we have

$$\begin{aligned} \left\langle \langle \kappa^{\mathcal{G}} \rangle_{[Y_{[p]}]}, \beta^t [Y_{[p]}, X_{[d]}] \right\rangle_{[X_{[d]}]} &= \left\langle \sum_{v \in \mathcal{U}} \epsilon_v [Y_{[p]}], \beta^t [Y_{[p]}, X_{[d]}] \right\rangle_{[X_{[d]}]} \\ &= \left\langle \sum_{v \in \mathcal{N}_{t,\nu}^{\mathcal{F}}} \epsilon_v [Y_{[p]}], \beta^t [Y_{[p]}, X_{[d]}] \right\rangle_{[X_{[d]}]} . \end{aligned}$$

Here we used in the second equation that only the vertices in $\mathcal{N}_{t,\nu}$ are in the image of t . It follows, that

$$\frac{1}{\langle \nu^{t,\mathcal{F}} [X_{[d]}] \rangle_{[\emptyset]}} \cdot \nu^{t,\mathcal{F}} [X_{[d]}] = \frac{\left\langle \langle \kappa^{\mathcal{G}} \rangle_{[Y_{[p]}]}, \beta^t [Y_{[p]}, X_{[d]}] \right\rangle_{[X_{[d]}]}}{\left\langle \langle \kappa^{\mathcal{G}} \rangle_{[Y_{[p]}]}, \beta^t [Y_{[p]}, X_{[d]}], \nu [X_{[d]}] \right\rangle_{[\emptyset]}}$$

and thus $\nu^{t,\mathcal{F}} [X_{[d]}] \in \Lambda^{t,\mathcal{G},\nu}$. □

Example 1.1 (Vertices) Vertices $\mathcal{F} = \{v [L]\}$ are proper faces of affine dimension 0, that is they consist of single vectors $v [L]$. Since each vertex is in the image $\mathcal{N}_{t,\nu}$, there exists an index tuple $x_{[d]} \in \mathcal{X}$ such that $\nu [X_{[d]} = x_{[d]}] = 1$ and

$$v [L] = \sigma^t [X_{[d]} = x_{[d]}, L] .$$

Then $\kappa^{\mathcal{F}} [Y_{[p]}]$ is the one-hot encoding of the vector $\sigma^t [X_{[d]} = x_{[d]}, L]$, that is

$$\kappa^{\mathcal{F}}[Y_{[p]}] = \epsilon_{\sigma^t}[X_{[d]=x_{[d]},L}] [Y_{[p]}] = \left\langle \epsilon_{\sigma^t}[X_{[d]=x_{[d]},L=\ell}] [Y_{\ell}] : \ell \in [p] \right\rangle_{[Y_{[p]}]} .$$

Such tensors are examples of elementary tensors corresponding to the elementary hypergraph (see Figure 1.1)

$$\text{EL} = ([p], \{\{\ell\} : \ell \in [p]\}) .$$

In particular, the activation core is elementary and the corresponding refined base measure is in $\Lambda^{t,\text{EL},\nu}$.

While vertices are the minimal non-vanishing faces in the face-lattice (see [10]), we now show that the maximal face, namely the polytope itself, is also representable with respect to the elementary hypergraph EL (see Figure 1.1).

Example 1.2 (Maximal face) The maximal face $\mathcal{F}^{\varnothing} = \mathcal{M}_{t,\nu}$ coincides with the mean polytope itself and is given by the choice $\theta_{\varnothing}[L] = 0[L]$. In this case the corresponding activation tensor to the face measure is

$$\kappa^{\varnothing}[Y_{[p]}] = \mathbb{I}[Y_{[p]}] = \bigotimes_{\ell \in [p]} \mathbb{I}[Y_{\ell}] ,$$

where by $\mathbb{I}[Y_{[p]}]$ we denote tensors with all coordinates 1. $\kappa^{\varnothing}Y_{[p]}$ is elementary and the maximal face is always representable by an elementary activation.




Fig. 1.1 Elementary tensor format consisting of the elementary hypergraph EL *a)*, and corresponding tensor networks *b)* being the tensor products of vectors.

1.2.2 Representation in the CP format

Extending Example 1.1 and Example 1.2, we can provide a coarse estimation of the hypergraph $\mathcal{G}$ required to decompose $\kappa^{\mathcal{F}}$ for generic faces $\mathcal{F}$. To this end, we use the CP format describing tensor networks on the CP hypergraph (see Figure 1.2)

$$\text{CP} = ([p] \cup \{I\}, \{\{\ell, I\} : \ell \in [p]\}) .$$

The dimension of the variable I is called the hidden rank of the format.

Lemma 1.4 *Any face $\mathcal{F}$ is representable by a CP graph with hidden rank*

$$r \leq \min \left(|\mathcal{N}_{t,\nu}^{\mathcal{F}}|, |\mathcal{N}_{t,\nu}| - |\mathcal{N}_{t,\nu}^{\mathcal{F}}| + 1 \right) .$$

Proof. We show the claim by constructing two face activating tensor networks to $\mathcal{F}$ in a CP hypergraph with hidden rank $|\mathcal{N}_{t,\nu}^{\mathcal{F}}|$ and in a CP hypergraph with hidden rank $|\mathcal{N}_{t,\nu}| - |\mathcal{N}_{t,\nu}^{\mathcal{F}}| + 1$. To show the first representation we enumerate the vertices by a variable I with dimension $r = |\mathcal{N}_{t,\nu}^{\mathcal{F}}|$, i.e. $\mathcal{N}_{t,\nu}^{\mathcal{F}} = \{v^i[L] : i \in [r]\}$. Then we define for $\ell \in [p]$ core tensors $\tau^{\ell}[Y_{\ell}, I]$

$$\tau^{\ell}[Y_{\ell}, I = i] = \epsilon_{v^i[L=\ell]} [Y_{\ell}] .$$

Then we have

$$\langle \{\tau^\ell [Y_\ell, I] : \ell \in [p]\} \rangle_{[Y_{[p]}]} = \sum_{v \in \mathcal{N}_{t,v}^{\mathcal{F}}} \epsilon_v [Y_{[p]}]$$

and thus have found an activation tensor network in a CP graph with hidden rank $|\mathcal{N}_{t,v}^{\mathcal{F}}|$ representing the face $\mathcal{F}$.

We continue with the second representation, for which we enumerate the set $\mathcal{N}_{t,v}/\mathcal{N}_{t,v}^{\mathcal{F}}$ by $v^i [L]$ where $i \in [|\mathcal{N}_{t,v}| - |\mathcal{N}_{t,v}^{\mathcal{F}}|]$. We define variable I with dimension $r = |\mathcal{N}_{t,v}| - |\mathcal{N}_{t,v}^{\mathcal{F}}| + 1$ and define for $\ell \in [p]$ core tensors

$$\tau^\ell [Y_\ell, I = i] = \begin{cases} -\epsilon_{v^i [L=\ell]} [Y_\ell] & \text{if } i < |\mathcal{N}_{t,v}| - |\mathcal{N}_{t,v}^{\mathcal{F}}| \\ \mathbb{I} [Y_\ell] & \text{if } i = |\mathcal{N}_{t,v}| - |\mathcal{N}_{t,v}^{\mathcal{F}}|. \end{cases}$$

We then have

$$\begin{aligned} \langle \{\tau^\ell [Y_\ell, I] : \ell \in [p]\} \rangle_{[Y_{[p]}]} &= \mathbb{I} [Y_{[p]}] - \sum_{v \in \mathcal{N}_{t,v}/\mathcal{N}_{t,v}^{\mathcal{F}}} \epsilon_v [Y_{[p]}] \\ &= \sum_{v \in \mathcal{N}_{t,v}^{\mathcal{F}}} \epsilon_v [Y_{[p]}] + \sum_{v \in (\times_{\ell \in [p]} [p\ell]) / \mathcal{N}_{t,v}} \epsilon_v [Y_{[p]}]. \end{aligned}$$

We have thus found a face activating tensor network for $\mathcal{F}$ in a CP format with hidden rank $r = |\mathcal{N}_{t,v}| - |\mathcal{N}_{t,v}^{\mathcal{F}}| + 1$. $\square$

We notice that the CP rank bound of Lem. 1.4 is tight in Example 1.1 and Example 1.2, since the elementary format is a restriction of the CP format to rank 1.




Fig. 1.2 The CP tensor format consist of the CP hypergraph a) and decorating tensor networks b). Tensors in this format constructed as in Lem. 1.4 represent arbitrary faces.

1.2.3 Examples of elementarily representable faces

We show in the following example that all faces of a hypercube and a simplex are representable by elementary activation tensors.

Example 1.3 (Hypercube) In cases where $X_{[d]}$ are Boolean and we have $\ell \in [p]$ features

$$f_\ell [X_{[d]} = x_{[d]}] = x_\ell$$

the mean polytope is the hypercube

$$\mathcal{M}_{\{f_\ell : \ell \in [p]\}, \mathbb{I}} = [0, 1]^p.$$

The non-empty faces in the face lattice $L([0, 1]^p)$ can be enumerated by subsets $\mathcal{U} \subset [p]$ and indices $y_{\mathcal{U}} \in \times_{\ell \in \mathcal{U}} [2]$ and represented by the cartesian products

$$\mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})} = \bigtimes_{\ell \in [p]} I^{\ell, (\mathcal{U}, y_{\mathcal{U}})}$$

where

$$\mathcal{I}^{l,(\mathcal{U},y_{\mathcal{U}})} = \begin{cases} [0, 1] & \text{if } \ell \notin \mathcal{U} \\ \{y_{\ell}\} & \text{if } \ell \in \mathcal{U} \end{cases}.$$

Each of these faces can be represented with respect to the elementary graph EL, namely by the tensor product of leg vectors

$$\kappa^{\ell,(\mathcal{U},y_{\mathcal{U}})} [Y_{\ell}] = \begin{cases} \mathbb{I} [Y_{\ell}] & \text{if } \ell \notin \mathcal{U} \\ \epsilon_{y_{\ell}} [Y_{\ell}] & \text{if } \ell \in \mathcal{U} \end{cases}.$$

Example 1.4 (Simplices) The $(p-1)$ -dimensional standard simplex $\mathcal{M}_{\Delta,p-1}$ in $[0, 1]^p$ is the convex hull of the vertex sets

$$\mathcal{N} = \{\epsilon_{\ell} [L] : \ell \in [p]\}.$$

The simplex is the mean polytope of the p -dimensional statistic with

$$t_{\ell} : [p] \rightarrow [2] \quad , \quad t_{\ell} [\tilde{\ell}] = \begin{cases} 1 & \text{if } \tilde{\ell} = \ell \\ 0 & \text{if } \tilde{\ell} \neq \ell \end{cases}.$$

The simplex represents all distributions with respect to the trivial base measure $\mathbb{I}$ which coordinates are all 1. When flattened to a vector, any tensor in $\Lambda^{\mathbb{I}}$ is an element of $\mathcal{M}_{\Delta,p-1}$ where $p = \prod_{k \in [d]} m_k$.

The face lattice of the simplex is enumerated by subsets of $[p]$ as

$$L(\mathcal{M}_{\Delta,p-1}) = \{\mathcal{F}^{\mathcal{U}} : \mathcal{U} \subset [p]\}$$

where the faces are

$$\mathcal{F}^{\mathcal{U}} = \text{conv}(\epsilon_{\ell} [L] : \ell \in \mathcal{U})$$

and are therefore themselves $(|\mathcal{U}| - 1)$ -dimensional standard simplices. The partial order of the faces coincides with the inclusion order of subsets $\mathcal{U}$.

1.2.4 Partition into relative interiors of faces

Let us now introduce relative interiors, which enable us to find disjoint partitions of the mean polytope.

Definition 1.4 (Relative Interior) The relative interior $(\mathcal{U})^{\circ}$ of an arbitrary set $\mathcal{U} \subset \mathbb{R}^p$ is the interior of $\mathcal{U}$ in the affine subspace $\text{conv}(\mathcal{U})$.

Lemma 1.5 *To each $\mu \in \mathcal{M}_{t,v}$ there is a unique face $\mathcal{F}$, which we denote by $\mathcal{F}(\mu)$, such that $\mu \in (\mathcal{F})^{\circ}$.*

Proof. For any mean vector μ on a face $\mathcal{F}$ we either have $\mu \in (\mathcal{F})^{\circ}$ or there is a face $\tilde{\mathcal{F}}$ of lower affine dimension with $\tilde{\mathcal{F}} \subset \mathcal{F}$ and $\mu \in \tilde{\mathcal{F}}$. Starting from the face $\mathcal{M}_{t,v}$ (for which by assumption $\mu \in \mathcal{M}_{t,v}$) we iteratively choose a face $\mathcal{F}$ of lower dimension containing μ until we reach a face with $\mu \in (\mathcal{F})^{\circ}$. This procedure is guaranteed to terminate, since the affine dimension is decreasing and any face of affine dimension 0 coincides with its relative interior. $\square$

1.3 Exponential families

It is well known that exponential families are specific maximum entropy distributions. However, not every maximum entropy distribution is an exponential family member. To prepare for the characterization of generic maximum entropy

distributions, we will in this section represent exponential family members as Computation-Activation Networks. We further show that a maximum entropy distribution is an exponential family member if and only if its mean parameter is in the relative interior of the mean polytope.

1.3.1 Representation as elementary Computation-Activation Networks

Let us first define exponential families as classes of probability distributions and then derive their representation by elementary Computation-Activation Networks.

Definition 1.5 (Exponential Family) Given a statistic function

$$t : \bigtimes_{k \in [d]} [m_k] \rightarrow \mathbb{R}^p$$

and a base measure $\nu [X_{[d]}]$ with $\langle \nu \rangle_{[\emptyset]} \neq 0$, the set $\Gamma^{t, \nu} = \{\mathbb{P}^{(t, \theta, \nu)} : \theta [L] \in \mathbb{R}^p\} \subset \Lambda^\nu$ of probability distributions

$$\mathbb{P}^{(t, \theta, \nu)} [X_{[d]}] = \frac{\exp \left[\langle \sigma^t [X_{[d]}, L], \theta [L] \rangle_{[X_{[d]}]} \right]}{\left\langle \exp \left[\langle \sigma^t [X_{[d]}, L], \theta [L] \rangle_{[X_{[d]}]} \right], \nu [X_{[d]}] \right\rangle_{[\emptyset]}}$$

is called the exponential family with respect to t .

To connect with standard notation of exponential families we notice that the coordinate of any distribution $\mathbb{P}^{(t, \theta, \nu)} [X_{[d]}]$ at some state $x_{[d]} \in \bigtimes_{k \in [d]} [m_k]$ is

$$\mathbb{P}^{(t, \theta, \nu)} [X_{[d]} = x_{[d]}] = \exp \left[\sum_{\ell \in [p]} \theta [L = \ell] \cdot t_\ell [x_{[d]}] - \psi_{t, \nu}(\theta) \right]$$

where

$$\psi_{t, \nu}(\theta) = \ln \left[\left\langle \exp \left[\langle \sigma^t [X_{[d]}, L], \theta [L] \rangle_{[X_{[d]}]} \right], \nu [X_{[d]}] \right\rangle_{[\emptyset]} \right]$$

is the cumulant generating function. The function $\psi_{t, \nu}(\theta)$ and its Legendre transform $\psi_{t, \nu}^*(\mu)$ define the dually flat Riemannian structure on the exponential family manifold [1]. We now continue with the notation of Def. 1.5 to find representations of exponential families by elementary Computation-Activation Networks.

Theorem 1.2 (Exponential Families are elementary Computation-Activation Networks) *Given any base measure ν and a sufficient statistic t we enumerate for each coordinate $\ell \in [p]$ the image $\text{im}(t_\ell)$ by a variable Y_ℓ taking values in $[[\text{im}(t_\ell)]]$, given an interpretation map*

$$I_\ell : [[\text{im}(t_\ell)]] \rightarrow \text{im}(t_\ell) .$$

For any canonical parameter vector $\theta [L] \in \mathbb{R}^p$ we build the activation cores $\alpha^{\ell, \theta} [Y_\ell]$ for each coordinate $y_\ell \in [[\text{im}(t_\ell)]]$ by

$$\alpha^{\ell, \theta} [Y_\ell = y_\ell] = \exp [\theta [L = \ell] \cdot I_\ell [y_\ell]]$$

and have (see Figure 1.3)

$$\mathbb{P}^{(t, \theta, \nu)} [X_{[d]}] = \frac{\langle \{\beta^t [Y_{[p]}, X_{[d]}]\} \cup \{\alpha^{\ell, \theta} [Y_\ell] : \ell \in [p]\} \rangle_{[X_{[d]}]}}{\langle \{\beta^t [Y_{[p]}, X_{[d]}]\} \cup \{\alpha^{\ell, \theta} [Y_\ell] : \ell \in [p]\} \cup \{\nu [X_{[d]}]\} \rangle_{[\emptyset]}} .$$

Proof. For each $x_{[d]} \in \bigtimes_{k \in [d]} [m_k]$ we have




Fig. 1.3 Representation of a member of an exponential family by a Computation-Activation Network with elementary activation. Here $Z \in \mathbb{R}$ is a normalization constant.

$$\begin{aligned}
 & \langle \{\beta^t [Y_{[p]}, X_{[d]}]\} \cup \{\alpha^{\ell, \theta} [Y_\ell] : \ell \in [p]\} \rangle_{[X_{[d]}=x_{[d]}]} \\
 &= \prod_{\ell \in [p]} \langle \beta^{\ell} [Y_\ell, X_{[d]}=x_{[d]}], \alpha^{\ell, \theta} [Y_\ell] \rangle_{[\emptyset]} \\
 &= \prod_{\ell \in [p]} \exp [\theta [L = \ell] \cdot I_\ell [t_\ell [x_{[d]}]]] \\
 &= \exp \left[\sum_{\ell \in [p]} \theta [L = \ell] \cdot I_\ell [t_\ell [x_{[d]}]] \right] \\
 &= \exp \left[\langle \sigma^t [X_{[d]}, L], \theta [L] \rangle_{[X_{[d]}=x_{[d]}]} \right].
 \end{aligned}$$

Therefore we have

$$\langle \{\beta^t [Y_{[p]}, X_{[d]}]\} \cup \{\alpha^{\ell, \theta} [Y_\ell] : \ell \in [p]\} \rangle_{[X_{[d]}]} = \exp \left[\langle \sigma^t [X_{[d]}, L], \theta [L] \rangle_{[X_{[d]}]} \right].$$

The claim follows, since this implies that also the contraction of both sides with $\nu [X_{[d]}]$ is equivalent. $\square$

We will use the following well known property (see e.g. [2]), that the mean parameters of the members of an exponential family are exactly the relative interior of the mean polytope.

Lemma 1.6 *The set of mean parameters of the members of an exponential family is the relative interior of the mean polytope.*

Proof. See The 3.3 in [9]. $\square$

Based on this property we define the forward and backward mappings to an exponential family.

Definition 1.6 The forward map of an exponential family is the map

$$F^{(t, \nu)} : \mathbb{R}^p \rightarrow (\mathcal{M}_{t, \nu})^\circ$$

defined as

$$F^{(t, \nu)}(\theta) = \langle \mathbb{P}^{t, \theta, \nu} [X_{[d]}], \sigma^t [X_{[d]}, L], \nu [X_{[d]}] \rangle_{[L]}$$

Any map $B^{(t, \nu)} : (\mathcal{M}_{t, \nu})^\circ \rightarrow \mathbb{R}^p$ with $F^{(t, \nu)}(B^{(t, \nu)}(\mu)) = \mu$ for all $\mu \in (\mathcal{M}_{t, \nu})^\circ$ is called a backward map.

1.3.2 Maximum entropy distributions

One universal property of exponential families amongst others (see [7]) is that they maximize the entropy among those with the same mean vector.

Lemma 1.7 *Any member of an exponential family $\Gamma^{t,\nu}$ is a maximum entropy distribution with respect to the moment constraint on t and the base measure ν .*

Proof. Let $\mathbb{P}^{t,\theta,\nu}$ be an exponential family member with canonical parameter θ and let $\mu[L]$ be the corresponding mean parameter. Let $\tilde{\mathbb{P}}[X_{[d]}]$ be any feasible distribution for the maximum entropy problem with respect to the moment constraint on t by $\mu[L]$ and the base measure ν . We also have $\langle \tilde{\mathbb{P}}[X_{[d]}], \sigma^t[X_{[d]}, L], \nu[X_{[d]}] \rangle_{[L]} = \mu[L]$ and thus

$$\begin{aligned} \mathbb{H}_\nu[\tilde{\mathbb{P}}, \mathbb{P}^{t,\theta,\nu}] &= -\langle \tilde{\mathbb{P}}[X_{[d]}], \ln[\mathbb{P}^{t,\theta,\nu}[X_{[d]}]], \nu[X_{[d]}] \rangle_{[\emptyset]} \\ &= -\langle \tilde{\mathbb{P}}[X_{[d]}], \sigma^t[X_{[d]}, L], \theta[L], \nu[X_{[d]}] \rangle_{[\emptyset]} + A^{(t,\nu)}(\theta) \\ &= -\langle \theta[L], \mu[L] \rangle_{[\emptyset]} + A^{(t,\nu)}(\theta) \\ &= \mathbb{H}_\nu[\mathbb{P}^{t,\theta,\nu}]. \end{aligned}$$

With the Gibbs inequality we have if $\tilde{\mathbb{P}} \neq \mathbb{P}^{t,\theta,\nu}$

$$\mathbb{H}_\nu[\mathbb{P}^{t,\theta,\nu}] - \mathbb{H}_\nu[\tilde{\mathbb{P}}] = \mathbb{H}_\nu[\tilde{\mathbb{P}}, \mathbb{P}^{t,\theta,\nu}] - \mathbb{H}_\nu[\tilde{\mathbb{P}}] > 0$$

and thus $\mathbb{H}_\nu[\tilde{\mathbb{P}}] < \mathbb{H}_\nu[\mathbb{P}^{t,\theta,\nu}]$. Therefore, any feasible $\tilde{\mathbb{P}}$ different from $\mathbb{P}^{t,\theta,\nu}$ has less entropy than $\mathbb{P}^{t,\theta,\nu}$ and $\mathbb{P}^{t,\theta,\nu}$ is a maximum entropy distribution. $\square$

Together with Lem. 1.6 stating the existence of an exponential family member for any mean parameter in the relative interior of the mean polytope we arrive at the following theorem.

Theorem 1.3 *If and only if $\mu[L]$ is in the relative interior of $\mathcal{M}_{t,\nu}$, then the unique solution of the maximum entropy problem is the distribution*

$$\mathbb{P}^{t,\theta,\nu}[X_{[d]}] \in \Gamma^{t,\nu}$$

where for some backward map $B^{t,\nu}$

$$\theta = B^{t,\nu}(\mu).$$

Proof. By Lem. 1.6 if

$$\mu[L] \in (\mathcal{M}_{t,\nu})^\circ,$$

there is at least one weight parameter $\theta[L] \in \mathbb{R}^p$ with

$$\langle \mathbb{P}^{t,\theta,\nu}[X_{[d]}], \sigma^t[X_{[d]}, L], \nu[X_{[d]}] \rangle_{[L]} = \mu[L]$$

and $B^{t,\nu}(\mu)$ with respect to any backward map $B^{t,\nu}$ is such one. With Lem. 1.7 we conclude that $\mathbb{P}^{(t,B^{t,\nu}(\mu),\nu)}$ is the maximum entropy distribution with respect to the moment constraint on t by $\mu[L]$ and the base measure ν . $\square$

While the maximum entropy distribution is unique, the weight vector θ representing the maximum entropy distribution is not always unique. This is reflected by multiple possible choices of backward mappings. Only if the exponential family is minimal (see [9]) then the weight vector is unique. However, we did not restrict to such situations, since exponential families on refined base measures are not minimal.

1.4 Generic maximum entropy distributions

In the previous section we have shown that the members of an exponential family are maximum entropy distributions. However, while their mean parameters constitute the relative interior of the mean polytope, they never lie on faces of lower dimension. We now exploit the tensor representation study of such faces (see Sect. 1.2) to characterize the maximum entropy distributions in the general case.

1.4.1 Main result

Our main result generalizes the maximum entropy characterization of Thm. 1.3 to arbitrary mean parameters.

Theorem 1.4 (Generic characterization of Maximum Entropy Solutions) *Let t be a statistic and ν a base measure. For any $\theta [L]$ and $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ the distribution*

$$\mathbb{P}^{t,\theta,\nu^{t,\mathcal{F}}} [X_{[d]}]$$

is a maximum entropy distribution.

For any $\mu [L]$ the maximum entropy problem has a feasible distribution, if and only if $\mu [L] \in \mathcal{M}_{t,\nu}$. If $\mu [L] \in \mathcal{M}_{t,\nu}$ the unique maximum entropy distribution is $\mathbb{P}^{t,\theta,\nu^{t,\mathcal{F}}}$ where $\mathcal{F} = \mathcal{F}(\mu)$ (see Lem. 1.5) and

$$\theta = B^{t,\nu^{t,\mathcal{F}}}(\mu)$$

where $\nu^{t,\mathcal{F}}$ is the refined base measure (see Def. 1.2) and $B^{t,\nu^{t,\mathcal{F}}}$ is any backward map to the exponential family $\Gamma^{t,\nu^{t,\mathcal{F}}}$ (see Def. 1.6).

For any hypergraph $\mathcal{G}$ and any face $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ we have that all maximum entropy distributions to mean parameters on $\mathcal{F}^\circ$ are in $\Lambda^{t,\mathcal{G},\nu}$ if $\mathcal{F}$ is representable with respect to $\mathcal{G}$ (see Def. 1.3).

To prepare for the proof of this theorem we first show in an auxiliary lemma that we can reduce the set of feasible distributions in Problem $P_{t,\mu,\nu}$.

Lemma 1.8 *For any $\mu [L] \in \mathcal{M}_{t,\nu}$ and a face $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ with $\mu [L] \in (\mathcal{F})^\circ$ the solutions of $P_{t,\mu,\nu}$ and $P_{t,\mu,\nu^{t,\mathcal{F}}}$ coincide.*

Proof. By Lem. 1.2 all feasible distributions are representable by the base measure refined to the face $\mathcal{F}$. We have that any for $P_{t,\mu,\nu}$ feasible distribution $\mathbb{P} [X_{[d]}]$ satisfies

$$\langle \mathbb{P} [X_{[d]}], \nu^{t,\mathcal{F}} \rangle_{[\emptyset]} = 1$$

and thus $\mathbb{P} [X_{[d]}] \in \Lambda^{\nu^{t,\mathcal{F}}}$. Conversely, any $\mathbb{P} [X_{[d]}] \in \Lambda^{\nu^{t,\mathcal{F}}}$ satisfies

$$\langle \mathbb{P} [X_{[d]}], \nu^{t,\mathcal{F}} \rangle_{[\emptyset]} = \langle \mathbb{P} [X_{[d]}], \nu \rangle_{[\emptyset]} = 1$$

and thus $\mathbb{P} [X_{[d]}] \in \Lambda^\nu$. Problem $P_{t,\mu,\nu}$ is thus equal to

$$\operatorname{argmax}_{\mathbb{P} [X_{[d]}] \in \Lambda^{\nu^{t,\mathcal{F}}}} \mathbb{H}_\nu [\mathbb{P} [X_{[d]}]] \quad \text{subject to} \quad \langle \mathbb{P} [X_{[d]}], \sigma^t [X_{[d]}, L], \nu [X_{[d]}] \rangle_{[L]} = \mu^* [L]$$

We further have that any $\mathbb{P} [X_{[d]}] \in \Lambda^{\nu^{t,\mathcal{F}}}$

$$\mathbb{H}_\nu [\mathbb{P} [X_{[d]}]] = \mathbb{H}_{\nu^{t,\mathcal{F}}} [\mathbb{P} [X_{[d]}]]$$

and arrive together with the above equivalence at the claim. $\square$

Proof of Thm. 1.4. To show that for any $\theta [L]$ and $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ the distribution $\mathbb{P}^{t,\theta,\nu^{t,\mathcal{F}}}$ is a maximum entropy distribution, we apply Lem. 1.8 and Thm. 1.3.

If $\mu [L] \notin \mathcal{M}_{t,\nu}$ then Problem $P_{t,\mu,\nu}$ is not feasible by definition. In case $\mu [L] \in \mathcal{M}_{t,\nu}$ we find $\mathcal{F} \in L(\mathcal{M}_{t,\nu})$ with $\mathcal{F} = \mathcal{F}(\mu)$ and by Lem. 1.8 the solution of maximum entropy problem Problem $P_{t,\mu,\nu}$ coincides with $P_{t,\mu,\nu^{t,\mathcal{F}}}$. Since by Lem. 1.1 the face is the polytope

$$\mathcal{F} = \mathcal{M}_{t,\nu^{t,\mathcal{F}}}$$

we find a weight parameter

$$\theta = B^{t,\nu^{t,\mathcal{F}}}(\mu).$$

and by Thm. 1.3 $\mathbb{P}^{t, \theta, \nu^{t, \mathcal{F}}}$ is the corresponding maximum entropy distribution.

To show the last claim, it is enough to show that for any face $\mathcal{F}$ representable by $\mathcal{G}$ and for any $\theta \in \mathbb{R}^p$ we have

$$\mathbb{P}^{t, \theta, \nu^{t, \mathcal{F}}} \in \Lambda^{t, \mathcal{G}, \nu}.$$

To this end, we construct activation cores as follows. By Def. 1.3 and Lem. 1.3 there is a tensor network $\kappa^{\mathcal{G}}$ such that

$$\nu^{t, \mathcal{F}} [X_{[d]}] = \langle \kappa^{\mathcal{G}} \cup \{ \nu [X_{[d]}] \} \rangle_{[X_{[d]}]}.$$

For each $\ell \in [p]$ we find an edge $e(\ell) \in \mathcal{E}$ with $\ell \in e(\ell)$. Now we define a tensor network $\tau^{\mathcal{G}}$ by the cores to $e \in \mathcal{E}$ as

$$\tau^e [X_e] = \langle \{ \kappa^e [X_e] \} \cup \{ \alpha^{\ell, \theta} [Y_{\ell}] : e = e(\ell) \} \rangle_{[X_e]}.$$

Then, we have

$$\langle \beta^t [Y_{[p]}, X_{[d]}], \langle \tau^{\mathcal{G}} \rangle_{[Y_{[p]}]} \rangle_{[X_{[d]}]} = \langle \beta^t [Y_{[p]}, X_{[d]}], \alpha^{\theta} [Y_{[p]}], \langle \kappa^{\mathcal{G}} \rangle_{[Y_{[p]}]} \rangle_{[X_{[d]}]}$$

and therefore

$$\mathbb{P}^{t, \theta, \nu^{t, \mathcal{F}}} = \frac{\langle \{ \beta^t [Y_{[p]}, X_{[d]}] \} \cup \tau^{\mathcal{G}} \rangle_{[X_{[d]}]}}{\langle \{ \beta^t [Y_{[p]}, X_{[d]}], \nu [X_{[d]}] \} \cup \tau^{\mathcal{G}} \rangle_{[\emptyset]}}.$$

We conclude $\mathbb{P}^{t, \theta, \nu^{t, \mathcal{F}}} \in \Lambda^{t, \mathcal{G}, \nu}$. □

1.4.2 Family of maximum entropy distributions

The family of maximum entropy distributions is the set

$$\Gamma^{t, \nu} = \{ \mathbb{P} [X_{[d]}] : \exists \mu \in \mathcal{M}_{t, \nu} : \mathbb{P} [X_{[d]}] \text{ is a solution of Problem } \mathbf{P}_{t, \mu, \nu} \}.$$

Thm. 1.4 suggests a parametrization of this family as tuples of corresponding faces $\mathcal{F}$ and weight vectors $\theta [L]$. As sketched in Figure 1.4 the activation tensor of each maximum entropy distribution is decomposed into a possibly non-elementary Boolean tensor corresponding to the base measure refinement on a face $\mathcal{F}$, and an elementary tensor dependent on the weight vector $\theta \in \mathbb{R}^p$. We thus define the canonical parameter space

$$\mathcal{P}_{t, \nu} := L(\mathcal{M}_{t, \nu}) \times \mathbb{R}^p$$

and have

$$\Gamma^{t, \nu} = \left\{ \frac{\langle \alpha^{\theta} [Y_{[p]}], \kappa^{\mathcal{F}} [Y_{[p]}], \beta^t [Y_{[p]}, X_{[d]}] \rangle_{[X_{[d]}]}}{\langle \alpha^{\theta} [Y_{[p]}], \kappa^{\mathcal{F}} [Y_{[p]}], \beta^t [Y_{[p]}, X_{[d]}], \nu [X_{[d]}] \rangle_{[\emptyset]}} : \theta [L] \in \mathbb{R}^p, \mathcal{F} \in \mathcal{P}_{t, \nu} \right\}.$$

Note that these families are disjoint, since each member of an exponential family has support by the support of the face measure.

1.5 Boolean statistics

We say that a statistic is Boolean, if $\text{im}(t_{\ell}) \subset \{0, 1\}$ for all $\ell \in [p]$. The mean polytope is then a 0/1-polytope [?], which vertices are exactly the contained Boolean vectors. The special case of Boolean statistics is especially interesting for two reasons. First, we can provide additional intuition on those faces representable by elementary face activating




Fig. 1.4 Tensor network decomposition of maximum entropy distributions to the constraint $\mu[L] = \langle \mathbb{P}[X_{[d]}], \sigma^t[X_{[d]}, L] \rangle_{[L]}$. Blue: Boolean tensor $\kappa^{\mathcal{F}}[Y_{[p]}]$ dependent on the face $\mathcal{F}(\mu)$, which represents the refinement of the base measure to the face, and is possibly non-elementary. Red: Elementary activation tensor $\alpha^{\theta}[Y_{[p]}] = \bigotimes_{s \in [n]} \alpha^{\ell, \theta}[Y_{\ell}]$ to the weight $\theta[L] = B^{t, \nu^{\mathcal{F}}}(\mu)$.

tensors, namely based on intersections with cube faces. Based on these properties we also show that any elementary Computation-Activation Network to a Boolean statistic is a maximum entropy distribution. Second, the statistic itself can be interpreted in terms of logical formulas. We exploit this property to provide examples and to characterize each 0/1-polytope as a mean polytope to a collection of propositional formulas.

1.5.1 Characterization of elementary representable faces

Orienting on Example 1.3 we now introduce cube-likeness of faces and polytopes.

Definition 1.7 We say that a face $\mathcal{F}$ of $\mathcal{M}_{t, \nu}$ is cube-like, if it is empty or there is $\mathcal{U} \subset [p]$ and $y_{\mathcal{U}} \in \times_{\ell \in \mathcal{U}} [2]$ such that

$$\mathcal{F} = \mathcal{M}_{t, \nu} \cap \mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})}.$$

Here we denote by $\mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})}$ the faces of the hypercube $[0, 1]^p$ (see Example 1.3). We further say that a polytope $\mathcal{M}_{t, \nu}$ is cube-like, if all faces $\mathcal{F} \in L(\mathcal{M}_{t, \nu})$ are cube-like.

We now show that a face is cube-like if and only if it is representable by an elementary tensor.

Theorem 1.5 Let t be a Boolean statistic, ν a base measure and $\mathcal{F}$ be a face of $\mathcal{M}_{t, \nu}$. Then the following are equivalent:

- (i) $\mathcal{F}$ is cube-like (see Def. 1.7).
- (ii) $\mathcal{F}$ is representable by an elementary tensor (see Def. 1.3).

Proof. If the face is empty, i.e. $\mathcal{F} = \emptyset$, it is by definition cube-like and has $0[Y_{[p]}]$ as an elementary activation tensor. We therefore assume in the following $\mathcal{F} \neq \emptyset$.

(i) $\Rightarrow$ (ii): Let us assume that $\mathcal{F}$ is cube-like, that is there is $\mathcal{U} \subset [p]$ and $y_{\mathcal{U}} \in \times_{\ell \in \mathcal{U}} [2]$ such that $\mathcal{F} = \mathcal{M}_{t, \nu} \cap \mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})}$. We use that $\mathcal{F}$, $\mathcal{M}_{t, \nu}$ and $\mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})}$ are the convex hulls of the cube vertex sets $\mathcal{N}_{t, \mathcal{M}}^{\mathcal{F}}$, $\mathcal{N}_{t, \mathcal{M}}$ and that for the set

$$\mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}} = \{v[L] : \forall \ell \in \mathcal{U} : v[L = \ell] = y_{\mathcal{U}}\}, \quad (1.1)$$

we have

$$\mathcal{N}_{t, \mathcal{M}}^{\mathcal{F}} = \mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}} \cap \mathcal{N}_{t, \mathcal{M}}.$$

Thus, we can choose $\mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}}$ for the representation of the face $\mathcal{F}$ (see Def. 1.3) and further have that

$$\sum_{v[L] \in \mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}}} \epsilon_{v[L]}[Y_{[p]}] = \bigotimes_{\ell \in [p]} \kappa^{\ell, (\mathcal{U}, y_{\mathcal{U}})}[Y_{\ell}],$$

where

$$\kappa^{\ell, (\mathcal{U}, y_{\mathcal{U}})} [Y_{\ell}] = \begin{cases} \epsilon_{y_{\ell}} [Y_{\ell}] & \text{if } \ell \in \mathcal{U} \\ \mathbb{I} [Y_{\ell}] & \text{if } \ell \notin \mathcal{U}. \end{cases}$$

We have thus found an elementary activation tensor for the face $\mathcal{F}$.

(ii)→(i) Conversely, let $\xi [Y_{[p]}]$ be an elementary activation tensor of the face $\mathcal{F}$ in $\mathcal{M}_{t, \nu}$. Since $\xi [Y_{[p]}]$ is the sum of different one-hot encodings it is Boolean and we find an elementary decomposition $\xi [Y_{[p]}] = \bigotimes_{\ell \in [p]} \xi^{\ell} [Y_{\ell}]$ such that the leg vectors $\xi^{\ell} [Y_{\ell}]$ are Boolean. Since $\mathcal{F} \neq \emptyset$ we further have for $\ell \in [p]$ that $\xi^{\ell} [Y_{\ell}] \neq 0 [Y_{\ell}]$, and thus $\xi^{\ell} [Y_{\ell}] \in \{\epsilon_0 [Y_{\ell}], \epsilon_1 [Y_{\ell}], \mathbb{I} [Y_{\ell}]\}$. We construct a set $\mathcal{U} \subset [p]$

$$\mathcal{U} = \{\ell : \xi^{\ell} [Y_{\ell}] \neq \mathbb{I} [Y_{\ell}]\}$$

and a tuple

$$y_{\ell} = \begin{cases} 1 & \text{if } \xi^{\ell} [Y_{\ell}] = \epsilon_1 [Y_{\ell}] \\ 0 & \text{if } \xi^{\ell} [Y_{\ell}] = \epsilon_0 [Y_{\ell}] \end{cases}.$$

By construction $\xi^{\ell} [Y_{\ell}] = \kappa^{\ell, (\mathcal{U}, y_{\mathcal{U}})} [Y_{\ell}]$ (see Example 1.3) follows for all $\ell \in [p]$ and therefore $\xi [Y_{[p]}] = \kappa^{(\mathcal{U}, y_{\mathcal{U}})} [Y_{[p]}]$. We therefore have for the set (1.1)

$$\xi [Y_{[p]}] = \sum_{v \in \mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}}} \epsilon_v [Y_{[p]}]$$

and since $\xi [Y_{[p]}]$ is an activation tensor for $\mathcal{F}$, that

$$\mathcal{N}_{t, \mathcal{M}}^{\mathcal{F}} = \mathcal{N}^{\mathcal{U}, y_{\mathcal{U}}} \cap \mathcal{N}_{t, \mathcal{M}}.$$

Taking convex hulls on both sides, this is equivalent to

$$\mathcal{F} = \mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})} \cap \mathcal{M}_{t, \nu}.$$

We conclude that $\mathcal{F}$ is cube-like. □

Thm. 1.5 implies that the maximum entropy distributions to hypercubes (see Example 1.3) and simplices (see Example 1.4) are in $\Lambda^{t, \text{EL}, \nu}$, since these polytopes are cube-like.

We can further show, that in case of Boolean statistics any distribution in $\Lambda^{t, \text{EL}, \nu}$ is a maximum entropy distribution.

Theorem 1.6 *Let t be a Boolean statistic. Any distribution in $\Lambda^{t, \text{EL}, \nu}$ is a maximum entropy distribution with respect to (t, μ, ν) where μ is its mean parameter.*

Proof. Let $\mathbb{P} \in \Lambda^{t, \text{EL}, \nu}$ be arbitrary. We then find an elementary tensor $\bigotimes_{s \in [n]} \tau^{\ell} [Y_{\ell}]$ and $\mathcal{Z} \in \mathbb{R}$ such that

$$\mathbb{P} [X_{[d]}] = \frac{1}{\mathcal{Z}} \left\langle \beta^t [Y_{[p]}, X_{[d]}], \bigotimes_{s \in [n]} \tau^{\ell} [Y_{\ell}] \right\rangle_{[X_{[d]}]}.$$

We then define for each $\ell \in [p]$ the vector $\kappa^{\ell} [Y_{\ell}]$ as the support of $\tau^{\ell} [Y_{\ell}]$ and

$$\theta [L = \ell] = \begin{cases} \ln \left[\frac{\tau^{\ell} [Y_{\ell}=1]}{\tau^{\ell} [Y_{\ell}=0]} \right] & \text{if } \tau^{\ell} [Y_{\ell}=0], \tau^{\ell} [Y_{\ell}=1] \neq 0 \\ 0 & \text{else} \end{cases}.$$

Since t is a Boolean statistic the $\bigotimes_{\ell \in [p]} \kappa^{\ell} [Y_{\ell}]$ is a face activation tensor to the face

$$\mathcal{F} := \mathcal{M}_{t, \mathbb{I}} \cap \mathcal{F}^{(\mathcal{U}, y_{\mathcal{U}})}$$

where $\mathcal{U} = \{\ell : \kappa^{\ell} [Y_{\ell}] \neq \mathbb{I} [Y_{\ell}]\}$ and for $\ell \in \mathcal{U}$

$$y_\ell = \begin{cases} 0 & \text{if } \kappa^\ell [Y_\ell = 1] = 0 \\ 1 & \text{else} \end{cases}.$$

Therefore, $\mathbb{P}$ is the maximum entropy distribution to the canonical parameters $\mathcal{F}$ and θ . $\square$

1.5.2 Interpretation by propositional formulas

Let us now assume that $m_k = 2$ for any $k \in [d]$. Following the formalism in [?] we understand each coordinate t_ℓ of the statistic as a propositional formula and the variables $X_{[d]}$ as atoms of a propositional theory. We use propositional connectives for syntactical representations of the formulas and denote by $\neg^1$ the logical negation and by $\neg^0$ the identity. A Boolean tensor $f[X_{[d]}]$ is called satisfiable, if there is at least one $x_{[d]} \in \times_{k \in [d]} [m_k]$ with $f[X_{[d]} = x_{[d]}] = 1$.

Theorem 1.7 *Let t be a Boolean statistic. Any Boolean vector $v[L] \in \{0, 1\}^P$ is a vertex of $\mathcal{M}_{t, \mathbb{I}}$ if and only if*

$$f_v[X_{[d]}] := \bigwedge_{\ell \in [p]} \neg^{1-v[L=\ell]} f_\ell[X_{[d]}]$$

is satisfiable.

Proof. Let $v[L] \in \{0, 1\}^P$ be arbitrary. If and only if $v[L]$ is a vertex of $\mathcal{M}_{t, \mathbb{I}}$ we find $x_{[d]} \in \times_{k \in [d]} [m_k]$ with $v[L] = \sigma^t X_{[d]} = x_{[d]}, L$. This is further equivalent to $f_\ell[X_{[d]} = x_{[d]}] = v[L = \ell]$ for any $\ell \in [p]$, $\neg^{1-v[L=\ell]} f_\ell[X_{[d]} = x_{[d]}] = 1$ for any $\ell \in [p]$, and $f_v[X_{[d]} = x_{[d]}] = 1$. $\square$

Since the presence of Boolean vectors in $\mathcal{M}_{t, \mathbb{I}}$ is related by Thm. 1.7 to the satisfiability of the conjunction $f_v[X_{[d]}]$, the absence is related to entailment statements. To be more precise, for any $v[L] \in \{0, 1\}^P$ and arbitrary $\mathcal{U} \subset [p]$ we have $v[L] \notin \mathcal{M}_{t, \mathbb{I}}$, if and only if

$$\left(\bigwedge_{\ell \in \mathcal{U}} \neg^{1-v[L=\ell]} f_\ell[X_{[d]}] \right) \models \left(\bigvee_{\ell \notin \mathcal{U}} \neg^{v[L=\ell]} f_\ell[X_{[d]}] \right).$$

By definition (see [?]) the latter holds if and only if

$$\left(\bigwedge_{\ell \in \mathcal{U}} \neg^{1-v[L=\ell]} f_\ell[X_{[d]}] \right) \wedge \neg \left(\bigvee_{\ell \notin \mathcal{U}} \neg^{v[L=\ell]} f_\ell[X_{[d]}] \right) = f_v[X_{[d]}]$$

is not satisfiable.

Let us now relate two well-studied sets of formulas, namely atomic formulas and minterm formulas, with two well-studied polytopes, namely hypercubes and simplices.

Example 1.5 (Atomic formulas) Let us consider the case of atomic formulas $f_\ell[X_{[d]}]$ defined with the coordinates at $x_{[d]} \in \times_{k \in [d]} [m_k]$ by

$$f_\ell[X_{[d]} = x_{[d]}] = x_\ell.$$

The mean polytope in this case is the d -dimensional hypercube (see Example 1.3)

$$\mathcal{M}_{\mathcal{F}_{[d]}, \mathbb{I}} = [0, 1]^P$$

which is called a simple polytope, since each vertex is contained in the minimal number of d facets.

Each face is characterized by the projections onto each variable, which is either $\{0\}$, $\{1\}$ or $[0, 1]$. The projections are represented by the tuple $(\mathcal{U}, y_{\mathcal{U}})$ defined in the following way:

- We define the set $\mathcal{U} \subset [d]$ of variables, such that the projection onto the variable is $\{0\}$ or $\{1\}$
- We define to each $\ell \in \mathcal{U}$ an index $y_\ell = 0$ if the projection is $\{0\}$ and $y_\ell = 1$ if the projection is $\{1\}$.

Trivially, each face of the hypercube is a cube face.

More generally, the mean polytope to a collection $f_\ell [X_{[d]}]$ of propositional formulas is the hypercube, if and only if for any Boolean vector $v [L]$ we have

$$\left\langle \bigwedge_{\ell \in [p]} \neg^{1-v[L=\ell]} f_\ell [X_{[d]}] \right\rangle_{[\emptyset]} \neq 0.$$

This means that no conjunction of possibly negated formulas from this statistic entails nor contradicts another formula.

Example 1.6 (Minterm formulas build simplices) The set of minterm formulas is indexed by $v [L] \in \{0, 1\}^p$ and given by

$$f_v [X_{[d]}] = \bigwedge_{\ell \in [d]} \neg^{1-v[X_{[d]}=x_{[d]}]} X_\ell = \epsilon_{v[L]} [X_{[d]}]$$

where by X_ℓ we denote the ℓ -th atomic formula (see Example 1.5). The mean polytope to the minterm statistic and base measure ν is the standard simplex of dimension

$$|\{v [L] : v \in \{0, 1\}^p, \nu [X_{[d]} = v [L]] \neq 0\}| - 1,$$

that is the set

$$\mathcal{M}_{\mathcal{F}_\wedge, \nu} = \text{conv} (\epsilon_{v[L]} [Y_{[p]}] : v \in \{0, 1\}^p, \nu [X_{[d]} = v [L]] \neq 0).$$

We notice that for Boolean base measures $\mathcal{M}_{\mathcal{F}_\wedge, \nu} = \Delta^\nu$. The simplex is further cube-like, since each face is indexed by a subset $\mathcal{U} \subset \{0, 1\}^p$ of vertices and the intersection of the simplex with the cube face $\mathcal{F}^{(\{0, 1\}^p \setminus \mathcal{U}, 1_{\{0, 1\}^p \setminus \mathcal{U}})}$.

More generally, the mean polytope to a Boolean statistic is a standard simplex, if and only if each target coordinate restriction $f_{\tilde{\ell}}$ contradicts all others, that is

$$\forall \ell \neq \tilde{\ell} \in [p] : f_\ell \models \neg f_{\tilde{\ell}}.$$

While hypercubes and simplices are cube-like, we now provide examples of statistics, which polytopes are not cube-like.

Example 1.7 (Maximum entropy distribution with non-elementary activation cores)

Consider two atomic variables X_0 and X_1 and a statistic $\mathcal{F}$ consisting of the formulas

$$f_0 [X_0, X_1] = (X_0 \wedge X_1) \quad , \quad f_1 [X_0, X_1] = (X_0 \Rightarrow X_1)$$

with the coordinatewise expressions

$$f_0 [X_0, X_1] = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} [X_0, X_1] \quad , \quad f_1 [X_0, X_1] = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} [X_0, X_1].$$

We can think of X_0 as a feature on an invoice, and X_1 as a feature on the accounting proposal.

From this we have

$$\begin{aligned} \beta^{(f_0, f_1)} [Y_0 = 0, Y_1 = 0, X_0, X_1] &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} [X_0, X_1] \quad , \\ \beta^{(f_0, f_1)} [Y_0 = 0, Y_1 = 1, X_0, X_1] &= \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} [X_0, X_1] \quad , \\ \beta^{(f_0, f_1)} [Y_0 = 1, Y_1 = 0, X_0, X_1] &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} [X_0, X_1] \quad \text{and} \\ \beta^{(f_0, f_1)} [Y_0 = 1, Y_1 = 1, X_0, X_1] &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} [X_0, X_1]. \end{aligned}$$

Since the only vanishing slice of $\beta^{\mathcal{F}}$ with respect to the head variables is that to $y_{0,1} = (1, 0)$, the vertices of the mean polytope are the vectors to the other head indices. The mean polytope is the convex hull of these vertices

$$\mathcal{M}_{(f_0, f_1)} = \text{conv} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix} [L], \begin{pmatrix} 0 \\ 1 \end{pmatrix} [L], \begin{pmatrix} 1 \\ 1 \end{pmatrix} [L] \right).$$

This polytope has a non cube-like face (sketched blue in Figure 1.5), which is the convex hull of the vertices $(00)^T, (11)^T$. This face is parametrized by the (CP-rank 2) hard activation core

$$\kappa^{(0,0),(1,1)} [Y_0, Y_1] = \epsilon_{(0,0)} [Y_0, Y_1] + \epsilon_{(1,1)} [Y_0, Y_1] = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} [Y_0, Y_1]$$

and has the refined base measure

$$\langle \kappa^{(0,0),(1,1)} [Y_0, Y_1], \beta^{\mathcal{F}} [Y_0, Y_1, X_0, X_1] \rangle_{[X_0, X_1]} = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix} [X_0, X_1].$$

Any mean parameter μ on the interior of that face can be parametrized by a scalar $\lambda \in (0, 1)$

$$\mu_{\lambda} [L] = (\lambda \ \lambda) [L].$$

With the canonical parameters $\theta [L] \in \mathbb{R}^2$ of the maximum entropy distributions on this face by

$$\mathbb{P} [X_0, X_1] = \frac{1}{1 + \exp [\theta [L = 0] + \theta [L = 1]]} \begin{pmatrix} 0 & 0 \\ 1 \exp [\theta [L = 0] + \theta [L = 1]] \end{pmatrix} [X_0, X_1]$$

we get the correspondence by the sigmoid

$$\lambda = \frac{1}{1 + \exp [-(\theta [L = 0] + \theta [L = 1])]}.$$

Note, that the hard activation core $\kappa^{(0,0),(1,1)} [Y_0, Y_1]$ to the blue face is the only non-elementary activation core. While the vertices have always elementary cores, the other non-vertex faces have elementary activation cores

$$\begin{aligned} \kappa^{(0,0),(1,0),(1,1)} [Y_0, Y_1] &= \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} [Y_0, Y_1] = \mathbb{I} [Y_0] \otimes \mathbb{I} [Y_1], \\ \kappa^{(0,0),(1,0)} [Y_0, Y_1] &= \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} [Y_0, Y_1] = \mathbb{I} [Y_0] \otimes \epsilon_0 [Y_1], \\ \kappa^{(1,0),(1,1)} [Y_0, Y_1] &= \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} [Y_0, Y_1] = \epsilon_0 [Y_0] \otimes \mathbb{I} [Y_1]. \end{aligned}$$

The maximum entropy distributions to mean parameters on the interior of all other faces than the blue face are represented by Computation-Activation Networks with only elementary activation cores. The corresponding lattice is sketched in Figure 1.6.

1.5.3 Faces activated in the TT format

The TT format is a well-studied tensor network format with favorable numerical properties [?] and applied across several scientific areas [?]. This format describes tensor networks on a hypergraph (see Figure 1.7)

$$\text{TT} = (X_{[d]} \cup I_{[d-1]}, \{ \{I_{k-1}, X_k, I_k\} : k \in \{1, \dots, d-2\} \} \cup \{ \{X_0, I_0\}, \{I_{d-2}, X_{d-1}\} \}),$$

where $I_{[d-1]}$ are auxiliary hidden variables. We now construct an example of a polytope with a face activatable with TT, in which all hidden variables are of dimension 2.




Fig. 1.5 a) Mean polytope of the statistic $t = (X_0 \wedge X_1, X_0 \Rightarrow X_1)$ (thick), as a subset of the cube $[0, 1]^2$ (dashed). The blue line is the face of the polytope, which is not cube-like, that is not an intersection of the polytope with the faces of the hypercube. We further define for $\lambda \in (0, 1)$ a mean parameter $\mu_\lambda [L] = [\lambda \lambda]^T$ which is on the interior of the blue face. b) Corresponding Computation-Activation Network being the maximum entropy distribution reproducing $\mu_\lambda [L]$, when λ is the sigmoid of $\theta [L = 0] + \theta [L = 1]$.




Fig. 1.6 Face lattice $L((X_0 \wedge X_1, X_0 \Rightarrow X_1))$ to Example 1.7. The directed arrows represent inclusion of the faces, which is a partial order of the faces. The face $\mathcal{F}^{(0,0),(1,1)}$ is the only face, which is not representable by an elementary hypergraph. It is representable in a CP with hidden rank 2




Fig. 1.7 TT format consisting of the TT hypergraph a) and decorating tensor networks b).

Example 1.8 (TT representation of diagonal faces) Consider the vertex set

$$\mathcal{N} := \{0, 1\}^p / v [L]$$

for some Boolean vector $v [L]$. We are interested in the face $\mathcal{F}^{\mathcal{N}, 1}$ with the normal $\theta [L] = \frac{1}{2}(2v [L] - \mathbb{I} [L])$. Its vertices are the p elements of $\{0, 1\}^p$, which differ from $v [L]$ in exactly one coordinate (i.e. those with Hamming distance of 1 from $v [L]$). We denote these vertices by v^ℓ for $\ell \in [p]$, which coordinates to $\tilde{\ell} \in [p]$ are

$$v^\ell [L = \tilde{\ell}] = \begin{cases} v [L = \tilde{\ell}] & \text{if } \tilde{\ell} \neq \ell \\ 1 - v [L = \tilde{\ell}] & \text{if } \tilde{\ell} = \ell \end{cases}.$$

We now represent their sum in an TT format. The Boolean hidden variables are denoted by $I_{[p-1]}$ and are interpreted as indicators, whether the coordinate flip has happened in $[\ell]$ coordinates. We construct a TT core for $\ell = 0$ by

$$\tau^0 [Y_0, I_0] = \epsilon_{v[L=0]} [Y_0] \otimes \epsilon_0 [I_0] + \epsilon_{1-v[L=0]} [Y_0] \otimes \epsilon_1 [I_0],$$

for $\ell \notin \{0, p-1\}$ the cores

$$\begin{aligned} \tau^\ell [I_{\ell-1}, Y_\ell, I_\ell] &= \epsilon_0 [I_{\ell-1}] \otimes \epsilon_{1-v[L=\ell]} [Y_\ell] \otimes \epsilon_1 [I_\ell] + \epsilon_0 [I_{\ell-1}] \otimes \epsilon_{v[L=\ell]} [Y_\ell] \otimes \epsilon_0 [I_\ell] \\ &\quad + \epsilon_1 [I_{\ell-1}] \otimes \epsilon_{v[L=\ell]} [Y_\ell] \otimes \epsilon_1 [I_\ell] \end{aligned}$$

and for $\ell = p - 1$

$$\tau^{p-1} [I_{p-1}, Y_{p-1}] = \epsilon_0 [I_{p-1}] \otimes \epsilon_{1-v[L=p-1]} [Y_{p-1}] + \epsilon_1 [I_{p-1}] \otimes \epsilon_{v[L=p-1]} [Y_{p-1}] .$$

For this tensor network in the TT format we then have

$$\sum_{\ell \in [p]} \epsilon_{v^\ell} [Y_{[p]}] = \langle \{ \tau^0 [Y_0, I_0], \tau^{p-1} [I_{p-2}, Y_{p-1}] \} \cup \{ \tau^\ell [I_{\ell-1}, Y_\ell, I_\ell] : \ell \in \{1, \dots, p-2\} \} \rangle_{[Y_{[p]}]} .$$

The dimension 2 to the hidden variables is furthermore minimal, since each matricification based on a partition of $[p]$ into non-empty sets has a matrix rank of 2 [?].

1.5.4 Construction of Boolean statistics to polytopes

We now show that any convex polytope with Boolean vertices in $\mathbb{R}^p$ is the mean polytope to a Boolean statistic. To construct the corresponding formulas we denote by $\mathcal{M}|_{\mathcal{U}}$ the restriction of the vectors in the set $\mathcal{M}$ to the coordinates in $\mathcal{U}$.

Theorem 1.8 *Let $\mathcal{M}$ be an arbitrary polytope with Boolean vertices in $\mathbb{R}^p$. Then we construct propositional formulas on atoms $X_{[p]}$ by*

$$f_0 [X_0] = \begin{cases} \top & \text{if } \mathcal{M}|_{\{0\}} = \{1\} \\ \perp & \text{if } \mathcal{M}|_{\{0\}} = \{0\} \\ X_0 & \text{if } \mathcal{M}|_{\{0\}} = [0, 1] \end{cases}$$

and iteratively for $\ell \in [p]$ with $\ell \geq 1$ by

$$\begin{aligned} f_\ell [X_{\ell+1}] &= \bigwedge_{v[L] \in (\mathcal{M} \cap \{0,1\}^p)|_{[\ell]}} \left(\left(\bigwedge_{\tilde{\ell} \in [\ell]} \neg^{1-v[L=\tilde{\ell}]} f_{\tilde{\ell}} [X_{\tilde{\ell}+1}] \right) \Rightarrow \right. \\ &\quad \left. \begin{cases} \top & \text{if } (\mathcal{M} \cap v \times \mathbb{R}^{p-\ell})|_{\{\ell\}} = \{1\} \\ \perp & \text{if } (\mathcal{M} \cap v \times \mathbb{R}^{p-\ell})|_{\{\ell\}} = \{0\} \\ X_\ell & \text{if } (\mathcal{M} \cap v \times \mathbb{R}^{p-\ell})|_{\{\ell\}} = [0, 1] \end{cases} \right). \end{aligned}$$

Here we denote by $\mathcal{M}|_V$ the projections of the vertices in $\mathcal{M}$ onto the subspaces V , and by 0_p the zero vector in $\mathbb{R}^p$.

Proof. We show per induction, that for any $\ell \in [p]$ the family of Hybrid Logic Networks with the statistic $f_{[\ell+1]}$ by the first $\ell + 1$ formulas has the mean polytope

$$\mathcal{M}_{f_{[\ell+1]}, \mathbb{I}} = \mathcal{M}|_{[\ell]} . \quad (1.2)$$

“ $\ell = 0$ ”: The polytope $\mathcal{M}|_{[\ell]} = \{1\}$ (respectively $\mathcal{M}|_{[\ell]} = \{0\}$) is reproduced by f_0 being a tautology (respectively a contradiction). In the case $\mathcal{M}|_{[\ell]} = [0, 1]$ the polytope is reproduced by the any formula, which is neither a tautology nor a contradiction, and the atomic formula X_0 is an example of such a contingency. Since the projection of a 0-1 polytope onto the first coordinates is itself a 0-1 polytope, these are the only possible cases and we conclude that in all

$$\mathcal{M}_{f_{[\ell]}, \mathbb{I}} = \mathcal{M}|_{[\ell]} .$$

“ $\ell \rightarrow \ell + 1$ ”: We define for any $v \in \mathcal{M}_{f_{[\ell]}, \mathbb{I}} \cap \{0, 1\}^\ell$ the formula

$$f_{\ell,v} [X_{[\ell+1]}] := \bigwedge_{\tilde{\ell} \in [\ell]} \neg^{1-v[L=\tilde{\ell}]} f_{\tilde{\ell}} [X_{[\tilde{\ell}+1]}] .$$

and notice that all formulas are satisfiable (since the corresponding vertices are in the $\mathcal{M}_{f_{[\ell],\mathbb{I}}}$) and have disjoint models. We have for any Boolean vector $v \in \{0, 1\}^\ell$ and $\lambda \in \{0, 1\}$ that $v \times \{\lambda\} \in \mathcal{M}_{[\ell+1]}$, if and only if

$$v \in \mathcal{M}_{[\ell]} \quad \text{and} \quad \lambda \in \mathcal{M} \cap v \times \mathbb{R}^{p-\ell}|_{[\ell]} .$$

With modus ponens on $f_{\ell+1} [X_{[\ell+2]}]$ this is further equivalent to the satisfaction of

$$f_{\ell,v} [X_{[\ell+1]}] \wedge \neg^{1-\lambda} f_{\ell+1} [X_{[\ell+2]}]$$

and thus to $v \times \{\lambda\} \in \mathcal{M}_{f_{[\ell+1],\mathbb{I}}}$. The vertices of the 0/1 polytopes $\mathcal{M}_{f_{[\ell+1],\mathbb{I}}}$ and $\mathcal{M}_{[\ell+1]}$ thus coincide and both polytopes are therefore equivalent.

By induction the equation (1.2) holds for arbitrary $\ell \in [p]$. For $\ell = p - 1$ this is the claim. $\square$

1.5.5 Examples in statistical physics and coding theory

Let us now investigate cube-like faces to standard examples in statistical physics and coding theory.

Example 1.9 (Ising model on 2 nodes) Consider two Boolean variables X_0, X_1 and the Ising statistic $t = (f_0, f_1, f_2)$ by three propositional formulas


$$f_0 [X_{[2]}] = X_0 \quad , \quad f_1 [X_{[2]}] = X_1 \quad \text{and} \quad f_2 [X_{[2]}] = X_0 \wedge X_1 .$$

In the Ising interpretation, the Boolean variables represent interacting spins at two locations. Their value is then measured by the first two formulas and their interaction by the third.

The vertices of the mean polytope to this statistic are

$$\begin{aligned} \sigma^t [X_{[2]} = (0, 0), L] &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} [L] \quad , \quad \sigma^t [X_{[2]} = (0, 1), L] = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} [L] \quad , \\ \sigma^t [X_{[2]} = (1, 0), L] &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} [L] \quad \text{and} \quad \sigma^t [X_{[2]} = (1, 1), L] = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} [L] \quad . \end{aligned}$$

The mean polytope is the convex hull of these, sketched as:



We notice, that the convex hull of any three out of the four vertices builds a facet. The only cube-like facet of these four (sketched in cyan in the above plot) is the convex hull of

$$\text{conv} \left(\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} [L], \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} [L], \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} [L] \right)$$


to which we have the parametrization by $(\mathcal{U}, y_{\mathcal{U}}) = (\{2\}, 0)$.

As a consequence, there are maximum entropy distributions of mean parameters in the Ising statistics, which do not have a representation by an elementary CompActNet in the Ising statistic.

Example 1.10 (Codeword Polytopes) Consider the codeword polytope with the vertices

$$\mathcal{N} = \left\{ y_{[3]} : \sum_{k \in [3]} y_k \mod 2 = 0 \right\} = \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} [L], \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} [L], \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} [L], \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} [L] \right\}$$

We interpret the vertices by the strings of length 3, which parity vanishes. Such constraints are common in the construction of codewords using parity-check codes (see [?]).



All facets are non-cube-like, whereas all other faces are cube-like (the edges are labeled by the 6 facets of $[0, 1]^3$).

Each facet is characterized by the Hamming distance of exactly 1 to a single of the 4 cube vertices, which are not codewords (see Example 3.9 in [9]). We can construct TT activation tensors with rank 2, which calculate the Hamming distance to these vertices via hidden ranks. Each core tensor in that format adds the contribution of a mode to the Hamming distance.

In more generality, Low-Density-Parity-Checking Codes (see [?]) construct codewords by a collection of parity constraints on subsets of variables. Such polytopes are also known as cycle polytopes, and studied e.g. in [4].

1.6 Mean as a Statistic

We in this section take the perspective of estimating a maximum entropy distribution given observed data.

- μ_D is an unbiased estimator of μ , i.e. $\mathbb{E}[\mu_D[L]] = \mu$
- μ_D is a consistent estimator of μ , i.e. $\mu_D[L] \rightarrow \mu$ by the law of large numbers coordinatewise almost everywhere.
- μ_D is the minimal variance unbiased estimator of μ .

The mean parameter μ_D given a dataset can be understood as a statistic of the dataset. We here show that for the family of maximum entropy distributions this statistic is a minimal sufficient statistic.

The family of maximum entropy distributions is the set

$$\{\mathbb{P}^\mu[X_{[d]}] : \mu \in \mathcal{M}_{t,v}\}$$

which has been characterized above by a union of exponential families with respect to face measures.

Taking a frequentist perspective we now understand datasets by random variables $X_{[d] \times [m]}$, where for $j \in [m]$ the variables $X_{[d],j}$ are drawn i.i.d. from a maximum entropy distribution. The mean statistic is then a tensor

$$\mu_D[X_{[d] \times [m]}, L]$$

with coordinates

$$\mu_D[X_{[d] \times [m]} = x_{[d] \times [m]}, L] = \frac{1}{m} \sum_{j \in [m]} \sigma^t[X_{[d],j} = x_{[d],j}, L].$$

Theorem 1.9 *The mean statistic is a sufficient statistic for the family of maximum entropy distributions (t, ν) . If the base measure has maximal support, the mean statistic is in addition minimal.*

Proof. It suffices to show that the likelihood is a function of μ_D . Let us choose a face $\mathcal{F}$ of $\mathcal{M}_{t,\nu}$, then the likelihood is different from 0 if and only if the empirical distribution is representable with respect to the face measure. This is the case if and only if μ_D is on the face. In case that μ_D is on the face, then the likelihood of any distribution on that face exponential family is

$$\exp \left[m \cdot \left(\langle \mu_D [X_{[d] \times [m], L}], \theta [L] \rangle_{[\emptyset]} - A^{(t,\nu)}(\theta) \right) \right]$$

We have thus shown that the likelihood is always a function of μ_D .

The minimality claim can be shown by the constant quotient criterion (see Thm. 6.2.13 in [3]). $\square$

1.7 Face Computation

We take the perspective of computation whether a vertex is in a face. Based on this perspective, we derive activation tensors to faces. Note, that we construct face activation tensors based on DFA computations here. Thus the decomposition studies of the circuits and automata chapter are relevant.

1.7.1 Examples

Example 1.11 (Generalization of Example 1.8 to larger TT ranks)

We understand this as the tensor representation of an automaton counting the cutted Hamming distance between two states.

We now generalize the construction of Example 1.8 by using the Hadamard distance $d(\cdot, \cdot)$ in $\{0, 1\}^p$, which counts the number of coordinates two vertices differ in. For $s \in \{1, \dots, p-1\}$ we define for a fixed $v [L] \in \{0, 1\}^p$ a polytope as the convex hull

$$\{\tilde{v}[L] : d(\tilde{v}, v) \geq s\}.$$

The face to the normal $\frac{1}{2}(2v [L] - \mathbb{I} [L])$ is the convex hull

$$\mathcal{F}^{a,s} := \text{conv}(\{\tilde{v}[L] : d(\tilde{v}, v) = s\})$$

containing $\binom{p}{s}$ vertices. We label these vertices by subsets $\mathcal{U} \subset [p]$ of cardinality s and define for $\ell \in [p]$

$$v^{\mathcal{U}} [L = \ell] = \begin{cases} v [\ell \in [p]] & \text{if } \ell \notin \mathcal{U} \\ 1 - v [\ell \in [p]] & \text{if } \ell \in \mathcal{U} \end{cases}$$

We now construct a TT with hidden variables I_ℓ and ranks $r_\ell = \min(\ell + 1, p - \ell + 1, s + 1)$ to represent the sum of their one-hot encodings. To this end, let there be TT cores for $\ell = 0$ by

$$\tau^0 [Y_0, I_0] = \epsilon_{v[L=0]} [Y_0] \otimes \epsilon_0 [I_0] + \epsilon_{1-v[L=0]} [Y_0] \otimes \epsilon_1 [I_0]$$

further for $\ell \notin \{0, p-1\}$ the cores

$$\tau^\ell [I_{\ell-1}, Y_\ell, I_\ell] = \sum_{\ell \in \{\max(s-p+\ell, 0), \dots, \max(\ell, s)\}} \epsilon_\ell [I_\ell] \otimes (\epsilon_{v[L=\ell]} [Y_\ell] \otimes \epsilon_\ell [I_{\ell+1}] + \epsilon_{1-v[L=\ell]} [Y_\ell] \otimes \epsilon_{\ell+1} [I_{\ell+1}])$$

and for $\ell = p-1$

$$\tau^{p-1} [I_{p-1}, Y_{p-1}] = \epsilon_0 [I_{p-1}] \otimes \epsilon_{1-v[L=p-1]} [Y_{p-1}] + \epsilon_1 [I_{p-1}] \otimes \epsilon_{v[L=p-1]} [Y_{p-1}].$$

Based on the interpretation, that the hidden variables I_ℓ count the Hamming distance of the vectors $v[L]_{|\mathbb{R}^\ell \times 0_{p-\ell}}$ and the respective $v^{\mathcal{U}}[L]_{|\mathbb{R}^\ell \times 0_{p-\ell}}$ one can show that

$$\sum_{\mathcal{U} \subset [p] : |\mathcal{U}|=s} \epsilon_{v^{\mathcal{U}}} [Y_{[p]}] = \langle \{ \tau^0 [Y_0, I_0], \tau^{p-1} [I_{p-1}, Y_{p-1}] \} \cup \{ \tau^\ell [I_{\ell-1}, Y_\ell, I_\ell] : \ell \notin \{0, p-1\} \} \rangle_{[Y_{[p]}]} .$$

Example 1.12 (Generalization of Example 1.11 to arbitrary HT formats) Instead of aligning the Hamming count variables linearly, one can find a representation of the activation tensor

$$\sum_{\mathcal{U} \subset [p] : |\mathcal{U}|=s} \epsilon_{v^{\mathcal{U}}} [Y_{[p]}]$$

in an arbitrary directed acyclic tree hypergraph format $\mathcal{G}$, which hidden ranks are bounded by the number of leafs in the subtree and $s+1$.

At any leaf of the tree we define a 2×2 matrix

$$\tau^\ell [Y_\ell, I_\ell] = \epsilon_{v[L=\ell]} [Y_\ell] \otimes \epsilon_0 [I_\ell] + \epsilon_{1-v[L=\ell]} [Y_\ell] \otimes \epsilon_1 [I_\ell] .$$

At each intermediate non-root hyperedge we choose an outgoing counting variable $I_{e^{\text{out}}}$ with dimension by $n_{e^{\text{out}}} = \min(\sum_{v \in e^{\text{in}}} n_{e^{\text{in}}}, s+1)$ and define a tensor with the slices

$$\tau^e [I_{e^{\text{out}}}, I_{e^{\text{in}}}] = \beta^+ [Y_+ = i_{e^{\text{out}}}, I_{e^{\text{in}}}] .$$

We further build at the root hyperedge e a tensor

$$\tau^e [I_e] = \sum_{i : \langle i \rangle_{[\emptyset]}=s} \epsilon_{i_e} [I_e] .$$

Using the counting variable interpretation one can now show that

$$\sum_{\mathcal{U} \subset [p] : |\mathcal{U}|=s} \epsilon_{v^{\mathcal{U}}} [Y_{[p]}] = \langle \{ \tau^e [I_e] : e \in \mathcal{E} \} \cup \{ \tau^\ell [Y_\ell, I_\ell] : v \in \mathcal{V} \} \rangle_{[Y_{[p]}]} .$$

1.7.2 Generic Construction

When the vertices are characterized by the sum of local terms, we can construct the face activation tensor in TT format, by summation cores. This is for example the case for the examples constructed based on Hamming distances.

1.7.3 Relation with Facet Complexity

Facet complexity [4, Definition 6.2.2] bounds the encoding length of the inequalities describing a polytope.

For any face $\mathcal{F}$ we find a normal $a \in \mathbb{R}^p$ and $b \in \mathbb{R}$ such that x

$$\mathcal{F} = \{ \mu [L] \in \mathcal{M} : \langle a [L], \mu [L] \rangle_{[\emptyset]} = b \} .$$

We can thus represent a face activating tensor by computing the contraction $\langle a [L], y [L] \rangle_{[\emptyset]}$.

To this end, let $\mathcal{U} \subset \mathbb{R}$ be a set with index interpretation function

$$I : [r] \rightarrow \mathcal{U} .$$

We demand $0, b \in \mathcal{U}$. We can for example choose $\mathcal{U}$ as the d -bit binary integer representation.

We then define for $\ell \in [p]$ the summation operations

$$+^{\ell, a[L=\ell]} : \mathcal{U} \times [n] \rightarrow \mathcal{U}$$

which satisfies for any $u \in \mathcal{U}$ and y_ℓ for which $u + a[L = \ell] \cdot y_\ell \in \mathcal{U}$ the equation

$$+^{\ell, a[L=\ell]}(u, y_\ell) = u + a[L = \ell] \cdot y_\ell.$$

We further assume, that for any tuple $y_{[p]} \in \times_{\ell \in [p]} [n_\ell]$ and any $\tilde{\ell} \in [p]$

$$\sum_{\ell \leq \tilde{\ell}} a[L = \ell] \cdot y_{\tilde{\ell}} \in \mathcal{U}.$$

Then we have

$$\sum_{v[L] \in \mathcal{N}_{t,v}^{\mathcal{F}}} \epsilon_{v[L]} [Y_{[p]}] = \left\langle \{\beta^{+^{\ell, a[L=\ell]}} [Y_{+^{\ell, a[L=\ell]}}, Y_\ell] : \ell \in [p]\} \cup \{\epsilon_{I^{-1}(0)} [Y_{+^{-1, a[L=-1]}}, \epsilon_{I^{-1}(b)} [Y_{+^{p-1, a[L=p-1]}}, \dots]\} \right\rangle_{[Y_{[p]}]}.$$

The variable $Y_{+^{-1, a[L=-1]}}$ is auxiliary to initialize the sum with 0.

We can weaken the assumptions on $\mathcal{U}$: When $a[L]$ is positive, then any partial sum which exceeds b can be represented by any member of $\mathcal{U}$ which is larger than b . This brings further sparsity, since no values larger than b are needed to represent the face (add for example the ∞ element to $\mathcal{U}$ and represent any partial sum exceeding b by it).

Definition 1.8 We say that a face $\mathcal{F}$ has *TT face complexity* r if there exists a normal $a \in \mathbb{R}^p$ and $b \in \mathbb{R}$ with

$$\mathcal{F} = \{\mu[L] \in \mathcal{M} : \langle a[L], \mu[L] \rangle_{[\emptyset]} = b\}.$$

and there is a subset $\mathcal{U}$ of cardinality at most r , for which $0, b \in \mathcal{U}$ and for any tuple $y_{[p]} \in \times_{\ell \in [p]} [n_\ell]$ and any $\tilde{\ell} \in [p]$

$$\sum_{\ell \leq \tilde{\ell}} a[L = \ell] \cdot y_{\tilde{\ell}} \in \mathcal{U}.$$

Comparison with facet complexity [4, Definition 6.2.2]:

- We define complexity here without the log (which would get the string length), to make a direct connection with TT ranks.
- Facet complexity considers only the representation of the equation, not the demand of computing it.
- Facet complexity considers just facets, not general faces, since it makes a statement about encoding the halfspace representation of the whole polytope.

Theorem 1.10 *Let $\mathcal{F}$ be a face with TT face complexity r . Then the face activation tensor has a TT rank of at most r .*

Proof. See above construction. $\square$

References

1. Shun-ichi Amari. *Information Geometry and Its Applications*, volume 194 of *Applied Mathematical Sciences*. Springer Japan, Tokyo, 2016.
2. Lawrence D. Brown. *Fundamentals of Statistical Exponential Families*. Institute of mathematical Statistics, Hayward, Calif, June 1987.
3. George Casella and Roger Berger. *Statistical Inference*. Cengage Learning, Pacific Grove, Calif, June 2001.
4. Robert G. Gallager. *Low-Density Parity-Check Codes*. The MIT Press, September 1963.
5. Alex Goessmann, Janina Schütte, Maximilian Fröhlich, and Martin Eigel. A tensor network formalism for neuro-symbolic AI, January 2026. arXiv:2601.15442 [cs].
6. Martin Grötschel, László Lovász, and Alexander Schrijver. *Geometric Algorithms and Combinatorial Optimization*, volume 2 of *Algorithms and Combinatorics*. Springer, Berlin, Heidelberg, 1993.
7. Sebastian Holtz, Thorsten Rohwedder, and Reinhold Schneider. On manifolds of tensors of fixed TT-rank. *Numerische Mathematik*, 120(4):701–731, April 2012.
8. Kevin P. Murphy. *Probabilistic Machine Learning: Advanced Topics*. The MIT Press, Cambridge, August 2023.

9. Stuart Russell and Peter Norvig. *Artificial Intelligence: A Modern Approach, Global Edition: A Modern Approach, Global Edition*. Pearson, Boston, 4 edition, May 2021.
10. Aaron Sander, Maximilian Fröhlich, Martin Eigel, Jens Eisert, Patrick Gelß, Michael Hintermüller, Richard M. Milbradt, Robert Wille, and Christian B. Mendl. Large-scale stochastic simulation of open quantum systems. *Nature Communications*, 16(1):11074, December 2025. arXiv:2501.17913 [quant-ph].
11. Martin J. Wainwright and Michael Irwin Jordan. *Graphical Models, Exponential Families, and Variational Inference*. Now Publishers Inc, 2008.
12. Günter M. Ziegler. Lectures on 0/1-Polytopes. In Gil Kalai and Günter M. Ziegler, editors, *Polytopes — Combinatorics and Computation*, pages 1–41. Birkhäuser, Basel, 2000.
13. Günter M. Ziegler. *Lectures on Polytopes*. Springer, New York, 1995th edition edition, October 2013.