



(12) **United States Plant Patent**
Tonies

(10) **Patent No.:** **US PP21,369 P2**
(45) **Date of Patent:** **Oct. 5, 2010**

(54) **PHLOX PLANT NAMED ‘DITOMFAV’**

(50) Latin Name: ***Phlox paniculata***
Varietal Denomination: **Ditomfav**

(75) Inventor: **Hubertus Josephus Tonies**,
Noordwijkerhout (NL)

(73) Assignee: **Kolster Beheer B.V.**, Boskoop (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 53 days.

(21) Appl. No.: **12/315,158**

(22) Filed: **Nov. 28, 2008**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./320**

(58) **Field of Classification Search** **Plt./320**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP19,600 P2 * 12/2008 Verschoor Plt./320

OTHER PUBLICATIONS

UPOV ROM GTITM Computer Database GTI Jouve Retrieval Soft-
ware 2009/04 Citation for ‘Ditomfav’.*

* cited by examiner

Primary Examiner—Wendy C. Haas

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Phlox* plant named ‘Ditomfav’,
characterized by its compact and upright plant habit; freely
branching and flowering habit; dark-colored stems and
leaves; light pink-colored flowers; and tolerance to Downy
Mildew.

3 Drawing Sheets

1

Botanical designation: *Phlox paniculata*.

Cultivar denomination: ‘DITOMFAV’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Phlox* plant, botanically known as *Phlox paniculata* and
hereinafter referred to by the name ‘Ditomfav’.

The new *Phlox* plant is a product of a planned breeding
program conducted by the Inventor in Noordwijkerhout, The
Netherlands. The objective of the breeding program is to
create new compact *Phlox* cultivars with attractive leaf and
flower coloration and disease resistance.

The new *Phlox* plant originated from a cross-pollination in
Noordwijkerhout, The Netherlands in July, 2000 of two
unnamed seedling selections of *Phlox paniculata*, not pat-
ented. The new *Phlox* was discovered and selected by the
Inventor as a single flowering plant from within the progeny
of the stated cross-pollination in a controlled environment in
Noordwijkerhout, The Netherlands in July, 2001.

Asexual reproduction of the new *Phlox* plant by cuttings in
a controlled environment in Noordwijkerhout, The Nether-
lands since 2001, has shown that the unique features of this
new *Phlox* plant are stable and reproduced true to type in
successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Phlox* have not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment and cultural prac-
tices such as temperature and light intensity without, how-
ever, any variance in genotype.

The following traits have been repeatedly observed and are
determined to be the unique characteristics of ‘Ditomfav’.
These characteristics in combination distinguish ‘Ditomfav’
as a new and distinct cultivar of *Phlox*:

2

1. Compact and upright plant habit.
2. Freely branching and flowering habit.
3. Dark-colored stems and leaves.
4. Light pink-colored flowers.
5. Tolerant to Downy Mildew.

Plants of the new *Phlox* differ primarily from plants of the
female parent selection in the following characteristics:

1. Plants of the new *Phlox* are more compact than plants of
the female parent selection.
2. Plants of the new *Phlox* have larger flowers than plants of
the female parent selection.
3. Plants of the new *Phlox* and the female parent selection
differ in flower color as plants of the female parent
selection have darker pink-colored flowers.

Plants of the new *Phlox* differ primarily from plants of the
male parent selection in the following characteristics:

1. Plants of the new *Phlox* have stronger stems than plants
of the male parent selection.
2. Plants of the new *Phlox* have darker-colored leaves than
plants of the male parent selection.
3. Plants of the new *Phlox* and the male parent selection
differ in flower color as plants of the male parent selec-
tion have pale lilac-colored flowers.

Plants of the new *Phlox* can also be compared to plants of
Phlox paniculata ‘Rosa Pastel’, not patented. In side-by-side
comparisons conducted in Noordwijkerhout, The Nether-
lands, plants of the new *Phlox* and ‘Rosa Pastel’ differed
primarily in the following characteristics:

1. Plants of the new *Phlox* were more compact than plants
of ‘Rosa Pastel’.
2. Plants of the new *Phlox* had darker-colored leaves than
plants of ‘Rosa Pastel’.
3. Flowers of plants of the new *Phlox* were darker in color
than flowers of plants of ‘Rosa Pastel’.
4. Plants of the new *Phlox* were tolerant to Downy Mildew
whereas plants of ‘Rosa Pastel’ were less tolerant to
Downy Mildew.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Phlox*, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Phlox*.

The photograph on the first sheet comprises a side perspective view of a typical plant of 'Ditomfav' grown in a container.

The photograph on the second sheet is a close-up view of typical leaves of 'Ditomfav'.

The photograph on the third sheet is a close-up view of a typical inflorescence of 'Ditomfav'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in Boskoop, The Netherlands, under commercial practice during the summer in containers in an outdoor nursery. During the production of the plants, day temperatures ranged from 12° C. to 30° C. and night temperatures ranged from 4° C. to 16° C. Plants had been growing for one year when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Botanical classification: *Phlox paniculata* 'Ditomfav'.

Parentage:

Female, or seed, parent.—Unnamed selection of *Phlox paniculata*, not patented.

Male, or pollen, parent.—Unnamed selection of *Phlox paniculata*, not patented.

Propagation:

Type.—By cuttings.

Time to initiate roots.—About two weeks at 15° C. to 25° C.

Time to produce a rooted young plant.—About 35 days at 15° C. to 25° C.

Root description.—Fibrous, thin; pale cream tan in color.

Rooting habit.—Moderate branching; dense.

Plant description:

Plant form/habit.—Compact and upright plant habit; moderately vigorous growth habit.

Branching habit.—Freely branching habit with about six lateral branches per plant; pinching enhances branching.

Plant height.—About 29.6 cm.

Plant width (spread).—About 31.9 cm.

Lateral branches.—Length: About 19.9 cm. Diameter: About 4 mm. Internode length: About 2.6 cm. Strength: Strong. Texture: Smooth, glabrous. Color: Close to N186C, near the nodes, close to 146C to 146D.

Foliage description:

Arrangement.—Opposite, simple.

Length.—About 10.5 cm.

Width.—About 3.7 cm.

Shape.—Narrowly ovate to elliptic.

Apex.—Acute to elongated acuminate.

Base.—Obtuse to attenuate.

Margin.—Very finely serrate.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close to 143B flushed with close to N186C. Developing leaves, lower surface: Close to 143B to 143C slightly flushed with close to N186C. Fully expanded leaves, upper surface: Between 137A and 139A; venation, close to 144B. Fully expanded leaves, lower surface: Close to 137B; venation, close to 144C to 144D.

Petioles.—Length: About 3 mm. Diameter: About 3 mm.

Texture, upper and lower surfaces: Smooth, glabrous.

Color, upper and lower surfaces: Close to 146B.

Flower description:

Flower type/habit.—Single rotate flowers arranged in terminal compound panicles; flowers face mostly upright or outwardly. Panicles rounded and hemispherical in shape. Freely flowering habit with about 80 flowers developing per inflorescence.

Fragrance.—Moderately fragrant; sweet and pleasant.

Natural flowering season.—Continuously flowering from July to September in The Netherlands.

Postproduction longevity.—Flowers last about ten days on the plant; flowers not persistent.

Flower buds.—Height: About 1.9 cm. Diameter: About 5 mm. Shape: Narrowly oblanceolate. Color: Close to 75C; lower half, close to N77B; base, close to 145C to 145D.

Inflorescence height.—About 10.2 cm.

Inflorescence diameter.—About 6.1 cm.

Flower diameter.—About 3 cm.

Flower depth.—About 2.7 cm.

Petals.—Quantity per flower: Typically five in a single whorl; petals fused at the base into a narrow tube. Length from throat: About 1.5 cm. Length fused: About 2.2 cm. Lobe width: About 1.5 cm. Lobe shape: Roughly spatulate. Apex: Rounded to retuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: Developing petals, upper surface: Close to 75A; towards the throat, spots, N155A; throat, close to 70A. Developing petals, lower surface: Close to 75A to 75B; tube, close to 70A. Fully expanded petals, upper surface: Close to 77C to 77D; towards the throat, close to N74B; throat, close to 70A. Fully expanded petals, lower surface: Close to 76B to 76C; tube, close to 70A.

Sepals.—Quantity per flower: Typically five in a single whorl, fused towards the base. Length: About 9 mm. Width: About 1.5 mm. Shape: Lanceolate. Apex: Narrowly apiculate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: Developing and fully expanded sepals, upper surface: Close to N186C; towards the base, close to 145C to 145D. Developing and fully expanded sepals, lower surface: Close to N186C; towards the base, close to 145C to 145D.

Peduncles.—Length: About 6.1 cm. Diameter: About 2.5 mm. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 145C to 145D tinged with close to N186C.

Pedicels.—Length: About 5 mm. Diameter: About 1 mm. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 177B.

Reproductive organs.—Stamens: Quantity per flower: Typically five. Filament length: About 1 mm. Anther shape: Oblong. Anther length: About 1.5 mm. Anther color: Close to 8D. Pollen amount: Scarce to moder-

ate. Pollen color: Close to 11D. Pistils: Quantity per flower: One. Pistil length: About 1.9 cm. Stigma shape: Three-parted. Stigma color: Close to 150D. Style length: About 1.8 cm. Style color: Close to 187D. Ovary color: Close to 143A. 5

Seed/fruit.—Seed and fruit development have not been observed.

Disease/pest resistance: Plants of the *Phlox* have been observed to be tolerant to Downy Mildew. Plants of the new *Phlox* have not been observed to be resistant to pests or 10 other pathogens common to *Phlox*.

Garden performance: Plants of the new *Phlox* have been observed to have good garden performance and tolerate rain, wind and temperatures ranging from about –20° C. to about 40° C.

It is claimed:

1. A new and distinct *Phlox* plant named ‘Ditomfav’ as illustrated and described.

* * * * *





