

US00PP22218P2

(12) **United States Plant Patent**  
**Ferguson**

(10) **Patent No.:** **US PP22,218 P2**  
(45) **Date of Patent:** **Nov. 1, 2011**

(54) **STRAWBERRY PLANT NAMED**  
**‘DRISSTRAWSEVENTEEN’**

(50) Latin Name: *Fragaria×ananassa*  
Varietal Denomination: **DrisStrawSeventeen**

(75) Inventor: **Michael D. Ferguson**, Moorpark, CA  
(US)

(73) Assignee: **Driscoll Strawberry Associates, Inc.**,  
Watsonville, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/803,582**

(22) Filed: **Jun. 30, 2010**

(51) **Int. Cl.**  
*A01H 5/00* (2006.01)

(52) **U.S. Cl.** ..... **Plt./209**

(58) **Field of Classification Search** ..... Plt./209  
See application file for complete search history.

*Primary Examiner* — Annette Para

(74) *Attorney, Agent, or Firm* — Jondle & Associates, P.C.

(57) **ABSTRACT**

This invention relates to a new and distinct variety of straw-  
berry plant named ‘DrisStrawSeventeen’. The new variety is  
primarily characterized by an upright and globose plant habit,  
very strong vigor, and very large-sized berries, is disclosed.

**3 Drawing Sheets**

**1**

Genus and species: *Fragaria×ananassa*.

Variety denomination: ‘DrisStrawSeventeen’.

**BACKGROUND OF THE NEW PLANT**

The present invention relates to a new and distinct straw-  
berry variety designated ‘DrisStrawSeventeen’ and botani-  
cally known as *Fragaria×ananassa*. This new strawberry  
variety was discovered in Ventura County, Calif. in October  
2006 and originated from a cross between the proprietary  
female parent ‘13H377’ (unpatented) and the proprietary  
male parent ‘119J176’ (unpatented). The original seedling of  
the new variety was first asexually propagated at a nursery in  
Shasta County, Calif. in October 2005.

‘DrisStrawSeventeen’ was subsequently asexually propa-  
gated at a nursery in Shasta County, Calif. and underwent  
further testing in Ventura County, Calif. for three years (2006-  
2009). The present invention has been found to retain its  
distinctive characteristics through successive asexual propa-  
gations via stolons and tissue culture.

Plant Breeder’s Rights for this variety have not been  
applied for. ‘DrisStrawSeventeen’ has not been made publicly  
available or sold more than one year prior to the filing date of  
this application.

**DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying color photographs show typical speci-  
mens of the new variety at various stages of development. The  
colors shown are as true as can be reasonably obtained by  
conventional photographic procedures. The photographs  
were taken from three-month-old plants.

FIG. 1 shows overall plant habit including fruit at various  
stages of development.

FIG. 2 shows leaves of the plant with three leaflets.

FIG. 3 shows both the upper surface and lower surface of  
several of the flowers.

FIG. 4 shows the whole fruit.

FIG. 5 shows the fruit in longitudinal cross-section.

**DESCRIPTION OF THE NEW VARIETY**

The following detailed descriptions set forth the distinctive  
characteristics of ‘DrisStrawSeventeen’. The data which

**2**

define these characteristics is based on observations taken in  
Ventura County, Calif. from 2006 to 2009. This description is  
in accordance with UPOV terminology. Color designations,  
color descriptions, and other phenotypical descriptions may  
deviate from the stated values and descriptions depending  
upon variation in environmental, seasonal, climatic, and cul-  
tural conditions. ‘DrisStrawSeventeen’ has not been observed  
under all possible environmental conditions. The botanical  
description of ‘DrisStrawSeventeen’ was taken from four-  
month-old plants. Color terminology follows The Royal Hor-  
ticultural Society Colour Chart, London (R.H.S.) (2001).

**DETAILED BOTANICAL DESCRIPTION OF THE  
PLANT**

**Classification:**

*Species.*—*Fragaria×ananassa*.

*Common name.*—Strawberry.

*Denomination.*—‘DrisStrawSeventeen’.

**Parentage:**

*Female parent.*—The proprietary female parent  
‘13H377’ (unpatented).

*Male parent.*—The proprietary male parent ‘119J176’  
(unpatented).

**Plant:**

*Height.*—30.8 cm.

*Diameter.*—50.7 cm.

*Number of crowns/plant.*—3.

*Habit.*—Upright and globose.

*Density of individual plant.*—Dense.

*Vigor.*—Very strong.

**Leaves:**

*Terminal leaflet length.*—9.8 cm.

*Terminal leaflet width.*—9.0 cm.

*Terminal leaflet length/width ratio.*—1.1.

*Number of teeth/terminal leaflet.*—24.

*Shape of teeth.*—Obtuse.

*Color.*—Upper surface: RHS 137A (Dark green). Lower  
surface: RHS 148B (Medium yellow-green).

*Leaf shape in cross section.*—Concave.

*Leaf blistering.*—Medium.

*Leaf glossiness.*—Medium.

*Number of leaflets.*—3 only.

*Terminal leaflet margin profile.*—Revolute.



*Terminal leaflet length/width ratio*.—As long as broad.  
*Terminal leaflet shape*.—Orbicular.  
*Terminal leaflet base shape*.—Rounded.  
*Terminal leaflet apex shape*.—Rounded.  
*Petiole*.—Length: 18.9 cm. Diameter: 0.316 cm. Pubes- 5  
 cence: Medium. Pose of hairs: Upwards. Color: RHS  
 145A (Medium yellow-green).  
*Petiolule*:  
*Color*.—RHS 145A (Medium yellow-green).  
*Length*.—1.361 cm.  
*Diameter*.—0.178 cm. 10  
*Stipule*:  
*Length*.—2.8 cm.  
*Width*.—0.793 cm.  
*Pubescence*.—Medium.  
*Stipule anthocyanin coloration*.—Weak; RHS 145C 15  
 (Light yellow-green).  
*Stolon*:  
*Number*.—Few.  
*Average number of daughter plants per plant*.—34.  
*Stolon anthocyanin*.—RHS 63B (Medium red-purple). 20  
*Thickness*.—Medium.  
*Pubescence*.—Medium.  
*Inflorescence*:  
*Position relative to foliage*.—Beneath.  
*Time of flowering (50% of plants at first flower)*.—Me- 25  
 dium.  
*Flower size*.—Medium.  
*Diameter*.—2.966 cm.  
*Petals*.—Shape: Orbicular. Apex: Rounded. Base: Con-  
 cave-convex. Margin: Entire. Spacing: Overlapping.  
 Length: 1.286 cm. Width: 1.330 cm. Length/width 30  
 ratio: 1.0; as long as broad. Typical and observed petal  
 number per flower: 6. Color (both surfaces): RHS  
 155C (White).  
*Calyx*.—Diameter: 3.234 cm. Diameter relative to  
 corolla: Same size. Inner calyx diameter relative to 35  
 outer: Same size.  
*Sepal*.—Shape: Elliptical. Apex: Convex. Margin:  
 Entire. Length: 1.244 cm. Width: 0.579 cm. Typical  
 and observed sepal number per flower: 13.  
*Receptacle color*.—RHS 1C (Light green-yellow). 40  
*Anther color*.—RHS 22A (Medium yellow-orange).  
*Fruiting Truss*:  
*Length*.—19.5 cm; long.  
*Diameter at base of truss*.—0.383 cm.  
*Number of berries per fruiting truss*.—5.  
*Attitude at first picking*.—Prostrate.  
*Color at base of truss*.—RHS 144B (Medium yellow-  
 green).  
*Fruit*:  
*Length*.—4.902 cm.  
*Width*.—4.690 cm.  
*Length/width ratio*.—1.0.  
*Fruit hollow length*.—1.876 cm.  
*Fruit hollow width*.—1.625 cm.  
*Fruit hollow length/width ration*.—1.2.  
*Fruit hollow center (size)*.—Medium.  
*Weight (per individual berry)*.—28.4 g.  
*Fruit ratio of length/maximum width*.—As broad as  
 long.  
*Relative fruit size*.—Very large.  
*Predominant fruit shape*.—Wedged.  
*Difference in shape between primary and secondary*  
*fruits*.—None or very slight.  
*Unevenness of fruit surface*.—Strong.  
*Fruit skin color*.—RHS 46B (Dark red).  
*Evenness of fruit color*.—Uneven.

*Fruit glossiness*.—Medium.  
*Insertion of achenes*.—Level with surface.  
*Achene coloration (sunward side of berry)*.—RHS  
 166D (Medium greyed-orange).  
*Achene coloration (shaded side of berry)*.—RHS 163C  
 (Light greyed-orange).  
*Achenes per berry*.—390.  
*Band without achenes*.—Narrow.  
*Insertion of calyx*.—Level.  
*Pose of calyx segments*.—Reflexed.  
*Size of calyx in relation to fruit*.—Between smaller and  
 same size.  
*Adherence of calyx*.—Strong.  
*Firmness of flesh*.—Firm.  
*Color of flesh*.—RHS N155D (White) and RHS N30A  
 (Medium orange-red).  
*Evenness of flesh color*.—Slightly uneven.  
*Distribution of flesh color*.—Marginal and central.  
*Sweetness*.—Medium.  
*Acidity*.—Medium.  
*Texture when tasted*.—Coarse.  
*Type of bearing*.—Fully everbearing.  
*Grams of fruit/plant*.—486 g.  
*Harvest interval*.—Early June to late December.  
*Harvest maturity*.—Mid-season.  
 Disease, pest, and stress resistance:  
*Botrytis fruit rot*.—Moderately susceptible.  
*Powdery mildew*.—Moderately susceptible.  
*Tetranychus urticae*.—Moderately resistant.  
*Tarsonemus pallidus*.—Moderately resistant.  
*Aphelencoides fragariae*.—Moderately resistant.  
*Pratylenchus penetrans*.—Moderately resistant.  
*Ditylenchus dipsac*.—Moderately resistant.  
*Anthonomus rubi*.—Moderately resistant.  
*Aphis spp. (Aphids)*.—Moderately resistant.  
*Lygus hesperus (Lygus bug)*.—Moderately resistant.  
*Wind*.—Moderately resistant.

#### COMPARISON WITH PARENTAL AND COMMERCIAL VARIETIES

When 'DrisStrawSeventeen' is compared to the propri-  
 etary female parent '13H377' (unpatented), 'DrisStrawSev-  
 enteen' has a more vigorous growth habit and a larger plant  
 size than '13H377'. In addition, 'DrisStrawSeventeen' has  
 lighter colored foliage, a larger fruit size, and a higher fruit  
 yield than '13H377'. 45

When 'DrisStrawSeventeen' is compared to the propri-  
 etary male parent '119J176' (unpatented), 'DrisStrawSeven-  
 teen' has a stronger everbearing habit and is more susceptible  
 to powdery mildew. In addition, 'DrisStrawSeventeen' has  
 lighter colored foliage, a larger fruit size, and a higher fruit  
 yield than '119J176'. 50

When 'DrisStrawSeventeen' is compared to the commer-  
 cial variety 'DrisStrawThree' (U.S. Plant Pat. No. 19,673),  
 the petiole pose of hairs for 'DrisStrawSeventeen' is upwards,  
 while the petiole pose of hairs for 'DrisStrawThree' is out-  
 wards to downwards. In addition, 'DrisStrawSeventeen' has  
 fruit with medium sweetness, coarse texture when tasted, and  
 an uneven fruit color, while 'DrisStrawThree' has fruit with a  
 strong sweetness, very fine texture when tasted, and a slightly  
 uneven fruit color. 55

I claim:

1. A new and distinct variety of strawberry plant as  
 described and shown herein.





FIG. 1




FIG. 2






FIG. 3




FIG. 4






FIG. 5