



(12) **United States Plant Patent**
Wood

(10) **Patent No.:** **US PP25,897 P2**
(45) **Date of Patent:** **Sep. 15, 2015**

(54) **VIBURNUM PLANT NAMED ‘SMVDBL’**

(50) Latin Name: *Viburnum bracteatum*
Varietal Denomination: **SMVDBL**

(71) Applicant: **Timothy D. Wood**, Spring Lake, MI
(US)

(72) Inventor: **Timothy D. Wood**, Spring Lake, MI
(US)

(73) Assignee: **Spring Meadow Nursery Inc.**, Grand
Haven, MI (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 132 days.

(21) Appl. No.: **13/987,980**

(22) Filed: **Sep. 20, 2013**

(51) **Int. Cl.**
A01H 5/08 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./226**

(58) **Field of Classification Search**

USPC Plt./226
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

UPOV hit CA PBR 12-7742 on *Viburnum bracteatum* ‘SMVDBL’,
published Oct. 31, 2012.*

* cited by examiner

Primary Examiner — Anne Grunberg

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Viburnum* plant named
‘SMVDBL’, characterized by its compact, upright and some-
what outwardly spreading plant habit; vigorous growth habit;
freely branching habit; glossy leaves; inflorescences with
numerous white-colored flowers; dark violet blue-colored
fruits; and good container and garden performance.

4 Drawing Sheets

1

Botanical designation: *Viburnum bracteatum*.
Cultivar denomination: ‘SMVDBL’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Viburnum* plant, botanically known as *Viburnum bractea-*
tum and hereinafter referred to by the name ‘SMVDBL’.

The new *Viburnum* plant is a product of a planned breeding
program conducted by the Inventor in Grand Haven, Mich.
The objective of the breeding program is to develop new
compact *Viburnum* plants with good container performance
and unique leaf types.

The new *Viburnum* plant originated from an open-pollina-
tion during the summer of 2007 of *Viburnum bracteatum*
‘Emerald Luster’, not patented, as the female, or seed, parent
with an unknown selection of *Viburnum bracteatum* as the
male, or pollen, parent. The new *Viburnum* plant was discov-
ered and selected by the Inventor in 2009 as a single flowering
plant from within the progeny of the stated open-pollination
in a controlled environment in Grand Haven, Mich.

Asexual reproduction of the new *Viburnum* plant by soft-
wood cuttings in a controlled environment in Grand Haven,
Mich. since June, 2009 has shown that the unique features of
this new *Viburnum* plant are stable and reproduced true to
type in successive generations of asexual reproduction.

SUMMARY OF THE INVENTION

Plants of the new *Viburnum* have not been observed under
all possible environmental conditions and cultural practices.
The phenotype may vary somewhat with variations in envi-
ronmental conditions such as temperature and light intensity
without, however, any variance in genotype.

2

The following traits have been repeatedly observed and are
determined to be the unique characteristics of ‘SMVDBL’.
These characteristics in combination distinguish ‘SMVDBL’
as a new and distinct *Viburnum* plant:

1. Compact, upright and somewhat outwardly spreading
plant habit.
2. Vigorous growth habit.
3. Freely branching habit.
4. Glossy leaves.
5. Inflorescences with numerous white-colored flowers.
6. Dark violet blue-colored fruits.
7. Good container and garden performance.

Plants of the new *Viburnum* can be compared to plants of
the female parent, ‘Emerald Luster’. Plants of the new *Vibur-*
num differ from plants of ‘Emerald Luster’ in the following
characteristics:

1. Plants of the new *Viburnum* are more compact than
plants of ‘Emerald Luster’.
2. Leaves of the new *Viburnum* are glossier than leaves of
plants of ‘Emerald Luster’.
3. Inflorescences of plants of the new *Viburnum* are more
compressed than inflorescences of ‘Emerald Luster’.

Plants of the new *Viburnum* can be compared to plants of
the *Viburnum bracteatum* ‘SMVDLS’, disclosed in a U.S.
Plant patent application filed concurrently having application
Ser. No. 13/987,981. Plants of the new *Viburnum* differ from
plants of ‘SMVDLS’ in the following characteristics:

1. Plants of the new *Viburnum* are more freely branching
than plants of ‘SMVDLS’.
2. Plants of the new *Viburnum* have smaller inflorescences
with fewer flowers per inflorescence than plants of
‘SMVDLS’.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the over-
all appearance of the new *Viburnum* plant showing the colors

as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Viburnum* plant.

The photograph on the first sheet is a side perspective view of a typical plant of 'SMVDBL' grown in a container.

The photograph on the second sheet is a close-up view of a typical lateral branch and leaves of 'SMVDBL'.

The photograph on the third sheet is a close-up view of typical inflorescences of 'SMVDBL'.

The photograph on the fourth sheet is a close-up view of typical fruits of 'SMVDBL'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants of the new *Viburnum* grown in three-gallon containers during the summer in an outdoor nursery in Grand Haven, Mich. and under cultural practices typical of commercial *Viburnum* production. Plants were three years old when the photographs and the description were taken. In the description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Viburnum bracteatum* 'SMVDBL'.

Parentage:

Female, or seed, parent.—*Viburnum bracteatum* 'Emerald Luster', not patented.

Male, or pollen, parent.—Unknown selection of *Viburnum bracteatum*, not patented.

Propagation:

Type.—By softwood cuttings.

Time to initiate roots, summer.—About 30 days at 24° C.

Time to produce a rooted young plant, summer.—About four months at 24° C.

Root description.—Fine to thick; white and brown in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Deciduous perennial shrub; compact, upright and somewhat outwardly spreading plant habit; vigorous growth habit.

Branching habit.—Freely branching habit with about 38 lateral branches developing per plant; pinching (removal of terminal apices) will enhance lateral branch development.

Plant height.—About 62 cm.

Plant diameter(area of spread).—About 60 cm.

Lateral branch description:

Length.—About 42 cm.

Diameter.—About 3 mm.

Internode length.—About 5 cm.

Texture.—Pubescent.

Strength.—Strong.

Aspect.—Erect to about 20° from vertical.

Color.—Close to 199B.

Leaf description:

Arrangement.—Opposite, simple.

Length.—About 8 cm.

Width.—About 6 cm.

Shape.—Ovate.

Apex.—Acute.

Base.—Obtuse.

Margin.—Crenate.

Texture, upper and lower surfaces.—Pubescent; coarse.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close to 146A. Developing leaves, lower surface: Close to 146B. Fully expanded leaves, upper surface: Close to 137A; venation, close to 146C; glossy. Fully expanded leaves, lower surface: Close to 137B; venation, close to 145C.

Petiole.—Length: About 1.2 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: Close to 146C.

Flower description:

Flower appearance and arrangement.—Single rotate flowers arranged in terminal cymes; freely flowering habit with usually about 228 flowers per inflorescence; flowers face upright to outwardly.

Natural flowering season.—Flowering commences in June in Mich.; flowers last about two to four weeks on the plant; flowers not persistent.

Fragrance.—Slightly fragrant; sweet, pleasant.

Inflorescence height.—About 3.8 cm.

Inflorescence diameter.—About 4 cm.

Flower diameter.—About 3.5 mm.

Flower length(height).—About 2 mm.

Flower buds.—Length: About 1 mm. Diameter: About 1 mm. Shape: Globose. Color: Close to 142B to 142C.

Petals.—Quantity per flower: Five petals in a single whorl. Length: About 1.8 mm. Width: About 1.5 mm. Shape: Obovate. Apex: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper and lower surfaces: Close to 155D. Fully opened, upper and lower surfaces: Close to 155D.

Sepals.—Sepal development has not been observed on plants of the new *Viburnum*.

Peduncles.—Length: About 2 cm. Diameter: About 4 mm. Strength: Strong. Aspect: Mostly erect. Texture: Smooth. Color: Close to 141C.

Pedicels.—Length: About 7.5 mm. Diameter: About 3 mm. Strength: Strong. Aspect: Erect to about 80° from peduncle axis. Texture: Smooth. Color: Close to 141C.

Reproductive organs.—Androecium: Quantity per flower: About five. Anther shape: Irregularly globular. Anther length: About 1 mm. Anther color: Close to 155A. Amount of pollen: Slight. Pollen color: Close to 155D. Gynoecium: Quantity per flower: One. Pistil length: About 2 mm. Style length: About 1.8 mm. Style color: Close to 143D. Stigma shape: Rounded. Stigma color: Close to 143D. Ovary color: Close to 143D. Fruits: Length: About 5 mm. Diameter: About 5 mm. Shape: Spherical. Texture: Smooth, glossy luster. Color, immature: Close to 143C. Color, mature: Close to 103B to 103C. Seeds: Length: About 4.8 mm. Diameter: About 3 mm. Texture: Smooth. Color: Close to 177C.

Garden performance: Plants of the new *Viburnum* have been observed to have good garden and container performance and to tolerate rain, wind and temperatures ranging from about -27° C. to about 40° C.

Pathogen & pest resistance: Plants of the new *Viburnum* have not been observed to be resistant to pathogens and pests common to *Viburnum* plants.

It is claimed:

1. A new and distinct *Viburnum* plant named 'SMVDBL' as illustrated and described.

* * * * *







