



(12) **United States Plant Patent**
Larse

(10) **Patent No.:** **US PP28,341 P3**
(45) **Date of Patent:** **Aug. 29, 2017**

(54) **STRAWBERRY PLANT NAMED ‘YOLI’**

(50) Latin Name: *Fragaria*×*ananassa*
Varietal Denomination: **Yoli**

(71) Applicant: **John Larse**, Aptos, CA (US)

(72) Inventor: **John Larse**, Aptos, CA (US)

(73) Assignee: **Sweet Darling Sales, Inc.**, Aptos, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/545,384**

(22) Filed: **Apr. 29, 2015**

(65) **Prior Publication Data**

US 2016/0324049 P1 Nov. 3, 2016

(51) **Int. Cl.**
A01H 5/08 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./209**

(58) **Field of Classification Search**
USPC Plt./209
See application file for complete search history.

Primary Examiner — Anne Grunberg

(74) Attorney, Agent, or Firm — Cooley LLP

(57) **ABSTRACT**

The present invention provides a new and distinct strawberry plant designated as ‘Yoli’ (a.k.a. ‘104522’).

6 Drawing Sheets

1

Latin name of the genus and species: *Fragaria*×*ananassa*.
Varietal denominations: ‘Yoli’ (a.k.a. ‘104522’).

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct strawberry variety designated as ‘Yoli’ (a.k.a. ‘104522’).

‘Yoli’ (a.k.a. ‘104522’) is the result of a controlled-cross between a female parent cultivar designated 1621 and a male parent cultivar designated 1285 (both unpatented, proprietary cultivars) made by the Inventor and was first fruited in Watsonville, Calif. growing fields. Following selection and during testing, the plant was originally designated ‘104522’ and subsequently named ‘Yoli’.

This new variety was asexually reproduced via runners (stolons) by the inventor at Watsonville, Calif. Asexual propagules from the original source have been tested in Watsonville growing fields and to a limited extent, grower fields in high elevation. The properties of this variety were found to be transmissible by such asexual reproduction. This cultivar is stable and reproduces true to type in successive generations of asexual reproduction.

BRIEF SUMMARY OF THE INVENTION

This invention relates to new and distinctive strawberry cultivar designated as ‘Yoli’. These cultivars are primarily adapted to the climate and growing conditions of the central coast of California. This region provides the necessary temperatures required for it to produce a strong vigorous plant and to remain in fruit production from March through October. The nearby Pacific Ocean provides the needed humidity and moderate day temperatures and evening chilling to maintain fruit quality for the production months.

The following traits and photographs in combination distinguish strawberry variety ‘Yoli’ from known strawberry varieties. In addition, this new cultivar was confirmed to be a unique strawberry germplasm when tested against the California Seed & Plant Lab, Inc. (Elverta, Calif.) database using Short Sequence Repeats (SSRs). Plants for the botani-

2

cal measurements in the present application were grown as annuals. Any color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The fruit produced by this new cultivar is attractive and of excellent quality.

DESCRIPTION OF THE DRAWINGS

The accompanying color photographs depict various characteristics of the cultivars as nearly true as possible to make color reproductions.

FIG. 1 shows ‘Yoli’ plants growing in the field.

FIG. 2 shows inflorescences and fruits of ‘Yoli’ plants.

FIG. 3 shows ‘Yoli’ plants with near-mature fruit.

FIG. 4 shows ‘Yoli’ plants with runners growing in the field.

FIG. 5 shows petiole leaf of ‘Yoli’ plants.

FIG. 6 shows the flower and calyx of ‘Yoli’ plants.

DETAILED DESCRIPTION OF THE INVENTION

‘Yoli’ (a.k.a. ‘104522’)

This invention relates to a new and distinctive day-neutral type cultivar designated as ‘Yoli’. It is primarily adapted to the climate and growing conditions of the central coast of California. This region provides the necessary temperatures required for it to produce a strong vigorous plant and to remain in fruit production from March through October. The nearby Pacific Ocean provides the needed humidity and moderate day temperatures and evening chilling to maintain fruit quality for the production months.

The following traits in combination distinguish strawberry variety ‘Yoli’ from the known strawberry varieties. Plants for the botanical measurements in the present application were grown as annuals. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

‘Yoli’ has not been observed under all possible environmental conditions, and the phenotype may vary significantly with variations in environment.

Hybrid Patent Data Sheet (planted: October Year 1; measurement: November, Year 2) ‘Yoh’ (a.k.a. ‘104522’)	
Plant	
Plant Growth Habit	semi-upright
Day length	neutral
Plant season	fall
Height (mm)	310
Diameter (mm)	430 (the entire plant structure, including the “superstructure” of petioles, leaves, flowers)
	60 (inner “base” or “heart” structure of the plant, i.e. the mature apical and axial crowns of the strawberry plant where petioles, flowers and runners are attached)
Density of foliage	medium
Plant vigor	moderate to high
Rain/weather tolerance	moderate
Harvest Ease	moderate
Terminal Leaf	
Length (mm)	90
Width (mm)	90
Terminal leaflet length/width ratio	1
Blistering	none
Terminal Leaflet shape	orbicular
Number of teeth per terminal leaflet	20-22
Shape of the terminal leaflet base	obtuse
Shape of teeth	obtuse
shape of terminal leaflet in cross-section	concave
margin description of the terminal	serrate
Color of upper side of leaves	136A
Color of lower side of leaves	137C
Leaf glossiness	medium to strong
Presence or absence of any leaf variegation	absence
No. of leaflets:	62
Terminal Leaflet margin	revolute
Terminal Leaflet shape of apex	rounded
Petiole	
Length (mm)	244
Width (mm)	3.9
Petiole pubescence	medium
Petiole pose of hairs	outwards
Petiole color	145B
Petiolule	
Length (mm)	28
Width(mm)	2.6
Stolon	
Count (mm)	43
Diameter (mm)	3
Widest diameter of stolon at point of attachment of leaf (mm)	4.1
Density of pubescence on the stolon	medium
Stolon number	8
Stolon thickness	medium to large
Stolon color	145A
Stolon anthocyanin	absent
Inflorescence	
Petal/flower	
Corolla (mm)	25
Calyx (mm)	25

-continued

Hybrid Patent Data Sheet (planted: October Year 1; measurement: November, Year 2) ‘Yoh’ (a.k.a. ‘104522’)		
5	Calyx diameter relative to corolla	1
	Position of calyx	even
	Color of calyx	137A
	Pose of calyx segments	spreading
10	Size of calyx in relation to fruit	smaller
	Petal length (mm)	13
	Petal width (mm)	13
	Petal length/width ratio	1
	Petal number per flower	5-6
15	Sepal length (mm)	9
	Sepal width (mm)	6
	Sepal number per flower	10-15
	Peduncle size	Medium to long
	Stipules length (mm)	35
	Stipules width (mm)	11
20	Stipule pubescence	medium
	Stipule anthocyanin	mild presence
	Stipule color	145B to 181D
	Pedicel color	145A
	Attitude of hairs on petiole and pedicel	horizontal
25	Inflorescence position relative to foliage	even to above foliage
	flower arrangement of petals	overlapping
	Number of flowers per plant	~20
	Flower size	medium
	Petal shape	orbicular
	Petal apex	rounded
30	Petal margin	entire
	Petal base shape	concave
	Petal spacing	overlapping
	Upper and lower Petal color	155C
	Sepal Shape	elliptical
	Sepal apex	convex
35	Number of stamen	22
	Receptacle color	4A
	Anther color	14A
	Fertility	Self-fertile
	Fruit	
40	Fruit/truss	2-5
	Fruit truss length (mm)	235
	Fruiting truss diameter (mm)	21.7-26.7
	Fruit truss attitude	Erect
	Truss color	145A
	Length (mm)	47
	Width (mm)	39
45	Fruit length/width ratio	1.2
	Core length (mm)	30
	Core width (mm)	14
	Glossiness	medium
	Fruit skin color	41A
	Fruit weight (g)	21
50	Predominant fruit shape	long-conical
	Difference in shape between primary & secondary fruits	none to very slight
	Width of band without of achenes (mm)	2
	Position of achenes	even to above
55	Achene color	145A
	Achenes per fruit	275
	Achene weight (g)	0.25
	Firmness of flesh	soft to firm
	Color of the flesh	41A
	Evenness of flesh color	even to slightly uneven
60	Grams of fruit/plant (During peak season, June)	497
	Appearance Score (from 1-5, with 5 = best)	4.5
	Storage longevity	9 to 13 days
	Cull rate: Non salable fruit production rate	<25%
65		

-continued

Hybrid Patent Data Sheet	
(planted: October Year 1; measurement: November, Year 2)	
‘Yoh’ (a.k.a. ‘104522’)	
Disease and pest resistance	Not tested
Acidity as measured by pH	NA
Sweetness as measured by brix	NA
(NA: Not Available)	

When ‘Yoli’ is compared to ‘Seascape’ (U.S. Plant Pat. No. 07,614), ‘Yoli’ has a margin of terminal leaflet that is

serrate, while the margin of the terminal leaflet of ‘Seascape’ is crenate. In addition, the leaf glossiness of ‘Yoli’ ranges from medium to strong, while the leaf glossiness of ‘Seascape’ is medium. The number of leaflets of ‘Yoli’ is 62, while the number of leaflets of ‘Seascape’ is 49. ‘Yoli’ also has a smaller fruit, higher yield when compared to ‘Seascape’, and the flowers of ‘Yoli’ are always presenting above foliage.

The invention claimed is:

1. A new and distinct cultivar of strawberry plant named ‘Yoli’ substantially as shown and described herein.

* * * * *

FIG. 1






FIG. 2

FIG. 3





FIG. 4



FIG. 5




FIG. 6.