



(12) **United States Plant Patent**
Korlipara

(10) **Patent No.:** **US PP28,908 P2**
(45) **Date of Patent:** **Jan. 30, 2018**

(54) **SCROPHULARIA MACRANTHA PLANT**
NAMED ‘TNSCRCR’

(50) Latin Name: *Scrophularia macrantha*
Varietal Denomination: **TNSCRCR**

(71) Applicant: **Terra Nova Nurseries, Inc.**, Canby,
OR (US)

(72) Inventor: **Harini Korlipara**, Canby, OR (US)

(73) Assignee: **Terra Nova Nurseries, Inc.**, Canby,
OR (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/330,804**

(22) Filed: **Nov. 7, 2016**

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./263.1**

(58) **Field of Classification Search**
USPC Plt./263.1
See application file for complete search history.

Primary Examiner — Annette H Para

(74) *Attorney, Agent, or Firm* — Klarquist Sparkman,
LLP

(57) **ABSTRACT**

A new and distinct form of *Scrophularia macrantha* plant
characterized by deep rose red flowers in dense panicle
cymes, blooms all summer, a short upright shrubby habit,
excellent branching with flower spikes at end of every
branch, and excellent vigor.

1 Drawing Sheet

1

Botanical denomination: *Scrophularia macrantha*.
Variety designation: ‘TNSCRCR’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Scrophularia macrantha*, given the name, ‘TNSCRCR’.
Scrophularia is in the family Scrophulariaceae. This new
cultivar was part of a planned breeding program for a shorter
Scrophularia macrantha. The parents were proprietary,
unreleased seedlings of *Scrophularia macrantha*. *Scrophu-*
laria macrantha, commonly known as Red Birds in a Tree
or New Mexican Figwort, is a rare native of the mountains
of New Mexico which blooms all summer with charming,
small reddish flowers that resemble birds. It is also drought
tolerant and loved by hummingbirds. Normal height for this
plant is three to four feet.

Compared to the representative members of the species
Scrophularia macrantha, the new cultivar is much shorter
with denser inflorescences of deep rose red. Compared to the
specific male and female *Scrophularia macrantha* parent
seedlings, the new cultivar is shorter, better branched, and
has larger flowers.

The new cultivar exhibits the following characteristics:

1. deep rose red flowers in dense panicle cymes,
2. blooms all summer,
3. a short upright shrubby habit,
4. excellent branching with flower spikes at end of every
branch, and
5. excellent vigor.

The new variety has been reproduced only by asexual
propagation (stem cuttings and micropropagation). Each of
the progeny exhibits identical characteristics to the original
plant. Asexual propagation by division and micropropaga-
tion as done in Canby, Oreg., shows that the foregoing
characteristics and distinctions come true to form and are
established and transmitted through succeeding propaga-
tions. The present invention has not been evaluated under all

2

possible environmental conditions. The phenotype may vary
with variations in environment without a change in the
genotype of the plant.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows 7 month-old *Scrophularia macrantha*
‘TNSCRCR’ (with a with background) in July grown in a
gallon container in full sun in Canby, Oreg.

FIG. 2 shows 7 month-old *Scrophularia macrantha*
‘TNSCRCR’ growing full sun in the trial field in July in
Canby, Oreg.

DETAILED PLANT DESCRIPTION

The following is a detailed description of the new *Scro-*
phularia macrantha cultivar based on observations of 7
month-old specimens grown outside in the sun in the trial
field in Canby, Oreg. Canby is Zone 8 on the USDA
Hardiness map. Temperatures range from a high of 95° F. in
August to an average of 32° F. in January. Normal rainfall
in Canby is 42.8 inches per year. The color descriptions are
all based on The Royal Horticultural Society Colour Chart,
5th edition.

Plant:

Form.—Upright and bushy.

Type.—Evergreen, subshrub/perennial.

Cold hardiness.—USDA Zone 5-10.

Size.—Grows to 50 cm wide and 55 cm tall, including
flowers.

Vigor.—Excellent.

Branching.—Extremely well-branched with about 5
main stems from the ground.

Roots.—Fibrous, freely branching, Greyed Orange
164D; plants root easily from stem cuttings.

Stem:

Size.—Grows to about 25 cm long and 4 mm wide.

Internodes.—4 cm to 5 cm long on main branches.

Surface.—Glandular pubescent.

Color.—Yellow Green 148B.

Leaf:

Type.—Simple, aromatic.

Arrangement.—Opposite.

Blade size.—Grow to 65 mm long and 38 mm wide.

Shape.—Deltoid to lanceolate.

Venation.—Pinnate, Yellow Green 146B on both sides.

Margins.—Coarsely serrate (sometimes biserrate) to dentate.

Apex.—Acute.

Base.—Truncate to cuneate.

Surface texture.—Both sides glabrous.

Petiole description.—6 mm to 25 mm long and 1 mm to 3 mm wide, glabrous, Yellow Green 146B.

Leaf color.—Top side Green N137D, bottom side closest to Yellow Green 147B.

Inflorescence:

Type.—Terminal paniculate cyme.

Number of flowers.—About 250 per cyme.

Size of inflorescence.—Grows to 24 cm long and 9 cm wide.

Peduncle.—Grows to 4.5 cm long from terminal leaves to base of inflorescence, Yellow Green 148B, glandular pubescent.

Pedicel.—3 mm long and 1 mm wide, glandular pubescent, Yellow Green 148B.

Bloom period.—Middle of May to October in Canby, Oreg.

Flower bud:

Size.—10 mm long and 5 mm wide at the widest point prior to opening.

Description.—Bulbous.

Surface texture.—Glandular pubescent.

Color.—Greyed Purple 185C.

Flower:

Type.—Perfect.

Shape.—Zygomorphic, tubular.

Corolla description.—7 mm wide and 19 mm long; glandular pubescent outside and glabrous inside, margins entire; tube 10 mm long and 7 mm wide; 4 main lobes in two lips, upper lip prominent, 5 mm wide and 5 mm long, with two overlapping lobes, ovate with tips obtuse; lower lip 3-lobed with two side lobes, 3 mm long and 4 mm wide, tip obtuse, bottom lobe slightly reflexed 2.5 mm long and 2 mm wide, tip obtuse; outside color between Red 46A and Red Purple 60A, inside closest to Greyed Red 181D.

Calyx description.—5-lobed, 5 mm wide and 4 mm deep, lobes ovate, 2 mm long and 2 mm wide, tips acute, margins entire, Yellow Green 147B on both sides, glandular pubescent outside and glabrous inside.

Pistil description.—1, 11 mm long, ovary conical, 3 mm long and 2 mm wide, Yellow Green 145C, style 8 mm long, Greyed Purple 186D, stigma White NN155A.

Stamen description.—4 and 1 staminode, White 155A, staminode attached dorsally, 12 mm long and 0.5 mm wide, filaments 10 mm long and 0.5 mm wide, anthers 1.5 mm long and 2 mm wide, White 155B, pollen White NN155A.

Fragrance.—None.

Lastingness.—A cyme blooms for about 4 weeks on the plant.

Fruit: A capsule, 10 mm long and 4 mm wide, Brown 200A, fertility good.

Seed: Oval, Brown N200A.

Pests and diseases: The new cultivar is typical to the genus. No known resistances to pests or diseases. No problems have been found in Canby, Oreg.

I claim:

1. A new and distinct form of *Scrophularia macrantha* plant substantially as shown and described.

* * * * *



FIG. 1 on left
FIG. 2 below

