

US00PP31650P2

(12) **United States Plant Patent**
Caster et al.

(10) **Patent No.:** **US PP31,650 P2**
(45) **Date of Patent:** **Apr. 14, 2020**

(54) **BLUEBERRY PLANT VARIETY NAMED
‘DRISBLUESEVENTEEN’**

(50) Latin Name: *Vaccinium corymbosum* L.
Varietal Denomination: **DrisBlueSeventeen**

(71) Applicant: **Driscoll’s, Inc.**, Watsonville, CA (US)

(72) Inventors: **Brian K. Caster**, Watsonville, CA
(US); **Jennifer K. Izzo**, Watsonville,
CA (US)

(73) Assignee: **Driscoll’s, Inc.**, Watsonville, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/350,021**

(22) Filed: **Sep. 13, 2018**

(51) **Int. Cl.**
A01H 6/36 (2018.01)
A01H 5/08 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./157**

(58) **Field of Classification Search**
USPC **Plt./156, 157**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP6,699 P 3/1989 Wood
PP11,807 P2 3/2001 Lyrene

PP12,783 P2 7/2002 Lyrene
PP20,436 P2 10/2009 Caster et al.
PP20,449 P2 11/2009 Caster et al.
PP20,488 P2 11/2009 Caster et al.
PP24,407 P3 4/2014 Caster et al.
PP24,489 P3 5/2014 Caster et al.
PP24,568 P3 6/2014 Caster et al.
PP24,569 P3 6/2014 Caster et al.
PP24,605 P3 7/2014 Caster et al.
PP26,287 P3 1/2016 Caster et al.
PP26,451 P3 3/2016 Caster et al.
PP26,537 P3 3/2016 Caster et al.
PP26,643 P3 4/2016 Caster et al.
PP26,748 P3 5/2016 Caster et al.
PP27,622 P3 1/2017 Caster et al.
PP28,933 P2 2/2018 Caster et al.

OTHER PUBLICATIONS

Caster et al., Unpublished U.S. Appl. No. 16/350,012, filed Sep. 12,
2018, titled “Blueberry Plant Variety Named ‘DrisBlueEighteen’”.
Mowrey et al., Unpublished U.S. Appl. No. 16/350,020, filed Sep.
13, 2018, titled “Blueberry Plant Variety Named ‘DrisBlueSixteen’”.

Primary Examiner — Susan McCormick Ewoldt

Assistant Examiner — Karen M Redden

(74) *Attorney, Agent, or Firm* — Morrison & Foerster
LLP

(57) **ABSTRACT**

A new and distinct variety of blueberry plant named ‘Dris-
BlueSeventeen’, particularly selected for its mid-range chill
requirement suitable for production in latitudes ranging from
30 to 40 degrees, its productiveness, as well as its extremely
large, firm fruit with a balanced sweet/tart flavor and an
attractive appearance, is disclosed.

6 Drawing Sheets

1

Latin name:
Botanical classification: *Vaccinium corymbosum* L.
Varietal denomination: The varietal denomination of the
claimed variety of blueberry plant is ‘DrisBlueSeventeen’.

BACKGROUND OF THE INVENTION

Blueberry plants are perennial flowering plants with
indigo-colored berries from the section *Cyanococcus* within
the genus *Vaccinium*. Many commercially sold species with
English common names, including blueberry, are currently
classified in section *Cyanococcus* of the genus *Vaccinium*
and come predominantly from North America. Many North
American native species of blueberries are grown commer-
cially in the Southern Hemisphere in Australia, New Zea-
land, and South American nations.

Vaccinium corymbosum, the northern highbush blueberry,
is a North American species of blueberry which has become
a food crop of significant economic importance. It is native
to eastern Canada and the eastern and southern United
States, from Ontario east to Nova Scotia and south as far as
Florida and eastern Texas. It has been naturalized in Europe,
Japan, New Zealand, and the Pacific Northwest of North

2

America. Other common names include blue huckleberry,
tall huckleberry, swamp huckleberry, high blueberry, and
swamp blueberry.

Blueberries are usually erect, prostrate shrubs that can
vary in size from approximately four inches to approxi-
mately 13 feet in height. In the commercial production of
blueberries, the smaller species are known as “lowbush
blueberries”, while the larger species are known as “high-
bush blueberries”.

Blueberry bushes typically bear fruit in the middle of the
growing season. However, fruiting times can be affected by
local conditions such as altitude and latitude. As such, peak
crop can vary from May to August in the northern hemi-
sphere, depending upon these conditions.

Blueberries are a popular fruit that is typically consumed
as fresh fruit, individually quick frozen (IQF) fruit, or in
prepared foods, such as purées, juices, jellies, jams, baked
goods, snack foods, and cereals.

Blueberry is an important and valuable fruit crop. Accord-
ingly, there is a need for new varieties of blueberry plant. In
particular, there is a need for improved varieties of blueberry
plant that are stable, high yielding, and agronomically
sound.

SUMMARY OF THE INVENTION

In order to meet these needs, the present invention is directed to an improved variety of blueberry plant. In particular, the invention relates to a new and distinct variety of blueberry plant (*Vaccinium corymbosum* L.), which has been denominated as 'DrisBlueSeventeen'.

Blueberry plant variety 'DrisBlueSeventeen' was discovered in Santa Cruz County, Calif. in September of 2006 and originated from a cross between the proprietary female parent blueberry plant '136D 2' (unpatented) and the proprietary male parent blueberry plant '8B 4' (unpatented). The original seedling of the new variety was first asexually propagated via cuttings in Monterey County, Calif. in 2007. 'DrisBlueSeventeen' was subsequently asexually propagated via cuttings and underwent further testing in Santa Cruz County, Calif. for ten years (2008 to 2017). The present blueberry variety has been found to be stable and reproduce true to type through successive asexual propagations via cuttings.

'DrisBlueSeventeen' exhibits the following distinguishing characteristics when grown under normal horticultural practices in Santa Cruz County, Calif.:

1. Large fruit size;
2. Firm fruit;
3. High fruit sweetness; and
4. Attractive fruit appearance.

The original seedling of the new variety was selected for its mid-range chill requirement suitable for production in latitudes ranging from 30 to 40 degrees, its productiveness, as well as its extremely large, firm fruit with a balanced sweet/tart flavor and an attractive appearance.

BRIEF DESCRIPTION OF THE DRAWINGS

This new blueberry plant variety is illustrated by the accompanying photographs. The colors shown are as true as can be reasonably obtained by conventional photographic procedures. The photographs are of plants that are seven years old.

FIG. 1 illustrates a section of a cane of variety 'DrisBlueSeventeen'.

FIG. 2 illustrates leaves of variety 'DrisBlueSeventeen'.

FIG. 3 illustrates flowers of variety 'DrisBlueSeventeen'.

FIG. 4 illustrates whole fruit of variety 'DrisBlueSeventeen'.

FIG. 5 illustrates whole fruit and cross-sections of the fruit of variety 'DrisBlueSeventeen'.

FIG. 6 illustrates plants of variety 'DrisBlueSeventeen'.

DETAILED BOTANICAL DESCRIPTION

The following description sets forth the distinctive characteristics of 'DrisBlueSeventeen'. The data which define these characteristics is based on observations taken in Santa Cruz County, Calif. from 2008 to 2017. This description is in accordance with UPOV terminology. Color designations, color descriptions, and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions. 'DrisBlueSeventeen' has not been observed under all possible environmental conditions. The botanical description of 'DrisBlueSeventeen' was taken from 7-year-old plants, unless indicated otherwise. The indicated values represent averages calculated from measurements of several plants. Color references are primarily

to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.) (2007 edition). Descriptive terminology follows the *Plant Identification Terminology, An Illustrated Glossary*, 2nd edition by James G. Harris and Melinda Woolf Harris, unless where otherwise defined.

Classification:

Family.—Ericaceae.

Botanical.—*Vaccinium corymbosum* L.

Common name.—Blueberry.

Variety name.—'DrisBlueSeventeen'.

Parentage:

Female parent.—The proprietary blueberry plant '136D 2'.

Male parent.—The proprietary blueberry plant '8B 4'.

Plant:

Height.—133.0 cm.

Width.—198.0 cm.

Length/width ratio.—0.7.

Growth habit.—Semi-upright.

One-year-old canes (young canes).—Length: 36 cm.

Diameter at the base: 6 mm. Diameter at the tip: 2 mm. Internode length on the upper half: 15.0 mm. Color: RHS 145B (Medium yellow-green). One-year-old cane texture: Smooth.

Five-year-old canes (mature canes).—Length: 73 cm.

Diameter at the base: 25 mm. Diameter at the tip: 7 mm. Five-year-old cane texture: Rough. Five-year-old cane color: RHS 199D (Light greyed-brown). Flower bud anthocyanin color: RHS N59C (Dark red-purple).

Leaves:

Length.—68.2 mm.

Width.—35.1 mm.

Length/width ratio.—1.9.

Internode length.—14.98 mm.

Shape.—Elliptic.

Margin.—Entire.

Color on upper side.—RHS 139A (Dark green).

Leaf arrangement.—Alternate.

Leaf venation pattern.—Reticulate.

Leaf vein color.—RHS 151B (Strong greenish yellow).

Shape of the leaf apex.—Cuspidate.

Shape of the leaf base.—Cuneate.

Leaflet texture of upper side.—Smooth.

Leaflet texture of lower side.—Smooth.

Leaflet color of lower side.—RHS N138C (Light green).

Petiole.—Length: 3.6 mm. Diameter: 1.45 mm. Petiole texture: Slightly pubescent. Petiole color: RHS 139D (Medium green).

Flowers:

Length (excluding peduncle).—9.60 mm.

Diameter.—7.85 mm.

Length/width ratio.—1.2.

Presence of flower fragrance.—Faint.

Flower bud.—Length: 7 mm. Width: 4 mm. Number of flowers per bud: 6.

Pedicel.—Length: 3.30 mm. Diameter: 1.09 mm. Pedicel color: RHS 142C (Light green).

Corolla.—Shape: Urceolate. Anthocyanin color of corolla tube: RHS N59C (Dark red-purple). Ridges on tube: Present. Petal width (ridge to ridge): 4.08 mm. Petal color: RHS 1570 (Green-white). Diameter of corolla aperture: 3.88 mm.

Reproductive organs.—Style length (including stigma): 7.01 mm. Ovary color: RHS 191B (Medium greyed-green). Anther color: RHS N172D (Medium greyed-orange). Pollen color: RHS 4D (Light yellow).

Flowering interval on one-year shoot.—March to April.

Pollinator requirement.—Insect pollinators such as honeybees or bumblebees are recommended.

Fruit:

Length.—15.06 mm.

Diameter.—18.99 mm.

Length/width ratio.—0.8.

Shape in longitudinal sections.—Round.

Attitude of sepals.—Semi-erect.

Type of sepals.—Incurving.

Calyx basin.—Diameter: 6.51 mm. Depth: 2.48 mm. Diameter/depth ratio: 2.6.

Weight.—3.4 g.

Number of berries per cluster.—6.20.

Peduncle length.—44.27 mm.

Diameter of pedicel.—1.09 mm.

Seed.—Length: 1.84 mm. Width: 0.99 mm. Length/width ratio: 1.9. Seed color: RHS N172C (Medium greyed-orange).

Color of unripe fruit.—RHS 144A (Medium yellow-green).

Color of skin (after removal of bloom).—RHS 95A (Dark violet-blue).

Color of flesh.—RHS 145C (Light yellow-green).

Fruit firmness.—Firm.

Fruiting type.—On one-year-old shoots.

Ripening interval on one-year-old shoot.—June to July.

Approximate date of first fruit pick.—June 1.

Approximate date of last fruit pick.—August 15.

Market use of fruit.—Fresh market.

Shipping and storage characteristics.—Following harvest, fruit have been stored for 21 days when maintained under cooled temperatures that are standard for blueberry storage.

Yield.—5000 kg/hectare to 10,000 kg/hectare of fruit per season from 60-month-old plants when grown in Santa Cruz County, Calif.

Resistance to abiotic stress, pests, and diseases:

Heat.—Moderately susceptible.

Spotted-wing drosophila (drosophila suzukii).—Moderately susceptible.

Botrytis fruit rot (botrytis cinerea).—Moderately resistant.

Botryosphaeria stem blight.—Moderately susceptible.

Phomopsis stem blight.—Moderately susceptible.

COMPARISONS TO PARENTAL AND COMMERCIAL BLUEBERRY VARIETIES

‘DrisBlueSeventeen’ differs from the proprietary female parent ‘136D 2’ (unpatented) in that fruit of ‘DrisBlueSeventeen’ are larger and have better appearance compared to fruit of ‘136D 2’.

‘DrisBlueSeventeen’ differs from the proprietary male parent ‘8B 4’ (unpatented) in that fruit of ‘DrisBlueSeventeen’ are larger and have better appearance compared to fruit of ‘8B 4’.

‘DrisBlueSeventeen’ differs from commercial blueberry plant variety ‘DrisBlueOne’ (U.S. Plant Pat. No. 20,449) in that flowers of ‘DrisBlueSeventeen’ have a small size of corolla tube, whereas flowers of ‘DrisBlueOne’ have a large size of corolla tube. Flowers of ‘DrisBlueSeventeen’ have strong anthocyanin coloration of corolla tube, whereas flowers of ‘DrisBlueOne’ have absent or very weak anthocyanin coloration of corolla tube. Further, fruit of ‘DrisBlueSeventeen’ have a dark blue color of skin after removal of bloom, whereas fruit of ‘DrisBlueOne’ have a medium blue color of skin after removal of bloom. Additionally, fruit of ‘DrisBlueSeventeen’ are high in sweetness, whereas fruit of ‘DrisBlueOne’ are medium in sweetness.

‘DrisBlueSeventeen’ differs from commercial blueberry plant variety ‘DrisBlueFive’ (U.S. Plant Pat. No. 24,489) in that plants of ‘DrisBlueSeventeen’ have medium vigor, whereas plants of ‘DrisBlueFive’ have strong vigor. Fruit of ‘DrisBlueSeventeen’ have a small size of corolla tube, whereas fruit of ‘DrisBlueFive’ have a medium size of corolla tube. Further, fruit of ‘DrisBlueSeventeen’ have strong anthocyanin coloration of corolla tube, whereas fruit of ‘DrisBlueFive’ have absent or very weak anthocyanin coloration of corolla tube. Additionally, fruit of ‘DrisBlueSeventeen’ have a round shape in longitudinal section, whereas ‘DrisBlueFive’ have an oblate shape in longitudinal section.

What is claimed is:

1. A new and distinct variety of blueberry plant designated ‘DrisBlueSeventeen’ as shown and described herein.

* * * * *




FIG. 1




FIG. 2




FIG. 3



FIG. 4



FIG. 5



FIG. 6