



(12) **United States Plant Patent**
Mendoza et al.

(10) **Patent No.:** **US PP34,072 P2**
(45) **Date of Patent:** **Mar. 29, 2022**

- (54) **STRAWBERRY PLANT NAMED
‘DRISSTRAWWEIGHTYNINE’**
- (50) Latin Name: *Fragaria x ananassa*
Varietal Denomination: **DrisStrawEightyNine**
- (71) Applicant: **Driscoll’s, Inc.**, Watsonville, CA (US)
- (72) Inventors: **Omar Carrillo Mendoza**, Watsonville,
CA (US); **Renae R. Robertson**,
Watsonville, CA (US); **Kevin Coons**,
Watsonville, CA (US); **Philip J.**
Stewart, Watsonville, CA (US)
- (73) Assignee: **Driscoll’s, Inc.**, Watsonville, CA (US)
- (*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **17/196,925**
- (22) Filed: **Mar. 9, 2021**
- (51) **Int. Cl.**
A01H 5/08 (2018.01)
A01H 6/74 (2018.01)
- (52) **U.S. Cl.**
USPC **Plt./209**
- (58) **Field of Classification Search**
USPC Plt./209, 208
CPC ... A01H 5/08; A01H 5/00; A01H 6/74; A01H
6/7409
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS

PP1,745 P 8/1958 Lang
PP3,981 P 11/1976 Bringhurst et al.
PP4,487 P 11/1979 Bringhurst et al.
PP4,538 P 5/1980 Bringhurst et al.
PP5,262 P 7/1984 Voth et al.
PP5,265 P 7/1984 Voth et al.
PP5,266 P 7/1984 Bringhurst et al.
PP5,300 P 10/1984 Johnson, Jr. et al.
PP5,840 P 12/1986 Johnson, Jr. et al.
PP6,191 P 5/1988 Johnson, Jr. et al.
PP6,231 P 7/1988 Johnson, Jr. et al.
PP6,578 P 1/1989 Voth et al.
PP6,579 P 1/1989 Bringhurst et al.
PP7,024 P 9/1989 Johnson, Jr. et al.
PP7,172 P 2/1990 Voth et al.
PP7,522 P 5/1991 Johnson, Jr. et al.
PP7,614 P 8/1991 Bringhurst et al.
PP7,615 P 8/1991 Bringhurst et al.
PP8,086 P 1/1993 Nelson et al.
PP8,205 P 4/1993 Nelson et al.
PP8,649 P 3/1994 Sjulín et al.
PP8,661 P 3/1994 Bringhurst et al.
PP8,708 P 5/1994 Voth et al.
PP8,745 P 5/1994 Sjulín et al.
PP9,130 P 5/1995 Sjulín et al.
PP9,909 P 6/1997 Ackerman et al.
PP10,221 P 2/1998 Sjulín et al.
PP10,534 P 8/1998 Sjulín et al.
PP10,642 P 10/1998 Amorao et al.
PP11,035 P 8/1999 Mowrey et al.
PP11,277 P 3/2000 Gilford et al.

PP11,279 P 3/2000 Gilford et al.
PP11,522 P 9/2000 Amorao et al.
PP11,548 P 10/2000 Amorao et al.
PP11,554 P 10/2000 Sjulín et al.
PP11,639 P 11/2000 Mowrey et al.
PP12,186 P2 11/2001 Gilford et al.
PP12,436 P2 3/2002 Amorao et al.
PP12,577 P2 4/2002 Amorao et al.
PP12,817 P2 7/2002 Gilford et al.
PP12,899 P2 9/2002 Mowrey et al.
PP13,386 P2 12/2002 Mowrey et al.
PP13,469 P3 1/2003 Larson et al.
PP14,005 P3 7/2003 Amorao et al.
PP14,062 P3 8/2003 Amorao et al.
PP14,109 P3 8/2003 Gilford et al.
PP14,771 P3 5/2004 Amorao et al.
PP15,145 P2 9/2004 Mowrey et al.
PP15,308 P2 11/2004 Sjulín et al.
PP15,375 P2 11/2004 Mowrey et al.
PP15,435 P2 12/2004 Sjulín et al.
PP15,596 P2 3/2005 Amorao et al.
PP15,731 P2 4/2005 Amorao et al.
PP15,752 P2 5/2005 Gilford et al.
PP16,070 P2 10/2005 Gilford et al.
PP16,238 P2 2/2006 Amorao et al.
PP16,241 P2 2/2006 Mowrey et al.
PP16,298 P2 2/2006 Gilford et al.
PP16,299 P2 2/2006 Gilford et al.
PP16,475 P2 4/2006 Gilford et al.
PP16,558 P3 5/2006 Lopez
PP18,000 P2 9/2007 Meulenbroek
PP18,040 P3 9/2007 Mowrey et al.
PP18,041 P3 9/2007 Gilford
PP18,458 P2 1/2008 Ferguson et al.
PP18,575 P3 3/2008 Amorao et al.
PP18,878 P2 6/2008 Mowrey et al.
PP19,240 P2 9/2008 Gilford et al.
PP19,673 P3 2/2009 Ferguson et al.
PP19,767 P2 2/2009 Shaw et al.
PP20,248 P3 9/2009 Rogers et al.
PP20,363 P2 9/2009 Chandler
PP20,701 P2 2/2010 Gilford et al.
PP20,731 P2 2/2010 Mowrey et al.
PP20,733 P2 2/2010 Mowrey et al.
PP20,735 P2 2/2010 Ferguson
PP20,775 P2 2/2010 Mowrey et al.
PP20,922 P2 4/2010 Gilford et al.
PP21,538 P2 11/2010 Gilford et al.
PP21,559 P2 12/2010 Ferguson et al.
PP21,762 P2 3/2011 Stewart et al.
PP22,040 P3 7/2011 Stewart et al.
PP22,218 P2 11/2011 Ferguson et al.
PP22,247 P2 11/2011 Ferguson
PP23,107 P2 10/2012 Ferguson et al.
PP23,148 P2 10/2012 Gilford et al.
PP23,377 P2 2/2013 Ferguson et al.
PP23,378 P2 2/2013 Pullen et al.
PP23,382 P2 2/2013 Ferguson et al.
PP23,383 P2 2/2013 Ferguson et al.
PP23,400 P2 2/2013 Ferguson et al.
PP23,401 P2 2/2013 Pullen et al.

(Continued)

Primary Examiner — June Hwu
(74) Attorney, Agent, or Firm — Morrison & Foerster
LLP

(57) **ABSTRACT**

A new and distinct variety of strawberry plant named ‘DrisStrawEightyNine’, particularly selected for the size, firmness, juiciness, shelf life, and shipping capability of its fruit, as well as its volume of production and moderate resistance to two-spotted spider mite, is disclosed.

6 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

PP23,459 P2

PP23,506 P3

PP23,517 P3

PP24,096 P3

PP24,317 P3

PP24,333 P3

PP24,395 P3

PP24,533 P3

PP24,745 P2

PP25,408 P3

PP25,437 P3

PP25,698 P3

PP25,699 P3

PP25,747 P3

PP25,866 P3

PP26,800 P3

PP26,801 P3

PP26,802 P3

PP27,442 P2

PP27,645 P3

PP27,682 P3

PP27,711 P3

PP27,813 P3

PP29,289 P3

PP29,728 P2

PP29,729 P2

PP29,730 P2

3/2013

4/2013

4/2013

12/2013

3/2014

3/2014

4/2014

6/2014

8/2014

4/2015

4/2015

7/2015

7/2015

7/2015

9/2015

6/2016

6/2016

6/2016

12/2016

2/2017

2/2017

2/2017

3/2017

5/2018

10/2018

10/2018

10/2018

Stewart et al.

Ferguson et al.

Ferguson et al.

Fear et al.

Ferguson et al.

Vitten et al.

Vitten et al.

Ferguson et al.

Vitten et al.

Vitten et al.

Vitten et al.

Ferguson et al.

Stewart et al.

Kibbe et al.

Ferguson et al.

Stewart et al.

Stewart et al.

Rodriguez Alcazar et al.

Kibbe et al.

Vitten et al.

Kibbe et al.

Vitten et al.

Ferguson et al.

Vitten et al.

Stewart et al.

Kibbe et al.

Kibbe et al.

PP29,731 P2

PP29,747 P2

PP29,748 P2

PP29,749 P2

PP30,775 P2

PP30,789 P2

PP30,818 P2

PP30,851 P2

PP30,936 P2

PP31,083 P2

PP31,233 P2

PP31,527 P3

PP31,655 P2

PP31,703 P2

PP31,827 P2

PP31,896 P2

PP31,935 P2

PP32,079 P2

PP32,080 P2

PP32,271 P2

PP32,305 P3

PP32,498 P2

PP32,499 P3

PP32,500 P3

PP32,801 P2

PP32,814 P2

PP32,824 P2

PP33,070 P2

PP33,090 P2

2003/0079263 P1

2013/0276182 P1

10/2018

10/2018

10/2018

10/2018

8/2019

8/2019

8/2019

8/2019

10/2019

11/2019

12/2019

3/2020

4/2020

4/2020

6/2020

6/2020

7/2020

8/2020

8/2020

10/2020

10/2020

11/2020

11/2020

11/2020

2/2021

2/2021

2/2021

5/2021

5/2021

4/2003

10/2013

Ferguson et al.

Vitten et al.

Vitten et al.

Stewart et al.

Carrillo Mendoza et al.

Ferguson et al.

Pakozdi et al.

Fear et al.

Stewart et al.

Carrillo Mendoza et al.

Pakozdi et al.

Carrillo Mendoza et al.

Carrillo Mendoza et al.

Stewart et al.

Ferguson et al.

Pakozdi et al.

Pakozdi et al.

Carrillo Mendoza et al.

Ferguson et al.

Ferguson et al.

Pakozdi et al.

Mendoza et al.

Jacobs et al.

Stewart et al.

Ferguson et al.

Stewart et al.

Stewart et al.

Mendoza et al.

Mendoza et al.

Gilford et al.

Fear et al.

1

STRAWBERRY PLANT NAMED
'DRISSTRAWIGHTYNINE'

Latin name:

Botanical classification: *Fragaria* x *ananassa*.

Varietal denomination: The varietal denomination of the claimed variety of strawberry plant is 'DrisStrawEighty-Nine'.

BACKGROUND OF THE INVENTION

Cultivated strawberry is a hybrid species of the genus *Fragaria* that is grown worldwide for its fruit. Modern strawberry was first bred in Brittany, France, in the 18th century by crossing *Fragaria virginiana* with *Fragaria chiloensis*. Strawberry fruit is an aggregate accessory fruit, with the fleshy part of the fruit being derived from the receptacle that holds the ovaries.

Strawberry varieties vary widely in color, size, shape, flavor, season of ripening, degree of fertility, and susceptibility to disease. Certain varieties vary in foliage, and some vary in the relative development of their reproductive organs. Typically, strawberry flowers appear hermaphroditic in structure, but function as either male or female. Generally, commercial production of strawberry plants involves propagation from runners and distribution as either plugs or bare root plants. Cultivation is either perennial or annual plasticulture. During the off season, strawberries can also be produced in greenhouses.

Strawberry fruit is widely appreciated for its characteristic bright red color, aroma, juicy texture, and sweetness. Strawberry fruit is a popular fruit that is generally consumed either fresh or in prepared foods, such as preserves and baked goods.

Strawberry is an important and valuable fruit crop. Accordingly, there is a need for new varieties of strawberry plants. In particular, there is a need for improved varieties of strawberry plant that are stable, high yielding, and agronomically sound.

SUMMARY OF THE INVENTION

In order to meet these needs, the present invention is directed to an improved variety of strawberry plant. In particular, the invention relates to a new and distinct variety of strawberry plant (*Fragaria* x *ananassa*), which has been denominated as 'DrisStrawEightyNine'.

Strawberry plant variety 'DrisStrawEightyNine' originated from a controlled cross between the proprietary female parent '88T205' (unpatented) and the proprietary male parent '107S 60' (unpatented). Progeny plants from this cross, including 'DrisStrawEightyNine', were asexually propagated via stolons in Shasta County, Calif. in the summer of 2012. Strawberry plant variety 'DrisStrawEightyNine' was later specifically identified and selected in Santa Barbara County, Calif. in May of 2013.

'DrisStrawEightyNine' was subsequently asexually propagated via stolons, and has undergone testing at test plots in Tapalpa, Mexico and Purepero, Mexico for five years (2015 to 2020). The present variety has been found to be stable and reproduce true to type through successive asexual propagations via stolons and tissue culture.

'DrisStrawEightyNine' was particularly selected for the size, firmness, juiciness, shelf life, and shipping capability of its fruit, as well as its volume of production and moderate resistance to two-spotted spider mite.

2

DESCRIPTION OF THE DRAWINGS

This new strawberry plant is illustrated by the accompanying photographs. The colors shown are as true as can be reasonably obtained by conventional photographic procedures. Unless otherwise indicated, the photographs are of plants that are six months old.

FIG. 1 illustrates whole fruit of variety 'DrisStrawEighty-Nine'.

FIG. 2 illustrates longitudinal sections of fruit of variety 'DrisStrawEightyNine'.

FIG. 3 illustrates the upper surface (top row) and lower surface (bottom row) of flowers of variety 'DrisStrawEightyNine'.

FIG. 4 illustrates the upper surface (top) and lower surface (bottom) of leaves of variety 'DrisStrawEightyNine'.

FIG. 5 illustrates a view of whole plants of variety 'DrisStrawEightyNine'.

FIG. 6 illustrates another view of whole plants of variety 'DrisStrawEightyNine'.

DETAILED BOTANICAL DESCRIPTION

The following detailed descriptions set forth the distinctive characteristics of 'DrisStrawEightyNine'. The data which define these characteristics is based on observations taken in Tapalpa, Mexico and Purepero, Mexico from 2015 to 2020. This description is in accordance with UPOV terminology. Color designations, color descriptions, and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic, and cultural conditions. 'DrisStrawEightyNine' has not been observed under all possible environmental conditions. The botanical description of 'DrisStrawEightyNine' was taken from plants that were six months old. The indicated values represent averages calculated from measurements of several plants. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.) (2015 edition). Descriptive terminology follows the *Plant Identification Terminology, An Illustrated Glossary*, 2nd edition by James G. Harris and Melinda Woolf Harris, unless where otherwise defined.

Classification:

Species.—*Fragaria* x *ananassa*.

Common name.—Strawberry.

Denomination.—'DrisStrawEightyNine'.

Parentage:

Female parent.—Proprietary strawberry plant '88T205' (unpatented).

Male parent.—Proprietary strawberry plant '107S 60' (unpatented).

Plant:

Height.—18 cm.

Diameter.—47 cm.

Number of crowns per plant.—5.

Growth habit.—Semi-upright.

Density of foliage.—Medium.

Vigor.—Medium.

Stolon:

Number of daughter plants per square foot.—2.

Diameter at bract.—3.8 mm.

Anthocyanin coloration.—Weak.

Anthocyanin color when present.—RHS 39B (Deep yellowish pink).

3

Density of pubescence.—Medium.

Length.—25.4 cm.

Overall color.—RHS 144A (Strong yellow green).

Leaf:

Number of leaflets.—Three only.

Color of leaf upper surface.—RHS NN137A (Greyish olive green).

Color of lower surface.—RHS 137D (Moderate yellowish green).

Blistering.—Medium.

Glossiness.—Medium.

Variation.—Absent.

Leaf length.—10.2 cm.

Leaf width.—6.7 cm.

Terminal leaflet.—Length: 8.4 cm. Width: 6.3 cm. Length/width ratio: 1.3. Number of teeth per terminal leaflet: 20. Shape of base: Obtuse. Shape of apex: Rounded. Margin: Serrate to crenate. Shape in cross section: Concave.

Petiole.—Length: 15.8 cm. Diameter: 3.3 mm. Attitude of hairs: Slightly outwards. Bract frequency (number present on each petiole): 2. Color: RHS 144C (Strong yellow green).

Petiolule.—Length: 6 mm. Diameter: 1.7 mm. Color: RHS 144C (Strong yellow green).

Stipule.—Length: 4.0 cm. Width: 13 mm. Anthocyanin coloration: Weak. Anthocyanin color: RHS 52C (Deep pink). Overall color: RHS 145A (Strong yellow green).

Inflorescence:

Number of flowers per plant.—20.

Position of inflorescence in relation to foliage.—Same level.

Pedicel.—Length: 9.9 cm. Color: RHS 144C (Strong yellow green). Attitude of hairs: Slightly outwards.

Flower.—Flower diameter (petal tip to petal tip on non-flattened flower): 21 mm. Arrangement of petals: Overlapping. Size of calyx in relation to corolla: Larger. Stamen: Present.

Petal.—Length: 12 mm. Width: 12 mm. Length/width ratio: 1.0. Number of petals per flower: 6. Color of upper side: RHS NN155C (White). Color of lower side: RHS NN155C (White). Overall shape: Orbicular. Base shape: Concavo-convex. Apex shape: Rounded. Margin type: Smooth.

Calyx.—Diameter (sepal tip to sepal tip, measured on back of flower): 59 mm.

Sepal.—Length: 25 mm. Width: 12 mm. Number of sepals per flower: 11. Color of upper side: RHS 137A (Moderate olive green). Color of lower side: RHS 137D (Moderate yellowish green). Overall shape: Elliptical. Base shape: Cuneate. Apex shape: Convex. Margin type: Smooth.

Flowering.—Flowering interval: Late September to late April.

Fruit:

Fruit size.—Length: 57 mm. Width: 46 mm. Length/width ratio: 1.2.

Fruit hollow.—Length: 24 mm. Width: 5 mm. Length/width ratio: 4.7.

Shape.—Conical.

Difference in shape of terminal and other fruits.—None or very slight.

Fruit color.—RHS 45A (Vivid red).

Evenness of color.—Slightly uneven.

Glossiness.—Medium.

Evenness of surface.—Even or very slightly uneven.

4

Width of band without achenes.—Medium.

Position of achenes.—Level with surface.

Position of calyx attachment.—Level with fruit.

Attitude of sepals.—Outwards.

Diameter of calyx in relation to diameter of fruit.—Slightly larger.

Adherence of calyx.—Medium.

Firmness.—Firm.

Color of flesh (excluding core).—RHS 42C (Strong reddish orange).

Evenness and distribution of flesh color.—Moderately even.

Color of core.—RHS NN155B (White).

Juiciness.—Moderate.

Fruit weight.—27.7 g.

Soluble solids (in ° brix).—10.4.

Fruiting.—Harvest interval: Late October to late April.

Type of bearing: Fully remontant. Productivity: 32,498 kg to 65,087 kg of fruit per hectare per season from six-month-old plants when grown in Tapalpa, Mexico. Market use of fruit: Fresh market. Shipping and storage characteristics: Following harvest, on average more than 39% of fruit is marketable after 9 days in storage.

Resistance to abiotic stress, pests, and diseases:

Two-spotted spider mite (tetranychus urticae).—Moderately resistant.

Botrytis fruit rot (botrytis cinerea).—Moderately susceptible.

COMPARISON WITH PARENTAL AND REFERENCE VARIETIES

‘DrisStrawEightyNine’ differs from the female parent proprietary strawberry plant ‘88T205’ (unpatented) in that plants of ‘DrisStrawEightyNine’ have a higher yield than plants of ‘88T205’.

‘DrisStrawEightyNine’ differs from the male parent proprietary strawberry plant ‘107S 60’ (unpatented) in that fruit of ‘DrisStrawEightyNine’ is larger and less misshapen than fruit of ‘107S 60’.

‘DrisStrawEightyNine’ differs from the reference variety ‘DrisStrawSixtyEight’ (U.S. Plant Pat. No. 32,079) in that ‘DrisStrawEightyNine’ has a conical fruit shape, no difference or a very slight difference in shapes of terminal and other fruits, and a slightly larger diameter of calyx in relation to diameter of fruit, whereas ‘DrisStrawSixtyEight’ has a cordate fruit shape, a slight difference in shapes of terminal and other fruits, and a same size diameter of calyx in relation to diameter of fruit. In addition, ‘DrisStrawEightyNine’ has a fully remontant type of bearing, while ‘DrisStrawSixtyEight’ has a partially remontant type of bearing.

‘DrisStrawEightyNine’ differs from the reference variety ‘DrisStrawThirtySix’ (U.S. Plant Pat. No. 25,698), in that ‘DrisStrawEightyNine’ has medium plant vigor, an obtuse terminal leaflet shape of base, a position of inflorescence at the same level as foliage, and a medium adherence of calyx to fruit, whereas ‘DrisStrawThirtySix’ has very strong plant vigor, an acute terminal leaflet shape of base, a position of inflorescence above foliage, and a very strong adherence of calyx to fruit. In addition, ‘DrisStrawEightyNine’ has a fully remontant type of bearing, while ‘DrisStrawThirtySix’ has a not remontant type of bearing.

We claim:

- 1. A new and distinct variety of strawberry plant named ‘DrisStrawEightyNine’ as shown and described herein.

* * * * *



FIG. 1



FIG. 2



FIG. 3




FIG. 4




FIG. 5



FIG. 6