



US00PP34655P2

(12) **United States Plant Patent**
Bisschop et al.

(10) **Patent No.:** **US PP34,655 P2**
(45) **Date of Patent:** **Oct. 11, 2022**

(54) **RUDBECKIA PLANT NAMED ‘ET RDB 949’**

(50) Latin Name: *Echinacea purpurea* X *Rudbeckia hirta*

Varietal Denomination: **ET RDB 949**

(71) Applicants: **Elisabeth Bisschop**, Boijl (NL); **Bart Noordhuis**, Boijl (NL)

(72) Inventors: **Elisabeth Bisschop**, Boijl (NL); **Bart Noordhuis**, Boijl (NL)

(73) Assignee: **ETERNAL PLANT BOIJL B.V.**, Boijl (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/674,694**

(22) Filed: **Feb. 17, 2022**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/14 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./474; Plt./428**

(58) **Field of Classification Search**

USPC Plt./474, 428
CPC A01H 5/02; A01H 5/00; A01H 6/14
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Garden Center Marketing *Rudbeckia hirta* ‘Becky Orange’, retrieved on May 2, 2022 at <https://www.gardencentermarketing.com/plantName/Rudbeckia-spp-Becky-Orange>, 3 pp. (Year: 2022).*

* cited by examiner

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — C. Anne Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Rudbeckia* plant named ‘ET RDB 949’, characterized by its relatively compact, upright to somewhat outwardly spreading and mounded plant habit; dense and bushy growth habit; freely flowering habit; large long-lasting single-type inflorescences with orange yellow-colored ray florets positioned above the foliar plane on strong peduncles; and good postproduction and garden performance.

2 Drawing Sheets

1

Botanical designation: *Echinacea purpurea* X *Rudbeckia hirta*.

Cultivar denomination: ‘ET RDB 949’.

CROSS-REFERENCED TO A RELATED APPLICATION & STATEMENT REGARDING PRIOR DISCLOSURES BY INVENTORS/APPLICANTS:

This application claims priority to a European Community Plant Breeders’ Rights application filed on Oct. 1, 2021, application number 2021/2439. There have been no offers for sale anywhere in the world prior to the effective filing date of this Application and no accessibility to one of ordinary skill in the art could have been derived from the printed Plant Breeder’s Rights documents.

The Inventors/Applicants assert that no publications nor advertisements relating to sales, offers for sale or public distribution occurred more than one year prior to the effective filing date of this application. Any information about the claimed plant would have been obtained from a direct or indirect disclosure from the Inventors/Applicants. Inventors/Applicants claim a prior art exception under 35 U.S.C. 102(b)(1) for disclosure and/or sales prior to the filing date but less than one year prior to the effective filing date.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Rudbeckia* plant, botanically known as *Echinacea purpurea* X *Rudbeckia hirta* and hereinafter referred to by the name ‘ET RDB 949’.

2

The new *Rudbeckia* plant is a product of a planned breeding program conducted by the Inventors in Boijl, The Netherlands. The objective of the breeding program is to create new compact and freely branching interspecific *Rudbeckia* plants with large long-lasting and attractive inflorescences on strong peduncles.

The new *Rudbeckia* plant originated from a cross-pollination made by the Inventors in Boijl, The Netherlands in August, 2017 of a proprietary selection of *Echinacea purpurea* identified as code designation 1950, not patented, as the female, or seed parent with a proprietary selection of *Rudbeckia hirta* identified as code designation RB 2503, not patented, as the male, or pollen, parent. The new *Rudbeckia* plant was discovered and selected by the Inventors as a single flowering plant from within the progeny of the stated cross-pollination in a controlled environment in Boijl, The Netherlands in July, 2018.

Asexual reproduction of the new *Rudbeckia* by in vitro meristem culture in a controlled greenhouse environment in Boijl, The Netherlands since August, 2018 has shown that the unique features of this new *Rudbeckia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Rudbeckia* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'ET RDB 949'. These characteristics in combination distinguish 'ET RDB 949' as a new and distinct *Rudbeckia* plant:

1. Relatively compact, upright to somewhat outwardly spreading and mounded plant habit.
2. Dense and bushy growth habit.
3. Freely flowering habit.
4. Large long-lasting single-type inflorescences with orange yellow-colored ray florets positioned above the foliar plane on strong peduncles.
5. Good postproduction and garden performance.

Plants of the new *Rudbeckia* differ from plants of the female parent selection in the following characteristics:

1. Plants of the new *Rudbeckia* are shorter than plants of the female parent selection.
2. Inflorescences of plants of the new *Rudbeckia* are larger and more uniformly mounding than inflorescences of plants of the female parent selection.

Plants of the new *Rudbeckia* differ from plants of the male parent selection in the following characteristics:

1. Plants of the new *Rudbeckia* are more compact than plants of the male parent selection.
2. Plants of the new *Rudbeckia* have smaller leaves than plants of the male parent selection.
3. Plants of the new *Rudbeckia* have smaller inflorescences than plants of the male parent selection.

Plants of the new *Rudbeckia* can be compared to *Rudbeckia hirta* X *Rudbeckia hybrida* 'ET RDB 18-30', disclosed in U.S. Plant Pat. No. 32,891. Plants of the new *Rudbeckia* differ primarily from plants of 'ET RDB 18-30' in the following characteristics:

1. Plants of the new *Rudbeckia* are slightly taller and narrower than plants of 'ET RDB 18-30'.
2. Plants of the new *Rudbeckia* have fewer and larger inflorescences per plant than plants of 'ET RDB 18-30'.
3. Inflorescences of plants of the new *Rudbeckia* have more ray florets than inflorescences of plants of 'ET RDB 18-30'.
4. Plants of the new *Rudbeckia* and 'ET RDB 18-30' differ in ray floret color as plants of the new *Rudbeckia* have orange yellow-colored ray florets whereas plants of 'ET RDB 18-30' have yellow orange and dark red bi-colored ray florets.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Rudbeckia* showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Rudbeckia* plant.

The photograph on the first sheet (FIG. 1) is a side perspective view of a typical flowering plant of 'ET RDB 949' grown in a container.

The photograph on the second sheet (FIG. 2) is a close-up view of typical inflorescences of 'ET RDB 949'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in 2-liter containers during the summer in an outdoor nursery in Boijl, The Netherlands and under cultural conditions

typical of commercial *Rudbeckia* production. During the production of the plants, day temperatures ranged from about 16° C. to about 24° C. and night temperatures ranged from about 8° C. to about 15° C. Plants were one year old when the photographs and the detailed description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Echinacea purpurea* X *Rudbeckia hirta* 'ET RDB 949'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Echinacea purpurea* identified as code designation 1950, not patented.

Male, or pollen, parent.—Proprietary selection of *Rudbeckia hirta* identified as code designation RB 2503, not patented.

Propagation:

Type.—By in vitro meristem culture.

Time to initiate roots, summer.—About seven to eight days at substrate temperatures about 15° C. and ambient temperatures about 20° C.

Time to produce a rooted young plants, summer.—About four weeks at substrate temperatures about 15° C. and ambient temperatures about 20° C.

Root description.—Medium in thickness; fleshy; color, typically close to 162C, actual color of the roots is dependent on substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant and growth habit.—Herbaceous perennial; compact, upright to somewhat outwardly spreading and mounded plant habit; dense and bushy appearance; vigorous growth habit and moderate growth rate.

Plant height.—About 65 cm to 70 cm.

Plant width.—About 50 cm to 55 cm.

Lateral branches (peduncles).—Length: About 35 cm to 50 cm. Diameter: About 6 mm to 8 mm. Internode length: About 4 cm to 7 cm. Angle: Upright to somewhat outwardly spreading. Strength: Strong. Texture: Pubescent. Color: Close to 143A sparsely covered with spots, close to 187A.

Leaf description:

Arrangement.—Alternate, simple; sessile.

Length.—About 10 cm to 17 cm.

Width.—About 5 cm to 7 cm.

Shape.—Oblanceolate to obovate.

Apex.—Acute to acuminate.

Base.—Attenuate to acute.

Margin.—Dentate; undulate.

Texture, upper and lower surfaces.—Pubescent; rough.

Venation pattern.—Pinnate.

Color.—Developing and fully developed leaves, upper surface: Close to 137A; venation, close to 145A. Developing and fully expanded leaves, lower surface: Close to 138A; venation, close to 145A.

Inflorescence description:

Type and arrangement.—Single-type inflorescence form with elliptic to oblanceolate-shaped ray florets and tubular disc florets; inflorescences borne on terminal and axillary peduncles above and beyond the foliar plane on strong peduncles; inflorescences

face upright to slightly outwardly; ray and disc florets arranged acropetally on a capitulum.

Fragrance.—None detected.

Flowering season.—Plants begin flowering about ten weeks after planting; long flowering period, plants flower continuously from mid-July until the end of October in The Netherlands.

Inflorescence longevity.—Good postproduction longevity with inflorescences lasting about eight to ten weeks on the plant; inflorescences persistent.

Quantity of inflorescences.—Freely flowering habit, typically more than 20 inflorescences per plant.

Inflorescence buds.—Height: About 1 cm to 2 cm. Diameter: About 1.5 cm. Shape: Round to ovoid. Color: Close to 143C.

Inflorescences.—Diameter: About 14 cm. Depth (height): About 3 cm to 5 cm. Diameter of disc: About 2.5 cm to 3 cm. Receptacle height: About 1 cm to 2 cm. Receptacle diameter: About 2 cm to 2.5 cm. Receptacle color: Close to 144A.

Ray florets.—Number of ray florets per inflorescence: About 20 in about one or two whorls. Length: About 5 cm to 6 cm. Width: About 2 cm to 3 cm. Shape: Elliptic to oblanceolate. Apex: Emarginate. Base: Attenuate. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Rough, glabrous. Color: When opening, upper surface: Close to 17A; towards the base, slightly darker than 17A. When opening, lower surface: More grey than 15B. Fully opened, upper surface: Close to 17A; towards the base, slightly darker than 17A; venation, similar to lamina colors; color does not change with subsequent development. Fully opened, lower surface: Close to 17A; venation, close to 17A; color does not change with subsequent development.

Disc florets.—Arrangement: Numerous disc florets massed at center of receptacle. Length: About 1 cm to 3 cm. Width: About 2 mm to 3 mm. Shape:

Tubular, elongated. Apex: Obtuse. Color, when opening and fully opened: Apex: Close to 200A to 200B. Mid-section: Whiter than 76D. Base: Close to 155C.

Phyllaries.—Number of phyllaries per inflorescence: About 15 to 20 arranged in about two to three whorls. Length: About 2 cm to 4 cm. Width: About 5 mm to 8 mm. Shape: Linear. Apex: Acute to obtuse. Base: Fused to receptacle. Margin: Entire; slightly undulate. Texture, upper and lower surfaces: Rough, pubescent. Color, upper surface: Close to 137A. Color, lower surface: Close to 138A.

Reproductive organs.—Androecium (present on ray and disc florets): Quantity per floret: Numerous. Filament length: About 2 mm to 5 mm. Filament color: Close to 200A to 200B. Anther shape: Round. Anther length: About 1 mm to 2 mm. Anther color: Close to 200A to 200B. Pollen amount: Abundant. Pollen color: Close to 21A. Gynoecium (present only on disc florets): Pistil length: About 2 mm to 3 mm. Stigma shape: Two-parted. Stigma color: Close to 200B. Style length: About 1 mm to 2 mm. Style color: Close to 200B.

Seeds and fruits.—To date, seed and fruit production has not been observed on plants of the new *Rudbeckia*.

Pathogen & pest resistance: To date, plants of the new *Rudbeckia* have not been observed to be resistant to pathogens and pests common to *Rudbeckia* plants.

Garden performance: Plants of the new *Rudbeckia* have been observed to have good garden performance and to tolerate wind, rain and temperatures from about -20° C. to about 40° C.

It is claimed:

1. A new and distinct *Rudbeckia* plant named 'ET RDB 949' as illustrated and described.

* * * * *



FIG. 1



FIG. 2