



US00PP35179P2

(12) **United States Plant Patent**
Blaker

(10) **Patent No.:** **US PP35,179 P2**
(45) **Date of Patent:** **May 23, 2023**

(54) **STRAWBERRY PLANT NAMED**
‘SB_13_164-042’

(50) Latin Name: *Fragaria ananassa*
Varietal Denomination: **SB_13_164-042**

(71) Applicant: **Strawberry Sciences, LLC,**
Watsonville, CA (US)

(72) Inventor: **Kendra M. Blaker,** Archer, FL (US)

(73) Assignee: **Strawberry Sciences, LLC,**
Watsonville, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/880,955**

(22) Filed: **Aug. 4, 2022**

(51) **Int. Cl.**
A01H 5/12 (2018.01)
A01H 6/74 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./208**

CPC *A01H 6/7409* (2018.05)

(58) **Field of Classification Search**
USPC Plt./208
CPC *A01H 6/7409*
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP20,394	P3	10/2009	Ackerman et al.
PP23,042	P3	9/2012	Chandler
PP30,564	P3	6/2019	Whitaker

Primary Examiner — Anne Marie Grunberg
(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP

(57) **ABSTRACT**

This invention relates to a new and distinct variety of strawberry plant named ‘SB_13_164-042’. This new strawberry plant named ‘SB_13_164-042’ is primarily adapted to the growing conditions of West Central Florida, and is primarily characterized by its achenes typically set even with the surface of the fruit, vigorous plant habit, high marketable yield, early time of first flower and fruit, and very large berry size.

4 Drawing Sheets

1

Latin name of the genus and species of the plant claimed:
Fragaria ananassa.
Variety denomination: ‘SB_13_164-042’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct strawberry variety named ‘SB_13_164-042’. This new variety is a result of a controlled cross made in 2013 in an ongoing breeding program between strawberry variety designated ‘FL 05-107’ (U.S. Plant Pat. No. 23,042) as the seed (female) parent, and strawberry variety designated ‘Valor’ (U.S. Plant Pat. No. 20,394) as the pollen (male) parent. The variety is botanically known as *Fragaria ananassa*.

The seedling resulting from the aforementioned cross was selected from a controlled breeding plot in Hillsborough County, Fla. in the fall/winter of 2014-2015. After its selection, the new variety was asexually propagated by stolons in both Siskiyou County, Calif. and San Joaquin County, Calif. The new variety was extensively tested over the next several years in fruiting fields in Hillsborough County, Fla. This propagation has demonstrated that the combination of traits disclosed herein as characterizing the new variety are fixed and remain true-to-type through successive generations of asexual reproduction.

BRIEF SUMMARY OF THE INVENTION

‘SB_13_164-042’ is primarily adapted to the climate and growing conditions of West Central Florida. The subtropical climate of West Central Florida provides the day length and

2

moderate temperatures needed to produce an early yielding, vigorous plant and maintain fruit quality during the fall and winter production months.

The following traits have been repeatedly observed and are determined to be unique characteristics of ‘SB_13_164-042’, which in combination distinguish this strawberry plant as a new and distinct variety:

1. Achenes typically set even with the surface of the fruit;
2. Vigorous plant habit;
3. High marketable yield;
4. Early time of first flower and fruit; and
5. Very Large berry size.

‘Florida Brilliance’ (U.S. Plant Pat. No. 30,564) is currently the dominant strawberry variety in Hillsborough County, Fla. The fruits of ‘SB_13_164-042’ are similar in color, flavor, and firmness to ‘Florida Brilliance’, but the achenes of ‘Florida Brilliance’ are slightly more sunken than those of ‘SB_13_164-042’ and its plant architecture is more open. In side-by-side comparisons from the 2021-2022 season (Nov. 23, 2021 to Feb. 25, 2022) ‘SB_13_164-042’ compares with ‘Florida Brilliance’ (U.S. Plant Pat. No. 30,564) in the following combination of characteristics as described in Table 1.

TABLE 1		
Characteristic	‘SB_13_164-042’	‘Florida Brilliance’ (U.S. Plant Pat. No. 30,564)
November marketable yield (gm/plt)	73	77
Season marketable yield (gm/plt)	517	818

TABLE 1-continued

Characteristic	'SB_13_164-042'	'Florida Brilliance' (U.S. Plant Pat. No. 30,564)
November average berry size (gm)	23.0	27.4
Season average berry size (gm)	31.7	27.7
Average runners/plant	1.1	5.5

For identification, a series of molecular markers have been determined for this new variety.

'SB_13_164-042' compares with its parents, 'FL 05-107' and 'Valor' by the following combination of characteristics as described in Tables 2 and 3.

TABLE 2

Characteristic	'SB_13_164-042'	'FL 05-107' (U.S. Plant Pat. No. 23,042)
Fruit: size	Very Large	Medium
Fruit: marketable yield	High	High
Fruit: firmness	Very Firm	Very Firm
Fruit: seed position	Even with surface	Slightly below to even with surface

TABLE 3

Characteristic	'SB_13_164-042'	'Valor' (U.S. Plant Pat. No. 20,394)
Fruit: size	Very Large	Medium to small
Fruit: marketable yield	High	Medium
Fruit: early yield	Very High	Medium
Fruit: seed position	Even with surface	Slightly above to even with surface

BRIEF DESCRIPTIONS OF THE PHOTOGRAPHS

The accompanying color photographs illustrate the overall appearance of typical specimens of the new strawberry variety 'SB_13_164-042' at various stages of development, as true as it is reasonably possible with color reproductions of this type. Color in the photographs may differ slightly from the color value cited in the botanical descriptions which accurately describe the color of 'SB_13_164-042'. The depicted plant and plant parts of the new strawberry variety 'SB_13_164-042' are approximately four months old. The photographs were taken in Hillsborough County, Fla.

FIG. 1 shows typical fruiting field characteristics of 'SB_13_164-042', taken in the month of February 2022;

FIG. 2 shows a close-up view of a typical plant of 'SB_13_164-042', taken in the month of February 2022;

FIG. 3 shows typical mature and immature field fruit of 'SB_13_164-042', taken in the month of February 2022; and

FIG. 4 shows typical internal and external mature fruit characteristics of 'SB_13_164-042', taken in the month of February 2022.

DETAILED BOTANICAL DESCRIPTION

The new variety 'SB_13_164-042' has not been observed under all possible environmental conditions. The characteristics of the new variety 'SB_13_164-042' may vary in

detail, depending upon variations in environmental factors, including weather (temperature, humidity and light intensity), day length, soil type and location. In addition, the characteristics of any parental variety or comparison variety included in Table 1 of the present invention may vary in detail, depending upon variations in environmental factors, including weather (temperature, humidity and light intensity), day length, soil type and location.

The aforementioned photographs, together with the following description of the new variety 'SB_13_164-042', unless otherwise noted, are based on observations taken during the 2021-2022 growing season in Hillsborough County, Fla. These measurements and ratings were taken from plants of 'SB_13_164-042' dug from a high-elevation nursery located in Siskiyou County, Calif. during mid-September 2021 and planted approximately four to five days later in Hillsborough County, Fla. The approximate age of the observed plants is four months. Yield observations including average weight and marketable yield, along with fruit quality characteristics including soluble solids, were measured during the 2021-2022 growing season. Flower measurements and characteristics are from secondary flowers unless otherwise noted. Fruit characteristics and measurements are from secondary fruit, unless otherwise noted.

Where noted, color terminology follows The Royal Horticultural Society Colour Chart, London (2015).

The following characteristics describe fruit, plant, stolon, foliage, fruiting truss, flower, reproductive organs and pest and disease characteristics of the new strawberry 'SB_13_164-042'.

Fruit characteristics:

Color of mature fruit.—RHS 185A (greyed-purple group).

Color of internal flesh (excluding core).—RHS 36D (red group).

Color of core.—RHS 45A (red group).

Average length (cm).—4.8.

Average width (cm).—4.2.

Size.—Very Large.

Average length/width ratio.—1.1 (ranges from as long as broad to slightly longer than broad).

Average calyx diameter (cm).—5.5.

Season average weight (gm).—31.7.

Achene color, shaded side.—RHS 178B (greyed-red group).

Achene color, sun-exposed side.—RHS 160A (greyed-yellow group).

Average achene weight (mg).—<2.5.

Average achenes per berry.—399.

Average achene length (mm).—1.8.

Average achene width (mm).—1.0.

Season marketable yield (gm/plant).—517.

Predominant shape.—Conical.

Difference in shape between primary and secondary fruit.—Ranges from moderate to large.

Band without achenes.—Very narrow.

Evenness of surface.—Even.

Evenness of color.—Even.

Glossiness.—Strong.

Insertion of achenes.—Even to surface.

Position of calyx attachment.—Inserted.

Attitude of sepals.—Outward.

Size of calyx in relation to fruit diameter.—Larger.

Adherence of calyx (when fully ripe).—Strong.

Firmness of flesh (gf).—425.

Distribution of red color of the flesh.—Heavily marginal and central.

Hollow center expression.—Mildly present.

Average cavity length (mm).—16.7.

Average cavity width (mm).—5.2.

Soluble solids (% brix).—6.2.

Time of first flowering.—Early (early to mid-October in Hillsborough County, Fla.).

Flowering season.—October-February.

Time of first fruit.—Early (mid-November in Hillsborough County, Fla.).

Fruiting season.—November-March.

Post-harvest fruit longevity.—9-11 days if stored according to industry standards.

Type of bearing.—Not remontant.

Plant characteristics:

Average height (cm) —29.3.

Average spread (cm).—42.2.

Size.—Large.

Habit.—Upright.

Density.—High.

Vigor.—Very High.

Stolon characteristics:

Color.—RHS N144D (yellow-green group).

Anthocyanin coloration.—RHS 182C (greyed-red group).

Anthocyanin intensity.—Medium.

Pubescence.—Dense.

Attitude of hairs.—Slightly Outward.

Average quantity in nursery (per square foot).—10.3 (medium).

Average diameter at the bract (mm).—2.7 (medium).

Average length (cm).—30.6.

Terminal leaflet characteristics:

Color of upper surface.—RHS 139A (green group).

Color of underside.—RHS 147B (yellow-green group).

Average length (cm).—7.3.

Average width (cm).—7.6.

Average area terminal (cm²).—55.5.

Average length/width ratio.—0.96 (slightly more broad than long).

Shape of base.—Obtuse.

Margins (shape of teeth).—Serrate to crenate.

Average serrations per leaf.—21.4.

Foliage characteristics:

Color of upper surface.—RHS 139A (green group).

Color of underside.—RHS 147B (yellow-green group).

Number of leaflets.—3.

Leaf size.—Large.

Average length (cm).—12.0.

Average width (cm).—14.0.

Average area foliage (cm²).—168.

Shape in cross section.—Concave.

Texture/interveinal blistering.—Light.

Leaf glossiness.—Mild.

Leaf variegation.—Absent.

Venation pattern.—Pinnate.

Apex descriptor.—Obtuse.

Petiole characteristics:

Petiole color.—RHS 144C (yellow-green group).

Average length (cm).—15.2.

Average diameter (mm).—3.9.

Petiolule color.—RHS 144B (yellow-green group).

Petiolule average length (mm).—16.2.

Average petiolule diameter (mm).—2.1.

Attitude of hairs.—Strongly outward.

Texture.—Moderate.

Frequency of bract leaflets.—Ranges from 0 to 2 (70% occurrence).

Size of bract leaflets.—Ranges from small to large.

Pubescence.—Light.

Stipule characteristics:

Color.—RHS 145A (yellow-green group).

Anthocyanin coloration.—RHS 182A (greyed-red group).

Anthocyanin intensity.—Mild.

Average length (mm).—39.8.

Average width (mm).—11.9.

Base descriptor.—Truncate.

Apex descriptor.—Obtuse.

Shape.—Triangular.

Margin.—Smooth.

Texture.—Smooth.

Fruiting truss characteristics:

Anthocyanin coloration.—N/A.

Anthocyanin intensity.—N/A.

Pubescence.—Moderate.

Attitude at first pick.—Prostrate.

Position relative to foliage.—Ranges from level with to below.

Flower quantity (average per plant season long).—29.9 (medium).

Average fruits per truss.—10.4.

Pedicel attitude of hairs.—Strongly outward.

Average pedicel length (cm).—16.3.

Average pedicel diameter (mm).—2.4.

Pedicel texture.—Moderate.

Pedicel color.—RHS N144B (yellow-green group).

Average peduncle length (cm).—4.0.

Average peduncle diameter (mm).—4.0.

Peduncle texture.—Smooth.

Peduncle color.—145C.

Flower characteristics:

Flower bud shape.—Pyriform.

Average flower bud length (mm).—16.5.

Average flower bud diameter (mm).—7.6.

Flower bud color.—RHS 144A (yellow-green group).

Flower depth (mm).—5.4.

Corolla (flower) average diameter (mm).—29.3 (ranges from medium to large).

Upper petal color.—RHS NN155D (white group).

Lower petal color.—RHS NN155D (white group).

Petal shape.—Orbicular.

Petal apex descriptor.—Rounded.

Petal margin.—Smooth.

Petal base.—Decurrent.

Petal texture.—Smooth.

Petal average length (mm).—11.3.

Petal average width (mm).—11.8.

Petal average length/width ratio.—0.96 (slightly longer than broad).

Average petals per flower.—5.7.

Relative position of petals (flowers with 5 or 6 petals).—Overlapping.

Upper sepal color.—RHS 137A (green group).

Lower sepal color.—RHS 138A (green group).

Sepal shape.—Cuneate.

Sepal apex descriptor.—Obtuse.

Sepal margin.—Serrate.

Sepal texture.—Smooth.

Sepal average length (mm).—14.6.
Sepal average width (mm).—6.9.
Sepal average length/width ratio.—2.1.
Average sepals per flower.—12.1.
Calyx average diameter (mm).—37.2.
Size of calyx relative to corolla.—Larger.
Size of inner calyx relative to outer calyx.—Smaller.
 Reproductive organs:
Receptacle color.—RHS N144A (yellow-green group).
Pollen color.—RHS 17A (yellow-orange group).
Stamen.—Present.
Average filament length (mm).—1.8.
Filament color.—RHS 149C.
Average anther length (mm).—1.3.
Anther shape.—Ovoid.
Anther color.—RHS 17A (yellow-orange color).
Average pistils per flower.—399.
Pistil length (mm).—1 to 2.
Style length (mm).—0.5 to 1 mm.

Style color.—RHS 154D.
Stigma diameter (mm).—<0.1.
Stigma shape.—Simple.
Ovary color.—RHS N144A (yellow-green group).
Pollen amount.—Abundant.
 5 Disease and pest reactions:
Colletotrichum crown rot (colletotrichum gloeosporioides).—Susceptible; (*Colletotrichum acutatem*) — Susceptible.
 10 *Fusarium wilt (fusarium oxysporum).*—Resistant.
Pestalotia leaf spot and fruit rot (neopestalotiopsis sp.).—Moderately Susceptible.
Phytophthora crown rot (phytophthora cactorum).— Moderately Susceptible.
 15 I claim:
 1. A new and distinct strawberry plant named ‘SB_13_164-042’, as herein described and illustrated by the characteristics set forth above.

* * * * *

FIG. 1



FIG. 2



FIG. 3




FIG. 4

