

IDC352 Jan-April 2014, Instructor : Professor Arvind

Name : Atul Singh Arora

Roll Number : MS11003

1 Title : Why a Quantum Theory of Gravity

Speaker : Prof. G. Date (IMSc)

Date : March 26, 2014

Venue : LH-3, LHC

Report :

The speaker opened with 'Quantum Gravity is the fiery marriage of General Relativity and Quantum Mechanics'. The pursuit he said is not purely hypothetical, nor a problem of unexplained data. Wheeler's description of the final state of a star is amongst one of the reasons why such a theory would be necessary.*

$$R_{\mu\nu}(g) - \frac{1}{2}R(g)g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}$$

There've been two revolutions in physics; Relativity and Quantum Mechanics. In General Relativity the notion of causality gets dynamical and metrical properties can also change. The combination of special relativity and Quantum Mechanics is Quantum Field Theory. QFT is consistent with fixed causality defined by Minkowski space-time, but assumes that causal structures can be defined at arbitrarily small lengths.

Next the speaker talked about the basic assumptions of GR.

1. space-time is defined by a 4d manifold and its metric $g_{\mu\nu}$ (-ve determinant).
2. Stress tensor, $T_{\mu\nu}$, which describes the matter distribution.

The metric imposes a causal structure on time like events in their local neighbourhood.

Remarks: Given a metric, at any point, you can define a set of tangent vectors. These can be classified as timelike, spacelike or null. Corresponding geodesics determine which events can influence each other causally.

Einstein's equation is only a local PDE and determines solution space-times only locally. Extended solutions are not guaranteed to be causal without imposing extra conditions. Then the speaker talked about predictability, that is events can be divided into cause and effect.

Next he talked about time orientability, strong causality and stable causality. He stated that causality is not automatic, so time-orientable space times are a subset of solutions of einstein's equation. An example is given in the next section.

Strong Causality: We can impose strong causality else its possible to have future curves that come arbitrarily close to the existing events.

Stable Causality: Imposing stable causality on the other hand would mean that spacelike and timelike do not interchange.

Global Hyperbolicity: space-times which admit spacelike hypersurface whose domain of dependence is the entire space-time are globally hyperbolic.

Thus, global hyperbolicity automatically ensures causal stability.

To have a space-time with predictability, we can construct globally hyperbolic space-time (take a spacelike hyper surface, define domain of dependence to be the full space-time, it abruptly ends. You extend it, how far? maximally possible). However, a predictable space-time should be an inextendable/globally hyperbolic solution of einstein's equation.

The speaker then went on to describe a slightly technical topic, 'Property of Globally Hyperbolic Space Times'. Consider a space $C(\Sigma, q)$ of causal curves, emanating from a hypersurface Σ and passing through a point q to its future. Its a theorem that a curve of maximum proper time is a geodesic without any 'conjugate point' between Σ and q .

Next the speaker digressed to talk about a 'Bundle of Time like Geodesics'. Imagine a cloud of particles freely 'falling' in a given space time. These will be described by a bungle of time-like geodesics. Their cross section in general can undergo 1. Shearing, 2. Twist or 3. Expansion.

'Raychaudhuri Equations and Conjugate Points' was discussed next. This I couldn't understand much however what is known that there arises a contradiction related to the existence of a singularity which is resolved by concluding that the geodesic must be incomplete, viz. the observer's world line must end abruptly. The details have been skipped.

Blackholes and Thermodynamics were discussed next. General relativity contains another class of intriguing solutions - Black holes - space-times which have event horizons. A stationary black hole can be described in terms of parameters like mass (M), angular momentum (J) and charge (Q). The event horizon is similarly described by the (A) area, (Ω) angular velocity and (Po-
tential ϕ electrostatic potential). They have a surface gravity (K) which is a

constant over the horizon.

If these are disturbed by any astrophysical process, they continue to remain black hole with changed values of their parameters, such that

$$\delta M = \frac{K}{8\pi} \delta A + \Omega \delta J + \phi \delta Q, \delta A \geq 0$$

This is a striking similarity to classical thermodynamics and there's no reason why blackholes should obey this. But continuing the analogy suggests that surface gravity \sim temperature and area \sim entropy. So by this analogy we expect that being a hot body, black-holes should radiate but they don't let anything escape.

Hawking pointed out that this is true only classically. Because of quantum fluctuations present in the horizon of the black hole, it has a temperature and entropy given by

$$T = \frac{kGh}{2\pi}, S = \frac{A}{4Gh}$$

The next part was rather interesting. Since we understand the origin of entropy from statistics of a finite number of entities, it seems to suggest that the horizons must have some atomic structure, and they must be purely geometric objects, viz. the 'atoms' must be atoms of geometry. Maybe space-time geometry is discrete.

This hints us to the conclusion that perhaps continuum geometry is inadequate near high curvature regions and that geometry may be discrete. This begins the framework of quantum. So what would be the goal of such a framework? It should reproduce the classical space-time approximately at the bare minimum. Further it should at least resolve the curvature singularities and provide an explanation of the blackhole entropy.

So where's the problem, well GR has a property of general covariance which makes it a gauge theory and results in loss of determinism. Further, if space-time is quantized, then 'notion of 'dynamics' is hard to define, since time evolution needs to be interpreted differently. Besides, the regimes where quantum rescue is sought, involves highly dynamical, strong curvatures, where perturbative tools are questionable.

The speaker ended the talk with discussing the two types of approaches taken to solve the problem of quantization. The first is based on discretizing points of geometry. The second involves promoting space to be an operator that acts on some state.

This I'll remember

It is possible to construct solutions to Einstein's equations such that locally they follow all the properties we expect but have peculiar global behaviour. Imagine making a light cone on a piece of paper and then folding it to make the future and past light cone match. Locally, everything's consistent, but globally, you're going to the past from the future!

Acknowledgements

I thank Ms. Prashansa Gupta (MS11021) for sharing her notes so that atleast our combined notes made sense.

2 Title : The Dance Language of the Bees

Speaker : Prof. Raghvendra Gadagkar (IISc)

Date : March 27, 2014 Venue : LH-5, LHC

Report :

Introduction

Apis florea is the species discussed in the talk. It is found only in Asia, and is open nesting (thus harder to domesticate). They are interesting as their social behaviour is not as sophisticated (compared to the other four major species). They communicate by transferring saliva and dancing, i.e. subject of the talk.

About these bees

Within these bees, there are drones (the males), worker (essentially sterile) and one queen (that lays eggs).

The said bees dance only on horizontal surfaces unlike the other species. There are two types of dances known. First is called a round dance, in which the bee makes a circular motion. The other's called a waggle dance in which the bee transcribes an 8 shape with vigorous vibrations half way.

Communication and Dance (the real deal)

Earlier it was thought that wagging related to pollen being found and round to nectar. Later experiments confirmed that the wagging dance corresponded to the food being farther than roughly 100 m and the round to less than that.

Experiments have confirmed that the waggle dance conveys information about the direction and distance of the resource found. They use the sun's

azimuthal as reference (as it is globally stable, unlike the surroundings) even though it changes its location periodically. The bees apparently correct for this. Further they even update their map of the surrounding everyday and use that to locate the sun.

In the waggle dance, during the middle part of 8, the direction in which the bee moves contains the direction information. Also during this part, the tempo of vibration codes for distance.

Although its not known how they find the angle, the distance is measured not using the energy requirement as was previously thought, but using optical flow.

Sophistication

These bees have evolved to use fallback mechanisms for defining the angle.

1. Sun is used when its bright enough
2. Polarization of light on the sky is used if the sky is overcast
3. Local landmarks can also be used. They learn it each day to compensate for their change with time.
4. Siesta; when the sun is overhead, they take time off

Experiments

Various experiments were done to test color vision, co-relation between sugar/nectar and waggle/round dance, co-relation between wiggle and flight direction (Fan like experiment), that optical information is used to find direction, use of Sun as reference (The lamp shift experiment | miscommunication) and if dancing is sufficient to communicate information about resources (The robot bee experiment).

3 Title : The Climate Energy Nexus

Speaker : Dr. SK Tandon (IIT Kanpur))

Date : April 16, 2014 Venue : LH-5, LHC

Report :

The talk was started by display of the greenhouse gasses data spanning the last 100 years. Some key estimates of CO₂ concentrations are as follows

- 180 ppm | glacial (cold) period
- 300 ppm | interglacial warm period

- 280 ppm | 18th century, Industrial Revolution
- 380 ppm | transportation revolution leading to urbanization

The speaker then discussed some of the techniques used for arriving at such conclusions

- Continental Ice sheets (also known as Climate Archive)
 - Cryosphere's annual layers: These are counted to establish the 'age'
 - Molten Ice: This is useful for identifying liquids present at the time
 - Ice Cracks: They are the molten ice for gas
 - Scintering: Scaling of air bubbles is used
- Annual Rings
- Radioactive
- Age Modelling

Next an attempt was made to relate CO₂ levels and human CO₂ emissions. The speaker claimed that a Nexus of sorts is formed with the following components:

Water, food, population control, health, energy, climate change, environment sanity, knowledge equity, sustainability and terrorism.

Next the speaker described the earth system as composed of

- Eco-sphere
- Human Factors
 - Anthroposphere: Aggregate of human activities
 - Metaphysical: Its a global subject and includes adoption of say international protocols

Coming to the most important point of the talk, the speaker justified how 0.6 degrees of change in temperature is unaccounted for, despite the removal of all known possible natural causes, including solar heating (primarily, believed to contribute ~0.2 degrees), orbit cooling and millenium warming. Thus by elimination, the only cause of the observed effects is anthropogenic

The speaker went on to say how the climate models in the past relied only on factors such as atmosphere, land and ocean while today additionally, the following are the base standard

1. Aerosols

2. Carbon Cycles
3. Dynamic Vegetation
4. Atmospheric Chemistry
5. Land Ice

He concluded that therefore today we are justified at having more confidence on these.

In conclusion the speaker pointed out that there's an urgent demand for affordable, reliable energy (it is he claimed pivotal to all our problems). He ended on optimistic terms, discussing two examples of possible solutions.

- Carbon Capture (involves capturing, liquefying, transporting and releasing the gas) [successful examples are known, although not precisely for environmental reasons]
- Shail Oil (the extraction technology is still in its infancy but the reserves are promising)

3.1 This I'll remember

There's something called shail oil, which will last us a very long time, but we haven't the technology to extract it yet.

3.2 Acknowledgements

I was unable to make proper notes for this seminar and Ms. Prashansa Gupta (MS11021) was kind enough to share hers.

4 Title : The Geometry of Physics

Speaker : Mr. Rahul Chajwa (Phy Maj, IISERM)

Date : April 16, 2014 **Venue :** LH-3, LHC

Report :

The speaker in essence talked about how one would characterize various physical systems using geometry. He started by asking the question, how one would characterize surfaces, after showing with a series of figures. One way of such characterization he said was counting the number of holes. He then showed how various surfaces like a cup, purse etc. may deformed to a surface having clearly one or two (or even no) holes. He went on to explain why such a characterization, though intuitive is not always easy. He talked about Euler's

polyhedron formula and the Gauss-Bonnet Theorem. He attempted to show some generalizations to higher dimensions, including the concept of derivatives. Specifically, he talked about Liouville's Theorem and discussed in some detail the tangent space and its dual, co-tangent space. The fascinating part of the talk was related to the double pendulum which we know is chaotic. He explicitly showed an animation proving his point. He then asked us what geometry we could associate with the co-ordinate phase space of the setup. The answer was a torus. He then stated a result which follows from the geometric analysis that the Hamilton's equations result in constraining the movement along geodesics of the torus geometry. This has a profound implication, the resulting motion will be periodic since geodesics are closed. This is almost impossible to see without the geometric considerations since we don't even have an analytic solution of the motion. Finally he demonstrated Euler's disc and talked about the infinite frequency before collapsing and discussed why this singularity is not a consequence of Hamiltonian evolution but instead that of dissipation. He concluded the discussion by attempting to convince the audience that we can't think of logic without invoking some sort of object (geometry) in our heads.

The speaker received criticism for not acknowledging the sources referred to, and for shoddy board work from the instructor.

4.1 This I'll remember

A double pendulum's (co-ordinate) phase space can be thought of as the points on the surface of a torus.

4.2 Acknowledgements

I thank Mr. Vivek Sagar (MS11017) for reminding me about the seminar, else it would've been missed by me.

5 Title : Magnetic Refrigeration

Speaker : Ms. Ayushi Singhania (Int PhD, IISER-M)

Date : April 23, 2014 **Venue :** LH-3, LHC

Report :

The speaker started with outlining her talk; introduction, history, magneto caloric effect, thermodynamics, advantages/disadvantages. She explained that the technology is based in magneto caloric effects (MCE). This was discovered by P. Weiss and it was suggested by P. Debye and W. Giaque. The MCE is based in changing the temperature of specific materials, by exposing it to

changing magnetic fields. The phenomenon hinges on the idea that at a given temperature, the magnetic field causes alignment and when it is removed, the disorientation leads to dropping of the temperature (assuming there's no heat exchange). This has similarities with the Curie temperature which has electric fields and ferroelectric domains as analogues to their magnetic counterparts.

The speaker motivated the functioning further by explain how thermal and magnetic entropy together function to drop the temperature. She broke down the process into four steps. In the first (adiabatic), magnetic field is increased. Since there's no exchange of heat, the net entropy is zero; the magnetic entropy drops and thus the temperature increases. In the next step, a liquid is passed that cools the system, taking away the heating that resulted. In the third step, the adiabatic demagnetization, magnetic entropy is increased, while the thermal entropy is decreased. This step leads to cooling! In the final step, whatever is intended to be cooled is passed through and the process is repeated.

The speaker also did some calculations on the board to arrive at an expression for change in temperature, in terms of the system's parameters.

$$\partial T = -\frac{\mu_0 T}{C_H} \left(\frac{\partial M}{\partial T} \right)_H dH$$

Candidate and prevalent working material for the refrigerator were discussed next which included alloys of gadolinium which are known to produce 3-4 K change, per Tesla of change in magnetic field.

The speaker concluded by drawing a parallel between magnetic and conventional refrigeration, pointing out the role of pressure being played by the magnetic field in the said case. She discussed some advantages, such as environment friendliness and high efficiency (60-70%) while also listing some drawbacks such as its high cost and requirement of rare earth materials.

The speaker was appreciated for her clarity and neat board work by the instructor.

5.1 Acknowledgements

I referred to the wikipedia article on the same topic for filling some gaps.

6 Title : Cooling by Dilution Refrigerator

Speaker : Mr. Subhendu Shekhar

Date : April 18, 2014

Venue : LH-3, LHC

Report :

The speaker started with outlining his talk. His first slide was about how cooling by evaporation of He-4 can only reach about 1.3 K. Below this temperature, the vapour pressure is very small and thus very little evaporates to facilitate the cooling. This issue is resolved by using mixtures of liquid He-3 and He-4. The dilution refrigerator works on this technique.

At 0.7K, the dilution refrigerator starts working. We start with two layers of almost pure He-3 and He-4 at 0.1K. He-3 starts diffusing into He-4. The reason for this was explained later. While diluting He-3 absorbs heat, thus the cooling action.

This next step can be understood in analogy with the evaporation process, wherein we need to remove vapour for evaporation to continue. By mixing into the lower layer, the He-3 layer above is effectively gets diluted. In order to continue cooling, the He-3 dissolved in dilute phase must somehow be removed.

Next the speaker talked about cooling power of the system. Vapour pressure of He falls exponentially with decreasing temperature. It is possible to increase the limiting concentration above 6.6% by increasing pressure, he asserted. He went on to derive an expression for change in entropy for such a system.

Then the realization of such a system was discussed. The basic principle may be described as follows. Like in evaporation, cooling takes place when Helium atoms move from liquid to vapour phase, in dilution, cooling takes place when He-3 atoms move from the concentrated to dilute phase.

Now coming to the reason for diffusion/dilution; it hinges on the statistical characteristics of the constituents. He-3 is a fermion while He-4 is a boson. Consequently, He-3 diffuses into He-4 to minimize its Fermi energy while He-4 tries to gain more super-fluid condensation energy.

Summary (has been written off of one of the slides):

1. The helium undergoes phase separation when cooled below 0.87K giving two phases.
2. The specific heat of He-3 atoms is higher in the dilute phase than in the concentration phase of the atom goes from the concentrated to dilute phase, results in the 'production of cold'.
3. Non-zero solubility of He-3 in He-4 even at 0K leads to cooling power which decreases with T^2 , is much higher than cooling power of evaporation which falls exponentially

6.1 What I'll remember

Even innocent looking phase diagrams can be true gems!
