

Contextuality in a Deterministic Theory

January 15, 2016

Thesis Problem

- ▶ Two theories: Quantum Mechanics (QM) & Bohmian Mechanics (BM)

Thesis Problem

- ▶ Two theories: Quantum Mechanics (QM) & Bohmian Mechanics (BM)
- ▶ Fundamental difference: BM is deterministic, viz. positions (q) and momenta (p) are well defined

Thesis Problem

- ▶ Two theories: Quantum Mechanics (QM) & Bohmian Mechanics (BM)
- ▶ Fundamental difference: BM is deterministic, viz. positions (q) and momenta (p) are well defined
- ▶ Aim: Construct a theoretical situation that defies determinism using QM and analyze using BM

Overview of the talk

- ▶ Bohmian Mechanics (BM)

Overview of the talk

- ▶ Bohmian Mechanics (BM)
- ▶ Determinism: The GHZ test & Contextuality

Overview of the talk

- ▶ Bohmian Mechanics (BM)
- ▶ Determinism: The GHZ test & Contextuality
- ▶ Its analysis using BM

Overview of the talk

- ▶ Bohmian Mechanics (BM)
- ▶ Determinism: The GHZ test & Contextuality
- ▶ Its analysis using BM
- ▶ [recent] Determinism in phase space (q,p)

Overview of the talk

- ▶ Bohmian Mechanics (BM)
- ▶ Determinism: The GHZ test & Contextuality
- ▶ Its analysis using BM
- ▶ [recent] Determinism in phase space (q,p)
- ▶ [in progress] Its analysis using BM

Bohmian Mechanics (BM)

Determinism & Contextuality

Its analysis using BM

(recent) Determinism in phase space (q, p)

(in progress) Its analysis using BM

Bohmian Mechanics (BM)

- ▶ QM: 'Copenhagen interpretation'; how to verify?

Bohmian Mechanics (BM)

- ▶ QM: 'Copenhagen interpretation'; how to verify?
- ▶ D. Bohm: 'hidden variable' theory
 - ▶ in principle known, in practice averaged

Bohmian Mechanics (BM)

- ▶ QM: 'Copenhagen interpretation'; how to verify?
- ▶ D. Bohm: 'hidden variable' theory
 - ▶ in principle known, in practice averaged
- ▶ Lessons:
 - ▶ Clarity: (a) description (b) deriving classical mechanics

Bohmian Mechanics (BM)

- ▶ QM: 'Copenhagen interpretation'; how to verify?
- ▶ D. Bohm: 'hidden variable' theory
 - ▶ in principle known, in practice averaged
- ▶ Lessons:
 - ▶ Clarity: (a) description (b) deriving classical mechanics
 - ▶ Accuracy of deductions: (a) Bell test (b) GHZ/Determinism test

Bohmian Mechanics (BM)

- ▶ QM: 'Copenhagen interpretation'; how to verify?
- ▶ D. Bohm: 'hidden variable' theory
 - ▶ in principle known, in practice averaged
- ▶ Lessons:
 - ▶ Clarity: (a) description (b) deriving classical mechanics
 - ▶ Accuracy of deductions: (a) Bell test (b) GHZ/Determinism test
- ▶ Immediate questions: (a) Uncertainty principle (b) Double slit, which is chosen? (c) Trajectories; observable? (d) Identical particles

BM | Formalism

A particle is associated with (1) (q, p) , which are precisely and continuously defined & (2) a wave (ψ). If the following are assumed, all results of QM are reproduced:

A particle is associated with (1) (q, p) , which are precisely and continuously defined & (2) a wave (ψ). If the following are assumed, all results of QM are reproduced:

- ▶ ψ satisfies the Schrödinger equation

A particle is associated with (1) (q, p) , which are precisely and continuously defined & (2) a wave (ψ). If the following are assumed, all results of QM are reproduced:

- ▶ ψ satisfies the Schrödinger equation
- ▶ $mv = p = \nabla S = \hbar \text{Im}(\nabla \psi / \psi)$, where $\psi = Re^{iS/\hbar}$

A particle is associated with (1) (q, p) , which are precisely and continuously defined & (2) a wave (ψ) . If the following are assumed, all results of QM are reproduced:

- ▶ ψ satisfies the Schrödinger equation
- ▶ $mv = p = \nabla S = \hbar \text{Im}(\nabla \psi / \psi)$, where $\psi = Re^{iS/\hbar}$
- ▶ in practice, we have a statistical ensemble with probability densities $\rho(q) = |\psi(q)|^2$

BM | Formalism

Comments:

- Observers not fundamental; $\hbar = 0$

Comments:

- ▶ Observers not fundamental; $\hbar = 0$
- ▶ Ready generalization to N particles. Non-locality becomes explicit; $p_i = \nabla_i S(q_1, q_2, \dots, q_N)$


Comments:


- ▶ Observers not fundamental; $\hbar = 0$
- ▶ Ready generalization to N particles. Non-locality becomes explicit; $p_i = \nabla_i S(q_1, q_2, \dots, q_N)$
- ▶ Extension to spins: Particle has (q, p) . The wavefunction has the spinor, say $\Psi \equiv (\psi_+, \psi_-)^T$; the generalization is

$$p = mv = \hbar \text{Im} \frac{(\Psi, \nabla \Psi)}{(\Psi, \Psi)}$$

where $(., .)$ is the inner product in the spin space \mathbb{C}^2 .

BM | Pictures





“Observing the Average Trajectories of Single Photons in a Two-Slit Interferometer”

Bohmian Mechanics (BM)

Determinism & Contextuality

Analysis using BM

(recent) Determinism in phase space (q, p)

(in progress) Its analysis using BM

Determinism

Defn: Determinism: Observables have values regardless of whether they are measured, viz. values are predefined.

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.
- ▶ Assume:

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.
- ▶ Assume:
 - ▶ Three particles; interact initially; given to three observers (one each)

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.
- ▶ Assume:
 - ▶ Three particles; interact initially; given to three observers (one each)
 - ▶ Two properties, \hat{X} & \hat{Y} , outcome is ± 1

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.
- ▶ Assume:
 - ▶ Three particles; interact initially; given to three observers (one each)
 - ▶ Two properties, \hat{X} & \hat{Y} , outcome is ± 1
- ▶ General Construction: $\hat{A} \equiv \hat{X} \otimes \hat{Y} \otimes \hat{Y}$, $\hat{B} \equiv \hat{Y} \otimes \hat{X} \otimes \hat{Y}$ and $\hat{C} \equiv \hat{Y} \otimes \hat{Y} \otimes \hat{X}$; outcome is $+1$ for each.
 $\hat{D} \equiv \hat{X} \otimes \hat{X} \otimes \hat{X}$ yields -1 .

Determinism: The GHZ Test (1/2)

- ▶ Objective: To show that the notion of determinism is incompatible with QM.
- ▶ Assume:
 - ▶ Three particles; interact initially; given to three observers (one each)
 - ▶ Two properties, \hat{X} & \hat{Y} , outcome is ± 1
- ▶ General Construction: $\hat{A} \equiv \hat{X} \otimes \hat{Y} \otimes \hat{Y}$, $\hat{B} \equiv \hat{Y} \otimes \hat{X} \otimes \hat{Y}$ and $\hat{C} \equiv \hat{Y} \otimes \hat{Y} \otimes \hat{X}$; outcome is $+1$ for each. $\hat{D} \equiv \hat{X} \otimes \hat{X} \otimes \hat{X}$ yields -1 .
- ▶ Explicit construction: $|\psi\rangle = (|000\rangle - |111\rangle)/\sqrt{2}$
($\sigma_z |0/1\rangle = \pm |0/1\rangle$; $\hat{X}, \hat{Y}, \hat{Z}$ are Pauli spin operators)

Determinism: The GHZ Test (2/2)

- ▶ Hypothesis: Assume that the world is deterministic. $\hat{A}\hat{B}\hat{C}$ must yield $+1$. Also, $\hat{A}\hat{B}\hat{C} = \hat{D}$ (because $Y^2 = 1$). But \hat{D} yields -1 . Thus we get $+1 = -1$.

Determinism: The GHZ Test (2/2)

- ▶ Hypothesis: Assume that the world is deterministic. $\hat{A}\hat{B}\hat{C}$ must yield $+1$. Also, $\hat{A}\hat{B}\hat{C} = \hat{D}$ (because $Y^2 = 1$). But \hat{D} yields -1 . Thus we get $+1 = -1$.
- ▶ Conclusion: The hypothesis is wrong. \implies can't have non-contextual determinism, where “non-contextual” is a subtle but necessary qualification.

Contextuality (1/2)

Consider: A and B are mutually compatible observables (commute).

Contextuality (1/2)

Consider: A and B are mutually compatible observables (commute).

Defn: “Non-contextual” and deterministic (NC): A theory that assigns pre-defined values to A and B , irrespective of which is measured.

Contextuality (1/2)

Consider: A and B are mutually compatible observables (commute).

Defn: “Non-contextual” and deterministic (NC): A theory that assigns pre-defined values to A and B , irrespective of which is measured.

- ▶ Thm: Kochen-Specker proved that NC theories are inconsistent with QM.

Contextuality (1/2)

Consider: A and B are mutually compatible observables (commute).

Defn: “Non-contextual” and deterministic (NC): A theory that assigns pre-defined values to A and B , irrespective of which is measured.

- ▶ Thm: Kochen-Specker proved that NC theories are inconsistent with QM.
- ▶ ‘Proof’ (Mermin’s):

$$\begin{array}{lll} A_{11} = \sigma_z \otimes I & A_{12} = I \otimes \sigma_z & A_{13} = \sigma_z \otimes \sigma_z \\ A_{12} = I \otimes \sigma_x & A_{22} = \sigma_x \otimes I & A_{23} = \sigma_x \otimes \sigma_x \\ A_{31} = \sigma_z \otimes \sigma_x & A_{32} = \sigma_x \otimes \sigma_z & A_{33} = \sigma_y \otimes \sigma_y \end{array}$$

Contextuality (1/2)

Consider: A and B are mutually compatible observables (commute).

Defn: “Non-contextual” and deterministic (NC): A theory that assigns pre-defined values to A and B , irrespective of which is measured.

- ▶ Thm: Kochen-Specker proved that NC theories are inconsistent with QM.
- ▶ ‘Proof’ (Mermin’s):

$$\begin{array}{lll} A_{11} = \sigma_z \otimes I & A_{12} = I \otimes \sigma_z & A_{13} = \sigma_z \otimes \sigma_z \\ A_{12} = I \otimes \sigma_x & A_{22} = \sigma_x \otimes I & A_{23} = \sigma_x \otimes \sigma_x \\ A_{31} = \sigma_z \otimes \sigma_x & A_{32} = \sigma_x \otimes \sigma_z & A_{33} = \sigma_y \otimes \sigma_y \end{array}$$

- ▶ NB: (1) Along a row, compatible; also along a column, (2) product along a row R_k or column C_k is 1, except for $C_3 = -1$. (Hint: $\sigma_z = -i\sigma_x\sigma_y$)

Contextuality (2/2)

- ▶ Thus, QM predicts: $\prod_{k=1,2,3} R_k C_k = -1$; NC theories would yield 1.

Contextuality (2/2)

- ▶ Thus, QM predicts: $\prod_{k=1,2,3} R_k C_k = -1$; NC theories would yield 1.
- ▶ Claim: NC theories yield $\langle \chi_{ks} \rangle = \langle R_1 \rangle + \langle R_2 \rangle + \langle R_3 \rangle + \langle C_1 \rangle + \langle C_2 \rangle - \langle C_3 \rangle \leq 4$ while QM yields $\langle \chi_{ks} \rangle = 6$.

Contextuality (2/2)

- ▶ Thus, QM predicts: $\prod_{k=1,2,3} R_k C_k = -1$; NC theories would yield 1.
- ▶ Claim: NC theories yield $\langle \chi_{ks} \rangle = \langle R_1 \rangle + \langle R_2 \rangle + \langle R_3 \rangle + \langle C_1 \rangle + \langle C_2 \rangle - \langle C_3 \rangle \leq 4$ while QM yields $\langle \chi_{ks} \rangle = 6$.
- ▶ Remarks: (1) Mermin's test is state independent (2) More suited for testing non-contextuality, as locality is not required.

Bohmian Mechanics (BM)

Determinism: The GHZ test & Contextuality

Analysis using BM

(recent) Determinism in phase space (q, p)

(in progress) Its analysis using BM

BM Analysis of the GHZ test (1/2)

- ▶ BM can explain this easily; Not surprising; BM never assigned spins.

BM Analysis of the GHZ test (1/2)

- ▶ BM can explain this easily; Not surprising; BM never assigned spins.
- ▶ Assmptn: (1) Stern Gerlach like measurements to measure spins; (2)
$$|\Psi(r_1, r_2, r_3, t = 0)\rangle = (\psi_{+++} |000\rangle - \psi_{---} |111\rangle)/\sqrt{2},$$
where r_i is the position vector in the frame of the i^{th} observer. (3) Assume a Gaussian particle distribution initially, propagating along the axes of their respective SG apparatus.

BM Analysis of the GHZ test (1/2)

- ▶ BM can explain this easily; Not surprising; BM never assigned spins.
- ▶ Assmptn: (1) Stern Gerlach like measurements to measure spins; (2)
$$|\Psi(r_1, r_2, r_3, t = 0)\rangle = (\psi_{+++} |000\rangle - \psi_{---} |111\rangle)/\sqrt{2},$$
where r_i is the position vector in the frame of the i^{th} observer. (3) Assume a Gaussian particle distribution initially, propagating along the axes of their respective SG apparatus.
- ▶ Claim: Time evolution of $\psi_{\pm\pm\pm}$ can be written as a product of 3 single particle solutions of the SG setup. (Bohm explicitly did the latter)

BM Analysis of the GHZ test (2/2)

- Claim: (1) If SG are setup to measure say $XY\bar{Y}$, then four attractor basins form: $(+++)$, $(+- -)$, $(- - +)$ and $(- + -)$. (2) If SG are setup to measure XXX , then the basins becomes $(- - -)$, $(- + +)$, $(+ - +)$ and $(+ + -)$.

BM Analysis of the GHZ test (2/2)

- ▶ Claim: (1) If SG are setup to measure say $XY\bar{Y}$, then four attractor basins form: $(+++)$, $(+--)$, $(- - +)$ and $(- + -)$. (2) If SG are setup to measure XXX , then the basins becomes $(---)$, $(-++)$, $(+ - +)$ and $(++-)$.
- ▶ Conclusion: Consistent with QM

BM Analysis of the GHZ test (2/2)

- ▶ Claim: (1) If SG are setup to measure say $XY\bar{Y}$, then four attractor basins form: $(+++)$, $(+--)$, $(- - +)$ and $(- + -)$. (2) If SG are setup to measure XXX , then the basins becomes $(---)$, $(-++)$, $(+ - +)$ and $(++-)$.
- ▶ Conclusion: Consistent with QM
- ▶ Remarks: Non locality causes attractor basins to depend on settings of *all* SG apparatus. Contextuality from this perspective is essentially the statement that the results of an experiment, depend on the experiment being performed.

Bohmian Mechanics (BM)


Determinism: The GHZ test & Contextuality

Analysis using BM

(recent) Determinism in phase space (q, p)

(in progress) Its analysis using BM

Determinism in phase space (0/3)



Determinism in phase space (1/3)

Was able to make good progress today. So here's what I did. First of all, consider the same states $|\psi_0\rangle, |\psi_1\rangle$ for $N = 8$, as those considered in my first paper. Recall that for \hat{Z} as defined there, viz. $\hat{Z} = Z(\hat{q} \bmod 2L)$, or more precisely as $\hat{Z} = Z(\hat{q}) = \text{sgn}(\sin(\hat{q}\pi/L))$, we had $\hat{Z}|\psi_0\rangle = |\psi_0\rangle$ and $\hat{Z}|\psi_1\rangle = -|\psi_1\rangle$. In addition to this, I define $\hat{X} = e^{-i\hat{p}L/\hbar}$ (as opposed to defining it to be hermitian). Now, we know that $|\psi_{\pm}\rangle \equiv \frac{|\psi_0\rangle \pm |\psi_1\rangle}{\sqrt{2}}$ is not an eigenstate of \hat{X} . So we optimize the observable \hat{X} to $\hat{X}' \equiv \hat{X} \hat{T}$, where $\hat{T} \equiv e^{i\hat{p}NL a(\hat{q})/2}$ where

$$a(q) = \begin{cases} 1 & 2L < q < 4L \\ 0 & \text{else} \end{cases}.$$

The idea is that you shift certain peaks to the right place, before applying the displacement operator \hat{X} .

Determinism in phase space (2/3)

To illustrate this, consider explicitly

$|\psi_0\rangle = (|\varphi_{-4}\rangle + |\varphi_{-2}\rangle + |\varphi_{-1}\rangle + |\varphi_{-3}\rangle) / \sqrt{4}$. The operation of \hat{T} is $\hat{T}|\varphi_4\rangle = |\varphi_{-5}\rangle$, $\hat{T}|\varphi_3\rangle = |\varphi_{-6}\rangle$ and $\hat{T}|\varphi_n\rangle = |\varphi_n\rangle$ for $n \in \{-4, -3, -2, -1, 1, 2\}$. It is now evident that $\hat{X}' = \hat{X}\hat{T}|\psi_0\rangle = |\psi_1\rangle$. Note also that $\hat{X}'^\dagger|\psi_0\rangle = |\psi_1\rangle$. Similarly $\hat{X}'|\psi_1\rangle = |\psi_0\rangle$ and \hat{X}'^\dagger does the same. So finally, now consider $|G\rangle \equiv (|\psi_0\psi_0\psi_0\rangle - |\psi_1\psi_1\psi_1\rangle) / \sqrt{2}$. With $\hat{A} \equiv \hat{X}' \otimes \hat{Y}' \otimes \hat{Y}'^\dagger$, where $\hat{Y}' \equiv i\hat{Z}\hat{X}'$, calculations yield $\hat{A}|G\rangle = |G\rangle$. With $\hat{B} \equiv \hat{Y}'^\dagger \otimes \hat{X}' \otimes \hat{Y}'$ and $\hat{C} \equiv \hat{Y}' \otimes \hat{Y}'^\dagger \otimes \hat{X}'$ also, by symmetry we get $\hat{B}|G\rangle = |G\rangle$ and $\hat{C}|G\rangle = |G\rangle$. Now $\hat{D} \equiv \hat{A}\hat{B}\hat{C} = \hat{X}' \otimes \hat{Y}'\hat{X}'\hat{Y}'^\dagger \otimes \hat{X}'$ and $\hat{E} \equiv \hat{X}' \otimes \hat{X}' \otimes \hat{X}'$ yield the paradox. If values were predefined, the value of \hat{D} and \hat{E} would return the same answer. However, a simple calculation yields $\hat{D}|G\rangle = |G\rangle$ (this can be seen directly by applying \hat{A} , \hat{B} and \hat{C} sequentially on $|G\rangle$ [was figured the next day]), while $\hat{E}|G\rangle = -|G\rangle$.

Determinism in phase space (3/3)

- ▶ The GHZ argument entails then, that no theory can have q, p NC deterministic.

Determinism in phase space (3/3)

- ▶ The GHZ argument entails then, that no theory can have q, p NC deterministic.
- ▶ Counterexample: BM, a theory which is NC deterministic, in q, p .

Bohmian Mechanics (BM)

Determinism: The GHZ test & Contextuality

Analysis using BM

(recent) Determinism in phase space (q, p)

(in progress) Its analysis
using BM

BM analysis of GHZ 2.0

- ▶ Measuring \hat{X}' is not trivial.

BM analysis of GHZ 2.0

- ▶ Measuring \hat{X}' is not trivial.
 - ▶ Wigner function

BM analysis of GHZ 2.0

- ▶ Measuring \hat{X}' is not trivial.
 - ▶ Wigner function
 - ▶ Functions of q, p directly from the particles q, p

BM analysis of GHZ 2.0

- ▶ Measuring \hat{X}' is not trivial.
 - ▶ Wigner function
 - ▶ Functions of q, p directly from the particles q, p
 - ▶ Formal measurement
- ▶ Formally, any Hermitian operator can be measured using the interaction Hamiltonian, $\hat{H} = -a\hat{Q}\hat{p}_y$, where \hat{Q} is the operator, and y is the coordinate of the pointer.










Conclusion

- ▶ New insight into relation between contextuality and non locality
- ▶ Fundamental difference between spins and (q, p)
- ▶ Meaning of measurement
- ▶ Pointed out and (almost) solved a paradox

The End

Questions

References

-  P. R. Holland, *The Quantum Theory of Motion* (Cambridge University Press, 1993) [cambridge Books Online](#).
-  D. Bohm, *Phys. Rev.* **85**, 166 (1952).
-  T. Durt and Y. Piereaux, *Phys. Rev. A* **66**, 052109 (2002).
-  A. Asadian, C. Budroni, F. E. S. Steinhoff, P. Rabl, and O. Gühne, *Phys. Rev. Lett.* **114**, 250403 (2015).
-  A. Peres, *Physics Letters A* **151**, 107 (1990).
-  E. S. Simon Kochen, *Indiana Univ. Math. J.* **17**, 59 (1968).
-  N. D. Mermin, *Phys. Rev. Lett.* **65**, 3373 (1990).
-  D. Dürr, S. Goldstein, and N. Zanghì, *Quantum Physics Without Quantum Philosophy* (Springer Berlin Heidelberg, 2012).
-  S. Massar and S. Pironio, *Phys. Rev. A* **64**, 062108 (2001)

<http://thisquantumworld.com/wp/wp-content/uploads/2015/01/surreal-1024x586-1024x586.png>