

$$C^{i_1 \dots i_p}_{j_1 \dots j_q}(\vec{x}')$$

$$= \partial_{k_1} x'^{i_1} \dots \partial_{k_p} x'^{i_p} \cdot \partial'_{q_1} x'^{j_1} \dots \partial'_{q_q} x'^{j_q} C^{k_1 \dots k_p}_{l_1 \dots l_q}(\vec{x})$$

Rank

$$(0,0) \text{ scalar } \phi'(\vec{x}') = \phi(\vec{x})$$

$$(1,0) \text{ contravariant vector}$$

$$(0,1) \text{ covariant vector}$$

$$A^{i_1 \dots i_p}_{j_1 \dots j_q} B^{k_1 \dots k_r}_{l_1 \dots l_s} \text{ — This transforms as a } (p+r, q+s) \text{ tensor}$$

$$A^{i_1 \dots i_p}_{j_1 \dots j_q}$$

$$A^{i_1 \dots i_p}_{j_1 \dots j_q}(\vec{x}') = \partial_{k_1} x'^{i_1} \dots \partial'_{q_1} x'^{j_1} A^{k_1 \dots k_p}_{l_1 \dots l_q}(\vec{x})$$

$$= \underbrace{\frac{\partial x'^{i_1}}{\partial x^{k_1}} \dots \frac{\partial x'^{j_1}}{\partial x^{l_1}}}_{\text{which is consistent}} A^{k_1 \dots k_p}_{l_1 \dots l_q}(\vec{x})$$

Claim: Take some point p on the manifold. \vec{x}_0 is the coordinate of p in some coordinate system.

Then \exists a coordinate transf $\vec{x} \rightarrow \vec{x}'$ s.t. $g'_{ij}(\vec{x}'_0) = \text{diag}(\pm 1, \pm 1, \dots, \pm 1)$

NB: For continuous metrics, the number of $+$ & $-$ entries will be constant.

$$x^i = f^i(\vec{x}') \Rightarrow x^i_0 \equiv f^i(\vec{x}'_0)$$

$$x^i = x^i_0 + A^i_k (x'^k - x'^k_0) + B^i_{kl} (x'^k - x'^k_0)(x'^l - x'^l_0) + \dots$$

Here we expand about x_0

$$\begin{aligned} g'_{kl}(\vec{x}'_0) &= \partial'_{k_1} x'^{i_1} \partial'_{l_1} x'^{j_1} g_{ij} \Big|_{\substack{\vec{x}' = \vec{x}'_0 \\ \vec{x} = \vec{x}_0}} \\ &= A^i_k A^j_l g_{ij}(\vec{x}_0) \quad (\text{take } \partial' \text{ \& should be clear}) \\ &= (A^T g A)_{kl} \quad (\text{in matrix notation}) \end{aligned}$$

Notation: Regard g_{ij} as matrix; i as row index, j as column index

Recall: S is symmetric; Thus can be diagonalized with orthogonal matrices.

$$\text{viz. } g = S^T g_d S \quad \text{s.t. } S^T S = \mathbb{1}$$

$$g_d = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix}$$

Non-singular metric only, i.e. eigenvalues are non-zero.

$$\text{Def } R \equiv R^T = \begin{pmatrix} \sqrt{|\lambda_1|} & & 0 \\ & \ddots & \\ 0 & & \sqrt{|\lambda_n|} \end{pmatrix} \Rightarrow R^T \eta R = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix} \quad \text{where } \eta = \begin{pmatrix} \frac{\lambda_1}{|\lambda_1|} & & 0 \\ & \ddots & \\ 0 & & \frac{\lambda_n}{|\lambda_n|} \end{pmatrix}$$

$$\text{Thus, } g = S^T g_d S = S^T R^T \eta R S$$

$$g'_{kl} = (A^T g A)_{kl} = (A^T S^T R^T \eta R S A)_{kl} \Rightarrow \text{for } A = (S)^{-1} \text{ we'll get } g'_{kl}(\vec{x}'_0) = \eta_{kl}$$

This is not unique; eg. $\eta = 1$, then $A = (URS)^{-1}$ will get the right form, where

Signature:

when all +, then euclidean
when one -, then Lorentzian

- NB: (1) Locally then, there's no more information
(2) This will not distinguish the plane & the surface of a sphere.
(3) This is more than diagonalization; A is more than just a $P^T P$ like matrix.

Def: $K_{ijk} = \partial_i g_{jk}$

$$\begin{aligned} K'_{i_1 i_2 i_3} &= \partial'_{i_1} (g'_{i_2 i_3}(\bar{x}')) \\ &= \partial'_{i_1} (\partial'_{i_2} x^{j_2} \partial'_{i_3} x^{j_3} g_{j_2 j_3}(\bar{x})) \\ &= \partial'_{i_1} \partial'_{i_2} x^{j_2} \partial'_{i_3} x^{j_3} g_{j_2 j_3} + \partial'_{i_2} x^{j_2} \partial'_{i_1} \partial'_{i_3} x^{j_3} g_{j_2 j_3} + \\ &\quad \partial'_{i_2} x^{j_2} \partial'_{i_3} x^{j_3} \partial'_{i_1} g_{j_2 j_3} \end{aligned}$$

This is the term we want (why? we'll see soon)

chain rule

$$= \partial'_{i_2} x^{j_2} \partial'_{i_3} x^{j_3} \partial'_{i_1} x^{j_1} \partial_{j_1} g_{j_2 j_3}$$

$$\left(\because \frac{\partial x^{j_1}}{\partial x'^{i_1}} = \frac{\partial g_{j_2 j_3}}{\partial x'^{i_1}} \right)$$

rearrange

$$= \underbrace{\partial'_{i_1} x^{j_1} \partial'_{i_2} x^{j_2} \partial'_{i_3} x^{j_3}}_{\text{this is how } K_{ijk} \text{ should transform if } K_{ijk} \text{ was a tensor}} \partial_{j_1} g_{j_2 j_3} \rightarrow K'_{i_1 i_2 i_3}$$

This is how K_{ijk} should transform if K_{ijk} was a tensor.

However, since the other terms are non-zero, $\Rightarrow K_{ijk}$ is not a tensor under general coordinate transformations.

Idea: Take a linear combination of K_{ijk} to construct a tensor.
: This is not possible. (or even polynomials of K_{ijk})

Proof sketch: Assume we can show that given a point, \exists a coordinate transformation (CT) s.t. K vanishes. Therefore, it must vanish in all coordinates (\because tensors transform in that way). Therefore it's impossible to find such a tensor (it'll be zero essentially), (if this argument is repeated for all points)

Recall: $x^i = x^i_0 + A^i_j (x'^j - x'^j_0) + B^i_{jk} (x'^j - x'^j_0)(x'^k - x'^k_0) + \dots$

We need 2 derivatives $\Rightarrow B^i_{jk}$ will be important

B^i_{jk} will be unique. π

We want $K'_{ijk} = 0$, so $\frac{n(n+1)}{2} \cdot n$ eg's are required to be zero.

constants is also $\frac{n(n+1)}{2} \cdot n$ ($\because B^i_{jk}$ also has similar symmetric aspects as K_{ijk})

Exercise: Find B^i_{jk} explicitly, to make $K'^i_{j,k} = 0$

-2-

Idea: Take 2 derivatives of g_{ij}

$$S_{ijkl} = \partial_i \partial_j g_{kl}$$

(Claim: This itself is not a tensor.)

: Use the old argument. See if it can be shown to be zero in some coordinate sys.

At \vec{x}_0 , # of components of $S_{ijkl} = \frac{n(n+1)}{2} \cdot \frac{n(n+1)}{2}$

Recall: $x^i = x^i_0 + A^i_j(\dots)^j + B^i_{jk}(\dots)^{jk} + C^i_{jkl}(\dots)^{jkl}$

Now # of parameters in $C^i_{jkl} = n \frac{n(n+1)(n+2)}{3!}$ & \leq (for large n)

\Rightarrow There're more constraints than parameters. This means one can't in general get

result. So $S_{ijkl} = 0$.
 $\frac{n^2(n+1)^2}{4} - \frac{n^2(n+1)(n+2)}{6} = \frac{1}{12} n^2(n^2-1) \begin{cases} 1 & \text{for } n=2 \\ 6 & \text{for } n=3 \\ 20 & \text{for } n=4 \end{cases}$

which means the tensor so constructed will have the aforesaid # of free parameters.

Defⁿ: $\Gamma^i_{jk} = \frac{1}{2} g^{il} (\partial_j g_{lk} + \partial_k g_{lj} - \partial_l g_{jk}) :=$ Christoffel symbol / connection

$R^i_{jkl} = \partial_l \Gamma^i_{jk} - \partial_k \Gamma^i_{jl} + \Gamma^m_{jk} \Gamma^i_{lm} - \Gamma^m_{jl} \Gamma^i_{km} :=$ Riemann tensor

Exercise: $\Gamma'^l_{mn}(\vec{x}') = \partial_i x'^l \partial'_m x^i \partial'_n x^k \Gamma^i_{jk}(\vec{x}) + \partial_k x'^l \partial'_m \partial'_n x^k$

NB: This is not a tensor \because of the second term.

: R^i_{jkl} is a tensor of rank (1,3)

: R^i_{jkl} has $\frac{1}{12} n^2(n^2-1)$ independent parameters.

$$x^i = f^i(\vec{x}')$$

$$x_0^i = f^i(\vec{x}_0')$$

$$x^i = x_0^i + \left(\frac{\partial f^i}{\partial x_j} \right) (x_0^j - x^j) + \dots$$

$$A^i_j (x_0^j - x^j)$$

$$ds^2 = g_{ij} x^i x^j$$

Doubt: If $\partial_i g_{jk} = 0$ then how can
 $\partial_m \partial_i g_{jk} \neq 0$ in general?

