

Bell Test with Continuous Variables

Atul Singh Arora

*University of Siegen, Germany and
Indian Institute of Science Education & Research, Mohali, India*

Ali Asadian

*University of Siegen, Germany
(Dated: May-July 2015)*

Abstract

We show the violation of the Bell inequality in a more classical setting than the usual quantum spin, viz. with continuous variables x, p . This is achieved using displacement operators, or equivalently modular momentum and position. Two possible experimental realizations of the scheme have also been delineated, one uses photons, while the other is based on electrons.

I. INTRODUCTION

The Bell inequality is a statement about the statistical outcomes of a certain class of experiments. The inequality was formulated by assuming the most general ‘local theory’. Any violation of the inequality would imply that local realism must be false. Quantum Mechanics predicts a violation which has been confirmed experimentally. The physical situation for which this violation is usually formulated, uses entangled spin states. Here we show that a violation can be achieved without having to use spins, which is a purely quantum phenomenon; measuring a particle’s position is sufficient.

II. VIOLATION OF BELL INEQUALITY

A. CHSH Inequality | prior art

Here the violation of Bell’s inequality for the typical spin case, is stated in a way that is easier to translate to the continuous variable case. We first state the Bell test in its modern form, known as the CHSH inequality.

Suppose: There are two observers[1], Alice and Bob.

There are two particles, which are allowed to interact initially and are then separated. Alice takes one, Bob takes the other. Alice can measure one of two properties of her particle, call them a_1, a_2 . Similarly Bob can measure b_1, b_2 of his particle.

Notatn: $\langle a_i b_j \rangle$ = Average value obtained when Alice chooses to measure a_i and Bob chooses to measure b_j

Statemnt: $|\langle a_1(b_1 + b_2) + a_2(b_1 - b_2) \rangle| \equiv |\langle C(a_i, b_i) \rangle| \leq 2$ if (a) result of measuring $a_i, b_i \in [-1, 1]$, (b) local realism is assumed. This result will be referred to as the CHSH inequality.

Quantum Mechanics predicts a violation of this in the following physical situation.

Defn: $|\psi\rangle \equiv \frac{|+-\rangle - |-+\rangle}{\sqrt{2}}$ where the $|\pm\rangle$ states are s.t. $\hat{\sigma}_x |\pm\rangle = \pm |\pm\rangle$.

	Alice		Bob	
Convention	a_1	a_2	b_1	b_2
Angle	$\theta_1 = \pi/2$	$\theta_2 = 0$	$\phi_1 = \pi/4$	$\phi_2 = \pi/4 + \pi/2$

Table I: Measurement Scheme

Consider: (a) Two particles are described by $|\psi\rangle$, (b) the first particle is with Alice, second with Bob, and (c) Alice measures \hat{z} or \hat{x} while Bob is allowed to choose between \hat{z}' and \hat{x}' , where $\hat{z}' = \frac{-\hat{z} - \hat{x}}{\sqrt{2}}$, $\hat{x}' = \frac{\hat{z} - \hat{x}}{\sqrt{2}}$.

Notatn: The Pauli matrices are referred to as $\hat{x}, \hat{y}, \hat{z}$.

Result: $\langle C((\hat{x}, \hat{z}), (\hat{x}', \hat{z}')) \rangle = 2\sqrt{2} \not\leq 2$, which is a violation.

This value can be reproduced in a slightly modified situation.

Modifctn: (i) Due to the symmetry in the setup, we expect the same results hold for \hat{x}, \hat{y} and \hat{x}', \hat{y}' and (ii) we demand that Alice and Bob can only measure \hat{x} but are allowed to perform a local unitary before the measurement.[2]

Explicitly then, consider: (a) The first qubit is with Alice and the second is with Bob. (b) Alice and Bob apply local unitaries $e^{i\hat{z}\theta/2} \otimes I$ and $I \otimes e^{i\hat{z}\phi/2}$ respectively and then both measure \hat{x} . (c) Alice can choose $\theta \in \{0, \pi/2\}$ and Bob can choose $\phi \in \{\pi/4, \pi/4 + \pi/2\}$; see Table I.[3]

Now it remains to show that this scheme also violates the Bell inequality (and by the same value). To that end, we state a result.

Claim(3): $\langle e^{-i\hat{z}\theta/2} \hat{x} e^{i\hat{z}\theta/2} \otimes e^{-i\hat{z}\phi/2} \hat{x} e^{i\hat{z}\phi/2} \rangle = -\cos(\phi - \theta)$

Given that, it is straight forward to evaluate

CHSH:

$$\begin{aligned} \langle C \rangle &= \langle a_1 b_1 \rangle + \langle a_1 b_2 \rangle + \langle a_2 b_1 \rangle - \langle a_2 b_2 \rangle \\ &= \left(-\frac{1}{\sqrt{2}} \right) + \left(-\frac{1}{\sqrt{2}} \right) + \left(-\frac{1}{\sqrt{2}} \right) - \left(+\frac{1}{\sqrt{2}} \right) \\ &= -2\sqrt{2} \end{aligned}$$

	Alice		Bob	
Convention	a_1	a_2	b_1	b_2
Angle	$\theta'_1 = 0$	$\theta'_2 = -\pi/2$	$\phi'_1 = 3\pi/8$	$\phi'_2 = 5\pi/4$

Table II: ‘Corrected’ Measurement Scheme

Remark: This is a violation since $|\langle C \rangle| = 2\sqrt{2} \not\leq 2$.

However, to convert this to a more conventional form, we simply redefine the angles θ and ϕ . [4]

Substn: $\phi \rightarrow \phi' = \phi + \pi/2$, $\theta \rightarrow \theta' = \theta - \pi/2$; see Table II

Now we have $\langle C \rangle = 2\sqrt{2}$.

Concln: The violation of the Bell spin state can be equivalently be obtained by the measurement of \hat{x} after applying a rotation along \hat{z} by the aforesaid angles.

B. Modular Variables and Displacement Operators | prior art

The observables admissible in the CHSH inequality must $\in [-1, 1]$. If we consider the *real variable* $p \in (-\infty, \infty)$, then one way of obtaining a bound variable is to use $p_{mod} \equiv p \bmod h/L$, where h/L has been introduced for dimensional reasons. This forces $p_{mod} \in [0, h/L)$. If we further consider, say $\cos(p_{mod}L/h)$, then we get precisely the bound needed for the CHSH inequality. We chose the factor so that for $p_{mod} = h/L$, the cos function also completes a period. Note that $\cos(p_{mod}L/h) = \cos(pL/h) = \frac{e^{ipL/h} + e^{-ipL/h}}{\sqrt{2}}$.

Next, recall that in quantum mechanics, the momentum that’s quantized, is one that generates infinitesimal translations. Formally, this means that $(1 - i\hat{p}\delta L/h)|x\rangle = |x + \delta L\rangle$. If we want a finite translation, we can do the following procedure: $\lim_{N \rightarrow \infty} \left(1 - i\hat{p}\frac{L}{hN}\right)^N |x\rangle = e^{-i\hat{p}L/h}|x\rangle = |x + L\rangle$. This operator is aptly called the displacement operator. We must add that $e^{i\hat{p}a}e^{i\hat{x}b} = e^{i\hbar ab}e^{i\hat{x}b}e^{i\hat{p}a}$. Thus for appropriate choices of a, b , the displacement operators can be made to commute or anti-commute. In the former case, it means one can simultaneously measure modular position and momentum (which is striking) and in the latter case, one can define pauli matrix like commutation.

The idea was that the obvious generalization for obtaining an appropriate operator for the CHSH inequality is $\cos(\hat{p}L/h) = \frac{e^{i\hat{p}L/h} + e^{-i\hat{p}L/h}}{\sqrt{2}}$. The action of the operator will guide us in creating the appropriate states in the next sub section.

C. Violation in Continuous Variable State

We start with constructing continuous variable kets that satisfy the same orthogonality and mutual relations

as $|0\rangle, |1\rangle, |+\rangle, |-\rangle$. [5][6] We then construct the relevant operators and finally evaluate $\langle C \rangle$ to show the violation.

To construct the states, we start with some definitions.

Forgt: $\hat{x}, \hat{y}, \hat{z}$ represent Pauli matrices

Defn: $L \equiv$ some length scale, $N \equiv$ number of slits

Consider: A state $|\varphi\rangle$ symmetric about $x = L/2$, with spread $\ll L$. Formally, $|\varphi\rangle \equiv \int dx \langle x - L/2 | \varphi \rangle |x\rangle$ where $|\varphi\rangle$ s.t. (a) $\langle x | \varphi \rangle = \langle -x | \varphi \rangle$ and (b) it’s spread is $\ll L$

Defn: $|\varphi_n\rangle \equiv \int dx \langle x - nL | \varphi \rangle |x\rangle$

Defn: $I_N = \{-\frac{N}{2}, \dots, \frac{N}{2} - 1, \frac{N}{2} - 1\}$, defined for even N only (for simplicity)

Defn:

$$|0'\rangle \equiv \frac{1}{\sqrt{N}} \sum_{n \in I_N} |\varphi_{2n+1}\rangle$$

$$|1'\rangle \equiv \frac{1}{\sqrt{N}} \sum_{n \in I_N} |\varphi_{2n}\rangle$$

Using these states, we construct the analogues of the $|+\rangle$ and $|-\rangle$ states.

Remark: Definition of $|0'\rangle$ and $|1'\rangle$ justify the word ‘slit’ in the definition of N .

Defn: $|+\rangle \equiv \frac{|0'\rangle + |1'\rangle}{\sqrt{2}}$, $|-\rangle \equiv \frac{|0'\rangle - |1'\rangle}{\sqrt{2}}$

NB: $\langle 0' | 1' \rangle = 0$, $\implies \langle + | - \rangle = 0$

Analogous to the spin case, we construct the entangled state

Defn: $|\Psi\rangle \equiv \frac{|+\rangle - \langle - | + \rangle}{\sqrt{2}}$

These states were constructed with a partial translational symmetry which is appropriate to the hermitian operator, $\cos(\hat{p}L/h)$ discussed earlier.

Defn:

$$\hat{X} \equiv \frac{e^{i\hat{p}L/h} + e^{-i\hat{p}L/h}}{2}$$

NB: Only in the limit $N \rightarrow \infty$,

$$\hat{X} |+\rangle = + |+\rangle$$

$$\hat{X} |-\rangle = - |-\rangle$$

just as was with $\hat{\sigma}_x$. In general however, the relation doesn’t hold.

We have almost all the cards, except for one that allows for setting $\{\phi, \theta\}$. In the spin case, this was achieved by operating $e^{i\hat{\sigma}_z \phi/2}$ on the state before measurement. Correspondingly, here we define this operator by it’s action on the relevant state.




Figure 1: Practically, the number of slits, N will be finite. The plot shows $\langle C \rangle$ as a function of N . To get a violation, we need merely 7 slits, while with 50 we are almost at $2\sqrt{2}$.

Defn: $\hat{U}(\phi)$ is defined to be s.t.

$$\begin{aligned}\hat{U}(\phi) |0'\rangle &= e^{i\phi/2} |0'\rangle \\ \hat{U}(\phi) |1'\rangle &= e^{-i\phi/2} |1'\rangle\end{aligned}$$

This can be achieved in principle using alternate glass slabs (capacitors) for photons (fermions) to adjust the relative phase as desired, since $|0'\rangle$ and $|1'\rangle$ are spatially separated. Alternatively, we could also define $\hat{U}(\phi)$ in its analytic form as

Alt Defn:

$$\hat{U}(\phi) = e^{iZ(\hat{x}_{\text{mod}2L})\phi/2}$$

where $Z(x') \equiv \begin{cases} 1 & 0 < x' \bmod 2L \leq L \\ -1 & L < x' \bmod 2L \leq 2L \end{cases}$ when Z acts on an eigenstate of \hat{x} . [7]

NB: $Z(\hat{x}_{\text{mod}2L}) |0'\rangle = |0'\rangle$ and $Z(\hat{x}_{\text{mod}2L}) |1'\rangle = -|1'\rangle$ viz. $Z(\hat{x}_{\text{mod}2L})$ plays the role of $\hat{\sigma}_z$

Notatn: $\hat{Z} \equiv Z(\hat{x}_{\text{mod}2L})$

To evaluate the CHSH inequality, we again need terms of the form $\langle \hat{U}^\dagger(\phi_i) \hat{X} \hat{U}(\phi_i) \otimes \hat{U}^\dagger(\theta_i) \hat{X} \hat{U}(\theta_i) \rangle$. If we assume $N \rightarrow \infty$, these can be evaluated easily to be the same as before $-\cos(\phi_i - \theta_i)$. Thus, the same analysis as before yields $\langle C \rangle = 2\sqrt{2}$ which is the same maximal violation, in a completely different setup. If N is not assumed to be arbitrarily large, then it can be shown that (see section V 5)

$$\begin{aligned}\langle \hat{U}^\dagger(\phi_i) \hat{X} \hat{U}(\phi_i) \otimes \hat{U}^\dagger(\theta_i) \hat{X} \hat{U}(\theta_i) \rangle &= \\ &= -\left(\frac{N-1}{N}\right)^2 \cos(\phi_i - \theta_i)\end{aligned}$$

Thus obviously then, $\langle C \rangle = \left(\frac{N-1}{N}\right)^2 2\sqrt{2}$. If we ask for the smallest N s.t. $\langle C \rangle > 2$, we get $N \geq 7$; see Figure 1.

D. Validating phase space behaviour

In the past, it has been shown that one can arrive at the violation of Bell inequalities in continuous variable systems. One issue concerning these is that the Wigner function corresponding to the observable operators, are unbounded. An example of this is parity. Note that the Wigner function for a physical state ρ has an upper bound. Here however, we are simply using the prescription to evaluate it for unitary operators.

We care because Wigner functions may be thought of as a specific case of the Wigner-Weyl correspondence, which relates classical functions to their corresponding quantum operators. Thus, in a phase-space description of quantum mechanics, it is important to show that it is not necessary to have unboundedness to show a violation of the Bell's inequality, and thereby non-locality.

Our set of permissible states is manifestly bounded. We must show that the observable in our case, given by $\hat{U}^\dagger(\phi) \hat{X} \hat{U}(\phi)$ is also bounded. To that end, we note that

$$\begin{aligned}W_1(q', p') &= \frac{1}{2\pi\hbar} \int dq e^{ip'q/\hbar} \left\langle q' - \frac{q}{2} \right| e^{-iZ(\hat{x}_{\text{mod}2L})\phi/2} \\ &\quad e^{ip'\frac{L}{\hbar}} e^{iZ(\hat{x}_{\text{mod}2L})\phi/2} \left| q' + \frac{q}{2} \right\rangle \\ &= \frac{1}{2\pi\hbar} e^{-iZ_1(q')\phi/2} e^{ip'\frac{L}{\hbar}} e^{iZ_2(q')\phi/2} \\ &= \frac{1}{2\pi\hbar} e^{ip'L/\hbar} e^{iZ_2(q')\phi}\end{aligned}$$

where $Z_{1/2}(q') = Z[(q' \mp \frac{L}{2}) \bmod 2L]$ and we used the fact that $Z(x') = -Z(x' + L)$ (omitting the $\bmod 2L$). Similarly one can evaluate W_2 and obtain

$$W = (W_1 + W_2)/2 = \frac{1}{2\pi\hbar} \cos(p'L/\hbar) e^{\pm i\phi}$$

which corresponds to the observable (we used that $Z_2 = \pm 1$). Clearly $|W| \leq (2\pi\hbar)^{-1}$.

E. Source of Violation

In the Bell test done with spins, the source of violation hinges on the non-commutativity of the Pauli matrices. It is known that $C^2 = 4\mathbb{I} + [a_1, a_2] \otimes [b_1, b_2]$, when $a_i, b_i \in \{-1, 1\}$. To obtain a violation in general (where the first term maybe less than $4\mathbb{I}$), it is therefore necessary that the commutations don't vanish. This is indeed true in our case. Explicitly, we must have $[\hat{U}^\dagger \hat{X} \hat{U}, \hat{U}'^\dagger \hat{X} \hat{U}'] \neq 0$. To arrive at the desired expression we first evaluate some simpler commutation and anti-commutation relations explicitly. Let's start with $[\hat{Z}, e^{ipL/\hbar}] = \hat{Z} e^{ipL/\hbar} - e^{ipL/\hbar} \hat{Z}$. To evaluate it, we multiply the second term with $\int dx |x\rangle \langle x|$ and obtain $Z e^{ipL/\hbar} + Z e^{ipL/\hbar}$ where we've used $Z(\hat{x}_{\text{mod}2L}) = -Z((\hat{x} \pm L)_{\text{mod}2L})$ and $e^{ipL/\hbar} |x\rangle = |x - L\rangle$. Thus we

have

$$[\hat{Z}, \hat{X}] = 2\hat{Z}\hat{X} = -2i\hat{Y}$$

where $\hat{Y} = i\hat{Z}\hat{X}$. Here i was introduced to ensure $\hat{Y}^\dagger = \hat{Y}$, since $\hat{X}^\dagger = \hat{X}$ and $\hat{Z}^\dagger = \hat{Z}$. Similarly it follows that $\{\hat{Z}, \hat{X}\} = 0$. From the definition of Y and the anti-commutation, $\{\hat{Y}, \hat{X}\} = 0$ and $\{\hat{Y}, \hat{Z}\} = 0$ also follow trivially. While we're at it, we may state that while $\hat{Z}^2 = 1$, it is not a sum of a 2 state projector and $\hat{X}^2 \neq 1$ in general. This manifests in the other commutation relations

$$\begin{aligned} [\hat{X}, \hat{Y}] &= -2i\hat{Z}\hat{X}^2 = -2i\hat{X}^2\hat{Z} \\ [\hat{Y}, \hat{Z}] &= -2i\hat{X} \end{aligned}$$

Further, note that $e^{-i\hat{Z}\phi/2}Xe^{i\hat{Z}\phi/2} = Xe^{i\hat{Z}\phi}$ and that $e^{i\hat{Z}\phi} = \cos\phi + i\hat{Z}\sin\phi$. With all these, it is easy to show that

$$\begin{aligned} [U^\dagger(\frac{\phi}{2})XU(\frac{\phi}{2}), U^\dagger(\frac{\phi'}{2})XU(\frac{\phi'}{2})] &= \\ (\cos\phi\sin\phi' - \sin\phi\cos\phi')\hat{Z}\hat{X}^2 &\neq 0 \end{aligned}$$

as was necessary. Ofcourse this is not sufficient, however we have shown explicitly the violation. This evaluation was meant to illuminate that the source of violation essentially hinges on the non-commutativity of $[\hat{x}_{mod2L}, \hat{p}_{mod2L}] \neq 0$. It is apparent that the complete pauli algebra is unnecessary for showing a violation.

F. Advantage of asymmetry in \hat{X} and \hat{Z}

Considering the operator (non-hermitian for simplicity) $\hat{X} = e^{i\hat{p}L/\hbar}$, defining $\hat{Z} = e^{i\hat{x}2\pi/2L}$ is more natural. They also follow the desired anti-commutation $\{\hat{X}, \hat{Z}\} = 0$ and we could define $\hat{Y} = i\hat{Z}\hat{X}$ to get a more natural generalization. The question is why did we choose $\hat{Z} = Z(\hat{x}_{mod2L})$. The cause of this asymmetry hinges on the preferential treatment of position space. We could have constructed states of the form $|0'\rangle = \sum_n |x + nd\rangle$ and used the natural definition of \hat{Z} to obtain the violation. The issue is that this forces us to choose a countable superposition of position eigenkets as our desired state. Such a state is strictly not even in the Hilbert space. In the scheme proposed, this issue is resolved by $Z(\hat{x}_{mod2L})$ which causes a larger set of physically realizable states to become admissible.

Thus while our construction is motivated by the spin formalism, it is not just a relabelling of states and operators achieved by a severe restriction of the Hilbert space.

III. PHYSICAL IMPLEMENTATION

We show that this scheme can be implemented rather easily using photons. We harness the two degrees of freedom of a photon, it's polarization and it's position to

construct the required state. With a slightly modified setup, it is possible to do the same with spin and position for matter waves (see section V 7). The overall setup (see figure 2) is such that we need only consider the quantum mechanical description along the x -axis.

A. Creation of the entangled state

The desired entangled state is $|\Psi\rangle \equiv \frac{|+-\rangle - |-+\rangle}{\sqrt{2}}$. We start with noting the triviality of constructing a

$$|+\rangle = \frac{1}{\sqrt{2N}} \sum_{n \in I_{2N}} |\varphi_n\rangle$$

state. One simply needs a source and a grating, viz. a screen with $2N$ slits of width $a \ll L$, separated by a distance L (centre to centre). The

$$|-\rangle = \frac{1}{\sqrt{2N}} \sum_{n \in I_{2N}} (-1)^{n+1} |\varphi_n\rangle$$

state can be similarly constructed by using glass slabs at alternate slits, such that the phase introduced is $\phi = \pi \implies e^{i\phi} = -1$. In figure 2, if you consider only one particle, and disregard everything after the grating, then the setup is expected to produce a $|+\rangle$ state, just after the grating. To produce the desired entangled state, we start with two entangled photons, such that their polarization state can be expressed as

$$|\chi\rangle = \frac{|HV\rangle - |VH\rangle}{\sqrt{2}}$$

Their spatial description (along x -axis) is initially assumed to be $|\zeta\rangle = |\gamma\rangle \otimes |\gamma\rangle = |\gamma\gamma\rangle$. The state $|\gamma\rangle$ maybe considered to be a gaussian with $\sigma \gg 2NL$ so that after the grating, the spatial state can be written as $|\zeta_{\text{grating}}\rangle = |++'\rangle$. Now comes the interesting part. To produce a $|-\rangle$ state, we already know we needed alternate glass slabs. If we had glass slabs, whose refractive index (given some orientation) was say $\eta_H = 1$ for a horizontally polarized beam and $\eta_V = \eta \neq 0$ for vertical polarization, then we could harness the entangled polarization state to create the required spatially entangled state. Birefringent crystals have such polarization dependent refractive indices. Assume that alternating birefringent crystals have been placed after both the gratings as shown in figure 2 (in the diagram, they're not alternating though). These have width such that the resultant phase it introduces is π . The combined initial state is given by

$$|\chi\rangle |\zeta\rangle = \frac{|HV\rangle - |VH\rangle}{\sqrt{2}} |\gamma\gamma\rangle$$

After the grating, the state is

$$|\chi\rangle |\zeta_{\text{grating}}\rangle = \frac{|HV\rangle - |VH\rangle}{\sqrt{2}} |++'\rangle$$

And finally after the crystals, the combined state becomes

$$|\chi\rangle |\zeta_{\text{birefringent}}\rangle = \frac{|HV\rangle |+'-' \rangle - |VH\rangle |-'+\rangle}{\sqrt{2}}$$

At this stage, if we were to trace out the polarization state, we'd end up with a mixed state. That is useless for our test. Ignoring the glass slabs at the moment in figure 2, we observe the action of the polarisers on the state. After the 45° polariser, the state becomes

$$|\chi_{45}\rangle |\zeta_{\text{birefringent}}\rangle \propto \frac{1}{\sqrt{2}} \left(\frac{|H\rangle + |V\rangle}{\sqrt{2}} \frac{|H\rangle + |V\rangle}{\sqrt{2}} |+'-' \rangle - \frac{|H\rangle + |V\rangle}{\sqrt{2}} \frac{|H\rangle + |V\rangle}{\sqrt{2}} |-'+\rangle \right)$$

Rewriting $(|H\rangle + |V\rangle)/\sqrt{2} = |H'\rangle$, we get

$$|\chi_{45}\rangle |\zeta_{\text{birefringent}}\rangle \propto |H'H'\rangle \left(\frac{|+'-' \rangle - |-'+\rangle}{\sqrt{2}} \right)$$

Now if we trace out the polarization state, we'd be left with the target entangled state. As a remark, it maybe be stated that although to arrive at this result we assumed that $\eta_H = 1$, which is unreasonable physically, we can compensate for $\eta_H \neq 0$ by putting appropriate glass slabs at the alternate empty slits, to produce zero relative phase when the polarization is horizontal.

B. Measurement Settings

In our scheme, Alice and Bob can choose $\phi \in \{\phi_1, \phi_2\}$ and $\theta \in \{\theta_1, \theta_2\}$ respectively. To implement this, we note that the action of $\hat{U}(\theta)$ is to introduce a phase difference between the $|0'\rangle$ and $|1'\rangle$ states. By construction, $|0'\rangle$ and $|1'\rangle$ are spatially disjoint; thus the operation of $\hat{U}(\theta)$ can be implemented by simply placing alternating glass slabs at the slits, with widths such that $|0'\rangle \rightarrow e^{i\theta/2} |0'\rangle$ and $|1'\rangle \rightarrow e^{-i\theta/2} |1'\rangle$. The same operation maybe done for $\hat{U}(\phi)$ for the second particle. In the overall scheme, as shown in figure 2, this is done by the glass slabs right after the birefringent crystals. The state after application of the unitaries is given by $|\Psi_f\rangle$, where f is used to index which of the four possible measurement settings was used.

C. Measurement

The scheme requires us to evaluate the expectation value of $\hat{X} \otimes \hat{X}$ (after the measurement settings have been applied by the appropriate unitaries). If from the experiment, we can obtain the probability $|\langle p_A, p_B | \Psi_f \rangle|^2$, then evaluating $\langle \hat{X} \otimes \hat{X} \rangle = \langle \cos(\hat{p}L/\hbar) \otimes \cos(\hat{p}L/\hbar) \rangle$ simply amounts to $\int dp_A dp_B \cos(p_A L/\hbar) \cos(p_B L/\hbar) |\langle p_A, p_B | \Psi_f \rangle|^2$.

Therefore it is sufficient to explain how to obtain the joint momentum probability distribution from the experiment, for a given measurement setting. We start with stating the result that

$$|\langle x_A, x_B | \Psi_{f \text{ screen}} \rangle|^2 \propto \left| \left\langle p_A = \frac{p_z x_A}{D}, p_B = \frac{p_z x_B}{D} | \Psi_f \right\rangle \right|^2$$

where $|\Psi_{f \text{ screen}}\rangle$ is the state of the system at the screen, D is the distance between the gratings and the screens and p_z is the z component of momentum of the particle. For a photon, $p_z = h/\lambda$ while for an electron, $p_z = m_e D/T$, where T is the time taken to arrive at the screen from the grating. The idea is simply that the momentum distribution at the grating can be recovered by observing the spatial distribution at a screen, sufficiently far away.

In an experiment then, there are two possibilities.

(a) Given f , Alice and Bob both note the position at which they obtain their particle. After repeating the experiment sufficiently many times, they share their list to create a sequence $\{(x_A^{(i)}, x_B^{(i)})\}$. From this sequence, they create a 2D histogram, by simply counting how many time they got (x'_A, x'_B) to lie inside a given cell. The normalized result is essentially $|\langle x_A, x_B | \Psi_{f \text{ screen}} \rangle|^2$ from which $\langle X \otimes X \rangle$ can be computed as discussed. f is changed to evaluate all the 4 terms to finally obtain $\langle C \rangle$ experimentally.

(b) The other simpler, equivalent and direct possibility is that Alice and Bob both obtain the position, $\{x_A^{(i)}, x_B^{(i)}\}$ for the i th repetition and evaluate $\cos(p_A^{(i)} L/\hbar)$ and $\cos(p_B^{(i)} L/\hbar)$. After sufficient iterations, they exchange their observations to simply evaluate the average, viz.

$$\frac{\sum_i \cos(p_A^{(i)} L/\hbar) \cos(p_B^{(i)} L/\hbar)}{\sum_i} = \langle C \rangle$$

IV. CONCLUSION

We have shown how to realize a Bell test in continuous variables position and momentum using specifically chosen and physically realizable states. In addition it has been demonstrated that the exact pauli commutation relations aren't necessary to capture the essence of non-locality; infact by relaxing the similarity, the set of states that show a violation is made considerably larger.

V. APPENDIX

Illustration (I)

This is trivial if one uses the Bloch sphere picture. Instead of measuring along an arbitrary axis, we rotate the Bloch sphere appropriately, and then measure x . To illustrate this, consider




Figure 2: Scheme for creating the state

Questn: \exists a U , s.t. if $|\chi\rangle \rightarrow |\chi'\rangle = U|\chi\rangle$ then $\langle\chi|x|\chi\rangle = \langle\chi'|y|\chi'\rangle$?
Explicitly, we have

$$\begin{aligned} y &= U^\dagger x U = e^{-iz\theta/2} x e^{iz\theta/2} \\ &= x e^{iz\theta} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{bmatrix} \\ &= \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \end{aligned}$$

for $\theta = \pi/2$ as one would guess geometrically.

Claims

1. If $|\psi\rangle \equiv \frac{|+-\rangle - |-+\rangle}{\sqrt{2}}$, then $|\psi\rangle = \frac{|10\rangle - |01\rangle}{\sqrt{2}}$
2. $\langle x \otimes x \rangle = -1$, for $|\psi\rangle$ in Claim(1)
3. $\langle e^{-iz\theta/2} x e^{iz\theta/2} \otimes e^{-iz\phi/2} x e^{iz\phi/2} \rangle = -\cos(\phi - \theta)$

Proof:

$$\begin{aligned} \text{LHS} &= \langle x e^{-iz\theta} \otimes x e^{iz\phi} \rangle \\ &= \left[\frac{\langle 10| - \langle 01|}{\sqrt{2}} \right] [x e^{iz\theta} \otimes x e^{iz\phi}] \left[\frac{|10\rangle - |01\rangle}{\sqrt{2}} \right] \\ &= \langle \psi | x \otimes x \\ &\quad \left[\frac{e^{i(\phi-\theta)} \left(\frac{|+-\rangle - |-+\rangle}{\sqrt{2}} \right) \left(\frac{|+-\rangle + |-+\rangle}{\sqrt{2}} \right)}{\sqrt{2}} \right. \\ &\quad \left. - \frac{e^{-i(\phi-\theta)} \left(\frac{|+-\rangle + |-+\rangle}{\sqrt{2}} \right) \left(\frac{|+-\rangle - |-+\rangle}{\sqrt{2}} \right)}{\sqrt{2}} \right] \end{aligned}$$

We define, $\delta \equiv \phi - \theta$ and using section V 1, it follows that only terms like $|+-\rangle$ or $|-+\rangle$; so

$$\begin{aligned} \text{LHS} &= \langle \psi | x \otimes x \\ &\quad \left[\frac{e^{i\delta} \left(\frac{|+-\rangle - |-+\rangle}{2} \right) - e^{-i\delta} \left(\frac{-|+-\rangle + |-+\rangle}{2} \right)}{\sqrt{2}} \right] \\ &= \langle \psi | x \otimes x \left[\frac{e^{i\delta} \left(\frac{|\psi\rangle}{\sqrt{2}} \right) + e^{-i\delta} \left(\frac{|\psi\rangle}{\sqrt{2}} \right)}{\sqrt{2}} \right] \\ &= - \frac{e^{i\delta} + e^{-i\delta}}{2} \\ &= - \cos(\phi - \theta) \end{aligned}$$

where we've used section V 2.

4. With definitions from section 3, without taking the large N limit,

$$\begin{aligned} \langle +'|X|+' \rangle &= \frac{N-1}{N}, \langle -'|X|-' \rangle = -\frac{N-1}{N} \\ \langle 0'|X|0' \rangle &= 0, \langle 1'|X|1' \rangle = 0 \\ \langle 1'|X|0' \rangle &= \frac{\frac{N-1}{2} + \frac{N}{2}}{2} = \frac{2N-1}{2N} = \langle 0'|X|1' \rangle \\ \langle -'|X|+' \rangle &= \frac{-\langle 1'|X|0' \rangle + \langle 0'|X|1' \rangle}{2} = 0 = \langle +'|X|-' \rangle \\ \langle \Psi | X \otimes X | \Psi \rangle &= \frac{\langle +'|X|+' \rangle \langle -'|X|-' \rangle + \langle -'|X|-' \rangle \langle +'|X|+' \rangle}{2} = \\ &= - \left(\frac{N-1}{N} \right)^2 \end{aligned}$$

$$\begin{aligned} 5. \quad \langle U^\dagger(\phi_i) X U(\phi_i) \otimes U^\dagger(\theta_i) X U(\theta_i) \rangle &= \\ - \left(\frac{N-1}{N} \right)^2 \cos(\phi_i - \theta_i) \end{aligned}$$

Proof: We start with defining $\phi \equiv \phi_i$, $\theta \equiv \theta_i$, $\delta \equiv \phi - \theta$, $\delta' \equiv \delta/2$. Next, we note that $\text{LHS} = \langle \Psi' | X \otimes X | \Psi' \rangle$

where $|\Psi'\rangle = U(\phi_i) \otimes U(\theta_i) |\Psi\rangle$.

$$\begin{aligned}
|\Psi'\rangle &= \frac{e^{i\delta'}}{\sqrt{2}} \left(\frac{|+\rangle - |-\rangle}{\sqrt{2}} \right) \left(\frac{|+\rangle + |-\rangle}{\sqrt{2}} \right) \\
&\quad - \frac{e^{-i\delta'}}{\sqrt{2}} \left(\frac{|+\rangle + |-\rangle}{\sqrt{2}} \right) \left(\frac{|+\rangle - |-\rangle}{\sqrt{2}} \right) \\
&= \frac{e^{i\delta'}}{2\sqrt{2}} (|++\rangle + |+-\rangle - |-+\rangle - |--\rangle) \\
&\quad - \frac{e^{-i\delta'}}{2\sqrt{2}} (|++\rangle - |+-\rangle + |-+\rangle - |--\rangle) \\
&= \frac{e^{i\delta'} - e^{-i\delta'}}{2\sqrt{2}} |++\rangle + \frac{e^{i\delta'} + e^{-i\delta'}}{2\sqrt{2}} |+-\rangle \\
&\quad - \left(\frac{e^{i\delta'} + e^{-i\delta'}}{2\sqrt{2}} \right) |-+\rangle - \left(\frac{e^{i\delta'} - e^{-i\delta'}}{2\sqrt{2}} \right) |--\rangle
\end{aligned}$$

Now using section V 4, we have

$$\begin{aligned}
\text{LHS} &= \langle \Psi' | X \otimes X | \Psi' \rangle \\
&= \frac{1}{2} \left(\frac{N-1}{N} \right)^2 \left[\left| \frac{e^{i\delta'} - e^{-i\delta'}}{2} \right|^2 \right. \\
&\quad \left. - \left| \frac{e^{i\delta'} + e^{-i\delta'}}{2} \right|^2 - \left| \frac{e^{i\delta'} + e^{-i\delta'}}{2} \right|^2 \right. \\
&\quad \left. + \left| \frac{e^{i\delta'} - e^{-i\delta'}}{2} \right|^2 \right] \\
&= - \left(\frac{N-1}{N} \right)^2 \frac{1}{2} [2(\cos^2 \delta/2 - \sin^2 \delta/2)] \\
&= - \left(\frac{N-1}{N} \right)^2 \cos(\delta)
\end{aligned}$$

6. Action of \hat{x}_{mod2L} can be defined explicitly

Proof: $\hat{x}_{mod2L} \equiv \int dx' x'_{mod2L} |x'\rangle \langle x'|$. To arrive at this more carefully, consider the operator $e^{i\hat{x}\frac{2\pi}{2L}}$. Note

that $e^{i\hat{x}\frac{2\pi}{2L}} |x'\rangle = e^{ix'\frac{2\pi}{2L}} |x'\rangle = e^{ix'_{mod2L}\frac{2\pi}{2L}} |x'\rangle$. Thus, $\hat{x}_{mod2L} |x'\rangle = x'_{mod2L} |x'\rangle$, consequently on the most general state $|f\rangle \equiv \int dx' f_{x'} |x'\rangle$ then, we'd have $\hat{x}_{mod2L} |f\rangle = \int dx' f_{x'} x'_{mod2L} |x'\rangle$.

Remark: One needn't necessarily consider eigenstates of \hat{x} to define the action. Eigenstates of $e^{i\hat{x}\frac{2\pi}{2L}}$ maybe considered instead; they can be expressed as (a) $|\varphi\rangle$, s.t. $\langle p + \frac{h}{2L} | \varphi \rangle = \langle p | \varphi \rangle$, $\forall p \in \mathbb{R}$ or (b) $|\bar{x}\rangle \propto \sum_{n \in \mathbb{Z}} |\bar{x} + n2L\rangle$. Using the second expression, we have $e^{i\hat{x}\frac{2\pi}{2L}} |\bar{x}\rangle = e^{i\bar{x}\frac{2\pi}{2L}} |\bar{x}\rangle$. Thus on a more general state, $|c\rangle \equiv \int_0^{2L} d\bar{x} c_{\bar{x}} |\bar{x}\rangle$, we have $\hat{x}_{mod2L} |c\rangle = \int_0^{2L} d\bar{x} c_{\bar{x}} \bar{x} |\bar{x}\rangle$.

7. Physical implementation with electrons is also possible

If we can show that the basic components used to describe the photon setup can be translated to the electron setup, then in principle we are through. (a) Glass slab: The equivalent is the electric AB effect. We need to simply put a capacitor after the slit and the two components will pick up a phase difference. (b) Polariser: The Stern Gerlach setup is the classic analogue. We simply block the orthogonal component. (c) Birefringent crystal: This is slightly tricky. It can be modelled by using a combination of gradient of magnetic field (as in Stern Gerlach) and a capacitor. We start with an equivalent superposition of spin states, $\frac{|\uparrow\rangle - |\downarrow\rangle}{\sqrt{2}} |++\rangle$. To construct the spin dependent $|-'\rangle$ state, we use the magnetic field gradient to spatially separate the $|\uparrow\rangle$ and $|\downarrow\rangle$ states. We place capacitors as described at the spatial position corresponding to $|\downarrow\rangle$ say. Thereafter, we remove the magnetic field gradient and allow the beams to meet again. This will effectively act as a Birefringent crystal, since the phase difference is spin dependent.

-
- [1] that are often chosen to be sufficiently far away that their measurements don't influence each other
 - [2] Note that (ii) is equivalent to measuring along any direction. Details in the Appendix.
 - [3] These values are motivated by the $\pi/2$ between z and x , & $\pi/4$ between z and z' in the conventional Bell case.
 - [4] so that $\phi - \theta \rightarrow \phi' - \theta' = \phi - \theta + \pi = (\phi + \pi/2) - (\theta - \pi/2)$

- [5] Defn: $|0\rangle, |1\rangle$ are s.t. $\hat{z}|0\rangle = |0\rangle$ and $\hat{z}|1\rangle = -|1\rangle$
- [6] We emphasise that the kets we construct, will not be a countable superposition of eigenstates of p or x since these are highly idealized and strictly, not even a part of the Hilbert space.
- [7] \hat{x}_{mod2L} is well defined, see section V 6