



Quantum Measurements and Relativity

Kishor Bharti

Centre for Quantum Technologies. Singapore

October 23, 2016

- Quantum mechanics not compatible with Relativity

- Quantum mechanics not compatible with Relativity
- What if it does?

- Quantum mechanics not compatible with Relativity
- What if it does? \implies Paradox


- Quantum mechanics not compatible with Relativity
- What if it does? \implies Paradox
 - What is the paradox?

- According to QM, measurement \Rightarrow instantaneous collapse

- According to QM, measurement \implies instantaneous collapse
- E.g.: Particle in momentum eigenstate

Collapse and Relativity


- According to QM, measurement \implies instantaneous collapse
- E.g.: Particle in momentum eigenstate



- In another frame of reference?

Collapse and Relativity

- In another frame of reference?




- What if particle confined in past light cone?

- What if particle confined in past light cone?
 - No longer momentum eigenstate

Collapse and Relativity

- What if particle confined in past light cone?
 - No longer momentum eigenstate



Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

- Who collapses state first?

Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

- Who collapses state first?
 - If Alice measures first, \exists inertial frame s.t. Bob measures first

Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$


- Who collapses state first?
 - If Alice measures first, \exists inertial frame s.t. Bob measures first
- Collapse at intersection of light cone?

Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

- Who collapses state first?
 - If Alice measures first, \exists inertial frame s.t. Bob measures first
- Collapse at intersection of light cone?




Collapse and Relativity

- E.g.: 2 distinguishable particle singlet state, space-like separated

$$|\Psi_{-}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

- Who collapses state first?
 - If Alice measures first, \exists inertial frame s.t. Bob measures first
- Collapse at intersection of light cone?



Collapse happens in the past
 \Rightarrow A, B measures product state \Rightarrow Obeys Bell's inequality
 \Rightarrow Paradox

Landau and Peierls:

Theorem (Relativistic Uncertainty Relation)

Measurement of momentum cannot be instantaneous. Momentum with accuracy Δp cannot take time less than Δt such that

$$\Delta p \Delta t \geq \frac{\hbar}{c}.$$

Landau and Peierls:

Theorem (Relativistic Uncertainty Relation)

Measurement of momentum cannot be instantaneous. Momentum with accuracy Δp cannot take time less than Δt such that

$$\Delta p \Delta t \geq \frac{\hbar}{c}.$$

- Measurement of non-local operator cannot be instantaneous.

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle .$$

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle .$$

- Suppose A and B can measure S^2 instantaneously.

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle .$$

- Suppose A and B can measure S^2 instantaneously.
- Rewrite state in total spin basis:

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle ,$$

where

$$|0, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) \quad \text{and} \quad |2, 0\rangle = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle) .$$

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:
 1. A and B measure S^2

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:

1. A and B measure S^2
2. A measures $S_z \implies \frac{1}{2}$ probability get \uparrow or \downarrow

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:
 1. A and B measure S^2
 2. A measures $S_z \implies \frac{1}{2}$ probability get \uparrow or \downarrow
- Measurement 2:

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:

1. A and B measure S^2
2. A measures $S_z \implies \frac{1}{2}$ probability get \uparrow or \downarrow

- Measurement 2:

1. B flips his spin

$$|\Psi'_{\alpha\beta}\rangle = \alpha |2, 1\rangle + \beta |2, -1\rangle$$

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:

1. A and B measure S^2
2. A measures $S_z \implies \frac{1}{2}$ probability get \uparrow or \downarrow

- Measurement 2:

1. B flips his spin

$$|\Psi'_{\alpha\beta}\rangle = \alpha |2, 1\rangle + \beta |2, -1\rangle$$

2. A and B measure S^2

Relativistic Constraint

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- Measurement 1:

1. A and B measure S^2
2. A measures $S_z \implies \frac{1}{2}$ probability get \uparrow or \downarrow

- Measurement 2:

1. B flips his spin

$$|\Psi'_{\alpha\beta}\rangle = \alpha |2, 1\rangle + \beta |2, -1\rangle$$

2. A and B measure S^2
3. A measures $S_z \implies |\alpha|^2$ probability get \uparrow , $|\beta|^2$ get \downarrow

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- B can send superluminal message by flipping in his lab!

E.g.: Measure total spin of 2 localised spin- $\frac{1}{2}$ particles with state

$$|\Psi_{\alpha\beta}\rangle = \frac{\alpha + \beta}{\sqrt{2}} |2, 0\rangle + \frac{\alpha - \beta}{\sqrt{2}} |0, 0\rangle$$

- B can send superluminal message by flipping in his lab!
- Unless $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$.

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measurement interaction

$$\mathcal{H}_{\text{int}} = g(t) [S_z^A P_z^A + S_z^B P_z^B]$$

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measurement interaction

$$\mathcal{H}_{\text{int}} = g(t) [S_z^A P_z^A + S_z^B P_z^B]$$

where

$$g(t) = \begin{cases} \frac{1}{T}, & 0 \leq t \leq T \\ 0, & \text{otherwise} \end{cases}$$

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measurement interaction

$$\mathcal{H}_{\text{int}} = g(t) [S_z^A P_z^A + S_z^B P_z^B]$$

where

$$g(t) = \begin{cases} \frac{1}{T}, & 0 \leq t \leq T \\ 0, & \text{otherwise} \end{cases}$$

- Measuring device state

$$Q_z^A + Q_z^B = 0 = P_z^A - P_z^B.$$

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measures $Q_z^A + Q_z^B \implies$ Measures $S_z = S_z^A + S_z^B$

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measures $Q_z^A + Q_z^B \implies$ Measures $S_z = S_z^A + S_z^B$
- Do this for S_x and S_y too.

Non-local Measurements

What if $|\alpha|^2 = |\beta|^2 = \frac{1}{2}$? \implies state-specific measurement

- Measures $Q_z^A + Q_z^B \implies$ Measures $S_z = S_z^A + S_z^B$
- Do this for S_x and S_y too.
- If $S_x = S_y = S_z$, then $S^2 = 0 \implies$ verified $|0, 0\rangle$

Measurable Non-local Operators

\exists operator measurable instantaneously? \implies Operator-specific measurement

Measurable Non-local Operators

\exists operator measurable instantaneously? \implies Operator-specific measurement

- Suppose \exists such non-degenerate operator W with eigenstate

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle \quad \text{with} \quad \alpha, \beta \in \mathbb{R}.$$

Measurable Non-local Operators

\exists operator measurable instantaneously? \implies Operator-specific measurement

- Suppose \exists such non-degenerate operator W with eigenstate

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle \quad \text{with} \quad \alpha, \beta \in \mathbb{R}.$$

- The other 3 eigenstates are $|W_1\rangle$, $|W_2\rangle$ and $|W_3\rangle$

Measurable Non-local Operators

\exists operator measurable instantaneously? \implies Operator-specific measurement

- Suppose \exists such non-degenerate operator W with eigenstate

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle \quad \text{with} \quad \alpha, \beta \in \mathbb{R}.$$

- The other 3 eigenstates are $|W_1\rangle$, $|W_2\rangle$ and $|W_3\rangle$
- The space is spanned by bases

$$|\Psi_{\alpha\beta}\rangle = \alpha |\uparrow\downarrow\rangle + \beta |\downarrow\uparrow\rangle; \quad |\Psi_{\alpha\beta}^\perp\rangle = \beta |\uparrow\downarrow\rangle - \alpha |\downarrow\uparrow\rangle; \quad |\uparrow\uparrow\rangle; \quad |\downarrow\downarrow\rangle.$$

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\psi_{\alpha\beta}^\perp\rangle$

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously
 - Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously
 - Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
 - Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously
 - Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
 - Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i
 2. A measures S_z^A

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously
 - Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
 - Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i
 2. A measures S_z^A
- For Alice, probability of getting up

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$
 1. A and B measure W instantaneously
 - Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
 - Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i
 2. A measures S_z^A
- For Alice, probability of getting up

$$P(\uparrow_A) = \frac{1 - 2\alpha\beta}{2} \alpha^2 + \frac{1 + 2\alpha\beta}{2} \sum_i |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \beta^2 |\langle W_i | \Psi_{\alpha\beta} \rangle|^2 \right]$$

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$

1. A and B measure W instantaneously

- Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
- Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i

2. A measures S_z^A

- For Alice, probability of getting up

$$P(\uparrow_A) = \frac{1 - 2\alpha\beta}{2} \alpha^2 + \frac{1 + 2\alpha\beta}{2} \sum_i |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \beta^2 |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \right]$$

- For Bob probability of getting up

Measurable Non-local Operators

- Measurement: Prepare state $|0, 0\rangle = \frac{\alpha - \beta}{\sqrt{2}} |\Psi_{\alpha\beta}\rangle + \frac{\alpha + \beta}{\sqrt{2}} |\Psi_{\alpha\beta}^\perp\rangle$

1. A and B measure W instantaneously

- Probability $\frac{|\alpha - \beta|^2}{2}$, getting $|\Psi_{\alpha\beta}\rangle$
- Probability $\frac{|\alpha + \beta|^2}{2}$, getting any of the other eigenstates W_i

2. A measures S_z^A

- For Alice, probability of getting up

$$P(\uparrow_A) = \frac{1 - 2\alpha\beta}{2} \alpha^2 + \frac{1 + 2\alpha\beta}{2} \sum_i |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \beta^2 |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \right]$$

- For Bob probability of getting up

$$P(\uparrow_B) = \frac{1 - 2\alpha\beta}{2} \beta^2 + \frac{1 + 2\alpha\beta}{2} \sum_i |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \alpha^2 |\langle W_i | \Psi_{\alpha\beta}^\perp \rangle|^2 \right]$$

Measurable Non-local Operator

What if Alice and Bob apply local unitary transformations (change relative phase)

Measurable Non-local Operator

What if Alice and Bob apply local unitary transformations (change relative phase)

$$|0, 0\rangle \mapsto |2, 0\rangle$$

Measurable Non-local Operator

What if Alice and Bob apply local unitary transformations (change relative phase)

$$|0, 0\rangle \mapsto |2, 0\rangle$$

- Repeat measurement:

Measurable Non-local Operator

What if Alice and Bob apply local unitary transformations (change relative phase)

$$|0, 0\rangle \mapsto |2, 0\rangle$$

- Repeat measurement:

$$P(\uparrow_A) = \frac{1+2\alpha\beta}{2}\alpha^2 + \frac{1-2\alpha\beta}{2} \sum_i \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \beta^2 \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \right]$$

$$P(\uparrow_B) = \frac{1+2\alpha\beta}{2}\beta^2 + \frac{1-2\alpha\beta}{2} \sum_i \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \alpha^2 \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \right]$$

Measurable Non-local Operator

What if Alice and Bob apply local unitary transformations (change relative phase)

$$|0, 0\rangle \mapsto |2, 0\rangle$$

- Repeat measurement:

$$P(\uparrow_A) = \frac{1+2\alpha\beta}{2}\alpha^2 + \frac{1-2\alpha\beta}{2}\sum_i \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \beta^2 \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \right]$$

$$P(\uparrow_B) = \frac{1+2\alpha\beta}{2}\beta^2 + \frac{1-2\alpha\beta}{2}\sum_i \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \left[|\langle W_i | \uparrow\uparrow \rangle|^2 + \alpha^2 \left| \langle W_i | \Psi_{\alpha\beta}^\perp \rangle \right|^2 \right]$$

- Demand probabilities to be the same

$$\implies \alpha\beta = 0 \quad \text{or} \quad \alpha = \pm\beta$$

Conclusion:

Conclusion:

A and B cannot measure W instantaneously unless each eigenstate of W is $|\Psi_{\alpha\beta}\rangle$ with $\alpha\beta = 0$ or $\alpha = \pm\beta$, up to local unitary transformation.

Conclusion:

A and B cannot measure W instantaneously unless each eigenstate of W is $|\Psi_{\alpha\beta}\rangle$ with $\alpha\beta = 0$ or $\alpha = \pm\beta$, up to local unitary transformation.

- This is necessary but not sufficient. Why? Think of S^2 .

Sufficient Conditions

- What about an operator with an eigenvector that is not entangled ($\alpha\beta = 0$)?

Sufficient Conditions

- What about an operator with an eigenvector that is not entangled ($\alpha\beta = 0$)?
- Can A and B simultaneously measure such an operator?

Sufficient Conditions

- What about an operator with an eigenvector that is not entangled ($\alpha\beta = 0$)?
- Can A and B simultaneously measure such an operator?
- Take two eigenvectors to be $\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ and $|\uparrow\uparrow\rangle$

Sufficient Conditions

- What about an operator with an eigenvector that is not entangled ($\alpha\beta = 0$)?
- Can A and B simultaneously measure such an operator?
- Take two eigenvectors to be $\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ and $|\uparrow\uparrow\rangle$
- Can Alice and Bob instantaneously measure an operator with these two states as eigenstates?

Sufficient Conditions

- If yes

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$
 - This is not an eigenstate of the chosen non-local operator!

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$
 - This is not an eigenstate of the chosen non-local operator!
 - The non-local operator has a chance of leaving the spins in the state $\frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$
 - This is not an eigenstate of the chosen non-local operator!
 - The non-local operator has a chance of leaving the spins in the state $\frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$
 - Now Alice has a chance of obtaining -1 when she measures σ_z

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$
 - This is not an eigenstate of the chosen non-local operator!
 - The non-local operator has a chance of leaving the spins in the state $\frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$
 - Now Alice has a chance of obtaining -1 when she measures σ_z
 - Bob can send superluminal signal to Alice!

Sufficient Conditions

- If yes \implies
 - A and B prepare their spins in the state $|\uparrow\uparrow\rangle$
 - A and B measure instantaneously
 - Alice measures σ_z on her spin and she is certain to obtain $\sigma_z = 1$
 - But, what if just before the measuring the non-local operator, Bob flips his spin?
 - $|\uparrow\uparrow\rangle$ changes to $|\uparrow\downarrow\rangle$
 - This is not an eigenstate of the chosen non-local operator!
 - The non-local operator has a chance of leaving the spins in the state $\frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$
 - Now Alice has a chance of obtaining -1 when she measures σ_z
 - Bob can send superluminal signal to Alice! \implies Alice and Bob can't measure such an operator instantaneously!

Conclusion:

A and B cannot measure W instantaneously unless each eigenstate of W is $|\Psi_{\alpha\beta}\rangle$ with $\alpha = \pm\beta$, up to local unitary transformation.

- Collapse is not Lorentz invariant \Rightarrow

- Collapse is not Lorentz invariant \implies Observers in different frames disagree about collapse

Countering Collapse

- Collapse is not Lorentz invariant \implies Observers in different frames disagree about collapse
- So what ?

Countering Collapse

- Collapse is not Lorentz invariant \implies Observers in different frames disagree about collapse
- So what ?
- This is not a contradiction

Countering Collapse

- Collapse is not Lorentz invariant \implies Observers in different frames disagree about collapse
- So what ?
- This is not a contradiction
- We have a contradiction only if observers in different frames verify incompatible account of collapse

Countering Collapse

- Collapse is not Lorentz invariant \implies Observers in different frames disagree about collapse
- So what ?
- This is not a contradiction
- We have a contradiction only if observers in different frames verify incompatible account of collapse
- Which they can't!

- State specific verifications disturb one another and so verifications are incompatible.

Countering C ollapse

- State specific verifications disturb one another and so verifications are incompatible.
- No contradiction over collapse!

Countering C ollapse

- State specific verifications disturb one another and so verifications are incompatible.
- No contradiction over collapse!
- In fact, disagreement is fine!

Countering C ollapse

- State specific verifications disturb one another and so verifications are incompatible.
- No contradiction over collapse!
- In fact, disagreement is fine!
- Observers in different frames disagree over length, temporal order etc.

- So what?

Countering Collapse

- So what?
- Is everything okay?

Countering Collapse

- So what?
- Is everything okay?
- What about measurements?

Countering Collapse

- So what?
- Is everything okay?
- What about measurements?
- Relativistic causality forbids an instantaneous collapse of almost all non-local operators!

Then what happens in our experiments?

