

MED9098 Homework 1

Autumn 2023

(Due 10/11 18:00)

1. The runtime T of an algorithm depends on its input data, which is characterized by three key parameters: k , m , and n (given the problem data). A simple and standard model that shows how T scales with k , m , and n has the form

$$\hat{T} = \alpha k^\beta m^\gamma n^\delta$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ are constants that characterize the approximate runtime model. Now suppose you are given measured runtimes for N **executions** of the algorithm, with different sets of input data. For each data record, you are given T_i (the runtime), and the parameters k_i, m_i , and n_i .

We wish to find values of α, β, γ , and δ for which our model fits our measurements (T_i). We define the fitting cost as

$$J = \sum_{i=1}^N (\log(\hat{T}_i / T_i))^2$$


where $\hat{T}_i = \alpha k_i^\beta m_i^\gamma n_i^\delta$ is the runtime predicted by our model, using the given parameter values. This fitting cost can be (loosely) interpreted in terms of relative or percentage fit. Please find constants $\alpha, \beta, \gamma, \delta$ that minimize J . (The file **hw1_prob1.csv** (**hw1_prob1.mat**) contains k, m, n, T, N)

- a) Explain how you design $Ax=b$. (Think of what factors are given as measurements, and what variables are unknown. Also, consider how to construct matrix A from the given information). (20 pt)
- b) Give the values of $\alpha, \beta, \gamma, \delta$ that minimize J , and the corresponding value of J . (20 pt)

2. We have a discrete signal $x \in \mathbb{R}^n$, and a measurement $y \in \mathbb{R}^n$, associated with a square matrix $A \in \mathbb{R}^{n \times n}$:

$$y_i = \sum_{j=1}^n a_{ij} x_j + w_i \quad \Rightarrow \quad y = Ax + w = y_{orig} + w$$

The matrix A plays a role as turning the signal x into y (smoothing x), which was perturbed by noise signal w . (The file **hw1_prob2.csv** (**hw1_prob2.mat**) contains A , x , y , y_{orig} (before perturbation)).



- Using inverse of A and pseudo-inverse (least-squares) methods, please find x and compare those with the original discrete signal x . (5 pt)
- Do not panic while obtaining the frustrated results from part (a). Let us take a look at a different approach based on the singular value decomposition (SVD). Take singular value decomposition on A . Plot the diagonal components (singular values of A), and show the 4 input (right) singular vectors (vector v). Explain what you observe in relation to the role of matrix A (smoothing). (15 pt)
- Recall [Low rank approximation] of SVD in Lecture 3.
 - Approximate A with the r largest singular values and corresponding singular vectors.
 - Do the least squares method to find x
 - Repeat (i)-(ii) while varying r from 1 to 50
 Find the value of r that minimizes the norm between reconstructed x and original x . Plot the reconstructed x with original x . (20 pt)

- Recall [Tychonov Regularization] in Lecture 4.

$$\|Ax - b\|_2^2 + \mu \|x\|_2^2$$

Find the value of μ that minimizes the norm between reconstructed x and original x . Plot the reconstructed x with original x . (Hint: Optimal μ that I found was less than 0.1) (20 pt)