

Physics 509 Theory of Measurements Course Notes

Tobias Faehndrich

This document was typeset on September 27, 2025

Introduction:

Notes written at UBC 2025W1 with Dr. Colin Gay.

Contents

1	Foundations of Probability Theory	3
1.1	Motivation: Stochastic Nature of Experimental Data	3
1.2	Probabilistic Interpretation of Experimental Results	3
1.3	Sample Spaces and Stochastic Variables	3
1.4	Events and Set Operations	4
1.5	Kolmogorov's Axioms of Probability	4
1.6	Consequences of the Probability Axioms	5
1.7	Uniform Probability on Finite Sample Spaces	5
1.8	Example: Probability of a Straight in Poker	6
1.9	Conditional Probability	6
2	Bayesian Inference and Random Variables	7
2.1	Bayes' Formula	7
2.2	Law of Total Probability	7
2.3	Independent Events	7
2.3.1	Example: Rolling Two Dice	8
2.4	Random Variables and Probability Distributions	8
2.5	Describing a Distribution	9
2.6	Cumulative Distribution Function (CDF)	9
2.7	Expectation Values	9
2.8	Characteristic Function	9
2.9	Central Moments	10
3	Bayesian Reasoning and Probability Distributions	11
3.1	Bayes Theorem and Its Applications	11
3.2	The Monty Hall Problem: A Bayesian Analysis	11
3.3	Alternate Monty Hall Formulations	12
3.4	Monty Hall Generalized to n Doors	12
3.5	Continuous Probability Distributions and Moments	13
3.6	Discrete Probability Distributions	13
3.7	Cumulative Distribution Functions	14
3.8	Multivariate Distributions and Covariance	14
4	Joint Distributions, Correlations, and Variable Transformations	15
4.1	Conditional Probability: A Simple Example	15
4.2	Distributions of Multiple Random Variables	15
4.3	Covariance Matrix and Correlation Coefficient	15
4.4	Independence and Uncorrelated Variables	16
4.5	Examples of Correlated and Uncorrelated Variables	16

4.6	Marginal Distributions	17
4.7	Change of Variables in Probability Densities	17
4.8	Change of Variables: Non One-to-One Case	18
4.9	Multivariate Transformations and the Jacobian	18
4.10	Example: Cartesian to Polar Transformation	18
5	Propagation of Uncertainty in Measurements	20
5.1	Propagation of Errors for a Single Variable	20
5.2	Variance Propagation for a Single Variable	20
5.3	Propagation of Errors for Multiple Variables	21
5.4	Covariance Propagation for Functions of Multiple Variables	21
5.5	Examples of Error Propagation in Measurements	22
5.6	Matrix Formulation of Linear Error Propagation	22
5.7	Variance of the Arithmetic Mean	22
5.8	Example: Measuring the Period of a Sine Wave	23
6	Covariance Transformations and the Binomial Distribution	24
6.1	Covariance Transformation Under Linear Transformations	24
6.2	The Binomial Distribution	25

1 Foundations of Probability Theory

This lecture covered the course structure and grading.

1.1 Motivation: Stochastic Nature of Experimental Data

- Stochastic processes:
 - muon decay
 - inherent stochasticity
 - quantum mechanics
- Mostly concerned with measurement devices — how do we measure?
- Example: a muon lifetime experiment
 - Take a cosmic muon, detect light, and discriminate.
 - Muon decays into an electron and neutrinos, and the electron produces light.
 - Measure the time between light pulses.
 - Many factors cause noise in the data — results change even if the same mechanism occurs twice.

1.2 Probabilistic Interpretation of Experimental Results

- Experiments are repeated trials.
- Probability (probabilistic interpretation):
 - Results are interpreted as the long-term average of repeating an experiment many times.
 - Example: coin flip

$$P(H) = \lim_{N \rightarrow \infty} \frac{n_H}{N}$$

$n(H)$ = number of heads in N trials

1.3 Sample Spaces and Stochastic Variables

- In modern probability theory:
 - 3 axioms (Kolmogorov)
 - Let X be a stochastic variable.
 - Define sample space S (Ω):

$$S = \{x_1, x_2, \dots\}$$

- Examples:

1. Coin flip:

$$S = \{H, T\}$$

2. Roll a die:

$$S = \{1, 2, 3, 4, 5, 6\}$$

3. Grade in this class:

$$S = \{0, 1, 2, \dots, 100\}$$

4. Decay time of a radioactive atom:

$$S = [0, \infty)$$

- S can be finite (Binomial), countable (Poisson), or infinite (Gaussian, Uniform).

1.4 Events and Set Operations

- Definition: An event E is a subset of S .
- Example: one die roll

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \text{rolling an even number} = \{2, 4, 6\}$$

- Example: E = atom decayed by time t_0

$$S = [0, t_0]$$

- Operations on events:
 - Union (OR) and Intersection (AND)
 - Let A, B be events in S :

$$E = A \cup B = \{e : e \in A \text{ or } e \in B \text{ (or both)}\}$$

- Example: flip a coin twice

$$S = \{HH, HT, TH, TT\}$$

$$A = \text{1st flip is H} = \{HH, HT\}$$

$$B = \text{2nd flip is H} = \{HH, TH\}$$

$$A \cup B = \{HH, HT, TH\}$$

$$A \cap B = \{e \mid e \in A \text{ and } e \in B\} = \{HH\}$$

$$AB = A \cap B$$

$$A^c = \{e \mid e \in S \text{ and } e \notin A\} = \{TH, TT\}$$

- Properties:

- Commutative:

$$A \cup B = B \cup A, \quad AB = BA$$

- Associative:

$$A \cup (B \cup C) = (A \cup B) \cup C, \quad (AB)C = A(BC)$$

- Distributive:

$$(A \cup B)C = AC \cup BC, \quad A(B \cup C) = AB \cup AC$$

- De Morgan's Laws:

$$(A \cup B)^c = A^c B^c, \quad (AB)^c = A^c \cup B^c$$

1.5 Kolmogorov's Axioms of Probability

- A function P on S is a probability measure if it satisfies:

1. $P(S) = 1$
2. $P(\emptyset) = 0$
3. For any countable sequence of disjoint events E_1, E_2, \dots in S :

$$E_i E_j = \emptyset \text{ for } i \neq j$$

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i)$$

1.6 Consequences of the Probability Axioms

•

$$P(\emptyset) = 0$$

Let

$$E_1 = S, \quad E_2 = \emptyset$$

$$E_1 E_2 = \emptyset$$

$$P(S \cup \emptyset) = P(S) + P(\emptyset) = 1 + P(\emptyset)$$

$$P(S) = 1, P(\emptyset) = 0$$

•

$$P(E^c) = 1 - P(E)$$

$$1 = P(S) = P(E \cup E^c) = P(E) + P(E^c)$$

• If $B \subset A$, then:

$$P(B) \leq P(A)$$

$$A = B \cup (B^c A)$$

$$P(A) = P(B \cup (B^c A))$$

$$P(B) = P(A) - P(B^c A) \leq P(A)$$

•

$$P(A \cup B) = P(A) + P(B) - P(AB)$$

If we let the areas of the Venn diagram be 1 (A), 2 (A+B), 3 (B), then:

$$A \cup B = 1 \cup 2 \cup 3$$

$$P(A \cup B) = P(1 \cup 2 \cup 3) = P(1) + P(2) + P(3)$$

$$P(A) = P(1) + P(2), \quad P(B) = P(2) + P(3)$$

$$P(A) + P(B) - P(2) = P(1) + P(2) + P(3) = P(A \cup B)$$

$$\text{equivalently } P(A) + P(B) - P(AB) = P(A) + P(B) - P(AB)$$

1.7 Uniform Probability on Finite Sample Spaces

•

$$E_i = S_i \text{ for } i = 1, 2, \dots, n$$

$$E_i E_j = \emptyset \text{ for } i \neq j$$

$$S = \bigcup_{i=1}^n E_i$$

$$P(S) = 1 = P(\bigcup_{i=1}^n E_i) = \sum_{i=1}^n P(E_i)$$

$$P(E_i) = P(E_j) \quad \text{all equally likely}$$

$$1 = \sum_{i=1}^N P(E_i) = NP(E_i)$$

$$P(E_i) = \frac{1}{N} = P(E_j)$$

$N = |S|$ = number of elements in (cardinality of) S

F be any event (set) in S with k elements $|F| = k$

$$P(F) = P(\bigcup_{S_i \in F} \{E_i\}) = \sum_{i=1}^k P(E_j) = \sum_{i=1}^k \frac{1}{N} = \frac{k}{N} = \frac{|F|}{|S|}$$

1.8 Example: Probability of a Straight in Poker

- Example: 5-card poker hand forming a straight

$$S = \{(AC, 2C, 3C, 4C, 5C), (2C, 3C, 4C, 5C, 6C), \dots\}$$

$$S = \binom{52}{5} = \frac{52!}{5!47!} = 2,598,960$$

- Event = straight = 5 consecutive cards, not of the same suit, any starting card.

$$10(4^5 - 4) = 10200$$

- Starting cards: Ace (A,2,3,4,5), 2 (2,3,4,5,6), ..., 10 (10,J,Q,K,A)
- Not all the same suit: $4^5 - 4$ (exclude all same suit)

$$P(\text{straight}) = \frac{10(4^5 - 4)}{\binom{52}{5}} = 0.00392465$$

1.9 Conditional Probability

- Given 2 events E, F , sample space S :

$$P(E) = \text{probability of a trial from } S \text{ in } E$$

$$P(F) = \text{probability of a trial from } S \text{ in } F$$

- Conditional probability of E given F has occurred:

$$P(E|F) = \text{probability of a trial from } S \text{ in } E, \text{ given the trial is in } F$$

- Note: $P(EF)$ is the probability of a trial from S in both E and F .
- Need to normalize by $P(F)$, so we define:

$$P(E|F) = \frac{P(EF)}{P(F)} \quad \text{if } P(F) > 0$$

$$P(EF) = P(E|F)P(F)$$

- Example: flip a coin 2 times

$$S = \{HH, HT, TH, TT\}$$

Conditional probability of $HH \equiv A$ given:

- First flip = $H \equiv B = \{HH, HT\}$
- Either flip is $H \equiv C = \{HH, HT, TH\}$

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{P(\{HH\})}{P(\{HH, HT\})} = \frac{\frac{1}{4}}{\frac{1}{2}} = \frac{1}{2}$$

$$P(A|C) = \frac{P(AC)}{P(C)} = \frac{P(\{HH\})}{P(\{HH, HT, TH\})} = \frac{\frac{1}{4}}{\frac{3}{4}} = \frac{1}{3}$$

2 Bayesian Inference and Random Variables

2.1 Bayes' Formula

- Let E, F be events:

$$E = EF \cup EF^c$$

$$P(E) = P(EF) + P(EF^c)$$

$$P(E) = P(E|F)P(F) + P(E|F^c)P(F^c)$$

$$P(E) = P(E|F)P(F) + P(E|F^c)(1 - P(F))$$

- Example:** Suppose a blood test is 95% effective in detecting a disease if the person has it. It also has a 1% false positive rate. Suppose 0.5% of the population has the disease.

D = person has disease

E = test is positive

- We want:

$$P(D|E) = \frac{P(ED)}{P(E)}$$

$$\begin{aligned} P(D|E) &= \frac{P(E|D)P(D)}{P(E|D)P(D) + P(E|D^c)(1 - P(D))} \\ &= \frac{0.95 \times 0.005}{0.95 \times 0.005 + 0.01 \times 0.995} = 0.32 \end{aligned}$$

- So even with a positive test, there is only a 32% chance of having the disease.

2.2 Law of Total Probability

- Let $\{F_i\}$ be mutually exclusive events such that:

$$\cup_{i=1}^n F_i = S$$

Then for any event E :

$$E = E \cap (\cup_{i=1}^n F_i) = \cup_{i=1}^n (EF_i)$$

$$P(E) = P(\cup EF_i) = \sum_{i=1}^n P(EF_i) = \sum_{i=1}^n P(E|F_i)P(F_i)$$

2.3 Independent Events

- Generally, $P(E|F) \neq P(E)$.
- If knowing F does not change the probability of E :

$$P(E|F) = \frac{P(EF)}{P(F)} = P(E)$$

$$\boxed{P(EF) = P(E)P(F)}$$

2.3.1 Example: Rolling Two Dice

- Let:

$$E_1 \equiv \text{sum} = 6$$

$$F \equiv \text{first die} = 4$$

$$E_1 : \{(1,5), (2,4), (3,3), (4,2), (5,1)\}$$

$$F : \{(4,1), (4,2), (4,3), (4,4), (4,5), (4,6)\}$$

$$E_1 F = \{(4,2)\}$$

$$P(E_1 F) = \frac{1}{36}$$

$$P(E_1) = \frac{5}{36}$$

$$P(F) = \frac{6}{36} = \frac{1}{6}$$

$$P(E_1)P(F) = \frac{5}{36} \times \frac{1}{6} = \frac{5}{216} \neq P(E_1 F)$$

- Let:

$$E_2 \equiv \text{sum} = 7$$

$$E_2 : \{(1,6), (2,5), (3,4), (4,3), (5,2), (6,1)\}$$

$$E_2 F = \{(4,3)\}$$

$$P(E_2) = \frac{6}{36} = \frac{1}{6}$$

$$P(F) = \frac{1}{6}$$

$$P(E_2 F) = \frac{1}{36}$$

2.4 Random Variables and Probability Distributions

- $S = \{\text{all possible outcomes of stochastic process } X\}$

$x = \text{random variable}$

$S = \text{finite or countable infinite: discrete random variable}$

$S = \text{uncountable infinite: continuous random variable}$

- Continuous case:

$$P(x_0, x_0 + dx) = p(x)dx$$

where $p(x)$ is the probability density function (pdf).

- Discrete case:

$$S = S_i$$

$p_i = \text{probability of } S_i \quad (\text{probability mass function, pmf})$

$$0 \leq P(S_i) \leq 1$$

$$1 = P(S)$$

$$0 \leq p(x)$$

$$\int_{-\infty}^{\infty} p(x)dx = 1$$

2.5 Describing a Distribution

- To describe $p(x)$ in general we specify:
 - **Mode** — peak value of $p(x)$
 - **Median** — 50% cumulative value
 - **Mean** — average value of x weighted by $p(x)$

2.6 Cumulative Distribution Function (CDF)

•

$$F(x) = \int_{-\infty}^x p(x') dx' = P(X \leq x)$$

$$F(-\infty) = 0, \quad F(\infty) = 1$$

2.7 Expectation Values

- Expectation of any function $f(x)$ over $p(x)$:

$$E(f) = \int_{\Omega} f(x) p(x) dx$$

$$E \text{ is a linear operator: } E(af + bg) = aE(f) + bE(g)$$

- Expectation of powers of x :

$$E(x^0) = E(1) = \int 1 \cdot p(x) dx = 1$$

$$E(x^1) = \int x p(x) dx \equiv \mu = \text{mean value of } x$$

$$E(x^2) = \int x^2 p(x) dx \equiv \sigma^2 = \text{variance of } x$$

2.8 Characteristic Function

- The characteristic function of $p(x)$:

$$\varphi(t) = \int_{-\infty}^{\infty} e^{itx} p(x) dx = E(e^{itx})$$

$$\varphi(t) = E \left(1 + itx + \frac{(itx)^2}{2!} + \dots \right)$$

$$= 1 + itE(x) + \frac{(it)^2}{2!} E(x^2) + \dots$$

$$= \sum_{k=0}^{\infty} \frac{(it)^k}{k!} \mu_{k'}$$

- Moments from $\varphi(t)$:

$$\left. \frac{d^n \varphi(t)}{dt^n} \right|_{t=0} = i^n \mu_{n'}$$

2.9 Central Moments

-

$$E((x - \mu)^n) = \int (x - \mu)^n p(x) dx \equiv \mu_n$$
$$\mu = E(x)$$

- 1st central moment:

$$E((x - \mu)^1) = E(x) - E(\mu) = \mu - \mu = 0$$

- 2nd central moment (variance):

$$E((x - \mu)^2) \equiv V(x) = \sigma^2$$

- 3rd central moment (skewness):

$$\text{skewness} = \frac{E((x - \mu)^3)}{\sigma^3}$$

- 4th central moment (kurtosis):

$$\text{kurtosis} = \frac{E((x - \mu)^4)}{\sigma^4} - 3$$

(The -3 ensures that the kurtosis of a normal distribution is 0.)

3 Bayesian Reasoning and Probability Distributions

Tuesday, September 16th 2025

3.1 Bayes Theorem and Its Applications

- Bayes Theorem: for events A and B , we have

$$P(AB) = P(A|B)P(B) = P(B|A)P(A) = P(BA)$$

- Usually it is given in this form:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

- People argued about when you are allowed to use this theorem.

3.2 The Monty Hall Problem: A Bayesian Analysis

- Example: Monty Hall Problem (Game show with host named Monty Hall)
 - There are 3 doors; behind one is a car, behind the other two are goats.
 - You select a door; if the car is behind it, you win.
 - Twist: after you select a door, Monty opens one of the other 2 doors to reveal a goat.
 - Question: stay or switch?
 - Solution: use Bayes theorem.
 - Sample space: $S = \{C_1 = \text{cgg}, C_2 = \text{gcg}, C_3 = \text{ggc}\}$
 - Event 2 = MH opens door 2.
 - Event 3 = MH opens door 3.
 - Number such that your choice is door 1.
 - Take case E_2 , then we want to know $P(C_1|E_2)$.

$$P(C_1|E_2) = \frac{P(E_2|C_1)P(C_1)}{P(E_2)}$$

- $P(C_1) = \frac{1}{3}$
- $P(E_2|C_1) = \frac{1}{2}$ because if the car is behind door 1, Monty can open either door 2 or 3.
- $P(E_2) = \frac{1}{2}$
- Law of total probability:

$$P(E_2) = P(E_2|C_1)P(C_1) + P(E_2|C_2)P(C_2) + P(E_2|C_3)P(C_3) = \frac{1}{2} \cdot \frac{1}{3} + 0 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} = \frac{1}{2}$$

- $P(C_1|E_2) = \frac{P(E_2|C_1)P(C_1)}{P(E_2)} = \frac{\frac{1}{2} \cdot \frac{1}{3}}{\frac{1}{2}} = \frac{1}{3}$
- $P(C_1|E_2) = \frac{1}{3}$
- $P(C_2|E_2) = 0$
- $P(C_3|E_2) = \frac{P(E_2|C_3)P(C_3)}{P(E_2)} = \frac{1 \cdot \frac{1}{3}}{\frac{1}{2}} = \frac{2}{3}$

3.3 Alternate Monty Hall Formulations

- Alternate version: E = MH shows you a goat from $\{2, 3\}$.

- We want to find $P(C_1|E)$.

- $P(C_1|E) = \frac{P(E|C_1)P(C_1)}{P(E)}$

- $P(C_1) = \frac{1}{3}$

- $P(E|C_1) = 1$ because if the car is behind door 1, Monty can open either door 2 or 3.

- $P(E) = 1$ by law of total probability:

$$P(E) = P(E|C_1)P(C_1) + P(E|C_2)P(C_2) + P(E|C_3)P(C_3) = 1 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} = 1$$

- $P(C_1|E) = \frac{1 \cdot \frac{1}{3}}{1} = \frac{1}{3}$

- Another version: What if MH does *not* know where the car is?

- E = MH opens $\{2, 3\}$ and reveals a goat.

- We want to find $P(C_1|E)$.

- $P(C_1|E) = \frac{P(E|C_1)P(C_1)}{P(E)}$

- $P(C_1) = \frac{1}{3}$ because we picked door 1.

- $P(E|C_1) = \frac{1}{2}$ because if the car is behind door 1, Monty can open either door 2 or 3 since he does not know where the car is.

- By law of total probability:

$$P(E) = P(E|C_1)P(C_1) + P(E|C_2)P(C_2) + P(E|C_3)P(C_3) = 1 \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{3} = \frac{2}{3}$$

- $P(C_1|E) = \frac{\frac{1}{3}}{\frac{2}{3}} = \frac{1}{2}$

3.4 Monty Hall Generalized to n Doors

- Now back to the standard version of the problem but with n doors.

- You pick door 1, MH opens any door with a goat behind it from 2 to n ($n - 1$ options).

- $P(C_1|E) = \frac{P(E|C_1)P(C_1)}{P(E)}$

- $P(E) = 1$ because he can always choose a door with a goat behind it (many options and he knows the answers).

- $P(C_1) = \frac{1}{n}$

- $P(E|C_1) = 1$ because if the car is behind door 1, Monty can open any of the other doors.

3.5 Continuous Probability Distributions and Moments

- Continuous probability distribution $p(x)$:
- Moments:

$$E(x^n) = \int_{-\infty}^{\infty} x^n p(x) dx$$

$$\begin{aligned} \text{mean: } & \mu = E(x) \\ \text{variance: } & V(x) = \sigma^2 = E((x - \mu)^2) = E(x^2) - \mu^2 \\ \text{std dev: } & \sigma = \sqrt{\sigma^2} \end{aligned}$$

- Central moments:

$$\begin{aligned} E(x - \mu) &= E(x) - \mu = 0 \\ E((x - \mu)^2) &= \sigma^2 \\ E((x - \mu)^3) &= \text{skewness} \\ E((x - \mu)^4) &= \text{kurtosis} \end{aligned}$$

- Characteristic function:

$$\Phi(t) = E(e^{itx}) = \int_{-\infty}^{\infty} e^{itx} p(x) dx \quad (3.1)$$

$$= \sum_{k=0}^{\infty} \frac{(it)^k}{k!} \mu_k \quad (3.2)$$

$$\Phi_{\mu}(t) = E(e^{it(x-\mu)}) = E(e^{itx})e^{-it\mu} = \Phi(t)e^{-it\mu}$$

$$V(x) = E((x - \mu)^2) \quad (3.3)$$

$$= E(x^2 - 2\mu x + \mu^2) \quad (3.4)$$

$$= E(x^2) - 2\mu E(x) + \mu^2 E(1) \quad (3.5)$$

$$= E(x^2) - 2\mu^2 + \mu^2 \quad (3.6)$$

$$= E(x^2) - \mu^2 = E(x^2) - (E(x))^2 \quad (3.7)$$

3.6 Discrete Probability Distributions

- The discrete case (e.g., rolling a die, picking a card) uses a probability mass function.
- Usually denote outcomes as r :
- p_r = probability of outcome r .
- $\sum_r p_r = 1$
- $E(r) = \sum_r p_r r$ = mean μ
- Variance: $V(r) = \sum_r (r - \mu)^2 p_r = E(r^2) - \mu^2$
- Coin flip example: $S = \{H, T\}$.
- Often map to 0 or 1: $H = 0, T = 1$.
- But in theory you can pick any two numbers a and b to map outcomes, just so you can calculate mean and variance.

$$E(r) = ap_H + bp_T$$

3.7 Cumulative Distribution Functions

- For continuous case:

$$F(x) = \int_{-\infty}^x f(x') dx'$$

- For discrete case:

$$F(r) = \sum_{r' \leq r} p_{r'}$$

- $F(x)$ is the cumulative distribution function (CDF).
- $F(x)$ is non-decreasing, $F(-\infty) = 0$, $F(\infty) = 1$.

3.8 Multivariate Distributions and Covariance

- Distribution of multiple variables:
- Elements belong to real vector space \mathbb{R}^n .
- $P(AB) \dots P(A, B)$
- $p(x_1, x_2, \dots, x_n) \geq 0$ is the joint probability distribution function (PDF).
- $\int_{\Omega} p(\vec{x}) d^n x = 1$
- $E(f(\vec{x})) = \int_{\Omega} f(\vec{x}) p(\vec{x}) d^n x$
- $\mu_i = \int x_i p(\vec{x}) d^n x$
- $V(x_i) = \sigma_i^2 = \int (x_i - \mu_i)^2 p(\vec{x}) d^n x$
- Covariance:
- $V_{i,j} = E((x_i - \mu_i)(x_j - \mu_j))$
- $V_{i,i} = \sigma_i^2 = E((x_i - \mu_i)^2)$ (variance)
- $V_{i,j} = V_{j,i}$ (symmetry)

4 Joint Distributions, Correlations, and Variable Transformations

Thursday, September 18th 2025

4.1 Conditional Probability: A Simple Example

- For fun, example that depends on cultural assumptions: A king comes from a family with two kids. What is the probability that the king's sibling is a sister?
- $S = \{(m, m), (m, f), (f, m), (f, f)\}$
- $P(S|K) = \frac{P(SK)}{P(K)} = \frac{\frac{1}{2}}{\frac{2}{3}} = \frac{3}{4}$

4.2 Distributions of Multiple Random Variables

- $p(x_1, x_2, \dots, x_n)$
- $S = \mathbb{R}^n$
- $\int p(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n = 1$
- For any function $f(\vec{x})$:

$$E(f) = \int f(\vec{x})p(\vec{x})d\vec{x}$$

- $E(x_1) = \int x_1 p(\vec{x})d\vec{x} = \mu_1$
- $E(x_i) = \mu_i$
- $V(x_i) \equiv \sigma_i^2 = \int (x_i - \mu_i)^2 p(\vec{x})d\vec{x}$

4.3 Covariance Matrix and Correlation Coefficient

- Define covariance:

$$V_{ij} = E((x_i - \mu_i)(x_j - \mu_j))$$

- $V_{ii} = \sigma_i^2$ (variance)
- $V_{ij} = V_{ji}$ (symmetry)
- $V_{ij} = 0$ for independent variables
- Expanding the covariance matrix:

$$\begin{aligned} V_{ij}(\vec{x}) &= E((x_i - \mu_i)(x_j - \mu_j)) \\ &= E(x_i x_j - \mu_i x_j - \mu_j x_i + \mu_i \mu_j) \\ &= E(x_i x_j) - \mu_i E(x_j) - \mu_j E(x_i) + \mu_i \mu_j \\ &= E(x_i x_j) - \mu_i \mu_j - \mu_j \mu_i + \mu_i \mu_j \\ &= E(x_i x_j) - \mu_i \mu_j \end{aligned}$$

- So we can say that $V_{ij} \geq 0$
- V_{ij} can be negative, zero, or positive

- Define the correlation coefficient:

$$\rho(x_i, x_j) = \rho_{ij} = \frac{V_{ij}}{\sqrt{V_{ii}}\sqrt{V_{jj}}} = \frac{V_{ij}}{\sigma_i\sigma_j}$$

- We find that $-1 \leq \rho_{ij} \leq 1$

4.4 Independence and Uncorrelated Variables

- Random variables x_1, \dots, x_n are independent if the joint pdf factorizes:

$$p(x_1, \dots, x_n) = p_1(x_1)p_2(x_2) \dots p_n(x_n)$$

- Independent variables are uncorrelated:

$$\begin{aligned} E(x_i x_j) &= \int x_i x_j p(\vec{x}) d\vec{x} \\ &= \int x_i x_j p_1(x_1) \dots p_n(x_n) dx_1 \dots dx_n \\ &= \int x_i p_i(x_i) dx_i \int x_j p_j(x_j) dx_j \int p_2(x_2) dx_2 \dots \int p_n(x_n) dx_n = \mu_i \mu_j \end{aligned}$$

$$V_{ij} = E(x_i x_j) - \mu_i \mu_j$$

In the case of independent variables:

$$V_{ij} = \mu_i \mu_j - \mu_i \mu_j = 0$$

- Independent variables are uncorrelated, but uncorrelated variables are not necessarily independent.

4.5 Examples of Correlated and Uncorrelated Variables

- 100% correlation example:
- $x = \text{Uniform}[-1, 1]$, plot distribution from -1 to 1 .
- $y = x$:
- $V_{ij} = E(xy) - E(x)E(y) = E(x^2) = \int_{-1}^1 x^2 \frac{1}{2} dx = \frac{1}{3} \neq 0$
- $y = |x|$:
- $E(xy) = \int_{-1}^0 x(-x)p(x)dx + \int_0^1 xxp(x)dx$
- $E(xy) = \int_0^1 x^2 \frac{1}{2} dx - \int_{-1}^0 x^2 \frac{1}{2} dx = \frac{1}{6} - \frac{1}{6} = 0$

4.6 Marginal Distributions

- For a joint pdf $p(x_1, x_2, \dots, x_n)$, the marginal probability density functions are:

$$f_1(x_1) = \int p(x_1, x_2, \dots, x_n) dx_2 dx_3 \dots dx_n$$

- If variables are independent:

$$\begin{aligned} f_1(x_1) &= \int p(x_1, x_2, \dots, x_n) dx_2 \dots dx_n \\ &= p_1(x_1) \int p_2(x_2) dx_2 \int p_3(x_3) dx_3 \dots \int p_n(x_n) dx_n \\ &= p_1(x_1) \cdot 1 \cdot 1 \cdot \dots \cdot 1 = p_1(x_1) \end{aligned}$$

4.7 Change of Variables in Probability Densities

- Something we need to know, because we do it all the time:
 - Change of variables of P
 - Calculate new V_{ij} under new variables
- Let x be a random variable with pdf $f(x)$ and let y be some function.
- First: y is one-to-one with f




Figure 4.1: 1-to-1 function

- What is the probability density of y , denoted $g(y)$?
- Conservation of probability:
- $f(x)dx = g(y)dy$
- $g(y) = f(x) \left| \frac{dx}{dy} \right|$

$$\boxed{f(x) \left| \frac{dx}{dy} \right| = g(y)}$$

4.8 Change of Variables: Non One-to-One Case

- If y is not one-to-one: sum over all segments that map to the same y .
- Example: $f(x)$ uniform on $[0, 1]$, $f(x) = 1$
- Let $y(x) = \frac{-1}{\lambda} \ln(x)$
- $\frac{dy}{dx} = \frac{-1}{\lambda x}$
- $\frac{dx}{dy} = -\lambda x$
- $-\lambda x = \ln x$
- $e^{-\lambda y} = x$
- $\lambda > 0 \Rightarrow \frac{dx}{dy} = -\lambda x = -\lambda e^{-\lambda y}$
- $g(y) = f(x) \left| \frac{dx}{dy} \right| = 1 \cdot \lambda e^{-\lambda y} = \lambda e^{-\lambda y}$

4.9 Multivariate Transformations and the Jacobian

- If we have variables $\{x_i\}$ and transform to new variables $\{y_i\}$:
- Region \mathbb{R} in x -space maps to region \mathbb{R}' in y -space.

$$\int_{\mathbb{R}} f(\vec{x}) d\vec{x} = \int_{\mathbb{R}'} f(\vec{x}(\vec{y}))(\vec{y}) \left| \frac{\partial \vec{x}}{\partial \vec{y}} \right| d\vec{y}$$

$$g(\vec{y}) = f(\vec{x}(\vec{y})) |J|$$

- Where $\left| \frac{\partial \vec{x}}{\partial \vec{y}} \right|$ is the Jacobian determinant of the transformation.
- Jacobian matrix J :

$$J = \begin{bmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} & \cdots & \frac{\partial x_2}{\partial y_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \frac{\partial x_n}{\partial y_2} & \cdots & \frac{\partial x_n}{\partial y_n} \end{bmatrix}$$

4.10 Example: Cartesian to Polar Transformation

- Change to polar coordinates:
- $x = r \cos \theta$
- $y = r \sin \theta$
- $P'(r, \theta) = ? = p(x, y) \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right|$
- $\frac{\partial x}{\partial r} = \cos \theta$
- $\frac{\partial y}{\partial r} = \sin \theta$

- $\frac{\partial x}{\partial \theta} = -r \sin \theta$
- $\frac{\partial y}{\partial \theta} = r \cos \theta$
- $J = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix}$
- $J = r \cos^2 \theta + r \sin^2 \theta = r$
- $p'(r, \theta) = \frac{r}{\pi} dr d\theta$

5 Propagation of Uncertainty in Measurements

Tuesday, September 20th 2025

5.1 Propagation of Errors for a Single Variable

- Given $f(x)$ pdf, $\mu \equiv E(x)$, $\sigma^2 \equiv V(x) = E(x^2) - \mu^2$
- Know $f(x) \rightarrow g(y)$, given $y(x)$.
- Taylor expand $y(x)$ about mean μ :

$$y(x) = y(\mu) + y'(\mu)(x - \mu) + \frac{1}{2!}y''(\mu)(x - \mu)^2 + \dots$$

$$E(y(x)) \equiv \mu_y$$

$$\begin{aligned} E(y(x)) &= E(y(\mu)) + y'(\mu)E(x - \mu) + \frac{1}{2!}y''(\mu)E((x - \mu)^2) + \dots \\ &= y(\mu) + y'(\mu) \cdot 0 + \frac{1}{2!}y''(\mu)V(x) + \dots \end{aligned}$$

- To the 1st order:

$$\mu_y = E(y(x)) = y(\mu) = y(E(x))$$

5.2 Variance Propagation for a Single Variable

- Variance of y :

$$V(y) = E((y(x) - E(y(x))))^2 \quad (5.1)$$

$$= E((y(x) - \mu_y)^2) \quad (5.2)$$

$$= E((y'(\mu)(x - \mu) + \frac{1}{2!}y''(\mu)(x - \mu)^2 + \dots)^2) \quad (5.3)$$

$$= E(y'(\mu)^2(x - \mu)^2 + y'(\mu)y''(\mu)(x - \mu)^3 + O((x - \mu)^4)) \quad (5.4)$$

$$= y'(\mu)^2V(x) + \dots \quad (5.5)$$

- Some relations:

$$E(x) \equiv \mu_x$$

$$V(x) \equiv \sigma_x^2$$

$$y = y(x)$$

$$E(y) \equiv \mu_y = y(\mu_x)$$

$$V(y) \equiv \sigma_y^2 = (y'(\mu_x))^2\sigma_x^2$$

$$\sigma_y = |y'(\mu_x)|\sigma_x$$

- Example: $y = \frac{1}{x}$, $\frac{dy}{dx} = -\frac{1}{x^2}$

$$\sigma_y^2 = \frac{1}{\mu_x^4}\sigma_x^2$$

5.3 Propagation of Errors for Multiple Variables

- Let us suppose we have n variables $\{x_i\}$, with pdf $f(\vec{x})$.
- Let $y_j = 1, 2, \dots, m$ be m functions of x_i .
- $y_j = y_j(x_1, x_2, \dots, x_n)$
- $V_{ij}(x)_{n \times n}(\vec{x}) = \text{covariance matrix of } \{x_i\}$
- $V_{ij}(\vec{x}) = E((x_i - \mu_i)(x_j - \mu_j))$
- Taylor expand each y_j : $\vec{\mu} = (\mu_1, \mu_2, \dots, \mu_n)$
- $y_j(\vec{x}) = y_j(\vec{\mu}) + \sum_i \frac{\partial y_j}{\partial x_i} \Big|_{\vec{\mu}} (x_i - \mu_i) + \frac{1}{2!} \sum_{i,k} \frac{\partial^2 y_j}{\partial x_i \partial x_k} \Big|_{\vec{\mu}} (x_i - \mu_i)(x_k - \mu_k) + \dots$
- $E(y_j(\vec{x})) = E(y_j(\vec{\mu})) + \sum \frac{\partial y_j}{\partial x_i} E(x_i - \mu_i) + \dots = y_j(\vec{\mu})$

5.4 Covariance Propagation for Functions of Multiple Variables

- Covariance between y_k and y_l :

$$\begin{aligned}
 & E((y_k - \mu_{y_k})(y_l - \mu_{y_l})) \\
 &= E((y_k - y_k(\mu))(y_l - y_l(\mu))) \\
 &= E\left(\sum_i \frac{\partial y_k}{\partial x_i} \Big|_{\mu} (x_i - \mu_i) \sum_j \frac{\partial y_l}{\partial x_j} \Big|_{\mu} (x_j - \mu_j)\right) \\
 &= \sum_{i,j} \frac{\partial y_k}{\partial x_i} \Big|_{\mu} \frac{\partial y_l}{\partial x_j} \Big|_{\mu} E((x_i - \mu_i)(x_j - \mu_j))
 \end{aligned}$$

$$\boxed{V_{kl}(\vec{y})_{m \times m} = \sum_{i,j} \frac{\partial y_k}{\partial x_i} \Big|_{\vec{\mu}} \frac{\partial y_l}{\partial x_j} \Big|_{\vec{\mu}} V_{ij}(\vec{x})_{n \times n}}$$

- Example: x, y random variables,

$$V(x, y) = \begin{bmatrix} V_{xx} & V_{xy} \\ V_{yx} & V_{yy} \end{bmatrix} = \begin{bmatrix} \sigma_x^2 & \rho_{xy} \sigma_x \sigma_y \\ \rho_{xy} \sigma_x \sigma_y & \sigma_y^2 \end{bmatrix}$$

- $z = x + y$
- $V(z) = \sigma_z^2 = \left(\frac{\partial z}{\partial x}\right)^2 V_{xx} + 2 \frac{\partial z}{\partial x} \frac{\partial z}{\partial y} V_{xy} + \left(\frac{\partial z}{\partial y}\right)^2 V_{yy}$
- $= \sigma_x^2 + 2\rho_{xy} \sigma_x \sigma_y + \sigma_y^2$
- If x_i are uncorrelated,

$$\begin{aligned}
 V_{ij} &= \sigma_{i,j} \sigma_i^2 = \begin{bmatrix} \sigma_1^2 & 0 \\ 0 & \sigma_2^2 \end{bmatrix} \\
 V_{kl}(\vec{y}) &= \sum_i \frac{\partial y_k}{\partial x_i} \Big|_{\mu} \frac{\partial y_l}{\partial x_i} \Big|_{\mu} V_{ii}(\vec{x}) \\
 \text{variance } V_{kk} &= \sum_i \left(\frac{\partial y_k}{\partial x_i}\right)^2 \sigma_i^2
 \end{aligned}$$

5.5 Examples of Error Propagation in Measurements

- Example: Measuring resistances. x_i independent, $z = x + y$, $x = R_1$ resistor value, $y = R_2$ resistor value, $z = R_{\text{tot}}$ total resistance.
- $R_1 \pm \sigma_{R_1}$
- Convention is to use $\sqrt{V(R)}$ as uncertainty.
- For a good measuring device, $E(R) = R_{\text{true}} \leftarrow$ unbiased.
- $V(R) = \text{small}$
- $R_1 \pm \sigma_{R_1}$, $R_2 \pm \sigma_{R_2}$, then $\sigma_{R_{\text{tot}}} = \sqrt{\sigma_{R_1}^2 + \sigma_{R_2}^2}$
- $R = R_{\text{tot}} = R_1 + R_2$
- $z = xy$, like I, R
- $\sigma_z^2 = \left(\frac{\partial z}{\partial x}\right)^2 \sigma_x^2 + \left(\frac{\partial z}{\partial y}\right)^2 \sigma_y^2 = y^2 \sigma_x^2 + x^2 \sigma_y^2$

$$\left(\frac{\sigma_z}{z}\right)^2 = \left(\frac{\sigma_x}{x}\right)^2 + \left(\frac{\sigma_y}{y}\right)^2$$

5.6 Matrix Formulation of Linear Error Propagation

- Formula is exact if transformation of variables is linear.
- $\vec{y} = A\vec{x}$, A is $m \times n$ matrix, \vec{x} is $n \times 1$, \vec{y} is $m \times 1$.
- $\frac{\partial y_k}{\partial x_i} = \text{constant} \Rightarrow$ higher order terms in Taylor expansion are 0
- $V_{kl}(\vec{y}) = \sum_{i,j} \frac{\partial y_k}{\partial x_i} \frac{\partial y_l}{\partial x_j} V_{ij}(\vec{x})$
- Matrix notation:
- $V_{kl}(\vec{y}) = \sum_{i,j} A_{ki} A_{lj} V_{ij}(\vec{x})$
- $= \sum_{i,j} A_{ki} V_{ij}(\vec{x}) A_{lj}$
- $= \sum_{i,j} A_{ki} V_{ij} (A^T)_{jl}$
- $= (AV(\vec{x})A^T)_{kl}$

$$V(\vec{y})_{m \times m} = A_{m \times n} V(\vec{x})_{n \times n} A_{n \times m}^T$$

5.7 Variance of the Arithmetic Mean

- Example: Arithmetic mean. Let $x_i = n$ identical independent variables with $V(x_i) = \sigma_x^2$
- Set $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
- Recall that $V(ax) = a^2 V(x)$
- $V(\bar{x}) = V\left(\frac{1}{n} \sum_{i=1}^n x_i\right) = \frac{1}{n^2} V\left(\sum_{i=1}^n x_i\right) = \frac{1}{n^2} \sum_{i=1}^n V(x_i) = \frac{1}{n^2} n \sigma_x^2 = \frac{\sigma_x^2}{n}$

- If variables are different σ_i^2 : n measurements
- $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
- $V(\bar{x}) = \frac{1}{n^2} \sum_{i=1}^n V(x_i) = \frac{1}{n^2} \sum_{i=1}^n \sigma_i^2$
- $\sigma_{\bar{x}} = \frac{1}{n} \sqrt{\sum_{i=1}^n \sigma_i^2}$

5.8 Example: Measuring the Period of a Sine Wave

- Example: Measure period of sine wave on scope.
- $T = \Delta t = t_2 - t_1$
- $\sigma_T^2 = \left(\frac{\partial \Delta t}{\partial t_1} \right)^2 \sigma_t^2 + \left(\frac{\partial \Delta t}{\partial t_2} \right)^2 \sigma_t^2 = \sigma_t^2 + \sigma_t^2 = 2\sigma_t^2$
- Measure N cycles, $T = \frac{1}{N} \Delta t$
- $\sigma_{T^2} = \frac{1}{N^2} \sigma_{\Delta t}^2 = \frac{2}{N^2} \sigma_t^2$

6 Covariance Transformations and the Binomial Distribution

Tuesday, September 25th 2025

6.1 Covariance Transformation Under Linear Transformations

- Linear transformation:

$$\vec{y} = A\vec{x}$$

$$V_{kl}(\vec{y}) = \sum_{i,j} \frac{\partial y_k}{\partial x_i} \frac{\partial y_l}{\partial x_j} V_{ij}(\vec{x})$$

- Linear $y_k = \sum A_{kj}x_j$

- then

$$V_{kl}(\vec{y}) = \sum_{i,j} A_{ki} A_{lj} V_{ij}(\vec{x})$$

- or in matrix form

$$V(\vec{y}) = \left(AV(\vec{x})A^T \right)_{kl}$$

Diagonalization via Eigenvectors

- If \hat{e}_i are the eigenvectors of V , then

$$V(\vec{x})\hat{e}_i = \lambda_i \hat{e}_i$$

- Form:

$$A = \begin{pmatrix} \hat{e}_1 \\ \dots \\ \hat{e}_n \end{pmatrix} = \begin{pmatrix} \hat{e}_{11} & \hat{e}_{12} & \dots & \hat{e}_{1n} \\ \dots & \dots & \dots & \dots \\ \hat{e}_{n1} & \hat{e}_{n2} & \dots & \hat{e}_{nn} \end{pmatrix}$$

- then

$$A^T A = I$$

- then:

$$VA^T = V \begin{pmatrix} \hat{e}_1 & \dots \\ \dots & \dots \\ \hat{e}_n & \dots \end{pmatrix} = \begin{pmatrix} \lambda_1 \hat{e}_{11} & \dots & \lambda_n \hat{e}_{n1} \\ \dots & \dots & \dots \\ \lambda_1 \hat{e}_{1n} & \dots & \lambda_n \hat{e}_{nn} \end{pmatrix}$$

- Then:

$$AVA^T = \begin{pmatrix} \hat{e}_{11} & \dots & \hat{e}_{1n} \\ \dots & \dots & \dots \\ \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} \lambda_1 \hat{e}_{11} & \dots \\ \dots & \dots & \dots \\ \lambda_1 \hat{e}_{1n} & \dots \end{pmatrix} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

- Then:

$$AVA^T = V(\vec{y}) = \begin{pmatrix} \sigma_1^2 & 0 & \dots & 0 \\ 0 & \sigma_2^2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \sigma_n^2 \end{pmatrix}$$

6.2 The Binomial Distribution

- Consider an experiment with two outcomes.
- E.g. coin flips, selecting a ball with 2 possible colours, etc.
- One trial is called a Bernoulli trial.

Bernoulli Trials and Sampling Methods

- Example – Method 1: You have an urn filled with N balls. Some are red (R), some are blue (B).
- (0) What is your estimate of n_R , n_B , or $f = n_R/N$ or p of drawing R?
- (1) You pick a ball: R. Q: estimate of $p = n_R/N$?
- (2) You pick another without replacing 1st ball: get R.
- (3) R
- (4) Get B
- This is a question about this ONE urn.
- Now Method 2: you draw red, and you PUT IT BACK. You repeat this several times.
- Now Method 3: We have an infinite source of balls with fraction p red and $(1 - p)$ blue.

$$\begin{aligned}P(R) &= p \\P(B) &= 1 - p\end{aligned}$$

Derivation of the Binomial Probability

- Make infinite number of urns all with N balls, with fraction p red and $(1 - p)$ blue.
- Open all, count n_R red balls, n_B blue balls.
- In our case we have N balls, prob $p = R$ and $1 - p = q = B$.
- Prob of getting sequence RRB is:

$$P(RRB) = p \cdot p \cdot (1 - p) = p^2(1 - p)$$

- If we don't care about order, then:

$$P(RRB) = P(RBR) = P(BRR) = p^2(1 - p)$$

- There are 3 ways of ordering RRB, so total probability is:

$$P(2R, 1B) = 3p^2(1 - p) = 3p^2q$$

- Number of ways to choose r items from N is:

$$\binom{N}{r} = \frac{N!}{r!(N - r)!}$$

- Probability of getting exactly r R out of N :

$$P_r = \binom{N}{r} p^r (1 - p)^{N - r} = B(r; N, p)$$

- This is called the Binomial distribution and applies to anything where there are 2 outcomes (A, \bar{A}).

Mean and Variance of the Binomial Distribution

- Want mean, σ

$$\begin{aligned}
 E(r) &= \sum_{r=0}^n r P_r = \sum_{r=0}^n r \binom{n}{r} p^r (1-p)^{n-r} \\
 &= \sum_{r=0}^n r \frac{n!}{r!(n-r)!} p^r (1-p)^{n-r} \\
 &= \sum_{r=1}^n \frac{n!}{(r-1)!(n-r)!} p^r (1-p)^{n-r} \\
 &= np \sum_{r=1}^n \frac{(n-1)!}{(r-1)!(n-r)!} p^{r-1} (1-p)^{n-r}
 \end{aligned}$$

- Change sum $r' = r - 1 \rightarrow n' = n - 1$

$$\begin{aligned}
 E(r) &= np \sum_{r'=0}^{n-1} \frac{(n-1)!}{r'!(n-1-r')!} p^{r'} (1-p)^{(n-1)-r'} \\
 &= np \sum_{r'=0}^{n-1} \binom{n-1}{r'} p^{r'} (1-p)^{n-1-r'} \\
 &= np \cdot 1 = np
 \end{aligned}$$

- from:

$$\begin{aligned}
 (p+q)^n &= \sum_{r=0}^n \binom{n}{r} p^r q^{n-r} \\
 (p+1-q)^n &= 1^n = 1 \\
 E(r) &= np
 \end{aligned}$$

- This is what we want!
- Now:

$$V(r) = \sum r^2 p_r - E(r)^2 = \sum r^2 p_r - n^2 p^2$$

- Slightly easier to calculate:

$$\begin{aligned}
 \sum r(r-1) p_r &= \sum_{r=0}^n r(r-1) \frac{n!}{r!(n-r)!} p^r (1-p)^{n-r} = \sum_{r=2}^n \frac{n!}{(r-2)!(n-r)!} p^r q^{n-r} \\
 &= n(n-1) p^2 \sum_{r=2}^n \frac{(n-2)!}{(r-2)!(n-r)!} p^{r-2} q^{n-r}
 \end{aligned}$$

- Sub $r' = r - 2$

$$\begin{aligned}
 &= n(n-1) p^2 \sum_{r'=0}^{n-2} \binom{n-2}{r'} p^{r'} q^{(n-2)-r'} \\
 &= n(n-1) p^2 \cdot 1 = n(n-1) p^2 \\
 &= n^2 p^2 - np^2
 \end{aligned}$$

- Such that:

$$\boxed{V(r) = \sum r^2 p_r - n^2 p^2} = np(1-p) = npq$$

Applications to Histograms and Counting Statistics

- Why is this important? Histograms are often Binomially distributed.
- Data either falls A : falls in bin, or \bar{A} : does not fall in bin.
- p = probability of falling in i th bin.
- $+n$ entries, e.g. students in class, histogram = grades.
- Expected number of entries is np .
- Plot of taking distribution several times and checking how many fall in bin i and then plotting that distribution is Binomial.
- Usually you have 1 histogram.
- Look at entries in bin i – n_i/n = fraction of entries in bin i .
- Estimator $p = n_i/n$.
- Expect if you repeated $\Rightarrow n_i$ would follow Binomial distribution with mean $\sigma_i = \sqrt{npq}$ and $V_i = \sigma_i^2 = np(1-p)$.

$$p = \frac{n_i}{n}$$

$$\sigma_i = \sqrt{n_i \left(1 - \frac{n_i}{n}\right)} \approx \sqrt{n_i} \text{ if } n \gg n_i$$

- Notes: we do know the total number n , how often is it in bin i .
- HW: given the distribution, how many times n do I need to do it to get that.
- r fixed n , vs. n fixed r .