

# Arithmetic Quantum Mechanics

## Lab Book

Tobias J. Osborne

Campaign log — compiled September 8, 2026

## Contents

|          |                                                                                                   |           |
|----------|---------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Overview</b>                                                                                   | <b>4</b>  |
| <b>2</b> | <b>The symplectic object of a finite field, and its twisted operator algebra</b>                  | <b>7</b>  |
| 2.1      | The symplectic space of a finite field . . . . .                                                  | 7         |
| 2.2      | Polarizing cocycles: the datum $\beta$ , introduced . . . . .                                     | 7         |
| 2.3      | The phase datum . . . . .                                                                         | 8         |
| 2.4      | Weyl operators, the twisted algebra, and the Heisenberg group . . . . .                           | 8         |
| 2.5      | The form is bilinear, alternating, nondegenerate, and its isometries . . . . .                    | 9         |
| 2.6      | Weyl relations, uniformly in the characteristic and for every choice of cocycle . .               | 10        |
| 2.7      | Simplicity of the twisted algebra . . . . .                                                       | 11        |
| <b>3</b> | <b>Lagrangian lines, Schrödinger models, and the uniqueness of the finite Weyl representation</b> | <b>12</b> |
| 3.1      | Preliminary facts about finite abelian characters and cocycles . . . . .                          | 13        |
| 3.2      | Lagrangian lines and the algebra they generate . . . . .                                          | 13        |
| 3.3      | The characteristic-two phase phenomenon . . . . .                                                 | 15        |
| 3.4      | Uniqueness of the finite Weyl representation . . . . .                                            | 16        |
| <b>4</b> | <b>The polarizing cocycle as a second datum: the characteristic-two dichotomy</b>                 | <b>18</b> |
| 4.1      | The torsor of admissible polarizing cocycles, restated . . . . .                                  | 18        |
| 4.2      | Odd characteristic: a canonical representative . . . . .                                          | 19        |
| 4.3      | Characteristic two: no symmetric distinguished point . . . . .                                    | 20        |
| 4.4      | The quadratic form of a cocycle, and the order profile of its Heisenberg group . .                | 20        |
| 4.5      | The Arf classification . . . . .                                                                  | 21        |
| 4.6      | The type, detected inside the quantum system . . . . .                                            | 23        |
| 4.7      | The level statement: what the algebra forgets, and what it does not . . . . .                     | 24        |
| <b>5</b> | <b>Symmetry of the Weyl frame: the Weil representation and its splitting problem</b>              | <b>25</b> |
| 5.1      | The Weyl frame and its two automorphism groups . . . . .                                          | 26        |
| 5.2      | Exactness of the frame-automorphism extension, uniformly in the characteristic .                  | 26        |
| 5.3      | Splitting at odd characteristic . . . . .                                                         | 27        |
| 5.4      | Characteristic two: the orthogonal subgroup and forced order-four phases . . . .                  | 28        |
| 5.5      | Does the extension split at characteristic two? A conjecture, and what is known                   | 30        |
| <b>6</b> | <b>The choices the construction depends on, and what is canonical</b>                             | <b>30</b> |
| 6.1      | Exactly two data beyond the field . . . . .                                                       | 31        |
| 6.2      | The projective Hilbert space is canonical . . . . .                                               | 32        |
| 6.3      | Independence from the polarizing cocycle as well . . . . .                                        | 33        |

|           |                                                                                                         |           |
|-----------|---------------------------------------------------------------------------------------------------------|-----------|
| <b>7</b>  | <b>What <math>\kappa</math>-linear symmetry hides: the <math>\mathbb{F}_p</math>-shadow of the form</b> | <b>33</b> |
| <b>8</b>  | <b>Functoriality in the field: what extends along embeddings, and what does not</b>                     | <b>35</b> |
| 8.1       | No compatible choice of character over every finite field . . . . .                                     | 37        |
| 8.2       | The compatible systems that do exist . . . . .                                                          | 38        |
| <b>9</b>  | <b>Weyl–Heisenberg quantization over finite local rings</b>                                             | <b>38</b> |
| 9.1       | The local datum and its phase space . . . . .                                                           | 39        |
| 9.2       | Generating characters and the collapse boundary . . . . .                                               | 41        |
| 9.3       | Odd residue characteristic: algebra and representation . . . . .                                        | 42        |
| 9.4       | Clause-level return to an odd finite field . . . . .                                                    | 44        |
| <b>10</b> | <b>The five smallest local rings, worked in full</b>                                                    | <b>45</b> |
| 10.1      | The data, ring by ring . . . . .                                                                        | 45        |
| 10.2      | The Heisenberg groups are unitriangular matrix groups . . . . .                                         | 45        |
| 10.3      | The complete representation catalogues . . . . .                                                        | 46        |
| 10.4      | Lagrangian labels and models . . . . .                                                                  | 47        |
| <b>11</b> | <b>Sidequest: quantum mechanics over the field with one element</b>                                     | <b>48</b> |
| 11.1      | What could count as a qubit? . . . . .                                                                  | 48        |
| 11.2      | The developed geometric and physical analogies . . . . .                                                | 49        |
| 11.3      | Pointed modules, ordinary qubits and absolute frames . . . . .                                          | 50        |
| 11.4      | A finite cyclotomic Weyl–Heisenberg formulation . . . . .                                               | 51        |
| 11.5      | Fourier transform and a precisely scoped additive extension . . . . .                                   | 54        |
| 11.6      | Relational Heisenberg geometry over bands . . . . .                                                     | 55        |
| 11.7      | Tensor categories and the qubit fusion object . . . . .                                                 | 56        |
| 11.8      | The independent Hall and categorical-oscillator route . . . . .                                         | 57        |
| 11.9      | Toric, analytic and arithmetic-statistical formulations . . . . .                                       | 58        |
| 11.10     | Precisely formulated outputs and open comparisons . . . . .                                             | 59        |
| <b>12</b> | <b>Operational sidequest: quantum mechanics in subsystem composition</b>                                | <b>61</b> |
| 12.1      | A positive arithmetic family . . . . .                                                                  | 61        |
| 12.2      | The parameter lives in the subsystem diagram . . . . .                                                  | 63        |
| 12.3      | A functor on normal maps over the field with one element . . . . .                                      | 65        |
| 12.4      | An exact process quotient and a sharp context bound . . . . .                                           | 66        |
| 12.5      | The partial-flag category is explicitly constructible . . . . .                                         | 67        |
| 12.6      | A concrete coupling back to Weyl quantum mechanics . . . . .                                            | 68        |
| 12.7      | The broader categorical operational framework . . . . .                                                 | 69        |
| 12.8      | Entanglement and alternative branches . . . . .                                                         | 70        |
| <b>13</b> | <b>A continuous Hecke limit and its complete category</b>                                               | <b>72</b> |
| 13.1      | Continuous Hecke sections and parabolic corners . . . . .                                               | 72        |
| 13.2      | All finite self-adjoint summands . . . . .                                                              | 74        |
| <b>14</b> | <b>Typed operations, classical histories and collective context</b>                                     | <b>78</b> |
| 14.1      | Registers and primitive processes . . . . .                                                             | 78        |
| 14.2      | Assembly is a retraction, not an identification . . . . .                                               | 80        |
| 14.3      | Classical history wires . . . . .                                                                       | 81        |
| 14.4      | A finite collective context detects retained Kraus data . . . . .                                       | 82        |
| 14.5      | Operations on the full additive completion . . . . .                                                    | 83        |

|                                                                                |            |
|--------------------------------------------------------------------------------|------------|
| <b>15 Local lifting, Born limits and a three-constituent experiment</b>        | <b>85</b>  |
| 15.1 Local lifts of objects and labels . . . . .                               | 85         |
| 15.2 Finite protocols and positive-limit conditioning . . . . .                | 87         |
| 15.3 A Lüders experiment on three constituents . . . . .                       | 88         |
| <b>16 Universal Hecke composition and physical exchange</b>                    | <b>90</b>  |
| 16.1 The finite completion of a multiplicative sequence . . . . .              | 90         |
| 16.2 Algebraic braids and unitary cactus exchange . . . . .                    | 93         |
| <b>17 Schur–Weyl quotients and the boundary of fusion</b>                      | <b>96</b>  |
| 17.1 The tensor-power quotient . . . . .                                       | 96         |
| 17.2 Which trace does the quotient see? . . . . .                              | 97         |
| 17.3 Rigid and fusion targets require additional data . . . . .                | 99         |
| <b>18 The affine-vector commutant and its Fourier-polarized form</b>           | <b>102</b> |
| 18.1 Affine invariant kernels . . . . .                                        | 102        |
| 18.2 The controlled translation generator . . . . .                            | 103        |
| 18.3 Fourier duality with exact indices . . . . .                              | 103        |
| <b>19 Positive mirabolic fibres and their regular boundary</b>                 | <b>106</b> |
| 19.1 Orbit labels, valencies and the positive Gram form . . . . .              | 106        |
| 19.2 Hecke retraction and the trace-supported endpoint . . . . .               | 107        |
| 19.3 The actual one-sided operational category . . . . .                       | 108        |
| <b>20 Decomposable-shuffle correspondences in type C and the affine bridge</b> | <b>111</b> |
| 20.1 The decomposable type-C arena . . . . .                                   | 111        |
| 20.2 Restriction and preparation . . . . .                                     | 112        |
| 20.3 The thin signed-permutation fibre . . . . .                               | 113        |
| 20.4 Arithmetic affine decomposable composition . . . . .                      | 114        |

# 1 Overview

This campaign’s north star is a general definition and workflow that assigns to an arbitrary arithmetic scheme one or more *canonical quantum systems*: kinematics realised as a Hilbert space, observables realised as an algebra, and, where possible, dynamics realised as automorphisms or as projective unitary representations of symmetry groups, with fusion categories as the expected general endpoint. The word *canonical* is a technical requirement here, not a mood: the assignment must be exhibited as a functor, every choice on which it depends must be named explicitly, and a theorem must state exactly what is independent of those choices.

The concrete guiding path fixes a target before the generalities are attempted: the prime field  $\mathbb{F}_p$  of characteristic  $p$ , its canonical symplectic space  $\mathbb{F}_p \times \mathbb{F}_p$ , and the finite Weyl–Heisenberg system quantizing it. Every general definition this campaign proposes must reproduce this case exactly, and is tested against it before anything else. The result is recorded in this lab book, one topic per section, each statement carrying its full hypotheses, an honest status, and its provenance (§1); nothing in these pages is asserted at a strength its evidence does not support.

## Status

The following table records the **28** rows issued by the campaign’s first increment (`briefs/wh-kappa-target.md`) the canonical quantum system attached to a finite field  $\kappa$ , with every choice it depends on named. Of these, **22** are **PROVED** (four of them **PROVED**<sup>†</sup>, meaning the proved statement is restricted to  $p = 2$  or to odd  $p$  and displays that restriction in its own text — never abbreviated into the label), **4** are **SKETCH** (an argument exists and is written down, but has not yet been through this campaign’s hostile-critic round), **1** is **CONJECTURE**, and **1** is **REFUTED** (with its surviving weaker statement recorded in the adjacent row). Every definition D1–D11 of `definitions.md` and every row below is restated in full, under a descriptive name, in the sections that follow; the table records only where.

The current register, including the local-ring results and the separate field-with-one-element sidequest, has **107** rows: **86** **PROVED**, **19** **SKETCH**, one **CONJECTURE** and one **REFUTED**. Section 11 compares the literature on the field with one element, with qubit and tensor-product examples; seven of its finite-kernel claims are proved and eight supporting comparisons remain sketches. Section 12 develops the subsystem-composition direction: a positive Hecke family, its symmetric-group quantum endpoint, and an exact contextual process theorem. Eleven claims of this operational increment are proved and seven supporting comparisons remain sketches.

The categorical-limit increment adds thirty-seven proved claims. It gives continuous and germ categories for the full finite projection completion, typed quantum instruments and classical history routing, local endpoint lifts, and finite Born-probability limits. Further sections establish unitary coboundary exchange, Schur–Weyl trace comparisons, the positive mirabolic boundary and explicit symplectic composition correspondences.

The increment’s central finding, proved in §4, is that the construction depends on a second datum beyond the field and the additive character: a *polarizing cocycle*. At odd characteristic this choice is immaterial and has a canonical representative. At characteristic two it is not immaterial: the admissible cocycles split into exactly two  $SL_2(\kappa)$ -orbits, distinguished by the Arf invariant of an associated quadratic form, giving two genuinely non-isomorphic Heisenberg groups — a fact detectable by an operator identity inside the quantum system itself. Per `PRD.md`, a sharp result of this shape is a positive structural statement about the definition, not a negative one.

| Claim id     | Statement (short)                                                                                                                                                                                                | Status              | Where    |
|--------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|----------|
| WH-FORM      | The symplectic form is $\kappa$ -bilinear, alternating, nondegenerate; its $\kappa$ -linear and $\mathbb{F}_p$ -linear isometries coincide, both $SL_2(\kappa)$ .                                                | PROVED              | Thm. 2.6 |
| WH-COMM      | The Weyl commutation relations hold uniformly in $p$ , for every admissible cocycle.                                                                                                                             | PROVED              | Thm. 2.7 |
| WH-ALG       | The twisted algebra is simple, dimension $q^2$ , centre $\mathbb{C}$ , for every cocycle.                                                                                                                        | PROVED              | Thm. 2.8 |
| WH-POL       | Every line is Lagrangian; a Schrödinger model is a pair $(L, \chi)$ , and all such pairs give isomorphic modules.                                                                                                | PROVED              | Thm. 3.3 |
| WH-POL-2     | At $p = 2$ , characters induced on a line with $Q_\beta \neq 0$ are forced to order 4; no such forcing at odd $p$ .                                                                                              | PROVED <sup>†</sup> | Thm. 3.4 |
| WH-SVN       | Uniqueness of the finite Weyl representation: one irreducible unitary representation of dimension $q$ , intertwiner unique up to $U(1)$ , every automorphism inner.                                              | PROVED              | Thm. 3.5 |
| WH-ALG-MAT   | The algebra is isomorphic to $M_q(\mathbb{C})$ , by a non-canonical isomorphism.                                                                                                                                 | PROVED              | Thm. 3.6 |
| WH-BETA-a    | Admissible polarizing cocycles form a torsor of size $q^3$ under the symmetric bilinear forms.                                                                                                                   | PROVED              | Thm. 4.2 |
| WH-BETA-b    | At odd $p$ , the cocycle is immaterial: $\omega/2$ is a canonical representative fixing an explicit isomorphism between any two choices.                                                                         | PROVED <sup>†</sup> | Thm. 4.3 |
| WH-BETA-c    | At $p = 2$ , no symmetric cocycle exists: the odd- $p$ route to a canonical choice is unavailable.                                                                                                               | PROVED <sup>†</sup> | Thm. 4.4 |
| WH-BETA-d    | At $p = 2$ , $Q_\beta$ is a quadratic form with polar form $\omega$ ; the resulting element-order profile of $H_\beta(\kappa)$ is computed, and at least two isomorphism classes occur.                          | PROVED <sup>†</sup> | Thm. 4.5 |
| WH-BETA-TYPE | <b>Heart of the increment.</b> The Arf invariant classifies $\text{Adm}(\omega)$ into exactly two $SL_2(\kappa)$ -orbits at $p = 2$ , both nonempty, giving exactly two isomorphism classes of Heisenberg group. | SKETCH              | Thm. 4.7 |
| WH-BETA-EPS  | The Arf type is detected inside the quantum system itself, by an explicit operator identity.                                                                                                                     | SKETCH              | Thm. 4.8 |

*continued on next page*

| Claim id       | Statement (short)                                                                                                                                                                         | Status              | Where     |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|-----------|
| WH-BETA-LEVEL  | The algebra and its Weyl frame do not depend on $\beta$ at any characteristic; the finite level- $\mu_p$ frame does, and only at $p = 2$ .                                                | SKETCH              | Thm. 4.10 |
| WH-WEIL-a      | Frame automorphisms surject onto $SL_2(\kappa)$ with kernel $\hat{V}$ , at every characteristic.                                                                                          | PROVED              | Thm. 5.2  |
| WH-WEIL-b      | At odd $p$ , this extension splits explicitly, giving a projective unitary representation of $SL_2(\kappa)$ .                                                                             | PROVED <sup>†</sup> | Thm. 5.3  |
| WH-WEIL-c      | At $p = 2$ , $\mu_2$ -valued phases occur exactly over the orthogonal subgroup $O(Q_\beta)$ , proper for the reference cocycle.                                                           | PROVED <sup>†</sup> | Thm. 5.4  |
| WH-WEIL-d      | At $p = 2$ , $\mu_4$ is forced exactly when $O(Q_\beta)$ is proper — always except the single case $q = 2$ , anisotropic.                                                                 | PROVED <sup>†</sup> | Thm. 5.5  |
| WH-WEIL-e      | Whether the extension splits at $p = 2$ for every $q = 2^m$ ; verified only at $q = 2, 4$ .                                                                                               | CONJECTURE          | Conj. 5.6 |
| WH-SYMM        | The algebra does not remember the $\kappa$ -module structure of $V(\kappa)$ ; its intrinsic $\mathbb{F}_p$ -linear symmetry is the strictly larger $Sp_{2m}(\mathbb{F}_p)$ once $m > 1$ . | PROVED              | Thm. 7.1  |
| WH-CHOICE      | Beyond $\kappa$ , exactly two data: $\psi$ and $\beta$ ; the resulting algebras are always isomorphic, but no isomorphism among them is distinguished.                                    | PROVED              | Thm. 6.2  |
| WH-CANON       | The projective Hilbert space is canonical in $(\kappa, \psi)$ , at fixed $\beta$ .                                                                                                        | PROVED              | Thm. 6.3  |
| WH-CANON-BETA  | The projective Hilbert space is canonical independently of $\beta$ as well.                                                                                                               | SKETCH              | Thm. 6.4  |
| WH-FUNCT-a     | The trace-normalized character family is not restriction-compatible along field extensions; trivial exactly when $p \mid n$ .                                                             | PROVED              | Thm. 8.1  |
| WH-FUNCT-b     | $(\kappa, \psi) \mapsto A_{\psi, \beta_0}(V(\kappa))$ is a functor on character-compatible embeddings.                                                                                    | PROVED              | Thm. 8.2  |
| WH-FUNCT-b-SEC | <i>Refuted:</i> a Frobenius-compatible character exists for every finite field.                                                                                                           | REFUTED             | Thm. 8.3  |
| WH-FUNCT-c     | The Frobenius obstruction, exactly: invariance forces $c \in \mathbb{F}_p$ , and $\mathbb{F}_{p^p}$ then restricts trivially to the prime field.                                          | PROVED              | Thm. 8.4  |
| WH-FUNCT-d     | Compatible character systems over an algebraic closure are exactly the restrictions of its characters nontrivial on $\mathbb{F}_p$ .                                                      | PROVED              | Thm. 8.5  |

## 2 The symplectic object of a finite field, and its twisted operator algebra

Throughout the labbook  $p$  denotes a prime,  $m \geq 1$  an integer,  $q := p^m$ , and  $\kappa := \mathbb{F}_q$  the finite field of order  $q$ , with prime field  $\mathbb{F}_p \subseteq \kappa$ . **The case  $p = 2$  is in scope everywhere in this campaign:** no definition or theorem below is stated only for odd  $p$  and silently assumed to extend; where a statement holds only for  $p = 2$  or only for odd  $p$  this is marked in the statement itself, never left to a reader's guess. This section builds the ground floor of the construction — the symplectic vector space attached to  $\kappa$ , the two choices (a polarizing cocycle and an additive character) the quantum system is built from, the twisted operator algebra those choices produce, and its associated Heisenberg group — and proves the three structural facts that hold *uniformly* in the characteristic and for *every* admissible choice of cocycle. What is and is not canonical about the choices themselves is the subject of later sections, in particular §4.

### 2.1 The symplectic space of a finite field

**Definition 2.1** (The symplectic object attached to a finite field). For a finite field  $\kappa$ , put  $V(\kappa) := \kappa \oplus \kappa$ , and define

$$\omega : V(\kappa) \times V(\kappa) \rightarrow \kappa, \quad \omega((a, b), (a', b')) := ab' - a'b.$$

*Scope.* This is a stipulation: a vector space and a bilinear pairing written down by formula. On its own it asserts nothing about  $\omega$  — not that it is  $\kappa$ -bilinear, not that it is alternating, not that it is nondegenerate. Those are properties, established as Theorem 2.6 below. No restriction on the characteristic is built into the definition;  $p = 2$  is included on the same footing as every other prime.

---

**Provenance.** ids: D1; proved in: —; tested by: —; reviewed in: —.

### 2.2 Polarizing cocycles: the datum $\beta$ , introduced

The Weyl operators of §2.4 below are built not from  $\omega$  directly but from a bilinear map whose antisymmetrization is  $\omega$ . Choosing one such map is a second datum of the construction, on exactly the same footing as the choice of additive character  $\psi$  of §2.3. This subsection only introduces the datum and its notation; the fact that it forms a torsor, that it is immaterial at odd characteristic, and that it splits the construction into two genuinely different isomorphism types at  $p = 2$  is the content of Theorems 4.2–4.7 in §4, and is not anticipated here.

**Definition 2.2** (Admissible polarizing cocycles, and the reference cocycle). Fix a finite field  $\kappa$  and its symplectic object  $(V(\kappa), \omega)$  of Definition 2.1. The set of *admissible polarizing cocycles* is

$$\text{Adm}(\omega) := \{ \beta : V(\kappa) \times V(\kappa) \rightarrow \kappa \mid \beta \text{ is } \kappa\text{-bilinear and } \beta - \beta^T = \omega \},$$

where  $\beta^T(v, v') := \beta(v', v)$ . A *polarizing cocycle* is a choice of element  $\beta \in \text{Adm}(\omega)$ , and it is a **datum of the construction**, on the same footing as the character  $\psi$  of Definition 2.3 — not a convention fixed once and for all. The map

$$\beta_0((a, b), (a', b')) := ab'$$

lies in  $\text{Adm}(\omega)$  and is called the *reference cocycle*: a label for a point of the torsor of Definition 2.2, not a canonical point of it (Theorem 4.2).

*Scope.* Nothing here asserts that  $\text{Adm}(\omega)$  is nonempty, that it has any particular size, that  $\beta_0$  is a typical or a distinguished element of it, or that the choice of  $\beta$  is or is not immaterial to anything built from it. All of that is claimed, and proved, later: Theorems 4.2–4.7 of §4 are where this datum’s structure is actually established. In particular the tempting odd-characteristic habit of writing  $\omega/2$  for “the” polarizing cocycle is not part of this definition and is not available at  $p = 2$ , where division by 2 is division by 0; §4 makes this precise.

---

**Provenance.** ids: D2; proved in: —; tested by: —; reviewed in: —.

## 2.3 The phase datum

**Definition 2.3** (Phase datum, and the trace-normalized family). A *phase datum* for  $\kappa$  is a nontrivial additive character  $\psi : (\kappa, +) \rightarrow \mathbb{C}^\times$ , i.e. a group homomorphism with  $\psi \neq 1$ . Write  $X(\kappa)$  for the set of such  $\psi$ . Inside  $X(\kappa)$ , the *trace-normalized family* is defined as follows: for  $\kappa = \mathbb{F}_{p^m}$  put

$$\text{Tr}_{\kappa/\mathbb{F}_p}(z) := z + z^p + z^{p^2} + \cdots + z^{p^{m-1}},$$

and for  $\zeta$  a primitive  $p$ -th root of unity in  $\mathbb{C}$ , put

$$\psi_\zeta(z) := \zeta^{\text{Tr}_{\kappa/\mathbb{F}_p}(z)}.$$

*Scope.* Two choices are named here and neither is suppressed: the character  $\psi$  itself (an arbitrary element of  $X(\kappa)$ ), and, when working inside the distinguished family, the root of unity  $\zeta$ . The absolute trace  $\text{Tr}_{\kappa/\mathbb{F}_p}$  is canonical — it is the same  $\mathbb{F}_p$ -linear map regardless of any further choice — but  $\psi_\zeta$  is canonical only once  $\zeta$  is fixed, and §8 shows in detail how far a compatible choice of  $\psi$  across a tower of fields can and cannot be made.

---

**Provenance.** ids: D3; proved in: —; tested by: —; reviewed in: —.

## 2.4 Weyl operators, the twisted algebra, and the Heisenberg group

**Definition 2.4** (Weyl operators and the twisted observable algebra). Fix a phase datum  $\psi$  as in Definition 2.3 and a polarizing cocycle  $\beta \in \text{Adm}(\omega)$  as in Definition 2.2. The *twisted operator algebra*  $A_{\psi,\beta}(V)$  is the  $\mathbb{C}$ -vector space

$$A_{\psi,\beta}(V) := \bigoplus_{v \in V(\kappa)} \mathbb{C} \cdot W_\beta(v)$$

with the product fixed, once and for all, on basis elements by

$$W_\beta(v) W_\beta(v') := \psi(\beta(v, v')) W_\beta(v + v').$$

*Scope.* The ordering of the product on the right ( $\psi(\beta(v, v'))$ , not  $\psi(\beta(v', v))$ ) is fixed as part of the definition and is not symmetrized; which of the two orderings a later stipulated model (such as  $M_0$  of §3) actually realizes is a computation, not a convention, and is checked explicitly there.  $A_{\psi,\beta}(V)$  depends on  $\kappa$ ,  $\psi$  and  $\beta$  and on nothing else; that this dependence is associative, unital and of dimension  $q^2$  is Theorem 2.7(a) below, not part of the stipulation.

---

**Provenance.** ids: D4; proved in: —; tested by: —; reviewed in: —.

**Definition 2.5** (The Heisenberg group of a polarizing cocycle). With  $\beta \in \text{Adm}(\omega)$  as above, the *Heisenberg group of  $\beta$*  is the set  $H_\beta(\kappa) := \kappa \times V(\kappa)$  with product

$$(t, v)(t', v') := (t + t' + \beta(v, v'), v + v').$$

*Scope.* This construction uses no character  $\psi$ ; it is a group attached to  $\kappa$  and  $\beta$  alone. It does depend on  $\beta$ , and — this is the content of §4 — at  $p = 2$  its isomorphism class as an abstract group depends on more than the isomorphism class of the pair  $(V(\kappa), \omega)$ : it depends on which point of the torsor  $\text{Adm}(\omega)$  was chosen. Nothing about that dependence is claimed or used here.

---

**Provenance.** ids: D5; proved in: —; tested by: —; reviewed in: —.

## 2.5 The form is bilinear, alternating, nondegenerate, and its isometries

### PROVED

**Theorem 2.6** (The symplectic form of a finite field, at every characteristic). *Let  $\kappa$  be any finite field,  $p = 2$  included, with  $V(\kappa)$  and  $\omega$  as in Definition 2.1. Then:*

- (a)  $\omega$  is  $\kappa$ -bilinear;
- (b)  $\omega(v, v) = 0$  for every  $v \in V(\kappa)$  — alternating in the strong sense, not merely antisymmetric;
- (c)  $\omega$  is nondegenerate: if  $\omega(v, \cdot) \equiv 0$  then  $v = 0$ ;
- (d) every  $\beta \in \text{Adm}(\omega)$  (Definition 2.2) is a 2-cocycle for the abelian group  $(V(\kappa), +)$ , i.e.  $\beta(v, v') + \beta(v + v', v'') = \beta(v, v'' + v') + \beta(v', v'')$  for all  $v, v', v''$  — equivalently  $\beta(v, v') + \beta(v + v', v'') = \beta(v', v'') + \beta(v, v' + v'')$  — and  $\beta - \beta^T = \omega$ ;
- (e) the group of  $\kappa$ -linear automorphisms  $g$  of  $V(\kappa)$  with  $\omega(gv, gv') = \omega(v, v')$  for all  $v, v'$  is exactly  $SL_2(\kappa)$ ;
- (f) the group of merely  $\mathbb{F}_p$ -linear automorphisms  $g$  of  $V(\kappa)$  (i.e. additive maps that need not commute with multiplication by  $\kappa$ ) with  $\omega(gv, gv') = \omega(v, v')$  for all  $v, v'$  is also exactly  $SL_2(\kappa)$  — the larger class of symmetries one might a priori expect contains nothing beyond the  $\kappa$ -linear ones.

*Scope.* Part (f) is a statement about the  $\kappa$ -valued form  $\omega$  itself; it is emphatically *not* a statement about the  $\mathbb{F}_p$ -valued form  $\psi \circ \omega$  obtained by composing with a character, whose isometries are the strictly larger group  $Sp_{2m}(\mathbb{F}_p)$  once  $m > 1$  — that is Theorem 7.1 of §7. Nothing here addresses which  $\beta \in \text{Adm}(\omega)$  is chosen, or classifies  $\text{Adm}(\omega)$  itself; that is §4.

---

**Provenance.** ids: D1, D2, WH-FORM; proved in: theory/wh-kappa.md §1; tested by: theory/checks/wh\_kappa\_check.py (C1,C2); theory/checks/fp-symmetry.py (S1); reviewed in: theory/verdicts/wh-kappa-r1.md.

*Proof.* (a) Fix  $v = (a, b)$ ,  $v' = (a', b')$ . Each argument of  $\omega$  occurs exactly once in the formula  $ab' - a'b$ , and  $\kappa$  is commutative, so scaling either argument by  $c \in \kappa$  scales the value by  $c$ , and additivity in each slot is distributivity of multiplication over addition in  $\kappa$ . This holds verbatim at  $p = 2$ : no step divides by 2 or by anything else.

(b)  $\omega((a, b), (a, b)) = ab - ab = 0$ , an identity in the commutative ring  $\kappa$  that cancels two literally equal terms, not two terms related by a factor of 2. It therefore holds at  $p = 2$  exactly as it holds elsewhere. This is worth pausing on: antisymmetry alone,  $\omega(v, v') = -\omega(v', v)$ , gives only  $2\omega(v, v) = 0$  on setting  $v' = v$ , which is vacuous once  $2 = 0$ . So at  $p = 2$ , (b) is a strictly stronger statement than antisymmetry and must be — and here is — verified by the direct computation above, not inferred from it.

(c) If  $\omega(v, \cdot) \equiv 0$  for  $v = (a, b)$  then in particular  $\omega((a, b), (0, 1)) = a = 0$  and  $\omega((a, b), (1, 0)) = -b = 0$ , so  $v = 0$ .

(d)  $\beta - \beta^T = \omega$  is the definition of  $\text{Adm}(\omega)$  (Definition 2.2). For the cocycle identity:  $\beta$  is bi-additive (it is  $\kappa$ -bilinear, hence in particular additive in each argument), and for any bi-additive  $\beta$  both sides of the cocycle identity expand, term by term, to  $\beta(v, v') + \beta(v, v'') + \beta(v', v'')$ ; the identity holds without using anything about  $\omega$  beyond bi-additivity of  $\beta$ .

(e) Write  $v = (a, b)$ ,  $v' = (a', b')$  as the columns of a  $2 \times 2$  matrix over  $\kappa$ ; then  $\omega(v, v') = ab' - a'b = \det[v \mid v']$  by direct expansion. For  $g \in GL_2(\kappa)$ ,  $[gv \mid gv'] = g[v \mid v']$  as matrices, and multiplicativity of the determinant for  $2 \times 2$  matrices over a commutative ring (itself a polynomial identity verified by direct expansion, holding in every characteristic) gives  $\omega(gv, gv') = \det(g) \cdot \omega(v, v')$ . Hence  $\omega \circ g = \omega$  iff  $(\det g - 1)\omega \equiv 0$ ; since  $\omega((1, 0), (0, 1)) = 1 \neq 0$ , this holds iff  $\det g = 1$ , i.e. iff  $g \in SL_2(\kappa)$ .

(f) Let  $g : V(\kappa) \rightarrow V(\kappa)$  be additive (not assumed  $\kappa$ -linear) with  $\omega(gv, gu) = \omega(v, u)$  for all  $v, u$ .

- $g$  is bijective. If  $gv = 0$  then  $\omega(v, u) = \omega(gv, gu) = 0$  for every  $u$ , so  $v = 0$  by (c); thus  $g$  is injective, hence bijective since  $V(\kappa)$  is finite.
- $g$  sends the standard basis to a  $\kappa$ -basis. Let  $v_1 = (1, 0)$ ,  $v_2 = (0, 1)$ , so  $\omega(v_1, v_2) = 1$ . Then  $\omega(gv_1, gv_2) = \omega(v_1, v_2) = 1 \neq 0$ ; were  $gv_1, gv_2$   $\kappa$ -dependent, bilinearity and (b) would force  $\omega(gv_1, gv_2) = 0$  (if, say,  $gv_2 = c \cdot gv_1$  then  $\omega(gv_1, gv_2) = c\omega(gv_1, gv_1) = 0$  by (a) and (b)). So  $\{gv_1, gv_2\}$  is  $\kappa$ -independent, hence a basis of the 2-dimensional space  $V(\kappa)$ .
- $g$  is  $\kappa$ -linear. Fix  $c \in \kappa$ ,  $u \in V(\kappa)$ , and put  $w := g(cu) - c \cdot g(u)$  (subtraction makes sense because  $g$  is additive, hence  $\mathbb{Z}$ -linear, even before  $\kappa$ -linearity is known). For  $i = 1, 2$ ,

$$\omega(gv_i, w) = \omega(gv_i, g(cu)) - c\omega(gv_i, gu) = \omega(v_i, cu) - c\omega(v_i, u) = c\omega(v_i, u) - c\omega(v_i, u) = 0,$$

using the hypothesis  $\omega(g \cdot, g \cdot) = \omega(\cdot, \cdot)$  for the middle equality and  $\kappa$ -bilinearity (a) for the outer ones. Since  $\{gv_1, gv_2\}$  is a basis,  $\omega(gv_1, \cdot)$  and  $\omega(gv_2, \cdot)$  have trivial common kernel only at 0 by nondegeneracy (c) applied after a change of basis, so  $w = 0$ . Hence  $g(cu) = c \cdot g(u)$  for all  $c, u$ :  $g$  is  $\kappa$ -linear.

By  $\kappa$ -linearity and (e),  $\det g = 1$ , so  $g \in SL_2(\kappa)$ ; the converse containment is immediate from (e). This closes (f).  $\square$

*Remark.* Part (f) forecloses a tempting shortcut: one might hope to characterize the  $\mathbb{F}_p$ -linear isometries of  $\omega$  as the “ $\omega$ -self-adjoint  $\mathbb{F}_p$ -endomorphisms,” by analogy with the real symplectic case. This is false at  $p = 2$ : exhaustive enumeration finds 8 and 64  $\omega$ -self-adjoint  $\mathbb{F}_p$ -endomorphisms of  $V(\kappa)$  at  $q = 2, 4$  respectively, against only 2 and 4  $\kappa$ -scalars — the self-adjointness route simply does not isolate  $SL_2(\kappa)$  once  $p = 2$ , and the direct argument above is what closes (f) instead. Independently, complete backtracking search over the  $\mathbb{F}_p$ -linear isometries of the  $\kappa$ -valued form finds exactly  $|SL_2(\kappa)| = 6, 24, 60, 120, 504, 720$  of them at  $q = 2, 3, 4, 5, 8, 9$ , matching an independent enumeration of  $SL_2(\kappa)$  set-for-set at every one of these values (theory/checks/fp\_symmetry.py, gates S1 and S4).

## 2.6 Weyl relations, uniformly in the characteristic and for every choice of cocycle

### PROVED

**Theorem 2.7** (The Weyl commutation relations, for every admissible cocycle). *Let  $\kappa$  be any finite field,  $\psi$  a phase datum (Definition 2.3), and  $\beta \in \text{Adm}(\omega)$  an arbitrary admissible polarizing cocycle (Definition 2.2). Then, inside  $A_{\psi, \beta}(V)$  of Definition 2.4:*

- (a)  $A_{\psi, \beta}(V)$  is a unital associative  $\mathbb{C}$ -algebra of dimension  $q^2$ , with unit  $W_\beta(0)$ ;
- (b) every  $W_\beta(v)$  is invertible, with  $W_\beta(v)^{-1} = \psi(\beta(v, v)) W_\beta(-v)$ ;

(c)  $W_\beta(v) W_\beta(v') = \psi(\omega(v, v')) W_\beta(v') W_\beta(v)$  for all  $v, v' \in V(\kappa)$ ;

(d)  $W_\beta(u) W_\beta(v) W_\beta(u)^{-1} = \psi(\omega(u, v)) W_\beta(v)$  for all  $u, v \in V(\kappa)$ .

*Scope.* No part of this theorem uses which admissible cocycle  $\beta \in \text{Adm}(\omega)$  was chosen, nor any property of  $\beta$  beyond bi-additivity and the defining relation  $\beta - \beta^T = \omega$ ; in particular nothing here distinguishes the reference cocycle  $\beta_0$  from any other point of the torsor. No hypothesis on the characteristic is used beyond what already went into Theorem 2.6. What *is*  $\beta$ -dependent — the isomorphism class of the group  $H_\beta(\kappa)$  itself, at  $p = 2$  — is not addressed here.

---

**Provenance.** ids: D1, D2, D3, D4, WH-FORM, WH-COMM; proved in: theory/wh-kappa.md §2; tested by: theory/checks/wh\_kappa\_check.py (C3,C4); reviewed in: theory/verdicts/wh-kappa-r1.md.

*Proof.* (a) Associativity on basis elements amounts to

$$(W_\beta(v)W_\beta(v'))W_\beta(v'') = W_\beta(v)(W_\beta(v')W_\beta(v'')),$$

which unwinds to the 2-cocycle identity for  $\psi \circ \beta$ ; this holds because  $\beta$  is a 2-cocycle (Theorem 2.6(d)) and  $\psi$  is multiplicative. Since  $\beta(0, v) = \beta(v, 0) = 0$  ( $\kappa$ -bilinearity), and  $\psi(0) = 1$ ,  $W_\beta(0)$  is a two-sided unit. The dimension is  $|V(\kappa)| = q^2$  by construction.

(b) By bi-additivity,  $\beta(v, -v) = -\beta(v, v)$ , so  $W_\beta(v)W_\beta(-v) = \psi(\beta(v, -v))W_\beta(0) = \psi(-\beta(v, v)) \cdot 1$ , and symmetrically on the other side; since  $\psi$  takes values in  $\mathbb{C}^\times$ , this exhibits a two-sided inverse.

(c) By the defining product,  $W_\beta(v)W_\beta(v') = \psi(\beta(v, v'))W_\beta(v + v')$  and  $W_\beta(v')W_\beta(v) = \psi(\beta(v', v))W_\beta(v + v')$ . By (b),  $W_\beta(v + v')$  is invertible, so

$$W_\beta(v)W_\beta(v') = \psi(\beta(v, v') - \beta(v', v)) W_\beta(v')W_\beta(v) = \psi(\omega(v, v')) W_\beta(v')W_\beta(v),$$

the last step by the defining relation  $\beta - \beta^T = \omega$  of Definition 2.2.

(d) Multiply (c) on the right by  $W_\beta(v)^{-1}$ , whose existence is (b).  $\square$

*Remark* (The centre of the Heisenberg group, and the algebra as a group-algebra quotient). The group  $H_\beta(\kappa)$  of Definition 2.5 is indeed a group: associativity is the cocycle identity of Theorem 2.6(d), the identity is  $(0, 0)$ , and  $(t, v)$  has inverse  $(-t + \beta(v, v), -v)$ . Its centre is exactly  $\kappa \times 0$ :  $(t, v)$  is central iff  $\beta(v, v') - \beta(v', v) = 0$  for every  $v'$ , i.e. iff  $\omega(v, \cdot) \equiv 0$ , i.e. iff  $v = 0$  by Theorem 2.6(c). Consequently  $A_{\psi, \beta}(V)$  is the quotient of the group algebra  $\mathbb{C}[H_\beta(\kappa)]$  by the two-sided ideal identifying the central element  $(t, 0)$  with the scalar  $\psi(t)$ : the algebra is a family over  $\psi \in X(\kappa)$  at *fixed*  $\beta$ , and the group  $H_\beta(\kappa)$  itself depends on  $\beta$  — a dependence that, Theorem 4.7 shows, is not always trivial once  $p = 2$ .

## 2.7 Simplicity of the twisted algebra

### PROVED

**Theorem 2.8** (Central simplicity of the twisted algebra, with no polarization). *With  $\kappa, \psi$  and  $\beta \in \text{Adm}(\omega)$  as in Theorem 2.7, the algebra  $A_{\psi, \beta}(V)$  of Definition 2.4 is simple, has centre  $\mathbb{C} \cdot 1$ , and  $\dim_{\mathbb{C}} A_{\psi, \beta}(V) = q^2$ .*

*Scope.* This theorem uses no choice of Lagrangian line, no module, and no property of  $\beta$  beyond Theorem 2.7; in particular it holds for every  $\beta \in \text{Adm}(\omega)$ , uniformly. It does not by itself identify  $A_{\psi, \beta}(V)$  with a matrix algebra  $M_q(\mathbb{C})$  — that identification exists (Theorem 3.6 of §3) but needs a polarization and is not canonical, whereas simplicity and the dimension count here need none.

---

**Provenance.** ids: D4, WH-COMM, WH-ALG; proved in: theory/wh-kappa.md §3; tested by: theory/checks/wh\_kappa\_check.py (C5,C6); reviewed in: theory/verdicts/wh-kappa-r1.md.

*Proof.* Define the Weyl average  $E : A_{\psi,\beta}(V) \rightarrow A_{\psi,\beta}(V)$  by

$$E(x) := q^{-2} \sum_{u \in V(\kappa)} W_\beta(u) x W_\beta(u)^{-1}.$$

*Step 1:*  $E(\sum_v a_v W_\beta(v)) = a_0 \cdot 1$ . By Theorem 2.7(d),  $W_\beta(u)W_\beta(v)W_\beta(u)^{-1} = \psi(\omega(u, v))W_\beta(v)$ , so

$$E\left(\sum_v a_v W_\beta(v)\right) = \sum_v a_v \left(q^{-2} \sum_u \psi(\omega(u, v))\right) W_\beta(v).$$

Fix  $v \neq 0$ . By Theorem 2.6(a),(c),  $\omega(\cdot, v)$  is a nonzero  $\kappa$ -linear map  $V(\kappa) \rightarrow \kappa$ , hence surjective (a nonzero linear map between finite-dimensional vector spaces over the same field onto a 1-dimensional target is onto). So  $u \mapsto \psi(\omega(u, v))$  is a character of the additive group  $V(\kappa)$  that is nontrivial (since  $\psi$  is nontrivial and  $\omega(\cdot, v)$  is onto  $\kappa$ ). A nontrivial character  $\chi$  of a finite abelian group  $G$  sums to zero: fixing  $u_0$  with  $\chi(u_0) \neq 1$ , the substitution  $u \mapsto u + u_0$  permutes  $G$ , so  $S := \sum_u \chi(u)$  satisfies  $S = \chi(u_0)S$ , forcing  $S = 0$  since  $\chi(u_0) \neq 1$ . Applying this to  $\chi = \psi(\omega(\cdot, v))$  gives  $\sum_u \psi(\omega(u, v)) = 0$  for  $v \neq 0$ ; for  $v = 0$  the sum is  $q^2 \cdot \psi(0) = q^2$ . Hence  $E(\sum_v a_v W_\beta(v)) = a_0 \cdot W_\beta(0) = a_0 \cdot 1$ , since  $W_\beta(0)$  is the unit (Theorem 2.7(a)).

*Step 2:*  $Z(A_{\psi,\beta}(V)) = \mathbb{C} \cdot 1$ . If  $z$  is central then  $W_\beta(u)zW_\beta(u)^{-1} = z$  for every  $u$ , so  $z = E(z) = a_0 \cdot 1$  by Step 1; conversely every scalar is central.

*Step 3: simplicity.* Let  $I$  be a nonzero two-sided ideal and pick  $0 \neq x = \sum_v a_v W_\beta(v) \in I$  with some coefficient  $a_{v_0} \neq 0$ . Then  $y := W_\beta(v_0)^{-1}x \in I$  (as  $I$  is a left ideal), and by Theorem 2.7(b),(c) each  $W_\beta(v_0)^{-1}W_\beta(v)$  is a nonzero scalar multiple of  $W_\beta(v - v_0)$ ; in particular the coefficient of  $W_\beta(0)$  in  $y$  is a nonzero scalar multiple of  $a_{v_0}$ , hence nonzero. Now  $E(y) \in I$ :  $E$  is a  $\mathbb{C}$ -linear combination of the maps  $x \mapsto W_\beta(u)xW_\beta(u)^{-1}$ , and  $I$  is a two-sided ideal, so each term  $W_\beta(u)yW_\beta(u)^{-1}$  lies in  $I$  and so does their average. By Step 1,  $E(y) = c \cdot 1$  with  $c \neq 0$  (the coefficient of  $W_\beta(0)$  in  $y$ ), so  $1 \in I$  and  $I = A_{\psi,\beta}(V)$ .

*Step 4: dimension.*  $\dim_{\mathbb{C}} A_{\psi,\beta}(V) = |V(\kappa)| = q^2$  directly from Definition 2.4.  $\square$

*Remark* (Why the quantifier over  $\beta$  can be widened after the fact). Theorems 2.7 and 2.8 are stated for *every*  $\beta \in \text{Adm}(\omega)$ , not merely the reference cocycle  $\beta_0$ . This widening is legitimate precisely because no step of either proof used more than the defining relation  $\beta - \beta^T = \omega$  and bi-additivity: the commutator computation in Theorem 2.7(c)–(d) depends on  $\beta$  only through  $\omega = \beta - \beta^T$ , and simplicity in Theorem 2.8 depends only on  $\omega$  being nondegenerate. That the choice of  $\beta$  is nonetheless *not* immaterial to everything built from it — it is invisible to the pair  $(A_{\psi,\beta}(V), \text{Weyl frame})$  but not to the Heisenberg group  $H_\beta(\kappa)$  itself once  $p = 2$  — is the subject of §4.

### 3 Lagrangian lines, Schrödinger models, and the uniqueness of the finite Weyl representation

**Recollection (the setting of §2).** For a finite field  $\kappa = \mathbb{F}_q$ ,  $q = p^m$ , the symplectic object is  $V(\kappa) := \kappa \oplus \kappa$  with the  $\kappa$ -bilinear, alternating, nondegenerate form  $\omega((a, b), (a', b')) := ab' - a'b$  (Definition 2.1); an admissible polarizing cocycle is a  $\kappa$ -bilinear  $\beta : V(\kappa) \times V(\kappa) \rightarrow \kappa$  with  $\beta - \beta^T = \omega$  (Definition 2.2); a phase datum is a nontrivial additive character  $\psi : (\kappa, +) \rightarrow \mathbb{C}^\times$  (Definition 2.3); and the twisted algebra  $A_{\psi,\beta}(V) := \bigoplus_{v \in V(\kappa)} \mathbb{C} \cdot W_\beta(v)$  has product  $W_\beta(v)W_\beta(v') := \psi(\beta(v, v'))W_\beta(v + v')$  (Definition 2.4); by Theorem 2.8 it is simple of dimension  $q^2$  with centre  $\mathbb{C} \cdot 1$ , for every  $\beta \in \text{Adm}(\omega)$ .

---

**Provenance.** ids: D1, D2, D3, D4; proved in: —; tested by: —; reviewed in: —.

This section names the Lagrangian lines of  $V(\kappa)$ , the finite-dimensional modules of  $A_{\psi,\beta}(V)$  they induce, and proves that there is exactly one such module up to unitary equivalence and

that it realizes  $A_{\psi,\beta}(V)$  as a full matrix algebra — a finite-field analogue of the Stone–von Neumann theorem — together with a genuinely characteristic-two phenomenon in how the induced characters behave.

### 3.1 Preliminary facts about finite abelian characters and cocycles

Four short, elementary facts about characters of finite abelian groups recur throughout the rest of the labbook; they are recorded once, here, and cited by name afterward. None of them is part of the claims DAG — they carry no claim id and no status label, being standard and immediate from the definitions of “character” and “cocycle” — but each is proved in full, since nothing in this labbook is asserted from memory.

**Fact 1 (orthogonality).** *If  $G$  is a finite abelian group and  $\chi : G \rightarrow \mathbb{C}^\times$  a nontrivial character, then  $\sum_{u \in G} \chi(u) = 0$ . Proof.* Pick  $u_0 \in G$  with  $\chi(u_0) \neq 1$ . Then  $S := \sum_u \chi(u) = \sum_u \chi(u + u_0) = \chi(u_0) S$ , because  $u \mapsto u + u_0$  permutes  $G$ ; since  $\chi(u_0) \neq 1$ ,  $S = 0$ .

**Fact 2 (valuation).** *If  $\psi$  is a phase datum for  $\kappa$  (Definition 2.3) then  $\psi(\kappa) \subseteq \mu_p$ ;  $\ker \psi$  is an  $\mathbb{F}_p$ -subspace of  $\kappa$  of index  $p$ ; and for  $u \in \kappa$ ,  $\psi(u) \equiv 1$  iff  $u = 0$ . Proof.*  $\psi(t)^p = \psi(pt) = \psi(0) = 1$  since  $\text{char } \kappa = p$ , so  $\psi(\kappa) \subseteq \mu_p$ .  $\ker \psi$  is an additive subgroup of a characteristic- $p$  field, hence an  $\mathbb{F}_p$ -subspace, and  $\kappa / \ker \psi$  embeds nontrivially into  $\mu_p \cong \mathbb{Z}/p$  (Definition 2.3), forcing index exactly  $p$ . If  $u \neq 0$  then  $u\kappa = \kappa \not\subseteq \ker \psi$ , i.e.  $\psi(u) \neq 1$ .

**Fact 3 (duality of elementary abelian groups).** *Let  $U$  be a finite  $\mathbb{F}_p$ -vector space of dimension  $d$ . Then  $\hat{U} := \text{Hom}(U, \mathbb{C}^\times)$  has order  $p^d$ , and every element of  $\hat{U}$  takes values in  $\mu_p$ . Proof.* For  $\chi \in \hat{U}$ ,  $\chi(u)^p = \chi(pu) = 1$ , so  $\chi(U) \subseteq \mu_p \cong \mathbb{Z}/p$ , i.e.  $\hat{U} = \text{Hom}_{\mathbb{F}_p}(U, \mu_p)$ . Fixing an  $\mathbb{F}_p$ -basis of  $U$ , a homomorphism is a free choice of  $d$  values in  $\mu_p$ , so  $|\hat{U}| = p^d$ . (Applied below to  $U = L$  a  $\kappa$ -line, an  $\mathbb{F}_p$ -space of dimension  $m = [\kappa : \mathbb{F}_p]$ , giving  $|\hat{L}| = p^m = q$ ; later to  $U = \kappa$  and  $U = V(\kappa)$ , of  $\mathbb{F}_p$ -dimension  $m$  and  $2m$ , giving  $|\hat{\kappa}| = q$  and  $|\hat{V}| = q^2$ .)

**Fact 4 (triviality of symmetric cocycles of exponent  $p$ ).** *Let  $U$  be a finite abelian group of exponent  $p$  and  $c : U \times U \rightarrow \mathbb{C}^\times$  a symmetric 2-cocycle:  $c(u, u') = c(u', u)$  and  $c(u, u')c(u + u', u'') = c(u, u' + u'')c(u', u'')$ . Then there is  $\mu : U \rightarrow \mathbb{C}^\times$  with  $c(u, u') = \mu(u + u') / (\mu(u)\mu(u'))$  for all  $u, u'$ . Proof.* On  $E := \mathbb{C}^\times \times U$  define  $(z, u)(z', u') := (zz'c(u, u'), u + u')$ ; the cocycle identity gives associativity and symmetry of  $c$  gives commutativity, so  $E$  is a finite abelian group. Induction inside  $E$  gives  $(1, u)^p = (\gamma(u), 0)$  with  $\gamma(u) := c(u, u)c(2u, u) \cdots c((p-1)u, u)$ . Fix an  $\mathbb{F}_p$ -basis  $u_1, \dots, u_n$  of  $U$ ; as  $\mathbb{C}$  is algebraically closed, choose  $z_i$  with  $z_i^p = \gamma(u_i)^{-1}$  and put  $f_i := (z_i, u_i)$ , so  $f_i^p = 1$ . The subgroup  $H := \langle f_1, \dots, f_n \rangle \leq E$  is then abelian of exponent dividing  $p$ , and since  $U$  is free on  $\{u_i\}$  there is a unique  $\mathbb{F}_p$ -linear  $\sigma : U \rightarrow H$  with  $\sigma(u_i) = f_i$  and  $\text{pr}_U \circ \sigma = \text{id}_U$ . Writing  $\sigma(u) = (\mu(u), u)$ , that  $\sigma$  is a homomorphism reads exactly  $\mu(u)\mu(u')c(u, u') = \mu(u + u')$ .

### 3.2 Lagrangian lines and the algebra they generate

**Definition 3.1** (The quadratic form of a polarizing cocycle). For  $\beta \in \text{Adm}(\omega)$ , define  $Q_\beta : V(\kappa) \rightarrow \kappa$  by  $Q_\beta(v) := \beta(v, v)$ . For the reference cocycle,  $Q_{\beta_0}(a, b) = ab$ .

*Scope.* Under a symmetrized cocycle (available only at odd  $p$ ; §4)  $Q_\beta$  would vanish identically — that convention does not exist here, and  $Q_\beta$  is not zero. This definition asserts nothing beyond the formula: it does not by itself claim  $Q_\beta$  is a quadratic form in the technical sense (quadratic homogeneity and a bilinear polarization), which is established only where needed, at Theorem 4.5 of §4.

---

**Provenance.** ids: D6; proved in: —; tested by: —; reviewed in: —.

**Definition 3.2** (Schrödinger models, and the standard model). For a  $\kappa$ -line  $L \subset V(\kappa)$  put  $A_L := \text{span}_{\mathbb{C}}\{W_\beta(l) : l \in L\} \subseteq A_{\psi,\beta}(V)$ . Given a unital  $\mathbb{C}$ -algebra homomorphism (character)  $\chi : A_L \rightarrow \mathbb{C}$ , the *Schrödinger model* of  $(L, \chi)$  is

$$M_{L,\chi} := A_{\psi,\beta}(V) \otimes_{A_L} \mathbb{C}_\chi.$$

The *standard model* (taken for  $\beta = \beta_0$ ) is  $M_0 := \bigoplus_{y \in \kappa} \mathbb{C} \cdot e_y$ , with  $\{e_y\}$  declared orthonormal and

$$W(a, b) e_y := \psi(-b(y + a)) e_{y+a}, \quad \text{i.e.} \quad W(a, b) = Z(-b)X(a),$$

where  $X(a)e_y := e_{y+a}$  and  $Z(b)e_y := \psi(by)e_y$ . **This is a stipulation.**

*Scope.* The sign in  $W(a, b) = Z(-b)X(a)$  is not cosmetic: the naive assignment  $W(a, b) = Z(b)X(a)$ , without the sign, realizes a *different* cocycle ( $-ab'$ , not  $\beta_0$ 's  $ab'$ ), and the discrepancy between the two is invisible at  $p = 2$  (where  $-1 = 1$ ) and visible only at odd  $p$  — a characteristic-dependent failure mode this definition is written to avoid by fixing the correct sign explicitly rather than leaving it to convention. That  $M_0$ , so defined, genuinely satisfies the multiplication law of Definition 2.4 is verified as part of the proof of Theorem 3.5 below (Step 1) and checked computationally by gate C11 of `theory/checks/wh_kappa_check.py`.

---

**Provenance.** ids: D8; proved in: —; tested by: —; reviewed in: —.

**PROVED**

**Theorem 3.3** (Lagrangian lines, and the Schrödinger models they carry). *Let  $\kappa$ ,  $V(\kappa)$ ,  $\omega$ ,  $\beta \in \text{Adm}(\omega)$ ,  $\psi$ ,  $A_{\psi,\beta}(V)$ ,  $Q_\beta$  (Definition 3.1) and  $A_L$ ,  $M_{L,\chi}$  (Definition 3.2) be as above, for any finite field  $\kappa$ ,  $p = 2$  included. Then:*

- (a) *every  $\kappa$ -line  $L \subset V(\kappa)$  is  $\omega$ -isotropic ( $\omega(v, v') = 0$  for all  $v, v' \in L$ ), hence maximal isotropic; so Lagrangian line and line coincide as conditions on  $L \subset V(\kappa)$ , at every characteristic;*
- (b) *there are exactly  $q + 1 = |\mathbb{P}^1(\kappa)|$  such lines;*
- (c)  *$A_L$  is commutative of dimension  $q$ , and carries exactly  $q$  characters, forming a torsor under  $\hat{L} = \text{Hom}(L, \mathbb{C}^\times)$ ;*
- (d)  $\dim_{\mathbb{C}} M_{L,\chi} = q$ ;
- (e) **(precision, binding on this row.)** *A Schrödinger model is a pair  $(L, \chi)$ : there are  $q(q + 1)$  such pairs,  $q$  per line — but by Theorem 3.5 below, all of them are isomorphic as  $A_{\psi,\beta}(V)$ -modules. The count  $q(q + 1)$  is therefore a count of labels, not of isomorphism classes of module. Among the  $q + 1$  lines, only those with  $Q_\beta|_L \equiv 0$  carry the trivial character  $\chi \equiv 1$  (since  $\chi \equiv 1$  is a character of  $A_L$  iff  $Q_\beta(v_0)\kappa \subseteq \ker \psi$  iff  $Q_\beta(v_0) = 0$ , for  $L = \kappa v_0$ ), and how many lines have  $Q_\beta|_L \equiv 0$  is itself  $\beta$ -dependent: exactly 2 for the reference cocycle  $\beta_0$  (§4 computes it in general).*

*Scope.* This theorem does not assert that distinct pairs  $(L, \chi)$  give distinct, let alone non-isomorphic, modules — to the contrary, part (e) is precisely the observation that they do not, once Theorem 3.5 is available. It does not address the characteristic-two phenomenon in the values a character of  $A_L$  can take; that is Theorem 3.4 below. Nothing here depends on which  $\beta \in \text{Adm}(\omega)$  was chosen beyond the formula for  $Q_\beta$ .

**Provenance.** ids: D1, D4, D6, D8, WH-FORM, WH-ALG, WH-POL; proved in: theory/wh-kappa.md §4; tested by: theory/checks/wh\_kappa\_check.py (C7); reviewed in: theory/verdicts/wh-kappa-r1.md.

*Proof.* (a) Write  $L = \kappa v_0$ . For  $c, c' \in \kappa$ ,  $\omega(cv_0, c'v_0) = cc' \omega(v_0, v_0) = 0$  by  $\kappa$ -bilinearity (Theorem 2.6(a)) and the computed alternating property (Theorem 2.6(b)) — at  $p = 2$  antisymmetry alone would not give this, so the computed form of (b) is exactly what is used. Since  $\dim_\kappa V(\kappa) = 2$  and  $\omega$  is nondegenerate, no subspace of dimension 2 (i.e. all of  $V(\kappa)$ ) is isotropic, so a line is a maximal isotropic subspace: *every* line is automatically Lagrangian, and the isotropy condition singles out nothing among lines.

(b) The nonzero vectors of  $V(\kappa)$  number  $q^2 - 1$ , and each line contains  $q - 1$  of them, so there are  $(q^2 - 1)/(q - 1) = q + 1 = |\mathbb{P}^1(\kappa)|$  lines.

(c) By Definitions 2.4 and 3.1,  $W(cv_0)W(c'v_0) = \psi(\beta(cv_0, c'v_0))W((c+c')v_0) = \psi(cc'Q_\beta(v_0))W((c+c')v_0)$ , using  $\beta(cv_0, c'v_0) = cc'\beta(v_0, v_0) = cc'Q_\beta(v_0)$  by  $\kappa$ -bilinearity. The phase is symmetric in  $(c, c')$ , so  $A_L$  is commutative;  $\dim_{\mathbb{C}} A_L = |L| = q$  since  $\{W(l) : l \in L\}$  is linearly independent (distinct basis elements of  $A_{\psi, \beta}(V)$ ). *Characters exist:*  $c(c, c') := \psi(cc'Q_\beta(v_0))$  is a symmetric bi-additive, hence symmetric 2-cocycle, on  $(L, +)$ , a group of exponent  $p$ ; Fact 3.1 supplies  $\mu : L \rightarrow \mathbb{C}^\times$  with  $c(c, c') = \mu(c + c')/(\mu(c)\mu(c'))$ , and  $\chi(W(cv_0)) := \mu(c)^{-1}$  is then an algebra character of  $A_L$  (it converts the twisted product into the untwisted one). *Torsor:* for two characters  $\chi, \chi'$  of  $A_L$ , the map  $l \mapsto \chi'(W(l))/\chi(W(l))$  is a homomorphism  $L \rightarrow \mathbb{C}^\times$ , and conversely composing a character with any element of  $\hat{L}$  gives another character; the action is free (a homomorphism fixing every  $\chi(W(l))$  is trivial), and  $|\hat{L}| = q$  by Fact 3.1 ( $U = L$ ,  $d = m$ ).

(d)  $A_{\psi, \beta}(V)$  is free as a right  $A_L$ -module on any set of coset representatives  $v_1, \dots, v_q$  of  $L$  in  $V(\kappa)$ : the elements  $\{W(v_i)W(l) : l \in L\}$  are, up to the nonzero scalars  $\psi(\beta(v_i, l))$ , exactly the basis elements  $\{W(v_i + l) : l \in L\}$  of the coset  $v_i + L$ . So  $A_{\psi, \beta}(V) \cong A_L^{\oplus q}$  as right  $A_L$ -modules, and  $M_{L, \chi} = A_{\psi, \beta}(V) \otimes_{A_L} \mathbb{C}_\chi$  has dimension  $q^2/q = q$ .

(e) There are  $q$  characters per line by (c) and  $q + 1$  lines by (b), giving  $q(q + 1)$  pairs  $(L, \chi)$ ; that they are all isomorphic as  $A_{\psi, \beta}(V)$ -modules is Theorem 3.5(c) below (which uses only parts (a)–(d) of this theorem, not this part, so there is no circular dependency). For  $L = \kappa v_0$ ,  $\chi \equiv 1$  is a character of  $A_L$  iff  $c(c, c') = \psi(cc'Q_\beta(v_0)) \equiv 1$  iff  $Q_\beta(v_0)\kappa \subseteq \ker \psi$ ; since  $Q_\beta(v_0)\kappa$  is either  $\{0\}$  or all of  $\kappa$  (it is a  $\kappa$ -subspace, being the image of scalar multiplication by  $Q_\beta(v_0)$ ) and  $\ker \psi \neq \kappa$  (Fact 3.1), this holds iff  $Q_\beta(v_0) = 0$ . For  $\beta = \beta_0$ ,  $Q_{\beta_0}(a, b) = ab$  (Definition 3.1) vanishes exactly on the two coordinate lines  $\{a = 0\}$  and  $\{b = 0\}$ , so exactly 2 of the  $q + 1$  lines carry the trivial character.  $\square$

### 3.3 The characteristic-two phase phenomenon

**PROVED** <sup>†</sup>

**Theorem 3.4** (Forced order-four phases in the induced characters, at characteristic two). *Let  $L = \kappa v_0 \subset V(\kappa)$  be a line with  $Q_\beta(v_0) \neq 0$  (Definition 3.1), and let  $\chi$  be any character of  $A_L$  (Definition 3.2, Theorem 3.3(c)).*

- [ $p = 2$ .]  $\chi$  takes a value of exact multiplicative order 4 on some  $W(l)$ ,  $l \in L$ : the phases of the model are forced out of  $\mu_2 = \psi(\kappa)$  into  $\mu_4$ . For the reference cocycle  $\beta_0$ , the lines with  $Q_{\beta_0}|_L \equiv 0$  are exactly the two coordinate axes (Theorem 3.3(e)), so this phenomenon applies to  $q - 1$  of the  $q + 1$  lines.
- [ $p$  odd.] No such phenomenon occurs: every character of  $A_L$  can be, and by the construction of Theorem 3.3(c) is,  $\mu_p$ -valued.

*Scope.* This theorem says nothing about lines with  $Q_\beta|_L \equiv 0$  (there the trivial character is available and no order-4 value is forced), and nothing about the algebra  $A_{\psi, \beta}(V)$  globally — only about the values a character of the specific commutative subalgebra  $A_L$  can take. It does

not identify *which*  $\mu_4$  this is inside a larger structure; that question, and its relation to the level- $\mu_p$  Weyl frame, is taken up in §4.

---

**Provenance.** ids: D6, D8, WH-POL, WH-POL-2; proved in: theory/wh-kappa.md §4 step ⟨1⟩5, ⟨1⟩6; tested by: theory/checks/beta\_census.py (B6, #Q=0); reviewed in: theory/verdicts/wh-kappa-r1.md.

*Proof.* Write  $Q_0 := Q_\beta(v_0) \neq 0$  and  $\nu(c) := \chi(W(cv_0))$  for the character  $\chi$  constructed in the proof of Theorem 3.3(c);  $\nu(0) = 1$ , and setting  $c' = c$  in the cocycle relation there,  $\nu(c)^2 = \psi(c^2 Q_0) \nu(2c)$ .

[ $p = 2$ .] Here  $2c = 0$ , so  $\nu(2c) = \nu(0) = 1$  and  $\nu(c)^2 = \psi(c^2 Q_0)$ . The Frobenius map  $c \mapsto c^2$  is injective, hence bijective, on the finite field  $\kappa$ , so  $\{c^2 Q_0 : c \in \kappa\} = Q_0 \kappa = \kappa$  (as  $Q_0 \neq 0$ ). Since  $\psi$  is nontrivial,  $\psi(c^2 Q_0) = -1$  for some  $c \in \kappa$  — because  $-1$  is the unique nontrivial element of  $\mu_2 = \{1, -1\}$  and  $\psi(\kappa) = \mu_2$  by Fact 3.1 — giving  $\nu(c)^2 = -1$ , i.e.  $\nu(c)$  has exact order 4. The count of lines follows from Theorem 3.3(e).

[ $p$  odd.] Put  $\mu(c) := \psi(c^2 Q_0/2)$  (division by 2 makes sense,  $p$  being odd). Then, directly,  $\psi((c^2 + c'^2)Q_0/2) \psi(cc'Q_0) = \psi((c + c')^2 Q_0/2)$ , i.e.  $\mu(c)\mu(c')\psi(cc'Q_0) = \mu(c + c')$ :  $\mu$  already solves the cocycle relation of Theorem 3.3(c) with values entirely inside  $\psi(\kappa) = \mu_p$ , so no order-4 (or any value outside  $\mu_p$ ) is ever forced.  $\square$

### 3.4 Uniqueness of the finite Weyl representation

#### PROVED

**Theorem 3.5** (Uniqueness of the finite Weyl representation). *Let  $\kappa, \beta \in \text{Adm}(\omega)$ ,  $\psi \in X(\kappa)$ , and  $A_{\psi, \beta}(V)$  be as above. Then:*

- (a) *the induced module  $M_0$  of  $(L_0, \chi_0)$ ,  $L_0 := 0 \oplus \kappa$ ,  $\chi_0 \equiv 1$ , has basis  $\{e_y\}_{y \in \kappa}$  with  $W_{\beta_0}(a, b)e_y = \psi(-b(y+a))e_{y+a}$  — i.e. the stipulated standard model of Definition 3.2 genuinely satisfies the multiplication law of Definition 2.4;*
- (b) *every simple  $A_{\psi, \beta}(V)$ -module is unitarily equivalent to  $M_0$  (equivalently, to any  $M_{L, \chi}$ ), and an  $A_{\psi, \beta}(V)$ -linear map  $M_0 \rightarrow M_0$  is a scalar;*
- (c) *consequently  $H_\beta(\kappa)$  has, up to unitary equivalence, exactly one irreducible unitary representation with central character  $\psi$ , of dimension  $q$ , and every  $\mathbb{C}$ -algebra automorphism of  $A_{\psi, \beta}(V)$  is inner;*
- (d) *for  $\{e_y\}$  orthonormal, every  $W_{\beta_0}(v)$  acts unitarily, and in general the unitary intertwiner between two irreducible unitary representations with the same central character is unique up to a scalar in  $U(1)$ .*

*Scope.* This theorem is stated and proved without invoking Maschke’s theorem, Artin–Wedderburn, Skolem–Noether, Jacobson density, or Pontryagin duality as black boxes: every step is elementary, using only Theorem 2.8 (simplicity) and Theorem 3.3. It is uniform in the characteristic: no clause is restricted to odd  $p$  or to  $p = 2$ , and no step below treats  $p = 2$  differently. It does not exhibit the isomorphism  $A_{\psi, \beta}(V) \cong M_q(\mathbb{C})$  as canonical — that non-canonicity is the separate content of Theorem 3.6 immediately below.

---

**Provenance.** ids: D1, D4, D8, WH-ALG, WH-POL, WH-SVN; proved in: theory/wh-kappa.md §5; tested by: theory/checks/wh\_kappa\_check.py (C3, C6, C11); reviewed in: theory/verdicts/wh-kappa-r1.md; theory/verdicts/wh-kappa-adjudication.md.

*Proof.* (a) *The standard model, verified.* Since  $Q_{\beta_0}(0, b) = 0 \cdot b = 0$ ,  $\chi_0 \equiv 1$  is a character of  $A_{L_0}$  by Theorem 3.3(e). Take coset representatives  $(y, 0)$ ,  $y \in \kappa$ , of  $L_0$  in  $V(\kappa)$  and put  $e_y := W_{\beta_0}(y, 0) \otimes 1 \in M_{L_0, \chi_0}$ . Then

$$W_{\beta_0}(a, b) e_y = \psi(\beta_0((a, b), (y, 0))) W_{\beta_0}(a + y, b) \otimes 1 = W_{\beta_0}(a + y, b) \otimes 1,$$

since  $\beta_0((a, b), (y, 0)) = a \cdot 0 = 0$ . To evaluate this against  $\chi_0$  the  $L_0$ -part must be moved to the *right*:

$$\begin{aligned} W_{\beta_0}(a + y, 0) W_{\beta_0}(0, b) &= \psi((a + y)b) W_{\beta_0}(a + y, b), \quad \text{so} \\ W_{\beta_0}(a + y, b) &= \psi(-b(y + a)) W_{\beta_0}(a + y, 0) W_{\beta_0}(0, b). \end{aligned}$$

Tensoring against  $\chi_0(W_{\beta_0}(0, b)) = 1$  gives  $W_{\beta_0}(a, b)e_y = \psi(-b(y + a))e_{a+y}$  as claimed, and  $\dim M_0 = q$  by Theorem 3.3(d). (The sign here is exactly Definition 3.2's stipulation, now derived rather than assumed: the naive un-signed assignment  $W(a, b) = Z(b)X(a)$  realizes the *different* cocycle  $-ab'$  and fails to satisfy Definition 2.4's multiplication law for  $\beta_0(v, v') = ab'$  at odd  $p$  — gate C11 of `theory/checks/wh_kappa_check.py` reports 0 violations for the signed model at all six tested  $q \in \{2, 3, 4, 5, 8, 9\}$ , against 36, 400, 3888 violations for the unsigned model at  $q = 3, 5, 9$  and, tellingly, 0 violations at  $q = 2, 4, 8$  — the defect is invisible at  $p = 2$ , where  $-1 = 1$ , and visible only once  $p$  is odd.)

(b) *Uniqueness and Schur.*  $\pi : A_{\psi, \beta}(V) \rightarrow \text{End}_{\mathbb{C}}(M_0)$  is a unital, nonzero algebra homomorphism, so  $\ker \pi$  is a proper two-sided ideal, hence 0 by simplicity (Theorem 2.8):  $\pi$  is injective. Since  $\dim \text{End}_{\mathbb{C}}(M_0) = q^2 = \dim A_{\psi, \beta}(V)$ ,  $\pi$  is also surjective, giving  $A_{\psi, \beta}(V) \cong \text{End}_{\mathbb{C}}(M_0)$ . Consequently  $A_{\psi, \beta}(V) \cong M_0^{\oplus q}$  as a left module over itself (the columns of  $\text{End}_{\mathbb{C}}(M_0)$ ), so *any* simple module  $N$  — generated by any nonzero  $n \in N$  — is a quotient of  $A_{\psi, \beta}(V) \cong M_0^{\oplus q}$ ; restricting the quotient map to one summand gives a nonzero map  $M_0 \rightarrow N$ , which is surjective (its image is a nonzero submodule of the simple  $N$ ) and injective (its kernel is a proper submodule of the simple  $M_0$ ), hence an isomorphism  $M_0 \cong N$ . For an  $A$ -endomorphism  $\varphi$  of  $M_0$ :  $\varphi$ , as a  $\mathbb{C}$ -linear endomorphism of a finite-dimensional  $\mathbb{C}$ -vector space, has an eigenvalue  $\lambda \in \mathbb{C}$  ( $\mathbb{C}$  is algebraically closed), and  $\ker(\varphi - \lambda)$  is then a nonzero  $A$ -submodule of the simple  $M_0$ , so  $\varphi = \lambda \cdot \text{id} : \text{End}_A(M_0) = \mathbb{C} \cdot \text{id}$ .

(c) *Representations of  $H_{\beta}(\kappa)$  and inner automorphisms.* A representation of  $H_{\beta}(\kappa)$  with central character  $\psi$  is exactly an  $A_{\psi, \beta}(V)$ -module (the remark after Theorem 2.7); irreducibility is simplicity of the module, so (b) applies: there is exactly one such representation up to isomorphism, of dimension  $q$ . For automorphisms: given  $\alpha \in \text{Aut}_{\mathbb{C}}(A_{\psi, \beta}(V))$ , twist the action on  $M_0$  by  $\alpha$ , i.e. put  $a \cdot_{\alpha} m := \pi(\alpha(a))m$ ; this twisted module  $M_0^{\alpha}$  is again simple of dimension  $q$ , hence  $\cong M_0$  by (b), via some invertible  $T$ ; unwinding,  $\alpha = \text{Ad}_T$ , with  $T$  unique up to a scalar by the Schur clause of (b). No Skolem–Noether theorem is invoked.

(d) *Unitarity.* Since  $|\psi(\cdot)| = 1$  (Fact 3.1), each  $W_{\beta_0}(a, b)$  of (a) sends the orthonormal  $\{e_y\}$  to  $\{\psi(-b(y + a))e_{a+y}\}$ , again orthonormal: every  $W_{\beta_0}(v)$  is unitary for this inner product, with no averaging needed. (For a general  $\beta$ , the Weyl operators generate the *finite* level- $\mu_p$  frame  $\mathcal{F}_{\psi, \beta}^{(p)}$  of Definition 4.9 in §4, of order  $pq^2$ ; averaging any inner product on the simple module over this finite group makes every  $\pi(W_{\beta}(v))$  unitary there too.) If  $T$  intertwines two unitary representations sharing central character  $\psi$ , then  $T^*T$  commutes with every operator of the first, so  $T^*T = c \cdot \text{id}$  with  $c > 0$  by the Schur clause of (b); then  $c^{-1/2}T$  is unitary, and two unitary intertwiners differ by a unimodular scalar, i.e. an element of  $U(1)$ .  $\square$

## PROVED

**Theorem 3.6** (The twisted algebra as a matrix algebra, non-canonically).  $A_{\psi, \beta}(V) \cong M_q(\mathbb{C})$ , by an isomorphism that depends on a choice of Lagrangian line and character  $(L, \chi)$  (Definition 3.2) and a choice of basis of  $M_{L, \chi}$ , and is **not** canonical.

*Scope.* This theorem asserts existence of an isomorphism to  $M_q(\mathbb{C})$  and the dependence of any such isomorphism on auxiliary choices; it does not assert, and Theorem 6.3 of §6 shows precisely how much of this data can be removed, that any two such isomorphisms agree, nor that a canonical one exists.

**Provenance.** ids: D4, D8, WH-ALG, WH-SVN, WH-ALG-MAT; proved in: `theory/wh_kappa.md` §5 *step* {1}2; tested by: `theory/checks/wh_kappa_check.py` (C5); reviewed in: `theory/verdicts/wh_kappa-r1.md`.

*Proof.* This is exactly the isomorphism  $\pi : A_{\psi,\beta}(V) \xrightarrow{\sim} \text{End}_{\mathbb{C}}(M_0)$  constructed in the proof of Theorem 3.5(b); choosing the basis  $\{e_y\}_{y \in \kappa}$  of  $M_0$  identifies  $\text{End}_{\mathbb{C}}(M_0)$  with  $M_q(\mathbb{C})$ . The construction visibly used a Lagrangian line  $L_0$ , a character  $\chi_0$  of  $A_{L_0}$ , and the basis  $\{e_y\}$  of the resulting module; a different line, character, or basis gives a different isomorphism, related to this one by an inner automorphism of  $M_q(\mathbb{C})$  (Theorem 3.5(c)) composed with the change of basis, and no choice among these is distinguished by the data  $(\kappa, \psi, \beta)$  alone.  $\square$

## 4 The polarizing cocycle as a second datum: the characteristic-two dichotomy

This section proves the increment’s central structural fact. The quantum system attached to a finite field  $\kappa$  is built not from the field alone, and not from the field together with a single choice of additive character  $\psi$ : it is built from *three* data —  $\kappa$ ,  $\psi$ , and a polarizing cocycle  $\beta$ , a choice inside the torsor  $\text{Adm}(\omega)$  of Definition 2.2 that is no more optional, and no more a matter of convention, than  $\psi$  itself. What this section shows is exactly how much that third choice matters. At odd characteristic it turns out not to matter at all: a canonical representative exists inside  $\text{Adm}(\omega)$ , and every other admissible cocycle produces an isomorphic Heisenberg group by an isomorphism fixing the centre. At characteristic two no canonical representative exists, and the choice is not immaterial: the torsor  $\text{Adm}(\omega)$  splits into exactly two orbits under the natural symmetry group  $SL_2(\kappa)$ , distinguished by the Arf invariant of an associated quadratic form, and the two orbits produce two genuinely non-isomorphic Heisenberg groups — a fact detectable by an operator identity inside the quantum system itself.

This is not a defect in the construction, and not a gap for a better convention to close. It is a theorem about what the construction actually depends on, carrying an intrinsic, computable invariant that says exactly when two choices of cocycle agree and when they do not. A sharp result of this shape — naming precisely the choice at stake and exhibiting the invariant that classifies it — is a positive structural statement about the definition, not a negative one, and it is this labbook’s principal new content.

### 4.1 The torsor of admissible polarizing cocycles, restated

**Definition 4.1** (Admissible polarizing cocycles, restated in full). For the symplectic object  $(V(\kappa), \omega)$  of Definition 2.1,

$$\text{Adm}(\omega) := \{ \beta : V(\kappa) \times V(\kappa) \rightarrow \kappa \mid \beta \text{ is } \kappa\text{-bilinear and } \beta - \beta^T = \omega \}.$$

A *polarizing cocycle* is a choice of element  $\beta \in \text{Adm}(\omega)$ , and it is a **datum** of the construction, on the same footing as the phase datum  $\psi$  — not a convention fixed once and for all. The *reference cocycle* is  $\beta_0((a, b), (a', b')) := ab'$ , a label for a point of the torsor below, not a canonical point of it. Write  $\text{Sym}(V)$  for the set of symmetric  $\kappa$ -bilinear forms on  $V(\kappa)$ .

*Scope.* This restates Definition 2.2 verbatim; nothing new is stipulated here. What is new in this section is everything that is *proved* about  $\text{Adm}(\omega)$ : its torsor structure (Theorem 4.2), the immateriality of the choice at odd  $p$  (Theorem 4.3), and its genuine two-way split at  $p = 2$  (Theorems 4.4–4.8).

---

**Provenance.** ids: D1, D2; proved in: —; tested by: —; reviewed in: —.

**PROVED**

**Theorem 4.2** (The admissible cocycles form a torsor of size  $q^3$ ).  $\text{Adm}(\omega)$  is a torsor under  $\text{Sym}(V)$  acting by  $\beta \mapsto \beta + s$ ,  $s \in \text{Sym}(V)$ ; consequently  $|\text{Adm}(\omega)| = q^3$ .

*Scope.* This is purely a statement about the set  $\text{Adm}(\omega)$  and its free transitive action; it does not yet distinguish any point of the torsor, does not use the characteristic in any way, and does not address isomorphism of the associated Heisenberg groups — those are the later theorems of this section.

---

**Provenance.** ids: D1, D2, WH-BETA-a; proved in: theory/wh-kappa-choice.md §6 step ⟨1⟩1; tested by: theory/checks/beta\_census.py (B1); theory/checks/wh\_kappa\_check.py (C10); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* If  $\beta, \beta' \in \text{Adm}(\omega)$ , put  $s := \beta' - \beta$ . Then  $s - s^T = (\beta' - \beta^{T'}) - (\beta - \beta^T) = \omega - \omega = 0$ , i.e.  $s \in \text{Sym}(V)$ . Conversely, for  $s \in \text{Sym}(V)$ ,  $(\beta + s) - (\beta + s)^T = (\beta - \beta^T) + (s - s^T) = \omega + 0 = \omega$ , so  $\beta + s \in \text{Adm}(\omega)$ . The action  $\beta \mapsto \beta + s$  is therefore well defined, free (if  $\beta + s = \beta + s'$  then  $s = s'$ ), and transitive (any  $\beta' \in \text{Adm}(\omega)$  equals  $\beta + (\beta' - \beta)$  for the  $s = \beta' - \beta \in \text{Sym}(V)$  just exhibited). A symmetric  $\kappa$ -bilinear form on  $V(\kappa) = \kappa e \oplus \kappa f$  is determined freely by the three values  $s(e, e), s(e, f), s(f, f) \in \kappa$  (symmetry forces  $s(f, e) = s(e, f)$ , and bilinearity determines everything else), so  $|\text{Sym}(V)| = q^3$ ; since  $\text{Adm}(\omega)$  is nonempty ( $\beta_0 \in \text{Adm}(\omega)$ ) and a torsor under  $\text{Sym}(V)$ , it has the same cardinality,  $q^3$ .  $\square$

## 4.2 Odd characteristic: a canonical representative

**PROVED**<sup>†</sup>

**Theorem 4.3** (Immateriality of the polarizing cocycle at odd characteristic). *[p odd.]  $\text{Adm}(\omega)$  contains exactly one antisymmetric member,  $\omega/2$ , and it is  $SL_2(\kappa)$ -invariant. For every  $s \in \text{Sym}(V)$ , the map*

$$\varphi_s(t, v) := \left( t + \frac{1}{2}s(v, v), v \right)$$

*is a group isomorphism  $H_\beta(\kappa) \rightarrow H_{\beta+s}(\kappa)$  fixing the centre  $\kappa \times 0$  pointwise, for every  $\beta \in \text{Adm}(\omega)$ . Hence at odd characteristic the choice of polarizing cocycle is immaterial: every admissible cocycle produces a Heisenberg group isomorphic, by an explicit isomorphism fixing the centre, to every other, and  $\omega/2$  is a canonical basepoint of the torsor.*

*Scope.* This theorem is restricted to odd  $p$  by hypothesis, not by omission: at  $p = 2$ , division by 2 is division by zero,  $\omega/2$  has no referent, and Theorem 4.4 shows the phenomenon this theorem describes simply does not occur. Nothing here compares  $H_\beta(\kappa)$  for different  $\beta$  as algebras or as sources of quantum systems beyond the group isomorphism exhibited; the algebra-level statement is Theorem 4.10.

---

**Provenance.** ids: D2, D5, WH-BETA-a, WH-BETA-b; proved in: theory/wh-kappa-choice.md §6 step ⟨1⟩2; tested by: theory/checks/beta\_census.py (B2,B3,B4); theory/checks/wh\_kappa\_check.py (C10); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof. Existence and uniqueness of the antisymmetric point.* Since  $p$  is odd,  $2 \in \kappa^\times$ , and  $\omega/2$  is  $\kappa$ -bilinear with  $(\omega/2) - (\omega/2)^T = \omega/2 + \omega/2 = \omega$  (using  $\omega^T = -\omega$ , i.e. antisymmetry of  $\omega$ , itself immediate from Theorem 2.6(b):  $\omega(v, v') + \omega(v', v) = \omega(v + v', v + v') - \omega(v, v) - \omega(v', v') = 0$ ), so  $\omega/2 \in \text{Adm}(\omega)$  and is manifestly antisymmetric. If  $\beta, \beta' \in \text{Adm}(\omega)$  are both antisymmetric, then  $s := \beta' - \beta \in \text{Sym}(V)$  (Theorem 4.2) is also antisymmetric, so  $2s = 0$ , so  $s = 0$  ( $p$  odd): the antisymmetric point is unique. *Invariance.*  $SL_2(\kappa)$  preserves  $\omega$  (Theorem 2.6(e)), hence preserves the scalar multiple  $\omega/2$ .

*The isomorphism  $\varphi_s$ .* Put  $f(v) := s(v, v)/2$ . By bilinearity and symmetry of  $s$ ,  $s(v + v', v + v') = s(v, v) + 2s(v, v') + s(v', v')$ , so  $f(v + v') = f(v) + f(v') + s(v, v')$ . Unwinding the group law of Definition 2.5 for  $H_\beta(\kappa)$  and  $H_{\beta+s}(\kappa)$ ,

$$\varphi_s((t, v)(t', v')) = \varphi_s(t + t' + \beta(v, v'), v + v') = (t + t' + \beta(v, v') + f(v + v'), v + v'),$$

while

$$\varphi_s(t, v) \varphi_s(t', v') = (t + f(v), v) (t' + f(v'), v') = (t + t' + f(v) + f(v') + (\beta + s)(v, v'), v + v'),$$

computed inside  $H_{\beta+s}(\kappa)$ . The two right-hand sides agree precisely because  $f(v + v') = f(v) + f(v') + s(v, v')$ . So  $\varphi_s$  is a group homomorphism; it is a bijection (identity on the second coordinate, an affine bijection  $t \mapsto t + f(v)$  on the first, for each fixed  $v$ ), hence an isomorphism, and  $\varphi_s(t, 0) = (t + f(0), 0) = (t, 0)$  since  $f(0) = s(0, 0)/2 = 0$ : the centre is fixed pointwise.  $\square$

### 4.3 Characteristic two: no symmetric distinguished point

**PROVED** <sup>†</sup>

**Theorem 4.4** (Absence of a symmetric distinguished cocycle at characteristic two). *[ $p = 2$ .]  $\text{Adm}(\omega)$  contains no antisymmetric member; the criterion that singles out a canonical basepoint at odd characteristic (Theorem 4.3) distinguishes nothing here.*

*Scope.* This theorem only rules out *this particular* route to a canonical choice (antisymmetry). It does not by itself assert that no other route to a canonical choice exists, nor that the resulting Heisenberg groups actually differ — those stronger, positive facts are Theorems 4.5–4.7 below.

---

**Provenance.** ids: D2, WH-BETA-a, WH-BETA-c; proved in: theory/wh-kappa-choice.md §6 step ⟨1⟩3; tested by: theory/checks/beta\_census.py (B2); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* At  $p = 2$ ,  $-1 = 1$ , so “antisymmetric” ( $\beta = -\beta^T$ ) coincides with “symmetric” ( $\beta = \beta^T$ ), i.e.  $\beta - \beta^T = 0$ . But every  $\beta \in \text{Adm}(\omega)$  has  $\beta - \beta^T = \omega$  (Definition 4.1), and  $\omega((1, 0), (0, 1)) = 1 \neq 0$  (Theorem 2.6(c)), so  $\omega \neq 0$ : no member of  $\text{Adm}(\omega)$  can be symmetric, hence none can be antisymmetric either.  $\square$

### 4.4 The quadratic form of a cocycle, and the order profile of its Heisenberg group

**PROVED** <sup>†</sup>

**Theorem 4.5** (The quadratic form of a polarizing cocycle, and the element-order profile of its Heisenberg group). *[ $p = 2$ .] Let  $\beta \in \text{Adm}(\omega)$ , with  $Q_\beta(v) := \beta(v, v)$  as in Definition 3.1. Then:*

- (i)  $Q_\beta(cv) = c^2 Q_\beta(v)$  for  $c \in \kappa$ ,  $v \in V(\kappa)$ , and  $Q_\beta(v + v') = Q_\beta(v) + Q_\beta(v') + \omega(v, v')$  for all  $v, v'$ :  $Q_\beta$  is a  $\kappa$ -quadratic form on  $V(\kappa)$  with polar form  $\omega$ ;
- (ii) in  $H_\beta(\kappa)$ ,  $(t, v)^2 = (Q_\beta(v), 0)$  for every  $(t, v)$ ; consequently, writing  $n_0 := |Q_\beta^{-1}(0)|$  (which is always  $\geq 1$ , since  $Q_\beta(0) = 0$ ), the multiset of element orders of  $H_\beta(\kappa)$  is

$$\{ 1 : 1, \quad 2 : q n_0 - 1, \quad 4 : q^3 - q n_0 \};$$

- (iii) hence  $\text{Adm}(\omega)$  meets at least two isomorphism classes of Heisenberg group: by direct enumeration at  $q = 2$ , 6 of the 8 admissible cocycles give the order profile  $\{1:1, 2:5, 4:2\}$  ( $n_0 = 3$ ) and the remaining 2 give  $\{1:1, 2:1, 4:6\}$  ( $n_0 = 1$ ), and these are different multisets, hence non-isomorphic groups.

*Scope.* Part (iii) establishes only that *at least* two isomorphism classes occur, by exhibiting two different order profiles at the single value  $q = 2$ ; it does not claim that exactly two classes occur for every  $q$ , nor identify the classes by name (this lane computes order profiles, not group names — that two of the eight cocycles at  $q = 2$  happen to give the quaternion group and the other six the dihedral group of order 8 is a fact the orchestrator’s independent census records, but it is not part of what this row proves). The full classification, valid at every  $q = 2^m$  and carrying an intrinsic invariant, is Theorem 4.7 below, which depends on this one.

---

**Provenance.** ids: D2, D6, WH-BETA-a, WH-BETA-d; proved in: theory/wh-kappa-choice.md §6 step ⟨1⟩4; tested by: theory/checks/beta\_census.py (B3,B6); theory/checks/wh\_kappa\_check.py (C10); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* (i) By  $\kappa$ -bilinearity,  $Q_\beta(cv) = \beta(cv, cv) = c^2\beta(v, v) = c^2Q_\beta(v)$ . Expanding  $Q_\beta(v + v') = \beta(v + v', v + v') = \beta(v, v) + \beta(v, v') + \beta(v', v) + \beta(v', v')$ ; at  $p = 2$ , addition and subtraction coincide, so  $\beta(v, v') + \beta(v', v) = \beta(v, v') - \beta(v', v) = \omega(v, v')$  (Definition 4.1), giving  $Q_\beta(v + v') = Q_\beta(v) + Q_\beta(v') + \omega(v, v')$ .

(ii) In  $H_\beta(\kappa)$ ,  $(t, v)(t, v) = (2t + \beta(v, v), 2v) = (Q_\beta(v), 0)$  at  $p = 2$ , since  $2t = 0$  and  $2v = 0$ . If  $(t, v) = (0, 0)$  this is the identity (order 1). If  $(t, v) \neq (0, 0)$  and  $Q_\beta(v) = 0$ , then  $(t, v)^2 = (0, 0)$ , so  $(t, v)$  has order exactly 2; the number of such elements is  $qn_0 - 1$  (for each of the  $n_0$  vectors  $v$  with  $Q_\beta(v) = 0$ , all  $q$  values of  $t$  occur, and the pair  $(t, v) = (0, 0)$  is excluded). If  $Q_\beta(v) \neq 0$ , then  $(t, v)^2 = (Q_\beta(v), 0) \neq (0, 0)$ , but  $(t, v)^4 = ((t, v)^2)^2 = (2Q_\beta(v), 0) = (0, 0)$ , so  $(t, v)$  has order exactly 4; there are  $q(q^2 - n_0) = q^3 - qn_0$  such elements ( $q^2 - n_0$  choices of  $v$ ,  $q$  choices of  $t$  each). These three counts sum to  $1 + (qn_0 - 1) + (q^3 - qn_0) = q^3 = |H_\beta(\kappa)|$ , as they must.

(iii) The order profile just derived is a group-isomorphism invariant (an isomorphism of groups preserves the multiset of element orders), so different profiles certify non-isomorphic groups. At  $q = 2$ , exhaustive enumeration over all 8 members of  $\text{Adm}(\omega)$  (theory/checks/beta\_census.py, gates B3 and B6) computes  $n_0$  for each and finds exactly the two values  $n_0 = 3$  (for 6 of the 8 cocycles, giving profile  $\{1:1, 2:5, 4:2\}$  by the formula of (ii)) and  $n_0 = 1$  (for the remaining 2, giving  $\{1:1, 2:1, 4:6\}$ ); these are distinct multisets, exhibiting two isomorphism classes.  $\square$

## 4.5 The Arf classification

**Definition 4.6** (The Artin–Schreier map, and the type of a polarizing cocycle). For  $\kappa$  of characteristic 2, define  $\wp : \kappa \rightarrow \kappa$  by  $\wp(x) := x^2 + x$ . The *type* of  $\beta \in \text{Adm}(\omega)$  is

$$\text{Arf}(Q_\beta) := Q_\beta(e)Q_\beta(f) \in \kappa/\wp(\kappa),$$

computed in any symplectic  $\kappa$ -basis  $(e, f)$  of  $V(\kappa)$ , i.e. a basis with  $\omega(e, f) = 1$ .

*Scope.* This is a stipulation, meaningful only at  $p = 2$ . That the value  $\text{Arf}(Q_\beta)$  does not depend on the choice of symplectic basis, that  $\kappa/\wp(\kappa)$  has exactly two elements, and that both values actually occur among the members of  $\text{Adm}(\omega)$ , are not part of the definition; they are exactly the content of Theorem 4.7 below, whose provenance names the argument DAG’s current row for this material (the single source’s own cross-reference for this content predates a later renaming in the DAG).

---

**Provenance.** ids: D10; proved in: —; tested by: —; reviewed in: —.

### SKETCH

**Theorem 4.7** (Classification of characteristic-two polarizing cocycles by the Arf invariant). *[p = 2.] Let  $\beta \in \text{Adm}(\omega)$ .*

- (i)  $\wp$  is additive with  $\ker \wp = \{0, 1\} = \mathbb{F}_2$ , so  $\kappa/\wp(\kappa)$  has exactly two elements;
- (ii)  $Q_\beta$  has a nonzero zero (some  $v \neq 0$  with  $Q_\beta(v) = 0$ ) if and only if  $\text{Arf}(Q_\beta) = 0$ ; in that case  $n_0 = |Q_\beta^{-1}(0)| = 2q - 1$ , and otherwise  $n_0 = 1$ . Consequently  $\text{Arf}(Q_\beta)$  does not depend on the choice of symplectic basis used to compute it, and is invariant under  $SL_2(\kappa)$ ;
- (iii) call  $\beta$  hyperbolic if  $\text{Arf}(Q_\beta) = 0$  and anisotropic otherwise. Two members of  $\text{Adm}(\omega)$  are carried to each other by an  $SL_2(\kappa)$ -isometry of their quadratic forms if and only if they

have the same type; both types occur, with  $q(q+1)/2$  hyperbolic and  $q(q-1)/2$  anisotropic quadratic forms compatible with  $\omega$ , and stabilizer orders  $|O(Q_\beta) \cap SL_2(\kappa)| = 2(q-1)$  (hyperbolic) and  $2(q+1)$  (anisotropic);

(iv)  $H_\beta(\kappa) \cong H_{\beta'}(\kappa)$  if and only if  $\text{Arf}(Q_\beta) = \text{Arf}(Q_{\beta'})$ . So  $\text{Adm}(\omega)$  meets exactly two isomorphism classes of Heisenberg group, indexed by  $\text{Arf}$ .

*Scope.* This theorem is marked SKETCH for a procedural reason recorded in `theory/verdicts/wh-kappa-adjudication.md`: it was written in the repair wave that followed this increment's single hostile-critic round and has not itself been through one, per this campaign's capped review loop (`PRD.md`) — not because a specific gap is known. It does not extend beyond  $p = 2$  (Theorem 4.3 covers odd  $p$  by an entirely different, and there, complete, argument), and it does not address whether the type can be chosen compatibly along field embeddings — that question is recorded as open in §8.

---

**Provenance.** ids: D6, D10, WH-BETA-d, WH-BETA-TYPE; proved in: `theory/wh-kappa-choice.md` §6 step  $\langle 1 \rangle 4, \langle 1 \rangle 5$ ; tested by: `theory/checks/beta_census.py` (B5, B6, B7); `theory/checks/wh_kappa_check.py` (C10); reviewed in: `theory/verdicts/wh-kappa-adjudication.md`.

*Proof.* (i)  $\wp(x+y) - \wp(x) - \wp(y) = (x+y)^2 - x^2 - y^2 = 2xy = 0$  at  $p = 2$ , so  $\wp$  is additive (a group homomorphism  $(\kappa, +) \rightarrow (\kappa, +)$ ). Its kernel consists of the roots of  $x^2 + x = 0$ , a degree-2 polynomial, so  $|\ker \wp| \leq 2$ ; both  $x = 0$  and  $x = 1$  are roots, so  $\ker \wp = \{0, 1\} = \mathbb{F}_2$  exactly. By rank-nullity for the  $\mathbb{F}_2$ -linear map  $\wp$  on the  $m$ -dimensional  $\mathbb{F}_2$ -space  $\kappa$ ,  $|\wp(\kappa)| = q/2$ , so  $|\kappa/\wp(\kappa)| = 2$ .

(ii) In a symplectic basis  $(e, f)$ , write  $v = ae + bf$ ; by Theorem 4.5(i),  $Q_\beta(ae + bf) = \alpha a^2 + ab + \delta b^2$  with  $\alpha := Q_\beta(e)$ ,  $\delta := Q_\beta(f)$  (using  $\omega(e, f) = 1$  for the cross term). If  $b = 0$ :  $Q_\beta = \alpha a^2$  vanishes for some  $a \neq 0$  iff  $\alpha = 0$ , and then  $\text{Arf} = \alpha\delta = 0 \in \wp(\kappa)$ . If  $b \neq 0$ , rescale to  $b = 1$ :  $\alpha a^2 + a + \delta = 0$ ; multiplying by  $\alpha$  (when  $\alpha \neq 0$ ) gives  $(\alpha a)^2 + (\alpha a) = \alpha\delta$ , i.e.  $\wp(\alpha a) = \alpha\delta$ , solvable for  $a$  iff  $\alpha\delta \in \wp(\kappa)$ ; when  $\alpha = 0$  the equation  $a + \delta = 0$  always has the solution  $a = \delta$ . So in every case,  $Q_\beta$  has a nonzero zero iff  $\alpha\delta \in \wp(\kappa)$ , i.e. iff  $\text{Arf}(Q_\beta) = 0 \in \kappa/\wp(\kappa)$ . When  $\text{Arf}(Q_\beta) = 0$ , the zero set of  $Q_\beta$  is a union of  $\kappa$ -lines (by homogeneity  $Q_\beta(cv) = c^2 Q_\beta(v)$  of Theorem 4.5(i)); the computation above produces exactly two such lines (the two roots of  $\wp(\alpha a) = \alpha\delta$  when  $\alpha \neq 0$ , or the lines  $b = 0$  and  $a = \delta$  when  $\alpha = 0$ ), giving  $n_0 = 2(q-1)+1 = 2q-1$  nonzero-vector zeros plus the origin; when  $\text{Arf}(Q_\beta) \neq 0$ ,  $n_0 = 1$  (only the origin). Since  $|\kappa/\wp(\kappa)| = 2$  by (i), the class  $\alpha\delta \bmod \wp(\kappa)$  is itself a basis-free condition — “ $Q_\beta$  is isotropic” — so  $\text{Arf}(Q_\beta)$  is independent of the symplectic basis chosen to compute it, and  $SL_2(\kappa)$ -invariant since an  $SL_2(\kappa)$ -image of a symplectic basis is again symplectic (Theorem 2.6(e)).

(iii) *Normal forms.* If  $\text{Arf}(Q_\beta) = 0$ , pick  $e \neq 0$  with  $Q_\beta(e) = 0$  (by (ii)) and any  $f$  with  $\omega(e, f) = 1$  (possible by nondegeneracy of  $\omega$ ); replacing  $f$  by  $f + Q_\beta(f)e$  changes  $Q_\beta(f)$  to  $Q_\beta(f) + Q_\beta(f)^2 Q_\beta(e) + Q_\beta(f)\omega(f, e)$ , which simplifies (using  $Q_\beta(e) = 0$  and  $\omega(f, e) = -\omega(e, f) = 1$  at  $p = 2$ ) to 0, giving  $Q_\beta(ae + bf) = ab$ : the hyperbolic normal form. If  $\text{Arf}(Q_\beta) \neq 0$  then  $Q_\beta(v) \neq 0$  for every  $v \neq 0$  (by (ii)); pick any  $e \neq 0$ , rescale by  $\lambda = Q_\beta(e)^{-1/2}$  (Frobenius is bijective on  $\kappa$ , so square roots exist and are unique) to arrange  $Q_\beta(e) = 1$ , then choose  $f$  with  $\omega(e, f) = 1$ ; replacing  $f$  by  $f + ae$  changes  $Q_\beta(f)$  by  $\wp(a)$  (direct computation from Theorem 4.5(i)), so  $Q_\beta(f)$  can be moved to any chosen representative of its class modulo  $\wp(\kappa)$ . Both normalizations send a symplectic basis to a symplectic basis, hence are realized by elements of  $SL_2(\kappa)$  (Theorem 2.6(e)): two quadratic forms of the same type are  $SL_2(\kappa)$ -isometric, and by (ii) two of different type cannot be (they have different  $n_0$ , an isometry invariant). *Counting.* Over a fixed symplectic basis, the  $\kappa$ -quadratic forms with polar form  $\omega$  are exactly the  $q^2$  maps  $\alpha a^2 + ab + \delta b^2$  (parametrized freely by  $(\alpha, \delta) \in \kappa^2$ , by Theorem 4.5(i)); among these,  $\#\{\alpha\delta \in \wp(\kappa)\} = q + (q-1)(q/2) = q(q+1)/2$  are hyperbolic (the case  $\alpha = 0$  contributes  $q$  choices of  $\delta$ , and each of the  $q-1$  nonzero  $\alpha$  pairs with the  $q/2$  elements of  $\wp(\kappa)$ ) and

the remaining  $q(q-1)/2$  are anisotropic — both classes nonempty for every  $q$ . By orbit-stabilizer inside  $SL_2(\kappa)$ , of order  $q(q^2-1)$ , acting transitively within each type: stabilizer orders  $q(q^2-1)/(q(q+1)/2) = 2(q-1)$  (hyperbolic) and  $q(q^2-1)/(q(q-1)/2) = 2(q+1)$  (anisotropic). *Groups.* If  $\text{Arf}(Q_\beta) = \text{Arf}(Q_{\beta'})$ , the normal-form argument gives  $Q_{\beta'} = Q_\beta \circ g$  for some  $g \in SL_2(\kappa)$ ; then  $r(v, v') := \beta(gv, gv') - \beta'(v, v')$  is symmetric with  $r(v, v) = 0$  for all  $v$  (both sides equal  $Q_{\beta'}(v)$  by construction of  $g$ ), so, fixing an  $\mathbb{F}_2$ -basis  $b_i$  of  $V(\kappa)$  and writing  $v = \sum c_i b_i$ ,  $h(v) := \sum_{i < j} c_i c_j r(b_i, b_j)$  satisfies  $h(v + v') - h(v) - h(v') = r(v, v')$  (a direct expansion using  $r$  symmetric and bilinear), and  $(t, v) \mapsto (t + h(v), gv)$  is then a group isomorphism  $H_{\beta'}(\kappa) \rightarrow H_\beta(\kappa)$  by the same computation as in the proof of Theorem 4.3. If instead  $\text{Arf}(Q_\beta) \neq \text{Arf}(Q_{\beta'})$ , the two groups have different values of  $n_0$  (by (ii)), hence different element-order profiles (Theorem 4.5(ii)), so no group isomorphism of any kind exists between them.  $\square$

*Remark.* Direct enumeration (`theory/checks/beta_census.py`, gates B5–B7) confirms every clause above at  $q = 2, 4, 8$ :  $\text{Arf}$  is constant on  $SL_2(\kappa)$ -orbits and takes exactly the two values across all  $q^3$  members of  $\text{Adm}(\omega)$ ;  $|O(Q_\beta) \cap SL_2(\kappa)| = 2, 6, 14$  (hyperbolic) and  $6, 10, 18$  (anisotropic), matching  $2(q \mp 1)$  exactly; the element-order profiles come in exactly two sizes,  $(6, 2)$ ,  $(40, 24)$ ,  $(288, 224)$ , at  $q = 2, 4, 8$  respectively; and the isomorphism constructed in the proof of (iii) is verified on all pairs within the hyperbolic class.

## 4.6 The type, detected inside the quantum system

### SKETCH

**Theorem 4.8** (The Arf type is visible inside the quantum system). *[ $p = 2$ .] Let  $\beta \in \text{Adm}(\omega)$  and  $\psi$  a phase datum. Then, inside  $A_{\psi, \beta}(V)$ :*

$$W_\beta(v)^2 = \psi(Q_\beta(v)) \cdot 1 \quad \text{for every } v \in V(\kappa),$$

and

$$q^{-1} \sum_{v \in V(\kappa)} W_\beta(v)^2 = \varepsilon \cdot 1, \quad \varepsilon = \begin{cases} +1 & \beta \text{ hyperbolic } (\text{Arf}(Q_\beta) = 0), \\ -1 & \beta \text{ anisotropic.} \end{cases}$$

Equivalently, identifying  $\kappa/\wp(\kappa) \cong \mathbb{F}_2$  via the trace  $\text{Tr}_{\kappa/\mathbb{F}_2}$  of Definition 2.3,  $\varepsilon = (-1)^{\text{Tr}(\text{Arf}(Q_\beta))}$ .

*Scope.* The identity above is an equation inside the algebra  $A_{\psi, \beta}(V)$  itself, so it exhibits  $\text{Arf}(Q_\beta)$  as an invariant not merely of the abstract group  $H_\beta(\kappa)$  but of the level- $\mu_p$  Weyl frame  $\mathcal{F}_{\psi, \beta}^{(p)}$  of Definition 4.9 below. As with Theorem 4.7, on which it depends, the SKETCH label reflects only that this material postdates the increment's single hostile-critic round, per `theory/verdicts/wh-kappa-adjudication.md`.

---

**Provenance.** ids: D3, D4, D6, D10, WH-BETA-EPS; proved in: `theory/wh-kappa-choice.md` §6 *step*  $\langle 1 \rangle 6$ ; tested by: `theory/checks/beta_census.py` (B6, B8); reviewed in: `theory/verdicts/wh-kappa-adjudication.md`.

*Proof. The square identity.* By Definition 2.4 and Definition 3.1,  $W_\beta(v)^2 = \psi(\beta(v, v))W_\beta(2v) = \psi(Q_\beta(v))W_\beta(0) = \psi(Q_\beta(v)) \cdot 1$ , using  $2v = 0$  at  $p = 2$ .

*The hyperbolic value.* For the hyperbolic normal form  $Q(a, b) = ab$  (Theorem 4.7(iii)),  $\sum_{a, b \in \kappa} \psi(ab) = \sum_a \left( \sum_b \psi(ab) \right)$ ; for  $a \neq 0$ ,  $b \mapsto \psi(ab)$  is a nontrivial character of  $\kappa$ , so the inner sum vanishes by Fact 3.1; for  $a = 0$  the inner sum is  $q$ . So the double sum is  $q$ , giving  $\varepsilon = +1$  for this normal form. Since  $\sum_v \psi(Q_\beta(v))$  is invariant under precomposing  $Q_\beta$  with any bijection of  $V(\kappa)$ , in particular under the  $SL_2(\kappa)$ -isometries of Theorem 4.7(iii),  $\varepsilon = +1$  for *every* hyperbolic  $\beta$ , not only the normal form.

*The anisotropic value.* Sum  $\sum_v \psi(Q(v))$  over all  $q^2$  quadratic forms  $Q(a, b) = \alpha a^2 + ab + \delta b^2$  compatible with  $\omega$  (Theorem 4.7(iii)) and exchange the order of summation:

$$\sum_{\alpha, \delta \in \kappa} \sum_{a, b \in \kappa} \psi(\alpha a^2 + ab + \delta b^2) = \sum_{a, b \in \kappa} \psi(ab) \left( \sum_{\alpha} \psi(\alpha a^2) \right) \left( \sum_{\delta} \psi(\delta b^2) \right).$$

By Fact 3.1,  $\sum_{\alpha} \psi(\alpha a^2) = q$  if  $a = 0$  (else the character  $\alpha \mapsto \psi(\alpha a^2)$  is nontrivial, since squaring is injective on  $\kappa$ , and the sum vanishes); likewise for the  $\delta$ -sum. So only the term  $a = b = 0$  survives, and the total is  $q^2 \cdot \psi(0) = q^2$ . By Theorem 4.7(iii) there are  $q(q+1)/2$  hyperbolic forms, each contributing  $q$  (just shown), and  $q(q-1)/2$  anisotropic forms, each contributing the same value  $S$  (by isometry-invariance, exactly as for the hyperbolic case). So  $\frac{q(q+1)}{2} \cdot q + \frac{q(q-1)}{2} \cdot S = q^2$ , giving  $S = \frac{q^2 - \frac{q^2(q+1)}{2}}{\frac{q(q-1)}{2}} = \frac{q^2(1 - \frac{q+1}{2})}{\frac{q(q-1)}{2}} = \frac{-q^2(q-1)/2}{q(q-1)/2} = -q$ . So  $\varepsilon = -1$  on the anisotropic class.

*Reading  $\varepsilon$  through the trace.*  $\text{Tr}_{\kappa/\mathbb{F}_2}$  is additive and  $\mathbb{F}_2$ -valued;  $\text{Tr}(\wp(x)) = \text{Tr}(x^2) + \text{Tr}(x) = \text{Tr}(x) + \text{Tr}(x) = 0$  (using  $\text{Tr}(x^2) = \text{Tr}(x)$ , since Frobenius permutes the terms of the trace sum, and  $2 \text{Tr}(x) = 0$  at  $p = 2$ ), so  $\text{Tr}$  kills  $\wp(\kappa)$ ; it is nonzero (e.g. on a normal-basis generator), hence onto  $\mathbb{F}_2$ , and  $\dim_{\mathbb{F}_2} \wp(\kappa) = m - 1 = \dim_{\mathbb{F}_2} \ker \text{Tr}$  by the rank computation of Theorem 4.7(i), so  $\wp(\kappa) = \ker \text{Tr}$  exactly:  $\text{Tr}$  descends to an isomorphism  $\kappa/\wp(\kappa) \xrightarrow{\sim} \mathbb{F}_2$ . Under this identification the hyperbolic class ( $\text{Arf} = 0$ ) corresponds to  $\text{Tr} = 0$  and  $\varepsilon = +1 = (-1)^0$ , and the anisotropic class to  $\text{Tr} = 1$  and  $\varepsilon = -1 = (-1)^1$ .  $\square$

## 4.7 The level statement: what the algebra forgets, and what it does not

**Definition 4.9** (The level- $\mu_p$  frame).

$$\mathcal{F}_{\psi, \beta}^{(p)} := \{ \zeta^j W_{\beta}(v) : j \in \mathbb{Z}/p, v \in V(\kappa) \} \subset A_{\psi, \beta}(V)^{\times},$$

the finite group of Weyl unitaries with phases in  $\mu_p = \psi(\kappa)$  (Fact 3.1), for  $\zeta$  a primitive  $p$ -th root of unity as in Definition 2.3.

*Scope.*  $\mathcal{F}_{\psi, \beta}^{(p)}$  is recorded separately from the Weyl frame  $\mathcal{F}_{\beta} := \{ \mathbb{C}^{\times} W_{\beta}(v) : v \in V(\kappa) \} \subset A_{\psi, \beta}(V)$  (the set of  $\mathbb{C}^{\times}$ -lines through Weyl operators; studied in §5 below) because the two behave differently under a change of  $\beta$ , exactly as Theorem 4.10 makes precise: the pair  $(A_{\psi, \beta}(V), \mathcal{F}_{\beta})$  turns out not to depend on  $\beta$  at all, while  $\mathcal{F}_{\psi, \beta}^{(p)}$  — equivalently the isomorphism class of  $H_{\beta}(\kappa)$  itself — does, and only at  $p = 2$ .

---

**Provenance.** ids: D11; proved in: —; tested by: —; reviewed in: —.

### SKETCH

**Theorem 4.10** (The algebra and its Weyl frame do not depend on  $\beta$ ; the level- $\mu_p$  frame does, at  $p = 2$ ). *For all  $\beta, \beta' \in \text{Adm}(\omega)$  there is an algebra isomorphism  $A_{\psi, \beta}(V) \rightarrow A_{\psi, \beta'}(V)$  of the form  $W_{\beta}(v) \mapsto \mu(v) W_{\beta'}(v)$  for some  $\mu : V(\kappa) \rightarrow \mathbb{C}^{\times}$  — a frame isomorphism, carrying  $\mathcal{F}_{\beta}$  to  $\mathcal{F}_{\beta'}$  — and this holds **at every characteristic**. Moreover:*

- $\mu$  can be chosen  $\mu_p$ -valued if and only if  $\psi \circ Q_{\beta} = \psi \circ Q_{\beta'}$ ;
- $[p = 2]$   $\mu$  can always be chosen  $\mu_4$ -valued, regardless of type.

Hence the pair  $(A_{\psi, \beta}(V), \mathcal{F}_{\beta})$  does not depend on  $\beta$  at all, while the level- $\mu_p$  frame  $\mathcal{F}_{\psi, \beta}^{(p)}$  — equivalently the isomorphism class of  $H_{\beta}(\kappa)$  (Theorem 4.7) — does depend on  $\beta$ , and does so only at  $p = 2$ .

*Scope.* This theorem does not contradict Theorems 4.4–4.8: it is a statement about the algebra  $A_{\psi,\beta}(V)$  and its  $\mathbb{C}^\times$ -frame  $\mathcal{F}_\beta$ , which turn out to be  $\beta$ -independent, while those theorems are about the finite group  $H_\beta(\kappa)$  (equivalently, the level- $\mu_p$  frame), which is not. The distinction between the two frames is exactly what makes both statements true simultaneously. This row is SKETCH for the same procedural reason as Theorems 4.7–4.8, and the explicit  $\mu_2/\mu_4$ -valued formulas at  $p = 2$  are presented at the level of detail the source shard gives them, verified computationally rather than re-derived symbol-by-symbol here.

**Provenance.** ids: D2, D4, D11, WH-BETA-a, WH-BETA-d, WH-BETA-LEVEL; proved in: theory/wh-kappa-choice.md §6 step {1}7; tested by: theory/checks/frame\_mu.py (M1,M2,M3,M4); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* Put  $s := \beta' - \beta \in \text{Sym}(V)$  (Theorem 4.2). Direct expansion of the multiplication law of Definition 2.4 shows  $v \mapsto \mu(v)W_{\beta'}(v)$  is an algebra map exactly when

$$\frac{\mu(v)\mu(v')}{\mu(v+v')} = \psi(-s(v, v')) \quad (\dagger)$$

for all  $v, v'$ , and any such map automatically carries  $\mathcal{F}_\beta$  to  $\mathcal{F}_{\beta'}$ , since it sends each line  $\mathbb{C}^\times W_\beta(v)$  to the line  $\mathbb{C}^\times W_{\beta'}(v)$ . The right-hand side  $\psi \circ (-s)$  is a symmetric 2-cocycle on  $(V(\kappa), +)$ , a group of exponent  $p$  (every element of the characteristic- $p$  field  $\kappa$ , hence of  $V(\kappa)$ , has additive order dividing  $p$ ), so Fact 3.1 supplies a solution  $\mu$  of  $(\dagger)$ : **the frame isomorphism exists at every characteristic**, with no restriction on  $\beta, \beta'$ .

*Necessity of the  $\mu_p$  criterion.* Normalize  $\mu(0) = 1$  (forced by  $(\dagger)$  at  $v = v' = 0$ , since  $\mu(0)^2/\mu(0) = \mu(0) = \psi(0) = 1$ ). At  $p = 2$ , setting  $v' = v$  in  $(\dagger)$  gives  $\mu(v)^2/\mu(0) = \mu(v)^2 = \psi(s(v, v)) = \psi(Q_{\beta'}(v) - Q_\beta(v))$ . So  $\mu$  is  $\mu_2$ -valued (i.e.  $\mu(v)^2 = 1$  for every  $v$ ) if and only if  $\psi(Q_{\beta'}(v)) = \psi(Q_\beta(v))$  for every  $v$ , i.e. iff  $\psi \circ Q_\beta = \psi \circ Q_{\beta'}$ ; at odd  $p$  the analogous computation with  $2 \in \kappa^\times$  gives the same criterion for  $\mu_p$ -valuedness in general.

*Sufficiency,  $p = 2$ .* Fix an  $\mathbb{F}_2$ -basis  $b_1, \dots, b_{2m}$  of  $V(\kappa)$  and write  $\psi(s(b_i, b_j)) = (-1)^{\sigma_{ij}}$ ,  $\sigma_{ij} \in \{0, 1\}$ , for the finitely many structure values of the symmetric form  $s$  on this basis (well defined since  $\psi(\kappa) = \mu_2 = \{\pm 1\}$  at  $p = 2$ , Fact 3.1). If  $\sigma_{ii} = 0$  for every  $i$  (the case  $\psi \circ Q_\beta = \psi \circ Q_{\beta'}$  just characterized), the  $\mathbb{F}_2$ -bilinear cochain  $\ell(v) := \sum_{i < j} c_i c_j \sigma_{ij}$ , for  $v = \sum_i c_i b_i$ , solves  $(\dagger)$  with values in  $\mu_2 = \{\pm 1\}$ , by a direct check against the defining bilinear expansion of  $s$ . In general — without assuming  $\sigma_{ii} = 0$  — the integer-valued cochain  $m(v) := 2 \sum_{i < j} c_i c_j \sigma_{ij} + \sum_i c_i \sigma_{ii} \pmod{4}$  solves  $(\dagger)$  with  $\mu(v) := i^{m(v)} \in \mu_4$  ( $i = \sqrt{-1}$ ): this is the explicit construction of theory/wh-kappa-choice.md §6, verified against all  $|V(\kappa)|^2$  pairs  $(v, v')$  for every one of the  $q^3$  members of  $\text{Adm}(\omega)$  at  $q = 2, 4, 8$  by gates M2 and M3 of theory/checks/frame\_mu.py (which finds  $\mu_4$ -valued solutions always, and  $\mu_2$ -valued solutions for exactly the  $q$  pairs  $(\beta, \beta')$  with  $\psi \circ Q_\beta = \psi \circ Q_{\beta'}$ , matching the necessity criterion above exactly).

*Sufficiency, odd  $p$ .*  $\mu(v) := \psi(s(v, v)/2)$  solves  $(\dagger)$ : since  $s$  is symmetric bilinear,  $s(v, v) + s(v', v') - s(v + v', v + v') = -2s(v, v')$ , so  $\mu(v)\mu(v')/\mu(v + v') = \psi([s(v, v) + s(v', v') - s(v + v', v + v')]/2) = \psi(-s(v, v'))$ , matching  $(\dagger)$  exactly, with values entirely inside  $\psi(\kappa) = \mu_p$ : this reproves, independently of Theorem 4.3, that nothing leaves the level- $\mu_p$  frame once  $p$  is odd — a second, algebra-level confirmation that the choice of  $\beta$  is immaterial there.

Gate M1 of theory/checks/frame\_mu.py additionally searches all  $2^{q^2}$  candidate  $\mu_2$ -valued phase functions exhaustively at  $q = 2$  and confirms solutions exist exactly for the pairs the necessity criterion predicts; gate M4 checks the odd- $p$  formula at  $q = 3, 5, 9$ .  $\square$

## 5 Symmetry of the Weyl frame: the Weil representation and its splitting problem

**Recollection.**  $\kappa = \mathbb{F}_q$ ,  $q = p^m$ ;  $V(\kappa) = \kappa \oplus \kappa$  with  $\omega((a, b), (a', b')) = ab' - a'b$  (Definition 2.1);  $SL_2(\kappa)$  is the group of  $\kappa$ -linear automorphisms of  $V(\kappa)$  preserving  $\omega$  (Theorem 2.6(e)); a polar-

izing cocycle  $\beta \in \text{Adm}(\omega)$  and phase datum  $\psi$  determine the algebra  $A_{\psi,\beta}(V) = \bigoplus_v \mathbb{C} \cdot W_\beta(v)$  (Definitions 4.1, 2.3, 2.4), and  $Q_\beta(v) := \beta(v, v)$  (Definition 3.1).

---

**Provenance.** ids: D1, D2, D3, D4, D6; proved in: —; tested by: —; reviewed in: —.

This section studies the symmetries of the Weyl frame itself: which permutations of the Weyl operators extend to algebra automorphisms, how large that symmetry group is, and whether  $SL_2(\kappa)$  — the natural symmetry group of  $(V(\kappa), \omega)$  — genuinely acts on the quantum system by honest algebra automorphisms (as opposed to merely projectively). The answer is uniform in one respect and characteristic-dependent in another: the group of frame automorphisms always surjects onto  $SL_2(\kappa)$  with kernel the dual group  $\hat{V}$ , at every characteristic; but whether this extension *splits* — whether  $SL_2(\kappa)$  itself, not just a projective shadow of it, acts by algebra automorphisms — is settled affirmatively at odd  $p$  and only partially resolved at  $p = 2$ , where it interacts with the Arf dichotomy of §4.

## 5.1 The Weyl frame and its two automorphism groups

**Definition 5.1** (The Weyl frame, and its  $\kappa$ -linear automorphisms). The *Weyl frame* is

$$\mathcal{F} := \{ \mathbb{C}^\times \cdot W_\beta(v) : v \in V(\kappa) \} \subset A_{\psi,\beta}(V),$$

the set of  $\mathbb{C}^\times$ -lines through Weyl operators.  $\text{Aut}_{\mathcal{F}}(A)$  is the group of  $\mathbb{C}$ -algebra automorphisms  $\alpha$  of  $A_{\psi,\beta}(V)$  with  $\alpha(\mathcal{F}) = \mathcal{F}$ .  $\text{Aut}_{\mathcal{F}}^\kappa(A) \subseteq \text{Aut}_{\mathcal{F}}(A)$  is the subgroup consisting of those  $\alpha$  whose induced permutation of  $V(\kappa)$  (well defined since  $\alpha$  permutes the lines of  $\mathcal{F}$ , which are indexed bijectively by  $V(\kappa)$ ) is  $\kappa$ -linear.

*Scope.* The  $\kappa$ -linearity clause in the definition of  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  is **imposed data**: nothing in  $A_{\psi,\beta}(V)$  or  $\mathcal{F}$  as built from  $(V, +)$  and the cocycle  $\psi \circ \beta$  forces an automorphism's induced permutation of  $V(\kappa)$  to respect scalar multiplication by  $\kappa$  rather than merely by  $\mathbb{F}_p$ . Precisely how much symmetry this clause discards is Theorem 7.1 of §7; nothing here addresses that question.

---

**Provenance.** ids: D7; proved in: —; tested by: —; reviewed in: —.

## 5.2 Exactness of the frame-automorphism extension, uniformly in the characteristic

Fix  $\beta \in \text{Adm}(\omega)$ . For  $g \in SL_2(\kappa)$ , write  $s_g(v, v') := \beta(gv, gv') - \beta(v, v')$  and  $Q_g(v) := s_g(v, v)$ ; note that  $Q_g(v) = Q_\beta(gv) - Q_\beta(v)$ .

**PROVED**

**Theorem 5.2** (Exactness of the frame-automorphism extension). *At every characteristic, the sequence*

$$1 \longrightarrow \hat{V} \longrightarrow \text{Aut}_{\mathcal{F}}^\kappa(A_{\psi,\beta}(V)) \xrightarrow{\alpha \mapsto g} SL_2(\kappa) \longrightarrow 1$$

*is exact, where  $\hat{V} := \text{Hom}((V(\kappa), +), \mathbb{C}^\times)$ ; the kernel consists exactly of the conjugations  $\text{Ad}_{W_\beta(u)}$ ,  $u \in V(\kappa)$ .*

*Scope.* This theorem does not assert that the extension splits — that  $SL_2(\kappa)$  itself embeds back into  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  as a subgroup — only that it is exact. Splitting is addressed separately, and with a genuinely different answer at odd and even characteristic, in Theorems 5.3 and 5.6 below. Nothing here concerns  $\text{Aut}_{\mathcal{F}}(A)$  without the imposed  $\kappa$ -linearity clause; that is Theorem 7.1.

---

**Provenance.** ids: D1, D4, D7, WH-ALG, WH-COMM, WH-WEIL-a; proved in: theory/wh-kappa-choice.md §8 step ⟨1⟩2, ⟨1⟩3; tested by: theory/checks/split24.py (X2); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.*  $s_g$  is symmetric bi-additive, and  $s_{gh}(v, v') = s_h(v, v') + s_g(hv, hv')$ . Symmetry:  $s_g(v, v') - s_g(v', v) = \omega(gv, gv') - \omega(v, v') = 0$ , since  $g \in SL_2(\kappa)$  preserves  $\omega$  (Theorem 2.6(e)); the telescoping identity is immediate by expanding  $\beta(ghv, ghv') - \beta(v, v')$  through the intermediate term  $\beta(hv, hv')$ . (For  $\beta = \beta_0$  and  $g = \begin{pmatrix} r & s \\ t & u \end{pmatrix}$ , direct expansion of  $\beta_0(gv, gv') = rt aa' + ru ab' + st a'b + su bb'$  gives the explicit formula  $s_g(v, v') = rt aa' + st(ab' + a'b) + su bb'$ , using  $ru - st = \det g = 1$ .)

*Description of  $\text{Aut}_{\mathcal{F}}^\kappa(A)$ .* Frame preservation gives  $\alpha(W_\beta(v)) = \lambda(v)W_\beta(gv)$  for some function  $\lambda : V(\kappa) \rightarrow \mathbb{C}^\times$  and some permutation  $g$  of  $V(\kappa)$ , imposed to be  $\kappa$ -linear by definition of  $\text{Aut}_{\mathcal{F}}^\kappa(A)$ . Applying  $\alpha$  to  $W_\beta(v)W_\beta(v') = \psi(\beta(v, v'))W_\beta(v + v')$  and comparing basis elements forces  $g$  additive (automatic here, being  $\kappa$ -linear) and

$$\lambda(v)\lambda(v')\psi(s_g(v, v')) = \lambda(v + v') \quad (*)$$

for all  $v, v'$ . Applying  $\alpha$  to the commutation relation of Theorem 2.7(c) and using  $\omega(gv, gv') = \det(g)\omega(v, v')$  (Theorem 2.6(e)'s proof) gives  $\psi((\det g - 1)\omega(v, v')) = 1$  for all  $v, v'$ ; since  $\omega$  is onto  $\kappa$  (Theorem 2.6(a),(c)), Fact 3.1 forces  $\det g = 1$ , i.e.  $g \in SL_2(\kappa)$ . So  $\alpha \mapsto g$  is a well-defined map  $\text{Aut}_{\mathcal{F}}^\kappa(A) \rightarrow SL_2(\kappa)$ , visibly a homomorphism (composition of automorphisms composes the induced permutations).

*The kernel.* If  $g = \text{id}$ ,  $(*)$  reads  $\lambda(v)\lambda(v') = \lambda(v + v')$ , i.e.  $\lambda \in \hat{V}$ , a group of order  $q^2$  by Fact 3.1 ( $U = V(\kappa)$ ,  $d = 2m$ ). The map  $u \mapsto \psi(\omega(u, \cdot))$ ,  $V(\kappa) \rightarrow \hat{V}$ , is injective (if  $\psi(\omega(u, \cdot)) \equiv 1$  then  $\omega(u, \cdot) \equiv 0$  by Fact 3.1, so  $u = 0$  by nondegeneracy, Theorem 2.6(c)), hence bijective ( $|V(\kappa)| = q^2 = |\hat{V}|$ ); so every kernel element has  $\lambda = \psi(\omega(u, \cdot))$  for a unique  $u \in V(\kappa)$ , and by Theorem 2.7(d) this is exactly  $\text{Ad}_{W_\beta(u)}$ .

*Surjectivity, uniformly in  $p$ .* For fixed  $g \in SL_2(\kappa)$ ,  $c(v, v') := \psi(s_g(v, v'))$  is a symmetric 2-cocycle on the exponent- $p$  group  $(V(\kappa), +)$  (symmetry of  $c$  from symmetry of  $s_g$ , just shown; the cocycle identity from bi-additivity of  $s_g$ ), so Fact 3.1 supplies  $\lambda$  solving  $(*)$ : every  $g \in SL_2(\kappa)$  is hit. This argument is uniform in the characteristic; nothing distinguishes  $p = 2$ .  $\square$

*Remark* (Every frame automorphism is projectively unitary, at every characteristic). Fix a model  $(M, \pi)$  as in Theorem 3.5. Taking absolute values in  $(*)$  makes  $v \mapsto |\lambda(v)|$  a homomorphism from the finite group  $V(\kappa)$  to the torsion-free group  $(\mathbb{R}_{>0}, \times)$ , hence trivial: every  $\lambda$  occurring in  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  is automatically unimodular. By Theorem 3.5(c),(d), every  $\alpha \in \text{Aut}_{\mathcal{F}}^\kappa(A)$  is  $\text{Ad}_T$  for some  $T$ , unique up to a scalar; since  $\alpha$  permutes the *unitary* Weyl operators (unimodular  $\lambda$  times unitary  $\pi(W_\beta(gv))$ ),  $T$  may be chosen unitary and is then unique up to  $U(1)$  (Theorem 3.5(d)). This gives an injective homomorphism  $\text{Aut}_{\mathcal{F}}^\kappa(A) \rightarrow PU(M)$ : a projective unitary representation of  $\text{Aut}_{\mathcal{F}}^\kappa(A)$ , at every characteristic. Composed with a section  $SL_2(\kappa) \rightarrow \text{Aut}_{\mathcal{F}}^\kappa(A)$  of Theorem 5.2's surjection — should one exist as an honest group homomorphism, i.e. should the extension split — this yields an honest (non-projective) representation of  $SL_2(\kappa)$  by algebra automorphisms, and hence, via  $\text{Ad}$ , a projective unitary representation of  $SL_2(\kappa)$  itself on  $M$ . What is *not* true is that this extends to all of  $\text{Aut}_{\mathbb{C}}(A) \cong PGL_q(\mathbb{C})$ : at  $q = 2$ ,  $T = \text{diag}(2, 1)$  conjugates  $\pi(W(1, 0)) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  to  $\begin{pmatrix} 0 & 2 \\ 1/2 & 0 \end{pmatrix}$ , whose singular values 2,  $1/2$  differ, so it is not a scalar times a unitary — the frame hypothesis in Definition 5.1 is exactly what is needed above.

### 5.3 Splitting at odd characteristic

**PROVED** <sup>†</sup>

**Theorem 5.3** (The Weil representation of  $SL_2(\kappa)$ , at odd characteristic). *[ $p$  odd.]  $\alpha_g(W_\beta(v)) := \psi(Q_g(v)/2)W_\beta(gv)$  defines a splitting  $g \mapsto \alpha_g$  of the extension of Theorem 5.2:  $SL_2(\kappa)$  acts on  $A_{\psi, \beta}(V)$  by  $\mathbb{C}$ -algebra automorphisms, with phases in  $\mu_p$ . Composed with Remark 5.2, this exhibits a projective unitary representation of  $SL_2(\kappa)$  on the model space of Theorem 3.5.*

*Scope.* This is an odd-characteristic statement only; Theorem 5.6 addresses whether an analogous honest (non-projective) action of  $SL_2(\kappa)$  by algebra automorphisms exists at  $p = 2$ , and the

answer there is open in general. Nothing here claims the splitting is unique, or canonical among splittings.

---

**Provenance.** ids: D4, D6, WH-WEIL-a, WH-SVN, WH-WEIL-b; proved in: theory/wh-kappa-choice.md §8 step ⟨1⟩4, ⟨1⟩7; tested by: theory/checks/split24.py (X4); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* Since  $s_g$  is symmetric bilinear,  $Q_g(v + v') = Q_g(v) + Q_g(v') + 2s_g(v, v')$  (the same polarization identity as in Definition 3.1's use elsewhere), so

$$\psi(Q_g(v + v')/2) = \psi(Q_g(v)/2) \psi(Q_g(v')/2) \psi(s_g(v, v')),$$

which is exactly relation (\*) of Theorem 5.2's proof for  $\lambda(v) := \psi(Q_g(v)/2)$ :  $\alpha_g$  is a well-defined element of  $\text{Aut}_{\mathcal{F}}^{\kappa}(A)$  over  $g$ . That  $g \mapsto \alpha_g$  is a homomorphism follows from the telescoping identity  $s_{gh}(v, v') = s_h(v, v') + s_g(hv, hv')$  of Theorem 5.2's proof, evaluated at  $v = v'$  to give  $Q_{gh}(v) = Q_h(v) + Q_g(hv)$ , whence  $\alpha_g(\alpha_h(W_{\beta}(v))) = \psi(Q_h(v)/2) \psi(Q_g(hv)/2) W_{\beta}(ghv) = \psi(Q_{gh}(v)/2) W_{\beta}(ghv) = \alpha_{gh}(W_{\beta}(v))$ . This is an honest algebra automorphism, so  $SL_2(\kappa)$  acts by algebra automorphisms with  $\mu_p$  phases (division by 2 makes sense since  $p$  is odd); the projective unitary statement is Remark 5.2 applied to this section.  $\square$

## 5.4 Characteristic two: the orthogonal subgroup and forced order-four phases

**PROVED** <sup>†</sup>

**Theorem 5.4** ( $\mu_2$ -phases occur exactly over the orthogonal subgroup of  $Q_{\beta}$ ). [ $p = 2$ .] Write  $O(Q_{\beta}) := \{g \in SL_2(\kappa) : Q_{\beta} \circ g = Q_{\beta}\}$ . A frame automorphism  $\alpha_{g,\lambda}$  (in the notation  $\alpha(W_{\beta}(v)) = \lambda(v)W_{\beta}(gv)$  of Theorem 5.2's proof) has some choice of  $\lambda$  landing entirely in  $\psi(\kappa) = \mu_2$  if and only if  $g \in O(Q_{\beta})$  — a condition on the chosen  $\beta$ , not on the characteristic alone. For the reference cocycle  $\beta_0$ ,  $O(Q_{\beta_0}) \cap SL_2(\kappa)$  consists exactly of the matrices  $\text{diag}(r, r^{-1})$  and  $\text{antidiag}(s, s^{-1})$ ,  $r, s \in \kappa^{\times}$ ; it has order  $2(q-1)$  (equal to 2, 6, 14 at  $q = 2, 4, 8$ ), and is a **proper** subgroup of  $SL_2(\kappa)$  for every  $q$ .

*Scope.* This theorem is about which  $g$  admit a  $\mu_2$ -valued lift, for a fixed  $\beta$ ; it says nothing yet about what happens for  $g \notin O(Q_{\beta})$  (Theorem 5.5), and the concrete description of  $O(Q_{\beta_0})$  is specific to the reference cocycle — for a general  $\beta$  of hyperbolic type the order is the same,  $2(q-1)$  (Theorem 4.7(iii)), but the concrete matrices differ with the normal-form basis chosen there.

---

**Provenance.** ids: D6, WH-WEIL-a, D7, WH-WEIL-c; proved in: theory/wh-kappa-choice.md §8 step ⟨1⟩5; tested by: theory/checks/beta\_census.py (B6); theory/checks/fp\_symmetry.py (S3); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* Setting  $v' = v$  in (\*) at  $p = 2$  ( $2v = 0$ ) gives  $\lambda(v)^2 = \psi(Q_g(v))$ . So a solution  $\lambda$  of (\*) is  $\mu_2$ -valued (i.e.  $\lambda(v)^2 = 1$  for every  $v$ ) iff  $Q_g(V(\kappa)) \subseteq \ker \psi$ , i.e. iff  $\psi \circ Q_g \equiv 1$ .

For every  $\beta$ , this is already the exact isometry condition.  $Q_g(v) = Q_{\beta}(gv) - Q_{\beta}(v)$  is additive in  $v$ : expanding  $Q_{\beta}(g(v + v')) - Q_{\beta}(v + v')$  via Definition 3.1's polarization identity on both terms, the  $\omega$ -cross-terms cancel because  $g$  preserves  $\omega$  (Theorem 2.6(e)), leaving  $Q_g(v + v') = Q_g(v) + Q_g(v')$ ; also  $Q_g(cv) = c^2 Q_g(v)$  since  $g$  is  $\kappa$ -linear. So, in a basis,  $Q_g(a, b) = \alpha a^2 + \delta b^2$  for some  $\alpha, \delta \in \kappa$  (no cross term). If  $\psi \circ Q_g \equiv 1$  then  $\psi(\alpha a^2) = 1$  for all  $a$  (setting  $b = 0$ ); since  $a \mapsto a^2$  is bijective on  $\kappa$ , this forces  $\alpha = 0$  by Fact 3.1, and likewise  $\delta = 0$ :  $Q_g \equiv 0$ , i.e.  $g \in O(Q_{\beta})$ . Conversely, if  $g \in O(Q_{\beta})$  then trivially  $\psi \circ Q_g \equiv 1$ , and moreover a  $\mu_2$ -valued  $\lambda$  solving (\*) then exists:  $\psi \circ s_g$  is a symmetric cocycle with trivial diagonal ( $\psi(s_g(v, v)) = \psi(Q_g(v)) = 1$ ), exactly the case handled by the  $\ell$ -construction in the proof of Theorem 4.10 (§4). So  $\psi \circ Q_g \equiv 1$  iff  $g \in O(Q_{\beta})$  iff a  $\mu_2$ -valued lift exists.

*The reference cocycle.* By the explicit formula of Theorem 5.2's proof,  $s_g(v, v') = rt aa' + st(ab' + a'b) + subb'$  for  $\beta_0$ , so  $Q_g(a, b) = rt a^2 + sub^2$  (the cross term drops at  $p = 2$ ). Thus  $g \in O(Q_{\beta_0})$  iff  $rt = 0$  and  $su = 0$ . Combined with  $\det g = ru - st = 1$ : if  $r = 0$  then  $st = -1 = 1$  (so  $s, t \neq 0$ ), and  $su = 0$  forces  $u = 0$ , giving  $g = \begin{pmatrix} 0 & s \\ s^{-1} & 0 \end{pmatrix}$  (using  $t = s^{-1}$  from  $st = 1$ ) — an antidiagonal matrix; if  $t = 0$  then  $ru = 1$  (so  $r, u \neq 0$ ,  $u = r^{-1}$ ), and  $su = 0$  forces  $s = 0$ , giving  $g = \text{diag}(r, r^{-1})$ ; the case  $r = t = 0$  contradicts  $\det g = 1$ . So  $O(Q_{\beta_0}) \cap SL_2(\kappa) = \{\text{diag}(r, r^{-1})\} \cup \{\text{antidiag}(s, s^{-1})\}$ , of order  $(q-1) + (q-1) = 2(q-1)$  (each family is parametrized bijectively by  $\kappa^\times$ ). Since  $Q_{\beta_0}(a, b) = ab$  is the hyperbolic normal form (Theorem 4.7(iii)), this matches the general formula  $2(q-1)$  there. Properness:  $2(q-1) < q(q^2-1) = |SL_2(\kappa)|$  for every  $q \geq 2$ , since  $q(q+1) \geq 6 > 2$ .  $\square$

**PROVED**<sup>†</sup>

**Theorem 5.5** ( $\mu_4$  is forced exactly when the orthogonal subgroup is proper). [ $p = 2$ .] *If  $g \notin O(Q_\beta)$ , every frame automorphism over  $g$  carries a phase of exact multiplicative order 4. Consequently  $\mu_4$  is forced somewhere in  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  exactly when  $O(Q_\beta) \cap SL_2(\kappa)$  is a proper subgroup of  $SL_2(\kappa)$ : this holds for  $\beta_0$  at every  $q$  (Theorem 5.4), and for every  $\beta \in \text{Adm}(\omega)$  at  $q = 4, 8$ , by direct enumeration. It does **not** hold in the single case  $q = 2$  with  $\beta$  anisotropic, where  $O(Q_\beta) \cap SL_2(\mathbb{F}_2) = SL_2(\mathbb{F}_2)$  (order  $6 = |O(Q_\beta) \cap SL_2(\kappa)|$  from the anisotropic formula  $2(q+1)$  of Theorem 4.7(iii) at  $q = 2$ ) and the entire group  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  admits  $\mu_2$ -valued phases throughout. So  $\mu_4$  is **not** a characteristic-two invariant of  $(V(\kappa), \omega, \psi)$  alone: whether it is forced depends on the type of  $\beta$ .*

*Scope.* This theorem uses only Theorem 5.4 and the comparison of  $2(q \mp 1)$  against  $|SL_2(\kappa)| = q(q^2 - 1)$ ; it does not itself re-derive the general- $q$  order  $2(q+1)$  for the anisotropic stabilizer (that is Theorem 4.7(iii), only cited here). The claim for *every*  $\beta$  at  $q = 4, 8$  rests on direct enumeration rather than a closed-form argument for general  $q$ .

---

**Provenance.** ids: D6, D10, WH-WEIL-c, WH-WEIL-d; proved in: theory/wh-kappa-choice.md §8 step <1>6; tested by: theory/checks/split24.py (X4, X6, and the anisotropic cross-run at  $q = 2$ ); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* By Theorem 5.4's proof, any  $\lambda$  solving  $(*)$  over  $g$  satisfies  $\lambda(v)^2 = \psi(Q_g(v))$  for every  $v$ . If  $g \notin O(Q_\beta)$  then  $Q_g \neq 0$ , and (by the same bijective-squaring argument as there)  $\psi(Q_g(v)) = -1$  for some  $v$  — since  $Q_g(v) = -1$ 's preimage under  $\psi$  being nontrivial requires only that  $\psi(\kappa) = \mu_2 = \{1, -1\}$  (Fact 3.1) and  $Q_g$  nonzero, hence onto a nonzero multiple of  $\kappa$ , hits every value including one where  $\psi = -1$ . Then  $\lambda(v)^2 = -1$ , i.e.  $\lambda(v)$  has exact order 4: *every*  $\lambda$  solving  $(*)$  over this  $g$  (not just one choice) is forced to take an order-4 value at this particular  $v$ , since the equation  $\lambda(v)^2 = \psi(Q_g(v)) = -1$  has no solution in  $\mu_2$ . Comparing  $2(q-1)$  and  $2(q+1)$  (Theorem 4.7(iii)) against  $|SL_2(\kappa)| = q(q^2-1) = q(q-1)(q+1)$ :  $2(q-1) < q(q-1)(q+1)$  always (as  $q(q+1) \geq 6$  for  $q \geq 2$ ), so the hyperbolic stabilizer — in particular  $O(Q_{\beta_0}) \cap SL_2(\kappa)$  — is always proper, giving some  $g \notin O(Q_\beta)$  and hence a forced order-4 phase, at every  $q$ . For the anisotropic stabilizer,  $2(q+1) < q(q-1)(q+1)$  iff  $2 < q(q-1)$ : this holds for  $q \geq 4$  but *fails* at  $q = 2$ , where  $q(q-1) = 2$  exactly, so  $2(q+1) = 6 = q(q-1)(q+1) = |SL_2(\mathbb{F}_2)|$ : the anisotropic stabiliser is the whole group at  $q = 2$ , so *no*  $g$  lies outside  $O(Q_\beta)$ , so no order-4 phase is forced by this argument, and — since  $\mu_2$ -valued lifts exist for every  $g \in O(Q_\beta)$  by Theorem 5.4's proof, and  $\hat{V}$  is automatically  $\mu_2$ -valued (Fact 3.1, exponent  $p = 2$ ) — the whole of  $\text{Aut}_{\mathcal{F}}^\kappa(A)$  admits  $\mu_2$ -valued phases in this one case. At  $q = 4, 8$ , exhaustive enumeration over every  $\beta \in \text{Adm}(\omega)$  (theory/checks/split24.py, gates X4 and X6) confirms  $O(Q_\beta) \cap SL_2(\kappa)$  is proper for both types, hence  $\mu_4$  is forced for every  $\beta$  there.  $\square$

## 5.5 Does the extension split at characteristic two? A conjecture, and what is known

### CONJECTURE

**Conjecture 5.6** (Existence of a Weil representation of  $SL_2(\kappa)$  at characteristic two). *[ $p = 2$ .] The extension of Theorem 5.2 splits for every  $q = 2^m$ :  $SL_2(\kappa)$  acts on  $A_{\psi,\beta}(V)$  by  $\mathbb{C}$ -algebra automorphisms, for some choice of splitting. This is verified for the reference cocycle  $\beta_0$  at  $q = 2$  (exhaustive search finds exactly 4 distinct complements of  $\hat{V}$  of order 6, over all generating pairs) and at  $q = 4$  (64 lifts of a  $(2, 3, 5)$ -presentation of  $SL_2(\mathbb{F}_4)$  satisfy the defining relations and generate a subgroup of order 60 meeting  $\hat{V}$  trivially); in both verified cases every complement necessarily uses phases of exact order 4 (no complement avoids  $\mu_4$ , since  $|\hat{V}| \cdot |O(Q_{\beta_0}) \cap SL_2(\kappa)|$  does not divide  $|\text{Aut}_{\mathcal{F}}^{\kappa}(A)|$ 's  $\mu_2$ -part suitably:  $6 \nmid 8$  at  $q = 2$ ,  $60 \nmid 96$  at  $q = 4$ ). No argument is known for general  $q$ .*

*Scope.* This is recorded as a conjecture, not a theorem, precisely because the evidence is two finite instances and no general argument: per this campaign's discipline, a passing computational instance promotes nothing. The verified cases are for the reference cocycle  $\beta_0$  only; splitting for other cocycles, and any relationship between splittings for cocycles of different Arf type, is not addressed. Should a splitting be found for general  $q$ , Remark 5.2 would then give a projective unitary representation of  $SL_2(\kappa)$  at  $p = 2$  exactly as Theorem 5.3 does at odd  $p$ .

---

**Provenance.** ids: D7, WH-WEIL-a, WH-WEIL-d, WH-WEIL-e; proved in: —; tested by: theory/checks/split24.py (X4,X5,X6); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*What is known, stated as evidence and not as a proof.* At  $q = 2$ ,  $|\text{Aut}_{\mathcal{F}}^{\kappa}(A)| = q^2 |SL_2(\kappa)| = 4 \cdot 6 = 24$ ; exhaustive search over generating pairs of order-6 subgroups meeting  $\hat{V} = \mathbb{Z}/2 \times \mathbb{Z}/2$  trivially finds exactly 4 complements, all using a phase of exact order 4 somewhere (split24.py gate X4). At  $q = 4$ , lifting a  $(2, 3, 5)$  generator-relation presentation of  $SL_2(\mathbb{F}_4) \cong A_5$  over all  $16 \times 16 = 256$  candidate phase choices on the two generators finds exactly 64 lifts satisfying  $x^2 = y^3 = (xy)^5 = 1$  and generating a subgroup of order 60 meeting  $\hat{V}$  trivially, again always with order-4 phases (gate X5). That no complement can avoid  $\mu_4$  in either case is confirmed by comparing group orders directly (gate X6): the  $\mu_2$ -phase part of  $\text{Aut}_{\mathcal{F}}^{\kappa}(A)$  has order  $q^2 \cdot |O(Q_{\beta_0}) \cap SL_2(\kappa)| = 4 \cdot 2 = 8$  at  $q = 2$  and  $16 \cdot 6 = 96$  at  $q = 4$ , and  $|SL_2(\kappa)| = 6$ , respectively 60, does not divide either. This settles the existence question affirmatively at these two finite values and settles that any splitting there must use order-4 phases; it establishes nothing for  $q = 2^m$ ,  $m \geq 3$ .

## 6 The choices the construction depends on, and what is canonical

**Recollection.**  $\kappa = \mathbb{F}_q$ ;  $\psi \in X(\kappa)$  a phase datum,  $\beta \in \text{Adm}(\omega)$  a polarizing cocycle (Definitions 2.3, 4.1);  $A_{\psi,\beta}(V)$  the resulting twisted algebra, simple of dimension  $q^2$  (Theorem 2.8), with a unique simple unitary module up to unitary equivalence (Theorem 3.5); by Theorem 4.10, for fixed  $\psi$  every two choices of  $\beta$  give algebras related by an explicit frame isomorphism.

---

**Provenance.** ids: D2, D3, D4, WH-ALG, WH-SVN; proved in: —; tested by: —; reviewed in: —.

Sections 4 and 5 named the choices the construction depends on and measured exactly how much they matter. This section assembles that information into a single statement: precisely which data beyond  $\kappa$  the construction needs, and precisely what survives canonically once those choices are made. The headline fact is that the *projective* Hilbert space is canonical — it does

not depend on which unitary model, which polarization, or (granting one more ingredient from §4) which polarizing cocycle was used to build it — even though the Hilbert space itself is not.

**Definition 6.1** (The model groupoid, and its projectivization).  $\text{Mod}_{\psi,\beta}(\kappa)$  is the category whose objects are pairs  $(M, \pi)$  with  $\pi : A_{\psi,\beta}(V) \rightarrow \text{End}_{\mathbb{C}}(M)$  a unital algebra homomorphism making  $M$  a simple module and every  $\pi(W_{\beta}(v))$  unitary, and whose morphisms  $(M, \pi) \rightarrow (M', \pi')$  are unitary intertwiners  $T : M \rightarrow M'$  ( $T\pi(a) = \pi'(a)T$  for all  $a$ ).  $\text{PMod}_{\psi,\beta}(\kappa)$  is the same category with each Hom-set replaced by  $\text{Hom}/U(1)$  (intertwiners identified when they differ by a unimodular scalar).

*Scope.* This definition alone asserts nothing about the size or connectivity of  $\text{Mod}_{\psi,\beta}(\kappa)$ ; that it is nonempty, connected, and has automorphism group  $U(1)$  at every object is Theorem 6.3(c) below, using Theorem 3.5, not part of the stipulation here.

---

**Provenance.** ids: D9; proved in: —; tested by: —; reviewed in: —.

## 6.1 Exactly two data beyond the field

### PROVED

**Theorem 6.2** (The construction depends on exactly two data beyond  $\kappa$ ). (a)  $X(\kappa) = \hat{\kappa} \setminus \{1\}$  has  $q - 1$  elements and is a free transitive  $\kappa^{\times}$ -torsor under  $(u \cdot \psi)(x) := \psi(ux)$  (a single point when  $q = 2$ , where  $\kappa^{\times}$  is trivial);

(b) beyond  $\kappa$ , the construction of  $A_{\psi,\beta}(V)$  uses **exactly two** further data: a nontrivial character  $\psi \in X(\kappa)$  and a polarizing cocycle  $\beta \in \text{Adm}(\omega)$  — given  $(\psi, \beta)$ ,  $A_{\psi,\beta}(V)$  (Definition 2.4) involves no further choice, while exhibiting a concrete Hilbert space (rather than the abstract algebra) needs a pair  $(L, \chi)$  in addition (Definition 3.2);

(c) distinct  $\psi, \psi' \in X(\kappa)$  give representations of the same group  $H_{\beta}(\kappa)$  (which, by Definition 2.5, does not involve  $\psi$  at all) with distinct central characters, hence inequivalent as representations of  $H_{\beta}(\kappa)$ ;

(d)  $A_{\psi,\beta}(V) \cong A_{\psi',\beta'}(V)$  for every  $\psi, \psi' \in X(\kappa)$  and  $\beta, \beta' \in \text{Adm}(\omega)$ ; but the frame-preserving isomorphisms between two such algebras form a torsor under  $\text{Aut}_{\mathcal{F}}^{\kappa}(A_{\psi,\beta}(V))$  (Theorem 5.2), a group of order  $q^2 |SL_2(\kappa)| \geq 24$  — so no single isomorphism among them is distinguished by the data  $(\kappa, \psi, \psi', \beta, \beta')$  alone.

*Scope.* Part (b) corrects an earlier working hypothesis of this campaign, that the construction depended on a single datum beyond  $\kappa$ : that hypothesis is refuted by Theorem 4.2 and Theorem 4.5 of §4, which show  $\beta$  is a second, independent choice. Part (d) asserts the algebras are abstractly isomorphic; it does not assert any *particular* isomorphism among them is preferred, and Theorem 6.3 below is precisely about what survives this non-uniqueness.

---

**Provenance.** ids: D2, D3, D4, WH-ALG, WH-BETA-a, WH-WEIL-a, WH-CHOICE; proved in: theory/wh-kappa-choice.md §7; tested by: theory/checks/wh\_kappa\_check.py (C8 in its E5 form, C10); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* (a)  $|\hat{\kappa}| = q$  by Fact 3.1 ( $U = \kappa$ ,  $d = m$ ), so  $|X(\kappa)| = q - 1$ . The action  $(u \cdot \psi)(x) := \psi(ux)$  preserves nontriviality: if  $\psi \not\equiv 1$  and  $u \neq 0$  then, as  $x \mapsto ux$  is a bijection of  $\kappa$ ,  $u \cdot \psi \equiv 1$  would force  $\psi \equiv 1$ , false. It is free:  $u \cdot \psi = \psi$  means  $\psi((u - 1)x) = 1$  for all  $x$ , i.e.  $(u - 1)\kappa \subseteq \ker \psi$ ; if  $u \neq 1$  then  $(u - 1) \in \kappa^{\times}$  so  $(u - 1)\kappa = \kappa \not\subseteq \ker \psi$  (Fact 3.1,  $\psi$  nontrivial), forcing  $u = 1$ . A free action of the order- $(q - 1)$  group  $\kappa^{\times}$  on the order- $(q - 1)$  set  $X(\kappa)$  has exactly one orbit of size  $q - 1$ , hence is transitive. At  $q = 2$ ,  $\kappa^{\times} = \{1\}$  and  $X(\kappa)$  is a single point — the  $q = 2$  choice, per Theorem 4.4, lives entirely in  $\beta$ , not in  $\psi$ .

(b) Immediate from Definitions 2.4 and 3.2: the structure constants of  $A_{\psi,\beta}(V)$  are  $\psi(\beta(v, v'))$ , a function of  $\kappa, \psi, \beta$  alone, while  $M_{L,\chi}$  additionally names a line  $L$  and a character  $\chi$  of  $A_L$ .

(c) A representation of  $H_\beta(\kappa)$  with central character  $\psi$  restricts on the centre  $\kappa \times 0$  to  $\psi \cdot \text{id}$  (the remark after Theorem 2.7); an intertwiner between representations with central characters  $\psi \neq \psi'$  would have to commute with the centre acting as  $\psi \cdot \text{id}$  on one side and  $\psi' \cdot \text{id}$  on the other, forcing it to be 0 on the difference  $(\psi(t) - \psi'(t)) \cdot \text{id} \neq 0$  for suitable  $t$  — no nonzero intertwiner exists, so the two representations of the fixed group  $H_\beta(\kappa)$  are inequivalent.

(d) *Existence, for every pair.* By Theorem 4.10,  $A_{\psi,\beta}(V) \cong A_{\psi,\beta_0}(V)$  for every  $\beta$  (via a frame isomorphism, taking  $\beta' = \beta_0$  there), and symmetrically  $A_{\psi',\beta'}(V) \cong A_{\psi',\beta_0}(V)$ . For the reference cocycle,  $W_\psi(a, b) \mapsto W_{\psi_u}(u^{-1}a, b)$  (for  $\psi' = u \cdot \psi = \psi_u$  as in (a)) is an isomorphism  $A_{\psi,\beta_0}(V) \rightarrow A_{\psi_u,\beta_0}(V)$ , since  $\beta_0(u^{-1}a, b') = u^{-1}\beta_0(v, v')$  makes the structure constants match:  $\psi_u(\beta_0(u^{-1}v, u^{-1}v')) = \psi(u \cdot u^{-2}\beta_0(v, v')) = \psi(u^{-1}\beta_0(v, v'))$ , consistently with the images  $W_{\psi_u}(u^{-1}v)$ . Composing the three maps ( $A_{\psi,\beta} \rightarrow A_{\psi,\beta_0} \rightarrow A_{\psi',\beta_0} \rightarrow A_{\psi',\beta'}$ , the middle map since  $X(\kappa)$  is a  $\kappa^\times$ -torsor by (a)) gives an isomorphism  $A_{\psi,\beta}(V) \rightarrow A_{\psi',\beta'}(V)$  for arbitrary  $\psi, \psi', \beta, \beta'$ . *Non-uniqueness.* The frame-preserving isomorphisms between two fixed such algebras, once nonempty, form a free transitive right  $\text{Aut}_{\mathcal{F}}^\kappa(A_{\psi,\beta}(V))$ -set under postcomposition (compose one such isomorphism with an automorphism of the source), and  $|\text{Aut}_{\mathcal{F}}^\kappa(A_{\psi,\beta}(V))| = q^2|SL_2(\kappa)|$  by Theorem 5.2, at least  $4 \cdot 6 = 24$ . More than one isomorphism exists whenever this group is non-trivial (always), and the data  $(\kappa, \psi, \psi', \beta, \beta')$  do not name a preferred one: e.g. a family  $\{h_u\}$ ,  $h_u := \text{diag}(u^{-j}, u^{j-1})$  for a further arbitrary choice of  $j \in \mathbb{Z}/(q-1)$ , gives infinitely many mutually inconsistent *coherent* systems of isomorphisms across the  $\kappa^\times$ -action of (a) — the family of isomorphisms is trivializable, but not canonically: choosing one is a section, not a theorem.  $\square$

## 6.2 The projective Hilbert space is canonical

### PROVED

**Theorem 6.3** (Canonicity of the projective Hilbert space, at fixed  $(\kappa, \psi, \beta)$ ). (a)  $\text{Mod}_{\psi,\beta}(\kappa)$  (Definition 6.1) is a nonempty, connected groupoid, with automorphism group  $U(1)$  at every object;

(b) consequently every  $\text{Hom}$ -set of  $\text{PMod}_{\psi,\beta}(\kappa)$  is a singleton, so every diagram in it commutes trivially, and the limit  $\varprojlim_{\text{PMod}_{\psi,\beta}(\kappa)} P(M)$  exists with every projection an isomorphism; define  $P(H_\psi(\kappa))$  to be this limit. The underlying Hilbert space  $H_\psi(\kappa)$  itself is canonical only up to a  $U(1)$  of identifications (one for each object of  $\text{Mod}_{\psi,\beta}(\kappa)$  chosen as a representative), but the projective Hilbert space  $P(H_\psi(\kappa))$  is canonical in  $(\kappa, \psi)$ , for fixed  $\beta$ .

*Scope.* This theorem fixes  $\beta$  throughout; that the resulting  $P(H_\psi(\kappa))$  does not, in fact, depend on the choice of  $\beta$  either is the separate, weaker-status content of Theorem 6.4 below, which is not needed for and does not feed back into this proof. Nothing here addresses canonicity of a specific *vector* in  $H_\psi(\kappa)$ , only of the isomorphism class of the pair  $(M, \pi)$  up to the  $U(1)$  ambiguity of unitary intertwiners.

---

**Provenance.** ids: D9, WH-SVN, WH-CHOICE, WH-CANON; proved in: theory/wh-kappa-choice.md §7 step  $\langle 1 \rangle 3, \langle 1 \rangle 4$ ; tested by: —; reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* (a) Nonempty: Theorem 3.5(a) exhibits the standard model  $M_0$ , unitary by Theorem 3.5(d). Connected: any two objects  $(M, \pi), (M', \pi')$  are unitarily isomorphic by Theorem 3.5(b)–(d) (uniqueness of the simple module up to unitary equivalence), so a morphism exists between any two objects, in either direction.  $\text{Aut}(M, \pi) = U(1)$ : by Theorem 3.5(b)'s Schur clause, any self-intertwiner is a scalar; a unitary one is then a scalar of modulus 1, and every element of  $U(1)$  is realized (scalars commute with everything, hence intertwine trivially).

(b) In a connected groupoid where every object has automorphism group  $U(1)$ , every Hom-set is a  $U(1)$ -torsor (fix one isomorphism between two objects; every other differs from it by an automorphism of the source, i.e. by an element of  $U(1)$ ). Quotienting each Hom-set by this  $U(1)$  therefore collapses it to a single point: every Hom-set of  $\text{PMod}_{\psi,\beta}(\kappa)$  is a singleton. Consequently, for any two morphisms  $f, g$  in  $\text{PMod}_{\psi,\beta}(\kappa)$  with the same source and target,  $f = g$  automatically (both are *the* unique element of that singleton Hom-set): every square of transition maps commutes, not by verification, but because there is only one morphism available to fill it. A category in which every Hom-set is a singleton is equivalent to the terminal (one-object, one-morphism) category, so the limit of the identity diagram  $P(M) \rightsquigarrow P(M)$  over  $\text{PMod}_{\psi,\beta}(\kappa)$  exists and every object, with its unique family of identifying isomorphisms to every other, computes it: this is  $P(H_\psi(\kappa))$ . Since the ordinary (non-projectivized) Hom-sets of  $\text{Mod}_{\psi,\beta}(\kappa)$  are  $U(1)$ -torsors and not singletons, no single Hilbert space among the objects is preferred over  $U(1)$ -many identifications with it — but every one of them projects isomorphically onto the same limit  $P(H_\psi(\kappa))$ .  $\square$

### 6.3 Independence from the polarizing cocycle as well

#### SKETCH

**Theorem 6.4** (The projective Hilbert space does not depend on  $\beta$  either).  *$P(H_\psi(\kappa))$ , as constructed in Theorem 6.3, is independent of the choice of polarizing cocycle  $\beta \in \text{Adm}(\omega)$  used to build it: the frame isomorphism  $A_{\psi,\beta}(V) \rightarrow A_{\psi,\beta'}(V)$  of Theorem 4.10 transports unitary simple modules and induces an equivalence  $\text{Mod}_{\psi,\beta}(\kappa) \rightarrow \text{Mod}_{\psi,\beta'}(\kappa)$ , which in turn induces the unique isomorphism between the two limits  $P(H_\psi(\kappa))$  constructed at  $\beta$  and at  $\beta'$ .*

*Scope.* This theorem is SKETCH for the same procedural reason as the  $p = 2$  rows of §4 it depends on (Theorem 4.10): it postdates this increment's single hostile-critic round. It asserts that the *limit* is independent of  $\beta$ , not that any specific object of  $\text{Mod}_{\psi,\beta}(\kappa)$  equals, on the nose, an object of  $\text{Mod}_{\psi,\beta'}(\kappa)$  — the identification goes through the transport functor described in the proof, consistently with the fact that the underlying group  $H_\beta(\kappa)$  genuinely changes isomorphism type with  $\beta$  at  $p = 2$  (Theorem 4.7).

---

**Provenance.** ids: WH-CANON, WH-BETA-LEVEL, WH-CANON-BETA; proved in: theory/wh-kappa-choice.md §7 step ⟨1⟩4.⟨2⟩1; tested by: —; reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* By Theorem 4.10 there is a frame isomorphism  $\Phi : A_{\psi,\beta}(V) \rightarrow A_{\psi,\beta'}(V)$ ,  $W_\beta(v) \mapsto \mu(v)W_{\beta'}(v)$ , with  $\mu$  unimodular (by the same argument as Remark 5.2: taking absolute values in the defining relation (†) of Theorem 4.10's proof makes  $|\mu|$  a homomorphism from a finite group to  $\mathbb{R}_{>0}$ , hence trivial). Transport of structure along  $\Phi$  sends an object  $(M, \pi)$  of  $\text{Mod}_{\psi,\beta}(\kappa)$  to  $(M, \pi')$  with  $\pi'(a') := \pi(\Phi^{-1}(a'))$  for  $a' \in A_{\psi,\beta'}(V)$ :  $M$  stays simple (an equivalence of module categories along an algebra isomorphism preserves simplicity), and each  $\pi'(W_{\beta'}(v)) = \pi(\Phi^{-1}(W_{\beta'}(v))) = \overline{\mu(v)}\pi(W_\beta(v))$  remains unitary (a unimodular scalar times a unitary operator). This is an equivalence of categories  $\text{Mod}_{\psi,\beta}(\kappa) \rightarrow \text{Mod}_{\psi,\beta'}(\kappa)$ , compatible with the projectivization functors to  $\text{PMod}_{\psi,\beta}(\kappa)$  and  $\text{PMod}_{\psi,\beta'}(\kappa)$  of Theorem 6.3(b). Both projectivized categories have singleton Hom-sets (Theorem 6.3(b), applied at  $\beta$  and at  $\beta'$  respectively), so the induced map between their limits  $P(H_\psi(\kappa))_\beta \rightarrow P(H_\psi(\kappa))_{\beta'}$  is again the unique morphism available between two terminal-equivalent categories, hence an isomorphism, and the unique one compatible with the transport functor.  $\square$

## 7 What $\kappa$ -linear symmetry hides: the $\mathbb{F}_p$ -shadow of the form

**Recollection.** For  $\beta \in \text{Adm}(\omega)$  and phase datum  $\psi$ , the Weyl frame is  $\mathcal{F} = \{\mathbb{C}^\times W_\beta(v) : v \in V(\kappa)\} \subset A_{\psi,\beta}(V)$ , and  $\text{Aut}_{\mathcal{F}}(A)$  is the group of  $\mathbb{C}$ -algebra automorphisms of  $A_{\psi,\beta}(V)$  preserving

$\mathcal{F}$  (Definition 5.1);  $\text{Aut}_{\mathcal{F}}^{\kappa}(A) \subseteq \text{Aut}_{\mathcal{F}}(A)$  is the further subgroup whose induced permutation of  $V(\kappa)$  is  $\kappa$ -linear — data *imposed* on top of  $\text{Aut}_{\mathcal{F}}(A)$ , not derived from it (Definition 5.1's scope). By Theorem 5.2,  $\text{Aut}_{\mathcal{F}}^{\kappa}(A)$  surjects onto  $SL_2(\kappa)$  with kernel  $\hat{V}$ , at every characteristic.

---

**Provenance.** ids: D4, D7, WH-WEIL-a; proved in: —; tested by: —; reviewed in: —.

This section makes good on the promissory note left in Definition 5.1: exactly how much symmetry the imposed  $\kappa$ -linearity clause discards. The answer is that  $A_{\psi,\beta}(V)$  and its frame  $\mathcal{F}$  do not, in fact, remember the  $\kappa$ -module structure of  $V(\kappa)$  at all — they are built from the bare abelian group  $(V(\kappa), +)$  and the cocycle  $\psi \circ \beta$  — so the *full* frame symmetry group is larger than  $SL_2(\kappa)$  whenever  $\kappa$  is a proper extension of its prime field, and the discarded symmetry is measured exactly.

**PROVED**

**Theorem 7.1** (The twisted algebra does not remember the  $\kappa$ -structure of  $V(\kappa)$ ). (a)  $A_{\psi,\beta}(V)$  and its frame  $\mathcal{F}$  are determined by the abelian group  $(V(\kappa), +)$  together with the  $\mathbb{C}^{\times}$ -valued function  $\psi \circ \beta$  on  $V(\kappa) \times V(\kappa)$  alone; nothing in either object refers to scalar multiplication by  $\kappa$ ;

(b) consequently, dropping the imposed  $\kappa$ -linearity clause of Definition 5.1, the frame automorphisms with merely  $\mathbb{F}_p$ -linear induced permutation  $g$  of  $V(\kappa)$  surject, with the same kernel  $\hat{V}$ , onto the isometry group  $Sp(V(\kappa), \psi \circ \omega) \cong Sp_{2m}(\mathbb{F}_p)$  of the  $\mathbb{F}_p$ -valued alternating form  $\psi \circ \omega$ ; so  $|\text{Aut}_{\mathcal{F}}(A)| = q^2 \cdot |Sp_{2m}(\mathbb{F}_p)|$ ;

(c)  $SL_2(\kappa) \subseteq Sp_{2m}(\mathbb{F}_p)$ , with index exactly 1 when  $m = 1$  and strictly greater than 1 as soon as  $m > 1$ : by complete enumeration,  $|SL_2(\kappa)| = 6, 24, 60, 120, 504, 720$  against  $|Sp_{2m}(\mathbb{F}_p)| = 6, 24, 720, 120, 1451520, 51840$  at  $q = 2, 3, 4, 5, 8, 9$  respectively — indices 1, 1, 12, 1, 2880, 72. So Definition 5.1's  $\kappa$ -linearity clause discards genuine symmetry the algebra has, whenever  $\kappa$  is a proper extension of its prime field.

*Scope.* This theorem does not claim  $Sp_{2m}(\mathbb{F}_p)$  acts by automorphisms preserving anything beyond the frame  $\mathcal{F}$  and the abelian group structure — in particular it says nothing about whether this larger symmetry is compatible with the  $\kappa$ -linear structure used elsewhere in this labbook (Theorem 2.6(f) shows, to the contrary, that the  $\kappa$ -valued form  $\omega$  itself sees only  $SL_2(\kappa)$  among  $\mathbb{F}_p$ -linear maps: the enlargement here is possible only because  $\text{Aut}_{\mathcal{F}}(A)$  is built from the strictly weaker  $\mathbb{F}_p$ -valued shadow  $\psi \circ \omega$ ). At the prime field,  $m = 1$ , the two groups coincide and nothing is discarded; this is the campaign's guiding path.

---

**Provenance.** ids: D1, D7, WH-WEIL-a, WH-FORM, WH-SYMM; proved in: theory/wh-kappa-choice.md §9; tested by: theory/checks/fp\_symmetry.py (S1,S2,S3); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* (a) By Definition 2.4, the structure constants of  $A_{\psi,\beta}(V)$  in the basis  $\{W_{\beta}(v)\}$  are exactly the values  $\psi(\beta(v, v'))$ , indexed by pairs of elements of the abelian group  $(V(\kappa), +)$ ; by Definition 5.1,  $\mathcal{F}$  is the set of  $\mathbb{C}^{\times}$ -lines through this same basis, again indexed only by  $(V(\kappa), +)$ . Neither construction refers to multiplication by elements of  $\kappa$ .

(b) Repeat the argument of Theorem 5.2's proof *without* imposing  $\kappa$ -linearity on  $g$ : frame preservation still gives  $\alpha(W_{\beta}(v)) = \lambda(v)W_{\beta}(gv)$  with  $g$  an (a priori only additive) bijection of  $V(\kappa)$  and  $\lambda$  satisfying relation (\*) there, since that derivation used only additivity of  $g$ , never  $\kappa$ -linearity. Applying  $\alpha$  to Theorem 2.7(c), an identity entirely in  $\psi \circ \omega$ , gives  $\psi(\omega(gv, gv')) = \psi(\omega(v, v'))$  for all  $v, v'$  — i.e.  $g \in Sp(V(\kappa), \psi \circ \omega) := \{g \text{ additive} : \psi \circ \omega \circ (g \times g) = \psi \circ \omega\}$  — rather than the stronger  $\omega \circ g = \omega$  used before. The surjectivity argument via Fact 3.1 applies verbatim to any such  $g$  (it never used  $\kappa$ -linearity, only that  $(V(\kappa), +)$  has exponent  $p$ ), so every  $g \in Sp(V(\kappa), \psi \circ \omega)$  lifts; the kernel computation ( $g = \text{id}$  forces  $\lambda \in \hat{V}$ , identified with

$V(\kappa)$  via  $u \mapsto \psi(\omega(u, \cdot))$  is unchanged, giving the same kernel  $\hat{V}$  of order  $q^2$ . Nondegeneracy of  $\psi \circ \omega$  as an  $\mathbb{F}_p$ -bilinear form on the  $2m$ -dimensional  $\mathbb{F}_p$ -space  $V(\kappa)$ : if  $\psi(\omega(v, \cdot)) \equiv 1$  then  $\omega(v, \cdot) \equiv 0$  (Fact 3.1, since  $\omega(v, \cdot)$  is either 0 or onto  $\kappa$ , being  $\kappa$ -linear, and  $\kappa \not\subseteq \ker \psi$ ), so  $v = 0$  by Theorem 2.6(c). A nondegenerate alternating form on a  $2m$ -dimensional space over  $\mathbb{F}_p$  has isometry group (by definition, matching the notation `notation.md` fixes for  $Sp_{2m}(\mathbb{F}_p)$ ) exactly  $Sp_{2m}(\mathbb{F}_p)$ .

(c) If  $g \in SL_2(\kappa)$  then  $\omega \circ (g \times g) = \omega$  (Theorem 2.6(e)), so certainly  $\psi \circ \omega \circ (g \times g) = \psi \circ \omega$ , i.e.  $SL_2(\kappa) \subseteq Sp(V(\kappa), \psi \circ \omega)$ . This containment is exactly the gap between Theorem 2.6(f) — which shows the  $\mathbb{F}_p$ -linear isometries of the  $\kappa$ -valued  $\omega$  are no larger than  $SL_2(\kappa)$  — and the present, strictly weaker condition of isometry for the  $\mathbb{F}_p$ -valued shadow  $\psi \circ \omega$  alone: an  $\mathbb{F}_p$ -linear  $g$  can preserve  $\psi \circ \omega$  without preserving  $\omega$  itself. Complete enumeration (`theory/checks/fp_symmetry.py`, gates S1 and S2) computes both orders directly at  $q = 2, 3, 4, 5, 8, 9$ , matching the closed form  $|Sp_{2m}(\mathbb{F}_p)| = p^{m^2} \prod_{i=1}^m (p^{2i} - 1)$  in every case, and confirms the indices 1, 1, 12, 1, 2880, 72; at  $m = 1$  (the prime field) the two group orders coincide, since  $Sp_2(\mathbb{F}_p) = SL_2(\mathbb{F}_p)$  by definition in dimension 2, and nothing is discarded.  $\square$

*Remark* (The characteristic-two orthogonal phenomenon survives the enlargement). The  $\mu_2$ -phase criterion of Theorem 5.4 ( $g \in O(Q_\beta)$ ) makes sense verbatim for merely  $\mathbb{F}_p$ -linear  $g$ , at  $p = 2$ :  $\psi \circ Q_\beta \circ g = \psi \circ Q_\beta$  is a condition on an  $\mathbb{F}_p$ -valued quadratic form  $\psi \circ Q_\beta$  with polar form  $\psi \circ \omega$ , so the relevant subgroup inside the enlarged symmetry group of Theorem 7.1 is  $O(\psi \circ Q_\beta) \cap Sp_{2m}(\mathbb{F}_p)$ . Direct enumeration (gate S3) finds its order to be 2, 72, 40320 at  $q = 2, 4, 8$ , of index 3, 10, 36 in  $Sp_{2m}(\mathbb{F}_p)$  — exactly the *same* index that  $|O(Q_\beta) \cap SL_2(\kappa)| = 2, 6, 14$  has inside  $SL_2(\kappa)$  (Theorem 5.4). So the characteristic-two orthogonal phenomenon of §5 is not an artefact of the imposed  $\kappa$ -linearity in Definition 5.1: it survives, at the same relative index, inside the larger, intrinsic symmetry group  $Sp_{2m}(\mathbb{F}_p)$  that Theorem 7.1 exhibits. This observation is recorded as part of the same source theorem as Theorem 7.1, under the same provenance (`theory/wh-kappa-choice.md` §9, step (1)4).

*Remark* (Reading). Two independent non-canoncities are now on record, and they are logically independent of each other. Theorem 4.7 of §4 shows the system does not remember which polarizing cocycle  $\beta$  was chosen, in a way that is invisible at odd characteristic and genuinely two-valued at  $p = 2$ . This theorem shows the system does not remember the  $\kappa$ -module structure of  $V(\kappa)$  itself, at every characteristic and every  $m > 1$ , regardless of  $\beta$ . Both are statements about what the definition actually depends on, in the sense of `PRD.md`'s “canonical” requirement: any assignment  $\text{Spec } R \mapsto \text{quantum system}$  general enough to restrict to this one on  $\text{Spec } \mathbb{F}_q$  carries both phenomena, not as defects, but as exactly the symmetry and the choice the construction is built from.

## 8 Functoriality in the field: what extends along embeddings, and what does not

**Recollection.** A phase datum for  $\kappa = \mathbb{F}_{p^m}$  is a nontrivial additive character  $\psi : (\kappa, +) \rightarrow \mathbb{C}^\times$ , and the trace-normalized family is  $\psi_\zeta(z) := \zeta^{\text{Tr}_{\kappa/\mathbb{F}_p}(z)}$  for  $\zeta$  a primitive  $p$ -th root of unity and  $\text{Tr}_{\kappa/\mathbb{F}_p}(z) := z + z^p + \cdots + z^{p^{m-1}}$  (Definition 2.3); by Theorem 6.2(a),  $X(\kappa)$  is a  $\kappa^\times$ -torsor, so every  $\psi \in X(\kappa)$  is  $\psi_c := \psi_\zeta(c \cdot)$  for a unique  $c \in \kappa^\times$ .

---

**Provenance.** ids: D3; proved in: —; tested by: —; reviewed in: —.

Earlier sections fixed a single finite field  $\kappa$  throughout. This section asks how the construction behaves as  $\kappa$  varies along field embeddings: whether the trace-normalized family is itself compatible with restriction, whether the assignment  $\kappa \mapsto A_{\psi, \beta_0}(V(\kappa))$  is a functor, and — the

question this campaign's earlier working hypothesis answered wrongly — whether a single compatible choice of character can be made once and for all, for every finite field simultaneously. It cannot; the precise obstruction is Frobenius, and the precise set of compatible choices that *does* exist is named exactly.

**PROVED**

**Theorem 8.1** (The trace-normalized family, restricted along an extension). *Let  $\kappa = \mathbb{F}_{p^m} \subseteq \kappa' = \mathbb{F}_{p^{mn}}$ . Then  $\psi'_\zeta|_\kappa = \psi_{\zeta^n}$  (both sides characters of  $\kappa$ , computed with the same  $\zeta$ ): the restriction of  $\kappa'$ 's trace-normalized character to  $\kappa$  is trivial exactly when  $p \mid n$  (smallest instance:  $\mathbb{F}_2 \subset \mathbb{F}_4$ ).*

*Scope.* This theorem is about the trace-normalized family specifically — a particular, and particularly natural, choice inside  $X(\kappa)$  for every  $\kappa$  — and asserts nothing about arbitrary families of characters; the statement that quantifies over *all* compatible systems is Theorem 8.4 below.

---

**Provenance.** ids: D3, WH-FUNCT-a; proved in: theory/wh-kappa-choice.md §10 step ⟨1⟩1; tested by: theory/checks/funct.sections.py (F1); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* For  $x \in \kappa$ ,  $x^{p^m} = x$  (elements of  $\mathbb{F}_{p^m}$  are exactly the fixed points of the  $m$ -th power of Frobenius), so the  $mn$  summands of  $\text{Tr}_{\kappa'/\mathbb{F}_p}(x) = \sum_{k=0}^{mn-1} x^{p^k}$  repeat with period  $m$ : grouping into  $n$  blocks of  $m$  consecutive terms, each block sums to  $\text{Tr}_{\kappa/\mathbb{F}_p}(x)$  (using  $x^{p^{k+m}} = (x^{p^m})^{p^k} = x^{p^k}$ ), so  $\text{Tr}_{\kappa'/\mathbb{F}_p}|_\kappa = n \cdot \text{Tr}_{\kappa/\mathbb{F}_p}$ . Hence  $\psi'_\zeta(x) = \zeta^{\text{Tr}_{\kappa'/\mathbb{F}_p}(x)} = \zeta^{n \cdot \text{Tr}_{\kappa/\mathbb{F}_p}(x)} = (\zeta^n)^{\text{Tr}_{\kappa/\mathbb{F}_p}(x)} = \psi_{\zeta^n}(x)$ , and  $\zeta^n = 1$  (so  $\psi_{\zeta^n} \equiv 1$ ) exactly when  $p \mid n$ , since  $\zeta$  has exact order  $p$ .  $\square$

**PROVED**

**Theorem 8.2** (The reference algebra is functorial in character-compatible embeddings). *Let  $\mathcal{FF}^\pm$  be the category with objects the pairs  $(\kappa, \psi)$ ,  $\psi \in X(\kappa)$ , and morphisms  $(\kappa, \psi) \rightarrow (\kappa', \psi')$  the field embeddings  $\iota : \kappa \hookrightarrow \kappa'$  with  $\psi' \circ \iota = \psi$  (character-compatible embeddings). Then*

$$(\kappa, \psi) \longmapsto A_{\psi, \beta_0}(V(\kappa))$$

*is a functor from  $\mathcal{FF}^\pm$  to the category of unital  $\mathbb{C}$ -algebras and injective algebra homomorphisms, respecting composition and identities.*

*Scope.* This functor uses the reference cocycle  $\beta_0$  throughout, on every object; it is not claimed that the resulting assignment is independent of this choice, nor that an analogous functor exists carrying a varying  $\beta_\kappa$  compatibly along embeddings — the reference cocycle is used here because it is a fixed, explicit stipulation (Definition 4.1), not because it is canonical (Theorem 4.4 of §4 shows at  $p = 2$  it is not). Nor is this a statement about the quantum *system* — the Hilbert space or the Heisenberg group — functorially in  $\kappa$ , only about the algebra.

---

**Provenance.** ids: D2, D4, WH-ALG, WH-FUNCT-b; proved in: theory/wh-kappa-choice.md §10 step ⟨1⟩2; tested by: theory/checks/funct.sections.py (F5); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* For a character-compatible embedding  $\iota : (\kappa, \psi) \rightarrow (\kappa', \psi')$ , put  $V(\iota)(a, b) := (\iota a, \iota b) : V(\kappa) \rightarrow V(\kappa')$ , additive since  $\iota$  is a field homomorphism. Since  $\iota$  is multiplicative,  $\beta_0(V(\iota)v, V(\iota)v') = \iota(a) \iota(b') = \iota(ab') = \iota(\beta_0(v, v'))$ . Define  $W_\beta(v) \mapsto W'_{\beta_0}(V(\iota)v)$  on basis elements of  $A_{\psi, \beta_0}(V(\kappa))$ . This sends distinct basis elements to distinct basis elements ( $V(\iota)$  is injective,  $\iota$  being a field embedding), and is multiplicative: the image of  $W_{\beta_0}(v)W_{\beta_0}(v') = \psi(\beta_0(v, v'))W_{\beta_0}(v + v')$  is  $\psi(\beta_0(v, v'))W'_{\beta_0}(V(\iota)(v + v'))$ , while the product of the images is  $\psi'(\beta_0(V(\iota)v, V(\iota)v'))W'_{\beta_0}(V(\iota)(v + v')) = \psi'(\iota(\beta_0(v, v')))W'_{\beta_0}(V(\iota)(v + v')) = \psi(\beta_0(v, v'))W'_{\beta_0}(V(\iota)(v + v'))$ , using  $\psi' \circ \iota = \psi$  (the compatibility condition). So the map is a unital, nonzero algebra homomorphism, hence injective by simplicity of the source (Theorem 2.8). It respects composition and identities because  $\iota \mapsto V(\iota)$  does:  $V(\iota' \circ \iota) = V(\iota') \circ V(\iota)$  and  $V(\text{id}) = \text{id}$ , directly from the definition of  $V(\iota)$ .  $\square$

## 8.1 No compatible choice of character over every finite field

### REFUTED

**Theorem 8.3** (Refuted: a Frobenius-compatible character exists for every finite field). (Claimed, and refuted.) *A section of the forgetful functor  $\mathcal{FF}^\pm \rightarrow \{\text{finite fields}\}$  exists: a choice  $\psi_\kappa \in X(\kappa)$  for every finite field  $\kappa$  such that  $\iota : \kappa \rightarrow \kappa'$  character-compatible (i.e.  $\psi_{\kappa'} \circ \iota = \psi_\kappa$ ) for every field embedding  $\iota$  — in particular, for the Frobenius self-embedding of every  $\kappa$ .*

*Scope.* This row is kept, refuted, rather than deleted, per this campaign’s status discipline: a refutation carrying the surviving weaker statement is a result, not a gap to be erased. The surviving statement — exactly which obstruction blocks a section, and exactly which weaker compatible systems *do* exist — is Theorem 8.4 immediately below and Theorem 8.5 thereafter. Nothing may depend on this row.

---

**Provenance.** ids: WH-FUNCT-b-SEC; proved in: —; tested by: theory/checks/funct\_sections.py (F2,F3); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Refutation.* Frobenius  $x \mapsto x^p$  is a field embedding  $\kappa \rightarrow \kappa$ , so a section would in particular have to satisfy  $\psi_\kappa \circ \text{Frob} = \psi_\kappa$  for every  $\kappa$ . Theorem 8.4 below shows this forces  $\psi_\kappa = \psi_c$  for some  $c$  in the prime field  $\mathbb{F}_p$ , and that for  $\kappa = \mathbb{F}_{p^p}$  every character of this Frobenius-invariant form restricts *trivially* to  $\mathbb{F}_p$  — contradicting  $\psi_{\mathbb{F}_p} \in X(\mathbb{F}_p)$ , which must be nontrivial by definition. So no section exists; the smallest instance of the obstruction is  $\mathbb{F}_2 \subset \mathbb{F}_4$ . A previously considered route to a section, by induction along the tower  $\mathbb{F}_{p^{n!}}$ , is insufficient in any case: a character of  $\mathbb{F}_{p^{n!}}$  compatible at that one stage can still restrict trivially to an intermediate field, so no such induction can repair the obstruction just exhibited.

### PROVED

**Theorem 8.4** (The Frobenius obstruction, stated exactly). *There is no choice of  $\psi_\kappa \in X(\kappa)$ , one for each finite field  $\kappa$ , compatible with all field embeddings. Precisely: writing  $\psi_c := \psi_\zeta(c \cdot)$  for  $c \in \kappa$  (Theorem 6.2(a)), a character  $\psi_c$  is invariant under the Frobenius self-embedding of  $\kappa = \mathbb{F}_{p^m}$  if and only if  $c \in \mathbb{F}_p$ ; and for  $\kappa = \mathbb{F}_{p^p}$ , every  $\mathbb{F}_p$ -valued  $c$  gives a character  $\psi_c$  that restricts trivially to the prime field  $\mathbb{F}_p \subset \mathbb{F}_{p^p}$ . The smallest instance is  $\mathbb{F}_2 \subset \mathbb{F}_4$ .*

*Scope.* This theorem exhibits the precise obstruction at the specific extension degree  $m = p$ ; it does not enumerate every extension degree at which some weaker compatibility fails, and it does not address (this is Theorem 8.5) which *restricted* systems of compatible characters — over a sub-poset of embeddings, or with the “for every embedding” requirement relaxed — do exist.

---

**Provenance.** ids: D3, WH-FUNCT-a, WH-FUNCT-c; proved in: theory/wh-kappa-choice.md §10 step <1>3; tested by: theory/checks/funct\_sections.py (F2,F3); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof.* *Frobenius-invariance forces  $c \in \mathbb{F}_p$ .* Write  $d := c^{1/p}$  (the unique  $p$ -th root, Frobenius being bijective on the finite field  $\kappa$ ). Since  $\text{Tr}_{\kappa/\mathbb{F}_p}(y^p) = \text{Tr}_{\kappa/\mathbb{F}_p}(y)$  for every  $y$  (Frobenius cyclically permutes the  $m$  summands of the trace, using  $y^{p^m} = y$ ), we get  $\psi_c(x^p) = \zeta^{\text{Tr}(cx^p)} = \zeta^{\text{Tr}((dx)^p)} = \zeta^{\text{Tr}(dx)} = \psi_d(x)$ , i.e.  $\psi_c \circ \text{Frob} = \psi_d$ . Since  $c \mapsto \psi_c$  is injective on  $\kappa$  (if  $\psi_c = \psi_{c'}$  then  $\text{Tr}((c - c')x) \equiv 0$  for all  $x$ , forcing  $c = c'$  by the same nondegeneracy of the trace pairing used in Theorem 6.2(a)’s proof),  $\psi_c$  is Frobenius-invariant iff  $d = c$ ; since  $d^p = c$  by definition of  $d$ , substituting  $d = c$  gives  $c^p = c$ , and conversely  $c^p = c$  gives  $d = c^{1/p} = c$  (Frobenius being bijective). So the condition is exactly  $c^p = c$ , i.e.  $c$  is a root of  $X^p - X$ , i.e. (this polynomial having at most  $p$  roots in a field, all  $p$  elements of  $\mathbb{F}_p$  among them by Fermat’s little theorem for finite fields)  $c \in \mathbb{F}_p$ .

*Trivial restriction at  $\kappa = \mathbb{F}_{p^p}$ .* Here  $m = p$ . For  $c \in \mathbb{F}_p$  and  $x \in \mathbb{F}_p \subset \kappa$ , every summand of  $\text{Tr}_{\kappa/\mathbb{F}_p}(cx) = \sum_{k=0}^{p-1} (cx)^{p^k}$  equals  $cx$  (as  $cx \in \mathbb{F}_p$  is already Frobenius-fixed), so  $\text{Tr}_{\kappa/\mathbb{F}_p}(cx) = p \cdot (cx) = 0$  in characteristic  $p$ . So  $\psi_c(x) = \zeta^0 = 1$  for every  $x \in \mathbb{F}_p$ : every Frobenius-invariant character of  $\mathbb{F}_{p^p}$  — the only kind a section could assign there — restricts trivially to the prime field, contradicting the requirement  $\psi|_{\mathbb{F}_p} \in X(\mathbb{F}_p)$  (nontrivial by Definition 2.3) forced by compatibility along the inclusion  $\mathbb{F}_p \subset \mathbb{F}_{p^p}$ . At  $p = 2$ ,  $\mathbb{F}_{p^p} = \mathbb{F}_4$ : the smallest instance.  $\square$

## 8.2 The compatible systems that do exist

### PROVED

**Theorem 8.5** (Compatible character systems over an algebraic closure). *Fix an algebraic closure  $\overline{\mathbb{F}}_p$ , and consider the inclusion poset of its finite subfields. The compatible systems  $(\psi_\kappa)_\kappa$  — one  $\psi_\kappa \in X(\kappa)$  for every finite subfield  $\kappa \subset \overline{\mathbb{F}}_p$ , with  $\psi_{\kappa'}|_\kappa = \psi_\kappa$  for every inclusion  $\kappa \subseteq \kappa'$  of finite subfields — are **exactly** the restrictions  $\psi_\kappa := \Psi|_\kappa$  of the additive characters  $\Psi$  of  $(\overline{\mathbb{F}}_p, +)$  with  $\Psi|_{\mathbb{F}_p} \neq 1$ .*

*Scope.* This theorem restricts attention to the inclusion poset of finite subfields of a *fixed* algebraic closure, respecting compatibility only along inclusions of subfields, not along all abstract field embeddings between isomorphic-but-not-identical finite fields; Theorem 8.3 shows the latter, stronger requirement admits no solution at all. It says nothing about compatible choices of a polarizing cocycle  $\beta$  along the same poset; whether the Arf type of §4 can be chosen compatibly there is not resolved by this labbook.

---

**Provenance.** ids: D3, WH-FUNCT-d; proved in: theory/wh-kappa-choice.md §10 step ⟨1⟩4; tested by: theory/checks/funct.sections.py (F4); reviewed in: theory/verdicts/wh-kappa-adjudication.md.

*Proof. Restrictions are compatible systems.* Given such a  $\Psi$ , put  $\psi_\kappa := \Psi|_\kappa$ ; compatibility along  $\kappa \subseteq \kappa'$  holds by construction ( $(\Psi|_{\kappa'})|_\kappa = \Psi|_\kappa$ ), and  $\psi_\kappa$  is nontrivial for every finite subfield  $\kappa$ : every such  $\kappa$  contains  $\mathbb{F}_p$ , and  $\Psi|_\kappa \equiv 1$  would force  $\Psi|_{\mathbb{F}_p} \equiv 1$  too, contradicting the hypothesis on  $\Psi$ .

*Compatible systems arise this way.* Given a compatible system  $(\psi_\kappa)$ , define  $\Psi(x) := \psi_\kappa(x)$  for any finite subfield  $\kappa \ni x$  (every  $x \in \overline{\mathbb{F}}_p$  lies in some finite subfield,  $\overline{\mathbb{F}}_p$  being their union). This is well defined: if  $x$  lies in two finite subfields  $\kappa, \kappa'$ , both lie inside a common finite subfield  $\kappa''$  (e.g. their compositum, again finite), and compatibility along  $\kappa \subseteq \kappa''$  and  $\kappa' \subseteq \kappa''$  gives  $\psi_\kappa(x) = \psi_{\kappa''}(x) = \psi_{\kappa'}(x)$ .  $\Psi$  is additive: any two elements lie in a common finite subfield, where  $\Psi$  agrees with the additive character  $\psi_\kappa$  of that subfield. And  $\Psi|_{\mathbb{F}_p} = \psi_{\mathbb{F}_p} \neq 1$ , by Definition 2.3's requirement on every  $\psi_\kappa \in X(\kappa)$  applied to  $\kappa = \mathbb{F}_p$ .

*The two constructions are mutually inverse*, and the condition “nontrivial on every finite subfield” (needed termwise, since each  $\psi_\kappa \in X(\kappa)$  by hypothesis) is equivalent to “nontrivial on the prime field alone”: the forward implication is immediate ( $\mathbb{F}_p$  is itself one of the subfields), and the reverse is exactly what the first paragraph showed.  $\square$

*Remark* (What this section does not claim). Theorem 8.2 is functoriality of the *algebra* in character-compatible embeddings, for the fixed reference cocycle  $\beta_0$  — not functoriality of the quantum system (Heisenberg group, Hilbert space) itself, and not a claim that any compatible choice of polarizing cocycle along embeddings exists. Whether the Arf type of Theorem 4.7 can be chosen compatibly along field embeddings at  $p = 2$  is recorded as an open question by this campaign's source shard, not resolved here.

## 9 Weyl–Heisenberg quantization over finite local rings

This section records the first extension from finite fields to finite commutative local rings. In the preferred three-stage pipeline (P1), the ring first produces the symplectic module  $R^2$ ; a

character and a polarizing cocycle then produce its Weyl algebra and Heisenberg group; finally the irreducible unitary representations are classified. The decisive condition is that the chosen character generate the character module. At odd residue characteristic this gives a single irreducible representation of dimension  $|R|$ .

There is a genuine fork at the first stage. Route A, used here, takes  $R \oplus R$  with an  $R$ -valued alternating form and then composes with a chosen additive character. Route B would take  $R \oplus \widehat{R}$  with its canonical circle-valued Pontryagin pairing; it needs no character to become nondegenerate. Route B and the comparison of the physics retained by the two routes are deferred; no result below ranks the routes.

## 9.1 The local datum and its phase space

**Definition 9.1** (Finite commutative local ring datum). A finite local ring datum is  $(R, \mathfrak{m}, \kappa, q)$  with  $R$  a finite commutative unital local ring with  $1 \neq 0$ ,  $\mathfrak{m}$  its unique maximal ideal,  $\kappa := R/\mathfrak{m}$ , and  $q := |\kappa|$ ; write  $R^\times$  for its unit group. For an ideal  $I \subseteq R$ , put

$$\text{Ann}(I) := \{r \in R : rI = 0\}, \quad \text{soc}(R) := \text{Ann}(\mathfrak{m}).$$

These are stipulations. Nilpotence of  $\mathfrak{m}$ , nonvanishing of the socle, and the other finite-local structure properties are established below, not assumed here. The definition is uniform in the residue characteristic.

*Scope.* Only finite commutative unital local rings with  $1 \neq 0$  are defined here. No Frobenius, characteristic, principal-ideal, or chain-ring hypothesis is part of the datum.

---

**Provenance.** ids: D12; proved in: —; tested by: —; reviewed in: theory/verdicts/fcr-local-r1.md.

**Definition 9.2** (Characters and generating characters). Let

$$\widehat{R} := \text{Hom}((R, +), \mathbb{C}^\times), \quad X(R) := \widehat{R} \setminus \{1\}.$$

For  $\psi \in X(R)$  and  $u \in R$ , put  $\psi_u(x) := \psi(ux)$ . Define

$$I_\psi := \sum \{I \subseteq R : I \text{ is an ideal and } I \subseteq \ker \psi\}, \quad \text{Gen}(R) := \{\psi \in X(R) : I_\psi = 0\}.$$

Thus  $I_\psi$  is the largest ideal contained in  $\ker \psi$ , and a member of  $\text{Gen}(R)$  is a *generating character*. This definition is uniform in the residue characteristic; existence is not stipulated.

*Scope.* The definition includes every nontrivial additive character, not only a trace-normalized family. It makes no assertion that generating characters exist for an arbitrary finite local ring.

---

**Provenance.** ids: D13; proved in: —; tested by: —; reviewed in: theory/verdicts/fcr-local-r1.md.

**Definition 9.3** (Symplectic module and phase perpendicularity). Put  $V(R) := R \oplus R$  and

$$\omega((a, b), (a', b')) := ab' - a'b.$$

Given  $\psi \in X(R)$ , write

$$B_\psi(v, w) := \psi(\omega(v, w)), \quad L^{\perp\psi} := \{v \in V(R) : B_\psi(v, l) = 1 \text{ for every } l \in L\}$$

for an  $R$ -submodule  $L \subseteq V(R)$ . A *Lagrangian* is an  $R$ -submodule satisfying  $L = L^{\perp\psi}$ , and  $\text{rad}(B_\psi) := V(R)^{\perp\psi}$ . Bilinearity, strong alternation,  $R$ -nondegeneracy of  $\omega$ , and nondegeneracy of  $B_\psi$  are claims, not stipulations; the character-valued pairing  $B_\psi$ , not  $R$ -nondegeneracy alone, is load-bearing. This definition is uniform in the residue characteristic.

*Scope.* Perpendicularity is defined only for  $R$ -submodules of  $R^2$  and is measured through the chosen character. No freeness condition is imposed on a Lagrangian.

---

**Provenance.** ids: D14; proved in: —; tested by: —; reviewed in: theory/verdicts/fcr-local-r1.md.

**Definition 9.4** (Admissible polarizing cocycles over a local ring). For the symplectic module above, set

$$\begin{aligned}\text{Adm}(\omega) &:= \{\beta : V(R) \times V(R) \rightarrow R : \beta \text{ is } R\text{-bilinear and } \beta - \beta^T = \omega\}, \\ \text{Sym}_R(V(R)) &:= \{s : V(R) \times V(R) \rightarrow R : s \text{ is } R\text{-bilinear and } s = s^T\}.\end{aligned}$$

A *polarizing cocycle* is a named choice  $\beta \in \text{Adm}(\omega)$ . The reference member is

$$\beta_0((a, b), (a', b')) := ab'.$$

This is the nonsymmetrized convention: no inverse of 2 is assumed. The definition is uniform in the residue characteristic.

*Scope.* The reference member is a stipulated coordinate formula, not a claim that the torsor has a canonical point. The symmetrized member  $\omega/2$  is used only under the explicit hypothesis  $2 \in R^\times$ .

---

**Provenance.** ids: D15; proved in: —; tested by: —; reviewed in: theory/verdicts/fcr-local-r1.md.

**Definition 9.5** (Local Weyl algebra, Heisenberg group, and models). For the data above define

$$A_{\psi, \beta}(V(R)) := \bigoplus_{v \in V(R)} \mathbb{C} W_\beta(v), \quad W_\beta(v) W_\beta(v') := \psi(\beta(v, v')) W_\beta(v + v'),$$

and

$$H_\beta(R) := R \times V(R), \quad (t, v)(t', v') := (t + t' + \beta(v, v'), v + v').$$

For an  $R$ -submodule  $L$ , put  $A_L := \text{span}_{\mathbb{C}}\{W_\beta(l) : l \in L\}$ . Given a unital algebra character  $\chi : A_L \rightarrow \mathbb{C}$ , put

$$M_{L, \chi} := A_{\psi, \beta}(V(R)) \otimes_{A_L} \mathbb{C}_\chi.$$

The fixed reference model is  $\ell^2(R) = \bigoplus_{y \in R} \mathbb{C} e_y$ , with  $\{e_y\}$  orthonormal,

$$X(a)e_y := e_{y+a}, \quad Z(b)e_y := \psi(by),$$

and, exactly,

$$W_{\beta_0}(a, b) := Z(-b)X(a), \quad W_{\beta_0}(a, b)e_y = \psi(-b(y + a))e_{y+a}.$$

The sign and ordering are stipulations. Every displayed formula is uniform in the residue characteristic and uses no half.

*Scope.* The algebra and group depend on the named cocycle, while the algebra and representation also depend on the named character. A submodule and its character label a realization; neither is added to the underlying algebra datum.

---

**Provenance.** ids: D16; proved in: —; tested by: theory/checks/fcr\_local\_check.py (G4); reviewed in: theory/verdicts/fcr-local-r1.md.

## 9.2 Generating characters and the collapse boundary

### PROVED

**Theorem 9.6** (Local Frobenius criterion and the unit torsor). *Let  $R$  be a finite commutative local ring with maximal ideal  $\mathfrak{m}$  and residue field  $\kappa$ . Then*

$$\text{Gen}(R) \neq \emptyset \iff \dim_{\kappa} \text{soc}(R) = 1.$$

*If  $\psi \in \text{Gen}(R)$ , then  $\psi_u$  is generating exactly when  $u \in R^\times$ , and  $R^\times$  acts freely and transitively on  $\text{Gen}(R)$ .*

*Scope.* No odd-characteristic hypothesis is used. For rings whose socle has  $\kappa$ -dimension greater than one, the theorem asserts nonexistence of a generating character; it does not propose a replacement quantization.

---

**Provenance.** ids: D12, D13, FCR-GEN; proved in: theory/fcr-local.md §1; tested by: theory/checks/fcr\_local\_check.py (G1,G8); reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* Finiteness makes  $R$  Artinian, so  $\mathfrak{m}$  is nilpotent. Wood's module socle  $S({}_R R)$  equals  $\text{Ann}(\mathfrak{m})$ : every simple ideal  $J$  has  $\mathfrak{m}J = 0$ , since  $\mathfrak{m}J = J$  would contradict nilpotence; conversely, if  $0 \neq x \in \text{Ann}(\mathfrak{m})$ , then  $Rx = \kappa x$  is one-dimensional over  $\kappa$  and is simple. Wood's local Frobenius condition is therefore  $\kappa \cong \text{soc}(R)$ .

Wood's character-module theorem and kernel-ideal criterion identify this condition with the existence of  $\psi$  whose kernel contains no nonzero ideal, namely  $I_\psi = 0$ . For such a  $\psi$ , multiplication by a unit preserves  $I_\psi = 0$ , whereas a nonunit lies in  $\mathfrak{m}$  and kills the nonzero socle. The character-module isomorphism then shows that every generating character is uniquely  $\psi_u$  for a unit  $u$ .  $\square$

### PROVED

**Theorem 9.7** (Radical of the character-valued symplectic pairing). *Let  $R$  be a finite commutative local ring and let  $\psi \in X(R)$ . The form  $\omega$  on  $R^2$  is  $R$ -bilinear, strongly alternating, and  $R$ -nondegenerate, while*

$$\text{rad}(B_\psi) = I_\psi \oplus I_\psi.$$

*Consequently  $B_\psi$  is nondegenerate if and only if  $\psi \in \text{Gen}(R)$ .*

*For  $R = \mathbb{F}_3[x, y]/(x, y)^2$ , write  $\psi_{a,b,c}(r + sx + ty) = \zeta_3^{ar+bs+ct}$ . The two nontrivial characters with  $(b, c) = (0, 0)$  have  $I_\psi = (x, y)$ , and each of the remaining 24 has  $I_\psi = \mathbb{F}_3(cx - by)$ . Thus all 26 nontrivial characters collapse the pairing through a proper quotient.*

*Scope.* The radical formula is uniform in the residue characteristic. The square-zero-ring census records the obstruction at one designated seed only; it does not construct a quantum system for a degenerate pairing and does not state a general quotient-collapse theorem.

---

**Provenance.** ids: D12, D13, D14, FCR-RAD; proved in: theory/fcr-local.md §2; tested by: theory/checks/fcr\_local\_check.py (G2,G3); exhaustive 26-character seed census in G1; reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* Testing  $\omega((a, b), -)$  against  $(0, r)$  and  $(r, 0)$  shows that a radical vector must have  $aR, bR \subseteq \ker \psi$ , hence  $a, b \in I_\psi$ . The reverse inclusion follows because  $I_\psi$  is an ideal. Bilinearity and strong alternation are immediate from the displayed determinant formula, and tests against  $(0, 1)$  and  $(1, 0)$  prove  $R$ -nondegeneracy. In the square-zero example every subspace of  $(x, y)$  is an ideal; restricting the character to that two-dimensional space gives exactly the displayed kernel line, or all of  $(x, y)$  when the restriction vanishes.  $\square$

### 9.3 Odd residue characteristic: algebra and representation

#### PROVED

**Theorem 9.8** (Weyl commutation over a finite local ring). *Let  $R$  be a finite commutative local ring, let  $\psi \in X(R)$ , and let  $\beta \in \text{Adm}(\omega)$ . For every  $v, v' \in V(R)$ ,*

$$W_\beta(v)W_\beta(v') = \psi(\omega(v, v'))W_\beta(v')W_\beta(v),$$

and

$$W_\beta(u)W_\beta(v)W_\beta(u)^{-1} = \psi(\omega(u, v))W_\beta(v).$$

*Scope.* The identities are uniform in the residue characteristic and do not require  $\psi$  to be generating. They do not by themselves imply simplicity of the Weyl algebra.

---

**Provenance.** ids: D14, D15, D16, FCR-COMM; proved in: theory/fcr-local.md §3; tested by: theory/checks/fcr\_local\_check.py (G2,G4); reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* The two products have the same basis term  $W_\beta(v + v')$ . Their scalar ratio is  $\psi(\beta(v, v') - \beta(v', v)) = \psi(\omega(v, v'))$ ; conjugation gives the second identity.  $\square$

#### PROVED<sup>†</sup>

**Theorem 9.9** (Odd-characteristic transport of polarizing cocycles). *Let  $R$  be a finite commutative local ring with  $2 \in R^\times$ . Then  $\text{Adm}(\omega)$  is a torsor under  $\text{Sym}_R(V(R))$  and has  $|R|^3$  elements. Its unique antisymmetric member is  $\omega/2$ . For every  $\beta \in \text{Adm}(\omega)$  and every symmetric  $s$ , the map*

$$\varphi_s(t, v) := \left( t + \frac{s(v, v)}{2}, v \right)$$

*is an isomorphism  $H_\beta(R) \rightarrow H_{\beta+s}(R)$  fixing the centre pointwise.*

*Scope.* The condition  $2 \in R^\times$  is essential to the displayed representative and transport. No symmetry group of the ring-side phase space is analyzed here.

---

**Provenance.** ids: D14, D15, D16, FCR-BETA-ODD; proved in: theory/fcr-local.md §4; tested by: theory/checks/fcr\_local\_check.py (G9); reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* The difference of two admissible forms is symmetric, and adding a symmetric form preserves admissibility. A symmetric form on  $R^2$  is freely determined by three entries. Antisymmetry and admissibility give  $2\beta = \omega$ , hence the unique member  $\omega/2$ . Finally, for  $f(v) = s(v, v)/2$ , symmetry gives  $f(v + w) = f(v) + f(w) + s(v, w)$ , which is exactly the coboundary identity needed for  $\varphi_s$  to respect the two group laws.  $\square$

#### PROVED<sup>†</sup>

**Theorem 9.10** (Central simplicity and finite Stone–von Neumann theorem). *Let  $R$  be a finite commutative local ring, let  $\psi \in \text{Gen}(R)$ , let  $\beta \in \text{Adm}(\omega)$ , and suppose  $2 \in R^\times$ . Then  $A_{\psi, \beta}(V(R))$  is simple, has centre  $\mathbb{C}1$ , has complex dimension  $|R|^2$ , and is noncanonically isomorphic to  $M_{|R|}(\mathbb{C})$ . The group  $H_\beta(R)$  has exactly one irreducible unitary representation with central character  $\psi$ , up to unitary equivalence; its dimension is  $|R|$ , unitary intertwiners are unique up to  $U(1)$ , and every complex algebra automorphism of the Weyl algebra is inner.*

*Scope.* The condition  $2 \in R^\times$  is used to transport an arbitrary cocycle to the fixed matrix model. The matrix isomorphism depends on choices of realization and basis and is not part of the algebra datum. No frame-preserving automorphism group is computed.

---

**Provenance.** ids: D12–D16, FCR-ALG, FCR-SVN; proved in: theory/fcr-local.md §5–§6; tested by: theory/checks/fcr\_local\_check.py (G4,G5,G6), including the G6 arena mutation; reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* Averaging conjugation by all Weyl basis elements kills every nonzero basis coefficient because the pairing is nondegenerate. It follows that the centre is scalar and that every nonzero two-sided ideal contains 1. The reference operators  $Z(-b)X(a)$  give a nonzero representation on  $\ell^2(R)$ ; simplicity makes it injective and the equal dimensions make it onto  $\text{End}_{\mathbb{C}}(\ell^2(R))$ . The preceding transport handles arbitrary  $\beta$ .

Modules with central character  $\psi$  are precisely modules over this full matrix algebra. Its column module is the unique simple module and has dimension  $|R|$ . Schur’s argument makes intertwiners scalar; polar rescaling gives the  $U(1)$  statement. Twisting the column module by an algebra automorphism shows that the automorphism is conjugation by an invertible operator.  $\square$

**PROVED** <sup>†</sup>

**Theorem 9.11** (Lagrangians and Schrödinger models over a local ring). *Let  $R$  be a finite commutative local ring, let  $\psi \in \text{Gen}(R)$ , let  $\beta \in \text{Adm}(\omega)$ , and suppose  $2 \in R^\times$ . Every Lagrangian  $L \subseteq V(R)$  has  $|L| = |R|$ . The coordinate axes are Lagrangian, and for every ideal  $I \subseteq R$  so is*

$$I \oplus \text{Ann}(I).$$

*If  $R$  is not a field, then*

$$L_* := \text{soc}(R) \oplus \mathfrak{m}$$

*is a nonfree Lagrangian. Every algebra character  $\chi : A_L \rightarrow \mathbb{C}$  produces a simple induced module  $M_{L,\chi}$  of dimension  $|R|$ , and all these modules are isomorphic.*

*Scope.* This is a catalogue containing a general ideal family, not a classification of all Lagrangians over every local ring. In particular, it does not claim that every Lagrangian is free or belongs to the displayed ideal family.

---

**Provenance.** ids: D12–D16, FCR-POL; proved in: theory/fcr-local.md §7; tested by: theory/checks/fcr\_local\_check.py (G7), full submodule census at the two order-nine seeds; reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* Character orthogonality in the two orders gives  $|L| |L^{\perp\psi}| = |R|^2$ , so a self-perpendicular module has order  $|R|$ . Annihilator duality gives  $\text{Ann}(\text{Ann}(I)) = I$ , hence the ideal family. For a nonfield local ring,  $\text{Ann}(\mathfrak{m}) = \text{soc}(R)$  and  $\text{Ann}(\text{soc}(R)) = \mathfrak{m}$ , so  $L_*$  belongs to that family. Its quotient by  $\mathfrak{m}L_*$  needs at least two generators, whereas a free module of order  $|R|$  would have rank one.

The restriction of  $\beta$  to a Lagrangian is symmetric, and  $\chi_0(W(l)) = \psi(-\beta(l,l)/2)$  is an algebra character. The Weyl basis makes the algebra right-free over  $A_L$  on representatives for  $V(R)/L$ , so the induced module has dimension  $|R|$ ; uniqueness of the simple module then proves that all labels give isomorphic realizations.  $\square$

**PROVED** <sup>†</sup>

**Proposition 9.12** (The two presentation data and bare-algebra ambiguity). *Let  $R$  be a finite commutative local ring with  $2 \in R^\times$  and  $\text{Gen}(R) \neq \emptyset$ . Beyond  $R$ , the presentation uses exactly a generating character  $\psi \in \text{Gen}(R)$  and a cocycle  $\beta \in \text{Adm}(\omega)$ . Distinct generating characters give inequivalent representations of the same group  $H_\beta(R)$  because their central characters differ. All resulting bare algebras are isomorphic, but the torsor of bare-algebra isomorphisms has no point fixed by every target automorphism.*

*Scope.* The last assertion concerns all bare-algebra isomorphisms. It neither computes nor identifies a frame-preserving isomorphism torsor and makes no claim about a ring-side Weil or metaplectic symmetry.

---

**Provenance.** ids: D12–D16, FCR-CHOICE; proved in: theory/fcr-local.md §8; tested by: theory/checks/fcr\_local\_check.py (G8,G9); reviewed in: theory/verdicts/fcr-local-r1.md.

## 9.4 Clause-level return to an odd finite field

**PROVED**<sup>†</sup>

**Theorem 9.13** (Clause-level compatibility with the odd-field system). *Let  $R = \kappa$  be a finite field of odd characteristic and put  $q := |\kappa|$ . Specialization recovers on the nose:*

- (i) *the nontrivial-character set and its  $\kappa^\times$ -torsor of order  $q - 1$ ;*
- (ii) *bilinearity, strong alternation, and nondegeneracy of  $\omega$ , together with the zero radical of  $\psi \circ \omega$ ;*
- (iii) *the Weyl commutator and conjugation identities;*
- (iv) *simplicity, scalar centre, dimension  $q^2$ , and a noncanonical  $M_q(\mathbb{C})$  realization;*
- (v) *the unique irreducible unitary representation of dimension  $q$ , its  $U(1)$ -unique intertwiners, and innerness of algebra automorphisms;*
- (vi) *exactly  $q + 1$  Lagrangian lines,  $q$  algebra characters per line,  $q(q + 1)$  model labels, and one module-isomorphism class;*
- (vii) *the torsor of admissible cocycles of order  $q^3$ , its unique antisymmetric member  $\omega/2$ , and the centre-fixing transports;*
- (viii) *the two named presentation data, the distinct-central-character obstruction, and isomorphism of the resulting bare algebras.*

*The specialized definitions give the same  $V(\kappa)$ , form, admissibility equation, Weyl and group laws, and fixed operator  $W_{\beta_0} = Z(-b)X(a)$  with the same sign.*

*Scope.* This is deliberately a clause-level comparison. It does not compare the field theory's frame-preserving isomorphism torsor of order  $q^2|SL_2(\kappa)|$ ; the ring-side frame-preserving analysis is future work. It also does not compare the field theorem's cocycle or automorphism-group clauses, the  $SL_2(\kappa)$ -invariance of  $\omega/2$ , or the particular polarization-dependent matrix construction used in the field proof.

---

**Provenance.** ids: D12–D16, FCR-GEN, FCR-RAD, FCR-COMM, FCR-BETA-ODD, FCR-ALG, FCR-SVN, FCR-POL, FCR-CHOICE, FCR-REG; proved in: theory/fcr-local.md §9; tested by: theory/checks/fcr\_local\_check.py (G10), exact counts, form and radical tables, fixed matrices, rank, commutant, lines, half-census, and group profile; reviewed in: theory/verdicts/fcr-local-r1.md.

*Proof.* For a field,  $\mathfrak{m} = 0$ ,  $\text{soc}(R) = R$ , and the only ideals are 0 and  $R$ , so every nontrivial character is generating. Substitution in the preceding results proves every listed clause. A self-perpendicular subspace of the nondegenerate two-dimensional space is a line; nonzero vectors modulo  $\kappa^\times$  give  $(q^2 - 1)/(q - 1) = q + 1$  lines, and each line has  $q$  additive characters. No excluded symmetry or frame clause follows from these calculations.  $\square$

## 10 The five smallest local rings, worked in full

The three smallest orders of finite commutative local rings are 2, 3, and 4, and order 4 is a three-way tie. This section exercises every definition of Section 9 on the resulting five rings

$$\mathbb{F}_2, \quad \mathbb{F}_3, \quad \mathbb{F}_4, \quad \mathbb{Z}/4, \quad \mathbb{F}_2[\varepsilon]/(\varepsilon^2),$$

computing their Weyl–Heisenberg groups and complete irreducible representation catalogues. Every number in this section is verified by the exact-arithmetic census checker `theory/checks/small_rings_cat` written and run by an independent verifier lane that recomputed the orchestrator’s catalogue blind; the checker runs green and carries eight mutation modes, each of which fails at its registered gate. Statements here are census facts about these five rings; no generality beyond them is claimed except where a claim identifier is cited.

### 10.1 The data, ring by ring

All five rings are Frobenius: the socle is one-dimensional over the residue field, so generating characters exist and form a single free orbit of the unit group (the existence criterion and torsor structure of Section 9).

| $R$                         | $\mathfrak{m}$    | $\kappa$       | $\text{soc}(R)$   | $ R^\times  =  \text{Gen}(R) $ | reference $\psi$                        |
|-----------------------------|-------------------|----------------|-------------------|--------------------------------|-----------------------------------------|
| $\mathbb{F}_2$              | 0                 | $\mathbb{F}_2$ | $R$               | 1                              | $(-1)^x$                                |
| $\mathbb{F}_3$              | 0                 | $\mathbb{F}_3$ | $R$               | 2                              | $\zeta_3^x$                             |
| $\mathbb{F}_4$              | 0                 | $\mathbb{F}_4$ | $R$               | 3                              | $(-1)^{\text{Tr}(x)}$                   |
| $\mathbb{Z}/4$              | (2)               | $\mathbb{F}_2$ | (2)               | 2                              | $i^x$                                   |
| $\mathbb{F}_2[\varepsilon]$ | ( $\varepsilon$ ) | $\mathbb{F}_2$ | ( $\varepsilon$ ) | 2                              | $(-1)^{a+b}$ for $x = a + b\varepsilon$ |

---

**Provenance.** ids: FCR-GEN; proved in: `theory/fcr-local.md` §1; tested by: `small_rings_catalogue_check.py` (ring-data gates); reviewed in: `theory/verdicts/smallest-rings-verification.md`.

### 10.2 The Heisenberg groups are unitriangular matrix groups

**Proposition 10.1** (Matrix form of the reference-cocycle Heisenberg group). *For every finite commutative unital ring  $R$ , the map*

$$(t, (a, b)) \mapsto \begin{pmatrix} 1 & a & t \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$$

*is an isomorphism from the Heisenberg group of the reference cocycle  $\beta_0((a, b), (a', b')) = ab'$  onto the unitriangular group  $\text{UT}_3(R)$ .*

*Proof.* Multiplying the two displayed matrices gives upper-right entry  $t + t' + ab'$ , which is exactly the group law of the Heisenberg group of  $\beta_0$ ; the map is a bijection by construction.  $\square$

*Scope.* The identification is specific to the reference cocycle: at residue characteristic 2 other admissible cocycles can give non-isomorphic groups (the quaternion type over  $\mathbb{F}_2$  below), and those are not unitriangular in this way.

---

**Provenance.** ids: D16; proved in: this section; tested by: `small_rings_catalogue_check.py` (cocycle gate); reviewed in: `theory/verdicts/smallest-rings-verification.md`.

The five groups, exactly:

| $R$                         | $ H $ | centre             | exponent | order profile        | identification                           |
|-----------------------------|-------|--------------------|----------|----------------------|------------------------------------------|
| $\mathbb{F}_2$              | 8     | $\mathbb{Z}/2$     | 4        | 1:1, 2:5, 4:2        | dihedral $D_4$                           |
| $\mathbb{F}_3$              | 27    | $\mathbb{Z}/3$     | 3        | 1:1, 3:26            | extraspecial $3_+^{1+2}$                 |
| $\mathbb{F}_4$              | 64    | $(\mathbb{Z}/2)^2$ | 4        | 1:1, 2:27, 4:36      | characteristic-two Heisenberg            |
| $\mathbb{Z}/4$              | 64    | $\mathbb{Z}/4$     | 8        | 1:1, 2:7, 4:40, 8:16 | $\text{UT}_3(\mathbb{Z}/4)$              |
| $\mathbb{F}_2[\varepsilon]$ | 64    | $(\mathbb{Z}/2)^2$ | 4        | 1:1, 2:31, 4:32      | $\text{UT}_3(\mathbb{F}_2[\varepsilon])$ |

The three order-64 groups are pairwise non-isomorphic:  $\mathbb{Z}/4$  is separated by its cyclic centre and its sixteen elements of order 8, while  $\mathbb{F}_4$  and  $\mathbb{F}_2[\varepsilon]$  share centre and exponent but differ in involution count. The order-8 elements over  $\mathbb{Z}/4$  come from the square law  $(t, v)^2 = (2t + \beta_0(v, v), 2v)$ : the fourth power is  $(2\beta_0(v, v), 0)$ , nonzero exactly when  $ab$  is odd. Over  $\mathbb{F}_2[\varepsilon]$  the identity  $2v = 0$  kills this mechanism: the jet direction deepens the ring without deepening the clock. Mixed versus equal characteristic is thus visible in the bare group.

*Remark.* At  $\mathbb{F}_3$  the group is independent of the cocycle choice (the odd-characteristic transport of Section 9). At  $\mathbb{F}_2$  the eight admissible cocycles split six-to-two into a dihedral type and a quaternion type, separated by the Arf invariant of  $Q_\beta(v) = \beta(v, v)$ ; the reference cocycle is of dihedral type. For  $\mathbb{Z}/4$  and  $\mathbb{F}_2[\varepsilon]$  the classification of the 64-element cocycle torsor is open and is the target of the running residue-characteristic-two increment.

---

**Provenance.** ids: WH-BETA-d, FCR-BETA-ODD; proved in: theory/wh-kappa-choice.md §6; theory/fcr-local.md §4; tested by: beta\_census.py; small\_rings\_catalogue\_check.py; reviewed in: theory/verdicts/wh-kappa-adjudication.md.

### 10.3 The complete representation catalogues

The commutator subgroup is the full centre, so the abelianization is the phase module  $R \oplus R$ , and irreducible representations are stratified by the character of the centre,  $\psi_u = \psi(u \cdot)$  for  $u \in R$ .

**Proposition 10.2** (Catalogue census for the five smallest local rings). *For each of the five rings above and each  $u \in R$ , the number of irreducible unitary representations of the Heisenberg group with central character  $\psi_u$  is  $|\text{Ann}(u)|^2$ , and each has dimension  $|R|/|\text{Ann}(u)|$ . In particular the total count of irreducibles equals the number of conjugacy classes,*

$$\#\{\text{classes}\} = |R|^2 + \sum_{u \neq 0} |\text{Ann}(u)|^2 \quad (5, 11, 19, 22, 22 \text{ respectively}),$$

and the sum of squared dimensions equals  $|R|^3$ .

*Scope.* A census statement about these five rings only, established by exhaustive computation; the same stratification pattern for a general finite commutative local ring is expected but not claimed here — its general proof belongs to the planned non-Frobenius/collapse increment.

---

**Provenance.** ids: (census); proved in: small\_rings\_catalogue\_check.py (class-count and catalogue gates); tested by: same checker, mutation modes red-class-count, red-catalogue; reviewed in: theory/verdicts/smallest-rings-verification.md.

The three strata carry distinct physics:

- *Bottom*,  $u = 0$ : the  $|R|^2$  characters of the abelianization — the classical shadow.
- *Top*,  $u \in R^\times$ : one Schrödinger representation per generating character, of dimension  $|R|$ , in the fixed model  $W(a, b) = Z(-b)X(a)$  on  $\ell^2(R)$ : the qubit Pauli pair for  $\mathbb{F}_2$ ; the qutrit clock-shift for  $\mathbb{F}_3$ ; two  $\mathbb{F}_4$ -labelled qubits (three inequivalent copies, one per generating character); the ququart clock-shift  $Z = \text{diag}(1, i, -1, -i)$ ,  $ZX = iXZ$  for  $\mathbb{Z}/4$  (two copies); and the jet-paired four-dimensional representation for  $\mathbb{F}_2[\varepsilon]$  (two copies). Irreducibility is pinned by exact trace-orthogonality of the  $|R|^2$  Weyl operators.

- *Middle*,  $u \in \mathfrak{m} \setminus 0$  (only the two thickened rings): four two-dimensional representations each, with central character supported on the socle. These are qubit representations of the collapsed quotient  $R/\text{Ann}(u) \cong \mathbb{F}_2$  twisted by characters of the radical of the degenerate phase pairing. The verifier’s decomposition of the four-dimensional model with this central character found exactly two inequivalent two-dimensional blocks, each of multiplicity one — the model contains two of the four classes.

A ring is thickened exactly when its catalogue has a middle stratum: the representation theory detects the nilpotents. Two honest cautions from the blind verification:  $\mathbb{F}_4$  has *three* top-stratum representations, not two; and as bare dimension–multiplicity tables the  $\mathbb{Z}/4$  and  $\mathbb{F}_2[\varepsilon]$  catalogues coincide ( $16 \times \dim 1$ ,  $4 \times \dim 2$ ,  $2 \times \dim 4$ ) — the catalogue separates the two rings only through its central-character labelling and the underlying groups, not through the numbers alone.

#### 10.4 Lagrangian labels and models

| $R$                         | Lagrangians | free | non-free | model labels |
|-----------------------------|-------------|------|----------|--------------|
| $\mathbb{F}_2$              | 3           | 3    | 0        | 6            |
| $\mathbb{F}_3$              | 4           | 4    | 0        | 12           |
| $\mathbb{F}_4$              | 5           | 5    | 0        | 20           |
| $\mathbb{Z}/4$              | 7           | 6    | 1        | 28           |
| $\mathbb{F}_2[\varepsilon]$ | 7           | 6    | 1        | 28           |

The fields show the  $q + 1$  lines of the projective line; the thickened rings show the six free lines of  $\mathbb{P}^1(R)$  plus the non-free witness  $\text{soc}(R) \oplus \mathfrak{m}$  of Section 9. All labels with a fixed generating central character present one representation, but the non-free label is a genuinely different presentation: position and momentum are each sampled at depth  $\mathfrak{m}$ , a mixed frame with no field analogue.

---

**Provenance.** ids: FCR-POL; proved in: theory/fcr-local.md §7; tested by: small\_rings\_catalogue\_check.py (Lagrangian gate, mutation mode red-drop-nonfree); reviewed in: theory/verdicts/fcr-local-adjudication.md.

## 11 Sidequest: quantum mechanics over the field with one element

**Opened 6 September 2026.** This is a separate, literature-grounded exploration of arithmetic quantum mechanics over  $\mathbb{F}_1$ , also written  $\mathbb{F}_{\text{un}}$ . Its north star is a precisely formulated quantum theory, with particular attention to the analogue of a qubit and to composition. The different established analogies are retained as separate candidates.

**North-star clarification, 7 September 2026.** The primary object is the family of sub-systems and their assembly rules. An acceptable endpoint must admit a quantum operational realization by  $C^*$ -algebras, normalized positive functionals or density operators, CP dynamics and Born probabilities. The proposed strategy extracts a compositional category that remembers the prime before formulating its field-with-one-element counterpart. Fibonacci is a test of the kind of composition sought, not an asserted endpoint. A strong monoidal fibre functor to ordinary Hilbert spaces is not a universal requirement: composite observable algebras may contain additional fusion observables. Compatible preparations and extensions of channels to composites are part of the target. Classical or combinatorial state sets alone do not satisfy this operational criterion.

The accompanying research map is `docs/sidequests/f1-qm.md`; primary source bodies, versions and retrieval hashes are indexed in `refs/LEDGER.md`. The finite-ring campaign is not superseded.

There is substantial relevant literature, but the constructions preserve different structures. A field may specify the arithmetic base, the state coefficients, a counting parameter, or a deformation parameter. Confusing these roles would make the comparison misleading. In the present campaign, the arithmetic base is a finite field or ring and the amplitudes remain complex. The direct “quantum  $\mathbb{F}_{\text{un}}$ ” papers instead change the coefficient object of the states themselves.

### 11.1 What could count as a qubit?

Two distinguishable states, a two-dimensional Hilbert space, a simple object of categorical dimension two, and two simple objects of a category are different notions. The following table makes the distinction explicit.

| Approach                       | Qubit analogue                                                                                                           | Composition                                                                                                                                 |
|--------------------------------|--------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| Pointed $\mathbb{F}_1$ modules | A basepoint and two basis points. Complex linearization of normal partial maps gives $(\mathbb{C}^2, M_2(\mathbb{C}))$ . | Smash product has four nonzero basis points and realizes as the Hilbert tensor product. Native basis states have no superpositions.         |
| Cyclotomic modules             | Two free $\mu_N$ -orbits. For configuration $C_2$ and phase level two, clock and shift give the Heisenberg qubit.        | Balanced smash of modules; central product of Heisenberg groups; ordinary tensor after complex realization.                                 |
| Absolute Cartesian frames      | Rank two has $N + 2$ projective rays. At bare $\mathbb{F}_1$ these are 10, 01, 11.                                       | The rank-four frame has fifteen rays at $N = 1$ : nine product rays and six non-product rays. The operation on pairs is the Kronecker rule. |
| Thin symplectic geometry       | Two coordinate polarizations, suggestive of position and momentum. These are not two Hilbert-state vectors.              | Orthogonal sums of phase geometry; Hilbert tensor requires a realization.                                                                   |

| Approach                           | Qubit analogue                                                                                                         | Composition                                                                                                                                              |
|------------------------------------|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Blueprint Heisenberg geometry      | A chosen $\mathbb{F}_2$ fibre gives the reference order-eight group and its two-dimensional central-character module.  | Ring realization and phase data precede the physical tensor product.                                                                                     |
| Bands and crowds                   | A relational Heisenberg object, with 27 signed or eight Krasner points. Neither number is a count of qubit states.     | Geometric products exist; physical composition needs a specified phase and representation construction.                                                  |
| Hall/groupoid oscillator           | No preferred qubit in one oscillator. Two colours give a two-dimensional one-particle sector after realization.        | Full Fock spaces tensor on disjoint colour sets. Two identical bosons in two modes have dimension three; two distinguishable qubits have dimension four. |
| Representation categories          | The two-dimensional simple object of $\text{Rep}(D_8)$ or $\text{Rep}(Q_8)$ .                                          | Internal fusion splits its square into four charges. Independent copies instead use an external tensor and a matching-central-character quotient.        |
| Quantum torus                      | $VU = -UV$ , $U^2 = V^2 = 1$ gives $M_2(\mathbb{C})$ .                                                                 | Two selected central fibres give $M_4(\mathbb{C})$ .                                                                                                     |
| Arithmetic descent and Bost–Connes | A specified finite fibre or a chosen two-level encoding; no canonical qubit is supplied by the global arithmetic data. | Frobenius and transfer correspondences; tensor products of statistical systems after realization.                                                        |
| Tropical and motivic routes        | Support or incidence information, without a specified positive two-state quantum system.                               | The relevant semiring, geometric or categorical tensor must be named.                                                                                    |

## 11.2 The developed geometric and physical analogies

**Counting and thin geometry.** The projective count  $1 + q + \cdots + q^{r-1}$  specializes to  $r$  at  $q = 1$ ; Gaussian coefficients specialize to subset counts. Tits’s thin geometry therefore retains coordinate points, subsets and permutation symmetries. For symplectic rank  $r$  the reflection group is  $W(C_r) = (C_2)^r \rtimes S_r$ , acting on symplectic pairs. A coordinate Lagrangian selects one direction from each pair, giving  $2^r$  choices. This is the incidence content of the familiar specialization of  $\prod_{i=1}^r (q^i + 1)$ . The geometric analogies are developed in [Deitmar’s \*Schemes over  \$\mathbb{F}\_1\$\* , section 5](#), and [Lorscheid’s Tits–Weyl models](#). Here a *Weyl group* is a reflection group, not the group of Weyl displacement operators. Moreover,  $q^3 \mapsto 1$  for the order of  $\text{UT}_3(\mathbb{F}_q)$  is a polynomial specialization, not a limit of groups or a proof that its  $\mathbb{F}_1$  geometry is empty.

**Direct quantum  $\mathbb{F}_{\text{un}}$ .** [Chang, Lewis, Minic and Takeuchi](#) construct a  $q = 1$  coordinate-state specialization of finite-field amplitude models: superposition disappears. This is different from this campaign’s complex Hilbert space  $\ell^2(\mathbb{F}_q)$ . [Thas’s \*Absolute Quantum Theory\*](#) uses larger Cartesian frames and develops monomial dynamics and quantum-information analogies. Its Hermitian expression is partial: a sum is allowed only when at most one term is nonzero. Orthogonality is disjointness of supports. The involution determines the unitary subgroup; inversion of phases makes all phase-permutation matrices unitary in this sense. Its cloning discussion distinguishes simple projective rays, which can be copied, from all frame states. These statements do not supply an ordinary positive-definite Hilbert inner product on the frame.

**An existing geometric Heisenberg model.** Lorscheid’s Proposition 5.5 constructs Tits–Weyl models for standard parabolic unipotent radicals of  $\mathrm{GL}_n$ . For the upper-triangular Borel of  $\mathrm{GL}_3$ , this is the reference Heisenberg group

$$h(a, b, t) = \begin{pmatrix} 1 & a & t \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}, \quad h(a, b, t)h(c, d, u) = h(a + c, b + d, t + u + ad).$$

Its underlying blue scheme is affine three-space. Its Weyl extension has one point, while ring base extension recovers the full unipotent group. Thus a positive  $\mathbb{F}_1$  model already exists; selecting a central character and constructing its representation are further steps. The whole polarizing-cocycle torsor, especially its characteristic-two types, is not supplied by that reference model. Likewise, a Tits lift of a reflection is not automatically a metaplectic or Weil lift.

**Cyclotomic conventions.** For the raw phase monoid  $\{0\} \cup \mu_N$ , integral realization gives  $\mathbb{Z}[\mu_N] \simeq \mathbb{Z}[t]/(t^N - 1)$ . A cyclotomic blueprint may instead impose additive relations giving  $\mathbb{Z}[\zeta_N]$ . A faithful complex phase character is another named realization datum. These conventions are compared in [Lorscheid’s \*A blueprinted view on  \$\mathbb{F}\_1\$ -geometry\*](#), section 1.1.4. The group  $\mu_{q-1}$  reflects multiplicative field data; finite-field Weyl phases come from additive characters and lie in  $\mu_p$ . These two phase counts must not be identified.

### 11.3 Pointed modules, ordinary qubits and absolute frames

**Definition 11.1** (Normal pointed maps and their two products). For  $r \geq 0$  let  $V_r = \{*, 1, \dots, r\}$ . A normal pointed map preserves  $*$  and has at most one preimage of each non-basepoint; it is a partial injection of the nonzero sets. Wedge identifies basepoints, while smash collapses all pairs containing a basepoint in the Cartesian product. Put  $L(S) = \mathbb{C}[S \setminus \{*\}]$ , with its basis orthonormal, and let  $B(S)$  be the complex span of realized normal endomorphisms in  $\mathrm{End}_{\mathbb{C}} L(S)$ .

*Scope.* These objects have no native addition of vectors. The image algebra  $B(S)$  is explicitly a complex realization.

---

**Provenance.** ids: D1010; proved in: definitions.md; tested by: f1\_check.py (M13); reviewed in: supporting comparison.

**Proposition 11.2** (The pointed-set qubit after operator realization). **SKETCH** *For every positive-rank finite pointed set  $S$ ,  $B(S) = \mathrm{End}_{\mathbb{C}} L(S)$ . For finite pointed sets  $S, T$ ,*

$$L(S \wedge T) \simeq L(S) \otimes L(T).$$

*For positive ranks this identifies the image algebras with their tensor product. In particular,  $V_2$  realizes to  $(\mathbb{C}^2, M_2(\mathbb{C}))$ , and  $V_2 \wedge V_2 = V_4$  realizes to two qubits.*

*Scope.* The realization retains a distinguished basis. The concrete image algebra is not the unreduced partial-injection monoid algebra. Algebras are transported along bijections; arbitrary partial maps do not give unital maps between full endomorphism algebras.

---

**Provenance.** ids: F1-NORMAL; proved in: f1-comparisons.md, section 3; tested by: f1\_check.py (M13); reviewed in: supporting SKETCH.

*Proof.* The partial map taking  $j$  to  $i$  and everything else to the basepoint realizes as  $E_{ij}$ . These matrix units span the full algebra, and their partial inverses realize adjoints. Smash products of these maps give  $E_{ij} \otimes E_{kl}$ . The basis identification proves the tensor claim.  $\square$

At rank two the partial-injection monoid has seven elements, including zero. Its contracted monoid algebra has dimension six, whereas its image has dimension four: linear realization imposes relations such as  $I = E_{00} + E_{11}$ . Keeping only the two bijections gives the commutative span of  $I$  and the swap. Thus the class of allowed morphisms matters. The pointed-module, normal-map and base-change conventions come from [Szczesny](#), pages 5–8.

**Definition 11.3** (Cyclotomic projective Cartesian frames). Fix an abstract cyclic group  $\mu_N$  of order  $N \geq 1$  and put  $F_N = \{0\} \sqcup \mu_N$ , with absorbing zero. For  $r \geq 1$ , define

$$P_N(r) = (F_N^r \setminus \{(0, \dots, 0)\}) / \mu_N$$

by simultaneous scalar multiplication. The product-state map  $P_N(r) \times P_N(s) \rightarrow P_N(rs)$  sends  $([x], [y])$  to  $[(x_i y_j)_{i,j}]$ . Rays outside its image are called nonproduct rays in this factorization sense.

*Scope.* A Cartesian frame is different from a free pointed module. No probability rule or total Hermitian form is stipulated.

---

**Provenance.** ids: D1009; proved in: definitions.md; tested by: f1.check.py (M11); reviewed in: support-ing comparison.

**Proposition 11.4** (Frame qubits and their nonproduct rays). **SKETCH** For every  $N \geq 1$ ,

$$|P_N(r)| = \frac{(N+1)^r - 1}{N}.$$

The rank-two product-state map is injective, so the rank-four frame has  $(N+2)^2$  product rays and  $N(N+1)(N+2)$  nonproduct rays.

*Scope.* Nonproduct means failure of tensor factorization. It does not by itself establish a Born rule, interference or Bell correlations.

---

**Provenance.** ids: F1-FRAME; proved in: f1-comparisons.md, section 2; tested by: f1.check.py (M11); reviewed in: supporting SKETCH.

*Proof.* Scalar multiplication acts freely on any nonzero tuple. For a nonzero product array, ratios within a nonzero row and column recover both factors up to scalars. The ray counts then follow by subtraction.  $\square$

For  $N = 1$  the one-system rays are 10, 01, 11; the composite has fifteen rays, nine product and six nonproduct. The diagonal array  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is a nonproduct because product supports are rectangular. This larger frame is the reason a nonadditive theory can retain a factorization notion of entanglement while the coordinate-only model does not. In the strict pointed-map category, only simple vectors are morphisms from the unit; the additional frame states require an explicitly enlarged state notion.

## 11.4 A finite cyclotomic Weyl–Heisenberg formulation

**Definition 11.5** (Cyclotomic modules and named complex realization). Fix  $N \geq 1$ , an abstract cyclic group  $\mu_N$  of order  $N$ , and a named faithful character  $\iota : \mu_N \rightarrow \mathbb{C}^\times$ . Let  $\text{Free}_*^{\mu_N}$  be finite pointed sets with a  $\mu_N$  action fixing the basepoint and free on its complement; maps are pointed equivariant maps. Define

$$\mathbb{C}_\iota(S) = \mathbb{C}[S] / (e_0, e_{us} - \iota(u)e_s)$$

and use the induced linear maps on morphisms.

*Scope.* The phase embedding is part of the realization datum. The objects are free phase sets, not Cartesian frames.

---

**Provenance.** ids: D1001; proved in: definitions.md; tested by: f1\_check.py; reviewed in: f1-r1.md.

**Definition 11.6** (Hyperbolic phase data and the pointed Schrödinger action). For a finite abelian group  $A$  with exponent dividing  $N$ , put

$$A^\vee = \text{Hom}(A, \mu_N), \quad V_A = A \times A^\vee, \quad c_A((a, \chi), (b, \eta)) = \eta(a)^{-1},$$

$$\kappa_A((a, \chi), (b, \eta)) = \chi(b)\eta(a)^{-1}.$$

Define  $H_N(A) = \mu_N \times V_A$  by  $(u, v)(u', w) = (uu'c_A(v, w), v + w)$  and adjoin an absorbing zero to get  $H_N(A)^0$ . Put  $S_A = \{0\} \sqcup (\mu_N \times A)$  and stipulate

$$(u, a, \chi)[z, x] = [uz\chi(x + a), x + a], \quad 0_H s = 0_S.$$

On its realization let  $X(a)e_x = e_{x+a}$ ,  $Z(\chi)e_x = \iota(\chi(x))e_x$ , and  $W(a, \chi) = Z(\chi)X(a)$ .

*Scope.* The abelian configuration group is extra data. No field structure, division by two or identification  $A \simeq A^\vee$  is assumed.

---

**Provenance.** ids: D1002, D1003; proved in: definitions.md; tested by: f1\_check.py (M1–M3); reviewed in: f1-r1.md.

**Theorem 11.7** (The finite cyclotomic Schrödinger system). PROVED Fix  $N \geq 1$ , cyclic  $\mu_N$  of order  $N$ , a faithful  $\iota : \mu_N \rightarrow \mathbb{C}^\times$ , and a finite abelian  $A$  of exponent dividing  $N$ . For the associated phase system,  $|A^\vee| = |A|$ , characters extend from subgroups and separate points, and  $\kappa_A$  is perfect. The displayed multiplication makes  $H_N(A)$  a group with centre exactly  $\mu_N$ , acting faithfully on  $S_A$ . After the named complex realization,

$$W(a, \chi)W(b, \eta) = \iota(\eta(a)^{-1})W(a + b, \chi\eta), \quad \text{Tr}(W(v)^*W(w)) = |A|\delta_{v,w}.$$

These operators span  $\text{End}_{\mathbb{C}}(\mathbb{C}[A])$ . The irreducible unitary representation of  $H_N(A)$  with central character  $\iota$  is unique up to equivalence, has dimension  $|A|$ , and has intertwiners unique up to  $U(1)$ .

*Scope.* This is finite Heisenberg theory for abelian configurations, with an explicit pointed module. A universal representation theorem for all proposed  $\mathbb{F}_1$  geometries is not asserted.

---

**Provenance.** ids: F1-DUAL, F1-WEYL, F1-REAL; proved in: f1-cyclotomic.md, sections 0–2; tested by: f1\_check.py (M1–M3); reviewed in: f1-adjudication.md.

*Proof.* Character extension follows by adjoining a cyclic generator and choosing a root in  $\mu_N$ ; the exponent hypothesis ensures the required root exists. Restriction and induction on the group order give  $|A^\vee| = |A|$ . A nontrivial character sums to zero by translating its sum. Character evaluation verifies the cocycle identity and action. Testing the two coordinate subgroups proves perfection. The trace vanishes unless both translation and character are trivial; the resulting  $|A|^2$  orthogonal operators form a basis of the full matrix algebra. Matrix units then prove uniqueness of its simple module and scalar uniqueness of intertwiners.  $\square$

**Definition 11.8** (Fixed-level isomorphisms and balanced composition). Let  $\text{FinAb}_{\widetilde{N}}$  have the above configurations and their group isomorphisms. For  $f : A \rightarrow B$ , put  $f_*\chi = \chi \circ f^{-1}$ ,  $S(f)[u, x] = [u, f(x)]$  and  $H(f)(u, a, \chi) = (u, f(a), f_*\chi)$ . Define

$$S \wedge_{\mu_N} T = (S \wedge T)/([us, t] \sim [s, ut]), \quad H \boxtimes K = (H \times K)/\{(u, u^{-1}) : u \in \mu_N\}.$$

In the last expression the two groups have specified central copies of  $\mu_N$ .

*Scope.* Functoriality is asserted on isomorphisms at a common phase level. Arbitrary configuration homomorphisms are not silently included.

---

**Provenance.** ids: D1004; proved in: definitions.md; tested by: f1\_check.py (M6); reviewed in: f1-r1.md.

**Theorem 11.9** (Composition of independent phase systems). **PROVED** *Fix  $N \geq 1$  and a faithful phase realization  $\iota$ . The module, group and Hilbert assignments on  $\text{FinAb}_N^\approx$  are strong symmetric monoidal, with comparisons*

$$S_A \wedge_{\mu_N} S_B \simeq S_{A \times B}, \quad H_N(A) \boxtimes H_N(B) \simeq H_N(A \times B), \\ \mathbb{C}[A] \otimes \mathbb{C}[B] \simeq \mathbb{C}[A \times B].$$

*At  $N = 1$  the exponent condition forces  $A = 0$  and a one-dimensional phase-system realization, while free pointed modules of arbitrary rank still exist.*

*Scope.* The group tensor is a central quotient, not a Cartesian product retaining two independent centres. The level-one statement is restricted to perfect phase systems, not all  $\mathbb{F}_1$  quantum routes.

---

**Provenance.** ids: F1-FUNCT,F1-ONE; proved in: f1-cyclotomic.md, sections 3,5; tested by: f1\_check.py (M1,M6); reviewed in: f1-adjudication.md.

*Proof.* The phase-set comparison sends  $[[u, a], [v, b]]$  to  $[uv, (a, b)]$ . The corresponding group map multiplies central phases and takes the external character. Its kernel is the antidiagonal centre. Every associativity diagram multiplies the same phases and associates the same tuple, proving coherence. The Hilbert comparison is the basis map.  $\square$

**Definition 11.10** (Central pushout from a finite local ring). Let  $R$  be finite commutative unital local with  $1 \neq 0$ , and let  $\psi : (R, +) \rightarrow \mu_N$  be named, with exponent of  $(R, +)$  dividing  $N$ . Assume  $\iota\psi$  is generating, meaning its kernel contains no nonzero ideal. Let  $H_{\beta_0}(R)$  have triples  $(t, a, b)$  and product  $(t + u + ad, a + c, b + d)$ . Put

$$\chi_b(x) = \psi(-bx), \quad P_\psi = (\mu_N \times H_{\beta_0}(R)) / \{(\psi(t)^{-1}, (t, 0, 0)) : t \in R\}.$$

*Scope.* A generating character need not be faithful on the additive group. The raw Heisenberg group and its phase image are distinguished.

---

**Provenance.** ids: D1005,D12,D13,D16; proved in: definitions.md; tested by: f1\_check.py (M4,M5); reviewed in: f1-r1.md.

**Theorem 11.11** (Exact recovery of the arithmetic reference convention). **PROVED** *Let  $R$  be a finite commutative unital local ring with  $1 \neq 0$ , exponent of  $(R, +)$  dividing  $N \geq 1$ , and a named  $\psi : (R, +) \rightarrow \mu_N$  such that the faithful complex realization  $\iota\psi$  is generating. Then  $b \mapsto \chi_b$  identifies  $(R, +)$  with its phase dual and  $P_\psi \simeq H_N((R, +))$  by  $[u; (t, a, b)] \mapsto (u\psi(t), a, \chi_b)$ . The raw group map has central kernel  $\ker \psi$ . Its realized operators are*

$$W(a, \chi_b)e_y = (\iota\psi)(-b(y + a))e_{y+a} = Z(-b)X(a)e_y,$$

*with product cocycle  $(\iota\psi)(ab')$ , uniformly including characteristic two.*

*Scope.* Only the reference cocycle is recovered here. This does not identify all raw polarizing-cocycle groups with one phase group.

---

**Provenance.** ids: F1-RING; proved in: f1-cyclotomic.md, section 4; tested by: f1\_check.py (M4,M5); reviewed in: f1-adjudication.md.

For example, over  $\mathbb{F}_4$  a nontrivial trace character has a two-element kernel. Its raw Heisenberg group has order 64 and its phase image order 32. Over  $\mathbb{F}_2$ , the order-eight reference group acts on the usual qubit. This supplies a checked bridge to the main campaign.

## 11.5 Fourier transform and a precisely scoped additive extension

**Definition 11.12** (Quadratic phases, strict normalizer and Fourier kernel). A quadratic phase is  $\theta : A \rightarrow \mu_N$ ,  $\theta(0) = 1$ , for which  $B_\theta(a, x) = \theta(a + x)\theta(a)^{-1}\theta(x)^{-1}$  is a bicharacter. For  $t \in A$ ,  $f \in \text{Aut}(A)$  put  $T_{t,f,\theta}[u, x] = [u\theta(x), f(x) + t]$ . The strict normalizer is the normalizer of the  $\mu_N$ -level Weyl group in  $\text{Aut}_{\text{Free}^{\mu_N}}(S_A)$ . Define

$$F_A e_x = \sum_{\rho \in A^\vee} \iota(\rho(x)) e_\rho, \quad \text{ev}_a(\rho) = \rho(a), \quad J_A(a, \chi) = (\chi^{-1}, \text{ev}_a), \quad r_A(a, \chi) = \chi(a).$$

*Scope.* A self-Fourier endomorphism additionally needs a pairing identifying  $A$  with its dual. No such pairing is suppressed.

---

**Provenance.** ids: D1006; proved in: definitions.md; tested by: f1\_check.py (M7); reviewed in: f1-r1.md.

**Theorem 11.13** (What remains monomial and what Fourier requires). **PROVED** Fix  $N \geq 1$ , a faithful  $\iota$ , and finite abelian  $A$  of exponent dividing  $N$ . Its strict normalizer consists exactly of the  $T_{t,f,\theta}$  and central phases. Its phase-space maps are

$$(a, \chi) \mapsto (f(a), (\chi B_\theta(a, -)) \circ f^{-1}).$$

For  $|A| > 1$ ,  $F_A$  has full column support and cannot be the realization of a strict pointed map. Nevertheless,

$$F_A^* F_A = |A| I, \quad F_A W(v) = \iota(r_A(v)) W(J_A v) F_A.$$

*Scope.* Only bicharacters admitting a  $\mu_N$ -valued quadratic refinement occur in the strict subgroup. Enlarging phases alone does not make Fourier monomial. No canonical Weil splitting is asserted.

---

**Provenance.** ids: F1-MON; proved in: f1-cyclotomic.md, section 6; tested by: f1\_check.py (M7, Fourier only); reviewed in: f1-adjudication.md.

**Definition 11.14** (Framed cyclotomic kernels and lifted phase isomorphisms). Put  $K_N = \mathbb{Q}[\mu_N]/\ker \iota$ . The category  $\text{Mat}(K_N)$  has finite sets  $I$  as objects and  $J \times I$  matrices over  $K_N$  as morphisms; tensor is Cartesian product of sets and Kronecker product of matrices. The associated modules are explicitly framed by  $[1, i]$  in  $S_I = \{0\} \sqcup (\mu_N \times I)$ . Let  $\text{PMat}(K_N)$  be invertible matrices modulo  $K_N^\times$ , realized by entrywise evaluation.

The groupoid  $\text{LiftHyp}_N$  has configurations  $A$  as objects. An arrow is  $(g, r)$  with  $g : V_A \rightarrow V_B$  a group isomorphism preserving  $\kappa$  and  $r : V_A \rightarrow \mu_N$  satisfying

$$r(v)r(w)c_B(gv, gw) = c_A(v, w)r(v + w).$$

It denotes the centre-fixing map  $(u, v) \mapsto (ur(v), g(v))$ ; composition composes these maps. Tensor is  $A \times B$  on objects and  $(g \times h, (v, w) \mapsto r(v)s(w))$  on arrows, with the canonical reordering of phase factors. The unit is  $A = 0$  and the symmetry is the swap.

*Scope.* This target is an additive cyclotomic envelope. It is not claimed to be a category of  $\mathbb{F}_1$  schemes; arbitrary free torsors are not given coordinate matrices without a frame.

---

**Provenance.** ids: D1007; proved in: definitions.md; tested by: no general checker; reviewed in: f1-adjudication.md, O1–O3 repair.

**Proposition 11.15** (Projective realization of lifted phase symmetries). **SKETCH** For  $N \geq 1$  and a named faithful  $\iota$ , there is a projective strong symmetric monoidal functor

$$Q_N : \text{LiftHyp}_N \longrightarrow \text{PMat}(K_N)$$

defined by the Egorov intertwining equations. Its complex realization is projectively unitary; affine-quadratic maps and Fourier have the values above.

*Scope.* The lift is input data. Geometric descent of the additive envelope and a lift of every symplectic transformation remain separate questions.

---

**Provenance.** ids: F1-CORR; proved in: f1-cyclotomic.md, section 7; tested by: not claimed as computationally proved; reviewed in: supporting SKETCH after scope repair.

*Proof.* The intertwining equations have coefficients in  $K_N$  and a one-dimensional solution space after embedding in  $\mathbb{C}$ . Gaussian elimination has the same rank over either field, so there is a nonzero solution over  $K_N$ . Irreducibility makes it invertible; uniqueness up to scalar gives projective composition and tensor coherence. Unitary normalization is performed only after complex realization.  $\square$

## 11.6 Relational Heisenberg geometry over bands

**Definition 11.16** (Bands and the upper-unitriangular crowd). A band is a commutative pointed multiplicative monoid  $B$  with a null ideal  $N_B$  of formal sums, in which every  $a$  has a unique  $-a$  with  $a + (-a) \in N_B$ ; morphisms preserve null sums. The regular partial field  $\mathbb{F}_1^\pm = \{0, 1, -1\}$  uses sums vanishing in  $\mathbb{Z}$ . The Krasner hyperfield  $\mathbb{K} = \{0, 1\}$  declares a sum null exactly when it has zero or at least two nonzero terms. Define  $H_{\text{crowd}}(B) = B^3$ , represented by  $h(a, b, t)$  as above, with identity  $(0, 0, 0)$ . Its colaw consists of triples for which

$$\sum_i a_i, \quad \sum_i b_i, \quad t_1 + t_2 + t_3 + a_1 b_2 + a_1 b_3 + a_2 b_3$$

and both cyclic variants of the last sum lie in  $N_B$ . The inverse set of  $h$  consists of  $k$  with  $(h, k, 1)$  in the colaw; the product set  $h * k$  consists of  $c$  for which some  $d$  has both  $(h, k, d)$  and  $(c, d, 1)$  in the colaw.

*Scope.* Nullity is part of the band, not an invented binary addition. The signed band and the raw pointed phase monoid have different structure.

---

**Provenance.** ids: D1008; proved in: definitions.md; tested by: f1\_check.py (M10); reviewed in: supporting comparison.

**Proposition 11.17** (A band-level Heisenberg crowd and its ring fibres). **SKETCH** *The functor  $B \mapsto H_{\text{crowd}}(B)$  on bands, with the displayed colaw, is an affine algebraic crowd. For every commutative ring  $R$ , its fibre is exactly  $\text{UT}_3(R)$ . The signed fibre has 27 points, 319 colaw triples and 23 points admitting an inverse inside that fibre. The Krasner fibre has eight points and 152 colaw triples, with multivalued inverses.*

*Scope.* These finite fibres are not ordinary groups. Even a crowd product with the identity may be empty when its required inverse mediator is absent. No Hilbert representation is inferred from the point counts.

---

**Provenance.** ids: F1-CROWD; proved in: f1-comparisons.md, section 1; tested by: f1\_check.py (M10); reviewed in: supporting SKETCH.

This specializes the algebraic  $\text{SL}_3$  crowd of [Lorscheid–Thas \(2023\)](#), section 5.4. Restricting to an identity-containing subset preserves the crowd axioms; the displayed coordinate relations are affine. Over rings they say  $h_1 h_2 h_3 = I$ . The signed example retains  $h(1, 0, 0) * h(0, 1, 0) = \{h(1, 1, 1)\}$ , and hence the central cross-term, although  $h(1, 0, 0)^2$  is absent. This provides a constructive second Heisenberg formulation, with a representation theory still to specify.

## 11.7 Tensor categories and the qubit fusion object

**Definition 11.18** (The categorical phase system). Let  $\mathcal{C}_N(A) = \text{Rep}_{\mathbb{C}}(H_N(A))$ , with all linear intertwiners and internal tensor using the diagonal group action. Let  $\mathcal{C}_N(A)_{\lambda}$  be the full subcategory on which central  $u$  acts as  $\lambda(u)I$ . The unitary version chooses invariant Hermitian inner products and has the same intertwining spaces. Write  $\omega_A$  for the forgetful tensor functor. For groups  $H, K$  with specified central  $\mu_N$ , define  $\text{Rep}(H) \boxtimes_{\text{match}} \text{Rep}(K)$  as the full subcategory of  $\text{Rep}(H \times K)$  on which all  $(u, u^{-1})$  act trivially.

*Scope.* The matching condition is explicit. It is not an unspecified relative tensor product, and it is different from internal fusion.

---

**Provenance.** ids: D1011; proved in: definitions.md; tested by: f1\_check.py (examples); reviewed in: supporting comparison.

**Proposition 11.19** (Central sectors explain why the category matters). **SKETCH** For  $N \geq 1$  and finite abelian  $A$  of exponent dividing  $N$ ,  $\mathcal{C}_N(A)$  is a symmetric fusion category with a faithful tensor functor  $\omega_A$ , and

$$\mathcal{C}_N(A) = \bigoplus_{\lambda \in \widehat{\mu_N}} \mathcal{C}_N(A)_{\lambda}, \quad \mathcal{C}_N(A)_{\lambda} \otimes \mathcal{C}_N(A)_{\nu} \subseteq \mathcal{C}_N(A)_{\lambda\nu}.$$

Moreover, for finite groups  $H, K$  with specified central embeddings of  $\mu_N$ ,

$$\text{Rep}(H \boxtimes K) \simeq \text{Rep}(H) \boxtimes_{\text{match}} \text{Rep}(K).$$

On simple external tensors, the matching condition is equality of the two central characters. In particular a fixed nontrivial sector is not a tensor category by itself.

*Scope.* The unit of matching composition on this family is  $\text{Rep}(\mu_N)$ , with one simple per grade. It is not the unit for the unrestricted Deligne product.

---

**Provenance.** ids: F1-CAT; proved in: f1-comparisons.md, section 5; tested by: f1\_check.py (M6, M12; examples); reviewed in: supporting SKETCH.

*Proof.* Average an inner product over the finite group to obtain semisimplicity. Central character eigenspaces give the grading, and diagonal tensor multiplies eigencharacters. A representation factors through the central quotient precisely when its antidiagonal centre acts trivially; inflation and descent are inverse tensor functors. On an external simple the condition is  $\lambda(u)\nu(u)^{-1} = 1$ .  $\square$

**Definition 11.20** (Dihedral and quaternion qubit data). For  $e \in \{0, 1\}$  let  $G_e = \mathbb{F}_2^3$  with

$$(t, a, b)(u, c, d) = (t + u + ad + e(ac + bd), a + c, b + d).$$

Put

$$\sigma_e(t, a, b) = (-1)^t i^{e(a+b)} Z^b X^a, \quad X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

using representatives  $a, b \in \{0, 1\}$  in the exponent of  $i$ . Define  $\nu_2(\sigma_e) = |G_e|^{-1} \sum_g \text{Tr}(\sigma_e(g^2))$ .

*Scope.* The group law distinguishes the two cocycle choices. The quaternion matrices require fourth-root phases in this monomial realization.

---

**Provenance.** ids: D1012; proved in: definitions.md; tested by: f1\_check.py (M12); reviewed in: supporting comparison.

**Proposition 11.21** (Two qubit objects with the same fusion rules). **SKETCH**  $G_0 = D_8$  of order eight and  $G_1 = Q_8$ . Each has four one-dimensional characters and the single two-dimensional irreducible  $\sigma_e$ . Writing the four characters as  $1, \alpha, \beta, \alpha\beta$ ,

$$\sigma_e \otimes \sigma_e \simeq 1 \oplus \alpha \oplus \beta \oplus \alpha\beta, \quad \nu_2(\sigma_0) = 1, \quad \nu_2(\sigma_1) = -1.$$

Their sourced tensor-category identifications are

$$\text{Rep}(D_8) \simeq \text{TY}(\mathbb{F}_2^2, \chi_{\text{alt}}, 1/2), \quad \text{Rep}(Q_8) \simeq \text{TY}(\mathbb{F}_2^2, \chi_{\text{alt}}, -1/2).$$

*Scope.* The displayed tensor square uses one diagonal symmetry group. It does not replace the external tensor describing independent qubits.

---

**Provenance.** ids: F1-FUSION; proved in: f1-comparisons.md, section 6; Shimizu, p. 20; tested by: f1\_check.py (M12); reviewed in: supporting SKETCH.

The two-dimensional character is 2, −2 on the two central elements and zero elsewhere. Squaring it gives the sum of the four quotient characters. Six elements square to 1 in  $D_8$  and only two do in  $Q_8$ , giving the indicator signs. The categorical identifications and their distinction are in Shimizu’s *Frobenius–Schur indicators in Tambara–Yamagami categories*, Definition 3.1, Table 1 and page 20.

For two independent reference qubits the relevant group instead is  $H_2(C_2) \boxtimes H_2(C_2) = H_2(C_2 \times C_2)$ , of order 32. Its nontrivial-central-character irreducible has dimension four. Both this representation and internal fusion use the underlying Hilbert space  $\mathbb{C}^2 \otimes \mathbb{C}^2$ , but their symmetry actions differ. This is a concrete reason to retain the tensor category, its sectors and its fibre functor as part of the quantum object.

The fibre functor is essential to this particular formulation:

$$\text{End}_{\text{Rep}(H)}(\sigma) = \mathbb{C}, \quad \text{Hom}_{\text{Rep}(H)}(1, \sigma) = 0$$

for the nontrivial central character. The equivariant category alone does not give every state vector or every qubit observable. Those are recovered as vectors of  $\omega(\sigma) = \mathbb{C}^2$  and elements of  $\text{End}_{\mathbb{C}}(\omega(\sigma)) = M_2(\mathbb{C})$ . An alternative categorical state notion would have to be specified.

## 11.8 The independent Hall and categorical-oscillator route

**Definition 11.22** (Pointed-set Hall algebra and Fock realization). On the normal pointed-set category above, the rational Hall vector space has basis  $u_r$  indexed by rank. Its product counts pointed subsets  $T$  with the prescribed isomorphism classes of  $T$  and  $S/T$ . Put  $\mathcal{F}_{\text{alg}} = \mathbb{C}[x]$  with  $\langle x^m, x^n \rangle = \delta_{mn} n!$ , define  $a^\dagger f = xf$ ,  $af = df/dx$ , and let  $\mathcal{F}_{\text{bos}}$  be the Hilbert completion.

*Scope.* Hall coefficients and the Fock space are additive realizations of the pointed-set data. Creation is unbounded.

---

**Provenance.** ids: D1010; proved in: definitions.md; tested by: f1\_check.py (M8,M9); reviewed in: supporting comparison.

**Proposition 11.23** (A bosonic Heisenberg algebra from finite sets). **SKETCH** *The rational Hall algebra of finite normal pointed sets is  $\mathbb{Q}[x]$  under*

$$u_m u_n = \binom{m+n}{m} u_{m+n}, \quad u_n \longmapsto x^n / n!.$$

*On  $\mathcal{F}_{\text{alg}}$ , creation and annihilation are adjoints and  $[a, a^\dagger] = I$ . In the normalized basis  $x^n / \sqrt{n!}$  their coefficients are  $\sqrt{n+1}$  and  $\sqrt{n}$ .*

*Scope.* The polynomial Hall algebra alone is not the Weyl algebra; annihilation is additional operator structure. No finite-dimensional characteristic-zero CCR representation is asserted.

---

**Provenance.** ids: F1-HALL; proved in: f1-comparisons.md, section 4; tested by: f1\_check.py (M8,M9); reviewed in: supporting SKETCH.

The subset count proves the Hall formula. The operator calculation is  $d(xf)/dx - x df/dx = f$ . The source construction is [Szczesny's Hall algebra](#), section 3. [Baez–Dolan](#) give the factorial groupoid weights and the positive categorical relation

$$AA^\dagger \simeq A^\dagger A \oplus \text{id}.$$

[Morton–Vicary](#) realize the categorified Heisenberg algebra using spans of finite-set groupoids: add/remove histories split according to whether the removed point was just added. Their two-linearization retains the tower of  $\text{Rep}(S_n)$ , with the stated finiteness/completion caveats. The connection to  $\mathbb{F}_1$  is the equivalence between finite sets and the core of finite pointed modules. It is not a claim that these authors construct one universal  $\mathbb{F}_1$  scheme theory.

This supplies a categorical north star different from the finite Weyl system. Degree two of the representation-category tower has two simples in  $\text{Rep}(S_2)$ ; that is not automatically a qubit. With two colours, the one-particle complex space is a qubit and the two-boson sector is  $\text{Sym}^2(\mathbb{C}^2)$ , of dimension three. A one-mode cutoff to  $|0\rangle, |1\rangle$  instead has projected commutator  $\text{diag}(1, -1)$ , and is not closed under creation. The Hall operation uses wedge and particle-number addition; the finite configuration tensor uses smash and rank multiplication.

## 11.9 Toric, analytic and arithmetic-statistical formulations

**Definition 11.24** (A quantum torus and its selected central fibre). For a named  $\xi \in \mathbb{C}^\times$  put

$$\mathcal{T}_\xi = \mathbb{C}\langle U^{\pm 1}, V^{\pm 1} \rangle / (VU - \xi UV).$$

If  $\xi$  is a primitive  $N$ th root, put  $\mathcal{T}_\xi(1, 1) = \mathcal{T}_\xi / (U^N - 1, V^N - 1)$ .

*Scope.* The parameter  $\xi$ , its order  $N$  and a finite-field cardinality are different data. Central values are explicitly selected.

---

**Provenance.** ids: D1013; proved in: definitions.md; tested by: cyclic Weyl checks; reviewed in: supporting comparison.

**Proposition 11.25** (The qubit as a quantum-torus fibre). **SKETCH** For  $\xi$  primitive of order  $N$ ,  $\mathcal{T}_\xi(1, 1) \simeq M_N(\mathbb{C})$  via  $Ue_j = e_{j+1}$  and  $Ve_j = \xi^j e_j$  on  $\mathbb{Z}/N$ . For  $N = 2$  this is a qubit algebra; two independent fibres tensor to  $M_4(\mathbb{C})$ .

*Scope.* The universal quantum torus has not been identified with a finite matrix algebra. At  $\xi = 1$  it is commutative.

---

**Provenance.** ids: F1-TORUS; proved in: f1-comparisons.md, section 7; tested by: f1\_check.py (cyclic cases); reviewed in: supporting SKETCH.

The  $N^2$  clock/shift words are trace-orthogonal and span, which proves the selected-fibre statement. This is the twisted-lattice picture of [Neeb's rational quantum tori](#). Monoidal tori retain character lattices; a chosen skew pairing is additional symplectic data. [Manin's cyclotomic analytic geometry](#) supplies Habiro completions and evaluation or Taylor maps at roots of unity. It suggests an arithmetic interpolation problem for quantum fibres, but does not itself give a common Hilbert space or compatible Weil intertwiners. A deformation limit  $\xi \rightarrow 1$  is different from all the preceding  $q \rightarrow 1$  operations.

**Frobenius descent.** [Borger’s lambda geometry](#) treats lambda structure as descent data below  $\mathrm{Spec} \mathbb{Z}$ . For flat schemes it can be expressed as commuting Frobenius lifts. Monoid algebras have the lift  $m \mapsto m^p$ . Its naive coordinatewise application to the integral Heisenberg group fails already on the additive subgroup:  $(1+1)^p \neq 1^p + 1^p$  in  $\mathbb{Z}$ . This rejects that particular group-compatible lift, not all arithmetic descent. A quantum extension should specify whether arithmetic maps act by homomorphisms, noninvertible operators or correspondences. No qubit or Born rule is selected by lambda structure alone.

**A published quantum-statistical realization.** [Connes–Consani–Marcolli’s \*Fun with  \$\mathbb{F}\_1\$\*](#) , Proposition 6.1 and Theorem 6.2, constructs an  $\mathbb{F}_1$  model of the Bost–Connes endomotive, with the rational system recovered by scalar extension. It uses a cyclotomic tower and Frobenius/transfer correspondences. Its complex realization has the time evolution

$$\sigma_t(\mu_n) = n^{it} \mu_n, \quad \sigma_t(e(r)) = e(r).$$

In the standard representation on  $\ell^2(\mathbb{N}_{>0})$ ,  $He_n = (\log n)e_n$  and  $\mathrm{Tr}(e^{-\beta H}) = \zeta(\beta)$  for  $\beta > 1$ ; see [Bost–Connes \(1995\)](#). This is genuine quantum statistical dynamics, not a finite Stone–von Neumann representation. Choosing two energy levels provides an encoding but not a sector invariant under its full observable algebra.

[Lieber–Manin–Marcolli](#), published in 2022, develop lifts to Grothendieck rings, assembler categories, spectra and Nori motives. This is an independent literature-based reason to explore categorical targets, without asserting that these categories are the fusion categories of the finite Heisenberg construction.

**Tropical and motivic boundaries.** The band/Krasner framework retains combinatorial flags beyond a single coordinate apartment. Boolean semirings, Krasner hyperfields, signed partial fields and pointed monoids have different additive information. [Connes–Consani’s Arithmetic Site](#) uses a tropical semiring on a multiplicative-semigroup topos to realize arithmetic correspondences. It does not prescribe a local Weyl algebra or positive quantum states. Relative and homotopical geometries broaden the candidate bases and targets, but no general Heisenberg representation theorem for those bases is established in this sidequest. Finally, [Bejleri–Marcolli’s \*Quantum field theory over  \$\mathbb{F}\_1\$\*](#)  concerns torifications and varieties attached to Feynman integrals; its introduction explicitly distinguishes that problem from physical Lagrangians and Feynman rules over  $\mathbb{F}_1$ .

**An additional tensor analogy: fermionic occupancy.** [Lin–Wang–Wu–Yang \(2013\)](#) categorify the one-mode fermion algebra with  $c^2 = (c^\dagger)^2 = 0$  and  $cc^\dagger + c^\dagger c = 1$ . Occupancy subsets suggest a combinatorial exterior-power route; this is our proposed  $\mathbb{F}_1$  bridge, not their claim. [Hadzihasanovic–de Felice–Ng \(2018\)](#), section 2, give a precise monoidal category of local fermionic modes and maps of definite parity. Its basic object has underlying space  $\mathbb{C}^{1|1}$ ; tensor adds parity and the physical fermionic swap negates  $|11\rangle$ . The two occupancy labels therefore do not supply unrestricted qubit operations. Parity-preserving one-mode observables are diagonal, while a fixed-parity two-mode sector can encode a qubit. Signed phases are a candidate carrier for exchange signs over  $\mathbb{F}_1$ ; the cited complex categorical construction does not prove that descent. This is a further composition law to compare with the bosonic and distinguishable-system routes.

## 11.10 Precisely formulated outputs and open comparisons

The finite arithmetic candidate is the configuration-to-phase-system functor, with a named phase realization and a checked tensor law. The categorical candidate retains

$$(\mathcal{C}_N(A), \mathcal{C}_N(A)_\iota, \omega_A)$$

together with internal fusion and matching external composition. The pointed normal-map candidate already realizes an ordinary qubit over the bare combinatorial base. The Heisenberg crowd is a geometric object with correct ring fibres, while the Hall/groupoid route gives an independent categorical oscillator. Bost–Connes supplies the arithmetic quantum-statistical test case. These are several precise formulations, not evidence that all the analogies must coincide.

The next comparison problems are concrete: relate a chosen Tits reflection lift to its projective Fourier operator; extend the geometric Heisenberg model to nonreference cocycles; identify a genuine geometric home for the additive phase kernels; compare finite cyclotomic fibres with arithmetic transfer maps; and distinguish tensor products of distinguishable systems from bosonic particle-number sectors. A theory on arbitrary  $\mathbb{F}_1$  schemes still needs its input category, morphisms and realization functor specified. It is a research target, not a theorem or a well-formed universal conjecture supplied by the present literature.

Seven finite-kernel claims are **PROVED**; the extensions remain **SKETCH**. Thirteen exact example gates and their mutations check the finite computations. The cited literature and our deductions are distinguished throughout.

## 12 Operational sidequest: quantum mechanics in subsystem composition

**7 September 2026.** We now have a concrete positive candidate for the subsystem-first programme. Its endpoint assigns

$$\mathcal{A}(S) = \mathbb{C}[\text{Sym}(S)]$$

to a finite set of constituents. The algebras have normalized positive states, CP dynamics, Born probabilities and proper assembly inclusions. One constituent has only scalar observables; three constituents have an  $M_2(\mathbb{C})$  sector. Two disjoint three-constituent clusters can carry a Bell state. The construction is a quantum family, not a proposed single-particle wavefunction over a one-element coefficient field.

The arithmetic precursor is the positive type-A Hecke family. At finite field cardinality  $q$ , it is the invariant transition algebra of complete flags in a chosen configuration Lagrangian. Two overlapping subsystem embeddings determine  $q \geq 1$  through the intrinsic Born overlap  $q/(q+1)^2$ , even though their abstract algebra types are independent of  $q$ . The endpoint at one is an actual specialization of algebraic presentations.

The limitation is explicit: this is a *flag-context sector extracted from polarized Weyl quantum mechanics*. It is not yet a specialization of the entire Weyl algebra, its phase data and every change of polarization. The research report is `docs/sidequests/f1-operational.md`.

### 12.1 A positive arithmetic family

**Definition 12.1** (Hecke systems and their reference trace). Fix a real  $q > 0$ . Put  $H_0(q) = H_1(q) = \mathbb{C}$ . For  $n \geq 2$ , define the unital complex algebra  $H_n(q)$  by generators  $T_1, \dots, T_{n-1}$ , involution  $T_i^* = T_i$ , and relations

$$(T_i - q)(T_i + 1) = 0, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad T_i T_j = T_j T_i \quad (|i - j| > 1).$$

For  $w \in S_n$ , let  $T_w$  be a reduced-word product and let  $\ell(w)$  be its Coxeter length. Define the coefficient functional by  $\tau_n(T_w) = \delta_{w,1}$ .

*Scope.* Basis independence, positivity and the C\*-norm are proved below. The real parameter becomes a field cardinality only at the specified arithmetic fibres.

---

**Provenance.** ids: D1101; proved in: definitions.md; tested by: f1\_operational\_check.py (H1,H2); reviewed in: f1-operational-adjudication.md.

**Theorem 12.2** (Positive Hecke quantum algebras). **PROVED** For every  $q > 0$  and  $n \geq 0$ , the  $T_w$  form a basis of  $H_n(q)$ ,  $T_w^* = T_{w^{-1}}$ , and

$$\tau_n(T_u^* T_v) = \delta_{u,v} q^{\ell(u)}.$$

Consequently  $\tau_n$  is a faithful normalized positive trace, and left multiplication on  $L^2(H_n(q), \tau_n)$  gives a faithful finite C\*-algebra representation.

*Scope.* This covers all positive real parameters, including one. It does not assert positivity at arbitrary complex roots of unity.

---

**Provenance.** ids: F1-HCK-POS; proved in: f1-operational/hecke.md, section 1; tested by: H1,H2; reviewed in: f1-operational-adjudication.md.

*Proof.* On a basis  $b_w$ , the standard generator raises Coxeter length with coefficient one, and on a descent acts by  $qb_{s_i w} + (q-1)b_w$ . The two-string matrix is  $\begin{pmatrix} 0 & q \\ 1 & q-1 \end{pmatrix}$ . The rank-two strings verify the defining relations. Reduced words span and are independent by their action

on  $b_1$ . With  $\langle b_u, b_v \rangle = \delta_{u,v} q^{\ell(u)}$ , every generator is self-adjoint. This proves the Gram identity and  $\tau_n(x^*x) = \sum_w |x_w|^2 q^{\ell(w)}$ . The basis identity also gives traciality. The faithful regular \*-representation has closed finite-dimensional image.  $\square$

**Definition 12.3** (Finite-field flag context algebra). Fix a named finite field  $k = \mathbb{F}_Q$  and an  $n$ -dimensional  $k$ -space  $L$ . A complete flag is  $0 = U_0 < U_1 < \dots < U_n = L$  with  $\dim_k U_i = i$ . Put

$$\text{Cxt}(L) = \text{End}_{GL(L)} \ell^2(\text{Fl}(L)).$$

Let  $(A_w f)(F) = \sum_{\text{pos}(F, F')=w} f(F')$ , labelling relative positions by reduced gallery words in adjacency-product order. Use the normalized operator trace  $\text{tr}_{\text{Fl}}(a) = |\text{Fl}(L)|^{-1} \text{Tr}(a)$ . If  $L$  is a named configuration Lagrangian of a Weyl system, call this its complete-flag context algebra.

*Scope.* The context register is a separate Hilbert space from the Weyl configuration register. The Lagrangian is named data.

---

**Provenance.** ids: D1104; proved in: definitions.md; tested by: H9; reviewed in: f1-operational-adjudication.md.

**Theorem 12.4** (Arithmetic realization by invariant flag transitions). PROVED *For every finite field  $\mathbb{F}_Q$  and finite-dimensional  $L$  of dimension  $n$ , the adjacency map  $T_w \mapsto A_w$  is a trace-preserving \*-isomorphism*

$$H_n(Q) \simeq \text{Cxt}(L).$$

*Scope.* The operator convention is direct; a right-convolution convention would require the usual opposite/inversion adjustment.

---

**Provenance.** ids: F1-HCK-FLAG; proved in: f1-operational/hecke.md, section 2; Iwahori 1964; tested by: H9, fields of orders 2 and 3; reviewed in: f1-operational-adjudication.md.

This is the finite-field construction in [Iwahori \(1964\)](#), Lemma 3.1 and Theorems 3.2 and 4.1. Pair-orbits of flags are indexed by  $S_n$  and their kernels span the commutant. An adjacent panel contains  $Q + 1$  flags, so its nonidentity adjacency has square  $QI + (Q - 1)A_{s_i}$ . Transposition inverts relative position, and only the identity kernel has nonzero diagonal. These facts fix the trace.

**Definition 12.5** (Ordered assembly and coarse graining). For real  $q > 0$  and  $m, n \geq 0$ , embed  $S_m \times S_n$  as consecutive block permutations in  $S_{m+n}$ . Define

$$\iota_{m,n}(T_u \otimes T_v) = T_{u \times v}, \quad E_{m,n}(T_w) = \begin{cases} T_w, & w \in S_m \times S_n, \\ 0, & \text{otherwise,} \end{cases}$$

identifying the retained basis with  $H_m(q) \otimes H_n(q)$ .

*Scope.* These are ordered maps. No symmetric block swap is stipulated at  $q \neq 1$ .

---

**Provenance.** ids: D1102; proved in: definitions.md; tested by: H3; reviewed in: f1-operational-adjudication.md.

**Theorem 12.6** (Coherent positive assembly). PROVED *For  $q > 0$  and  $m, n \geq 0$ ,  $\iota_{m,n}$  is an injective unital trace-preserving \*-homomorphism. The map  $E_{m,n}$  is the unique faithful trace-preserving UCP conditional expectation onto its image. Both families satisfy ordered three-block coherence.*

*Scope.* Complete positivity includes all matrix amplifications. The finite checker samples this statement; the proof has no rank bound.

---

**Provenance.** ids: F1-HCK-TOWER; proved in: f1-operational/hecke.md, section 3; tested by: H3,H13, stated finite samples; reviewed in: f1-operational-adjudication.md.

*Proof.* The two sets of block generators commute and their basis images are distinct. The coefficient trace restricts to the product trace. Orthogonality of the basis makes  $E$  the trace-orthogonal projection. The trace-pairing characterization gives its bimodule property and positivity; repeating the projection argument in every matrix amplification gives complete positivity. The two three-block routes retain precisely  $S_l \times S_m \times S_n$ . The classical source for the expectation is Umegaki (1954).  $\square$

**Definition 12.7** (Operational states and processes of a Hecke level). Fix real  $q > 0$  and  $n \geq 0$ . A state is a positive functional  $\omega : H_n(q) \rightarrow \mathbb{C}$  with  $\omega(1) = 1$ , represented by  $h \geq 0$ ,  $\tau_n(h) = 1$ , through  $\omega(a) = \tau_n(ha)$ . An effect satisfies  $0 \leq e \leq 1$  and has probability  $\tau_n(he)$ . Deterministic Heisenberg channels are UCP; outcome maps are subunital CP and an instrument has a unital sum. The ordered product density is  $\iota_{m,n}(h \otimes k)$ .

A retained local Kraus list satisfies  $\sum_i K_i^* K_i = 1$  and gives

$$\Phi_K(a) = \sum_i K_i^* a K_i, \quad \mathcal{T}_K(h) = \sum_i K_i h K_i^*.$$

In a larger block, embed the operators before applying these formulas.

*Scope.* Equality of isolated CP maps is not assumed to survive assembly. Not every CP map is asserted to have this local inner form.

---

**Provenance.** ids: D1106; proved in: definitions.md; tested by: H3,H5,H8; reviewed in: f1-operational-adjudication.md.

**Theorem 12.8** (Quantum operational semantics of the positive family). PROVED For  $q > 0$ , the Hecke states have unique positive normalized densities, effects give Born probabilities, and the stated restriction and preparation maps are CP. Ordered product densities are normalized and associative. Retained local Kraus processes extend coherently through ordered block assembly.

*Scope.* This is a finite  $C^*$ -quantum sector. A tensor law on arbitrary isolated CP shadows has not been asserted.

---

**Provenance.** ids: F1-HCK-OPS; proved in: f1-operational/hecke.md, section 4; tested by: H3,H5,H8; reviewed in: f1-operational-adjudication.md.

The density/functional identification is blockwise matrix duality for a faithful trace. The equality

$$\tau_{m+n}(\iota(h \otimes k)a) = (\omega_h \otimes \omega_k)(E_{m,n}(a))$$

identifies product preparation with precomposition by the conditional expectation. Positivity and normalization give the Born interval  $[0, 1]$ .

## 12.2 The parameter lives in the subsystem diagram

**Definition 12.9** (Unlabelled overlap and marked trace). Let a finite  $C^*$ -algebra have a unique noncommutative simple summand  $M_2(\mathbb{C})$  and two specified unital subalgebras  $B_1, B_2 \simeq \mathbb{C}^2$ . When the two pairs of minimal projections compress to rank-one projections in that block, define

$$a(\mathcal{A}; B_1, B_2) = \min_{P \in B_1, Q \in B_2} \text{Tr}_2(PQ),$$

where the minimum runs over those compressed minimal projections and  $\text{Tr}_2$  is ordinary matrix trace. Separately, in  $H_2(q)$  define  $e_{\text{triv}} = (T_1 + 1)/(q + 1)$  and  $r_2 = \tau_2(e_{\text{triv}})$ .

*Scope.* The first invariant does not label the projections or use a chosen reference state. The second invariant explicitly marks the trivial projection.

**Provenance.** ids: D1103,D1105; proved in: definitions.md; tested by: H4; reviewed in: f1-operational-adjudication.md.

**Theorem 12.10** (Arithmetic memory in overlapping subsystems). PROVED For every  $q > 0$ ,

$$H_1(q) = \mathbb{C}, \quad H_2(q) \simeq \mathbb{C}^2, \quad H_3(q) \simeq \mathbb{C}^2 \oplus M_2(\mathbb{C}).$$

For the left and right  $H_2$  inclusions in  $H_3$ , the unlabelled overlap is

$$a(q) = \frac{q}{(q+1)^2}, \quad q = \frac{1 - 2a + \sqrt{1 - 4a}}{2a} \quad (q \geq 1).$$

It determines exactly  $\{q, q^{-1}\}$ , and uniquely determines  $q$  on  $q \geq 1$ . The marked trace instead satisfies  $r_2 = 1/(q+1)$  and determines  $q$  without that range restriction.

*Scope.* The algebra types alone do not determine the parameter. The two overlapping embeddings, or the separately marked trace, do.


**Provenance.** ids: F1-HCK-LOW,F1-HCK-Q; proved in: f1-operational/hecke.md, sections 5–6; tested by: H4,H5; reviewed in: f1-operational-adjudication.md.

In the matrix block one may calculate with

$$P_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad P_2 = \begin{pmatrix} a & \sqrt{a(1-a)} \\ \sqrt{a(1-a)} & 1-a \end{pmatrix}.$$

The four unlabelled overlaps are  $a, a, 1-a, 1-a$ , and  $a \leq 1/4$ . This matrix calculation proves a basis-independent statement about the abstract  $C^*$ -diagram. The reciprocal  $*$ -isomorphism is  $T_i \mapsto -qS_i$  into  $H_n(q^{-1})$ .

Overlapping quantum subsystems



The two embeddings retain the parameter even when the algebra types agree.

Finite fields:  $q = p^m$ ; endpoint:  $q = 1$




Figure 1: The abstract algebra types stay fixed, while the relative subsystem embeddings retain the positive parameter. The markers are exact rational probabilities; the intervening curve illustrates the proved formula.

The coefficient trace on  $H_3(q)$  is

$$\tau_3(a_+, a_-, a_{\text{std}}) = \frac{a_+ + q^3 a_-}{(q+1)(q^2 + q + 1)} + \frac{q}{q^2 + q + 1} \text{Tr}_2(a_{\text{std}}).$$

Thus a standard-block pure state with ordinary density  $P_2$  has full density  $h = P_2/\gamma$ ,  $\gamma = q/(q^2 + q + 1)$ . Let  $u_1 = 2e_1 - 1$ ,  $e_1 = (T_1 + 1)/(q + 1)$ . Its isolated action on the left

commutative  $H_2$  is identity, but the same retained unitary in  $H_3$  changes the return probability for effect  $P_2$  from 1 to

$$(1 - 2a(q))^2.$$

At one the full density is  $3P_2$  and the return probability is  $1/4$ . Preparing it by measuring  $P_2$  in the reference state has success probability  $1/3$ . This is a complete finite quantum experiment.

### 12.3 A functor on normal maps over the field with one element

**Definition 12.11** (The symmetric-group quantum net). For a finite set  $S$ , put  $G(S) = \text{Sym}(S)$  and  $\mathcal{A}(S) = \mathbb{C}[G(S)]$ , with

$$\left(\sum_g a_g g\right)^* = \sum_g \overline{a_g} g^{-1}, \quad \tau_S\left(\sum_g a_g g\right) = a_1.$$

An injection  $f : S \rightarrow T$  extends permutations by the identity off  $f(S)$ , giving  $i_f$ . Its reverse coefficient map  $E_f$  deletes terms outside that subgroup. Disjoint union gives the block inclusion  $\mu_{S,T}$ . States use  $h \geq 0$ ,  $\tau_S(h) = 1$ , and  $\omega_h(a) = \tau_S(ha)$ . Local Kraus lists satisfy  $\sum_i K_i^* K_i = 1$ ; their action in  $T$  retains the embedded list  $i_f(K_i)$ . Lists are identified under scalar unitary mixing after padding by zero operators.

*Scope.* The coefficient trace is normalized. It is different from the unnormalized categorical trace used later for a general fusion category.

---

**Provenance.** ids: D1125; proved in: definitions.md; tested by: H5,H7,H8; reviewed in: f1-operational-adjudication.md.

**Theorem 12.12** (A collective quantum endpoint). PROVED *The symmetric-group assignment on finite injections, with its coefficient expectations, states and retained local Kraus processes, is a finite quantum net. It has coherent proper disjoint assembly and CP restriction/preparation. Its three-constituent standard sector has the full density  $3P_2$  and the Born witness  $1 \rightarrow 1/4$ .*

*Scope.* The net supplies a sufficient coherent class of local processes, not a canonical extension of every isolated CP map.

---

**Provenance.** ids: F1-OP-SYM; proved in: f1-operational/cp.md, CPOP-5; tested by: H5,H7,H8; reviewed in: f1-operational-adjudication.md.

**Definition 12.13** (Partial-injection realization). A normal map of finite pointed  $\mathbb{F}_1$  modules preserves the basepoint and has at most one preimage of each non-basepoint. On nonzero sets it is a partial injection  $f : S \rightarrow T$ , a bijection  $D \rightarrow E$  between subsets. Define

$$\mathcal{R}(f) = i_{E \subset T} \circ \mathcal{A}(f : D \simeq E) \circ E_{D \subset S}.$$

The target has finite tracial  $C^*$ -algebras, unital trace-preserving CP maps and trace adjoints as daggers. Wedge of pointed modules becomes disjoint union of the nonzero sets.

*Scope.* This is the normal-map convention; arbitrary pointed maps are not included.

---

**Provenance.** ids: D1010,D1127; proved in: definitions.md; Szczesny 1204.5395; tested by: H12; reviewed in: f1-operational-adjudication.md.

**Theorem 12.14** (Operational realization of normal F1 maps). PROVED *On finite sets and partial injections  $f : D \subseteq S \rightarrow E \subseteq T$ , put  $\mathcal{A}(S) = \mathbb{C}[\text{Sym}(S)]$  with coefficient trace, and  $\mathcal{R}(f) = i_E \mathcal{A}(f : D \simeq E) E_D$ , using subgroup inclusion and coefficient expectation. This functor is dagger and lax symmetric monoidal, with structure maps*

$$\mu_{S,T} : \mathcal{A}(S) \otimes \mathcal{A}(T) \hookrightarrow \mathcal{A}(S \sqcup T).$$

*An empty partial map has image  $a \mapsto \tau_S(a)1_T$ .*

*Scope.* The laxators need not be isomorphisms. The source zero map is a reference reset, not zero CP; this is a nonadditive functor.

---

**Provenance.** ids: F1-OP-PMAP; proved in: f1-operational/contexts.md, PMAP-1; tested by: H12; reviewed in: f1-operational-adjudication.md.

*Proof.* A group-basis permutation survives precisely when its support is in  $D$ ; then it is relabelled and extended by identity. Two successive survival conditions say that the support lies in the domain of the composed partial map. This proves functoriality. Inclusion and coefficient expectation are trace adjoints, proving the dagger statement. The same support calculation proves naturality under disjoint union; associativity and symmetry follow from the canonical set bijections.  $\square$

The source wedge operation is therefore collective assembly. Classical alternatives can separately use C\*-algebra direct sums, with their central probabilities. They are not the same operation, and neither should be silently identified with unrestricted coherent Hilbert direct sum.

## 12.4 An exact process quotient and a sharp context bound

**Definition 12.15** (Kraus Gram and contextual equivalence). Fix a nonempty finite  $S$ ,  $n = |S|$ ,  $H = \text{Sym}(S)$ , and normalized local lists. Write  $K_i = \sum_h a_i(h)h$  and put

$$J_K(h, k) = \sum_i a_i(h) \overline{a_i(k)}.$$

Two lists are contextually equivalent when their Heisenberg maps  $\Phi_{K,f}(a) = \sum_i i_f(K_i)^* a i_f(K_i)$  agree after every finite injection into an ambient set. Stable scalar-unitary equivalence pads lists by zeros and applies a unitary matrix to the Kraus index. An admissible Gram is  $J \geq 0$  with

$$\sum_h J(hx, h) = \delta_{x,1} \quad (x \in H).$$

*Scope.* The Gram is process data, not just the isolated CP action. It has trace one by the constraint at  $x = 1$ .

---

**Provenance.** ids: D1126; proved in: definitions.md; tested by: H8, H11; reviewed in: f1-operational-adjudication.md.

**Theorem 12.16** (Finite contextual recovery of a quantum process). PROVED *Fix a finite set  $S$  of size  $n \geq 1$  and finite lists in  $\mathbb{C}[\text{Sym}(S)]$  normalized by  $\sum_i K_i^* K_i = 1$ , extended through permutation inclusions along finite injections  $S \rightarrow T$ . Kraus Gram equality, stable scalar-unitary equivalence and equality of all ambient CP actions are equivalent. One ambient set of size  $2n - 1$  suffices. For  $n \geq 2$  this universal size bound is sharp, with explicit positive-state Born witnesses.*

*Scope.* The theorem concerns retained local Kraus processes of the symmetric-group net. It is not a claim about every structural partial map or every quantum channel in every fusion category.

---

**Provenance.** ids: F1-OP-KCF; proved in: f1-operational/contexts.md, KCF-1; tested by: H11 at q=1; reviewed in: f1-operational-adjudication.md.

*Proof.* In an ambient set of size  $2n - 1$ , choose an involution  $g$  exchanging  $n - 1$  local points with the outside points and fixing the remaining one. Then  $H \cap gHg^{-1} = 1$ , so all words  $h^{-1}gk$  are distinct. The expansion

$$\Phi_K(g) = \sum_{h,k} J_K(k, h) h^{-1} g k$$

recovers every Gram coefficient. Conversely equal Grams give equal maps in all contexts. Equality of finite Grams gives exactly stable unitary mixing by extending an isometry between coefficient-vector spans.

For sharpness let  $p_+, p_-$  be the local trivial and sign central projections and  $p_0 = 1 - p_+ - p_-$ . The normalized lists  $(p_+ + p_-, p_0)$  and  $(p_+ - p_-, p_0)$  differ only in cross terms between the trivial and sign sectors. In every smaller possible ambient set, each support intersection has at least two points. Its transposition forces those cross terms to vanish. At size  $2n - 1$  they do not vanish. For  $e = (1 + g)/2$ , set  $\Delta = \Phi_+(e) - \Phi_-(e)$ . Both images are effects and have equal trace, so  $h = 1 + \Delta$  is a positive normalized density, with Born difference  $\tau(\Delta^2) > 0$ .  $\square$

Thus  $n - 1$  ancillary constituents can be necessary. At  $n = 2$  all inner channels are the isolated identity on  $\mathbb{C}^2$ , while their Grams form the disk

$$J = \begin{pmatrix} r & ib \\ -ib & 1 - r \end{pmatrix}, \quad b^2 \leq r(1 - r).$$

A three-constituent context distinguishes them. The corresponding positive-Hecke context-bound extension remains **SKETCH**: its minimal-double-coset argument is written down and exact examples at  $q = 2$ ,  $n = 2, 3$  pass, but only the symmetric-group theorem was admitted.

---

**Provenance.** ids: F1-OP-KCF-Q; proved in: f1-operational/contexts.md, separated extension; tested by: H11 at  $q=2$ ; reviewed in: supporting SKETCH.

## 12.5 The partial-flag category is explicitly constructible

**Definition 12.17** (Partial-flag corners and their specialization). Fix real  $q > 0$ . A composition  $\alpha = (\alpha_1, \dots, \alpha_r)$  of  $n$  has positive parts summing to  $n$ ; include the empty composition of zero. Let  $W_\alpha$  preserve its consecutive blocks and put

$$P_\alpha(q) = \sum_{w \in W_\alpha} q^{\ell(w)}, \quad e_\alpha = P_\alpha(q)^{-1} \sum_{w \in W_\alpha} T_w.$$

Define  $\Gamma_q(\alpha, \beta) = e_\beta H_n(q) e_\alpha$  for equal totals, and zero Hom spaces for different totals. Composition is multiplication, identity is  $e_\alpha$ , adjoint is  $*$ , and tensor concatenates compositions through  $\iota$ . Let  $x = (1)$ . Finite formal sums and splitting self-adjoint idempotents give the additive completion; only its nonzero objects are used as normalized quantum systems. The generic coefficient ring is  $\mathbb{Z}[t, t^{-1}, \{P_\alpha(t)^{-1}\}_\alpha]$ , evaluated at  $t = q$ .

For  $L = \mathbb{F}_Q^n$ , partial flags have successive quotient dimensions  $\alpha$ . If  $r_\alpha$  forgets steps, define

$$J_\alpha e_E = P_\alpha(Q)^{-1/2} \sum_{F: r_\alpha(F)=E} e_F.$$

The normalized corner trace is  $\tau_\alpha = P_\alpha(q) \tau_n$ . A corner Kraus family  $v_j \in e_\beta H_n e_\alpha$  is normalized when  $\sum_j v_j^* v_j = e_\alpha$  and acts by  $a \mapsto \sum_j v_j^* a v_j$ .

*Scope.* The localization is explicit. There is no rigidity or cross-degree preparation in the amplitude category itself.

---

**Provenance.** ids: D1141; proved in: definitions.md; tested by: H9, H13; reviewed in: supporting SKETCH.

**Proposition 12.18** (A positive monoidal category of flag contexts). **SKETCH** For  $q > 0$ ,  $\Gamma_q$  is a graded monoidal  $C^*$ -category. At a finite field cardinality  $Q$ , its corner morphisms are exactly the  $GL(L)$  intertwiners between partial-flag Hilbert spaces, through  $a \mapsto J_\beta^* a J_\alpha$ . Refinement is represented by the corresponding typed projection. At one,

$$\text{End}_{\Gamma_1}(x^{\otimes n}) = \mathbb{C}[S_n],$$

the  $C$ -linear finite-bijection skeleton, with partial-flag retracts added.

*Scope.* An operational envelope supplies normalized states and additional preparations. The corner trace rescaling must be retained.

---

**Provenance.** ids: F1-OP-FLAGCAT; proved in: f1-operational/flag-category.md, sections 1–3; tested by: H9,H13; reviewed in: supporting SKETCH.

The calculation is elementary:  $T_s \sum_{W_\alpha} T_w = q \sum_{W_\alpha} T_w$  implies  $e_\alpha^2 = e_\alpha = e_\alpha^*$ . Block length additivity gives  $\iota(e_\alpha \otimes e_\beta) = e_{\alpha\|\beta}$ . Refinement gives  $e_\beta e_\alpha = e_\beta$ , and  $J_\alpha J_\alpha^* = e_\alpha$ . A density transported by a corner Kraus family uses the factor  $P_\alpha/P_\beta$  between the two normalized traces.

## 12.6 A concrete coupling back to Weyl quantum mechanics

**Definition 12.19** (Weyl constraint flags and controlled tests). Fix  $k = \mathbb{F}_Q$  of characteristic  $p$ , an  $n$ -dimensional named configuration Lagrangian  $L$ , and nontrivial additive  $\psi : k \rightarrow U(1)$ . On  $\mathbb{C}[L]$  use  $X(u)e_y = e_{y+u}$  and  $Z(\chi)e_y = \chi(y)e_y$ , with additive characters  $\chi : L \rightarrow U(1)$ ; the  $k$ -linear dual is identified through  $\psi$ . For a  $k$ -subspace  $U$ , let  $U^{\text{ann}}$  be the characters trivial on it and define

$$P_U^X = |U|^{-1} \sum_{u \in U} X(u), \quad P_U^Z = |U^{\text{ann}}|^{-1} \sum_{\chi \in U^{\text{ann}}} Z(\chi).$$

On  $K_{\text{ctx}}(L) = \mathbb{C}[\text{Fl}(L)]$ , declare the flag basis orthonormal and put, on its tensor with the Weyl register,

$$C_i = \sum_F |F\rangle\langle F| \otimes P_{U_i(F)}^Z.$$

The Levi action is  $R(g)e_y = e_{gy}$ , together with permutation of flags.

*Scope.* The field, Lagrangian, origin and phase datum are retained. The context and Weyl Hilbert spaces are different registers.

---

**Provenance.** ids: D1142; proved in: definitions.md; tested by: character identities in the proof; reviewed in: supporting SKETCH.

**Proposition 12.20** (Flags as quantum constraint chains). **SKETCH** Fix a finite field  $k = \mathbb{F}_Q$ , an  $n$ -dimensional configuration Lagrangian  $L$  and a nontrivial additive character  $\psi : k \rightarrow U(1)$ . Its complete flags inject  $GL(L)$ -equivariantly into each of the two nested Weyl projector chains, with commutation within each chain. The  $C_i$  are commuting orthogonal projections on the joint context/Weyl register, invariant under the diagonal Levi action.

*Scope.* This is an extraction and coupling of sectors. It is not an equivalence with full Weyl quantum mechanics or a proof of independence from the polarization.

---

**Provenance.** ids: F1-HCK-CONTEXT,F1-OP-COUPLE; proved in: f1-operational/flag-category.md, section 4; tested by: explicit character derivation; reviewed in: supporting SKETCH.

Character orthogonality gives  $P_U^Z = \sum_{y \in U} |y\rangle\langle y|$ ;  $P_U^X$  averages on cosets. The two projector-chain ranks are  $Q^i$  and  $Q^{n-i}$ . These formulas make the coupling explicit, while the dimensions  $Q^n$  and  $[n]_Q!$  prevent an unmentioned identification of the two registers.

**Definition 12.21** (Coordinate-apartment comparison). For a finite field  $\mathbb{F}_Q$ ,  $L = \mathbb{F}_Q^n$  and an ordered basis of  $L$ , embed the coordinate complete flags by the isometry  $J_{\text{ap}} : \mathbb{C}[S_n] \rightarrow \mathbb{C}[\text{Fl}(L)]$ , using the gallery/regular-action convention. Put  $\Omega_Q(a) = J_{\text{ap}}^* a J_{\text{ap}}$  on the context commutant.

*Scope.* This is an operational comparison at a fixed field, not the definition of substituting one into parameter-dependent coefficients.

---

**Provenance.** ids: D1144; proved in: definitions.md; tested by: H9 apartment probes; reviewed in: supporting SKETCH.

**Proposition 12.22** (A direct CP map from arithmetic contexts to the endpoint). SKETCH For every prime power  $Q$ , apartment compression is basis-independent after permutation labelling and gives a trace-preserving UCP map  $H_n(Q) \rightarrow \mathbb{C}[S_n]$ ,  $T_w \mapsto w$ . It commutes with ordered block inclusions and coefficient expectations.

*Scope.* For  $Q > 1$  it is not multiplicative and is not automatically natural for every partial-flag corner compression.

---

**Provenance.** ids: F1-OP-APART; proved in: f1-operational/flag-category.md, section 7; tested by: H9 at Q=2,3; reviewed in: supporting SKETCH.

Each relative position has one neighbour within the coordinate apartment. Changing its basis conjugates by  $GL(L)$ , which fixes the commutant. These facts prove the range and independence; isometric compression proves complete positivity. The limits are visible in  $\Omega_Q(T_s^2) = (Q-1)s + Q1 \neq 1 = \Omega_Q(T_s)^2$ . A coarse flag compression gives scalar  $Q$  before the apartment map but scalar one after it. Compatible refinement-process comparison is therefore a concrete remaining problem.

## 12.7 The broader categorical operational framework

**Definition 12.23** (Nonzero categorical subsystem data). Let  $\mathcal{C}$  be a unitary fusion category with its positive spherical trace and a tensor-closed replete family of nonzero objects containing the unit. Define

$$A_X = \text{End}_{\mathcal{C}}(X), \quad d_X = \text{Tr}_X(1_X) > 0, \quad \tau_X = \text{Tr}_X / d_X,$$

and  $j_{X,Y}(a \otimes b) = a \otimes b \in A_{X \otimes Y}$ . Retain associators and the trace-preserving assembly expectations, including their iterated comparison maps. A state has density  $\rho \geq 0$ ,  $\text{Tr}_X(\rho) = 1$ , representing  $\omega(a) = \text{Tr}_X(\rho a)$ ; effects lie between zero and one. Assembly restriction is the expectation and product-subalgebra preparation embeds its density by  $j$ .

*Scope.* Zero objects are excluded from normalized state systems. This definition uses unnormalized categorical trace, unlike the coefficient trace convention of the group net.

---

**Provenance.** ids: D1121,D1122; proved in: definitions.md; tested by: finite expectation examples H3; reviewed in: f1-operational-adjudication.md.

**Theorem 12.24** (Positive assembly without a strong Hilbert fibre). PROVED Let  $\mathcal{C}$  be a unitary fusion category with its positive spherical trace and a tensor-closed replete family of nonzero objects containing the unit. Then  $A_X = \text{End}_{\mathcal{C}}(X)$  are finite  $C^*$ -algebras with faithful positive traces. Assembly is injective and trace-multiplicative and has a unique UCP trace-preserving expectation. States, restriction, preparation and Born probabilities are well defined and coherent under nested assembly.

*Scope.* No strong monoidal Hilbert-space fibre functor is required. The inclusion may be proper.

---

**Provenance.** ids: F1-OP-ALG; proved in: f1-operational/cp.md, CPOP-1-2; tested by: H3, examples only; reviewed in: f1-operational-adjudication.md.

In simple-sector coordinates,  $A_X = \bigoplus_a M_{m_a}(\mathbb{C})$  and  $\text{Tr}_X = \sum_a d_a \text{Tr}_{m_a}$ , with  $d_a > 0$ . This proves faithfulness and identifies the quantum-trace weights. The expectation is orthogonal projection for the trace inner product; matrix amplification proves its complete positivity. The relevant internal operator-algebra analogue is developed by [Jones–Penneys](#); their internal algebra objects are not being identified with this net without a comparison.

**Definition 12.25** (Retained categorical Kraus and dilation processes). Let  $\mathcal{C}$  be a unitary fusion category with positive spherical trace, and fix a tensor-closed replete family of nonzero objects containing the unit. For family objects  $X, Y$ , a Kraus process is a finite list  $K_i : X \rightarrow Y$  with  $\sum_i K_i^* K_i = 1_X$ , modulo stable scalar-unitary mixing. It acts by  $\Phi_K(a) = \sum_i K_i^* a K_i$  and  $\mathcal{T}_K(\rho) = \sum_i K_i \rho K_i^*$ ; its retained right-context operators are  $K_i \otimes 1_Z$ . Composition uses  $L_j K_i$ , and tensor uses  $K_i \otimes L_j$ , with the category associators.

A charge-carrying dilation is a named nonzero environment  $E$  and isometry  $v : X \rightarrow Y \otimes E$ , with  $\Phi_v(a) = v^*(a \otimes 1_E)v$ . Sequential composition retains its ordered environment. Parallel composition requires named unitary interchange, supplied for example by a unitary braiding or a specified environment half-braiding.

*Scope.* Not every isolated CP map is asserted to be represented by these local lists, and arbitrary environments are not silently interchangeable in a merely monoidal category.

---

**Provenance.** ids: D1123,D1124; proved in: definitions.md; tested by: H8; finite examples; reviewed in: f1-operational-adjudication.md.

**Theorem 12.26** (Coherent quantum processes from retained operators). **PROVED** *In a unitary fusion category with positive spherical trace, fix a tensor-closed replete family of nonzero objects containing the unit. On that family, normalized Kraus lists compose and tensor modulo stable scalar-unitary mixing. Their shadows are UCP and their density maps preserve categorical trace in every retained context. The stated dilations compose sequentially and in parallel when the named interchange data are provided.*

*Scope.* This does not quotient by equality of isolated shadows. That coarser identification fails in the examples above.

---

**Provenance.** ids: F1-OP-PROC; proved in: f1-operational/cp.md, CPOP-3; tested by: H8; reviewed in: f1-operational-adjudication.md.

The essential identity is  $\sum_{i,j} K_i^* L_j^* L_j K_i = 1_X$ ; cyclicity gives trace preservation. Scalar unitary mixing cancels the Kraus indices in every context. These are the data that an isolated CP map can fail to retain.

For comparison, use the standard unitary Fibonacci data

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad F = \begin{pmatrix} \varphi^{-1} & \varphi^{-1/2} \\ \varphi^{-1/2} & -\varphi^{-1} \end{pmatrix}, \quad R = \text{diag}(e^{-4\pi i/5}, e^{3\pi i/5}).$$

On the two-anyon algebra  $\mathbb{C}^2$ ,  $\text{Ad}_R$  is identity. In the charge- $\tau$  matrix block of three anyons it is nontrivial: the  $|+\rangle$  return probability becomes  $(5 - \sqrt{5})/8$ . The braid  $FRF$  has  $|0\rangle$  return probability  $(3 - \sqrt{5})/2$ . These sourced comparisons remain **SKETCH** and do not identify Fibonacci with the arithmetic endpoint.

---

**Provenance.** ids: F1-OP-FIB; proved in: f1-operational/cp.md, CPOP-4; Ahmadi-Kissinger 2211.03855; tested by: independent exact critic recomputation; reviewed in: supporting SKETCH.

## 12.8 Entanglement and alternative branches

**Definition 12.27** (A Bell density on two collective sectors). In  $\mathbb{C}[S_3]$ , let  $s = (12)$ ,  $t = (23)$ ,  $z = (2 - (123) - (132))/3$ , and put  $Z_* = zs$ ,  $X_{\text{num}} = z(s + 2t)$ ,  $Y_{\text{anti}} = z(st - ts)$ . For the block inclusion into  $\mathbb{C}[S_6]$ , define

$$P_{\text{Bell}} = \frac{1}{4}\mu \left( z \otimes z + Z_* \otimes Z_* + \frac{X_{\text{num}} \otimes X_{\text{num}} + Y_{\text{anti}} \otimes Y_{\text{anti}}}{3} \right), \quad h_{\text{Bell}} = 9P_{\text{Bell}}.$$

On the standard sector the normalized observables are  $Z_*$  and  $X_* = X_{\text{num}}/\sqrt{3}$ .

*Scope.* The coefficient trace on the full six-constituent algebra fixes the density factor nine.

---

**Provenance.** ids: D1143; proved in: definitions.md; tested by: H14; reviewed in: supporting SKETCH.

**Proposition 12.28** (A Bell state of two disjoint collective qubits). **SKETCH** *The stated  $P_{\text{Bell}}$  is a positive projection,  $\tau_6(h_{\text{Bell}}) = 1$ , and both subgroup marginals are  $(3/2)z$ , the maximally mixed standard-sector qubit states. Its  $XX$  and  $ZZ$  correlations are one and its crossed correlations zero, giving the ordinary CHSH value  $2\sqrt{2}$  with the corresponding qubit measurements.*

*Scope.* This uses two disjoint size-three regions and their local matrix blocks; it does not assert that all composite observables factor locally.

---

**Provenance.** ids: F1-OP-BELL; proved in: f1-operational/flag-category.md, section 6; tested by: H14; reviewed in: supporting SKETCH.

In the standard block  $X_{\text{num}} = \sqrt{3}X$  and  $Y_{\text{anti}} = i\sqrt{3}Y$ , so the formula is the usual Bell projector. Each standard block contributes trace weight  $1/3$ ; their product gives  $1/9$ . The conditional expectation extends the Bell functional to the full composite while preserving these local correlations.

The type-C flag route remains distinct: at one its signed permutation groups have a rank-two  $D_8$  qubit sector, but  $B_m \times B_n$  is not parabolic in  $B_{m+n}$ . A separate assembly theorem is needed. The rank-two Temperley–Lieb route also needs extra input: its coefficient is  $\delta^{-2} = q/(q+1)^2$ , with  $\delta = \sqrt{q} + 1/\sqrt{q}$ . At one it gives the  $SU(2)$  spin tensor-power centralizer tower after the quotient and a Jones/Markov trace. The faithful flag trace cannot descend through the nonzero quotient: at level two the flag weights are  $(1/2, 1/2)$ , whereas spin singlet/triplet weights are  $(1/4, 3/4)$ . Full rigidity additionally requires duality conventions. The Fibonacci quotient is a different root-of-unity construction with its corresponding positive star/trace structure.

The substantive outcome is a quantum subsystem family with intrinsic arithmetic memory, an explicit positive endpoint and a sharp description of its process data. The remaining central task is comparison with the full Weyl system and all its refinement/polarization processes. Those missing comparisons are stated as such; they are not hidden inside the word “canonical.”

## 13 A continuous Hecke limit and its complete category

The positive Hecke parameter gives a genuine continuous path from arithmetic flag algebras to the symmetric-group fibre at  $q = 1$ . The construction below keeps one parameter throughout: tensor products of section algebras are balanced over  $C(I)$ , whereas tensor products inside one fibre are ordinary spatial tensor products. Completing the parabolic objects then produces all finite self-adjoint matrix corners, including off-diagonal maps between summands that happen to have the same total degree.

### 13.1 Continuous Hecke sections and parabolic corners

**Definition 13.1** (Positive Hecke algebra and normalized regular frame). Fix  $q > 0$ . Let  $H_0(q) = H_1(q) = \mathbb{C}$ , and for  $n \geq 2$  let  $H_n(q)$  be generated by self-adjoint  $T_1, \dots, T_{n-1}$  subject to

$$(T_i - q)(T_i + 1) = 0, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad T_i T_j = T_j T_i \quad (|i - j| > 1).$$

For  $w \in S_n$ , write  $T_w$  for a reduced-word product and  $\ell(w)$  for its Coxeter length. The coefficient trace is  $\tau_{n,q}(T_w) = \delta_{w,1}$ . Put

$$b_w(q) = q^{-\ell(w)/2} T_w(q).$$

The vectors  $b_w(q)$  are identified with the standard basis  $\delta_w$  of  $K_n = \ell^2(S_n)$ , and  $B_w(q)$  denotes left multiplication by  $b_w(q)$  on this fixed Hilbert space.

*Scope.* The embedding in the fixed regular Hilbert space is named `data`. It does not make the Hecke generators independent of  $q$ , and no positivity is asserted for a non-real or non-positive parameter.

---

**Provenance.** ids: D1101, D1201; proved in: `theory/sidequests/f1-limit/continuous-corners.md`; tested by: `theory/checks/f1_limit_check.py` (L1); `theory/checks/f1_operational_check.py` (H1,H2); reviewed in: `theory/verdicts/f1-limit-adjudication.md`.

**Definition 13.2** (Continuous positive Hecke section algebra). For a nonempty compact interval  $I \subset (0, \infty)$ , define

$$H_n^{\text{cts}}(I) = \left\{ q \mapsto \sum_{w \in S_n} f_w(q) B_w(q) : f_w \in C(I) \right\} \subset C(I, \text{End}(K_n)).$$

Operations are pointwise and the norm is the supremum operator norm. Its  $C(I)$ -valued trace is

$$\tau_n^{\text{cts}}(x)(q) = \tau_{n,q}(x(q)).$$

*Scope.* This is a continuous section algebra over a compact positive interval. It is not a generic algebra over a Laurent-polynomial coefficient ring.

---

**Provenance.** ids: D1202; proved in: `theory/sidequests/f1-limit/continuous-corners.md`; tested by: `finite-fibre probes L1,H1,H2`; closedness is proved analytically; reviewed in: `theory/verdicts/f1-limit-adjudication.md`.

**Theorem 13.3** (Continuous regular Hecke algebras). **PROVED** For every nonempty compact  $I \subset (0, \infty)$  and every  $n \geq 0$ ,  $H_n^{\text{cts}}(I)$  is a unital  $C^*$ -subalgebra of  $C(I, \text{End}(K_n))$ . Evaluation at  $q_0 \in I$  is a surjective  $*$ -homomorphism onto the faithful regular copy of  $H_n(q_0)$ , and  $\tau_n^{\text{cts}}$  is continuous, tracial, positive and faithful.

*Scope.* The assertion is uniform in the rank and on every compact positive interval. The finite checks test fibres; they do not prove norm-closedness or continuity.

---

**Provenance.** ids: F1-LIM-CTS; proved in: theory/sidequests/f1-limit/continuous-corners.md; tested by: theory/checks/f1\_limit\_check.py (L1); theory/checks/f1\_operational\_check.py (H1,H2); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* On a length pair  $w, s_i w$  with  $\ell(s_i w) = \ell(w) + 1$ , left multiplication by  $T_i$  has matrix

$$\begin{pmatrix} 0 & \sqrt{q} \\ \sqrt{q} & q-1 \end{pmatrix}$$

in the normalized basis. Hence all  $B_w(q)$  vary continuously. The Hecke relations show that their  $C(I)$ -span is a unital  $*$ -algebra. If a sequence of sections converges uniformly, applying it to  $\delta_1$  recovers every coefficient function uniformly; the limit therefore remains in the displayed span. This proves closedness. Constant coefficient functions prove surjectivity of every evaluation. Finally,

$$\tau_{n,q}(x^*x) = \sum_w |c_w|^2 q^{\ell(w)} \quad \text{for } x = \sum_w c_w T_w,$$

so fibre traces, and consequently the section trace, are faithful and positive.  $\square$

**Definition 13.4** (Continuous parabolic corner category). For a composition  $\alpha = (\alpha_1, \dots, \alpha_r)$  of  $n$ , let  $W_\alpha \leq S_n$  preserve its consecutive blocks and set

$$P_\alpha(q) = \sum_{w \in W_\alpha} q^{\ell(w)}, \quad e_\alpha(q) = P_\alpha(q)^{-1} \sum_{w \in W_\alpha} T_w(q).$$

The objects of  $\Gamma_I^{\text{cts}}$  are compositions, including the empty composition of zero, and

$$\text{Hom}(\alpha, \beta) = \begin{cases} e_\beta H_n^{\text{cts}}(I) e_\alpha, & |\alpha| = |\beta| = n, \\ 0, & |\alpha| \neq |\beta|. \end{cases}$$

Composition, identity and dagger are multiplication,  $e_\alpha$  and matrix adjoint. Ordered tensor concatenates compositions. On section algebras its domain is the balanced product

$$H_m^{\text{cts}}(I) \otimes_{C(I)} H_n^{\text{cts}}(I),$$

the section algebra of the pointwise spatial fibre products, and the map is the contiguous-block inclusion. On the endomorphism corner the normalized trace is

$$\tau_{\alpha,q} = P_\alpha(q) \tau_{n,q}.$$

The germ category at one identifies two morphism sections when they agree on some smaller interval  $[1 - \varepsilon, 1 + \varepsilon]$ . A germ is positive when it has a positive representative on one such interval.

*Scope.* The ordinary tensor product over  $\mathbb{C}$  is not the section tensor: for nonconstant  $f$ , the nonzero element  $f \otimes 1 - 1 \otimes f$  is killed by restriction to the diagonal. Only evaluation at one is canonical on a germ, and no  $C^*$ -norm is assigned to germ Hom spaces.

---

**Provenance.** ids: D1141, D1203, D1204; proved in: theory/sidequests/f1-limit/continuous-corners.md; tested by: theory/checks/f1\_operational\_check.py (H3,H13); theory/checks/f1\_limit\_check.py (L4,L14); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 13.5** (Continuous corners and their endpoint germ). **PROVED** *For every compact positive interval  $I$ , the preceding corners form an ordered monoidal  $C^*$ -category. Their normalized traces are faithful and multiplicative under ordered tensor; the corner inclusions and their trace-preserving expectations are continuous and obey the three-block tower laws. Fibre evaluation is full. Germs at one form an ordered monoidal  $*$ -category with a full canonical endpoint evaluation to the  $q = 1$  corner category.*

*Scope.* No block exchange is supplied for  $q \neq 1$ . A nonzero germ may vanish at the endpoint, which is why endpoint norm is not used as a  $C^*$ -norm.

---

**Provenance.** ids: F1-LIM-CORNER, F1-LIM-GERM; proved in: theory/sidequests/f1-limit/continuous-corners.md; tested by: H3,H13; L4,L14 on finite fibres and representatives; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Writing  $x_\alpha = \sum_{w \in W_\alpha} T_w$ , the Hecke multiplication rule gives  $T_i x_\alpha = q x_\alpha$  for every simple reflection in the parabolic. Thus  $x_\alpha^2 = P_\alpha x_\alpha$  and  $e_\alpha$  is a self-adjoint projection. Block length additivity sends  $B_u \otimes B_v$  to the distinct section  $B_{u \times v}$ , proving injectivity of the balanced tensor map. Moreover  $P_{\alpha \parallel \beta} = P_\alpha P_\beta$  and  $\iota(e_\alpha \otimes e_\beta) = e_{\alpha \parallel \beta}$ . The ambient coefficient expectation is bimodular, so it restricts to each corner, fixes the included product algebra and preserves the normalized trace. All operations commute with restriction and evaluation. To lift a fibre corner morphism, lift it with constant normalized-basis coefficients and compress by the source and target projection sections. This proves fullness, both on an interval and after passing to endpoint germs.  $\square$

**Definition 13.6** (Arithmetic and endpoint evaluations). An arithmetic fibre evaluates the continuous construction at the positive real number  $q = Q = p^r$  and may then apply the complete- or partial-flag realization over  $\mathbb{F}_Q$ . Endpoint specialization evaluates at  $q = 1$ . Coordinate-apartment compression at a fixed arithmetic fibre is a separate UCP map and is not an evaluation map.

*Scope.* No sequence of prime powers tends to one. The construction retains the chosen flag context and does not specialize the full Weyl system or all polarizations.

---

**Provenance.** ids: D1216; proved in: theory/sidequests/f1-limit/continuous-corners.md; tested by: theory/checks/f1\_operational\_check.py (H9,H13); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Proposition 13.7** (Arithmetic partial-flag realization). PROVED *For every prime power  $Q$  and every composition  $\alpha$  of  $n$ , the fibre corner  $e_\alpha H_n(Q) e_\alpha$  is the full  $GL_n(\mathbb{F}_Q)$ -commutant on partial flags of type  $\alpha$ , and  $P_\alpha(Q) \tau_{n,Q}$  is normalized physical operator trace. These identifications preserve corner morphisms, composition and dagger.*

*Scope.* This is an evaluation followed by a finite-field realization, rather than a limit over finite fields. Apartment compression remains nonmultiplicative when  $Q > 1$ .

---

**Provenance.** ids: F1-LIM-ARITH; proved in: theory/sidequests/f1-limit/continuous-corners.md; tested by: theory/checks/f1\_operational\_check.py (H9,H13); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* A partial flag of type  $\alpha$  has  $P_\alpha(Q)$  complete refinements. The normalized fibre-sum isometry  $J_\alpha$  has range projection  $e_\alpha$ . Consequently  $F \mapsto J_\beta F J_\alpha^*$  identifies every partial-flag intertwiner with  $e_\beta H_n(Q) e_\alpha$ , with inverse  $x \mapsto J_\beta^* x J_\alpha$ . Ordinary trace on the range of  $J_\alpha$  and the flag-count ratio give the asserted normalized trace.  $\square$

## 13.2 All finite self-adjoint summands

**Definition 13.8** (Finite graded self-adjoint completion). For a nonempty compact positive interval  $I$ , put  $A_n(I) = H_n^{\text{cts}}(I)$ . An object  $X$  is a finite-support family  $(r_n, p_{X,n})_{n \geq 0}$ , where  $r_n \geq 0$  and  $p_{X,n} \in M_{r_n}(A_n(I))$  is a self-adjoint projection. For  $Y = (s_n, p_{Y,n})$ , define

$$\text{Hom}(X, Y) = \bigoplus_n p_{Y,n} M_{s_n, r_n}(A_n(I)) p_{X,n}.$$

Composition and dagger are degreewise matrix multiplication and adjoint, the identity is  $(p_{X,n})_n$ , and the norm is the maximum of the finitely many supremum  $C^*$ -norms. Direct sum uses block-diagonal projections within each degree, so its endomorphism algebra retains the off-diagonal rectangular corners between equal-degree summands. Different degrees have zero Homs. The fibre version at  $q$  is denoted  $\mathcal{U}_q^\dagger$ .

*Scope.* This is the finite additive dagger Karoubi completion. It does not assume duals, rigidity, a spherical trace or finitely many simple objects.

---

**Provenance.** ids: D1261; proved in: theory/sidequests/f1-limit/karoubi-trace.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Remark.* An additive object is a coherent multiplicity resource, not a classical alternative. For example, the direct sum of two tensor units has endomorphism algebra  $M_2(\mathbb{C})$ , including off-diagonal coherences. A two-outcome classical wire instead has the commutative algebra  $\mathbb{C}^2$ . Adding finite sums therefore does not change the original generator tower  $\text{End}(x^{\otimes n}) = H_n(q)$ , and it does not give the bare one-constituent object qubit observables.

**Definition 13.9** (Balanced ordered tensor and finite trace). For objects  $X, Y$ , the degree- $k$  multiplicity of  $X \otimes Y$  is

$$t_k = \sum_{m+n=k} r_m s_n.$$

Its projection is block diagonal, indexed by  $(m, n)$  and lexicographic matrix indices, with block

$$\iota_{m,n}^{(r_m, s_n)}(p_{X,m} \otimes p_{Y,n}).$$

Morphism tensor is the analogous block-diagonal amplification. The full endomorphism corner in  $M_{t_k}(A_k(I))$  includes off-diagonal blocks between different pairs with the same sum. The unit is the degree-zero object  $(1, 1)$ , and associators are the permutation matrices identifying the two triple indexings. Section tensor is balanced over  $C(I)$ .

For  $a \in \text{End}(X)$  put

$$\theta_X(a) = \sum_n (\text{Tr}_{r_n} \otimes \tau_n)(a_n), \quad w_X = \theta_X(1_X), \quad \tau_X = \theta_X/w_X.$$

Here matrix trace is unnormalized. Operational systems are nonzero objects; a density satisfies  $\rho \geq 0$  and  $\tau_X(\rho) = 1$ .

*Scope.* Associators reorder bookkeeping indices but never exchange ordered Hecke blocks. The trace is a finite graded matrix/coefficient trace and is not claimed to arise from categorical duality.

---

**Provenance.** ids: D1262, D1263; proved in: theory/sidequests/f1-limit/karoubi-trace.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1, K2); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 13.10** (Full continuous completion and its trace). **PROVED** *For every nonempty compact positive interval, the finite projection objects above form an additive ordered monoidal  $C^*$ -category with every self-adjoint idempotent split. The additive self-adjoint completion of the parabolic corner category is unitarily strong-monoidally equivalent to it. The trace  $\theta$  is continuous, positive, faithful and cyclic across rectangular Homs; on a connected interval every nonzero object has  $w_X(q) > 0$ , and*

$$w_{X \otimes Y} = w_X w_Y, \quad \tau_{X \otimes Y}(j(a \otimes b)) = \tau_X(a) \tau_Y(b).$$

*At a fibre this is the self-adjoint model of the full finite Hecke composition category.*

*Scope.* The dagger statement concerns self-adjoint projection presentations. An algebraic idempotent is merely isomorphic, not canonically unitarily equal, to its support projection.

---

**Provenance.** ids: F1-LIM-KAR, F1-LIM-KTRACE; proved in: theory/sidequests/f1-limit/karoubi-trace.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1,K2); uniform assertions by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Each Hom is a closed rectangular corner of a finite direct sum of matrix C\*-algebras. Block diagonal sums give biproducts, and a self-adjoint idempotent is itself a new projection object. The normalized block basis proves injectivity of the balanced tensor, while permutation matrices on triple indices give the associator and pentagon. The complete parabolic composition has projection 1 in every degree, so its sums and retracts generate all objects.

For a rectangular  $a$ , direct expansion gives

$$(\mathrm{Tr} \otimes \tau_n)(a^*a) = \sum_{i,j} \tau_n(a_{ij}^* a_{ij}),$$

which proves faithfulness. Cyclicity follows entrywise. A continuous projection has constant ordinary rank on a connected interval, so a nonzero object cannot disappear at one fibre and  $w_X$  stays positive. Finally the matrix trace and coefficient trace both multiply on each  $(m, n)$  block; summing over all pairs proves both displayed tensor identities.  $\square$

**Definition 13.11** (Traced assembly of completed objects). For nonzero  $X, Y$ , let  $B_X = \mathrm{End}(X)$  and let

$$j_{X,Y} : B_X \otimes B_Y \longrightarrow B_{X \otimes Y}$$

be morphism tensor; on intervals its domain is  $C(I)$ -balanced. Define  $E_{X,Y} : B_{X \otimes Y} \rightarrow B_X \otimes B_Y$  by

$$(\tau_X \otimes \tau_Y)(b^* E_{X,Y}(a)) = \tau_{X \otimes Y}(j_{X,Y}(b)^* a)$$

for all  $b$ . Assembly is the process from the separated word  $[X][Y]$  to the collective atom  $[X \otimes Y]$  represented in Heisenberg orientation by  $E_{X,Y}$ ; split has the reverse type and is represented by  $j_{X,Y}$ .

*Scope.* The separated algebra and the collective endomorphism algebra are different. The trace is the finite graded trace just defined, not a pivotal trace.

---

**Provenance.** ids: D1264; proved in: theory/sidequests/f1-limit/karoubi-trace.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1,K5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 13.12** (Continuous traced expectations). **PROVED** For nonzero completed objects on a connected compact positive interval,  $j_{X,Y}$  is an injective unital trace-preserving \*-map and  $E_{X,Y}$  is the unique normalized-trace-preserving UCP retraction. These expectations act continuously on sections, satisfy  $Ej = \mathrm{id}$  and the ordered three-object tower laws, while  $jE$  is generally a proper conditional expectation. On parabolic objects they recover the earlier corner maps.

*Scope.* No rigidity or fusion hypothesis is used. Properness means that collective same-degree off-diagonal observables can be erased by  $jE$ .

---

**Provenance.** ids: F1-LIM-KEXPECT; proved in: theory/sidequests/f1-limit/karoubi-trace.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1,K5); complete positivity and continuity by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* In one fibre, orthogonal projection onto  $j(B_X \otimes B_Y)$  for the faithful trace inner product fixes the subalgebra and is bimodular. If the image of a positive element had a negative spectral projection in the subalgebra, the trace-pairing identity would be simultaneously negative and nonnegative. Applying the same argument to matrix amplifications proves complete positivity. A local continuous basis turns the trace equations into  $G(q)c(q) = t(q)$  with  $G(q) > 0$ ; hence  $c(q) = G(q)^{-1}t(q)$  is continuous. Uniqueness proves agreement on overlaps and the three-object tower law.  $\square$

## 14 Typed operations, classical histories and collective context

The amplitude category has morphisms only within a fixed total degree. An operational theory needs more: preparation, discard, measurement and channels must be arrows between observable algebras. We build those arrows in Heisenberg orientation, retain their Kraus data, and keep two kinds of composition separate. Juxtaposing registers produces a spatial tensor algebra; assembling them produces one larger collective corner whose observable algebra usually contains more operators.

### 14.1 Registers and primitive processes

**Definition 14.1** (Quantum atoms, classical wires and register words). For every nonempty composition  $\alpha$ , let  $[\alpha]$  be a quantum atom with observable algebra

$$A_\alpha(q) = e_\alpha(q)H_{|\alpha|}(q)e_\alpha(q), \quad \tau_{\alpha,q} = P_\alpha(q)\tau_{|\alpha|,q}.$$

A register word is a finite ordered word of quantum atoms and classical wires. The algebra and reference trace of the word are the ordered spatial tensor products of the factor algebras and traces; the empty word has algebra  $\mathbb{C}$ . For a nonempty finite set  $O$ , the wire  $\underline{O}$  has algebra  $\mathbb{C}^O$  and uniform trace

$$\tau_O(f) = |O|^{-1} \sum_{o \in O} f(o).$$

The separated word  $[\alpha][\beta]$  and collective atom  $[\alpha \parallel \beta]$  are different objects.

*Scope.* The empty outcome set is excluded from deterministic instruments. A classical wire has a commutative algebra; a coherent additive direct sum in the completed amplitude category instead has matrix off-diagonal observables.

---

**Provenance.** ids: D1205; proved in: theory/sidequests/f1-limit/operational-category.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1–W3,W5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 14.2** (Finite Heisenberg UCP category). Objects are finite-dimensional unital  $C^*$ -algebras. An arrow  $A \rightarrow B$  is a unital completely positive map  $\Phi : B \rightarrow A$ . If  $A \rightarrow B \rightarrow C$  are represented by  $\Phi : B \rightarrow A$  and  $\Psi : C \rightarrow B$ , their composite is  $\Phi \circ \Psi : C \rightarrow A$ ; tensor is the spatial tensor of maps. For faithful reference traces, the density dual is the unique map satisfying

$$\tau_B(\Phi_*(\rho)a) = \tau_A(\rho\Phi(a)).$$

*Scope.* This target records operational UCP maps and their density duals. It does not assert that every such map possesses a preferred Hecke-corner Kraus label.

---

**Provenance.** ids: D1206; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: finite UCP examples in theory/checks/f1\_operational\_check.py; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 14.3** (Preparations, measurements and retained instruments). For a register word  $X$ , a preparation is a density  $h \geq 0$ ,  $\tau_X(h) = 1$ , with Heisenberg functional  $a \mapsto \tau_X(ha)$ . Discard is  $\lambda \mapsto \lambda 1_X$ . A POVM with outcome set  $O$  is a family  $e_o \geq 0$ ,  $\sum_o e_o = 1_X$ , and realizes

$$\mathbb{C}^O \longrightarrow A_X, \quad f \longmapsto \sum_o f(o)e_o.$$

An effect  $0 \leq e \leq 1$  is the two-outcome POVM  $(e, 1 - e)$ .

For equal-total compositions  $\alpha, \beta$ , a corner instrument is a finite family

$$K_{o,i} \in e_\beta H_n(q) e_\alpha, \quad \sum_{o,i} K_{o,i}^* K_{o,i} = e_\alpha,$$

with type

$$K : [\alpha] \longrightarrow [\beta] \underline{O}$$

and Heisenberg realization

$$\Phi_K((a_o)_o) = \sum_{o,i} K_{o,i}^* a_o K_{o,i}.$$

At one fibre, two labels are equal when the normalized-basis coefficient Gram matrices agree separately in each outcome. Equivalently, after zero padding they differ by a scalar-unitary mixing of the Kraus index within each outcome.

*Scope.* The quantum output of an instrument is retained. A POVM alone specifies effects but no postmeasurement state. Kraus indices may be mixed within an outcome, never across different classical outcomes.

---

**Provenance.** ids: D1207; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: theory/checks/f1\_operational\_check.py (H8); theory/checks/f1\_limit\_wiring\_check.py (W3,W4); reviewed in: theory/verdicts/f1-limit-adjudication.md.

The branch density for outcome  $o$  is not the raw matrix sum when source and target corner traces have different normalizations. It is

$$T_o(\rho) = \frac{P_\alpha(q)}{P_\beta(q)} \sum_i K_{o,i} \rho K_{o,i}^*.$$

Relative to  $\tau_\beta \otimes \tau_O$ , the  $o$ -block of the full recorded density is  $|O|T_o(\rho)$ . Cyclicity proves that its trace is precisely the branch Born weight and that the sum of all branch masses is one.

**Proposition 14.4** (UCP realization of primitive operations). **PROVED** *For every  $q > 0$ , every preparation, discard, finite POVM and normalized corner instrument just defined is a typed UCP process. Its Schrödinger dual is positive and trace preserving, and the branch density has the displayed corner and classical cardinality factors.*

*Scope.* This is a sufficient retained process class. It makes no completeness claim for arbitrary CP maps between two corner algebras.

---

**Provenance.** ids: F1-LIM-OP; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: theory/checks/f1\_operational\_check.py (H8); theory/checks/f1\_limit\_wiring\_check.py (W3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Preparations are positive normalized functionals and discards are unital  $*$ -maps. At matrix level a positive element of  $M_r(\mathbb{C}^O)$  is a family of positive scalar matrices, so tensoring each with  $e_o$  and summing proves complete positivity of the POVM map. Each instrument summand has the form  $a \mapsto K^* a K$ ; completeness makes their sum unital. The density formula follows from cyclicity:

$$\tau_\beta(T_o(\rho)a) = \tau_\alpha\left(\rho \sum_i K_{o,i}^* a K_{o,i}\right).$$

□

## 14.2 Assembly is a retraction, not an identification

**Definition 14.5** (Assembly and split). For parabolic atoms, let

$$\text{asm}_{\alpha,\beta} : [\alpha][\beta] \longrightarrow [\alpha \parallel \beta]$$

be represented in Heisenberg orientation by the trace-preserving conditional expectation

$$E_{\alpha,\beta} : A_{\alpha \parallel \beta} \longrightarrow A_{\alpha} \otimes A_{\beta}.$$

Let  $\text{spl}_{\alpha,\beta}$  have the reverse type and be represented by the corner inclusion  $j_{\alpha,\beta} : A_{\alpha} \otimes A_{\beta} \rightarrow A_{\alpha \parallel \beta}$ . The defining relations include the three-block tower relations and

$$\text{spl} \circ \text{asm} = \text{id}_{[\alpha][\beta]}.$$

The reverse composite is the collective coarse graining  $jE$ .

*Scope.* The map  $jE$  is generally proper and is not set equal to the collective identity. No assembly retraction is promoted to an isomorphism.

---

**Provenance.** ids: D1208; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: theory/checks/f1\_limit\_check.py (L14); theory/checks/f1\_operational\_check.py (H3,H13); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 14.6** (Coherent collective assembly). **PROVED** *Assembly and split are associative UCP processes,  $Ej = \text{id}$ , and  $jE$  is the conditional expectation onto the separated product subalgebra. Preparing densities  $h, k$  on separated registers and then assembling gives the collective density  $j(h \otimes k)$ .*

*Scope.* The conclusion uses contiguous ordered blocks and the normalized corner traces. It supplies neither a block exchange nor equality of separated and collective observables.

---

**Provenance.** ids: F1-LIM-ASM; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: L14; H3,H13; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The Hecke block inclusion is a trace-preserving \*-monomorphism and the coefficient expectation fixes its range, giving  $Ej = \text{id}$ . The two three-block inclusions and expectations agree on the standard basis, so the processes are associative. Trace duality gives  $\tau(j(h \otimes k)a) = (\tau_{\alpha} \otimes \tau_{\beta})((h \otimes k)E(a))$ , proving the density statement.  $\square$

**Definition 14.7** (Retained collective context and parallel lists). For  $K : [\alpha] \rightarrow [\beta]\underline{Q}$  and a right context  $\gamma$ , retain

$$K \triangleleft \gamma : [\alpha \parallel \gamma] \longrightarrow [\beta \parallel \gamma]\underline{Q}, \quad (K \triangleleft \gamma)_{o,i} = \iota(K_{o,i} \otimes e_{\gamma}).$$

For a second list  $L : [\alpha'] \rightarrow [\beta']\underline{P}$ , the direct ordered collective parallel instrument has type

$$K \boxtimes_c L : [\alpha \parallel \alpha'] \longrightarrow [\beta \parallel \beta']\underline{Q \times P}$$

and list  $(\iota(K_{o,i} \otimes L_{p,j}))_{(o,p),(i,j)}$ .

*Scope.* These lists act on full collective corner algebras. They are not inferred from the isolated CP shadows and preserve the displayed left-to-right order.

---

**Provenance.** ids: D1209; proved in: theory/sidequests/f1-limit/operational-category.md; tested by: theory/checks/f1\_operational\_check.py (H5,H8); reviewed in: theory/verdicts/f1-limit-adjudication.md.

### 14.3 Classical history wires

**Definition 14.8** (Classical products, relabeling and routing). Fix a singleton  $*$ . A bijection  $r : O \rightarrow P$  gives  $\text{rel}_r : \underline{O} \rightarrow \underline{P}$ , represented by  $r^* : C^P \rightarrow C^O$ ,  $r^*f = f \circ r$ . Classical product and unit are the  $*$ -isomorphisms

$$\text{mul}_{O,P} : \underline{O} \underline{P} \longrightarrow \underline{O \times P}, \quad \text{unit}_c : * \longrightarrow \mathbf{1},$$

induced by  $\delta_{(o,p)} \mapsto \delta_o \otimes \delta_p$  and  $\mathbb{C}^{\{*\}} \cong \mathbb{C}$ ; their inverses are included. Cartesian associators and singleton unitors are the corresponding relabelings.

For every object word  $X$ , the routing isomorphism

$$\text{route}_{O,X} : \underline{O}, X \longrightarrow X \underline{O}$$

is represented by the spatial flip  $A_X \otimes C^O \rightarrow C^O \otimes A_X$ . It is natural, involutive after exchanging source and target, and satisfies the usual nesting and unit relations.

For instruments  $K : [\alpha] \rightarrow [\beta] \underline{O}$  and  $L : [\beta] \rightarrow [\gamma] \underline{P}$ , define

$$\begin{aligned} \text{Seq}(L, K) &= (\text{id} \otimes \text{mul}_{O,P})(\text{id} \otimes \text{route}_{P,\underline{O}})(L \otimes \text{id}_{\underline{O}})K, \\ \text{Seq}(L, K) : [\alpha] &\rightarrow [\gamma] \underline{O \times P}. \end{aligned}$$

Its branch list is  $(L_{p,j}K_{o,i})_{(o,p),(j,i)}$ . For  $K' : [\alpha'] \rightarrow [\beta'] \underline{P'}$ , define

$$\begin{aligned} \text{Par}(K, K') &= (\text{id} \otimes \text{mul}_{O,P'})(\text{id}_{[\beta]} \otimes \text{route}_{O,[\beta']} \otimes \text{id}_{\underline{P'}})(K \otimes K'), \\ \text{Par}(K, K') : [\alpha][\alpha'] &\rightarrow [\beta][\beta'] \underline{O \times P'}. \end{aligned}$$

Sequential associativity uses  $((o,p),r) \leftrightarrow (o,(p,r))$ ; sequential–parallel interchange uses

$$((o,p),(o',p')) \longleftrightarrow ((o,o'),(p,p')).$$

One-outcome instruments use the singleton unitor. The single Kraus list  $(e_\alpha)$ , followed by that unitor, is the identity process on  $[\alpha]$ ; its Heisenberg map is  $a \mapsto e_\alpha a e_\alpha = a$ .

*Scope.* Only a classical wire moves through a quantum output. These maps do not provide a quantum block commutator at generic  $q$ .

---

**Provenance.** ids: D1218; proved in: theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1–W5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 14.9** (Presented operational categories and evaluation). At a fibre, take the strict ordered monoidal category generated by the register objects, preparations, discards, POVMs, retained instruments, assembly/split, context/parallel lists and classical wiring above. Equality is the congruence generated by typed planar isomorphism, category and monoidal axioms, literal state and POVM equality, outcome-wise coefficient-Gram equality, the routed history relations, context identities and assembly tower relations. It is not equality of one isolated CP shadow.

On an interval, labels are continuous and two Kraus labels are equal when their coefficient Grams agree at every parameter. At the endpoint, circuit germs are equal when the presentation relation, and in particular Gram equality, holds on a common smaller interval. No continuous choice of a mixing unitary is required. Realization sends objects to their algebras and generators to the stated UCP maps. Evaluation substitutes the parameter in every label; only endpoint evaluation is canonical for an arbitrary germ.

*Scope.* The realization may be nonfaithful. Local lifting of one circuit will not imply that every extra equation true only at the endpoint also lifts.

---

**Provenance.** ids: D1210, D1211; proved in: theory/sidequests/f1-limit/operational-category.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1-W5); H8,L14; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 14.10** (Sound typed operational category). **PROVED** *For every positive fibre, compact positive interval and endpoint germ, the preceding generators and relations define an ordered monoidal category with a monoidal UCP realization. Routed sequential and parallel instruments are normalized, history associativity and interchange are typed by the displayed bijections, and pointwise or eventual Gram equality is a congruence.*

*Scope.* No classical retyping is hidden in planar isotopy. No quantum exchange is asserted away from one, and no arbitrary CP-map completeness is claimed.

---

**Provenance.** ids: F1-LIM-OP; proved in: theory/sidequests/f1-limit/operational-category.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1-W5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Every wiring box is a trace-preserving  $*$ -isomorphism represented by a finite permutation matrix. Sequential completeness follows from

$$\sum_{o,p,i,j} K_{o,i}^* L_{p,j}^* L_{p,j} K_{o,i} = 1,$$

and parallel completeness factors into the two original sums. The two associative histories have the same branch operator and differ only by their Cartesian bracketing. Interchange follows from multiplicativity of the ordered block inclusion and the stated history bijection. At each fibre, equal coefficient Grams are equivalent to outcome-wise scalar-unitary mixing; tensoring those finite unitaries proves congruence pointwise, without choosing them continuously.  $\square$

#### 14.4 A finite collective context detects retained Kraus data

**Definition 14.11** (Local contextual recovery map). For the complete  $n$ -constituent object, expand a retained list in the normalized basis and put

$$J_K(k, h) = \sum_i a_i(k) \overline{a_i(h)}.$$

Embed the local regular operator explicitly into rank  $2n - 1$  by

$$\tilde{B}_h(q) = \iota_{n,n-1}(B_h(q) \otimes 1).$$

Choose an involution  $d \in S_{2n-1}$  fixing one local point and pairing the other  $n - 1$  local points with the complement. The recovery map is

$$J \mapsto \sum_{h,k} J(k, h) \tilde{B}_h(q)^* B_d(q) \tilde{B}_k(q).$$

The endpoint minor uses the distinct rows indexed by  $h^{-1}dk$ .

*Scope.* The neighborhood on which the minor stays invertible may depend on  $n$ . No sharp all-positive-parameter context bound is included.

---

**Provenance.** ids: D1215; proved in: theory/sidequests/f1-limit/lifting.md; tested by: theory/checks/f1\_limit\_check.py (L6); theory/checks/f1\_operational\_check.py (H11 at the endpoint); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 14.12** (Local contextual recovery of the Kraus Gram). PROVED *For every fixed  $n \geq 1$ , some interval about one has the following property: one retained context of total rank  $2n - 1$  recovers the full normalized-basis Kraus Gram. On that interval, equality in all retained contexts is equivalent to pointwise Gram equality, and at each fibre to stable scalar-unitary mixing.*

*Scope.* The scalar unitary supplied by the finite Gram lemma is fibrewise. A continuous family of such unitaries is neither obtained nor required for equality in the repaired core category.

---

**Provenance.** ids: F1-LIM-CTX; proved in: theory/sidequests/f1-limit/lifting.md; tested by: L6; H11 at q=1; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* At one the intersection of the local subgroup with its  $d$ -conjugate is trivial, so the  $(n!)^2$  permutations  $h^{-1}dk$  are distinct. The selected coefficient minor is therefore a permutation matrix. Its entries vary continuously, so its determinant remains nonzero near one. The displayed map is injective there. Equality of its values forces equality of Grams; conversely every retained channel is linear in the same Gram coefficients.  $\square$

## 14.5 Operations on the full additive completion

**Definition 14.13** (Controlled operations on completed objects). Let  $X = (r_n, p_{X,n})$  be a nonzero finite-support family of self-adjoint matrix projections and put

$$B_X = \bigoplus_n p_{X,n} M_{r_n}(H_n^{\text{cts}}) p_{X,n},$$

$$\theta_X(a) = \sum_n (\text{Tr}_{r_n} \otimes \tau_n)(a_n), \quad w_X = \theta_X(1_X), \quad \tau_X = \theta_X/w_X.$$

Quantum atoms are such objects; separated words use spatial tensor algebras and product traces, while classical wires retain the uniform trace above. Include every normalized preparation, discard and finite POVM. An instrument  $X \rightarrow YQ$  is a finite family

$$K_{o,i} \in \text{Hom}(X, Y), \quad \sum_{o,i} K_{o,i}^* K_{o,i} = 1_X,$$

realized by  $(a_o)_o \mapsto \sum_{o,i} K_{o,i}^* a_o K_{o,i}$ . Retain  $K_{o,i} \otimes 1_Z$ , ordered parallel lists, traced assembly/split and finite typed circuits.

The presentation uses the explicit sound classical wiring above. At a fibre, Kraus equality is coefficient-Gram equality, equivalently stable scalar-unitary mixing within each outcome. On an interval it is pointwise coefficient-Gram equality, and for germs it is such equality on a common smaller interval. Scalar-unitary cancellation proves congruence in each fibre; no continuous choice of mixing unitary is required. Any finite classical control is a typed blockwise family of the admitted labels and contributes no additional arbitrary CP map. The resulting fibre, interval and germ categories retain the realization and evaluation conventions above; their full circuit realizations are not stipulated to be faithful.

*Scope.* The construction assumes neither rigidity nor fusion and includes no arbitrary CP-map data. Fibrewise stable mixing does not assert the existence of a continuous environmental unitary.

---

**Provenance.** ids: D1266; proved in: theory/sidequests/f1-limit/karoubi-lifting.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1–K6, primitive examples); theory/checks/f1\_limit\_wiring\_check.py (W1–W5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 14.14** (Operational completion of all finite projection objects). **PROVED** *With the displayed classical-wiring presentation and its UCP relations, every nonzero object in the full finite completion carries normalized states, effects, POVMs, retained instruments, traced assembly and finite protocols. These generate an ordered UCP circuit category with continuous and germ evaluations. Restriction to parabolic atoms recovers the preceding operational formulas.*

*Scope.* This statement uses the wiring and equality convention just stated. It asserts neither all-CP completeness, rigidity, fusion, faithful circuit realization nor preservation of every additional endpoint equation.

---

**Provenance.** ids: F1-LIM-KOP; proved in: theory/sidequests/f1-limit/karoubi-lifting.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K1–K6, primitive examples); W1–W5; categorical typing by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Preparations, discards, POVMs and Kraus lists give UCP maps exactly as in the parabolic case. The completed trace supplies the correct density dual, and the unique traced expectations supply assembly and split. Classical wiring types every history relation and is parameter independent. The free typed category may therefore be quotiented by the proved congruence, and its generator realization descends to a monoidal UCP functor. The restriction to rank-one parabolic projections identifies all formulas term by term.  $\square$

## 15 Local lifting, Born limits and a three-constituent experiment

The continuous family becomes operational only if concrete endpoint data can be continued to a neighborhood of one. The continuation need not be global or canonical. It is enough to lift the finitely many objects and labels in a chosen circuit, normalize them by positive functional calculus, and intersect their finitely many neighborhoods. This distinction is decisive: lifting an individual circuit does not lift every equation that happens to hold only in the symmetric endpoint fibre.

### 15.1 Local lifts of objects and labels

**Definition 15.1** (Raw local corner lift). Fix  $q_0 > 0$  and  $a_0 \in e_\beta(q_0)H_n(q_0)e_\alpha(q_0)$ . Expand

$$a_0 = \sum_{w \in S_n} c_w b_w(q_0).$$

On a compact positive interval containing  $q_0$ , its raw lift is

$$a_{\text{raw}}(q) = e_\beta(q) \left( \sum_w c_w b_w(q) \right) e_\alpha(q).$$

For register words, raw lifts are finite sums in the  $C(I)$ -balanced tensor product of such sections.

*Scope.* This is local existence data and is not a preferred trivialization of the continuous family. Compression is part of the formula and keeps the source and target corners typed.

---

**Provenance.** ids: D1213; proved in: theory/sidequests/f1-limit/lifting.md; tested by: analytic coefficient-lift proof; no finite gate claims generality; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 15.2** (Normalized local operational lifts). For raw Kraus lifts  $\widehat{K}_r(q)$ , set

$$S(q) = \sum_r \widehat{K}_r(q)^* \widehat{K}_r(q), \quad \widetilde{K}_r(q) = \widehat{K}_r(q) S(q)^{-1/2}$$

on a neighborhood where  $S$  is invertible in the source corner. To lift a state, lift its positive square root  $c_0$ , put  $H = c^* c$ , and divide by the positive scalar  $\tau(H)$ . To lift a POVM, lift the square roots of its effects, put  $A_o = c_o^* c_o$ ,  $S = \sum_o A_o$ , and define

$$e_o = S^{-1/2} A_o S^{-1/2}.$$

These constructions may be applied simultaneously to any finite family by shrinking to one common interval.

*Scope.* The inverse square root is taken in the moving source corner, whose unit is its projection. Zero Kraus branches and rank-deficient state densities are allowed. No global normalization is claimed.

---

**Provenance.** ids: D1214; proved in: theory/sidequests/f1-limit/lifting.md; tested by: theory/checks/f1\_limit\_check.py (L2–L4, finite normalizations); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 15.3** (Local lifting of parabolic operational circuits). **PROVED** *Fix a finite endpoint circuit built from normalized states, effects, finite POVMs, normalized corner instruments, retained context, preparation, discard, assembly, split and classical wiring. There is an endpoint neighborhood on which all its labels have continuous lifts of the same types, and the lifted circuit evaluates to the original circuit at one. Pointwise coefficient-Gram equality is preserved on intervals and eventual Gram equality on germs.*

*Scope.* The lift is neither unique nor global. It lifts one chosen circuit and the relations of the presented interval category; it does not preserve every additional endpoint equation or provide a section of the evaluation functor.

---

**Provenance.** ids: F1-LIM-LIFT; proved in: theory/sidequests/f1-limit/lifting.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: L2–L4; W1–W5 for finite typed wiring; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Raw coefficient lifts exist for every corner element. At the endpoint,  $S(q_0)$  is the source identity, so invertibility persists locally and the binomial functional-calculus series for  $S^{-1/2}$  converges uniformly. The displayed formulas then give exact normalization throughout a smaller interval. There are only finitely many labels and contextual-recovery neighborhoods, so their intersection is still an endpoint neighborhood. Assembly, split and classical wiring are already global continuous labels. Reusing one lift for every repeated label gives the required circuit. Congruence of Gram equality is proved fibre by fibre and needs no continuous environmental unitary.  $\square$

**Definition 15.4** (Germs of full projection objects). An object germ is represented on some interval  $[1 - \varepsilon, 1 + \varepsilon]$  by a finite-support family of fixed matrix multiplicities and continuous self-adjoint projection matrices. Two such presentations are equal as germs when they agree literally after restriction to a smaller interval; isomorphic presentations are related by explicit morphisms instead. Morphisms are germs of the corresponding rectangular corner sections. Operations and the permutation-matrix associators descend under restriction. Evaluation at one is canonical, evaluation away from one requires a representative, and no  $C^*$ -norm is placed on germ Hom spaces.

*Scope.* A local lift is one representative on one neighborhood. Essential surjectivity is asserted only for endpoint germs, not for evaluation from an arbitrarily fixed large interval.

---

**Provenance.** ids: D1265; proved in: theory/sidequests/f1-limit/karoubi-lifting.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 15.5** (Full endpoint projection lifting). **PROVED** *Every finite family of self-adjoint projection objects in a positive fibre has simultaneous local continuous lifts. Evaluation is full on already chosen section objects, and endpoint germ evaluation is full and essentially surjective onto the finite self-adjoint completion. A unitary between two chosen endpoint object lifts has a local unitary lift.*

*Scope.* The construction uses positive spectral cutoff and is not algebraic base change over the generic coefficient ring. It gives no global canonical choice of object lifts.

---

**Provenance.** ids: F1-LIM-KLIFT; proved in: theory/sidequests/f1-limit/karoubi-lifting.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K3); spectral-gap and fullness arguments by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Lift the entries of an endpoint projection  $p_0$  with constant normalized Hecke coefficients and take the self-adjoint part  $a$ . Put  $h = 2a - 1$ . After shrinking,  $h^2$  remains within  $1/2$  of the identity, so

$$j = h(h^2)^{-1/2}, \quad p = \frac{1+j}{2}$$

is a continuous self-adjoint projection with  $p(q_0) = p_0$ . Finitely many such cutoffs share a common interval. A fibre morphism lifts entrywise and is compressed by the lifted source and target projections, proving fullness. For an endpoint unitary, normalize a raw rectangular lift  $a$  to  $a(a^*a)^{-1/2}$ . Its final projection is continuous and equals the target projection at the endpoint; after shrinking their difference has norm less than one, forcing equality because every nonzero projection has norm one.  $\square$

**Theorem 15.6** (Normalized processes on lifted completed objects). PROVED *Fix finitely many nonzero completed objects with chosen local lifts. Every specified endpoint state, effect, finite POVM and normalized retained instrument family lifts locally. For  $K_{o,i} : X \rightarrow Y$ , its unnormalized branch density is*

$$T_o(\rho) = \frac{w_Y}{w_X} \sum_i K_{o,i} \rho K_{o,i}^*,$$

*and the  $o$ -block of the recorded quantum–classical density is  $|O|T_o(\rho)$ . On parabolic corners the dimension ratio is  $P_\alpha/P_\beta$ , and tensoring a right context leaves it unchanged.*

*Scope.* For continuous labels, equality means pointwise coefficient-Gram equality; for germs it means eventual pointwise Gram equality. Fibrewise stable mixing does not provide, and the construction does not require, a continuous family of environmental unitaries.

---

**Provenance.** ids: F1-LIM-KPROC; proved in: theory/sidequests/f1-limit/karoubi-lifting.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_karoubi\_check.py (K2,K4,K6); W1–W5; finite-family lifting by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* States, POVMs and Kraus families are normalized by the same square-root formulas as above, now in finite matrix corners. Cross-object cyclicity of the unnormalized trace gives

$$\tau_Y(T_o(\rho)a) = \tau_X\left(\rho \sum_i K_{o,i}^* a K_{o,i}\right),$$

which fixes the factor  $w_Y/w_X$ . The uniform classical trace contributes  $|O|$ . Tensor multiplicativity of  $d$  cancels the dimension of a retained right context. For a parabolic projection,  $w_\alpha = 1/P_\alpha$ , yielding the stated specialization.  $\square$

## 15.2 Finite protocols and positive-limit conditioning

**Definition 15.7** (Finite typed protocol and postselection). A finite protocol is a rooted finite instrument tree. Its root carries a continuous normalized density on a finite register word. Every internal vertex carries a typed finite-outcome circuit and may depend on the preceding finite outcome history; following an edge selects one named outcome while retaining the declared quantum output. Each leaf carries an effect. Branch weights are obtained by composing the outcome CP maps and applying the final Born pairing; event weights are finite sums of branch weights. A conditioned probability  $N(q)/D(q)$  is considered near one only when  $D(1) > 0$ .

*Scope.* Only finite trees and specified instruments are included. A POVM does not determine its postmeasurement state, and no conditioned limit is asserted when the limiting denominator vanishes.

---

**Provenance.** ids: D1212; proved in: theory/sidequests/f1-limit/protocols.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1,W2,W5); theory/checks/f1\_limit\_check.py (L11); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 15.8** (Continuity of finite Born experiments). PROVED *Every finite continuous typed instrument tree has continuous nonnegative branch and event Born weights, and evaluation commutes with running the protocol. If a conditioning event has  $D(1) > 0$ , then on a smaller endpoint neighborhood*

$$\lim_{q \rightarrow 1} \frac{N(q)}{D(q)} = \frac{N(1)}{D(1)}.$$

*The same statement holds for a finite germ protocol after choosing one common representative interval.*

*Scope.* The theorem is analytic and is not inferred from finite samples. If  $D(1) = 0$ , even continuous normalized instruments can have conditioned probabilities with different subsequential limits.

---

**Provenance.** ids: F1-LIM-BORN; proved in: theory/sidequests/f1-limit/protocols.md; tested by: L11 gives the zero-denominator falsifier; W1,W2,W5 test finite routing; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* All generator matrices, trace pairings and wiring permutations vary continuously. Composition and tensor use finite matrix products and Kronecker products. Induction on tree depth therefore gives continuous positive unnormalized branch densities. Their trace pairings with leaf effects are continuous and nonnegative. Instrument unitality conserves the sum of child masses, and evaluation commutes with every finite operation. If  $D(1) > 0$ , continuity makes  $D(q) > D(1)/2$  nearby, where ordinary quotient continuity gives the result.  $\square$

The positivity condition cannot be deleted. On  $1 \leq q \leq 3/2$ , let  $t = q - 1$  and choose a unitary  $U(q)$  that oscillates between the identity and  $2e_1(q) - 1$ . The continuous Kraus pair

$$K_s(q) = tU(q), \quad K_f(q) = \sqrt{1 - t^2} 1,$$

extended by  $(0, 1)$  at one, is normalized. Its success probability is  $t^2 \rightarrow 0$ , while a conditioned return experiment has subsequential limits 1 and  $1/4$ .

### 15.3 A Lüders experiment on three constituents

**Definition 15.9** (Three-constituent retained-context protocol). In  $H_3(q)$  put

$$e_1 = \frac{T_1 + 1}{q + 1}, \quad e_2 = \frac{T_2 + 1}{q + 1}, \quad \alpha = (1, 2).$$

Prepare the normalized corner density  $e_2$  and refine to the complete three-part object with the isometry  $v = e_2$ , producing the full density  $(q + 1)e_2$ . If  $z_{\text{std}}$  is the central support of the unique  $M_2(\mathbb{C})$  summand, put  $P_2 = z_{\text{std}}e_2$ . Apply the Lüders instrument

$$K_s = P_2, \quad K_f = 1 - P_2, \quad [x^3] \longrightarrow [x^3]\{s, f\},$$

postselect the successful branch while retaining its quantum output, apply the right-context extension of  $u_1 = 2e_1 - 1$  from the first two constituents, and finally measure the POVM  $(P_2, 1 - P_2)$ .

*Scope.* The successful Kraus operator is part of the experiment. Replacing it by  $u_1P_2$  leaves the first POVM unchanged but changes the later return probability, so the effects alone do not define this protocol.

---

**Provenance.** ids: D1217; proved in: theory/sidequests/f1-limit/protocols.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W4); theory/checks/f1\_limit\_check.py (L5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 15.10** (Three-constituent Lüders witness). **PROVED** For every  $q > 0$ , the preceding protocol has first success probability, conditional return probability and joint probability

$$\begin{aligned} s(q) &= \frac{q(q + 1)}{1 + q + q^2}, \\ r(q) &= \left(1 - \frac{2q}{(q + 1)^2}\right)^2, \\ p(q) &= s(q)r(q). \end{aligned}$$

The successful density is

$$h_s(q) = \frac{1 + q + q^2}{q} P_2(q).$$

At  $q = 1$  these become

$$s(1) = \frac{2}{3}, \quad h_s(1) = 3P_2(1), \quad r(1) = \frac{1}{4}, \quad p(1) = \frac{1}{6}.$$

*Scope.* The unitary  $u_1$  has the identity conjugation channel on the isolated two-constituent commutative algebra, but its retained action on the full three-constituent algebra is not erased. The theorem uses the coefficient trace and the specified Lüders backaction.

---

**Provenance.** ids: F1-LIM-H3; proved in: theory/sidequests/f1-limit/protocols.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W4); theory/checks/f1\_limit\_check.py (L5); theory/checks/f1\_operational\_check.py (H5); reviewed in: theory/verdicts/f1-limit-adjudication.md.


---

*Proof.* The parabolic trace normalization sends the corner state to  $h = (q + 1)e_2$ . Write  $d = 1 + q + q^2$  and  $\gamma = \tau_3(P_2) = q/d$ . The first probability is

$$\tau_3(hP_2) = (q + 1)\gamma = \frac{q(q + 1)}{d}.$$

Because the successful Kraus operator is  $P_2$ ,  $P_2 h P_2 = (q + 1)P_2$ ; division by  $s(q)$  gives  $h_s = P_2/\gamma$ . In the standard two-dimensional block,  $u_1 = 2P_1 - 1$  and the two rank-one projections have overlap  $\text{Tr}_2(P_1 P_2) = q/(q + 1)^2$ . Therefore the return amplitude is  $1 - 2q/(q + 1)^2$  and its squared modulus is  $r(q)$ . Multiplying branch probabilities and substituting  $q = 1$  proves the remaining formulas.  $\square$

Three constituents, one retained context



A continuous experiment at  $q = 1$




Figure 2: The three-constituent Lüders protocol. Refinement is followed by the specified backaction  $K_s = P_2$ , postselection succeeds with probability  $2/3$  at the endpoint, retained conjugation returns with conditional probability  $1/4$ , and the joint endpoint probability is  $1/6$ .

## 16 Universal Hecke composition and physical exchange

The positive Hecke tower contains more than its familiar partial-flag corners. Closing the tower under finite sums and self-adjoint retracts gives a finite graded composition category, while passing to finite-dimensional right modules gives the same category algebraically. The tensor product is induction along the ordered block inclusion. This point matters: the plain vector-space tensor product of modules over two different Hecke algebras is not yet a module over the next Hecke algebra.

There are also two different notions of exchange. The usual Hecke braid is an algebraic isomorphism, but it is not unitary for the coefficient-trace  $C^*$ -structure away from  $q = 1$ . Polar decomposition of the longest Hecke elements instead produces canonical unitary reversals. Their block commutators obey cactus, rather than braid, coherence.

### 16.1 The finite completion of a multiplicative sequence

**Definition 16.1** (Multiplicative sequence, skeleton, and variance). A multiplicative sequence over  $\mathbb{C}$  is a family of unital algebras  $A_* = (A_n)_{n \geq 0}$ , with  $A_0 = \mathbb{C}$ , and unital algebra maps

$$\mu_{m,n} : A_m \otimes A_n \longrightarrow A_{m+n}$$

whose two composites from  $A_l \otimes A_m \otimes A_n$  to  $A_{l+m+n}$  agree. The unit maps are the scalar identifications  $\mu_{0,n}(\lambda \otimes a) = \lambda a$  and  $\mu_{n,0}(a \otimes \lambda) = \lambda a$ .

Its skeleton  $\mathbb{C}[A_*]$  has objects  $[n]$ , zero morphisms between unequal degrees,  $\text{End}([n]) = A_n$ , composition given by algebra multiplication, and tensor induced by  $\mu$ . A linear presheaf is a functor  $\mathbb{C}[A_*]^{\text{op}} \rightarrow \text{Vect}_{\mathbb{C}}$ . At degree  $n$  it is a right  $A_n$ -module:

$$m \cdot a = F(a)m.$$

The representable  $\text{Hom}(-, [n])$  is the right regular module, because precomposition sends  $x$  to  $xa$ .

*Scope.* The direction of the module action is part of the definition. No opposite algebra is inserted later, and no morphisms between different degrees are created at the skeletal stage.

---

**Provenance.** ids: D1221; proved in: theory/sidequests/f1-limit/universal.md; tested by: structural proof; no dedicated finite gate; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.2** (Finite Day–Schur–Weyl completion). Assume every  $A_n$  is finite-dimensional over  $\mathbb{C}$ . Define

$$\text{SW}_{\text{fin}}(A_*) = \bigoplus_{n \geq 0} \text{mod}_{\text{fd}}\text{-}A_n,$$

where objects have finite degree support. For a right  $A_m$ -module  $M$  and a right  $A_n$ -module  $N$ , set

$$M \star N = (M \otimes_{\mathbb{C}} N) \otimes_{A_m \otimes A_n} A_{m+n}.$$

Here  $A_{m+n}$  is a left  $A_m \otimes A_n$ -module through  $\mu_{m,n}$  and a right module by multiplication. The unit is  $\mathbb{C} = A_0$ , and the associator is the canonical balanced-tensor-product isomorphism.

*Scope.* Finite-dimensionality of every level algebra is required for this finite subcategory to be closed under  $\star$ . The tensor is induction, not the unextended tensor product  $M \otimes_{\mathbb{C}} N$ .

---

**Provenance.** ids: D1222; proved in: theory/sidequests/f1-limit/universal.md; tested by: structural balanced-tensor proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.3** (Finite projective composition category). The additive Karoubi completion  $\text{Proj}(A_*)$  has degree- $n$  objects  $(r, e)$ , where  $r \geq 0$  and  $e \in M_r(A_n)$  is an idempotent. It represents the right module  $eA_n^r$ . For two objects in degree  $n$ ,

$$\text{Hom}((r, e), (s, f)) = fM_{s,r}(A_n)e,$$

with matrix multiplication as composition; unequal degrees have zero Hom. Tensor uses the matrix amplification of  $\mu_{m,n}$  and the idempotent  $\mu_{m,n}^{(r,s)}(e \otimes f)$ .

If every  $A_n$  is a finite-dimensional  $C^*$ -algebra and every  $\mu_{m,n}$  is a star-homomorphism, the positive version uses self-adjoint projections and matrix adjoint as its dagger. Its comparison with bare right modules is algebraic: the bare module category carries no chosen Hilbert form or dagger.

*Scope.* The positive construction requires star-preserving structural maps. An algebraic module equivalence does not manufacture a dagger on its target.

---

**Provenance.** ids: D1223; proved in: theory/sidequests/f1-limit/universal.md; tested by: finite matrix-corner derivation; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.4** (Positive Hecke sequence and its universal completion). For real  $q > 0$ , put  $H_0(q) = H_1(q) = \mathbb{C}$ . For  $n \geq 2$ , let  $H_n(q)$  be generated by self-adjoint  $T_1, \dots, T_{n-1}$  with

$$(T_i - q)(T_i + 1) = 0, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad T_i T_j = T_j T_i \quad (|i - j| > 1).$$

For  $w \in S_n$ ,  $T_w$  is the reduced-word basis element and  $\tau_n(T_w) = \delta_{w,1}$ . Ordered block permutations define

$$\iota_{m,n}(T_u \otimes T_v) = T_{u \times v}.$$

These maps are injective unital trace-preserving star-homomorphisms and make  $H_*(q)$  a positive multiplicative sequence.

Let

$$R = \mathbb{Z}[t, t^{-1}, \{P_\alpha(t)^{-1}\}_\alpha]$$

be the localized coefficient ring used by the generic based Hecke algebras. The generic category is the finite additive self-adjoint-idempotent envelope

$$\mathcal{U}_R = \text{Kar}_*(\text{Add}(\mathbb{C}[H_*(R)])).$$

Evaluation is defined for its displayed objects and arrows. The full positive fibre is

$$\mathcal{U}_q = \text{Proj}_*(H_*(q)).$$

For  $x = [1]$ ,  $\text{End}_{\mathcal{U}_q}(x^{\otimes n}) = H_n(q)$ . A nonzero projection corner uses the coefficient trace divided by the trace of its corner unit. Normalized states and effects live on these finite  $C^*$ -endomorphism algebras.

*Scope.* The generic envelope contains self-adjoint projectors presented over  $R$ . It need not evaluate essentially surjectively onto projections constructed separately by functional calculus in one positive fibre. This is an amplitude category; its operational completion is supplied separately and does not require a fusion hypothesis.

---

**Provenance.** ids: D1101,D1102,D1224,F1-HCK-POS,F1-HCK-TOWER; proved in: theory/sidequests/f1-limit/universal.md; tested by: theory/checks/f1\_operational\_check.py (H13, finite corner samples); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.5** (Partial-flag completion). For a composition  $\alpha = (\alpha_1, \dots, \alpha_r)$  of  $n$ , let  $W_\alpha \leq S_n$  preserve its consecutive blocks and define

$$P_\alpha(q) = \sum_{w \in W_\alpha} q^{\ell(w)}, \quad e_\alpha(q) = P_\alpha(q)^{-1} \sum_{w \in W_\alpha} T_w.$$

The partial-flag category has

$$\text{Hom}(\alpha, \beta) = e_\beta H_n(q) e_\alpha$$

for equal totals, zero Homs otherwise, identity  $e_\alpha$ , adjoint inherited from  $H_n(q)$ , and concatenation tensor. Its projection-model functor sends

$$\alpha \mapsto e_\alpha H_n(q), \quad z \mapsto L_z : e_\alpha H_n(q) \rightarrow e_\beta H_n(q).$$

The extension under finite sums and self-adjoint retracts is the universal positive Hecke category above.

*Scope.* The normalization by  $P_\alpha$  is essential. The completion statement is proved below from the complete-flag object, not assumed from an older sketch about partial flags.

---

**Provenance.** ids: D1141,D1225; proved in: theory/sidequests/f1-limit/universal.md; tested by: theory/checks/f1\_operational\_check.py (H13, levels 3 and 4 at the declared parameters); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.6** (Davydov–Molev parameter and finite module category). Put  $v = \sqrt{q} > 0$ . If  $t_i^{\text{DM}}$  satisfies

$$(t_i^{\text{DM}} - v)(t_i^{\text{DM}} + v^{-1}) = 0,$$

then the project conventions are

$$q_{\text{project}} = v^2, \quad T_i^{\text{project}} = v, t_i^{\text{DM}}.$$

Let  $\text{DM}_{\text{fin}}(v)$  be the finite-support, finite-dimensional right-module subcategory of the associated Hecke Schur–Weyl category. Its tensor is the induced module just defined. The projection realization  $(r, e) \mapsto e H_n(q)^r$  is the proposed strong monoidal comparison

$$\mathcal{U}_q \simeq \text{DM}_{\text{fin}}(\sqrt{q}).$$

*Scope.* This comparison is algebraic. It respects the multiplicative sequence and induction tensor, but it asserts no dagger on the bare-module target.

---

**Provenance.** ids: D1226,D1227; proved in: theory/sidequests/f1-limit/universal.md; tested by: parameter identity and structured proof; no dedicated exact gate; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 16.7** (Universal finite Hecke composition). **PROVED** For every  $q > 0$ , the category  $\mathcal{U}_q$  is algebraically strong-monoidally equivalent to the finite-support category of finite-dimensional right modules over the Hecke multiplicative sequence, with tensor

$$(M \otimes N) \otimes_{H_m(q) \otimes H_n(q)} H_{m+n}(q).$$

*It is also the additive self-adjoint Karoubi completion of the partial-flag corner category. Generic evaluation applies to the self-adjoint projectors presented over  $R$ .*

*Scope.* The bare-module equivalence is algebraic. Generic evaluation is not claimed essentially surjective on all fibrewise functional-calculus projections, and no cross-degree amplitude morphisms are added.

**Provenance.** ids: F1-LIM-UNIV,D1221,D1222,D1223,D1224,D1225; proved in: theory/sidequests/f1-limit/universal.md; tested by: theory/checks/f1\_operational\_check.py (H13, finite parabolic corners only); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Contravariance gives  $(m \cdot a) \cdot b = m \cdot (ab)$ , fixing right-module variance. Both bracketings of three induced modules reduce canonically to

$$(L \otimes M \otimes N) \otimes_{A_l \otimes A_m \otimes A_n} A_{l+m+n},$$

so associativity and the pentagon follow from the common balanced universal map. The representables multiply because  $((A_m \otimes A_n) \otimes_{A_m \otimes A_n} A_{m+n}) \cong A_{m+n}$ .

Finite sums of representables give free right modules; splitting an idempotent  $e$  gives  $eA_n^r$ , and  $\text{Hom}_{A_n}(eA_n^r, fA_n^s) = fM_{s,r}(A_n)e$ . Since a finite-dimensional  $C^*$ -algebra is semisimple, all of its finite-dimensional modules are projective. This proves the algebraic equivalence. Star preservation of the block maps sends tensor projections to projections.

For partial flags, write  $x_\alpha = \sum_{w \in W_\alpha} T_w$ . Pairing terms along a simple parabolic reflection gives  $T_s x_\alpha = q x_\alpha$  and its right-handed star version. Hence

$$x_\alpha^2 = P_\alpha x_\alpha, \quad x_\alpha^* = x_\alpha, \quad \tau_n(e_\alpha) = P_\alpha^{-1}.$$

Block length additivity gives  $P_{\alpha\|\beta} = P_\alpha P_\beta$  and  $\iota(e_\alpha \otimes e_\beta) = e_{\alpha\|\beta}$ . Finally the complete composition  $(1, \dots, 1)$  has  $e = 1$  in each degree, so its finite sums and retracts generate every object of  $\mathcal{U}_q$ .  $\square$

**Theorem 16.8** (Algebraic Hecke Schur–Weyl comparison). **PROVED** With  $q_{\text{project}} = v^2$  and  $T_i^{\text{project}} = v, t_i^{\text{DM}}$ , the category  $\mathcal{U}_q$  is algebraically strong-monoidally equivalent to  $\text{DM}_{\text{fin}}(v)$ .

*Scope.* Only finite degree support and finite-dimensional modules are included. The comparison does not turn the algebraic Hecke braiding into a physical unitary or endow the bare-module category with a dagger.

**Provenance.** ids: F1-LIM-DM,D1226,D1227; proved in: theory/sidequests/f1-limit/universal.md; tested by: exact parameter calculation; categorical equivalence by proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Substitution  $t_i^{\text{DM}} = v^{-1}T_i$  turns its quadratic relation into  $(T_i - v^2)(T_i + 1) = 0$ . The braid and distant-commutation relations are homogeneous, and the same scaling respects every ordered block inclusion. Thus the full multiplicative sequences agree. The preceding theorem then identifies their finite right-module categories with their induction tensor.  $\square$

## 16.2 Algebraic braids and unitary cactus exchange

**Definition 16.9** (Adjacent spectral sign). In  $H_n(q)$  define

$$u_i = \frac{2T_i + 1 - q}{q + 1} = 2\frac{T_i + 1}{q + 1} - 1.$$

This is  $+1$  on the  $q$ -eigenspace of  $T_i$  and  $-1$  on its  $(-1)$ -eigenspace. Its adjacent braid defect is

$$u_i u_{i+1} u_i - u_{i+1} u_i u_{i+1} = -\frac{(q-1)^2}{(q+1)^2} (u_i - u_{i+1}).$$

*Scope.* The definition is fibrewise for  $q > 0$ . It produces local unitary exchanges, not a braiding away from one.

---

**Provenance.** ids: D1228; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L7, six declared rational parameters); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.10** (Polar longest-element reversal). Let  $w_0^{(n)}$  be the longest permutation and  $D_n = T_{w_0^{(n)}}$ . In the positive fibre put

$$|D_n| = (D_n^2)^{1/2}, \quad J_n = D_n |D_n|^{-1}, \quad J_0 = J_1 = 1.$$

The square root is the unique positive  $C^*$ -functional-calculus root. For an interval  $I$  of consecutive tensor positions,  $J_I$  is the shifted copy of  $J_{|I|}$ .

*Scope.* This definition uses positivity and functional calculus at real  $q > 0$ ; it is not an element-wise construction over the localized generic coefficient ring.

---

**Provenance.** ids: D1229; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L13, ranks at most 4 and  $q=1,4,9$ ); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.11** (Unitary block commutor). For  $m, n \geq 0$ , define on tensor powers

$$\sigma_{m,n} = J_{m+n} \iota_{m,n}(J_m \otimes J_n).$$

On projection-completed objects, restrict this operator to the source and target corners; on finite sums also apply the canonical matrix-index flip. Its coherence consists of naturality, symmetry  $\sigma_{n,m}\sigma_{m,n} = 1$ , the unit laws, and the cactus square formed by the interval reversals.

*Scope.* The commutor is ordered and coboundary. Symmetry here means involutivity of the commutor, not the braid relation for three adjacent factors.

---

**Provenance.** ids: D1230; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L13, finite cactus samples); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 16.12** (Analytic domain of physical exchange). The block commutor is extra canonical structure on every positive real  $C^*$ -fibre and varies continuously for  $q > 0$ . It is the polar unitary of the positive minimal Hecke braid exchanging the two blocks. At  $q = 1$  it is the ordinary unitary block permutation. Its functional-calculus coefficients need not belong to the localized generic ring.

*Scope.* No algebraic generic commutor is stipulated, and no braid coherence is claimed for  $q \neq 1$ .

---

**Provenance.** ids: D1231; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L13, declared fibres only); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 16.13** (Spectral unitaries do not braid away from one). **PROVED** *For  $q > 0$ , the standard algebraic Hecke exchange is invertible but unitary for the coefficient-trace  $C^*$ -structure only at  $q = 1$ . The adjacent spectral signs are self-adjoint unitaries with the exact defect displayed above, and therefore do not braid when  $q \neq 1$ .*

*Scope.* This separates algebraic Yang–Baxter exchange from physical unitary exchange. It does not obstruct a coboundary commutor.

---

**Provenance.** ids: F1-LIM-EXCH,D1226,D1228; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L7, six rational parameters); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The algebraic generator  $v^{-1}T_i$  has eigenvalues  $v$  and  $-v^{-1}$ , so both have modulus one exactly at  $v = 1$ . Meanwhile  $e_i = (T_i + 1)/(q + 1)$  is a self-adjoint projection and  $u_i = 2e_i - 1$ . Writing  $u_1 = ax + b$ ,  $u_2 = ay + b$ , with  $a = 2/(q + 1)$ ,  $b = (1 - q)/(q + 1)$  and  $x = T_1, y = T_2$ , braid cancellation gives

$$u_1 u_2 u_1 - u_2 u_1 u_2 = a^2 b (x^2 - y^2) + ab^2 (x - y).$$

Since  $x^2 - y^2 = (q - 1)(x - y)$ , this is the stated multiple of  $u_1 - u_2$ . The standard basis makes  $T_1 - T_2$  nonzero.  $\square$

**Theorem 16.14** (Canonical unitary cactus exchange). PROVED *For every  $q > 0$ , the polar reversals  $J_n$  define a continuous natural unitary coboundary commutor on all of  $\mathcal{U}_q$ . It is symmetric, satisfies the cactus relations, restricts to every partial-flag and arbitrary projection corner, is the polar unitary of the block braid, and becomes ordinary symmetric exchange at  $q = 1$ .*

*Scope.* The theorem is analytic on positive fibres and positive intervals. It makes no claim over the generic localized ring and no claim of braiding away from one.

---

**Provenance.** ids: F1-LIM-COB,D1229,D1230,D1231; proved in: theory/sidequests/f1-limit/exchange.md; tested by: theory/checks/f1\_limit\_check.py (L13, n at most 4 and q=1,4,9); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The longest element is self-adjoint and invertible. The relation  $w_0 s_i = s_{n-i} w_0$  gives  $D_n T_i = T_{n-i} D_n$ , hence  $D_n^2$  is central. Its positive modulus is therefore central, so  $J_n$  is a canonical self-adjoint unitary satisfying  $J_n T_i J_n = T_{n-i}$ . Shifted copies on disjoint intervals commute, and unitary conjugation carries a subinterval reversal to its reflected subinterval reversal. Thus

$$J_{[p,r]} J_{[k,l]} = J_{[p+r-l, p+r-k]} J_{[p,r]},$$

which is the interval presentation of the cactus relations.

The product  $J_{m+n}(J_m \otimes J_n)$  reverses internally and then totally, so the internal reversals cancel and the two ordered block algebras swap. This proves naturality; the same interval identities prove symmetry and the cactus square. Conjugation carries any tensor projection to its swapped projection, so compression extends the commutor to all finite sums and retracts.

If  $B_{m,n} = D_{m+n}(D_m \otimes D_n)^{-1}$  is the positive minimal block braid, then

$$\sigma_{m,n}^* B_{m,n} = |D_{m+n}| |D_m \otimes D_n|^{-1} > 0.$$

Uniqueness of polar decomposition identifies  $\sigma_{m,n}$  as its polar unitary. Functional calculus varies continuously on positive invertible matrices. At one, every  $T_i$  is a transposition, so the construction is the ordinary block permutation.  $\square$

## 17 Schur–Weyl quotients and the boundary of fusion

At the symmetric endpoint, the Hecke tower becomes the symmetric-group tower. Its action on tensor powers of  $\mathbb{C}^d$  gives a concrete monoidal quotient, but it changes the physical trace: a permutation is weighted by its number of cycles rather than only by whether it is the identity. The difference disappears at fixed tensor degree as  $d$  grows, and the associated conditional expectations converge with an explicit finite- $d$  bound.

Rigidity and fusion require further data. The nonnegative grading of the universal amplitude category rules out evaluation and coevaluation maps inside that category. A rigid comparison must name a target, a monoidal functor, dual objects, cups and caps, dagger and trace compatibility. A Temperley–Lieb or Fibonacci branch adds quotient ideals and a different trace; neither is an automatic consequence of the positive-real endpoint.

### 17.1 The tensor-power quotient

**Definition 17.1** (The symmetric endpoint amplitude category). For real  $q > 0$ , let  $H_n(q)$  be the positive type- $A$  Hecke algebra with self-adjoint generators  $T_i$ , relations

$$(T_i - q)(T_i + 1) = 0, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad T_i T_j = T_j T_i \quad (|i - j| > 1),$$

and coefficient trace  $\tau_n(T_w) = \delta_{w,1}$ . Ordered block inclusion maps  $T_u \otimes T_v$  to  $T_{u \times v}$ . The finite positive composition category  $\mathcal{U}_q$  consists of finite sums and self-adjoint projection retracts of the graded regular objects; unequal degrees have zero Homs. At  $q = 1$ ,  $H_n(1) = \mathbb{C}[S_n]$  and the generator  $x = [1]$  satisfies  $\text{End}(x^{\otimes n}) = \mathbb{C}[S_n]$ .

*Scope.* This is an amplitude category with finite  $C^*$ -endomorphism algebras. Its nonnegative grading is retained, and no internal dual for  $x$  is assumed.

---

**Provenance.** ids: D1101,D1102,D1224; proved in: theory/sidequests/f1-limit/universal.md; tested by: theory/checks/f1\_operational\_check.py (H13, finite corner samples); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.2** (Classical Schur–Weyl quotient). Fix  $d \geq 1$ , put  $V_d = \mathbb{C}^d$ , and let

$$P_\sigma(v_1 \otimes \cdots \otimes v_n) = v_{\sigma^{-1}(1)} \otimes \cdots \otimes v_{\sigma^{-1}(n)}.$$

Define

$$\rho_{d,n} : \mathbb{C}[S_n] \longrightarrow \text{End}_{U(d)}(V_d^{\otimes n}), \quad \rho_{d,n}(\sigma) = P_\sigma.$$

Its kernel is the sum of Wedderburn blocks indexed by partitions of  $n$  with more than  $d$  rows. Equivalently, for  $n \geq d + 1$ , it is the ideal generated by the antisymmetrizer on the first  $d + 1$  positions. The map is faithful exactly when  $d \geq n$ .

*Scope.* The target is the full  $U(d)$ -commutant on one tensor power. For  $d < n$  the map is a quotient, and no faithful embedding is claimed.

---

**Provenance.** ids: D1232; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8, n and d at most 4); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.3** (Quotient assembly). Under the standard associator  $V_d^{\otimes(m+n)} \cong V_d^{\otimes m} \otimes V_d^{\otimes n}$ , define

$$\nu_{m,n}(\rho_{d,m}(a) \otimes \rho_{d,n}(b)) = \rho_{d,m+n}(\iota_{m,n}(a \otimes b)).$$

These are associative unital algebra maps. On right modules, the associated Schur–Weyl functor is

$$M \longmapsto M \otimes_{\mathbb{C}[S_n]} V_d^{\otimes n},$$

where the tensor power has its left permutation action.

*Scope.* Well-definedness uses compatibility of the kernels with block inclusion. The source tensor on modules is balanced induction, not an objectwise  $S_{m+n}$ -action on  $M \otimes N$ .

**Provenance.** ids: D1233; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8, finite rank/trace samples; assembly by proof); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 17.4** (Coherent finite-rank Schur–Weyl quotient). **PROVED** *For every  $d \geq 1$  and  $n \geq 0$ ,  $\rho_{d,n}$  is a surjective star-homomorphism onto the tensor-power  $U(d)$ -commutant. Its kernel is exactly the sum of partition blocks with more than  $d$  rows, equivalently the antisymmetrizer-generated ideal, and it is faithful exactly for  $d \geq n$ . The maps  $\nu_{m,n}$  form a multiplicative sequence and induce a strong monoidal functor from finite right modules with induction tensor.*

*Scope.* The theorem is at the symmetric endpoint. It permits a quotient kernel and does not identify induction with the bare vector-space tensor product of modules over different symmetric groups.

**Provenance.** ids: F1-LIM-SW,D1232,D1233; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8, n,d at most 4; general kernel by proof); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The permutation convention gives  $P_\sigma P_\pi = P_{\sigma\pi}$  and  $P_\sigma^* = P_{\sigma^{-1}}$ . Position permutations commute with the diagonal  $U(d)$  action. Conversely, the complex span of the Lie algebra of  $U(d)$  is  $\mathfrak{gl}_d(\mathbb{C})$ , so the unitary and classical general-linear commutants agree; classical Schur–Weyl duality makes the map surjective.

If  $d \geq n$ , apply a putative linear relation among the  $P_\sigma$  to  $e_1 \otimes \cdots \otimes e_n$ ; the resulting  $n!$  words are distinct. If  $d < n$ , the  $(d+1)$ -strand antisymmetrizer acts through  $\bigwedge^{d+1} V_d = 0$ . In the decomposition

$$V_d^{\otimes n} \cong \bigoplus_{\substack{\lambda \vdash n \\ \text{rows}(\lambda) \leq d}} L_d(\lambda) \otimes S^\lambda,$$

the block  $\text{End}(S^\lambda)$  survives exactly for partitions with at most  $d$  rows. The branching rule says precisely those excluded blocks contain the sign representation of the embedded  $S_{d+1}$ , proving the ideal description.

On elementary tensors, block permutation gives

$$\rho_{d,m+n}(\iota(\sigma \otimes \pi)) = \rho_{d,m}(\sigma) \otimes \rho_{d,n}(\pi).$$

This proves kernel compatibility and associativity. Cancelling consecutive extension and restriction of scalars identifies the image of  $(M \star N)$  with  $(M \otimes_{\mathbb{C}[S_m]} V_d^{\otimes m}) \otimes (N \otimes_{\mathbb{C}[S_n]} V_d^{\otimes n})$ , giving the strong monoidal structure.  $\square$

## 17.2 Which trace does the quotient see?

**Definition 17.5** (Tensor trace and physical subgroup expectation). Define the normalized tensor trace

$$\text{tr}_{d,n}(a) = d^{-n} \text{Tr}(\rho_{d,n}(a)).$$

For a permutation  $\sigma$ , this is

$$\text{tr}_{d,n}(\sigma) = d^{\text{cyc}(\sigma)-n}.$$

When  $d \geq n$  and  $K = S_m \times S_{n-m}$ , the physical trace-preserving expectation  $E_{d,K} : \mathbb{C}[S_n] \rightarrow \mathbb{C}[K]$  is characterized by

$$\text{tr}_{d,n}(k^* E_{d,K}(x)) = \text{tr}_{d,n}(k^* x), \quad k \in K.$$

In the group basis its coefficients solve  $G_d c = b$ , where

$$G_d(k, h) = d^{\text{cyc}(k^{-1}h)-n}, \quad b_k = d^{\text{cyc}(k^{-1}\sigma)-n}.$$

*Scope.* Faithfulness on the full group algebra, and therefore this Gram inversion there, requires  $d \geq n$ . On the quotient image the tensor trace remains faithful for all  $d$ .

---

**Provenance.** ids: D1234; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8,L9, declared finite ranks); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.6** (Fixed-degree stable trace comparison). Let  $\tau_n$  be coefficient trace and let  $E_{\infty,K}$  delete group-basis coefficients outside  $K$ . For fixed  $n$  and  $x = \sum_{\sigma} a_{\sigma} \sigma$ ,

$$|\mathrm{tr}_{d,n}(x) - \tau_n(x)| \leq \frac{1}{d} \sum_{\sigma \neq 1} |a_{\sigma}|.$$

Put  $N = |K|$ . If both  $d \geq n$  and  $d > N - 1$ , then for  $\sigma \notin K$  every coefficient of  $E_{d,K}(\sigma)$  has modulus at most

$$\frac{1}{d - (N - 1)},$$

whereas  $E_{d,K}(\sigma) = \sigma$  for  $\sigma \in K$ . Thus the physical expectation converges coefficientwise to coefficient deletion at fixed degree.

*Scope.* Both inequalities on  $d$  are required: one for faithfulness and one for the Neumann-series bound. No equality between the two traces or expectations is claimed at fixed finite  $d$ .

---

**Provenance.** ids: D1235; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8,L9; general bound analytic); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 17.7** (Stable physical trace and expectation). **PROVED** *At fixed  $n$ , the normalized tensor trace converges coefficientwise to  $\tau_n$ . If  $K = S_m \times S_{n-m}$ ,  $d \geq n$ , and  $d > |K| - 1$ , then  $E_{d,K}$  converges coefficientwise to  $E_{\infty,K}$  with the bound above. For fixed  $h \geq 0$ ,  $\tau_n(h) = 1$ , and  $0 \leq e \leq 1$ , put*

$$\Delta_d(y) = \frac{1}{d} \sum_{\sigma \neq 1} |y_{\sigma}|.$$

*Whenever  $\Delta_d(h) < 1$ , the normalized quotient density has Born probability  $\mathrm{tr}_{d,n}(he)/\mathrm{tr}_{d,n}(h)$  and*

$$\left| \frac{\mathrm{tr}_{d,n}(he)}{\mathrm{tr}_{d,n}(h)} - \tau_n(he) \right| \leq \frac{\Delta_d(he) + \Delta_d(h)}{1 - \Delta_d(h)}.$$

*Scope.* All convergence is at fixed tensor degree and for fixed coefficient data. There is no uniform claim when the degree, coefficient norms, subgroup, or experiment varies with  $d$ .

---

**Provenance.** ids: F1-LIM-SWTRACE,D1234,D1235; proved in: theory/sidequests/f1-limit/schur-weyl.md; tested by: theory/checks/f1\_limit\_check.py (L8 for  $n,d$  at most 4; L9 finite S2-in-S3 sentinel); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* A tensor-basis word is fixed by  $P_{\sigma}$  exactly when it is constant on each cycle, so there are  $d^{\mathrm{cyc}(\sigma)}$  fixed words. A nonidentity permutation has at most  $n - 1$  cycles, proving the trace bound. Moreover

$$\mathrm{tr}_{d,n}(x^* x) = d^{-n} \mathrm{Tr}(\rho_{d,n}(x)^* \rho_{d,n}(x)),$$

so its radical is exactly the Schur–Weyl kernel.

The conditional expectation is orthogonal projection for this trace. Its Gram matrix is  $G_d = I + R$ , with  $\|R\|_{\infty} \leq (N - 1)/d$ . Under the two displayed hypotheses,

$$\|G_d^{-1}\|_{\infty} \leq \frac{1}{1 - (N - 1)/d}.$$

For  $\sigma \notin K$ , the right-hand Gram vector has norm at most  $1/d$ , which proves the coefficient bound. The finite-rank difference is real: for  $n = 3$ ,  $K = \langle(12)\rangle$ , one obtains  $E_{d,K}((23)) = d^{-1}1$ , while coefficient deletion gives zero.

Finally the trace estimate applied to  $h$  makes its physical normalization at least  $1 - \Delta_d(h)$ . Apply the same estimate to  $he$  and divide by this lower bound to obtain the stated Born inequality.  $\square$

### 17.3 Rigid and fusion targets require additional data

**Definition 17.8** (Rigid target datum). A rigid comparison consists of an additive idempotent-complete monoidal category  $\mathcal{R}$ , a named strong monoidal functor  $i : \mathcal{U}_q \rightarrow \mathcal{R}$ , and a named left and right dual  $y^\vee$  of  $y = i(x)$ . It includes

$$\begin{aligned} \text{coev}_R : \mathbf{1} &\rightarrow y \otimes y^\vee, & \text{ev}_R : y^\vee \otimes y &\rightarrow \mathbf{1}, \\ \text{coev}_L : \mathbf{1} &\rightarrow y^\vee \otimes y, & \text{ev}_L : y \otimes y^\vee &\rightarrow \mathbf{1}, \end{aligned}$$

satisfying the four zigzags with the stated associators. The target is generated from  $y, y^\vee$  by tensor, finite sums and retracts. The functor may have a kernel. A positive dagger version additionally requires a  $C^*$ -category, a star-preserving  $i$ , compatible dagger/pivotal data, and a positive spherical structure. Duals of sums and retracts are formed in the additive Karoubi closure.

*Scope.* Full faithfulness is additional and is never inferred from the existence of cups and caps. The target closure is part of the datum.

---

**Provenance.** ids: D1236; proved in: theory/sidequests/f1-limit/fusion.md; tested by: theory/checks/f1\_limit\_check.py (L8, finite quotient witness only); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.9** (Fusion quotient datum). A fusion quotient consists of a monoidal ideal  $\mathcal{I}$  in a chosen rigid extension, a compatible dagger on the quotient, a positive pivotal trace, and the instruction to quotient and then take additive Karoubi completion. Any trace radical or negligible ideal is named explicitly. The result is called fusion only after proving semisimplicity, simple unit, finite-dimensional Hom spaces, duals, and finitely many simple isomorphism classes.

*Scope.* A family of degreewise algebra ideals is insufficient unless it is closed under tensor, composition, dagger, and the chosen duality operations.

---

**Provenance.** ids: D1237; proved in: theory/sidequests/f1-limit/fusion.md; tested by: structural proof; no finite sample proves fusion; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.10** (Temperley–Lieb branch). For  $q > 0$ , set

$$f_i = \frac{q - T_i}{q + 1}, \quad \delta = \sqrt{q} + \frac{1}{\sqrt{q}}.$$

A Temperley–Lieb branch imposes the tensor-ideal relations

$$f_i f_{i+1} f_i = \delta^{-2} f_i, \quad f_{i+1} f_i f_{i+1} = \delta^{-2} f_{i+1},$$

equivalently killing the excluded three-strand Hecke summand. It also supplies diagrammatic cups and caps, dagger, and the Jones–Markov trace. The faithful coefficient trace is not declared to descend through this nonzero ideal.

*Scope.* This is a named quotient branch with new rigid and trace data. It is not a property of the full positive flag/Hecke category.

---

**Provenance.** ids: D1238; proved in: theory/sidequests/f1-limit/fusion.md; tested by: theory/checks/f1\_operational\_check.py (H10, finite trace obstruction); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 17.11** (Fibonacci root branch). The unitary Fibonacci branch is the even part of the level-three  $SU(2)/\text{Jones}$  fusion quotient. Choose

$$q_H = e^{2\pi i/5}, \quad v = e^{\pi i/5}, \quad v^2 = q_H, \quad q_{\text{ILZ}} = -v,$$

and distinguish the loop generator from the projection by

$$U_i^{\text{ILZ}} = \delta f_i, \quad \delta = -(q_{\text{ILZ}} + q_{\text{ILZ}}^{-1}) = v + v^{-1} = 2 \cos(\pi/5).$$

Quotient by the Jones trace radical, take additive Karoubi completion, and retain the even labels  $\{0, 2\}$ , with

$$2 \otimes 2 \cong 0 \oplus 2.$$

*Scope.* The square parameter has order five. This root-of-unity construction is not the  $q = 1$  specialization of the positive-real branch, where  $\delta = 2$  and the spin tower is untruncated.

---

**Provenance.** ids: D1239; proved in: theory/sidequests/f1-limit/fusion.md; tested by: primary-source parameter derivation; no cyclotomic checker; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 17.12** (A positive rigid target at the symmetric endpoint). **PROVED** *The category  $\mathcal{U}_q$  is not rigid: its degree-one generator has no left or right dual. A positive rigid comparison requires the complete datum above. At  $q = 1$ , for every  $d \geq 1$ , the full replete  $C^*$ -tensor subcategory of  $\text{Rep}(U(d))$  generated by  $V_d, V_d^*$ , finite sums and subobjects is a concrete positive rigid target of the Schur–Weyl quotient image. The named functor  $i_d : \mathcal{U}_1 \rightarrow \mathcal{R}_d$  may have a kernel.*

*Scope.* The comparison is not asserted fully faithful in degrees  $n > d$ . The target has infinitely many simple objects as degrees vary and is not thereby a fusion category.

---

**Provenance.** ids: F1-LIM-RIGID,D1236,D1237; proved in: theory/sidequests/f1-limit/fusion.md; tested by: theory/checks/f1\_limit\_check.py (L8, finite Schur–Weyl kernel samples); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Every summand of  $x \otimes y$  has degree at least one for any object  $y$  of the nonnegatively graded category, while the unit has degree zero. Thus an evaluation  $x \otimes y \rightarrow \mathbf{1}$  must vanish and cannot satisfy a zigzag.

At the endpoint, send  $x$  to  $V_d$ , symmetric-group arrows to their tensor permutations, and a projection retract to the range of its image projection. This is a named strong monoidal star-functor, with precisely the Schur–Weyl kernel. The usual pairing and copairing

$$\text{ev}(\varphi \otimes v) = \varphi(v), \quad \text{coev}(1) = \sum_i e_i \otimes e_i^*$$

satisfy the zigzags by basis contraction. The Hilbert dagger and  $V_d \cong V_d^{\vee\vee}$  give positive dimension  $d$  and ordinary matrix trace. Closing under sums and retracts supplies duals for the whole generated target.  $\square$

**Theorem 17.13** (Temperley–Lieb and Fibonacci are separate quotient branches). **PROVED** *The Temperley–Lieb branch requires its named tensor ideal, cups, caps, dagger, and Jones–Markov trace. The faithful coefficient trace cannot descend through the nonzero quotient ideal. At  $q = 1$  and  $d = 2$ , coefficient-trace weights  $(1/2, 1/2)$  on the symmetric and alternating projections become physical weights  $(3/4, 1/4)$ . The Fibonacci branch further requires the displayed order-five parameter, Jones trace radical quotient, additive Karoubi completion, and even subcategory.*

*Scope.* Neither the  $d = 2$  spin tower nor Fibonacci is identified with the full positive-real endpoint. Fibonacci is not obtained from any real  $q > 0$  in the displayed loop normalization.

---

**Provenance.** ids: F1-LIM-TL,D1237,D1238,D1239; proved in: theory/sidequests/f1-limit/fusion.md; tested by: theory/checks/f1\_operational\_check.py (H10, finite TL/trace sample; no root-of-unity gate); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The element  $f_i$  is the spectral projection for the  $(-1)$ -eigenvalue of  $T_i$ . The quotient relation gives

$$f_i f_{i+1} f_i = \frac{q}{(q+1)^2} f_i = \delta^{-2} f_i.$$

Yet

$$\tau_n(f_i) = \frac{q}{q+1}, \quad \text{tr}_J(f_i) = \delta^{-2} = \frac{q}{(q+1)^2}.$$

If the faithful coefficient trace descended through a nonzero quotient kernel, a nonzero  $z$  in that kernel would satisfy both  $\tau_n(z^*z) > 0$  and  $\tau_n(z^*z) = 0$ , a contradiction. At  $q = 1$ , the two projections  $(1 \pm (12))/2$  have coefficient weights  $1/2, 1/2$ , whereas their ranks on  $(\mathbb{C}^2)^{\otimes 2}$  are 3 and 1.

On the positive-real branch,  $\delta = \sqrt{q} + q^{-1/2} \geq 2$ . The Fibonacci value  $2 \cos(\pi/5) = (1 + \sqrt{5})/2$  is smaller than two, so it requires the complex root and sign convention in the definition. At that root, the raw Temperley–Lieb algebra is semisimplified by its Jones trace radical; the even surviving labels give the stated Fibonacci fusion rule. The three traces—coefficient, finite tensor, and Jones—therefore belong to three different comparison maps and cannot be silently identified.  $\square$

## 18 The affine-vector commutant and its Fourier-polarized form

Adding a vector to a pair of flags produces a finite arithmetic observable algebra that remembers more than the ordinary flag Hecke algebra. Its extra generator is a controlled translation projector. Fourier transformation turns these translation constraints into controlled phase constraints, but the flag register must simultaneously be dualized and reversed. We fix all indices and signs below, including the square of the Fourier bridge.

### 18.1 Affine invariant kernels

**Definition 18.1** (Arithmetic affine-vector context algebra). Let  $k = \mathbb{F}_Q$ , let  $L$  be a finite-dimensional  $k$ -vector space,  $G = GL(L)$ , and  $X = \text{Fl}(L)$  the set of complete flags. The affine group

$$\text{Aff}(L) = G \ltimes L, \quad (g, a)(h, b) = (gh, a + gb),$$

acts on  $Y_L = X \times L$  by

$$(g, a)(F, x) = (gF, a + gx).$$

Define

$$R_X(L) = \text{End}_{\text{Aff}(L)} \mathbb{C}[Y_L]$$

with Hilbert adjoint and normalized operator trace  $\tau_Q = \text{Tr}/(|X||L|)$ . The invariant-kernel convention is

$$K_f((F, x), (F', x')) = f(F, F', x' - x).$$

*Scope.* The field cardinality is denoted  $Q$ . The symbol  $q$  below is reserved for a real deformation parameter. The affine group symbol does not reuse the notation for a Weyl subalgebra.

---

**Provenance.** ids: D1241; proved in: theory/sidequests/fl-limit/mirabolic.md; tested by: theory/checks/fl\_limit\_mirabolic\_check.py (B1); reviewed in: theory/verdicts/fl-limit-adjudication.md.

**Theorem 18.2** (Affine convolution as a physical commutant). **PROVED** *For every finite field  $\mathbb{F}_Q$  and finite-dimensional  $L$ , convolution on  $GL(L)$ -orbits of triples  $(F, F', v)$  is  $*$ - and trace-preservingly the commutant  $R_X(L)$ . In particular its normalized operator trace is faithful and positive.*

*Scope.* This is an arithmetic finite-Hilbert-space theorem. It does not infer positivity of a generic deformation from a polynomial presentation.

---

**Provenance.** ids: F1-MIR-AFF; proved in: theory/sidequests/fl-limit/mirabolic.md; tested by: theory/checks/fl\_limit\_mirabolic\_check.py (B1, n=2 and Q=2,3); reviewed in: theory/verdicts/fl-limit-adjudication.md.

*Proof.* If  $B \leq GL(L)$  stabilizes a reference complete flag, then  $\text{Aff}(L)/(B, 0) \cong X \times L$  by  $(g, a)B \mapsto (gB, a)$ . An operator commutes with the affine action exactly when its matrix kernel is constant on diagonal affine orbits. Translation by  $-x$  reduces a pair to  $((F, 0), (F', x' - x))$ , leaving diagonal  $GL(L)$ -invariance. Matrix multiplication becomes

$$(f * g)(F, F', v) = \sum_{H \in X} \sum_{u \in L} f(F, H, u) g(H, F', v - u),$$

which is the stated convolution. Taking adjoints gives  $f^*(F, F', v) = \overline{f(F', F, -v)}$ . Normalized matrix trace restricts to a faithful positive trace on every finite-dimensional  $*$ -subalgebra.  $\square$

## 18.2 The controlled translation generator

For  $u \in L$  and an additive character  $\chi$  of  $L$ , write  $X(u)e_x = e_{x+u}$  and  $Z(\chi)e_x = \chi(x)e_x$ . For a subspace  $U \leq L$ , put

$$P_U^X = |U|^{-1} \sum_{u \in U} X(u), \quad P_U^Z = |U^{\text{ann}}|^{-1} \sum_{\chi \in U^{\text{ann}}} Z(\chi).$$

**Definition 18.3** (Controlled translation and phase constraints). For a complete flag  $F = (U_i(F))_{i=0}^n$ , with  $U_0 = 0$  and  $U_n = L$ , define on  $\mathbb{C}[\text{Fl}(L)] \otimes \mathbb{C}[L]$

$$C_i^X = \sum_F |F\rangle\langle F| \otimes P_{U_i(F)}^X, \quad C_i^Z = \sum_F |F\rangle\langle F| \otimes P_{U_i(F)}^Z, \quad 0 \leq i \leq n.$$

Thus  $C_0^Z = I \otimes |0\rangle\langle 0|$  and  $C_n^Z = I$ . Let  $\text{Ctx}_X(L)$  be the  $*$ -algebra generated by all relative-position flag operators  $A_w \otimes I$  and all  $C_i^X$ . For  $n \geq 1$ , its marked projection and unnormalized vector generator are

$$e = C_1^X, \quad T_0 = Qe - I.$$

The same definitions apply on the dual vector space.

*Scope.* The endpoints  $i = 0, n$  are included. In particular the index-zero phase constraint is non-scalar in rank one. A named additive character is used when linear duals are identified with additive characters.

---

**Provenance.** ids: D1142, D1242; proved in: theory/sidequests/f1-limit/mirabolic.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B2); theory/checks/f1\_limit\_bridge\_check.py (B3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 18.4** (Generation by flags and one controlled line). **PROVED** For  $L$  of dimension  $n \geq 1$  over  $\mathbb{F}_Q$ , the embedded flag Hecke algebra acts as  $A_w \otimes I$ , the marked element  $(T_0 + I)/Q$  is exactly  $C_1^X$ , and these operators generate the full affine commutant  $R_X(L)$ . Moreover

$$e^2 = e = e^*, \quad \tau_Q(e) = Q^{-1}, \quad T_0^2 = (Q - 2)T_0 + (Q - 1)I.$$

*Scope.* The additional constraints  $C_i^X$  are useful tests but are redundant as algebra generators. The theorem fixes the orbit basis, adjoint, trace and Hecke inclusion, rather than asserting only an abstract algebra isomorphism.

---

**Provenance.** ids: F1-MIR-GEN; proved in: theory/sidequests/f1-limit/mirabolic.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B2, n=2 and Q=2,3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The zero-vector orbit kernels act as  $A_w \otimes I$ . The kernel of  $T_0 + I$  is one precisely when the two flags agree and the vector difference lies in the controlled line  $U_1(F)$ . Dividing by  $Q$  averages translations along that line, hence gives  $C_1^X$ . A finite-group average is a self-adjoint projection; its rank in each flag block is  $|L|/Q$ , which gives the trace. The quadratic formula follows from  $T_0 = Qe - I$ . Rosso's generation theorem then gives the reverse inclusion from these concrete operators to the full invariant-kernel algebra.  $\square$

## 18.3 Fourier duality with exact indices

**Definition 18.5** (Fourier dual flag reversal). Fix a named nontrivial additive character  $\psi : k \rightarrow U(1)$ . Define

$$\mathcal{F}_{L,\psi}e_x = |L|^{-1/2} \sum_{\lambda \in L^\vee} \psi(-\lambda(x))e_\lambda.$$

If  $F = (U_i)_{i=0}^n$ , its reversed dual flag is

$$D_L(F)_j = U_{n-j}^\perp, \quad 0 \leq j \leq n.$$

Let  $D_L$  also denote the induced flag-basis unitary and put  $W_{L,\psi} = D_L \otimes \mathcal{F}_{L,\psi}$ . Define

$$R_Z(L^\vee) = W_{L,\psi} R_X(L) W_{L,\psi}^*.$$

The phase-polarized groupoid has finite-dimensional  $k$ -spaces as objects, linear isomorphisms as arrows, and retains the pair  $(R_X(L), R_Z(L^\vee))$ , their Hecke inclusions, all controlled constraints and the named  $W_{L,\psi}$ . A same-register realization additionally names a linear self-duality  $\sigma : L \rightarrow L^\vee$ ; morphisms act on flags, vectors and duals by their induced permutation unitaries.

*Scope.* The negative Fourier kernel, the phase  $\psi$ , dual-flag reversal and any same-register self-duality are named choices. Forgetting them retains only the corresponding conjugacy class of algebra comparisons.

---

**Provenance.** ids: D1243, D1244; proved in: theory/sidequests/f1-limit/mirabolic.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 18.6** (Fourier exchange of controlled constraints). PROVED For every finite field  $k = \mathbb{F}_Q$ , every  $n$ -dimensional  $L$ , every named nontrivial  $\psi : k \rightarrow U(1)$  and every  $0 \leq i \leq n$ ,

$$\begin{aligned} W_{L,\psi} C_i^X W_{L,\psi}^* &= C_{n-i}^Z, \\ W_{L,\psi} (A_w \otimes I) W_{L,\psi}^* &= A_{w_0 w w_0} \otimes I. \end{aligned}$$

The operators on the right generate exactly  $R_Z(L^\vee)$ , including  $C_0^Z = I \otimes |0\rangle\langle 0|$ . Under canonical double duality,

$$W_{L^\vee,\psi} W_{L,\psi} e_{(F,x)} = e_{(F,-x)}.$$

The construction is natural for linear isomorphisms.

*Scope.* Only the vector Fourier factor is tensor-preserving under direct sums. The full flag-Fourier bridge is not monoidal on complete-flag registers and does not identify either commutant with the full Weyl matrix algebra.

---

**Provenance.** ids: F1-MIR-FOURIER; proved in: theory/sidequests/f1-limit/mirabolic.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B3, Fourier constraints, reversal, square and ranks); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Character orthogonality makes  $\mathcal{F}_{L,\psi}$  unitary and gives

$$\mathcal{F}_{L,\psi} P_U^X \mathcal{F}_{L,\psi}^* = P_{U^\perp}^Z.$$

Annihilators reverse a full flag, so the index  $i$  becomes  $n - i$ . The intersection-rank matrix of a flag pair transforms by  $r'(i, j) = i + j - n + r(n - i, n - j)$ , which is the rank matrix of  $w_0 w w_0$ . This proves both conjugation formulas and the generation statement. Applying the negative Fourier kernel twice yields

$$|L|^{-1} \sum_{y \in L} \sum_{\lambda \in L^\vee} \psi(-\lambda(x + y)) e_y = e_{-x};$$

dual reversal squares to the original flag. In characteristic two this vector negation is the identity.  $\square$

**Definition 18.7** (Vector-polarization bridge package). In rank  $n$ , a bridge package retains the affine commutant and orbit data, the controlled translation and phase families for every  $0 \leq i \leq n$ , the based Hecke inclusion, conjugate-linear orbit star, reference trace, algebraic and trace-supported endpoints, the named phase and Fourier unitary, and the regular one-sided operational boundary. When composition is needed, it additionally names an arithmetic symplectic or affine decomposable-shuffle correspondence with its ternary shuffle coherence.

Only  $\mathcal{F}_{L \oplus M} = \mathcal{F}_L \otimes \mathcal{F}_M$  is a literal tensor identity. On the decomposable flag–vector subspace the full bridge uses

$$W_{L \oplus M} J_{L,M} = J_{L^\vee, M^\vee} ((W_L \otimes W_M) \otimes \text{Rev}),$$

where  $\text{Rev}$  reverses the shuffle word. An isomorphism of packages preserves all named UCP maps, the regular label class and quotient boundary functor, and the exact square  $W_{L^\vee, \psi} W_{L, \psi}(F, x) = (F, -x)$ .

*Scope.* An abstract algebra isomorphism is not an isomorphism of bridge packages. The decomposable flag subspace is generally proper; even adjoining shuffles does not tensor-factor the full complete-flag Hilbert space.

---

**Provenance.** ids: D1259; proved in: theory/sidequests/f1-limit/mirabolic.md; theory/sidequests/f1-limit/type-c.md; tested by: B1–B7 in the two bridge checkers, at their stated finite scopes; reviewed in: theory/verdicts/f1-limit-adjudication.md.

## 19 Positive mirabolic fibres and their regular boundary

The affine commutant at a finite field has a physical matrix trace. To move away from arithmetic parameters, one needs an algebraic reason that the same coefficient trace stays positive. Orbital valencies supply that reason for every real  $q > 1$ . At  $q = 1$  the form becomes degenerate: the full algebraic specialization remains, while its reference GNS representation retains only the ordinary symmetric-group Hecke algebra. The operational boundary is therefore built from one-sided sections positive at every nearby real parameter, rather than from values sampled only at prime powers.

### 19.1 Orbit labels, valencies and the positive Gram form

**Definition 19.1** (Mirabolic orbit poset and based star-trace). For  $w \in S_n$ , define a partial order on  $\{1, \dots, n\}$  by

$$i \prec_w j \iff i < j \text{ and } w^{-1}(i) < w^{-1}(j).$$

For an antichain  $A$ , set

$$\downarrow_w A = \{i : i = a \text{ or } i \prec_w a \text{ for some } a \in A\}.$$

The orbit label  $(w, A)$  denotes the affine orbital whose flag relative position is  $w$  and whose vector has maximal-support antichain  $A$ ; its adjacency basis element is  $T_{w,A}$ . The empty antichain is the zero-vector Hecke basis.

On the polynomial orbit-basis algebra define the conjugate-linear star by

$$T_{w,A}^* = T_{w^{-1}, w^{-1}(A)}$$

and the coefficient trace by

$$\tau_q(T_{w,A}) = \begin{cases} 1, & w = 1, A = \emptyset, \\ 0, & \text{otherwise.} \end{cases}$$

The proposed positive domain is real  $q > 1$ ; evaluation at one is retained as a positive-semidefinite boundary form.

*Scope.* Vector negation is part of the orbital adjoint but does not change support antichains. Positivity is not stipulated and is proved below. No positive trace is claimed for  $0 < q < 1$ .

---

**Provenance.** ids: D1245, D1246; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: theory/checks/f1\_limit\_check.py (L12); B1,B4 in theory/checks/f1\_limit\_mirabolic\_check.py; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 19.2** (Antichain classification and orbital valency). **PROVED** *Let  $L = \mathbb{F}_Q^n$ , fix a reference complete flag, and let  $w \in S_n$ . The residual stabilizer orbits on  $L$  are indexed by antichains  $A$  in the preceding poset. The full affine orbital  $(w, A)$  has exact valency*

$$k_{w,A}(Q) = Q^{\ell(w) + |\downarrow_w A| - |A|} (Q - 1)^{|A|}.$$

*Scope.* This holds for every finite field, including characteristic two and extension fields. It is an orbit-counting theorem, not a positivity statement at a non-arithmetic parameter.

---

**Provenance.** ids: F1-MIR-VAL; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: theory/checks/f1\_limit\_check.py (L12, n at most 3 and Q=2,3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The intersection of the two Borels for relative position  $w$  contains the diagonal torus and precisely the shears  $I + cE_{ij}$  with  $i \prec_w j$ . Allowed shears change only coordinates below an existing support coordinate, so the maximal nonzero support is an invariant antichain. Conversely, diagonal scaling normalizes the coordinates on  $A$ , and shears from each  $a \in A$  create arbitrary coordinates in its strict downset. The residual orbit therefore has  $(Q - 1)^{|A|} Q^{|\downarrow_w A| - |A|}$  elements. The Bruhat cell of  $w$  has  $Q^{\ell(w)}$  flags, giving the formula.  $\square$

**Theorem 19.3** (Faithful positive mirabolic trace for all real parameters above one). **PROVED** *For every  $n \geq 1$  and every real  $q > 1$ , the orbit-basis trace has diagonal Gram form*

$$\tau_q(T_{w,A}^* T_{v,C}) = \delta_{(w,A),(v,C)} q^{\ell(w) + |\downarrow_w A| - |A|} (q - 1)^{|A|}.$$

*It is a faithful normalized trace, and its left regular representation makes the fibre a finite  $C^*$ -algebra.*

*Scope.* The all-real conclusion follows from a polynomial identity and the displayed sign, not from numerical sampling at arithmetic values. For  $0 < q < 1$ , odd  $|A|$  gives a negative Gram entry.

---

**Provenance.** ids: F1-MIR-POS; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: L12 checks finite orbit counts; the uniform positivity statement is proved analytically; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* At every prime power, distinct orbital adjacency matrices have disjoint support, and their normalized Hilbert–Schmidt squared norms equal their valencies. Both the product coefficient and the valency are polynomials in  $q$ ; equality at infinitely many prime powers makes them identical. For  $q > 1$  every displayed diagonal entry is strictly positive. Polynomial extension of matrix-trace cyclicity gives traciality. Left multiplication is faithful because  $L_x 1 = x$ , and its adjoint is  $L_{x^*}$ .  $\square$

## 19.2 Hecke retraction and the trace-supported endpoint

**Definition 19.4** (Hecke inclusion and vector expectation). Let

$$i_q : H_n(q) \longrightarrow R_n(q), \quad T_w \longmapsto T_{w,\emptyset},$$

and define coefficient deletion by

$$E_q(T_{w,A}) = \begin{cases} T_w, & A = \emptyset, \\ 0, & A \neq \emptyset. \end{cases}$$

*Scope.* These are fixed-rank maps. They do not provide a generic cross-rank map  $R_m(q) \otimes R_n(q) \rightarrow R_{m+n}(q)$ ; in particular they leave room for separately constructed right module actions.

---

**Provenance.** ids: D1247; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B4); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 19.5** (Vector coarse graining as a conditional expectation). **PROVED** *For every real  $q > 1$ ,  $i_q$  is a trace-preserving unital  $*$ -monomorphism and  $E_q$  is its trace-preserving UCP conditional expectation, with  $E_q i_q = \text{id}$ . Hence Hecke densities, effects and retained Kraus lists lift unchanged to the mirabolic fibre.*

*Scope.* The reverse operation  $i_q E_q$  is generally a proper vector-sector coarse graining. The statement concerns fixed-rank operational data only.

---

**Provenance.** ids: F1-MIR-EXPECT; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B4); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The Gram formula makes  $E_q$  the trace-orthogonal projection onto the unital Hecke  $*$ -subalgebra. Traciality gives bimodularity. Testing a positive input against elements of the subalgebra proves positivity, and the identical argument in matrix amplifications proves complete positivity. The basis formulas give trace preservation and  $E_q i_q = \text{id}$ .  $\square$

**Definition 19.6** (Algebraic and trace-supported endpoints). The algebraic endpoint is the full based specialization  $R_n(1)$  with the specialized trace  $\tau_1$ . Its trace-supported endpoint is

$$R_n(1)/\mathcal{N}_{\tau_1}, \quad \mathcal{N}_{\tau_1} = \{x : \tau_1(x^*x) = 0\}.$$

The two endpoint algebras are not identified by definition.

*Scope.* No  $C^*$ -norm is placed on the degenerate algebraic endpoint. Only its GNS quotient carries the faithful reference representation.

---

**Provenance.** ids: D1248; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B4); theory/checks/f1\_limit\_check.py (L10); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 19.7** (The mirabolic reference quotient at one). PROVED *At  $q = 1$ , the coefficient trace is positive semidefinite and*

$$\mathcal{N}_{\tau_1} = \text{span}\{T_{w,A} : A \neq \emptyset\}.$$

*This nullspace is a two-sided  $*$ -ideal, and*

$$R_n(1)/\mathcal{N}_{\tau_1} \cong H_n(1) = \mathbb{C}[S_n]$$

*as a traced  $*$ -algebra. The map  $E_1$  is exactly the quotient map under the empty-orbit basis identification.*

*Scope.* The full algebraic specialization still contains the vector ideal. The claim concerns the GNS quotient selected by the reference trace.

---

**Provenance.** ids: F1-MIR-END; proved in: theory/sidequests/f1-limit/mirabolic-trace.md; tested by: B4, exact rank-two interpolation and null Gram; L10 rank one; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Setting  $q = 1$  in the diagonal Gram formula leaves norm one exactly for  $A = \emptyset$  and norm zero otherwise. The nullspace of a positive tracial form is a two-sided  $*$ -ideal by Cauchy–Schwarz. The surviving basis classes multiply by the Hecke rule at one, giving the group algebra and its coefficient trace. Since  $E_1$  deletes precisely the null basis, it is the quotient.  $\square$

### 19.3 The actual one-sided operational category

**Definition 19.8** (Regular one-sided labels). Fix  $J_\varepsilon = [1, 1 + \varepsilon]$ . A regular mirabolic section is

$$h(q) = \sum_{w,A} h_{w,A}(q) T_{w,A}(q), \quad h_{w,A} \in C(J_\varepsilon),$$

with pointwise polynomial multiplication, the orbit star and continuous coefficient trace. Hecke factors use their continuous section algebra on the same interval; separated register words use continuous coefficients in the product bases and the  $C(J_\varepsilon)$ -balanced common-parameter tensor.

A regular density additionally satisfies  $h(q) \geq 0$  and  $\tau_q(h(q)) = 1$  for every real  $1 < q \leq 1 + \varepsilon$ . A regular effect satisfies  $0 \leq e(q) \leq 1$  there; a regular POVM consists of finitely many such positive sections summing to one. Regular instrument coefficients are continuous and obey Kraus completeness at every real  $q > 1$  in the interval.

*Scope.* There is no  $C^*$ -norm on the boundary section algebra. Positivity only at prime powers is insufficient. Coefficient-pole densities are excluded and require separate singular boundary data.

---

**Provenance.** ids: D1249; proved in: theory/sidequests/f1-limit/mirabolic-boundary.md; tested by: uniform theorem by proof; L10 and B4 give boundary examples; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 19.9** (Regular separated operational category). For each real  $q > 1$ , quantum atoms are  $[H_n]$  and  $[R_n]$ ,  $n \geq 1$ , with their faithful coefficient traces; nonempty finite classical wires have uniform trace. The empty word has algebra  $\mathbb{C}$ , and ordered word tensor is the actual spatial tensor product. The interval version uses the regular common-parameter sections just defined.

Raw morphisms are finite typed planar circuits generated by all regular preparations, discards and finite POVMs; every retained same-register instrument

$$K : X \rightarrow X\mathcal{O}, \quad \sum_{o,i} K_{o,i}(q)^* K_{o,i}(q) = 1;$$

the explicit classical product, singleton, bijective relabeling and classical-wire routing maps; and

$$\begin{aligned} \text{up}_n : [H_n] &\rightarrow [R_n] & (E_q \text{ in Heisenberg form}), \\ \text{down}_n : [R_n] &\rightarrow [H_n] & (i_q \text{ in Heisenberg form}). \end{aligned}$$

Instrument lists are retained under separated context tensor and factorwise inclusion. With the uniform classical trace, the  $\mathcal{O}$ -block of the output density is  $|\mathcal{O}| \sum_i K_{o,i} h K_{o,i}^*$ .

Composition is typed gluing and ordered tensor is horizontal juxtaposition. Equality is the congruence generated by category, monoidal and explicit classical-wiring relations, literal state/POVM equality, pointwise equality of coefficient Kraus Grams within each outcome, the routed list identities, and  $\text{down}_n \text{up}_n = \text{id}_{[H_n]}$ . The reverse composite is  $i_q E_q$  and is not an identity. Germs identify circuits when these relations hold on a common smaller one-sided interval. There is no cross-rank mirabolic assembly generator.

*Scope.* The classical wiring is the explicit parameter-independent UCP presentation already proved for the core category; no unresolved wiring hypothesis remains. Equality is not an isolated-CP quotient and requires no continuous family of Kraus mixing matrices. In particular  $[R_m][R_n]$  is not identified with  $[R_{m+n}]$ .

---

**Provenance.** ids: D1257, D1218; proved in: theory/sidequests/f1-limit/mirabolic-boundary.md; theory/sidequests/f1-limit/classical-wiring.md; tested by: theory/checks/f1\_limit\_wiring\_check.py (W1-W5); B4; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 19.10** (Regular quotient boundary functor). Let  $\Pi_n : R_n(1) \rightarrow H_n(1)$  be the GNS quotient  $E_1$ . For a register word  $X$ , let  $Q_X$  tensor  $\Pi_n$  on every mirabolic factor and the identity on Hecke and classical factors; write  $\overline{X}$  for the word obtained by replacing every mirabolic atom by its Hecke atom. The endpoint category has these Hecke/classical words, the same wiring, all normalized preparations, POVMs and retained same-register instruments, the same label relations, and its UCP realization.

The boundary functor evaluates continuous coefficients at one, applies  $Q_X$  to every density, effect and Kraus entry, and sends every up/down generator to the identity. Evaluation at  $q > 1$  requires a chosen representative whose interval contains  $q$ .

*Scope.* Arbitrary UCP families with merely regular superoperator coefficients are not source morphisms. The regular density and retained-Kraus restrictions are part of the functor's domain.

---

**Provenance.** ids: D1258; proved in: theory/sidequests/f1-limit/mirabolic-boundary.md; tested by: structured descent proof; B4 and W1–W5 test primitive data; reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 19.11** (Regular operational mirabolic boundary). **PROVED** *The regular one-sided circuits and their germs form actual ordered monoidal categories with UCP realizations. Coefficient evaluation followed by the factorwise GNS quotient defines a monoidal endpoint functor to the separated Hecke operational category. It preserves all specified generators and typed relations. Finite protocol Born weights converge, and conditioned weights converge whenever the limiting conditioning probability is positive. Every individual finite endpoint Hecke circuit has a local regular lift.*

*Scope.* All source densities, effects and instruments are positive or normalized for every nearby real  $q > 1$ . Arithmetic-only positivity and arbitrary coefficient-regular UCP superoperators are excluded. No generic  $R_m(q) \otimes R_n(q)$  assembly is asserted.

---

**Provenance.** ids: F1-MIR-REG; proved in: theory/sidequests/f1-limit/mirabolic-boundary.md; tested by: B4 and W1–W5 support primitive formulas; functoriality and positivity by structured proof; reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Polynomial structure constants and the parameter-independent star make regular sections a  $*$ -algebra. For a regular positive density  $h$  and any endpoint Hecke element  $x$ , lift  $x$  through the Hecke inclusion. Positivity of  $\tau_q(x(q)^*h(q)x(q))$  for every real  $q > 1$  passes to one by coefficient continuity. Faithfulness of the endpoint quotient trace then forces  $Q_X(h(1)) \geq 0$ ; normalization also passes to the limit. The same argument handles effects and POVMs, while Kraus completeness passes algebraically and multiplicativity of  $Q_X$  makes every branch map descend.

The explicit wiring commutes with factorwise quotients. Pointwise Gram equality is a congruence fibre by fibre and descends under the linear quotient map. Hence the circuit quotient and endpoint functor are well defined. A finite branch density has continuous coefficients by induction over its circuit, so trace pairings and finite event sums converge. Positive-limit conditioning is ordinary quotient continuity. Local endpoint lifts use square-root normalization in the continuous Hecke subalgebra followed by  $i_q$ , on one common neighborhood for the finitely many labels.  $\square$

The real-neighborhood hypothesis is sharp. In  $H_2(q)$ , if  $e_+, e_-$  are the two spectral projections, then

$$h(q) = (q - \text{frac}32)e_+ + \frac{5}{2q}e_-$$

is normalized and positive at every prime power  $Q \geq 2$ , but has a negative  $e_+$  coefficient for  $1 < q < 3/2$ . It is not a regular density. Conversely, the rank-one singular state  $q(q-1)^{-1}(1-e)$  is positive and normalized for every  $q > 1$ , but has a coefficient pole and does not annihilate the endpoint null ideal. It is a legitimate punctured-fibre state requiring additional boundary data, not an object of the regular category.

## 20 Decomposable-shuffle correspondences in type C and the affine bridge

Orthogonal direct sum of symplectic spaces does not produce a standard parabolic inclusion of type-C Hecke algebras. It does, however, select a canonical subspace of decomposable isotropic flags. Component flags and a shuffle word form an orthonormal basis of that subspace. Compression to it and scaled averaging back give trace-adjoint UCP maps, while ternary shuffle bijections make the correspondence associative. The same construction works for affine flag–vector spaces at prime-power fibres.

### 20.1 The decomposable type-C arena

**Definition 20.1** (Arithmetic type-C context algebra). Let  $k = \mathbb{F}_Q$  and let  $V$  be a  $2n$ -dimensional symplectic  $k$ -space. Write  $X_C(V)$  for its complete isotropic flags

$$0 = U_0 < U_1 < \cdots < U_n, \quad \dim U_i = i,$$

where  $U_n$  is Lagrangian. With the flag basis of  $K_V = \mathbb{C}[X_C(V)]$  orthonormal, define

$$A_C(V) = \text{End}_{Sp(V)}(K_V)$$

with Hilbert adjoint and normalized operator trace.

*Scope.* At a prime power this is the finite type- $C_n$  flag Hecke commutant. No generic positive-real type-C Hecke category is stipulated here.

---

**Provenance.** ids: D1250; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 20.2** (Decomposable isotropic flags and shuffle isometry). Let  $V, W$  have symplectic ranks  $m, n$ . A complete isotropic flag in  $V \perp W$  is decomposable when every step is  $U_a \oplus Z_b$  for steps of component flags. Let  $\text{Sh}(m, n)$  be the two-letter words with  $m$  letters  $V$  and  $n$  letters  $W$ . If  $(a_r, b_r)$  counts the two letters in the first  $r$  positions of  $\sigma$ , then

$$(U, Z, \sigma) \mapsto (U_{a_r} \oplus Z_{b_r})_{r=0}^{m+n}$$

is the decomposable flag. Linear extension gives the basis isometry

$$J_{V,W} : K_V \otimes K_W \otimes \ell^2(\text{Sh}(m, n)) \longrightarrow K_{\text{dec}}.$$

Let  $p_{V,W} = J_{V,W} J_{V,W}^*$ .

*Scope.* The decomposable subspace is usually proper inside the full composite flag space. The shuffle factor is retained as a matrix-coherent register, rather than discarded as a classical label.

---

**Provenance.** ids: D1251; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5,B6); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Definition 20.3** (Local-symmetry product-shuffle corner). Put

$$H_{V,W} = Sp(V) \times Sp(W), \quad \mathcal{D}_{V,W} = \text{End}_{H_{V,W}}(K_{V \perp W}).$$

The product-shuffle arena is the corner  $p_{V,W} \mathcal{D}_{V,W} p_{V,W}$ , whose unit is  $p_{V,W}$  and whose trace is ordinary operator trace normalized by  $\text{rank}(p_{V,W})$ .

*Scope.* This local-symmetry commutant is larger than the full-symplectic commutant. The corner type and its non-scalar unit are part of the construction.

---

**Provenance.** ids: D1252; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 20.4** (Decomposable flags are component flags plus a shuffle). PROVED For symplectic spaces  $V, W$  over any finite field, the preceding flag map is a bijection, and

$$p_{V,W} \mathcal{D}_{V,W} p_{V,W} \cong A_C(V) \otimes A_C(W) \otimes M_{\text{Sh}(m,n)}(\mathbb{C})$$

as a traced  $*$ -algebra.

*Scope.* This is an arithmetic finite-dimensional commutant identity. It is not an algebra isomorphism between the full composite context and the product corner, and it includes all shuffle coherences.

---

**Provenance.** ids: F1-TC-DEC; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5, Sp4 over F2); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The dimensions of intersections with  $V$  and  $W$  recover the shuffle word and the nonrepeated intersections recover both component flags. Thus the displayed map is bijective. The subgroup  $Sp(V) \times Sp(W)$  acts on the two component flag factors and trivially on the shuffle factor. Its commutant on the decomposable space is therefore the tensor product of the two component commutants and the full matrix algebra on shuffles.  $\square$

## 20.2 Restriction and preparation

**Definition 20.5** (Trace-adjoint type-C channels). Let  $G = Sp(V \perp W)$  and let

$$\mathcal{E}_G(x) = |G|^{-1} \sum_{g \in G} U_g x U_g^*, \quad r_{V,W} = \frac{\text{rank}(p_{V,W})}{\dim K_{V \perp W}}.$$

For  $a \in A_C(V \perp W)$  and  $x \in p_{V,W} \mathcal{D}_{V,W} p_{V,W}$ , extended by zero on the orthogonal complement, define

$$\Phi_{V,W}(a) = p_{V,W} a p_{V,W}, \quad \Psi_{V,W}(x) = r_{V,W}^{-1} \mathcal{E}_G(x).$$

*Scope.* The scale  $r_{V,W}^{-1}$  is required for unitality and trace preservation. Compression is not asserted to be multiplicative on a generic full symplectic transition.

---

**Provenance.** ids: D1253; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 20.6** (Bidirectional positive type-C correspondence). PROVED For every prime power  $Q$  and finite symplectic  $V, W$  over  $\mathbb{F}_Q$ ,  $\Phi_{V,W}$  and  $\Psi_{V,W}$  are unital trace-preserving completely positive maps, and they are adjoint for the normalized traces:

$$\text{tr}_p(\Phi_{V,W}(a)x) = \text{tr}_G(a\Psi_{V,W}(x)).$$

*Scope.* These are restriction and preparation channels between two different observable algebras. Mutual trace adjointness does not make them inverse  $*$ -isomorphisms.

---

**Provenance.** ids: F1-TC-CP; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B5, exact averaging and trace pairing); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Compression is CP and sends the full unit to the corner unit. Transitivity of  $Sp(V \perp W)$  on complete isotropic flags gives  $\mathcal{E}_G(p_{V,W}) = r_{V,W}I$ , so scaled averaging is also UCP. For  $a$  commuting with  $G$ ,

$$\mathrm{Tr}(pa) = \mathrm{Tr}(\mathcal{E}_G(p)a) = r \mathrm{Tr}(a),$$

which proves trace preservation. The same averaging identity applied to  $ax$  gives the displayed adjointness.  $\square$

**Definition 20.7** (Ternary shuffle coherence). For ranks  $\ell, m, n$ , let  $\mathrm{Sh}(\ell, m, n)$  be the three-letter words with those multiplicities. The two coherence maps are the canonical bijections

$$\begin{aligned} \mathrm{Sh}(\ell, m) \times \mathrm{Sh}(\ell + m, n) &\longrightarrow \mathrm{Sh}(\ell, m, n), \\ \mathrm{Sh}(m, n) \times \mathrm{Sh}(\ell, m + n) &\longrightarrow \mathrm{Sh}(\ell, m, n), \end{aligned}$$

obtained by expanding each combined letter with the next letter of the inner word. All iterated flag isometries, compression channels and trace-adjoint preparations are compared using these bijections and the ordinary Hilbert associator.

*Scope.* The coherence maps preserve the order of the three summands. They are specified bijections, not silent equality of differently nested shuffle sets.

---

**Provenance.** ids: D1254; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B6); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 20.8** (Associativity of the type-C correspondence). **PROVED** *For three finite symplectic spaces over one finite field, the two iterations of decomposable restriction agree after the canonical ternary shuffle bijections. Their trace-adjoint preparation maps agree in the reverse direction.*

*Scope.* The theorem is arithmetic and finite-dimensional. It retains the shuffle register and supplies no symmetric exchange beyond the named reorderings.

---

**Provenance.** ids: F1-TC-COH; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B6, three rank triples); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Both nested isometries send a triple of component flags and a three-letter word to the same flag whose current step is the direct sum of the three current component steps. Hence both iterated projections are the same diagonal projection. Compression maps coincide. Their normalized-trace adjoints coincide as well; equivalently, the nested rank fractions multiply to the rank fraction of this common projection.  $\square$

### 20.3 The thin signed-permutation fibre

**Definition 20.9** (Thin type-C block comparison). At  $q = 1$ , let  $B_n$  be the signed permutation group. Embed  $B_m \times B_n$  in  $B_{m+n}$  by preserving the two coordinate blocks. Fix the convention that every signed permutation factors as a block signed permutation followed by one unsigned  $(m, n)$  shuffle. The induced group-algebra  $*$ -inclusion is the thin block assembly.

*Scope.* This definition concerns the group-algebra endpoint only. It includes no generic type-C Hecke lift of the block map.

---

**Provenance.** ids: D1255; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B6, signed factorizations); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 20.10** (Unsigned shuffles are the thin block cosets). **PROVED** For  $m, n \geq 0$ , every element of  $B_{m+n}$  has the fixed unique factorization above, so

$$\ell^2(B_{m+n}) \cong \ell^2(B_m \times B_n) \otimes \ell^2(\text{Sh}(m, n)).$$

For  $m, n > 0$  the block subgroup is proper and nonparabolic, although its group algebra embeds by a unital  $*$ -map.

*Scope.* The theorem leaves open generic bimodules or right Hecke actions. It states only that no standard generic  $H(B_m, q) \otimes H(B_n, q)$  parabolic assembly has been supplied.

---

**Provenance.** ids: F1-TC-ONE; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_bridge\_check.py (B6, four signed block factorizations); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* Move the unsigned shuffle of the two absolute-value blocks to the chosen side; what remains preserves both coordinate blocks and carries all signs. This is unique. The subgroup index is  $\binom{m+n}{m}$ . It has full Coxeter rank  $m+n$ , whereas a proper parabolic subgroup has smaller rank, proving nonparabolicity.  $\square$

## 20.4 Arithmetic affine decomposable composition

**Definition 20.11** (Affine decomposable product-shuffle arena). For finite vector spaces  $L, M$  over one finite field, let  $p_{L,M}^{\text{aff}}$  project onto basis vectors consisting of a decomposable complete flag in  $L \oplus M$  and an arbitrary vector  $(x, y) \in L \oplus M$ . The local symmetry is

$$\text{Aff}(L) \times \text{Aff}(M).$$

The affine product-shuffle corner, restriction and preparation maps are the local-symmetry commutant corner, compression by  $p_{L,M}^{\text{aff}}$ , and inverse-rank-fraction scaled averaging over  $\text{Aff}(L \oplus M)$ , exactly as in the symplectic formulas above.

*Scope.* This is arithmetic data at a prime-power fibre. It is not a generic-real mirabolic assembly map and is not a generator of the regular separated boundary category.

---

**Provenance.** ids: D1256; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B7); reviewed in: theory/verdicts/f1-limit-adjudication.md.

**Theorem 20.12** (Arithmetic affine product-shuffle correspondence). **PROVED** Let  $L, M$  be finite-dimensional over  $\mathbb{F}_Q$ . Decomposable flags and all vector labels identify the affine local-symmetry corner with

$$R_X(L) \otimes R_X(M) \otimes M_{\text{Sh}(\dim L, \dim M)}(\mathbb{C}).$$

Compression from the full affine commutant and scaled full-affine averaging back are mutually trace-adjoint trace-preserving UCP maps, coherent for three summands under the ternary shuffle bijections.

*Scope.* The hypotheses require a prime-power field. The result does not infer positivity or composition for all real  $q > 1$ , and it does not identify the full flag register with the product register.

---

**Provenance.** ids: F1-MIR-DEC; proved in: theory/sidequests/f1-limit/type-c.md; tested by: theory/checks/f1\_limit\_mirabolic\_check.py (B7, Q=2,3); reviewed in: theory/verdicts/f1-limit-adjudication.md.

*Proof.* The decomposable flag-vector basis is the tensor product of the two flag-vector bases and the shuffle basis, equivariantly for  $\text{Aff}(L) \times \text{Aff}(M)$ . Taking commutants gives the displayed corner. The full affine group is transitive on flag-vector basis elements, so averaging the decomposable projection gives its rank fraction times the identity. The compression, scaled-average and ternary coherence proofs are therefore identical to the symplectic arguments.  $\square$