# cft-anyons Lab Book

From category data to provably rigorous CFTs

cft-anyons contributors

October 27, 2026

## Contents

**1 Frontmatter, North Star, and Claim Status** **20**
1.1 The Two Success Criteria . . . 20
1.2 Claim Status . . . 20
1.3 Evidence Chain . . . 20

**2 Programme Map** **21**
2.1 The Pipeline . . . 21
2.2 Stage 1 — Category Input . . . 21
2.3 Stage 2 — Microscopic Models . . . 21
2.4 Stage 3 — Lattice Symmetry Generators . . . 21
2.5 Stage 4 — Provable Continuum Limit . . . 21
2.6 Stage 5 — Rigorous (C)FT Target . . . 22
2.7 First Decisions . . . 22

**3 Many-Particle Hilbert Spaces: the AND/OR/Vacuum Grammar** **23**
3.1 Sources and Dependencies . . . 23
3.2 The Three Primitives . . . 23
3.3 Two Monoidal Structures and Their Units . . . 23
3.4 The Grammar and its Carrier . . . 24
3.5 Particle-Number Grading . . . 24
3.6 Lamport Dependency Skeleton . . . 24
3.7 Scope and Deferral . . . 25

**4 Indefinite Particle Number: the Full Fock Completion** **26**
4.1 Sources and Dependencies . . . 26
4.2 Two Direct Sums — the Load-Bearing Distinction . . . 26
4.3 Completed Tensor Powers . . . 26
4.4 The Full Fock Space . . . 26
4.5 The Grammar Generates the Indefinite-Particle Space . . . 27
4.6 Exchange Sectors Come Later . . . 27
4.7 Lamport Dependency Skeleton . . . 28
4.8 Scope and Pitfalls . . . 28

**5 A Hilbert-Space Compiler Contract** **29**
5.1 Status and Local Anchors . . . 29
5.2 Mini North Star . . . 29
5.3 Input Record . . . 29
5.4 Output Record . . . 30
5.5 Observable-Algebra Policies . . . 30
5.6 Lamport Dependency Skeleton . . . 30
5.7 Boundary . . . 30

**6 Symmetry-Decorated Many-Particle Grammar** **31**
6.1 Sources and Dependencies . . . 31
6.2 Decorated Atoms . . . 31
6.3 Propagation Through OR and AND . . . 31
6.4 The Derived Full-Fock Representation . . . 31
6.5 Intertwiners and Symmetry-Preserving Compilation . . . 32
6.6 Lamport Dependency Skeleton . . . 32
6.7 Boundary . . . 32

**7 Sectors, Projections, Subspaces, and Quotients** **33**
7.1 Sources and Dependencies . . . 33
7.2 The Triangle . . . 33
7.3 Finite-Group Invariants . . . 33
7.4 Coinvariant Quotient . . . 34
7.5 Observable-Algebra Distinctions . . . 34
7.6 Checked Example . . . 34
7.7 Lamport Dependency Skeleton . . . 34
7.8 Boundary . . . 35

**8 Exchange Symmetry and Statistics as Sector Selection** **36**
8.1 Sources and Dependencies . . . 36
8.2 Two Symmetry Layers . . . 36
8.3 Irrep-Labeled Exchange Projectors . . . 36
8.4 Quotient Presentations . . . 37
8.5 Para and Anyonic Boundaries . . . 37
8.6 Lamport Dependency Skeleton . . . 37
8.7 Compiler Consequence . . . 37

**9 Scoping the Hilbert-Space Compiler** **38**
9.1 Status, Sources, and Dependencies . . . 38
9.2 Compile Judgement . . . 38
9.3 Surface Language for the First Compiler . . . 38
9.4 Constructor Rules . . . 38
9.5 Example 1: A Maybe Qubit . . . 39
9.6 Example 2: Two Distinguishable Qubits . . . 40
9.7 Example 3: Derived Tensor-Power Closure . . . 40
9.8 Example 4: Two-Qubit Exchange-Irrep Sectors . . . 40
9.9 Example 5: A Finite Invariant Quotient . . . 41
9.10 Example 6: A Required Compile Error . . . 41

9.11 Lamport Dependency Skeleton . . . 41
9.12 Acceptance Criteria and Boundary . . . 41

**10 Fibonacci Kinematics as a Fusion-Rule Sector** **42**
10.1 Status, Sources, and Dependencies . . . 42
10.2 One Physical Atom, Two Charge Labels . . . 42
10.3 Ambient Path Space and Admissibility Projector . . . 42
10.4 Dimension Recurrence . . . 43
10.5 Small Examples . . . 43
10.6 Compiler Consequence . . . 43
10.7 Lamport Dependency Skeleton . . . 44
10.8 Boundary . . . 44

**11 Why Lattice Symmetry Generators Are a Problem** **45**
11.1 Status, Sources, and Dependencies . . . 45
11.2 The Minimal Quantum-Mechanical Input . . . 45
11.3 What the Lattice Adds . . . 45
11.4 The Continuum Target . . . 45
11.5 Intuition to Be Checked . . . 46
11.6 Source Boundary . . . 46

**12 The Continuum Bulk Symmetry Target** **47**
12.1 Status, Sources, and Dependencies . . . 47
12.2 Group-Level Target . . . 47
12.3 Vector-Field Derivation . . . 47
12.4 Boost-Time Relation . . . 48
12.5 Boundary . . . 48

**13 A One-Dimensional Boost Produces a Momentum Current** **49**
13.1 Status, Sources, and Dependencies . . . 49
13.2 Setup . . . 49
13.3 Commutator Calculation . . . 49
13.4 Uniform Spacing . . . 50
13.5 Interpretation . . . 50
13.6 Boundary . . . 50

**14 Position-Dependent Bulk Generators** **51**
14.1 Status, Sources, and Dependencies . . . 51
14.2 Continuum Picture . . . 51
14.3 Scalar Ramps . . . 51
14.4 Boost Candidates . . . 51
14.5 Rotation Candidates . . . 52
14.6 Compiler Moral . . . 52
14.7 Boundary . . . 52

**15 Koo–Saleur as the Prototype** **53**
15.1 Status, Sources, and Dependencies . . . 53
15.2 Continuum Side . . . 53
15.3 Lattice Side . . . 53
15.4 Scaling Representation Matters . . . 53
15.5 What This Foreshadows . . . 54
15.6 Boundary . . . 54

**16 Acceptance Tests for Lattice Symmetry Candidates** **55**
16.1 Status, Sources, and Dependencies . . . 55
16.2 Local Finite-Scale Checks . . . 55
16.3 Algebraic Checks . . . 55
16.4 Scaling Checks . . . 55
16.5 Observable Covariance Checks . . . 55
16.6 Acceptance Payload . . . 56
16.7 Boundary . . . 56

**17 A Compiler Interface for Lattice Symmetry Candidates** **57**
17.1 Status, Sources, and Dependencies . . . 57
17.2 Input Contract . . . 57
17.3 Candidate Generation . . . 57
17.4 Output Contract . . . 57
17.5 Failure Modes . . . 58
17.6 Compiler Principle . . . 58

**18 Small Examples and the Lattice-Symmetry Research Queue** **59**
18.1 Status, Sources, and Dependencies . . . 59
18.2 Example 1: Black-Box Quantum System . . . 59
18.3 Example 2: Open Uniform Chain . . . 59
18.4 Example 3: Nonuniform Chain . . . 59
18.5 Example 4: Periodic Chain . . . 59
18.6 Example 5: Square Lattice . . . 60
18.7 Example 6: Koo–Saleur Modes . . . 60
18.8 Research Queue . . . 60
18.9 Landing Point . . . 60

**19 Why Galilean Symmetry Is a Different Lattice Target** **61**
19.1 Status, Sources, and Dependencies . . . 61
19.2 The New Target . . . 61
19.3 Immediate Consequence for Lattices . . . 61
19.4 Compiler Implication . . . 61
19.5 Why Projective Data Matter Earlier . . . 62
19.6 Boundary for CA-18–CA-22 . . . 62

**20 The Galilei Algebra as Newtonian Vector Fields** **63**
20.1 Status, Sources, and Dependencies . . . 63
20.2 Generators . . . 63
20.3 Translation and Boost Brackets . . . 63
20.4 Rotation on Vectors . . . 63

20.5 Rotation-Rotation Bracket . . . 64
20.6 Checked Representatives . . . 64
20.7 Boundary . . . 64

**21 Projective Galilean Symmetry and the Mass Central Term 65**
21.1 Status, Sources, and Dependencies . . . 65
21.2 Why This Belongs in the Grammar . . . 65
21.3 Canonical One-Particle Calculation . . . 65
21.4 Compatibility with Time Evolution . . . 65
21.5 Central Extension Versus Vector Fields . . . 66
21.6 Source Gap . . . 66
21.7 Compiler Payload . . . 66

**22 Lattice Galilean Boosts from Mass-Density First Moments 67**
22.1 Status, Sources, and Dependencies . . . 67
22.2 Input Data . . . 67
22.3 Discrete Continuity Assumption . . . 67
22.4 First-Moment Derivation . . . 67
22.5 Relation to Center of Mass . . . 68
22.6 What Is and Is Not Proven . . . 68

**23 Compiler Rules and Examples for Galilean Targets 69**
23.1 Status, Sources, and Dependencies . . . 69
23.2 Target Declaration . . . 69
23.3 Required Input Slots . . . 69
23.4 Generated Candidates . . . 69
23.5 Example 1: One Free Particle . . . 70
23.6 Example 2: Number-Conserving Open Chain . . . 70
23.7 Example 3: Hamiltonian Density Only . . . 70
23.8 Example 4: Spin Chain with No Particle Number . . . 70
23.9 Example 5: Direct Sum of Mass Sectors . . . 70
23.10Next Obligations . . . 71

**24 Gaussian Boson Lorentz Generator Roadmap 72**
24.1 Status, Sources, and Dependencies . . . 72
24.2 Problem Statement . . . 72
24.3 North Star . . . 72
24.4 Shard Roadmap . . . 73
24.5 Side Quests . . . 74
24.6 Numerical Verification Plan . . . 74
24.7 Acceptance Boundary . . . 74

**25 Gaussian Boson Symbol Calculus and the Boost-Time Residual 75**
25.1 Status, Sources, and Dependencies . . . 75
25.2 Coefficient Tier . . . 75
25.3 Nearest-Neighbour Klein-Gordon Example . . . 75
25.4 Momentum-Space Position . . . 76
25.5 Boost-Time Residual . . . 76
25.6 Lattice Residual . . . 76

25.7 Rotation Residual . . . 77
25.8 Boundary . . . 77

**26 Gaussian Boson Diagonalization and the One-Particle Symbol 78**
26.1 Status, Sources, and Dependencies . . . 78
26.2 Scalar Coefficient Tier . . . 78
26.3 Mode Decoupling . . . 78
26.4 Complex Structure and Fock Space . . . 79
26.5 What Is Not Yet Included . . . 79
26.6 Compiler Consequence . . . 79

**27 One-Particle Generator Algebra for Gaussian Bosons 80**
27.1 Status, Sources, and Dependencies . . . 80
27.2 Momentum Core . . . 80
27.3 Exact Translation and Rotation Brackets . . . 80
27.4 Boost-Momentum Bracket . . . 81
27.5 Boost-Time Residual . . . 81
27.6 Rotation-Hamiltonian Residual . . . 81
27.7 Boost-Boost Residual . . . 82
27.8 Second Quantization . . . 82

**28 Lorentz Residual Conditions on Gaussian Coefficients 83**
28.1 Status, Sources, and Dependencies . . . 83
28.2 Gradient Residual . . . 83
28.3 Low-Energy Hessian Condition . . . 83
28.4 Nearest-Neighbour Klein-Gordon . . . 84
28.5 Anisotropic Counterexample . . . 84
28.6 Hessian-Passing Instability Witness . . . 84
28.7 Doubler Counterexample . . . 85
28.8 Compiler Rule . . . 85

**29 Numerical Verification Suite for Gaussian Boson Examples 86**
29.1 Status, Sources, and Dependencies . . . 86
29.2 Why These Tests . . . 86
29.3 Numerical Tolerance Policy . . . 86
29.4 Structural Coefficient Validation . . . 87
29.5 Periodic Stiffness Matrix . . . 87
29.6 Implemented Example Systems . . . 87
29.7 Current Julia Surface . . . 89
29.8 Next Numerical Milestones . . . 90

**30 Gaussian Boson Real-Space Energy Density 91**
30.1 Status, Sources, and Dependencies . . . 91
30.2 Cell Volume and Notation . . . 91
30.3 Equal-Endpoint Bond Split . . . 91
30.4 Boundary Policy . . . 92
30.5 Vacuum-Energy Policy . . . 92
30.6 Improvement Policy . . . 92
30.7 Compiler Consequence . . . 92

**31 One-Dimensional Gaussian Boson Stress-Energy Candidates** **93**
31.1 Status, Sources, and Dependencies . . . 93
31.2 Continuum Motivation and Boundary . . . 93
31.3 Finite Open Chain Setup . . . 93
31.4 Candidate Dictionary . . . 93
31.5 Boundary, Subtraction, and Improvement . . . 94
31.6 Compiler Payload and Acceptance . . . 95

**32 Gaussian Current-Symbol Equivalence in One Dimension** **96**
32.1 Status, Sources, and Dependencies . . . 96
32.2 Real-Space Current Orientation . . . 96
32.3 Klein-Gordon Equality . . . 96
32.4 Julia Witness . . . 97
32.5 Open Obligations . . . 97

**33 Higher-Dimensional Gaussian Cell-Current Proposal** **98**
33.1 Status, Sources, and Dependencies . . . 98
33.2 Compiler Input: Oriented Support Data . . . 98
33.3 Energy Density and Energy Flux . . . 98
33.4 Momentum Densities and Stress Currents . . . 99
33.5 Boost and Symbol Diagnostics . . . 99
33.6 Rotation and Antisymmetric-Stress Residual . . . 99
33.7 Trace, Improvement, and Boundary Alternatives . . . 100

**34 First Gaussian Stress-Energy Numerical Suite** **101**
34.1 Status, Sources, and Dependencies . . . 101
34.2 Finite Open-Chain Continuity . . . 101
34.3 Current-Symbol Slope Residual . . . 101
34.4 What This Falsifies . . . 102
34.5 Open Obligations . . . 102

**35 Qubit Nearest-Neighbour Symmetry Quest** **103**
35.1 Status, Sources, and Dependencies . . . 103
35.2 Input Tier . . . 103
35.3 Question . . . 103
35.4 Investigation Plan . . . 104
35.5 Immediate Rule-Outs . . . 104

**36 One-Dimensional Qubit Boost-Current Obstructions** **105**
36.1 Status, Sources, and Dependencies . . . 105
36.2 Adjacent Current in Pauli Coefficients . . . 105
36.3 Fully On-Site Terms . . . 105
36.4 Other Immediate Rejections . . . 106
36.5 The Next Residual . . . 106
36.6 Boundary . . . 107

**37 Two-Dimensional Qubit Plaquette Symmetry Diagnostics** **108**
37.1 Status, Sources, and Dependencies . . . 108
37.2 Edge-Density Input . . . 108
37.3 Patch Diagnostics . . . 108
37.4 Proposed Rejection Filters . . . 108
37.5 Rotation Is Not Free . . . 109
37.6 Rule-Out Classes . . . 109

**38 Qubit Nearest-Neighbour Witt/Virasoro Diagnostics** **110**
38.1 Status, Sources, and Dependencies . . . 110
38.2 Fourier Density and Current Modes . . . 110
38.3 First Failure Witness . . . 110
38.4 Chiral-Rank Diagnostic . . . 110
38.5 Compressed Witt Residuals . . . 111
38.6 Boundary . . . 111

**39 Qubit Local-Algebra Equation Framework** **112**
39.1 Status, Sources, and Dependencies . . . 112
39.2 Pauli Word Algebra . . . 112
39.3 Bulk Equality Modulo Divergence . . . 112
39.4 What "Necessary" Means Here . . . 113

**40 One-Dimensional Qubit Poincare Necessary Equations** **114**
40.1 Status, Sources, and Dependencies . . . 114
40.2 Input and Current . . . 114
40.3 Translation Conservation . . . 114
40.4 Boost-Translation Relation . . . 115
40.5 Boundary . . . 115

**41 Computer Algebra for One-Dimensional Qubit Residuals** **116**
41.1 Status, Sources, and Dependencies . . . 116
41.2 Operator and Coefficient Roundtrip . . . 116
41.3 Checked Residual Helpers . . . 116
41.4 Sentinels . . . 116
41.5 Current Gap . . . 117

**42 Two-Dimensional Qubit Poincare Equation Schema** **118**
42.1 Status, Sources, and Dependencies . . . 118
42.2 Edge Densities and Vertex Currents . . . 118
42.3 Divergence-Relaxed Residuals . . . 118
42.4 Necessary Bulk Equations . . . 119
42.5 Implementation Boundary . . . 119

**43 Qubit Witt/Virasoro Necessary Equations** **120**
43.1 Status, Sources, and Dependencies . . . 120
43.2 Exact Density-Mode Algebra . . . 120
43.3 Chiral Residuals . . . 120
43.4 Necessary Filter . . . 121

**44 Vacuum Moment Constraints for Qubit Residuals** **122**
44.1 Status, Sources, and Dependencies . . . 122
44.2 Finite Moment Data . . . 122
44.3 Zero Expectations . . . 122
44.4 GNS Relation Constraints . . . 122
44.5 Highest-Weight Caveat . . . 123

**45 SDP Exclusion Hierarchy for Qubit Hamiltonians** **124**
45.1 Status, Sources, and Dependencies . . . 124
45.2 Level Data . . . 124
45.3 SDP Constraints . . . 124
45.4 Exclusion Statement . . . 124
45.5 Non-Implications . . . 125

**46 Roadmap for Qubit Symmetry Exclusion** **126**
46.1 Status, Sources, and Dependencies . . . 126
46.2 What We Have Now . . . 126
46.3 Implementation Steps . . . 126
46.4 Acceptance for an Exclusion Claim . . . 126
46.5 Open Gaps . . . 127

**47 Qubit SDP Implementation Contract** **128**
47.1 Status, Sources, and Dependencies . . . 128
47.2 Package Split . . . 128
47.3 Verdicts . . . 128

**48 Pauli Word Moment Basis for Infinite Qubit Chains** **129**
48.1 Status, Sources, and Dependencies . . . 129
48.2 Positioned Words . . . 129
48.3 Translation Quotient . . . 129

**49 Moment Matrices and Positivity Constraints** **130**
49.1 Status, Sources, and Dependencies . . . 130
49.2 Affine Moment Matrix . . . 130
49.3 Tiny Feasibility Tests . . . 130

**50 Compiling Poincare Residuals into Moment Equations** **131**
50.1 Status, Sources, and Dependencies . . . 131
50.2 Algebraic Witnesses . . . 131
50.3 Moment Relations . . . 131
50.4 Important Boundary . . . 131

**51 SDP Levels and the Mosek Backend** **132**
51.1 Status, Sources, and Dependencies . . . 132
51.2 Level Data . . . 132
51.3 Solver Status . . . 132
51.4 Smoke Run . . . 132

**52 Sentinel Hamiltonians for the SDP Hierarchy** **133**
52.1 Status, Sources, and Dependencies . . . 133
52.2 Matrices . . . 133
52.3 Gate Outcomes . . . 133

**53 Candidate Hamiltonian Selection Roadmap** **134**
53.1 Status, Sources, and Dependencies . . . 134
53.2 Screening Order . . . 134
53.3 Next Implementation Steps . . . 134

**54 Qubit Candidate Scan Contract** **135**
54.1 Status, Sources, and Dependencies . . . 135
54.2 Contract . . . 135
54.3 Verdict Boundary . . . 135

**55 Pauli Coefficients for Scan Families** **136**
55.1 Status, Sources, and Dependencies . . . 136
55.2 Implemented Maps . . . 136
55.3 Synthetic Families . . . 136

**56 Locally Sourced Spin-Chain Inputs** **137**
56.1 Status, Sources, and Dependencies . . . 137
56.2 Allowed Physical Labels . . . 137
56.3 Excluded From Physical Labeling . . . 137

**57 Synthetic and Stress-Test Hamiltonian Grids** **138**
57.1 Status, Sources, and Dependencies . . . 138
57.2 Grids . . . 138
57.3 Purpose . . . 138

**58 Current, Conservation, and Boost Witness Gates** **139**
58.1 Status, Sources, and Dependencies . . . 139
58.2 Gates . . . 139
58.3 Interpretation . . . 139

**59 Optional SDP Levels for Scan Survivors** **140**
59.1 Status, Sources, and Dependencies . . . 140
59.2 Level Data . . . 140
59.3 First-Run Boundary . . . 140

**60 Reproducible Scan Bundle Format** **141**
60.1 Status, Sources, and Dependencies . . . 141
60.2 Files . . . 141
60.3 Acceptance . . . 141

**61 First Candidate Scan Results** **142**
61.1 Status, Sources, and Dependencies . . . 142
61.2 Headline Table . . . 142
61.3 Notable Points . . . 142

**62 From Scan Failures to Scaling Work** **143**
62.1 Status, Sources, and Dependencies . . . 143
62.2 Queue . . . 143
62.3 SDP Upgrade . . . 143

**63 The Categorical Borchers–Uhlmann Pipeline: a Programme Refinement** **144**
63.1 Status, Sources, and Dependencies . . . 144
63.2 Refined North Star . . . 144
63.3 Five Stages . . . 144
63.4 Two Load-Bearing Constraints . . . 145
63.5 Functoriality and Fidelity Assessment . . . 146
63.6 Boundary . . . 146

**64 All-Level Moment Feasibility Gives a Qubit Reference State** **147**
64.1 Status, Sources, and Dependencies . . . 147
64.2 Canonical Full-Window Levels . . . 147
64.3 Finite Moments Are Finite States . . . 147
64.4 Compactness . . . 148
64.5 Residual Relations in the GNS Representation . . . 148
64.6 Exact Alternative . . . 148
64.7 Non-Implications . . . 149
64.8 Boundary . . . 149

**65 Quantified Relaxations of the Qubit Symmetry Gates** **150**
65.1 Status, Sources, and Dependencies . . . 150
65.2 Coefficient Norm and Gauge . . . 150
65.3 Algebraic Profiles . . . 150
65.4 State-Level Residuals . . . 151
65.5 Scaling Reading . . . 151
65.6 Sentinel Expectations . . . 151
65.7 Boundary . . . 151

**66 Anyonic Word Algebra from a Site Object** **152**
66.1 Status, Sources, and Dependencies . . . 152
66.2 Site Object First . . . 152
66.3 Placed Matrix Blocks . . . 152
66.4 Inclusions, Shift, and AF Limit . . . 153
66.5 The BU-Style Free Pre-Layer . . . 153
66.6 Defect Algebra Is a Different Layer . . . 154
66.7 Functoriality Target . . . 154
66.8 Lamport Dependency Skeleton . . . 154
66.9 Boundary . . . 154

**67 Anyonic States and Variable-$N$ Fock Space** **155**
67.1 Status, Sources, and Dependencies . . . 155
67.2 The Finite Hard-Core Carrier . . . 155
67.3 Action of the Word Algebra . . . 155
67.4 Birth and Death Operators . . . 156
67.5 Hollands' Dense Chain as a Corner . . . 156

67.6 Fibonacci Test Targets . . . 157
67.7 Lamport Dependency Skeleton . . . 157
67.8 Boundary . . . 158

**68 Refinement-Map Requirements and Obstructions 159**
68.1 Status, Sources, and Dependencies . . . 159
68.2 Strict OAR Requirements . . . 159
68.3 Fallback and Soft Relaxations . . . 159
68.4 The No-Go Mechanism as a Checklist . . . 159
68.5 Where Fixed Anyon Number Enters the Braiding RG . . . 160
68.6 Boundary . . . 160

**69 Refinement Maps for the Variable-$N$ Anyonic Word Algebra 161**
69.1 Status, Sources, and Dependencies . . . 161
69.2 Vacuum Insertion . . . 161
69.3 Algebra Map . . . 161
69.4 State Consistency . . . 161
69.5 Density Reading . . . 162
69.6 Virasoro Compatibility Target . . . 162
69.7 Dead Prior Art: the Deficit/Pair-Creation Precedent . . . 162
69.8 Fibonacci $H_2 \to H_4$ Test . . . 162
69.9 Comparison and Boundary . . . 163

**70 The Dilute Temperley–Lieb Algebra and the Anyonic Word Algebra 164**
70.1 Status, Sources, and Dependencies . . . 164
70.2 The Dilute Algebra . . . 164
70.3 The Evaluation Map . . . 164
70.4 The Obstruction: No Surjection for Fibonacci . . . 165
70.5 Consequences for the Programme . . . 165
70.6 Boundary . . . 165

**71 The Dilute Koo–Saleur Ansatz 166**
71.1 Status, Sources, and Dependencies . . . 166
71.2 What the Dense Template Requires . . . 166
71.3 The Dilute Ansatz . . . 166
71.4 Two Sourced Critical Anchors . . . 166
71.5 Proof Obligations . . . 167
71.6 Validation Plan (CA-71 Tier) . . . 167
71.7 Boundary . . . 167

**72 The Refinement Placement Category and Exact Composition 168**
72.1 Status, Sources, and Dependencies . . . 168
72.2 Placement Injections and Bare Refinements . . . 168
72.3 Exact Composition . . . 168
72.4 Cellwise Refinement and Cell-Shift Compatibility . . . 169
72.5 The Forest-Functor Reading . . . 169
72.6 The Ore Dichotomy and Fallback Homes . . . 169
72.7 Julia Invariants . . . 170
72.8 Boundary . . . 170

**73 GNS Descent and the Corner Calculus of Placement Refinements** **171**
73.1 Status, Sources, and Dependencies . . . 171
73.2 Setting . . . 171
73.3 The Corner Calculus . . . 171
73.4 The GNS Descent Lemma . . . 171
73.5 Residual Descent: the Contract for the Categorical Residual Set . . . 172
73.6 The Unital CP Completion and Its Exact Defect . . . 172
73.7 The BU-Level Square . . . 173
73.8 Julia Invariants . . . 173
73.9 Boundary . . . 173

**74 The Categorical Residual Set on the Anyonic Word Tower** **174**
74.1 Status, Sources, and Dependencies . . . 174
74.2 Setting . . . 174
74.3 The Ported Local Residual Calculus . . . 174
74.4 Open-Window Mode Calculus . . . 174
74.5 Witt/Virasoro Residuals with Two Central Constants . . . 175
74.6 The Categorical Residual Set . . . 175
74.7 Exact Transport and the Stretched Dichotomy . . . 175
74.8 Julia Invariants . . . 176
74.9 Boundary . . . 176

**75 The Jones–Wenzl Kernel Decision for the Dilute Evaluation Map** **178**
75.1 Status, Sources, and Dependencies . . . 178
75.2 Setting and the Decision Data . . . 178
75.3 Parity-Refined Multiplicities . . . 178
75.4 The Kernel Theorem . . . 179
75.5 The Upgraded Conjecture . . . 179
75.6 Julia Invariants . . . 180
75.7 Gram-Rank Deferral and Boundary . . . 180

**76 Integration Map: One Model-Independent Pipeline** **181**
76.1 Status, Sources, and Dependencies . . . 181
76.2 The Pipeline as Typed Layers . . . 181
76.3 Unwritten Links . . . 183
76.4 One Acceptance Contract . . . 183
76.5 Status of the Six Explorations . . . 184
76.6 Decisions Taken (2026-10-22): the Layering Principle . . . 185
76.7 Examples (Instances Only) . . . 185
76.8 Boundary . . . 186

**77 Corner-to-Wilson Bridge: Refinement Meets Dynamics** **187**
77.1 Status, Sources, and Dependencies . . . 187
77.2 Setting and Notation . . . 187
77.3 Corrections to CA-72 and CA-73 . . . 187
77.4 Corner Weight to Wilson Drift . . . 188
77.5 Dynamics: Vacuum Residual and Gap . . . 189
77.6 Polar Heat Refinement . . . 190

77.7 The Dichotomy: CA-75 Link (b3) . . . 191

**78 Pointed Carriers: Occupation Grading, Dual-Pair Blocking, and Corner Qualifications 192**
78.1 Status, Sources, and Dependencies . . . 192
78.2 Setting and the Two Towers . . . 192
78.3 Occupation Grading and Parity . . . 192
78.4 Dual-Pair Blocking Is Decidable . . . 193
78.5 Vacuum Insertion, States, and Qualifications of CA-71–CA-73 . . . 194
78.6 The Configuration Carrier . . . 194
78.7 Conjecture, with Caveat . . . 195
78.8 What This Does Not Claim . . . 195
78.9 Examples (calibration only) . . . 196

**79 Finite Selection Equations for a Categorical Nearest-Neighbour Hamiltonian 197**
79.1 Status, Sources, and Dependencies . . . 197
79.2 Setting and the Coefficient Compiler . . . 197
79.3 Blocking, Leakage, and Metric Response . . . 197
79.4 Additive Yang–Baxter as a Coefficient System . . . 198
79.5 Defect Symmetry . . . 199
79.6 Interfaces to WT-1 and VR-1 . . . 200
79.7 The Sufficiency Conjecture and Its Attackable Core . . . 200

**80 Koo–Saleur Tiles: Spectral-Shell Certificate and Refinement Defect 202**
80.1 Status, Sources, and Dependencies . . . 202
80.2 The KS Prescription Is a Tile Prescription . . . 202
80.3 A Spectral-Shell Certificate for VR-1 . . . 202
80.4 KS Modes under Categorical Refinement . . . 204
80.5 Extended Fields Are Declared Data . . . 204
80.6 Headline Conjecture and First Attackable Lemma . . . 205

**81 Exact Lattice Cells, Temporal Reflection Positivity, and a Two-Sided Fidelity Certificate 206**
81.1 Status, Sources, and Dependencies . . . 206
81.2 Setting and Notation . . . 206
81.3 Exact Cells and Sewing . . . 206
81.4 Temporal Reflection Positivity . . . 207
81.5 Defect Operators Do Not Certify Fidelity . . . 207
81.6 The Two-Sided Certificate . . . 208
81.7 What This Does Not Claim . . . 209

**82 The CFT Machine and Its Quantitative OAR Contract 210**
82.1 Sources, Dependencies, and Status . . . 210
82.2 The Actual Construction Target . . . 210
82.3 A Quantitative Wilson-State Theorem . . . 210
82.4 The Generator Contract and Its Boundary . . . 211

**83 A Supported Ising Modular-Data Compiler** **212**
83.1 Sources, Dependencies, and Scope . . . 212
83.2 Exact Input and Concrete Output . . . 212
83.3 The Local Observable Theory . . . 212
83.4 Symmetry and Modular Fidelity . . . 213

**84 Coset Lattice Stress Generators with Exact Core Stabilization** **214**
84.1 Sources, Dependencies, and Scope . . . 214
84.2 An Explicit Microscopic Polynomial . . . 214
84.3 Exact Stabilization on Free-Energy Cores . . . 214
84.4 Exact Witnesses and the Remaining Interface . . . 215

**85 Coset Positivity and a Scalar-Augmented OAR Tower** **216**
85.1 Sources, Dependencies, and Scope . . . 216
85.2 The Raw Finite Zero Mode Is Not Positive . . . 216
85.3 Computable Vacuum-Descendant Spaces . . . 216
85.4 An Independent Scalar Gives Genuine Unital OAR . . . 217
85.5 What the Construction Does Not Identify . . . 217

**86 Smooth Coset Symmetry Limits and Local Reconstruction** **218**
86.1 Sources and Scope . . . 218
86.2 Three Cutoffs and an Ambient Operator Theorem . . . 218
86.3 Local Fields in the Descendant OAR Representation . . . 219
86.4 The Local Fusion-Space Question . . . 219

**87 String-Net Histories as Trotter Expansions: the Spacetime Dictionary** **220**
87.1 Status, Sources, and Dependencies . . . 220
87.2 The Levin–Wen Rules and What They Fix . . . 220
87.3 The Trotter Dictionary . . . 220
87.4 Dense Calibration . . . 221
87.5 Dilute Calibration . . . 221
87.6 What the Calibration Says About the Principle . . . 222
87.7 Beyond Loops: the Fibonacci Net Gas (Proposal) . . . 222

**88 The Net Case: the Dilute Fibonacci Chain Is the Golden Chain Blocked** **224**
88.1 Status, Sources, and Dependencies . . . 224
88.2 Blocking: When Dilute Is Dense in Disguise . . . 224
88.3 The Temperley–Lieb Generators in Tile Form . . . 224
88.4 The Isotropic Net Gas Is the $Q = \varphi + 1$ Potts Model . . . 225
88.5 What This Does to the Principle . . . 226

**89 The Layered Target Grid: Setting, Input Data, and Target Rungs** **227**
89.1 Status, Sources, and Dependencies . . . 227
89.2 Three Axes . . . 227
89.3 Existing Results on the Grid . . . 228
89.4 The Six Headline Conjectures, Restated . . . 229
89.5 Round-2 Verdicts (2026-10-22) . . . 231
89.6 How to Read a Statement under the Layering Principle . . . 231
89.7 What This Does Not Claim . . . 232

89.8 Examples (Instances Only) . . . 232

**90 Ward-to-Koo–Saleur Transfer: Graph-Norm Interfaces, Derivative Matching and the Contact Cocycle 233**
90.1 Status, Sources, and Dependencies . . . 233
90.2 Setting and the Conjecture as Posed . . . 233
90.3 Graph-Norm Calculus and the Choice of Control . . . 234
90.4 Transfer of A2 and A4: Theorem 89.1(i) . . . 234
90.5 A3 from Differentiated Matching: Theorem 89.1(ii) . . . 235
90.6 The Contact Cocycle: Theorem 89.1(iii) . . . 235
90.7 The First Gap and the Next Lemma . . . 236
90.8 What This Does Not Claim . . . 236

**91 Local Unital Lifts from Relative Half-Braidings 238**
91.1 Status, Sources, and Dependencies . . . 238
91.2 Setting and the Transport Datum . . . 238
91.3 The One-Gap Lift . . . 239
91.4 The Dyadic Lift . . . 239
91.5 Wilson Consequences . . . 240
91.6 Non-Braided Input: Finite Equations and the One-Gap Obstruction . . . 240
91.7 Circuit Depth and Dressing . . . 241
91.8 Integration Status and the Next Lemma . . . 242
91.9 What This Does Not Claim . . . 242
91.10Examples (Instances Only) . . . 242

**92 Parity, Mixed Koo–Saleur Brackets, and the Chiral/Full Delineation 243**
92.1 Status, Sources, and Dependencies . . . 243
92.2 Two Corrections Recorded First . . . 243
92.3 Setting . . . 243
92.4 Parity: Anomaly Balance without Convergence . . . 244
92.5 Mixed Brackets Reduce to One Zero-Mode Commutator . . . 244
92.6 The Delineation . . . 245
92.7 Finite Ingredients for Parity Breaking . . . 246
92.8 What This Says about Rule 17's Chirality Layer . . . 246
92.9 What This Does Not Claim . . . 247

**93 Spatial Seams, Reflection Pairings and a Finite Osterwalder–Schrader Lemma 248**
93.1 Status, Sources, and Dependencies . . . 248
93.2 Setting and the Seam Pairing . . . 248
93.3 The Bent Kernel and Single-Tile Spatial RP . . . 249
93.4 The Linear History: an Exact Schur Criterion . . . 249
93.5 Negative Results and Pairing Dependence . . . 250
93.6 A Finite Osterwalder–Schrader Lemma (Category-Free) . . . 250
93.7 Two Directions: What Fails as Posed and What Survives . . . 251
93.8 The First Gap: Reflection Coherence . . . 251
93.9 What This Does Not Claim . . . 252
93.10Examples (Instances Only) . . . 252

**94 Normal Limit Expectations and the Repaired Two-Sided Fidelity Certificate 253**
94.1 Status, Sources, and Dependencies . . . 253
94.2 Scoped Notation . . . 253
94.3 The Normal Limit Expectation . . . 253
94.4 Sharpness of the Hypotheses . . . 254
94.5 The Two-Sided Fidelity Theorem at its Lowest Layer . . . 255
94.6 The Full-Setting Corollary and a False Reading . . . 255
94.7 What a Lattice Construction Must Supply . . . 256
94.8 What This Does Not Claim . . . 256

**95 Number-Changing Collision Dynamics on the Pointed Carrier: a Tuned Continuum Theorem and Three Obstructions 258**
95.1 Status, Sources, and Dependencies . . . 258
95.2 Setting and Shard-Local Conventions . . . 258
95.3 Fixed Densities Fail, and the Corrected Scaling . . . 259
95.4 The Closed Collision Form . . . 259
95.5 The Tuned Continuum Theorem . . . 260
95.6 Obstructions, Naturality, and Range . . . 261
95.7 Next Lemma: Contact Trichotomy . . . 262
95.8 What This Does Not Claim . . . 262
95.9 Examples (Instances Only) . . . 263

**96 Probe-Word Ward Lifting: Uniform Bounds, the Geometric Gap, and a False Converse 264**
96.1 Status, Sources, and Dependencies . . . 264
96.2 Physical Probe Bounds and the Conditional Geometric Interface . . . 264
96.3 Two Obstructions and the Precise Analytic Converse . . . 265
96.4 Spectral Calibration and the First Categorical Input . . . 266
96.5 Effect on other shards . . . 267
96.6 What This Does Not Claim . . . 267
96.7 Examples (Instances Only) . . . 268
96.8 Single Next Target . . . 268

**97 Blocked Dyadic Lifts: Finite-Power Transport and the Failure of Single-Site Centrality 269**
97.1 Status, Sources, and Dependencies . . . 269
97.2 Setting and the Refuted Equivalence . . . 269
97.3 Vacancy Shuffle and the General Block Construction . . . 269
97.4 Two Obstructions to the Proposed Characterization . . . 270
97.5 Bare Fusion-Data Existence and Naturality . . . 271
97.6 Effect on other shards . . . 272
97.7 What This Does Not Claim . . . 273
97.8 Examples (Instances Only) . . . 273
97.9 Next Target: Finite-Power Transport Extraction . . . 273

**98 What the Tower Remembers: Rooted AF Invariants and Marked Refinements 274**
98.1 Status, Sources, and Dependencies . . . 274
98.2 Setting and Rooted Invariant . . . 274
98.3 Trace and Classification . . . 275
98.4 Information Loss . . . 276
98.5 What the Refinement Record Adds . . . 277
98.6 Effect on other shards . . . 278
98.7 What This Does Not Claim . . . 278
98.8 Examples (Instances Only) . . . 278
98.9 Single Next Target: Intermediate-Record Faithfulness . . . 279

**99 Canonical Seams, Reflection Defects, and Conditional Spatial Positivity 280**
99.1 Status, Sources, and Dependencies . . . 280
99.2 Algebraic Seams and Oriented Cut Spaces . . . 280
99.3 Reflection Obstructions . . . 281
99.4 Conditional Histories and Finite Transfer Operators . . . 282
99.5 Effect on other shards . . . 283
99.6 What This Does Not Claim . . . 283
99.7 Examples (Instances Only) . . . 283
99.8 Next Target: Real-Self-Duality Two-Tile Reflection . . . 284

**100 Signed Contact Classification on the Pointed Carrier: Robin Limits, Dirichlet Limits and Full-Fock Boundaries 285**
100.1 Status, Sources, and Dependencies . . . 285
100.2 Exact Family and Classification . . . 285
100.3 Finite Contacts and the Weak Generator . . . 286
100.4 Both Divergent Signs and the Refuted Branch . . . 287
100.5 Full-Fock Results and General-Density Boundaries . . . 288
100.6 Effect on other shards . . . 289
100.7 What This Does Not Claim . . . 289
100.8 Examples (Instances Only) . . . 289
100.9 Single Next Target . . . 289

**101 Zero Modes and Lattice Chirality: Parity-Free KS Ledger Balance 290**
101.1 Status, Sources, and Dependencies . . . 290
101.2 Common-System Hypotheses and Transport . . . 290
101.3 Parity-Free Balance on the Common Domain . . . 291
101.4 Two Exact Obstructions to Automatic Zero-Mode Arguments . . . 292
101.5 Commutation Range, Cut Current and Spatial Support . . . 292
101.6 A Supplied-Target Construction Outside the KS Class . . . 293
101.7 Effect on other shards . . . 294
101.8 What This Does Not Claim . . . 294
101.9 Examples (Instances Only) . . . 294
101.10 Single Next Target . . . 294

**102 Ancilla Transport Lifts and the Failure of Finite-Power Extraction** **295**
102.1 Status, Sources, and Dependencies . . . 295
102.2 Setting and Ancilla Transport . . . 295
102.3 Gauge Invariants and Extraction Boundaries . . . 296
102.4 Finite Equations and Further Obstructions . . . 297
102.5 Effect on other shards . . . 297
102.6 What This Does Not Claim . . . 297
102.7 Examples (Instances Only) . . . 298
102.8 Single Next Target . . . 299

**103 The Interval/Dilation Record Forgets the Point** **300**
103.1 Status, Sources, and Dependencies . . . 300
103.2 The General Record and Cut Information . . . 300
103.3 An Exact Counterexample Family . . . 301
103.4 Named Points and the Remaining Marks . . . 302
103.5 Effect on other shards . . . 303
103.6 What This Does Not Claim . . . 303
103.7 Examples (Instances Only) . . . 303
103.8 Single Next Target: Pointed-Record Faithfulness . . . 304

# 1 Frontmatter, North Star, and Claim Status

This is a reproducible research lab book. Its guiding goal — the *north star* — is a single, concrete construction problem:

> Given a (unitary) fusion or modular tensor category $\mathcal{C}$, together with whatever additional data is needed (OPE coefficients / conformal data), construct a family of microscopic (lattice) models and their symmetry generators whose continuum limit *provably* yields a mathematically rigorous QFT/CFT that (i) realises the input category data and (ii) carries a full projective unitary representation of the symmetries.

The working belief is that this is achievable at least for *rational* CFTs, and perhaps for all QFTs. This is a programme, not a settled theory: every step toward the goal is a question, proposal, or hypothesis until a local source, a local derivation, or a reproducible run supports it.

## 1.1 The Two Success Criteria

A construction is only a candidate solution if it meets both, non-negotiably:

1. **Provability.** The continuum limit is established rigorously (an operator-algebraic / wavelet renormalisation limit with theorems, not a heuristic scaling argument).
2. **Fidelity.** The realised continuum content faithfully reproduces the input category data $\mathcal{C}$, and the symmetry representation is *full* and projective-unitary.

## 1.2 Claim Status

No claim is treated as established until it is marked *Source-local* or *Checked* and linked to a local source, derivation, test, or run artifact. The status labels are fixed in `CONVENTIONS.md`: *Question*, *Proposal*, *Source-local*, *Checked*, and *Rejected*. These labels are part of the scientific record; they separate programme-level conjecture from results.

## 1.3 Evidence Chain

Every substantive assertion should be traceable through:

1. a local source manifest under `references/` (the kept papers are the ground truth),
2. a convention entry in `CONVENTIONS.md`,
3. a derivation, script, test, or run bundle when computation is involved,
4. an index row in `INDEX.md`, and
5. a report shard in this lab book.

Missing links are allowed during exploration only when they are explicit.

# 2 Programme Map

This shard is a map of questions, not a list of results. The north star (§1 / CA-00) factors into a pipeline of stages; each stage is an open problem with its own conventions, sources, and checkable invariants.

## 2.1 The Pipeline

$$\underbrace{\mathcal{C}\ (+\,\text{OPE/conformal data})}_{\text{input}} \longrightarrow \underbrace{\text{microscopic models}}_{\text{lattice Hilbert space + Hamiltonian}} \longrightarrow \underbrace{\text{lattice symmetry generators}}_{\text{e.g. Koo–Saleur } L_n}$$

$$\longrightarrow \underbrace{\text{provable continuum limit}}_{\text{OAR / wavelets}} \longrightarrow \underbrace{\text{rigorous (C)FT}}_{\text{conformal net / VOA / Wightman}}$$

## 2.2 Stage 1 — Category Input

- Which categories are in initial scope: multiplicity-free unitary fusion categories (Fibonacci, Ising/TL), then modular tensor categories?
- What auxiliary data beyond $\mathcal{C}$ is genuinely required (OPE coefficients, central charge, conformal weights), and for which target CFTs?
- Which gauge and indexing conventions are fixed before any F/R-symbol is written (see `CONVENTIONS.md`)?

## 2.3 Stage 2 — Microscopic Models

- Which family realises $\mathcal{C}$ on the lattice: anyon chains, string-nets, or another construction — and how is the family indexed by scale?
- What is the lattice Hilbert space (fusion-tree basis) and the Hamiltonian, and which invariants pin them (fusion-rule consistency, pentagon, projector idempotence)?

## 2.4 Stage 3 — Lattice Symmetry Generators

- Which lattice operators approximate the continuum symmetry generators (Koo–Saleur lattice Virasoro $L_n$, $\bar{L}_n$, and kin)?
- What convergence statement is the target, and what is checkable at finite size (commutator algebra, central charge extraction)?

## 2.5 Stage 4 — Provable Continuum Limit

- Which rigorous renormalisation framework carries the limit (operator- algebraic renormalisation, wavelet / multi-scale entanglement)?
- What must be proved: existence of the limit, the limiting Hilbert space, and convergence of the symmetry generators to a projective unitary representation?

### 2.6 Stage 5 — Rigorous (C)FT Target

- What is the target object: a conformal net, a vertex operator algebra, a Wightman theory, or an Osterwalder–Schrader theory?
- What is the fidelity statement: that the target's superselection / representation category recovers $\mathcal{C}$?

### 2.7 First Decisions

The first technical session should not start with formulas. It should register the sources for one worked input category (Fibonacci or Ising/TL), fix the minimum conventions (F/R-symbol gauge, vacuum index, fusion-tree ordering), and identify the smallest invariant that can be checked end-to-end on one stage of the pipeline.

# 3 Many-Particle Hilbert Spaces: the AND/OR/Vacuum Grammar

## 3.1 Sources and Dependencies

This is a pre-topic foundation for the programme (see CA-01, the programme map): how to build the Hilbert space of an indefinite number of particles of arbitrary types. The guiding idea is that two logical connectives map cleanly onto two Hilbert-space constructions, and the no-particle system is the complex line.

This shard treats the *finite* grammar and its semantics; the rigorous *indefinite*-particle (Hilbert-completed) case is CA-03. It harmonises with, and re-proves nothing from, the finite "AND/OR many-particle grammar" developed by the author in the sibling project (notation $\mathcal{V}$, $\mathcal{V}_n$, AND $= \otimes$, OR $= \oplus$, $\mathbf{1} = \mathbb{C}$); see that project's shard `AQM-70`. Statistics (Bose / Fermi / para) are deferred throughout.

## 3.2 The Three Primitives

Let particle *types* be (separable, complex) Hilbert spaces; a type's space is the state space of one particle of that type. Three primitives generate composite systems:

- **OR = direct sum $\oplus$.** "Either a particle of type 1 or of type 2" is $\mathcal{H}_1 \oplus \mathcal{H}_2$.
- **AND = tensor product $\otimes$.** "A particle of type 1 *and* a particle of type 2" is $\mathcal{H}_1 \otimes \mathcal{H}_2$.
- **vacuum $= \mathbb{C}$.** The no-particle system is the complex line.

Crucially, this grammar fixes *indefinite particle number*; exchange symmetry (statistics) is a *separate, later* step, realised as a quotient or subspace (CA-03, §"Symmetries come later"). Keeping the two apart is the whole point.

## 3.3 Two Monoidal Structures and Their Units

On the category Hilb of complex Hilbert spaces (with the obvious unitary coherence isomorphisms) there are two symmetric monoidal structures:

$$(\text{Hilb},\ \oplus,\ \{0\}) \qquad \text{and} \qquad (\text{Hilb},\ \otimes,\ \mathbb{C}),$$

with the two units *distinct*:

$$A \oplus \{0\} \ \cong\ A \qquad (\oplus\text{-unit is the } \textit{zero space}\ \{0\}),$$

$$A \otimes \mathbb{C} \ \cong\ A \qquad (\otimes\text{-unit is the } \textit{vacuum}\ \mathbb{C}).$$

The two are linked by a *distributivity* natural isomorphism, which on Hilbert spaces is *unitary*:

$$\delta :\ A \otimes (B \oplus C) \ \xrightarrow{\cong}\ (A \otimes B) \oplus (A \otimes C), \qquad A \otimes \{0\} \ \cong\ \{0\}.$$

Together $(\text{Hilb}, \oplus, \{0\}, \otimes, \mathbb{C})$ is a symmetric *rig* (bimonoidal) category. Its coherence — that all diagrams built from associators, unitors, symmetries, and $\delta$ commute — is standard (Laplaza; Kelly) and is *not* re-proved here; downstream we use only the elementary, manifestly unitary associativity/unitor isomorphisms and the degree-0 identification $\mathbb{C} \otimes A \cong A$.

**Vacuum $= \mathbb{C}$, not $\{0\}$.** The vacuum is the $\otimes$-unit $\mathbb{C}$ (the empty tensor product, one particle's worth of "nothing"), a *one-dimensional* space carrying the unit vector $\Omega$. It is *not* the $\oplus$-unit $\{0\}$ (the empty direct sum, the impossible system). Empty $\otimes$ is $\mathbb{C}$; empty $\oplus$ is $\{0\}$. Conflating them is the first pitfall this grammar guards against (cf. CONVENTIONS.md (a), which fixes the vacuum *object* as $\mathbb{C}$).

### 3.4 The Grammar and its Carrier

Fix a set $\{K_x\}_{x\in X}$ of single-particle type spaces (the *atoms*). Grammar expressions are

$$E ::= 0 \;\mid\; \mathbf{1} \;\mid\; K \;\mid\; \mathrm{AND}(E,F) \;\mid\; \mathrm{OR}(E,F),$$

where 0 is the zero atom, $\mathbf{1}$ is the vacuum atom ($\mathbf{1} = \mathbb{C}$), and $K$ ranges over the atoms. The *carrier* $\mathcal{V}(E)$ is defined recursively:

$$\mathcal{V}(0) = \{0\}, \qquad \mathcal{V}(\mathbf{1}) = \mathbb{C}, \qquad \mathcal{V}(K) = K,$$
$$\mathcal{V}(\mathrm{AND}(E,F)) = \mathcal{V}(E) \otimes \mathcal{V}(F), \quad \mathcal{V}(\mathrm{OR}(E,F)) = \mathcal{V}(E) \oplus \mathcal{V}(F).$$

Thus AND is independent coexistence and OR is alternatives. Parenthesisation and order are part of the written expression; the rig coherence above guarantees that re-bracketing or re-ordering a *fixed* expression changes $\mathcal{V}(E)$ only by a canonical unitary isomorphism, so $\mathcal{V}$ is well defined up to canonical unitary isomorphism. (Morally $\mathcal{V}$ is the evaluation of the free symmetric rig category on $\{K_x\}$ into Hilb; we use this only as motivation, not as a load-bearing theorem.)

### 3.5 Particle-Number Grading

Declare each atom $K$ to be *one* particle and $\mathbf{1}$ to be *zero* particles. The carrier then carries a particle-number grading $\mathcal{V}_n$:

$$\mathcal{V}_0(\mathbf{1}) = \mathbb{C}, \quad \mathcal{V}_n(\mathbf{1}) = \{0\}\ (n \neq 0), \qquad \mathcal{V}_1(K) = K, \quad \mathcal{V}_n(K) = \{0\}\ (n \neq 1), \quad \mathcal{V}_n(0) = \{0\},$$
$$\mathcal{V}_n(\mathrm{OR}(E,F)) = \mathcal{V}_n(E) \oplus \mathcal{V}_n(F), \qquad \mathcal{V}_n(\mathrm{AND}(E,F)) = \bigoplus_{i+j=n} \mathcal{V}_i(E) \otimes \mathcal{V}_j(F).$$

For a *finite* expression the parse tree is finite, so only finitely many sectors are nonzero and $\mathcal{V}(E) = \bigoplus_{n\geq 0} \mathcal{V}_n(E)$ is a *finite* direct sum. The integer $n$ is a grammar count of one-particle atoms; it is not a dynamics, a statistics rule, or (yet) an indefinite-particle space.

**Worked atoms.** A one-particle "menu" over a finite type set $X$ is $P = \mathrm{OR}_{x\in X} K_x$, with $\mathcal{V}(P) = \bigoplus_{x\in X} K_x$ (the infinite, $\ell^2$-completed menu is CA-03). The expression $\mathbf{1}\,\mathrm{OR}\,K$ has carrier $\mathbb{C} \oplus K$ — a "maybe" particle (either zero or one particle of type $K$), as in the sibling shard's $\mathbb{C} \oplus \mathbb{C}^d$. Tensoring $N$ "maybe" atoms, $\mathrm{AND}_{j=1}^{N}(\mathbf{1}\,\mathrm{OR}\,K)$, has carrier $(\mathbb{C} \oplus K)^{\otimes N}$, whose grade-$n$ sector (by distributivity) is $\bigoplus_{|S|=n} \bigotimes_{x\in S} K$ over $n$-subsets of the $N$ slots — the finite "occupation" picture.

### 3.6 Lamport Dependency Skeleton

$D1$ : atoms $\{K_x\}_{x\in X}$ (single-particle Hilbert spaces); $\mathbf{1} = \mathbb{C}$ (vacuum), $0 = \{0\}$.
$D2$ : $E ::= 0 \mid \mathbf{1} \mid K \mid \mathrm{AND}(E,F) \mid \mathrm{OR}(E,F)$.
$D3$ : $\mathcal{V}$ : $\mathcal{V}(0) = \{0\}$, $\mathcal{V}(\mathbf{1}) = \mathbb{C}$, $\mathcal{V}(\mathrm{AND}) = \otimes$, $\mathcal{V}(\mathrm{OR}) = \oplus$.
$D4$ : $\mathcal{V}_0(\mathbf{1}) = \mathbb{C}$, $\mathcal{V}_1(K) = K$; $\mathcal{V}_n(\mathrm{OR}) = \oplus$, $\mathcal{V}_n(\mathrm{AND}) = \bigoplus_{i+j=n} \mathcal{V}_i \otimes \mathcal{V}_j$.
$D5$ : $(\mathrm{Hilb}, \oplus, \{0\}, \otimes, \mathbb{C})$ a symmetric rig category, with
$\delta : A \otimes (B \oplus C) \xrightarrow{\cong} (A \otimes B) \oplus (A \otimes C)$ unitary and natural.
$T1$ : (D5) coherence (Laplaza; Kelly) $\Rightarrow \mathcal{V}$ well defined up to canonical unitary iso.
$T2$ : (D4) finite $E$ : $\mathcal{V}(E) = \bigoplus_{n\geq 0} \mathcal{V}_n(E)$ a finite sum (finite particle number).

### 3.7 Scope and Deferral

Two extensions are deliberately out of this shard:

- **Indefinite particle number.** A genuine superposition over *unboundedly* many particle numbers has infinite support and does not live in the finite direct sum above; it requires the Hilbert ($\ell^2$) completion. That is the content of CA-03, and is the new rigorous part of this pre-topic.
- **Statistics.** AND and OR choose no exchange symmetry. Bosonic / fermionic / para sectors arise later, by acting with the permutation group on tensor slots and passing to an invariant subspace or quotient (CA-03). This grammar only sets up the carrier those symmetries act on.

**Pitfalls.** (i) The $\oplus$-unit is $\{0\}$; the $\otimes$-unit (the vacuum) is $\mathbb{C}$ — distinct objects. (ii) Distributivity is an *isomorphism*, not an equality; all sector identities below hold up to canonical unitary iso. (iii) Everything here is *finite*; do not silently read the grading as an indefinite-particle space.

# 4 Indefinite Particle Number: the Full Fock Completion

## 4.1 Sources and Dependencies

This shard completes CA-02's finite AND/OR/vacuum grammar to a genuine *indefinite*-particle Hilbert space. It depends on CA-02 (the two monoidal structures, the units $\{0\}$ and $\mathbb{C}$, the carrier $\mathcal{V}$ and grading $\mathcal{V}_n$). The named Fock construction is anchored locally:

The sibling project's finite grammar (`AQM-70`) is recovered as the finite-truncation of the object built here.

## 4.2 Two Direct Sums — the Load-Bearing Distinction

Let $\{W_n\}_{n\geq 0}$ be a countable family of Hilbert spaces.

- The **algebraic direct sum** $\bigoplus_n^{\mathrm{alg}} W_n$ consists of finite-support sequences $(w_n)$ ($w_n \in W_n$, $w_n = 0$ for all but finitely many $n$). It is a pre-Hilbert space, generally *incomplete*.
- The **Hilbert ($\ell^2$) direct sum** $\widehat{\bigoplus}_n W_n$ consists of sequences with $\sum_n \|w_n\|_{W_n}^2 < \infty$, with inner product $\langle (w_n), (v_n)\rangle = \sum_n \langle w_n, v_n\rangle_{W_n}$ (absolutely convergent by Cauchy–Schwarz). It is *complete*, hence a Hilbert space.

$\bigoplus_n^{\mathrm{alg}} W_n$ is a *dense* subspace of $\widehat{\bigoplus}_n W_n$, and the latter is its completion (standard functional analysis).

**Why indefinite particle number forces $\widehat{\bigoplus}$.** A state with nonzero amplitude in *unboundedly* many particle-number sectors (a genuine superposition over indefinitely many particles) has infinite support: it lies in $\widehat{\bigoplus}$ but not in $\bigoplus^{\mathrm{alg}}$. "Indefinite particle number" $\Leftrightarrow$ "not confined to finitely many sectors" $\Rightarrow$ the $\ell^2$ completion is *required*. This is exactly what CA-02's finite grammar lacks.

## 4.3 Completed Tensor Powers

For Hilbert spaces $H, K$, the *completed* (Hilbert) tensor product $H \mathbin{\widehat{\otimes}} K$ is the completion of the algebraic tensor product under the inner product determined by $\langle a\otimes b,\, c\otimes d\rangle = \langle a, c\rangle_H \langle b, d\rangle_K$ (well defined and positive definite). For finite-dimensional factors $\widehat{\otimes} = \otimes$; the completion bites only for infinite-dimensional one-particle spaces. Define the *completed tensor powers*

$$H^{\widehat{\otimes} 0} := \mathbb{C} \ \text{ (the vacuum sector)}, \qquad H^{\widehat{\otimes} n} := \underbrace{H \mathbin{\widehat{\otimes}} \cdots \mathbin{\widehat{\otimes}} H}_{n},$$

the bracketing immaterial up to the canonical unitary associativity isomorphism.

## 4.4 The Full Fock Space

**Definition.** For a one-particle Hilbert space $H$, the *full Fock space* is

$$\boxed{\mathfrak{F}(H) \;:=\; \widehat{\bigoplus_{n\geq 0}} H^{\widehat{\otimes} n}}, \qquad H^{\widehat{\otimes} 0} = \mathbb{C},$$

i.e. $\psi = (\psi_n)_{n\geq 0}$ with $\psi_n \in H^{\widehat{\otimes} n}$ and $\|\psi\|^2 = \sum_n \|\psi_n\|^2 < \infty$. The unit vector $\Omega = (1, 0, 0, \dots)$ in the degree-0 sector $\mathbb{C}$ is the *vacuum*. This is the *distinguishable* Fock space (the author's term in the local source `1910.10619`); in operator algebra it is also called the *free* or *Boltzmann* Fock

space (standard terminology, not anchored in a local source), namely the Hilbert completion of the tensor algebra $T(H) = \bigoplus_n H^{\otimes n}$. It is *not* the symmetric (bosonic) or antisymmetric (fermionic) Fock space; see "Exchange sectors come later" below.

**Compiler status.** In the compiler grammar this is not an additional primitive constructor. It is the Hilbert completion of the ordinary tensor-power expressions $\mathbf{1}, H, H^{\widehat{\otimes}2}, \ldots$: a countable OR over all finite AND powers. If a one-particle symmetry representation is present, CA-05 makes this same object a graded direct-sum representation.

**Theorem (Fock is a Hilbert space).** $\mathfrak{F}(H)$ is a Hilbert space; (i) the algebraic direct sum $\bigoplus_n^{\mathrm{alg}} H^{\widehat{\otimes}n}$ (the finite-particle vectors) is *dense*; (ii) distinct-$n$ sectors are *mutually orthogonal*, $\langle \psi, \varphi \rangle = \sum_n \langle \psi_n, \varphi_n \rangle$, so $H^{\widehat{\otimes}m} \perp H^{\widehat{\otimes}n}$ for $m \neq n$; (iii) on the degree-0 sector the inner product is the standard one on $\mathbb{C}$ and $\Omega$ is a unit vector. *Proof.* An instance of the $\ell^2$-direct-sum facts above with $W_n = H^{\widehat{\otimes}n}$: completeness, density of the algebraic sum, and summand orthogonality are exactly those statements. □

**The $1/N!$ caveat (a convention trap).** Schottenloher's inner product on $D = \bigoplus_N H_N$ weights sector $N$ by $1/N!$; that factor is a *symmetric*-Fock normalisation correcting the $N!$ overcounting of symmetric tensors. The full Fock space above uses the *plain* $\ell^2$ sum with *no* combinatorial weight. Do not copy the $1/N!$ into the distinguishable case.

### 4.5 The Grammar Generates the Indefinite-Particle Space

Let the one-particle menu over a countable type set $X$ be $P = \mathrm{OR}_{x \in X} K_x$, with carrier $\mathcal{V}(P) = \widehat{\bigoplus}_{x \in X} K_x$ (the $\ell^2$ direct sum; an ordinary finite direct sum when $X$ is finite). The indefinite-particle space over this menu is $\mathfrak{F}(\mathcal{V}(P))$. Expanding each completed tensor power by distributivity (CA-02's $\delta$, extended unitarily to the completions) gives the $(n, \text{type-word})$-graded decomposition

$$\mathcal{V}(P)^{\widehat{\otimes}n} \;\cong\; \widehat{\bigoplus}_{w \in X^n} K_w, \quad K_w := K_{x_1} \mathbin{\widehat{\otimes}} \cdots \mathbin{\widehat{\otimes}} K_{x_n}, \qquad \Longrightarrow \qquad \mathfrak{F}(\mathcal{V}(P)) \;\cong\; \widehat{\bigoplus}_{n \geq 0} \widehat{\bigoplus}_{w \in X^n} K_w.$$

Each of the $n$ tensor slots independently chooses a type $x_i \in X$ (two slots may choose the same type; nothing is symmetrised). The grading is $\mathcal{V}_n = \widehat{\bigoplus}_{w \in X^n} K_w$ ($n$ = particle number, $w \in X^n$ = ordered type-word) — the $\ell^2$-completed, all-$n$ analogue of CA-02's finite grade-$n$ sector. The sibling project's finite expressions ($P_X^{\mathrm{AND}\,m}$, the truncation $T_{\leq M}$) are precisely the finite-support / finite-$M$ restrictions of this object; that is the exact harmonisation: `AQM-70` is the $\bigoplus^{\mathrm{alg}}$/finite-$M$ truncation of $\mathfrak{F}$.

### 4.6 Exchange Sectors Come Later

The full Fock space already carries symmetry representations when $H$ does: internal symmetries act sectorwise by tensor powers (CA-05), and for each $n$ the permutation group $S_n$ acts unitarily on $H^{\widehat{\otimes}n}$ by permuting tensor slots. What is deferred is the *sector selection*. Bosonic and fermionic Fock spaces are the images of the per-sector orthogonal projectors

$$\Gamma_{\mathrm{s}}(H) = P_{\mathrm{sym}}\, \mathfrak{F}(H) \quad \text{(symmetric / Bose)}, \qquad \Gamma_{\mathrm{a}}(H) = P_{\mathrm{anti}}\, \mathfrak{F}(H) \quad \text{(antisymmetric / Fermi)},$$

with para-statistics the other Young-symmetry isotypic components. This is the "impose an equivalence relation under particle exchange" step. We define no projector and prove nothing here:

the construction of these symmetry quotients/subspaces, and the genuinely anyonic case (where standard Fock space is not the right object), are deferred to later shards.

### 4.7 Lamport Dependency Skeleton

$D6$ : $\bigoplus^{\mathrm{alg}}$(finite support) $\subset \widehat{\bigoplus}(\ell^2)$ : $\langle (w_n),(v_n)\rangle = \sum_n \langle w_n, v_n\rangle$.
$D7$ : completed tensor power $H^{\widehat{\otimes} n}$, $H^{\widehat{\otimes} 0} = \mathbb{C}$ (vacuum).
$D8$ : $\mathfrak{F}(H) := \widehat{\bigoplus}_{n\geq 0} H^{\widehat{\otimes} n}$ (full / distinguishable Fock); $\Omega = (1, 0, \dots)$.
$D9$ : $P = \mathrm{OR}_{x\in X} K_x$; $\quad \mathfrak{F}(\mathcal{V}(P)) = \widehat{\bigoplus}_{n\geq 0} \mathcal{V}(P)^{\widehat{\otimes} n}$.
$T3$ : (D6,D7) $\mathfrak{F}(H)$ Hilbert; $\bigoplus^{\mathrm{alg}}$ dense; $H^{\widehat{\otimes} m} \perp H^{\widehat{\otimes} n}$ $(m \neq n)$.
$T4$ : indefinite particle no. $\Leftrightarrow$ infinite support $\Rightarrow \widehat{\bigoplus}$ (not $\bigoplus^{\mathrm{alg}}$).
$T5$ : (D8,$\delta$) $\mathfrak{F}(\mathcal{V}(P)) \cong \widehat{\bigoplus}_{n\geq 0} \widehat{\bigoplus}_{w\in X^n} K_w$ ($(n$, type-word) grading).
$T6$ : $\Gamma_{\mathrm{s}} = P_{\mathrm{sym}}\mathfrak{F}$, $\Gamma_{\mathrm{a}} = P_{\mathrm{anti}}\mathfrak{F}$ deferred (statistics later).

### 4.8 Scope and Pitfalls

**In scope:** the indefinite-particle direction, i.e. countable OR (the Fock completion). **Out of scope:** countable AND (an *infinite tensor product*), which is analytically a different object (von Neumann's infinite tensor product, needing a stabilising reference sequence) — not developed here.

Pitfalls: (i) *algebraic* $\bigoplus^{\mathrm{alg}}$ (finite particle number, dense, incomplete) vs. *completed* $\widehat{\bigoplus}$ (indefinite, complete) — the distinction this shard exists to make. (ii) For infinite-dimensional one-particle $H$, the AND of CA-02 must mean the *completed* $\widehat{\otimes}$, not the algebraic $\otimes$. (iii) The $1/N!$ symmetric-Fock weight must not be imported into the distinguishable case. (iv) All sector identities hold on dense subspaces and extend by continuity; states are $\ell^2$-summable and cross-sector inner products vanish.

# 5 A Hilbert-Space Compiler Contract

## 5.1 Status and Local Anchors

**Status: Proposal / contract shard.** This shard introduces no new mathematical classification theorem. It records a design target for the pre-work sequence around CA-02 and CA-03: a typed grammar of many-particle expressions should be compilable to Hilbert-space data and observable-algebra data with explicit evidence status.

Local anchors are CA-02 for the finite AND/OR/vacuum grammar, CA-03 for the Hilbert direct-sum Fock completion, and `CONVENTIONS.md` entries (f) and (g) for symmetry and sector semantics. The first executable check for this sequence is the finite averaging projector in `test/runtests.jl`, backed by `src/CftAnyons.jl`.

## 5.2 Mini North Star

Proposal: build a *Hilbert-space compiler*. The user supplies an expression in a typed grammar, together with the atom table and required policies; the compiler returns the Hilbert carrier, its observable algebra policy, its symmetry representation data, and any sector projection or quotient presentation that was requested.

The motivating analogy is "regular expressions for quantum mechanics": a small syntax has compositional semantics. The analogy is only a prompt. Here the syntax is typed and semantics-bearing; every constructor must say what it does to Hilbert carriers, particle gradings, observable algebras, symmetries, and evidence metadata.

## 5.3 Input Record

A compiled expression is not just a string. It has an input record:

$$\mathsf{Input} = (\mathsf{Atoms},\ E,\ \mathsf{Completion},\ \mathsf{ObsPolicy},\ \mathsf{Symmetry},\ \mathsf{SectorPolicy}).$$

**Atoms.** A table of atoms. At minimum, each atom names a Hilbert space $K_x$. Later entries may include a group representation, a particle degree, a source locator, or category data.

**$E$.** A grammar expression built from $0, \mathbf{1}, K_x, \mathrm{AND}, \mathrm{OR}$, and later sector constructors. Fock-style all-$n$ notation is treated as a derived completion of tensor powers, not as a primitive grammar constructor.

**Completion.** Whether $\oplus, \otimes$ are finite Hilbert operations, Hilbert completions $\widehat{\bigoplus}, \widehat{\otimes}$, or a finite cutoff. CA-03 fixes the first infinite completion of the derived all-tensor-power construction used here.

**ObsPolicy.** The observable-algebra choice. There is no silent default: full $\mathcal{B}(H)$, block-preserving, tensor-local, symmetry-invariant, and compressed sector algebras are different outputs.

**Symmetry.** Optional unitary or projective-unitary representation data, handled according to CONVENTIONS.md (f).

**SectorPolicy.** Optional projection, subspace, or quotient data; see CONVENTIONS.md (g).

### 5.4 Output Record

The intended output is a typed record, not only a Hilbert-space dimension:

$$\mathsf{Compile}(E) = (H_E,\ \{H_{E,n}\}_{n\geq 0},\ \mathcal{A}_E,\ U_E,\ P_E,\ Q_E,\ \mathsf{Evidence}_E).$$

$H_E$. The Hilbert carrier. CA-02 and CA-03 define the first carrier semantics.

$\{H_{E,n}\}$. The particle-number grading when the atom table declares particle degrees.

$\mathcal{A}_E$. The selected observable algebra. This slot is mandatory once any observable claim is made.

$U_E$. The propagated symmetry representation, if symmetry data were supplied.

$P_E$. The selected sector projection, if a sector constructor was used.

$Q_E$. The quotient presentation, if one was requested and its closed relation subspace was named.

$\mathsf{Evidence}_E$. Source-local, checked, proposal, question, or rejected evidence tags for the pieces above.

### 5.5 Observable-Algebra Policies

The carrier alone does not determine the observable algebra relevant to a model. For example, $H \oplus K$ supports at least:

$$\mathcal{B}(H \oplus K), \qquad \mathcal{B}(H) \oplus \mathcal{B}(K), \qquad \text{off-diagonal creation/annihilation blocks between } H \text{ and } K.$$

All three are legitimate in different contexts; none is forced by the word "OR" alone. Similarly, $H \otimes K$ supports the full algebra $\mathcal{B}(H \otimes K)$ and the tensor-local subalgebra generated by $\mathcal{B}(H) \otimes I$ and $I \otimes \mathcal{B}(K)$.

Compiler rule: no report statement may infer an observable algebra merely from a carrier constructor. The statement must name the observable policy, or remain a carrier-only statement.

### 5.6 Lamport Dependency Skeleton

$D10$ : $\mathsf{Input} = (\mathsf{Atoms}, E, \mathsf{Completion}, \mathsf{ObsPolicy}, \mathsf{Symmetry}, \mathsf{SectorPolicy})$.
$D11$ : $\mathsf{Compile}(E) = (H_E, \{H_{E,n}\}, \mathcal{A}_E, U_E, P_E, Q_E, \mathsf{Evidence}_E)$.
$D12$ : $\mathsf{ObsPolicy} \in \{\text{carrier-only}, \mathcal{B}(H), \text{block}, \text{tensor-local}, \text{invariant}, \text{compressed}\}$.
$T7$ : A carrier claim needs $H_E$; an observable claim also needs $\mathcal{A}_E$.
$T8$ : The compiler analogy is a proposal, not a theorem of canonical quantization.

### 5.7 Boundary

This contract does not yet compile fusion-category labels, anyonic braid statistics, Hamiltonians, continuum limits, or CFT data. It only fixes the pre-work target: future shards should be able to say exactly which input record they compile and which output slots they have actually justified.

# 6 Symmetry-Decorated Many-Particle Grammar

## 6.1 Sources and Dependencies

This shard extends CA-02 and CA-03 by decorating atoms with symmetry data. It uses CONVENTIONS.md (f).

## 6.2 Decorated Atoms

Fix a topological group $G$. A symmetry-decorated atom is a pair

$$(K_x, U_x), \qquad U_x : G \to U(K_x),$$

where $K_x$ is the atomic Hilbert space and $U_x$ is a unitary representation. If the intended symmetry is projective, the pre-work convention is to choose a central extension

$$1 \to U(1) \to E \to G \to 1$$

and use an honest unitary representation $U_x : E \to U(K_x)$. The expression is then compiled over $E$, with a note that the induced physical symmetry is projective over $G$.

The vacuum atom is $(\mathbb{C}, \mathbf{1})$, where $\mathbf{1}(g)z = z$. The zero atom carries the unique representation on $\{0\}$.

## 6.3 Propagation Through OR and AND

If $E, F$ have compiled carriers and representations $(H_E, U_E)$ and $(H_F, U_F)$ for the same group or chosen central extension, define

$$H_{\mathrm{OR}(E,F)} = H_E \oplus H_F, \qquad U_{\mathrm{OR}(E,F)}(g) = U_E(g) \oplus U_F(g),$$

and

$$H_{\mathrm{AND}(E,F)} = H_E \mathbin{\widehat{\otimes}} H_F, \qquad U_{\mathrm{AND}(E,F)}(g) = U_E(g) \mathbin{\widehat{\otimes}} U_F(g).$$

For finite-dimensional factors, $\widehat{\otimes} = \otimes$. In infinite-dimensional Hilbert spaces, the completed tensor product is the one already required by CA-03.

These formulas are the Hilbert-space version of the monoidal structure on $\mathrm{Rep}^{\dagger}(G)$: direct sums are block actions and tensor products are diagonal actions. If two atoms use different groups, different central extensions, or incompatible projective multipliers, the compiler must stop and ask for a common symmetry object before forming a single $U_E$.

## 6.4 The Derived Full-Fock Representation

For an ordinary unitary representation $U : G \to U(H)$, the full-Fock object is itself a graded representation:

$$\mathfrak{F}(U)(g) = \bigoplus_{n \geq 0} U(g)^{\widehat{\otimes} n} \quad \text{on} \quad \mathfrak{F}(H) = \widehat{\bigoplus}_{n \geq 0} H^{\widehat{\otimes} n},$$

with $U(g)^{\widehat{\otimes} 0} = \mathrm{id}_{\mathbb{C}}$. This preserves the particle-number grading. This is not a new grammar constructor: it is the direct sum of the tensor-power representations already produced by OR and AND, completed as in CA-03. In this pre-work shard, no Shale–Stinespring-type implementability theorem is claimed for general canonical field algebras. Implementing a one-particle unitary on a chosen CAR/CCR observable algebra is a separate field-algebra question.

## 6.5 Intertwiners and Symmetry-Preserving Compilation

A morphism between decorated atoms or compiled expressions is symmetry-preserving only when it is an intertwiner:

$$TU_E(g) = U_F(g)T \quad \text{for all } g.$$

This matters for compiler optimisations. Rebracketing and distributivity isomorphisms used by CA-02 are acceptable in the decorated grammar only when they also intertwine the propagated actions. The standard Hilbert-space associators, unitors, symmetries, and distributivity maps do so for diagonal direct-sum and tensor-product actions.

## 6.6 Lamport Dependency Skeleton

$D13$ : decorated atom $(K_x, U_x),\ U_x : G \to U(K_x)$.
$D14$ : projective symmetry over $G$ is compiled via $U_x : E \to U(K_x)$ for a named central extension $E$.
$D15$ : $U_{E\,\mathrm{OR}\,F} = U_E \oplus U_F, \quad U_{E\,\mathrm{AND}\,F} = U_E \mathbin{\widehat{\otimes}} U_F$.
$D16$ : $\mathfrak{F}(U) = \widehat{\bigoplus}_{n\geq 0} U^{\widehat{\otimes} n}, \quad U^{\widehat{\otimes} 0} = 1_{\mathbb{C}}$.
$T9$ : The propagated symmetry preserves the CA-02/CA-03 particle grading.
$T10$ : A single compiled $U_E$ requires a common group or central extension.

## 6.7 Boundary

This shard does not yet impose invariance, select charge sectors, form quotient spaces, or define symmetry-invariant observable algebras. Those are sector operations, not mere symmetry propagation, and are treated in CA-06.

# 7 Sectors, Projections, Subspaces, and Quotients

## 7.1 Sources and Dependencies

This shard depends on CA-04 and CA-05 and fixes the first sector semantics for the Hilbert-space compiler. It uses CONVENTIONS.md (g).

## 7.2 The Triangle

Let $H$ be a Hilbert space. The compiler treats a sector first as an orthogonal projection

$$P = P^* = P^2 \in \mathcal{B}(H).$$

It may then expose either of the equivalent Hilbert-space presentations

$$H_P := \operatorname{Ran}(P) \subset H, \qquad H/\ker(P) = H/\operatorname{Ran}(1-P).$$

The map

$$[v] \in H/\ker(P) \longmapsto Pv \in \operatorname{Ran}(P)$$

is a well-defined unitary isomorphism when the quotient has the Hilbert quotient norm. Thus the safe slogan is not "symmetry equals quotient" but:

$$\text{projection} \iff \text{closed subspace} \iff \text{quotient by the named orthogonal complement.}$$

## 7.3 Finite-Group Invariants

Let $G$ be finite and $U : G \to U(H)$ a unitary representation. Define

$$P_G = \frac{1}{|G|} \sum_{g \in G} U(g).$$

Then $P_G$ is the orthogonal projection onto the invariant subspace

$$H^G = \{v \in H : U(g)v = v \ \forall g \in G\}.$$

*Proof.* Idempotence follows by reindexing:

$$P_G^2 = \frac{1}{|G|^2} \sum_{g,h} U(gh) = \frac{1}{|G|} \sum_{k} U(k) = P_G.$$

Self-adjointness follows from unitarity and $g \mapsto g^{-1}$:

$$P_G^* = \frac{1}{|G|} \sum_{g} U(g)^* = \frac{1}{|G|} \sum_{g} U(g^{-1}) = P_G.$$

For every $k$, $U(k)P_G = P_G$, so $\operatorname{Ran}(P_G) \subset H^G$. If $v \in H^G$, then $P_G v = v$, so $H^G \subset \operatorname{Ran}(P_G)$. □

### 7.4 Coinvariant Quotient

For the same finite action, let

$$R_G = \operatorname{span}\{U(g)v - v : g \in G,\ v \in H\}.$$

Then $R_G \subseteq \ker(P_G)$ because $P_G U(g) = P_G$. Conversely, if $w \in \ker(P_G)$, then

$$w = \frac{1}{|G|} \sum_{g \in G} \big(w - U(g)w\big) \in R_G.$$

Therefore $R_G = \ker(P_G) = (H^G)^\perp$, and the coinvariant quotient

$$H_G := H/R_G$$

is canonically unitarily isomorphic to the invariant subspace $H^G$. For compact non-finite groups the analogous statement requires a Haar-integral and analytic hypotheses; this shard records only the finite checked case.

### 7.5 Observable-Algebra Distinctions

There are three different symmetry-related algebras that the compiler must not identify silently:

$$\mathcal{B}(H), \qquad \mathcal{B}(H)^G := \{A : U(g)AU(g)^* = A\ \forall g\}, \qquad P_G\mathcal{B}(H)P_G \cong \mathcal{B}(H^G).$$

The first is the full algebra before imposing symmetry. The second is the algebra of invariant observables on the original carrier. The third is the compressed algebra acting on the invariant sector. A model may choose one of these, but the choice is an observable policy in the sense of CA-04, not a consequence of the word "symmetry" alone.

### 7.6 Checked Example

The current Julia check uses $G = C_2$ acting on $\mathbb{C}^2$ by the swap matrix $S$. The averaging projection is

$$P = \frac{1}{2}(I + S) = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}.$$

The test checks $P^* = P = P^2$, checks $I - P$ is the complementary orthogonal projection, verifies $P(I - P) = 0$, and verifies that $(1, 1)$ is fixed while $(1, -1)$ is killed. This is the first executable invariant for the sector compiler.

### 7.7 Lamport Dependency Skeleton

$D17$ : sector projection $P = P^* = P^2$; $\quad H_P = \operatorname{Ran}(P)$.
$D18$ : $H/\ker(P) \xrightarrow{[v]\mapsto Pv} \operatorname{Ran}(P)$.
$D19$ : $P_G = |G|^{-1} \sum_g U(g) \quad (G \text{ finite})$.
$D20$ : $R_G = \operatorname{span}\{U(g)v - v\}$.
$L1$ : $P_G = P_G^* = P_G^2$, $\quad \operatorname{Ran}(P_G) = H^G$.
$L2$ : $R_G = \ker(P_G) = (H^G)^\perp$, $\quad H/R_G \cong H^G$.
$T11$ : Projection, subspace, and quotient are equivalent only after the relation subspace is named.
$T12$ : $\mathcal{B}(H)^G$ and $P_G\mathcal{B}(H)P_G$ are different observable policies.

### 7.8 Boundary

The finite averaging theorem is not a statement about spontaneous symmetry breaking, gauge constraints, BRST cohomology, braid statistics, or superselection sectors in a continuum CFT. It is the first Hilbert-space sector rule needed by the compiler.

# 8 Exchange Symmetry and Statistics as Sector Selection

## 8.1 Sources and Dependencies

This shard discharges part of the deferral in CA-03: ordinary Bose/Fermi statistics are treated as choices of irreducible exchange-symmetry sectors on the full distinguishable Fock carrier. It depends on CA-05 for internal symmetry propagation and CA-06 for projection/subspace/quotient semantics.

## 8.2 Two Symmetry Layers

There are two distinct actions on many-particle carriers:

$$\text{internal symmetry } G, \qquad \text{exchange symmetry } S_n.$$

If $U : G \to U(H)$ is an internal one-particle representation, then on the $n$-particle distinguishable sector $H^{\widehat{\otimes} n}$ it acts diagonally:

$$U_n(g) = U(g)^{\widehat{\otimes} n}.$$

The permutation group $S_n$ acts by permuting tensor slots:

$$\Pi_n(\sigma)(v_1 \otimes \cdots \otimes v_n) = v_{\sigma^{-1}(1)} \otimes \cdots \otimes v_{\sigma^{-1}(n)}.$$

The two actions commute:

$$\Pi_n(\sigma) U_n(g) = U_n(g) \Pi_n(\sigma).$$

Thus internal charge sectors and exchange-statistics sectors can be imposed in either order in the ordinary tensor-slot setting.

## 8.3 Irrep-Labeled Exchange Projectors

The representation $\Pi_n : S_n \to U(H^{\widehat{\otimes} n})$ is the primary datum. Sector projections are supplied by irreducible representation data of $S_n$, not by specialised grammar constructors. For a one-dimensional unitary character $\chi : S_n \to U(1)$, CA-06 applied to the twisted representation $\overline{\chi}\,\Pi_n$ gives

$$P_{\chi,n} = \frac{1}{n!} \sum_{\sigma \in S_n} \overline{\chi(\sigma)}\, \Pi_n(\sigma),$$

whose range is the $\chi$-sector

$$\{ w \in H^{\widehat{\otimes} n} : \Pi_n(\sigma) w = \chi(\sigma) w \ \forall \sigma \in S_n \}.$$

For the trivial character 1, this is the symmetric projector:

$$P_{\mathrm{s},n} = \frac{1}{n!} \sum_{\sigma \in S_n} \Pi_n(\sigma),$$

and for the sign character sgn, this is the antisymmetric projector:

$$P_{\mathrm{a},n} = \frac{1}{n!} \sum_{\sigma \in S_n} \mathrm{sgn}(\sigma) \Pi_n(\sigma).$$

Their ranges are the symmetric and antisymmetric tensor powers:

$$\mathrm{Ran}(P_{\mathrm{s},n}) = H^{\widehat{\otimes}_s n}, \qquad \mathrm{Ran}(P_{\mathrm{a},n}) = \bigwedge^n H.$$

The corresponding Fock sectors are

$$\Gamma_{\mathrm{s}}(H) = \widehat{\bigoplus}_{n\geq 0} \mathrm{Ran}(P_{\mathrm{s},n}), \qquad \Gamma_{\mathrm{a}}(H) = \widehat{\bigoplus}_{n\geq 0} \mathrm{Ran}(P_{\mathrm{a},n}).$$

Thus "Bose" and "Fermi" name the trivial- and sign-irrep choices for ordinary exchange symmetry. The compiler should record the selected irrep/character and the resulting projection, not a special Bose/Fermi syntax node.

## 8.4 Quotient Presentations

For a one-dimensional character $\chi$, the sector can also be presented as a quotient of the full tensor algebra by the closed linear span of relations

$$\Pi_n(\sigma)v - \chi(\sigma)v.$$

The trivial character recovers symmetric relations; the sign character recovers antisymmetric signed relations. In the compiler, these quotient presentations are secondary: the primary sector object is the orthogonal projector, because it fixes the inherited Hilbert structure and the compressed observable algebra.

## 8.5 Para and Anyonic Boundaries

Higher-dimensional irreducible $S_n$-sectors require central idempotents or matrix-block projectors for the group algebra of $S_n$. This is exactly the same representation/irrep principle, but this shard records it as a proposal rather than a completed construction; a later shard must source the convention and projector normalization before using it.

Genuinely anyonic exchange is not an $S_n$-projection problem on ordinary tensor slots. It involves braid-group or categorical data and, in this project, must be handled together with the fusion-category conventions for associators, braiding, and $F/R$-symbols. No braid-statistics projector is defined here.

## 8.6 Lamport Dependency Skeleton

$D21$ : $U_n(g) = U(g)^{\widehat{\otimes}n}$ (internal symmetry on $H^{\widehat{\otimes}n}$).
$D22$ : $\Pi_n : S_n \to U(H^{\widehat{\otimes}n})$ permutes tensor slots.
$D23$ : $P_{\chi,n} = n!^{-1} \sum_\sigma \overline{\chi(\sigma)} \Pi_n(\sigma)$ ($\chi$ a one-dimensional $S_n$ character).
$L3$ : $\Pi_n(\sigma) U_n(g) = U_n(g) \Pi_n(\sigma)$.
$L4$ : $P_{\chi,n}$ is an orthogonal projection by CA-06 finite averaging.
$T13$ : $\Gamma_{\mathrm{s}}, \Gamma_{\mathrm{a}}$ are the trivial/sign irrep sectors inside full Fock.
$T14$ : Anyonic exchange remains deferred to categorical braid data.

## 8.7 Compiler Consequence

The compiler now has its first nontrivial sector family: given the exchange representation $\Pi_n : S_n \to U(H^{\widehat{\otimes}n})$, a selected irrep character $\chi$ supplies $P_{\chi,n}$, its range, its quotient relations, and the chosen compressed observable algebra. There is no primitive Bose, Fermi, or Fock grammar node in this account; Fock is the derived graded representation of CA-05, and Bose/Fermi are names for particular irrep choices of exchange symmetry.

# 9 Scoping the Hilbert-Space Compiler

## 9.1 Status, Sources, and Dependencies

**Status: specification shard.** This shard turns CA-04's contract into a first partial compiler specification. Its scope is the pre-work grammar of CA-02–CA-07: finite Hilbert carriers, derived all-tensor-power completions, unitary/projective symmetries via central extensions, and irrep/projection sectors.

## 9.2 Compile Judgement

The compiler is a *partial* map. A successful judgement has the form

$$\Gamma; \Pi \vdash E \Downarrow (H_E, \{H_{E,n}\}, \mathcal{A}_E, U_E, P_E, Q_E, \mathsf{Ev}_E),$$

where $\Gamma$ is the atom table and $\Pi$ is the policy table. A failed judgement has the form

$$\Gamma; \Pi \vdash E \Uparrow \mathsf{reason}.$$

The failure result is part of the specification: incompatible symmetry data, missing observable policy, or an ill-typed sector constructor must stop compilation.

The atom table entry for $x$ has at least a Hilbert space $K_x$, a degree $d_x \in \mathbb{N}$ (usually 1), evidence metadata, and optionally symmetry data $U_x : G \to U(K_x)$ or a named projective central extension as in CA-05.

## 9.3 Surface Language for the First Compiler

The first surface language is deliberately small:

$$E ::= 0 \mid \mathbf{1} \mid x \mid \mathrm{OR}(E, F) \mid \mathrm{AND}(E, F) \mid \mathrm{Sector}_{G,\chi}(E).$$

Here $G$ acts on the compiled carrier and $\chi$ is a one-dimensional unitary character; invariants are the case $\chi = 1$. Fock, finite truncation, symmetric sectors, and antisymmetric sectors are derived from OR/AND, representation propagation, and the generic sector operation.

For a one-particle atom $x$, define tensor-power expressions recursively by

$$x^{[0]} := \mathbf{1}, \qquad x^{[m+1]} := \mathrm{AND}(x^{[m]}, x).$$

Then

$$\mathsf{T}_{\le M}(x) := \mathrm{OR}_{m=0}^{M} x^{[m]}, \qquad \mathsf{T}_{\infty}(x) := \text{Hilbert completion of } \mathrm{OR}_{m\ge 0}\, x^{[m]}.$$

$\mathsf{T}_{\infty}(x)$ is the usual full/distinguishable Fock space $\widehat{\bigoplus}_{m\ge 0} K_x^{\widehat{\otimes} m}$, but as a derived graded representation.

## 9.4 Constructor Rules

The carrier, grading, and symmetry rules are recursive:

$$H_0 = \{0\}, \quad H_{\mathbf{1}} = \mathbb{C}, \quad H_x = K_x,$$

$$H_{\mathrm{OR}(E,F)} = H_E \oplus H_F, \qquad H_{\mathrm{AND}(E,F)} = H_E \mathbin{\widehat{\otimes}} H_F.$$

The grading rules are CA-02's:

$$H_{\mathrm{OR}(E,F),n} = H_{E,n} \oplus H_{F,n}, \qquad H_{\mathrm{AND}(E,F),n} = \bigoplus_{i+j=n} H_{E,i} \mathbin{\widehat{\otimes}} H_{F,j}.$$

If $U_E, U_F$ are representations of the same group or named central extension, the symmetry rules are

$$U_{\mathrm{OR}}(g) = U_E(g) \oplus U_F(g), \qquad U_{\mathrm{AND}}(g) = U_E(g) \mathbin{\widehat{\otimes}} U_F(g).$$

If the symmetry objects are not the same, the carrier may still compile, but the symmetry output is rejected unless the requested output was explicitly carrier-only.

For the derived tensor-power closures,

$$H_{\mathsf{T}_\infty(x)} = \widehat{\bigoplus}_{m\geq 0} K_x^{\widehat{\otimes} m}, \qquad H_{\mathsf{T}_{\leq M}(x)} = \bigoplus_{m=0}^{M} K_x^{\otimes m}.$$

For finite examples the compiler may report dimensions:

$$\begin{aligned} \dim \mathrm{OR} &= \dim H_E + \dim H_F, \\ \dim \mathrm{AND} &= (\dim H_E)(\dim H_F), \\ \dim \mathsf{T}_{\leq M}(x) &= \sum_{m=0}^{M} (\dim K_x)^m. \end{aligned}$$

These follow from choosing orthonormal bases: disjoint union gives direct-sum bases, ordered pairs give tensor-product bases, and truncation is a finite sum.

Sector constructors return projection witnesses supplied by representation data. For finite $G$, a unitary character $\chi : G \to U(1)$, and representation $U_E : G \to U(H_E)$, set

$$P_\chi = |G|^{-1} \sum_{g\in G} \overline{\chi(g)}\, U_E(g), \qquad H_{\mathrm{Sector}_{G,\chi}(E)} = \mathrm{Ran}(P_\chi).$$

The quotient relation is generated by $U_E(g)v - \chi(g)v$. The invariant sector is $\chi = 1$. For exchange sectors on $K^{\widehat{\otimes} n}$, $G = S_n$ and $U_E = \Pi_n$; the trivial and sign characters give

$$P_{\mathrm{s},n} = n!^{-1} \sum_{\sigma\in S_n} \Pi_n(\sigma), \qquad P_{\mathrm{a},n} = n!^{-1} \sum_{\sigma\in S_n} \mathrm{sgn}(\sigma)\Pi_n(\sigma).$$

The observable algebra is never inferred from these formulas. It is the policy slot $\Pi_{\mathrm{obs}}$: carrier-only, full, block, tensor-local, invariant, or compressed.

### 9.5 Example 1: A Maybe Qubit

Let $Q = \mathbb{C}^2$ with basis $e_0, e_1$. Compile

$$E_1 = \mathrm{OR}(\mathbf{1}, Q).$$

The rule gives

$$H_{E_1} = \mathbb{C} \oplus \mathbb{C}^2, \qquad H_{E_1,0} = \mathbb{C}, \quad H_{E_1,1} = \mathbb{C}^2, \quad \dim H_{E_1} = 3.$$

With full observable policy, $\mathcal{A}_{E_1} = \mathcal{B}(\mathbb{C} \oplus \mathbb{C}^2)$. With block policy,

$$\mathcal{A}_{E_1}^{\mathrm{block}} = \mathcal{B}(\mathbb{C}) \oplus \mathcal{B}(\mathbb{C}^2).$$

Thus the same carrier has two different compiled observable outputs.

### 9.6 Example 2: Two Distinguishable Qubits

Compile $E_2 = \mathrm{AND}(Q, Q)$. The carrier is

$$H_{E_2} = Q \otimes Q, \qquad \{e_0 \otimes e_0, e_0 \otimes e_1, e_1 \otimes e_0, e_1 \otimes e_1\} \text{ is a basis,}$$

so $\dim H_{E_2} = 4$, all in degree 2. The tensor-local policy is the subalgebra generated by

$$\mathcal{B}(Q) \otimes I, \qquad I \otimes \mathcal{B}(Q),$$

whereas the full policy is $\mathcal{B}(Q \otimes Q)$. The compiler records which one was requested; it does not identify them by syntax alone.

### 9.7 Example 3: Derived Tensor-Power Closure

Compile the macro-expanded expression

$$E_3 = \mathsf{T}_{\leq 2}(Q) = \mathrm{OR}(\mathbf{1}, \mathrm{OR}(Q, \mathrm{AND}(Q, Q))).$$

The output carrier is

$$H_{E_3} = \mathbb{C} \oplus Q \oplus (Q \otimes Q),$$

with sector dimensions

$$\dim H_0 = 1, \qquad \dim H_1 = 2, \qquad \dim H_2 = 4, \qquad \dim H_{E_3} = 1 + 2 + 4 = 7.$$

If $U : G \to U(Q)$ is supplied, symmetry propagation gives

$$U_{E_3}(g) = 1_{\mathbb{C}} \oplus U(g) \oplus (U(g) \otimes U(g)).$$

No Bose/Fermi relation has been imposed yet; this is still the distinguishable graded representation.

### 9.8 Example 4: Two-Qubit Exchange-Irrep Sectors

Let $S$ be the swap on $Q \otimes Q$. Since

$$S(e_0e_0) = e_0e_0, \quad S(e_1e_1) = e_1e_1, \quad S(e_0e_1) = e_1e_0, \quad S(e_1e_0) = e_0e_1,$$

where $e_ie_j$ abbreviates $e_i \otimes e_j$, the exchange representation of $S_2$ has two one-dimensional irrep characters, trivial and sign, supplying projectors with ranges:

$$P_{\mathrm{s}} = \frac{1}{2}(I + S), \qquad P_{\mathrm{a}} = \frac{1}{2}(I - S)$$

$$\mathrm{Ran}(P_{\mathrm{s}}) = \mathrm{span}\{e_0e_0,\ (e_0e_1 + e_1e_0)/\sqrt{2},\ e_1e_1\},$$
$$\mathrm{Ran}(P_{\mathrm{a}}) = \mathrm{span}\{(e_0e_1 - e_1e_0)/\sqrt{2}\}.$$

Thus the compiler reports dimensions 3 and 1 for the trivial- and sign-irrep sectors. The Julia test suite checks the projection, orthogonality, trace/dimension, and basis-vector claims for this example.

### 9.9 Example 5: A Finite Invariant Quotient

Let $C_2 = \{1, s\}$ act on $H = \mathbb{C}^2$ by the swap matrix. Then

$$P_{C_2} = \frac{1}{2}(I + S) = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}.$$

Its range is $\mathrm{span}\{(1,1)\}$, and its kernel is $\mathrm{span}\{(1,-1)\}$. The quotient presentation is therefore

$$\mathbb{C}^2/\,\mathrm{span}\{(1,-1)\} \cong \mathrm{span}\{(1,1)\}.$$

This is $\mathrm{Sector}_{C_2,1}(H)$, the trivial-irrep sector of the $C_2$ representation. The invariant observable algebra on the original carrier consists of the operators commuting with $S$, while the compressed algebra on the invariant line is one-dimensional. These are different observable policies even in this small example.

### 9.10 Example 6: A Required Compile Error

Let $A, B$ be atoms with projective symmetries over the same physical group $G$, but represented in the input by unrelated central extensions $E_A \to G$ and $E_B \to G$. The carrier of $\mathrm{OR}(A, B)$ is still $K_A \oplus K_B$. However, a request for one compiled symmetry representation must fail:

$$\Gamma; \Pi \vdash \mathrm{OR}(A, B) \Uparrow \text{“no common symmetry group or central extension specified”}.$$

This is not a missing feature; it is the strict semantics required by CA-05 and CONVENTIONS.md (f).

### 9.11 Lamport Dependency Skeleton

$D24$ : $\Gamma; \Pi \vdash E \Downarrow (H_E, \{H_{E,n}\}, \mathcal{A}_E, U_E, P_E, Q_E, \mathsf{Ev}_E)$ or $\Gamma; \Pi \vdash E \Uparrow \mathsf{reason}$.
$D25$ : $E ::= 0 \mid \mathbf{1} \mid x \mid \mathrm{OR} \mid \mathrm{AND} \mid \mathrm{Sector}_{G,\chi}$.
$D26$ : $\mathsf{T}_{\le M}(x) = \mathrm{OR}_{m=0}^{M} x^{[m]}, \quad H_{\mathsf{T}_{\le M}} = \bigoplus_{m=0}^{M} K_x^{\otimes m}$.
$D27$ : $P_\chi = |G|^{-1} \sum_g \overline{\chi(g)} U_E(g), \quad H_{\mathrm{Sector}_{G,\chi}} = \mathrm{Ran}(P_\chi)$.
$T15$ : Finite dimensions obey sum, product, and $\sum_{m=0}^{M} d^m$ by explicit basis constructions.
$T16$ : Irrep/character choices supply projection/subspace/quotient witnesses.
$T17$ : Compiler failure is a specified outcome when requested structure is ill typed or unsupported.

### 9.12 Acceptance Criteria and Boundary

Future implementation work must preserve these acceptance criteria: every constructor fills typed output slots; every nontrivial mathematical output has an evidence tag; every quotient names its relation subspace; every observable claim names its observable policy; and every executable example asserts an invariant, not merely successful execution.

This compiler does not yet parse fusion-category simples, anyonic braid data, Hamiltonians, lattice limits, conformal nets, or VOA data. Those enter only after their source conventions and invariants are fixed. CA-08 is the finite Hilbert-space kernel that later shards must extend without weakening the output record, the evidence discipline, or the failure semantics.

# 10 Fibonacci Kinematics as a Fusion-Rule Sector

## 10.1 Status, Sources, and Dependencies

**Status: checked kinematic shard.** This shard confirms the side conjecture: before braiding, Hamiltonians, or continuum limits, the Fibonacci fusion-path Hilbert spaces are a sector construction from one physical atom $\tau$. The vacuum $\mathbf{1}$ is still present as the tensor unit and as a charge label; it is not a second physical particle atom.

Dependencies: CA-02 for AND/OR/vacuum, CA-06 for projection/subspace semantics, CA-08 for compiler output slots, and CONVENTIONS.md (c) for the left-associated kinematic path basis.

## 10.2 One Physical Atom, Two Charge Labels

The physical input word is $n$ copies of one anyon type:

$$\tau, \tau, \ldots, \tau.$$

The charge labels used to describe intermediate fusion outcomes are

$$L = \{\mathbf{1}, \tau\}.$$

Thus the proposed slogan is correct only with this distinction:

$$\text{one physical atom } \tau \quad + \quad \text{charge labels } \{\mathbf{1}, \tau\} \quad + \quad \text{fusion-rule sector selection.}$$

The vacuum $\mathbf{1}$ is required because it is both the tensor unit and a possible total charge. It is not an additional non-vacuum particle species.

## 10.3 Ambient Path Space and Admissibility Projector

Use the left-associated path basis fixed in CONVENTIONS.md (c). For $n$ physical $\tau$-atoms and total charge $c \in \{\mathbf{1}, \tau\}$, define the ambient word space

$$W_{n,c} = \ell^2\{(x_0, \ldots, x_n) : x_i \in \{\mathbf{1}, \tau\},\ x_0 = \mathbf{1},\ x_n = c\}.$$

The basis vector $|x_0, \ldots, x_n\rangle$ records the cumulative charge after successively fusing the first $i$ copies of $\tau$.

The Fibonacci fusion rules needed here are

$$\mathbf{1} \otimes \tau = \tau, \qquad \tau \otimes \mathbf{1} = \tau, \qquad \tau \otimes \tau = \mathbf{1} \oplus \tau.$$

Let $N_{ab}^{c} \in \{0, 1\}$ be these multiplicity-free fusion coefficients. Define the diagonal operator

$$P_{n,c}|x_0, \ldots, x_n\rangle = \left(\prod_{i=1}^{n} N_{x_{i-1},\tau}^{x_i}\right)|x_0, \ldots, x_n\rangle.$$

Since every diagonal entry is 0 or 1, $P_{n,c} = P_{n,c}^{*} = P_{n,c}^{2}$. The kinematic Fibonacci Hilbert space is therefore the sector

$$H_{n,c}^{\mathrm{Fib}} := \mathrm{Ran}(P_{n,c}) \subseteq W_{n,c}.$$

This is exactly the CA-06 projection/subspace semantics, but the projector is not supplied by a group character. It is supplied by categorical fusion-rule admissibility.

## 10.4 Dimension Recurrence

Let

$$a_n = \dim H^{\mathrm{Fib}}_{n,\mathbf{1}}, \qquad b_n = \dim H^{\mathrm{Fib}}_{n,\tau}.$$

At $n = 0$, no physical $\tau$-atom has been fused, so

$$a_0 = 1, \qquad b_0 = 0.$$

Adding one more $\tau$ gives the transition equations

$$a_{n+1} = b_n, \qquad b_{n+1} = a_n + b_n.$$

The first equation says total charge $\mathbf{1}$ can only come from $\tau \otimes \tau \to \mathbf{1}$. The second says total charge $\tau$ can come from $\mathbf{1} \otimes \tau \to \tau$ or from $\tau \otimes \tau \to \tau$. Hence

$$b_{n+1} = b_n + b_{n-1}, \qquad b_0 = 0, \quad b_1 = 1.$$

Thus the total-$\tau$ sector dimensions are Fibonacci numbers:

$$0, 1, 1, 2, 3, 5, 8, \ldots.$$

The Julia test suite checks the pair sequence

$$(a_n, b_n) = (1,0), (0,1), (1,1), (1,2), (2,3), (3,5), (5,8), (8,13)$$

for $0 \le n \le 7$, including the recurrence.

## 10.5 Small Examples

**$n = 1$.** For total charge $\tau$, the ambient basis is $|\mathbf{1}, \tau\rangle$, and it is admissible. Thus $H^{\mathrm{Fib}}_{1,\tau} \cong \mathbb{C}$.

**$n = 2$.** The admissible paths are

$$|\mathbf{1}, \tau, \mathbf{1}\rangle \in H^{\mathrm{Fib}}_{2,\mathbf{1}}, \qquad |\mathbf{1}, \tau, \tau\rangle \in H^{\mathrm{Fib}}_{2,\tau}.$$

This is the kinematic content of $\tau \otimes \tau = \mathbf{1} \oplus \tau$.

**$n = 3$.** The total-$\tau$ ambient basis has two admissible paths:

$$|\mathbf{1}, \tau, \mathbf{1}, \tau\rangle, \qquad |\mathbf{1}, \tau, \tau, \tau\rangle.$$

The forbidden word $|\mathbf{1}, \mathbf{1}, \tau, \tau\rangle$, for example, is killed because $N^{\mathbf{1}}_{\mathbf{1},\tau} = 0$. Therefore $\dim H^{\mathrm{Fib}}_{3,\tau} = 2$.

## 10.6 Compiler Consequence

This is the first example where the Hilbert-space compiler wants category data. At the carrier level, it can compile the ambient finite word space $W_{n,c}$ and the diagonal sector projection $P_{n,c}$. But producing $P_{n,c}$ from the expression "$n$ copies of $\tau$" requires the fusion coefficients $N^c_{ab}$. That is already tensor-category input, not ordinary Hilbert-space grammar.

Moreover, the source makes clear that different fusion orders give different fusion-tree bases and are related by $F$-moves. Therefore a full Fibonacci compiler must eventually output not only

$$H^{\mathrm{Fib}}_{n,c}$$

but also the coherent recoupling data that identifies different bracketings: the $F$-symbols satisfying the pentagon equations, and later the $R$-symbols for braiding. The current shard is only kinematic: it builds the admissible sector and counts it.

## 10.7 Lamport Dependency Skeleton

$D28 : L = \{\mathbf{1}, \tau\}, \quad \mathbf{1} \otimes \tau = \tau, \quad \tau \otimes \tau = \mathbf{1} \oplus \tau.$
$D29 : W_{n,c} = \ell^2\{(x_0, \ldots, x_n) : x_0 = \mathbf{1},\ x_n = c,\ x_i \in L\}.$
$D30 : P_{n,c}|x\rangle = \prod_i N^{x_i}_{x_{i-1},\tau}|x\rangle.$
$D31 : H^{\mathrm{Fib}}_{n,c} = \mathrm{Ran}(P_{n,c}).$
$L5 : P_{n,c} = P^*_{n,c} = P^2_{n,c}.$
$T18 : \dim H^{\mathrm{Fib}}_{n,\tau}$ obeys $b_{n+1} = b_n + b_{n-1},\ b_0 = 0,\ b_1 = 1.$
$T19 :$ Full Fibonacci compilation needs $F$-coherent tensor-category data.

## 10.8 Boundary

This shard does not fix the Fibonacci $F$-matrix gauge, any $R$-symbols, braid representations, golden-chain Hamiltonians, or a continuum limit. It only realises the fusion-path Hilbert spaces kinematically as admissibility sectors.

# 11 Why Lattice Symmetry Generators Are a Problem

## 11.1 Status, Sources, and Dependencies

**Status: proposal shard with sourced boundary.** The claim that a lattice boost should look like a first moment of the Hamiltonian density is an intuition until a local source, derivation, or test is attached. This shard sets the boundary for CA-10–CA-17.

Dependencies: CA-05 for symmetry as unitary/projective-unitary representation, CA-08 for compiler output slots, and CONVENTIONS.md (h) for lattice candidate generator signs.

## 11.2 The Minimal Quantum-Mechanical Input

An unadorned quantum system gives a Hilbert space and, once dynamics is chosen, a self-adjoint Hamiltonian. By Stone's theorem this is the same kind of datum as a strongly continuous unitary representation of the additive group $\mathbb{R}$:

$$U(t) = \exp(-itH).$$

Thus the one symmetry forced by the bare dynamical presentation is time translation. Everything beyond this requires extra structure: spatial coordinates, locality, a metric, a lattice graph, or a continuum target.

## 11.3 What the Lattice Adds

A lattice Hamiltonian is not merely a self-adjoint operator. It usually comes as a decomposition

$$H = \sum_j h_j$$

where each $h_j$ is local near a site, bond, cell, or other geometric support. The cited lattice-fermion source uses exactly this pattern: a Hamiltonian is a sum of self-adjoint finite-support terms, together with an observable algebra whose generators are intended as discretisations of continuum fields.

This decomposition is extra data. Two different decompositions of the same operator $H$ can lead to different candidate densities and hence different candidate symmetry generators. Therefore the compiler target is not

$$\text{Hilbert space plus one operator } H,$$

but rather

$$(\text{Hilbert space},\ \text{lattice geometry},\ \{h_j\},\ \text{scaling map}).$$

## 11.4 The Continuum Target

If the intended continuum theory is Lorentz-invariant in $d+1$ dimensions, the target symmetry is not just $\mathbb{R}$-time translation. It is a projective-unitary representation of the Poincare group, lifted as necessary to a unitary representation of a covering group. Its translation subgroup has self-adjoint generators

$$P_0 = H, \qquad P_1, \ldots, P_d,$$

where the $P_a$ are momenta.

**Motivating question.** If the microscopic input only supplies a lattice Hamiltonian density $H = \sum_j h_j$, how do we guess the lattice analogues of

$$P_a, \qquad M_{ab} \text{ rotations}, \qquad M_{0a} \text{ boosts?}$$

## 11.5 Intuition to Be Checked

**Intuition 1:** a rotation is a position-dependent spatial translation. In the plane, the vector field for an infinitesimal rotation about the origin is

$$v(x, y) = (-y, x).$$

Thus if we can identify local momentum densities, rotations should be spatial first moments of those densities.

**Intuition 2:** a boost is a position-dependent time translation. Since time translation is generated by energy, the one-dimensional boost candidate is a first moment of the energy density,

$$K_x \ \propto \ \sum_j x_j h_j.$$

**Intuition 3:** the commutator of a boost with time translation should produce spatial translation, up to sign and normalization. On the lattice this suggests that

$$i[H, K_x]$$

is a candidate momentum generator. CA-12 derives the nearest-neighbour formula from this ansatz and records the sign convention explicitly.

## 11.6 Source Boundary

Weinberg QFT Vol. 1 is now registered with targeted OCR anchors, and the original Koo–Saleur source is now registered from arXiv with source-TeX and PDF text anchors. The present shard still cites Schottenloher for continuum symmetry definitions, OAR/wavelet sources for scaling-limit translation recovery, and the local lattice-fermion source for rigorous free-fermion Koo–Saleur convergence. CA-14 cites the original Koo–Saleur paper for the source-local lattice Virasoro prototype.

# 12 The Continuum Bulk Symmetry Target

## 12.1 Status, Sources, and Dependencies

**Status: sourced definitions plus checked local derivation.** The Poincare group and the projective-unitary implementation are sourced. The commutator table below is a local vector-field derivation from the sourced metric-preserving description, and its signs are pinned by a Julia check.

Dependencies: CA-05, CA-10, and CONVENTIONS.md (h), (k).

## 12.2 Group-Level Target

For a $D = d + 1$ dimensional Lorentzian target, Schottenloher describes the Poincare group as the semidirect product of the Lorentz group and translations:

$$P = SO_0(1, d) \ltimes \mathbb{R}^{d+1},$$

with the usual replacement by the relevant covering group when the quantum symmetry is implemented projectively. The translation subgroup is abelian and has commuting self-adjoint generators $P_\mu$. In the source convention,

$$P_0 = H, \qquad P_a \text{ are spatial momenta } (a = 1, \ldots, d).$$

For Euclidean two-dimensional CFT language, the Euclidean motion group is generated by rotations and translations. This gives the Wick-rotated analogue: rotations and translations are the geometric target before scale/conformal extensions are added.

## 12.3 Vector-Field Derivation

Let $\eta = \mathrm{diag}(1, -1, \ldots, -1)$, and lower indices by $x_\mu = \eta_{\mu\nu} x^\nu$. Use vector fields

$$P_\mu = \partial_\mu, \qquad M_{\mu\nu} = x_\mu \partial_\nu - x_\nu \partial_\mu, \qquad M_{\mu\nu} = -M_{\nu\mu}.$$

The source gives the metric-preserving Lie algebra condition $\omega^T \eta + \eta\omega = 0$. The $M_{\mu\nu}$ form the corresponding coordinate vector fields, while $P_\mu$ are translations.

Compute the Lie bracket of vector fields. Since the $P_\mu$ have constant coefficients,

$$[P_\mu, P_\nu] = 0.$$

For the mixed bracket,

$$\begin{aligned}[M_{\mu\nu}, P_\rho] &= -\partial_\rho(x_\mu)\partial_\nu + \partial_\rho(x_\nu)\partial_\mu \\ &= \eta_{\nu\rho} P_\mu - \eta_{\mu\rho} P_\nu.\end{aligned}$$

For Lorentz-Lorentz brackets,

$$\begin{aligned}[M_{\mu\nu}, M_{\rho\sigma}] = \eta_{\nu\rho} M_{\mu\sigma} - \eta_{\mu\rho} M_{\nu\sigma} \\ + \eta_{\mu\sigma} M_{\nu\rho} - \eta_{\nu\sigma} M_{\mu\rho}.\end{aligned}$$

The Julia check evaluates representative brackets using this convention. For example in $3 + 1$ dimensions,

$$[M_{01}, P_0] = -P_1, \qquad [M_{01}, P_1] = -P_0.$$

### 12.4 Boost-Time Relation

With $K_a := M_{0a}$, the vector-field convention gives

$$[K_a, P_0] = -P_a.$$

Equivalently, $[P_0, K_a] = P_a$. The passage from this vector-field statement to self-adjoint commutator notation is the separate sign convention in CONVENTIONS.md (k): Stone generators use $U_X(t) = \exp(-itA_X)$, the spacetime flow is implemented passively, and the resulting target is

$$i[H, K_a] = P_a, \qquad i[P_b, K_a] = \delta_{ab} H.$$

Spatial rotations carry the additional convention $J_{ab} = -A_{M_{ab}}$, which is why the Gaussian one-particle table has $i[K_a, K_b] = J_{ab}$ even though the active vector fields obey $[M_{0a}, M_{0b}] = -M_{ab}$. CA-12 keeps finite-lattice signs explicit under CONVENTIONS.md (h).

### 12.5 Boundary

This shard does not claim that a finite lattice has an exact Poincare representation. It only identifies the continuum algebraic target whose relations a sequence of lattice candidates should approximate after scaling.

# 13 A One-Dimensional Boost Produces a Momentum Current

## 13.1 Status, Sources, and Dependencies

**Status: checked local derivation.** The identity in this shard is not quoted from a source. It is derived from the local Hamiltonian-density ansatz and checked by a small coefficient invariant.

Dependencies: CA-10, CA-11, and CONVENTIONS.md (h).

## 13.2 Setup

Let a finite open one-dimensional lattice have sites or bonds labelled $j = 1, \ldots, N$, real positions $x_j$, and a self-adjoint local-density decomposition

$$H = \sum_{j=1}^{N} h_j.$$

Assume the density terms commute at distance greater than one:

$$[h_j, h_k] = 0 \qquad \text{if } |j-k| > 1.$$

This is the finite-range simplification appropriate for the first derivation; longer-range densities only add longer-range edge terms.

Define the position-weighted energy moment

$$K_x = \sum_{j=1}^{N} x_j h_j.$$

This is the lattice boost candidate in the sense of CA-10: a boost is treated as a position-dependent time translation, and time translation is generated by energy density.

## 13.3 Commutator Calculation

Compute

$$\begin{aligned} i[H, K_x] &= i\left[\sum_j h_j, \sum_k x_k h_k\right] \\ &= i\sum_{j,k} x_k [h_j, h_k]. \end{aligned}$$

The terms with $j = k$ vanish, and by the nearest-neighbour assumption only pairs $(j, j+1)$ survive. For each adjacent pair,

$$i x_{j+1}[h_j, h_{j+1}] + i x_j [h_{j+1}, h_j] = i(x_{j+1} - x_j)[h_j, h_{j+1}].$$

Hence

$$i[H, K_x] = \sum_{j=1}^{N-1} (x_{j+1} - x_j)\, i[h_j, h_{j+1}].$$

Each $i[h_j, h_{j+1}]$ is self-adjoint because $h_j, h_{j+1}$ are self-adjoint.

### 13.4 Uniform Spacing

If $x_j = aj$, then $x_{j+1} - x_j = a$ and

$$i[H, K_x] = a \sum_{j=1}^{N-1} i[h_j, h_{j+1}].$$

For unit spacing this gives the candidate bulk momentum

$$P_{\text{bulk}}^{\text{lat}} := \sum_{j=1}^{N-1} i[h_j, h_{j+1}],$$

with endpoint terms deliberately visible. Periodic chains require an extra boundary convention because the position function $j$ is not single-valued on a circle.

### 13.5 Interpretation

The continuum bracket in CA-11 says that boost plus time translation gives a spatial translation, up to sign convention. The lattice calculation says that the same operation applied to the first energy moment produces the commutator of neighbouring energy densities. Therefore $i[h_j, h_{j+1}]$ is not guessed from thin air: it is the local current forced by the first-moment ansatz.

### 13.6 Boundary

This is not yet a theorem that $P_{\text{bulk}}^{\text{lat}}$ converges to continuum momentum. The missing ingredients are a scaling map, a domain for the candidate operators, a state/sector sequence, and convergence of the commutator relations. CA-15 turns these into acceptance checks.

# 14 Position-Dependent Bulk Generators

## 14.1 Status, Sources, and Dependencies

**Status: proposal shard with local derivation pattern.** The sourced continuum content is geometric: vector fields, translations, and rotations. The higher-dimensional lattice formulas are compiler targets, not established convergence claims, and this shard does not define a stress-energy tensor or a lattice momentum density.

Dependencies: CA-10–CA-12 and CONVENTIONS.md (h).

## 14.2 Continuum Picture

Translations are flows of constant vector fields. In the Euclidean plane, the infinitesimal motion

$$X(z) = c + i\theta z$$

contains both translations ($c$) and rotations ($i\theta z$). Written in real coordinates, the rotation part is the position-dependent vector field

$$(x, y) \mapsto (-\theta y, \theta x).$$

This is the precise continuum content of the slogan:

$$\text{rotation} = \text{position-dependent translation}.$$

## 14.3 Scalar Ramps

Suppose lattice density terms are attached to positions $r \in \Lambda \subset \mathbb{R}^d$. For a scalar function $f : \Lambda \to \mathbb{R}$, define

$$K_f := \sum_{r \in \Lambda} f(r) h_r.$$

The same derivation as CA-12 gives, for commuting non-overlapping densities,

$$i[H, K_f] = \sum_{\{r,s\}} (f(s) - f(r))\, i[h_r, h_s],$$

where the sum is over unordered overlapping pairs after a choice of orientation. This is a discrete-gradient response. Linear ramps

$$f_a(r) = r^a$$

are therefore the first candidates for spatial momentum components.

## 14.4 Boost Candidates

Boosts can be attempted directly from energy density:

$$K_a^{\text{lat}} := \sum_{r \in \Lambda} r^a h_r.$$

Then $i[H, K_a^{\text{lat}}]$ gives the $a$-direction current candidate. This is the higher-dimensional version of the one-dimensional derivation, but it depends on the chosen density supports and orientations.

## 14.5 Rotation Candidates

Rotations require one more layer. In continuum notation,

$$J_{ab} = \int (x_a p_b(x) - x_b p_a(x))\, dx.$$

Here this formula is only a continuum mnemonic for why rotations should involve an angular-momentum density. The Schottenloher anchors above support the underlying source vector fields and rotations, not a QFT stress-energy tensor, not a momentum-density formula, and not a lattice dictionary for $p_a(x)$. Thus an actual lattice rotation candidate must be routed through a later momentum-density/current convention rather than built directly from $h_r$. Once such a convention is fixed, the provisional compiler shape is:

$$J^{\mathrm{lat}}_{ab} \approx \sum_r \big(r_a p^{\mathrm{lat}}_b(r) - r_b p^{\mathrm{lat}}_a(r)\big),$$

where $p^{\mathrm{lat}}_a$ must come with its own source, derivation, or checked continuity-equation evidence.

## 14.6 Compiler Moral

The lattice Hilbert-space compiler cannot infer these operators from the carrier alone. It needs a lattice geometry, a local Hamiltonian-density decomposition, and a target vector-field algebra. The output is not an exact symmetry by default; it is a candidate list with proof obligations.

## 14.7 Boundary

No stress-energy convention is fixed here. No cell-orientation convention is fixed here. No periodic wrapping convention is fixed here. No higher-dimensional convergence theorem is fixed here. Those choices must be made in a later shard before any numerical or categorical lattice model uses these candidates as checked symmetries.

# 15 Koo–Saleur as the Prototype

## 15.1 Status, Sources, and Dependencies

**Status: source-local prototype plus later rigorous precedent.** The original Koo–Saleur paper is registered here and supports the lattice Virasoro prototype as a conjectural construction with numerical and analytic checks for XXZ/RSOS/Temperley–Lieb settings. This shard keeps that separate from the local lattice-fermion/OAR treatment, which states and proves a free-fermion Koo–Saleur convergence result.

Dependencies: CA-10–CA-13.

## 15.2 Continuum Side

In a two-dimensional CFT on the circle, the Hamiltonian is expressed as an integral of a Hamiltonian density,

$$H = \int h(x)\, dx.$$

The source identifies $h(x)$ as a local quantum field built from the chiral and anti-chiral energy-momentum tensors $T, \bar{T}$. Its Fourier modes $H_k$ are linear combinations of the Virasoro generators $L_k, \bar{L}_k$.

Thus in this special setting the desired spacetime symmetries are not abstract: they are generated by modes of the stress tensor.

## 15.3 Lattice Side

The Koo–Saleur move is to imitate the continuum construction using the lattice Hamiltonian density. From a lattice density

$$h^{(N)} : \Lambda_N \to \mathcal{A}_N$$

one forms Fourier modes $H_k^{(N)}$. In the original Temperley–Lieb presentation, the Hamiltonian-limit density is represented by ground-state-subtracted $e_j - e_\infty$ terms, while the second stress-tensor component is represented by nearest-neighbour commutators $[e_j, e_{j+1}]$. The lattice Virasoro candidates $l_n, \bar{l}_{-n}$ are therefore Fourier sums of

$$(e_j - e_\infty) \;\pm\; \text{constant} \cdot [e_j, e_{j+1}],$$

with the $c/24$ zero-mode shift. The later free-fermion/OAR Definition 4.1 uses the same structural pattern: density modes plus commutator corrections against the lattice Hamiltonian mode $H_0^{(N)}$ produce lattice analogues of $L_k$ and $\bar{L}_k$.

The important structural point for our programme is:

$$\text{lattice symmetry generator} = \text{weighted Hamiltonian density} + \text{commutator correction}.$$

This is the same motif as CA-12, but now with Fourier weights and a Virasoro target rather than a first moment and a Poincare translation target.

## 15.4 Scaling Representation Matters

The original source already builds in ground-state subtraction, normal ordering, and a double-limit process; it also warns that finite-$L$ lattice quantities do not close as the Virasoro algebra and that a scaling limit of a commutator is not generally the commutator of scaling limits. The

later rigorous free-fermion/OAR source similarly makes convergence depend on the correct scaling-limit representation and on subtracting ground-state expectation values, corresponding to normal ordering. Therefore the compiler must not output only formal sums. It must also output:

state sequence,
renormalisation/scaling maps,
vacuum subtraction policy,
operator topology target.

### 15.5 What This Foreshadows

Koo–Saleur is a prototype for the lattice-symmetry compiler:

$$\{h_j\} \longmapsto \{H_k^{(N)}, L_k^{(N)}, \bar{L}_k^{(N)}\} \longrightarrow \{L_k, \bar{L}_k\}.$$

The first arrow is algebraic and finite-scale. The second is analytic and requires a scaling theorem.

### 15.6 Boundary

The original Koo–Saleur source supports the prototype as a conjectural and numerically checked construction for XXZ/RSOS/Temperley–Lieb lattice models; it does not prove a general rigorous convergence theorem for arbitrary microscopic models. The convergence theorem cited here is still the later free-fermion/OAR result, and arbitrary anyon chains remain outside both sourced claims.

# 16 Acceptance Tests for Lattice Symmetry Candidates

## 16.1 Status, Sources, and Dependencies

**Status: specification shard.** This shard does not prove convergence for new models. It states the checks required before a candidate generator is promoted from proposal to checked continuum symmetry data.

Dependencies: CA-10–CA-14.

## 16.2 Local Finite-Scale Checks

At fixed lattice size $N$, a candidate generator $G_N$ must first pass finite operator checks:

$$G_N = G_N^*, \qquad G_N \text{ is built from named local density terms,} \qquad \operatorname{supp}(G_N) \text{ policy is explicit.}$$

For unbounded thermodynamic-limit candidates, the finite check is only the first layer; domains must later be named.

## 16.3 Algebraic Checks

Given a target Lie algebra with generators $X_a$, the lattice candidates $G_a^{(N)}$ should satisfy commutator closure in a controlled sense:

$$i[G_a^{(N)}, G_b^{(N)}] \approx f_{ab}{}^c G_c^{(N)} + \text{central or boundary terms.}$$

The approximation relation must be specified: exact finite-size equality, matrix-element convergence, strong resolvent convergence, strong operator convergence on a core, or convergence after compression to a low-energy sector.

## 16.4 Scaling Checks

The OAR source shows that continuous spatial translations can be reconstructed from discrete dyadic translations in a suitable scaling limit, with the usual momentum operators as generators. It also states that analogous convergence for Lorentz or conformal transformations requires further work. This is the main warning for this programme:

$$\text{translation recovery} \not\Rightarrow \text{Poincare recovery.}$$

Therefore every candidate boost or rotation needs its own scaling statement. For a sequence $G_N$, the report must say:

$$G_N \xrightarrow{?} G$$

and name the topology, state sequence, renormalisation maps, and dense domain or core on which the statement is intended.

## 16.5 Observable Covariance Checks

The target is not only an operator on states. It is a symmetry action on the observable system. The evidence payload should include

$$\alpha_N^G(A) = e^{itG_N} A e^{-itG_N}$$

or the infinitesimal commutator $i[G_N, A]$, together with the expected transformation of local algebras. For translations this means moving supports. For rotations it means rotating supports. For boosts it means mixing time and space in the continuum representation.

### 16.6 Acceptance Payload

A lattice generator candidate is accepted into the compiler only with:

1. density provenance: the exact $h_j$ and supports,
2. finite algebra checks: self-adjointness and commutators,
3. renormalisation maps and state sequence,
4. operator topology or matrix-element convergence target,
5. observable covariance statement,
6. boundary/central/vacuum-subtraction policy.

### 16.7 Boundary

Time-zero net axioms and Lieb-Robinson bounds give important locality control, but local causality is not the same as full Poincare covariance. The compiler must keep these evidence levels separate.

# 17 A Compiler Interface for Lattice Symmetry Candidates

## 17.1 Status, Sources, and Dependencies

**Status: compiler specification shard.** This shard extends CA-08 at the level of output slots. It is not yet an implementation.

Dependencies: CA-08 and CA-10–CA-15.

## 17.2 Input Contract

The lattice symmetry compiler receives a compiled Hilbert/QM system plus:

| | |
|---|---|
| $\Lambda_N$ | : finite lattice, graph, or cell complex, |
| $x$ | : $\Lambda_N \to \mathbb{R}^d$ position map, |
| $\mathcal{A}_N$ | : observable algebra with locality map, |
| $H_N = \sum_r h_r$ | : self-adjoint local Hamiltonian density, |
| $T$ | : target symmetry algebra or group, |
| $S_N$ | : state/sector sequence and scaling maps, if known. |

If the Hamiltonian is supplied only as a black-box matrix, the compiler must fail for boost/rotation generation: the density decomposition is part of the mathematical data.

## 17.3 Candidate Generation

For one-dimensional open chains, the first candidate layer is deterministic:

$$K_x = \sum_j x_j h_j, \qquad P_x = i[H, K_x].$$

For uniform nearest-neighbour density this reduces to the CA-12 current sum.

For higher-dimensional lattices, scalar ramps $f_a(r) = r^a$ generate candidate momentum components

$$P_a^{\mathrm{lat}} := i[H, K_{f_a}].$$

Rotation candidates are then built only after the momentum-density/current layer has its own convention and evidence:

$$J_{ab}^{\mathrm{lat}} \approx \sum_r (r_a p_b^{\mathrm{lat}}(r) - r_b p_a^{\mathrm{lat}}(r)).$$

Until that layer is fixed, this is a deferred proposal slot rather than a formula emitted by the compiler. Koo–Saleur-style modes use Fourier weights instead of polynomial weights and include the commutator corrections specified by the cited Definition 4.1.

## 17.4 Output Contract

The compiler output for each candidate generator must include:

| | |
|---|---|
| name | : for example $P_a, K_a, J_{ab}, L_k, \bar{L}_k$, |
| formula | : finite expression in $h_r$ and commutators, |
| domain policy | : finite matrix, core, or unbounded deferral, |
| locality support | : site/bond/cell support accounting, |
| status | : Proposal, Source-local, or Checked, |
| evidence | : source lines, derivation, test, or run artifact, |
| continuum target | : target generator and expected bracket. |

### 17.5 Failure Modes

The compiler must fail loudly when:

1. $H$ has no density decomposition,
2. $\Lambda_N$ has no position map but a boost/rotation is requested,
3. periodic boundary positions are requested without a branch convention,
4. the target algebra requires a source/convention not yet fixed,
5. a convergence claim is requested without scaling maps or states.

### 17.6 Compiler Principle

The finite expression and the continuum theorem are different compilation products. The first can often be generated algebraically from $(\Lambda_N, x, \{h_r\})$. The second is a proof obligation involving scaling maps, states, and operator topology. The compiler should output both slots, but it must never fill the proof slot with intuition.

# 18 Small Examples and the Lattice-Symmetry Research Queue

## 18.1 Status, Sources, and Dependencies

**Status: worked examples and queue.** The examples are deliberately small. They illustrate compiler behaviour, not new convergence theorems.

Dependencies: CA-10–CA-16 and CONVENTIONS.md (h), (j).

## 18.2 Example 1: Black-Box Quantum System

Input: a Hilbert space and a self-adjoint Hamiltonian $H$.

Output: time translation $U(t) = e^{-itH}$.

Failed request: infer boosts, rotations, or spatial translations. There is no lattice geometry, no position map, and no Hamiltonian density. The compiler must stop.

## 18.3 Example 2: Open Uniform Chain

Input:

$$H = \sum_{j=1}^{N} h_j, \qquad x_j = j, \qquad [h_j, h_k] = 0 \text{ for } |j-k| > 1.$$

Candidate boost:

$$K = \sum_j j h_j.$$

Candidate momentum:

$$P = i[H, K] = \sum_{j=1}^{N-1} i[h_j, h_{j+1}].$$

Status: checked finite derivation; no continuum convergence yet.

## 18.4 Example 3: Nonuniform Chain

Input: positions $x_1 < \cdots < x_N$. The same derivation gives

$$P_x = i[H, K_x] = \sum_{j=1}^{N-1} (x_{j+1} - x_j) i[h_j, h_{j+1}].$$

The coefficients are discrete spacings. The Julia test pins this invariant for uniform and nonuniform examples.

## 18.5 Example 4: Periodic Chain

Input: a circle with periodic density terms.

Problem: $x_j = j$ is not a single-valued function on the circle. A periodic boost-like first moment requires a branch cut, a sawtooth, Fourier modes, or a different construction. The same warning applies to finite periodic momentum-coordinate tests: a finite grid supports periodic symbols and Fourier eigenvalue checks, not an exact differential operator $X = i\partial_k$ with a canonical commutator. Koo–Saleur Fourier modes are the better sourced periodic prototype.

### 18.6 Example 5: Square Lattice

Input: positions $r = (m, n)$. Scalar ramps $f_x(r) = m$, $f_y(r) = n$ give two candidate momentum components via

$$P_a^{\text{lat}} = i[H, \sum_r r^a h_r].$$

Rotation about the origin should then be built as a position-dependent translation from the two momentum-density components. Status: proposal until an orientation and density-localisation convention is fixed.

### 18.7 Example 6: Koo–Saleur Modes

Input: periodic one-dimensional critical Hamiltonian density. Fourier modes of the density and commutator corrections give candidates for $L_k, \bar{L}_k$. In the cited free-fermion/OAR setting, smeared approximants converge to smeared Virasoro generators. Status outside that setting: source-local precedent, not a general theorem.

### 18.8 Research Queue

1. Use registered Weinberg anchors before promoting exact Lorentz-generator formulas.
2. Use the registered original Koo–Saleur anchors before promoting model-specific KS claims.
3. Source or derive a general local energy-current theorem.
4. Fix periodic branch/sawtooth/Fourier first-moment conventions.
5. Define higher-dimensional oriented density supports.
6. Add numerical examples checking commutator closure in small chains.
7. Connect anyon-chain Hamiltonian densities to the same candidate layer.

### 18.9 Landing Point

The next compiler upgrade is therefore not only a Hilbert-space compiler and not yet only a tensor-category compiler. It is a lattice-QM compiler layer:

$$(\text{carrier}, \mathcal{A}_N, \Lambda_N, \{h_r\}, x, S_N) \longmapsto (\text{candidate bulk symmetries}, \text{evidence obligations}).$$

Only after those obligations are discharged can the continuum symmetry representation be claimed.

# 19 Why Galilean Symmetry Is a Different Lattice Target

## 19.1 Status, Sources, and Dependencies

**Status: sourced boundary plus proposal shard.** The Galilei group as a classical nonrelativistic symmetry group is sourced locally. The general projective-representation and central-extension mechanism is sourced locally. The specific mass central extension of the Galilei group is not yet sourced in this repository; CA-20 treats its coefficient as a checked canonical-commutator derivation, not as a literature claim about a named group.

Dependencies: CA-05 for projective symmetries via central extensions, CA-10–CA-17 for the Lorentzian lattice-symmetry programme, and CONVENTIONS.md (i).

## 19.2 The New Target

The previous block targeted Lorentz-invariant continuum systems. Its first lattice intuition was:

$$\text{Lorentz boost} \simeq \text{position-dependent time translation}, \qquad K_a^{\mathrm{Lor}} \sim \sum_x x_a h_x.$$

That is appropriate only when the continuum target mixes space and time in the Lorentzian way and the boost is controlled by the energy density.

The nonrelativistic target is different. Schottenloher lists the Galilei group as the symmetry group for free particles in classical nonrelativistic mechanics. For the lab-book compiler this means that a Galilean target is not obtained by renaming the Poincare block. It has absolute time, spatial translations, rotations, and boosts whose quantum implementation may be projective.

## 19.3 Immediate Consequence for Lattices

The Lorentzian boost ansatz used only the Hamiltonian density $h_x$. A Galilean boost needs more input. At time $t = 0$, the familiar nonrelativistic one-particle expression is controlled by the mass-weighted position, not by the energy-weighted position:

$$G_a(0) \sim m X_a.$$

For a many-body lattice this suggests a mass-density or particle-number-density first moment:

$$G_a^{\mathrm{lat}}(0) \sim \sum_{x \in \Lambda} x_a \, \mu_x, \qquad \mu_x = m \, n_x \quad \text{when the particles have common mass } m.$$

This is a proposal-level lattice formula until the model supplies the density $\mu_x$, the conservation law, and the continuity/current relation.

## 19.4 Compiler Implication

The compiler rule is therefore stricter than in CA-16:

$$\begin{aligned} &\text{Lorentzian candidate layer:} && (\Lambda, x, \{h_x\}) \text{ may generate } K_a \sim \textstyle\sum_x x_a h_x, \\ &\text{Galilean candidate layer:} && (\Lambda, x, \{h_x\}) \text{ is not enough.} \end{aligned}$$

For Galilean boosts the input must also name:

$$\{\mu_x\} \text{ or } \{n_x\}, \qquad M = \sum_x \mu_x, \qquad [H, M] = 0,$$

plus, for a momentum candidate, a current witness for the discrete continuity equation. If those data are absent, the compiler must fail loudly rather than silently using the energy first moment.

## 19.5 Why Projective Data Matter Earlier

In the Hilbert-space grammar, projective symmetries were handled by passing to a central extension. Galilean symmetry is the first lattice target where that choice becomes operational: the mass parameter is expected to sit in the projective/central part of the quantum representation. Locally we can support only this precise statement:

projective quantum symmetries are represented by unitary representations of central extensions.

The stronger statement identifying the Galilei extension by its standard literature name remains a source-acquisition task.

## 19.6 Boundary for CA-18–CA-22

The follow-up block does four things:

| | |
|---|---|
| CA-19 | derives the unextended Galilei vector-field algebra, |
| CA-20 | derives the mass central coefficient on a canonical core, |
| CA-21 | derives the lattice first-moment continuity formula, |
| CA-22 | turns the data into compiler rules and small examples. |

No shard in this block claims that a finite lattice has exact Galilean symmetry unless the corresponding finite-dimensional representation and commutator checks are actually supplied.

# 20 The Galilei Algebra as Newtonian Vector Fields

## 20.1 Status, Sources, and Dependencies

**Status: checked local derivation.** The source anchor is that the Galilei group is the nonrelativistic free-particle symmetry group. The bracket table below is derived here from vector fields and checked by the Julia test suite.

Dependencies: CA-18 and CONVENTIONS.md (i).

## 20.2 Generators

Use Newtonian spacetime coordinates

$$(t, x_1, \ldots, x_d).$$

The convention fixed in CONVENTIONS.md (i) is:

$$\begin{aligned} H &= \partial_t, && \text{time translation,} \\ P_a &= \partial_a, && \text{spatial translation,} \\ G_a &= t\,\partial_a, && \text{Galilean boost,} \\ J_{ab} &= x_a\partial_b - x_b\partial_a, \quad a<b, && \text{spatial rotation.} \end{aligned}$$

Here $a, b, c \in \{1, \ldots, d\}$. The boost vector field says that a boost changes position by a velocity times time while leaving absolute time fixed.

## 20.3 Translation and Boost Brackets

The translation fields have constant coefficients, hence

$$[H, P_a] = 0, \qquad [P_a, P_b] = 0.$$

Since $G_a = t\partial_a$,

$$\begin{aligned} [H, G_a] &= [\partial_t, t\partial_a] \\ &= \partial_t(t)\partial_a \\ &= P_a. \end{aligned}$$

The reverse bracket is therefore $[G_a, H] = -P_a$. Spatial translations do not differentiate the coefficient $t$, so

$$[P_a, G_b] = 0, \qquad [G_a, G_b] = 0.$$

This is the unextended vector-field algebra. It is not yet the projective quantum algebra; CA-20 adds the central term that is invisible to vector fields on spacetime.

## 20.4 Rotation on Vectors

For $J_{ab} = x_a\partial_b - x_b\partial_a$, compute against a translation:

$$\begin{aligned} [J_{ab}, P_c] &= -\partial_c(x_a)\partial_b + \partial_c(x_b)\partial_a \\ &= \delta_{bc}P_a - \delta_{ac}P_b. \end{aligned}$$

The same calculation applies to boosts because $J_{ab}$ does not act on $t$:

$$[J_{ab}, G_c] = \delta_{bc}G_a - \delta_{ac}G_b.$$

Thus both $P_c$ and $G_c$ transform as spatial vectors.

## 20.5 Rotation-Rotation Bracket

For two rotations, expanding the vector-field bracket gives

$$\begin{aligned}[J_{ab}, J_{cd}] = \delta_{bc} J_{ad} - \delta_{ac} J_{bd} \\ + \delta_{ad} J_{bc} - \delta_{bd} J_{ac},\end{aligned}$$

with the convention $J_{ba} = -J_{ab}$ and $J_{aa} = 0$. In $d = 3$, one representative checked bracket is

$$[J_{12}, J_{23}] = J_{13}.$$

## 20.6 Checked Representatives

The test suite pins the signs with the following representative outcomes:

$$\begin{aligned}[H, G_1] &= P_1, \\ [G_1, H] &= -P_1, \\ [J_{12}, P_1] &= -P_2, \\ [J_{12}, P_2] &= P_1, \\ [J_{12}, G_1] &= -G_2, \\ [J_{12}, J_{23}] &= J_{13}.\end{aligned}$$

It also checks that $[P_1, G_2] = 0$ and $[G_1, G_2] = 0$ in the unextended vector-field algebra.

## 20.7 Boundary

The signs here are signs for vector fields. A quantum model uses self-adjoint generators and unitary flows, so translating this table into operator commutators requires the Stone convention and the chosen $i[A, B]$ observable action. The report therefore cites this shard for the geometric algebra and CA-20 for the central quantum coefficient.

# 21 Projective Galilean Symmetry and the Mass Central Term

## 21.1 Status, Sources, and Dependencies

**Status: sourced projective framework plus checked local derivation.** Schottenloher supplies the representation-theoretic framework: projective symmetries need not lift to the original group, but lift through a central extension. The coefficient $m\delta_{ab}$ below is a local canonical commutator derivation checked in Julia. A local source specifically identifying the Galilei mass central extension by its standard name has not yet been registered.

Dependencies: CA-05, CA-18, CA-19, and CONVENTIONS.md (f), (i).

## 21.2 Why This Belongs in the Grammar

CA-05 made a conservative grammar choice: a projective symmetry is represented by an honest unitary representation of a named central extension

$$1 \longrightarrow U(1) \longrightarrow E \longrightarrow G \longrightarrow 1.$$

For Galilean symmetry this is not a decorative detail. The central parameter is physically interpreted as mass in the canonical nonrelativistic model, and it controls how boosts and momenta compose at the quantum level.

## 21.3 Canonical One-Particle Calculation

Work formally on a common invariant core where the position and momentum operators obey

$$[X_a, P_b] = i\delta_{ab} I.$$

For a particle of mass $m$, the time-zero boost generator is represented by

$$G_a(0) = mX_a.$$

Then

$$\begin{aligned} [G_a(0), P_b] &= [mX_a, P_b] \\ &= m[X_a, P_b] \\ &= i\, m\, \delta_{ab} I. \end{aligned}$$

The Julia check records exactly the real coefficient $m\delta_{ab}$.

## 21.4 Compatibility with Time Evolution

For the free Hamiltonian

$$H = \frac{1}{2m} \sum_b P_b^2,$$

the same canonical commutator gives

$$\begin{aligned} [H, mX_a] &= \frac{1}{2} \sum_b [P_b^2, X_a] \\ &= \frac{1}{2} \sum_b \big( P_b [P_b, X_a] + [P_b, X_a] P_b \big) \\ &= \frac{1}{2} \big( -iP_a - iP_a \big) = -iP_a. \end{aligned}$$

Thus

$$i[H, G_a(0)] = P_a.$$

This matches the CA-19 vector-field relation $[H, G_a] = P_a$ after applying the self-adjoint-generator convention used for observable evolution.

### 21.5 Central Extension Versus Vector Fields

The unextended vector-field algebra had $[G_a, P_b] = 0$. The canonical quantum calculation has instead

$$[G_a, P_b] = i\, m\, \delta_{ab} I.$$

The additional term is central: it is a scalar multiple of the identity on a fixed-mass sector. This is the algebraic reason the compiler must attach a central-extension witness to Galilean symmetry data rather than store only the spacetime vector fields.

### 21.6 Source Gap

The local source collection includes Bargmann's theorem about lifting projective representations under a Lie-algebra cohomology hypothesis, and it includes the general central-extension theorem for projective representations. It does not yet include a registered source for the standard mass central extension of the Galilei group. Until that source is acquired, this shard uses the cautious name *mass central term* and cites the checked canonical calculation as its local evidence.

### 21.7 Compiler Payload

A Galilean projective symmetry record must therefore include:

| | |
|---|---|
| base group/algebra | Galilei target from CA-19, |
| central generator | $M$ or $I$ on a fixed-mass sector, |
| central charge | $m$ or a mass operator, |
| raw commutator witness | $[G_a, P_b] = i\delta_{ab} M$, |
| sign convention | conversion to $i[A, B]$ stated explicitly. |

An OR of sectors with different masses can still be a Hilbert direct sum, but the central element must then be represented as a block-diagonal operator rather than a single scalar charge.

# 22 Lattice Galilean Boosts from Mass-Density First Moments

## 22.1 Status, Sources, and Dependencies

**Status: proposal plus checked local derivation.** Local sources support Galilean invariance as a physical assumption in nonrelativistic fermion models and give examples where conservation laws produce continuity equations. The finite open-chain first-moment formula below is a local derivation checked by the Julia test suite.

Dependencies: CA-18–CA-20 and CONVENTIONS.md (h), (i).

## 22.2 Input Data

Let $\Lambda = \{1, \ldots, N\}$ be a finite open chain with positions $x_j$. A Galilean lattice candidate needs, in addition to the Hamiltonian, a conserved mass or particle-number density:

$$M = \sum_j \mu_j, \qquad [H, M] = 0.$$

For identical particles of mass $m$, a common special case is

$$\mu_j = m n_j,$$

where $n_j$ is a number density.

The time-zero Galilean boost candidate is the first mass moment

$$G^{\mathrm{lat}}(0) = \sum_j x_j \mu_j.$$

This is the Galilean analogue of $mX$, not the Lorentzian energy moment $\sum_j x_j h_j$.

## 22.3 Discrete Continuity Assumption

Assume a nearest-neighbour open-chain continuity equation with bond currents $J_{j+1/2}$:

$$i[H, \mu_j] = J_{j-1/2} - J_{j+1/2},$$

where endpoint currents are either absent or treated as boundary terms. This assumption is model data. It does not follow merely from writing $H = \sum_j h_j$.

## 22.4 First-Moment Derivation

Multiply the continuity equation by $x_j$ and sum:

$$\begin{aligned} i\left[H, \sum_j x_j \mu_j\right] &= \sum_j x_j (J_{j-1/2} - J_{j+1/2}) \\ &= \sum_{j=1}^{N-1} (x_{j+1} - x_j) J_{j+1/2} \quad + \quad \text{boundary terms.} \end{aligned}$$

For a unit-spaced open chain away from endpoints,

$$i[H, G^{\mathrm{lat}}(0)] \simeq \sum_{j=1}^{N-1} J_{j+1/2}.$$

This is the lattice momentum candidate associated with the Galilean boost. For nonuniform positions the coefficients are the spacings

$$\Delta x_j = x_{j+1} - x_j,$$

and the test suite records a coefficient check for this shard.

### 22.5 Relation to Center of Mass

The local paired-fermion source uses Galilean invariance to explain a response in terms of center-of-mass acceleration while relative motion is unaffected. That is exactly the intuition this shard turns into compiler data:

$$\text{Galilean boost} \quad \rightsquigarrow \quad \text{center-of-mass / mass-first-moment generator.}$$

The source does not prove the lattice formula above. It supports the physical role of center-of-mass motion under Galilean invariance; the lattice identity is the checked local derivation.

### 22.6 What Is and Is Not Proven

The finite identity proves only that, given a local continuity equation, the time derivative of the mass first moment is a current sum. It does not prove:

1. that the current sum converges to continuum momentum,
2. that the finite lattice has exact Galilei symmetry,
3. that the mass current equals momentum density in every model,
4. that boundary terms vanish without a stated boundary condition.

Those are acceptance-test obligations for later shards or model-specific examples.

# 23 Compiler Rules and Examples for Galilean Targets

## 23.1 Status, Sources, and Dependencies

**Status: compiler specification shard.** This shard does not implement the compiler. It specifies the extra fields a Galilean target requires, using the checked algebra and first-moment derivations from CA-19–CA-21.

Dependencies: CA-08, CA-16, and CA-18–CA-21.

## 23.2 Target Declaration

A Galilean compiler target is a structured request:

$$\text{Target}(\text{Gal}(d),\ \text{projective},\ \text{mass}).$$

It asks for:

| | |
|---|---|
| $H$ | time translation, |
| $P_a$ | spatial translations / momenta, |
| $J_{ab}$ | rotations, |
| $G_a$ | Galilean boosts, |
| $M$ | central mass generator. |

The finite candidate layer is allowed to output formulas and witnesses. It is not allowed to assert continuum Galilean covariance without scaling-limit evidence.

## 23.3 Required Input Slots

Compared with CA-16, the Galilean target adds the following required slots:

| | |
|---|---|
| $\mu_x$ or $n_x$ | local mass or particle-number density, |
| $M = \sum_x \mu_x$ | conserved total mass/number, |
| $[H, M] = 0$ | finite commutator witness, |
| $J_{x,e}$ | current witness for a discrete continuity equation, |
| $m$ or $M$ | central charge/operator for the projective representation, |
| boundary policy | open, periodic with branch, or thermodynamic-limit handling. |

If any of the first four are absent, boosts and momenta for a Galilean target are underdetermined.

## 23.4 Generated Candidates

For an open chain, the deterministic candidate formulas are

$$G^{\text{lat}}(0) = \sum_j x_j \mu_j,$$

and, when a continuity witness is supplied,

$$P^{\text{lat}} := i[H, G^{\text{lat}}(0)] = \sum_{j=1}^{N-1} (x_{j+1} - x_j) J_{j+1/2} \quad + \quad \text{boundary terms.}$$

Rotations in higher dimension require momentum or current-density data:

$$J^{\text{lat}}_{ab} \approx \sum_x (x_a p_b(x) - x_b p_a(x)).$$

Thus rotations are second-stage candidates: first compile current/momentum density, then form the spatial first moment.

### 23.5 Example 1: One Free Particle

Input:

$$\mathcal{H} = L^2(\mathbb{R}^d), \qquad H = \frac{P^2}{2m}, \qquad [X_a, P_b] = i\delta_{ab} I.$$

Output at $t = 0$:

$$G_a = mX_a, \qquad [G_a, P_b] = im\delta_{ab} I, \qquad i[H, G_a] = P_a.$$

Status: checked formal derivation on a common core, with unbounded-operator domain work deferred.

### 23.6 Example 2: Number-Conserving Open Chain

Input:

$$H, \quad n_j, \quad [H, \sum_j n_j] = 0, \quad i[H, n_j] = J_{j-1/2} - J_{j+1/2}.$$

For identical mass $m$,

$$\mu_j = mn_j, \qquad G^{\text{lat}}(0) = m \sum_j x_j n_j.$$

The compiler may output the current-sum momentum candidate with coefficient $m(x_{j+1} - x_j)$. It must still mark continuum Galilean covariance as a proof obligation.

### 23.7 Example 3: Hamiltonian Density Only

Input:

$$H = \sum_j h_j$$

with no $n_j$, no $\mu_j$, and no current witness. For a Lorentzian target, CA-12 permits the energy-moment proposal. For a Galilean target, the compiler must return:

```
CompileError:  missing mass/number density and continuity witness.
```

Using $\sum_j x_j h_j$ here would be a convention bug.

### 23.8 Example 4: Spin Chain with No Particle Number

A spin chain may have a perfectly good Hamiltonian density and exact lattice translations. Unless a conserved density is explicitly declared as mass or particle number, it does not compile as a Galilean particle system. The correct output is either a failed Galilean target or a different target declaration.

### 23.9 Example 5: Direct Sum of Mass Sectors

For

$$\mathcal{H} = \mathcal{H}_{m_1} \oplus \mathcal{H}_{m_2},$$

the central generator is not one scalar if $m_1 \neq m_2$. It is

$$M = m_1 I_{\mathcal{H}_{m_1}} \oplus m_2 I_{\mathcal{H}_{m_2}}.$$

The compiler can carry the OR carrier, but an irreducible fixed-mass sector selection is a separate projection/sector step in the sense of CA-06.

### 23.10 Next Obligations

The next Galilean work needs:

1. a registered source for the standard mass central extension,
2. model examples with explicit lattice number-current continuity,
3. finite commutator tests for those models,
4. scaling maps proving that current sums converge to continuum momenta,
5. a decision on periodic boundary branch conventions.

Only after these are present should the compiler promote Galilean lattice candidates from formulas to continuum symmetry evidence.

# 24 Gaussian Boson Lorentz Generator Roadmap

## 24.1 Status, Sources, and Dependencies

**Status: roadmap shard with checked seed.** This shard scopes a multi-shard programme. CA-24 supplies the first checked symbol calculation. The stronger continuum conclusion is not yet proved for a general coefficient class.

Dependencies: CA-10–CA-17 for lattice Lorentz motivation and acceptance tests, and CONVENTIONS.md (j). This shard starts the Gaussian-boson block.

## 24.2 Problem Statement

We now restrict the lattice-symmetry problem to a solvable class: translation-invariant Gaussian bosonic lattice systems in spatial dimensions

$$d = 1, 2, 3, \qquad D = d + 1.$$

The physical target is Lorentz/Poincare covariance in the scaling limit. The expected endpoint is the scalar free field:

$$\omega(k)^2 = m^2 + |k|^2$$

after the correct normalization of lattice spacing, velocity, field strength, and mass.

The inputs are not merely a matrix Hamiltonian. The compiler receives:

| | |
|---|---|
| $\Lambda_{\epsilon,L}$ | periodic hypercubic lattice or infinite lattice, |
| $q_x, p_x$ | canonical bosonic field variables, |
| $V_r$ | translation-invariant quasi-local quadratic coefficients, |
| $\epsilon$ | lattice spacing, |
| $S_\epsilon$ | state/scaling-map data, when a continuum claim is requested. |

The first coefficient tier is the scalar canonical Hamiltonian

$$H = \frac{1}{2}\sum_x p_x^2 + \frac{1}{2}\sum_{x,r} q_x V_r q_{x+r}.$$

Full bosonic BdG/pairing systems are a later tier, not silently included in the first convention.

## 24.3 North Star

For each admissible coefficient family $V_r^{(\epsilon)}$, compute the candidate generators as functions of the coefficients:

$$H, \quad P_a, \quad J_{ab}, \quad K_a,$$

then compute the residuals in the Poincare algebra. The residuals become conditions on the symbol

$$\omega_\epsilon(k)^2 = \sum_r V_r^{(\epsilon)} e^{i\epsilon k\cdot r}.$$

The end goal is a theorem-shaped statement:

Gaussian lattice coefficients satisfying locality, positivity,
single low-energy minimum, and isotropic Klein-Gordon Taylor data
$\Longrightarrow$ scaling limit is the free scalar/Klein-Gordon Poincare system.

**CA-15 proof boundary.** This is a theorem-shaped target, not a theorem already obtained from the symbol checks. A continuum claim still needs the CA-15 payload: scaling maps, state sequence, topology, operator domains, local observable covariance, and boundary/renormalization policy. The value of the Gaussian block is that the algebraic residuals can be made explicit before attempting that proof.

### 24.4 Shard Roadmap

The current Gaussian block is:

**CA-24.** Symbol calculus for the scalar coefficient tier, including the boost-time residual and the checked nearest-neighbour Klein-Gordon seed.

**CA-25.** Diagonalization into one-particle symbols, with the sourced free-scalar Fock construction separated from the local coefficient-symbol derivation and BdG pairing deferred.

**CA-26.** Smooth-patch one-particle generator algebra for $H, P, J, K$, exact brackets, dispersion-dependent residuals, and only formal $\mathrm{d}\Gamma$ bookkeeping on the algebraic finite-particle core.

**CA-27.** Lorentz residual conditions on coefficients: Taylor constraints, anisotropic failures, and doubler counterexamples.

**CA-28.** Numerical invariant suite for scalar symbols and finite periodic stiffness matrices, covering Klein-Gordon, anisotropic, and doubler examples.

**CA-29.** Real-space energy-density convention for the scalar Gaussian Hamiltonian: cell volume, bond sharing, boundary policy, vacuum subtraction, and improvement terms before any position-weighted use.

**CA-30.** One-dimensional stress-energy candidate dictionary: $T_{00}$ from CA-29, checked finite-open-chain $T_{01}$ from A4, and proposal-level $T_{10}$ and $T_{11}$ witnesses left for later work.

**CA-31.** One-dimensional current-symbol bridge: the translation-invariant integrated $T_{01}$ current symbol equals the nearest-neighbour Klein-Gordon boost-time symbol, while finite open-chain $\mathrm{d}\Gamma(k)$ equivalence remains open.

**CA-32.** Higher-dimensional cell-current proposal: oriented $d = 2, 3$ support data, energy-flux equations, momentum-density witnesses, stress-current witnesses, and rotation/boundary diagnostics as compiler inputs rather than convergence claims.

The remaining queued follow-up block is proposal work, not completed stress-energy material:

**Open-chain momentum equivalence.** Upgrade the CA-31 symbol bridge to finite open-chain wave-packet and $\mathrm{d}\Gamma(k)$ comparison data, including boundary residuals.

**Stress-energy numerical suite.** Add numerical checks only after candidate formulas and source-backed expected identities exist.

**Source acquisition.** Use the newly registered discrete Gaussian/free-field Virasoro source only within its square-grid dGFF change-of-measure scope. Separate source anchors are still required for harmonic-chain energy-current formulas and any continuum stress-energy/Poincare-generator claim.

## 24.5 Side Quests

**Source acquisition.** The registered local OAR sources cover the free scalar lattice scaling limit and spacetime translations. They explicitly warn that Lorentz/conformal convergence requires further work. The discrete Gaussian free-field Virasoro source is now local; it supports lattice-level change-of-measure Virasoro claims for the square-grid dGFF, not a continuum stress-energy or Poincare-generator formula.

**Generator ambiguity.** A local Hamiltonian density is not unique:

$$h_x \mapsto h_x + \nabla \cdot b_x$$

changes first moments by boundary or current terms. The Gaussian block must track which density convention produced $K_a = \sum_x x_a h_x$.

**Periodic positions.** Numerical tests use periodic lattices and finite Brillouin grids. First moments, position coordinates, and finite-grid momentum-coordinate generators need a branch, sawtooth, Fourier interpolation, or another explicit periodic coordinate convention. Until that is fixed, early finite tests are symbol/eigenvalue checks on the periodic grid, or open-chain first-moment checks.

**BdG pairing.** A general quasi-local Hamiltonian in creation and annihilation operators may include pairing terms. The scalar $q, p$ tier captures Klein-Gordon directly; BdG requires symplectic positivity and a stable bosonic Bogoliubov convention before generator formulas are promoted.

## 24.6 Numerical Verification Plan

The Julia layer should stay at the invariant level:

1. symbol from coefficients equals the analytic dispersion,
2. $\frac{1}{2}\nabla\omega_\epsilon(k)^2$ equals the smooth-patch boost-time residual,
3. $\frac{\sin(\epsilon k_a)}{\epsilon} \to k_a$ for the KG lattice,
4. anisotropic Hessians fail the Lorentz residual checks,
5. finite Fourier matrices diagonalize periodic stiffness matrices.

The first three checks are implemented with CA-24.

## 24.7 Acceptance Boundary

The roadmap rows above remain scoped by the CA-15 proof boundary stated next to the theorem-shaped goal. Passing symbol or finite-matrix tests is not yet a proof of Lorentz invariance.

# 25 Gaussian Boson Symbol Calculus and the Boost-Time Residual

## 25.1 Status, Sources, and Dependencies

**Status: sourced model class plus checked flat local derivation.** The standard lattice scalar-field Hamiltonian and dispersion are sourced. The boost-time residual calculation is a local one-particle derivation on flat $L^2(dk)$, checked by Julia on positive-dispersion patches. It is not yet a sourced OAR or mass-shell Poincare representation theorem, and massless zero-mode examples are coefficient-level tests only.

Dependencies: CA-11, CA-12, CA-23, and CONVENTIONS.md (h), (j), (k).

## 25.2 Coefficient Tier

Fix $d \in \{1, 2, 3\}$. On a periodic hypercubic lattice with spacing $\epsilon$, use canonical fields

$$[q_x, q_y] = 0, \qquad [p_x, p_y] = 0, \qquad [q_x, p_y] = i\delta_{xy}.$$

The first scalar Gaussian tier is

$$H = \frac{1}{2}\sum_x p_x^2 + \frac{1}{2}\sum_{x,r} q_x V_r q_{x+r}, \qquad V_r = V_{-r} \in \mathbb{R}.$$

Quasi-locality means that $V_r$ has enough decay, or finite range in the first examples, so that the Fourier symbol and its derivatives used below exist.

Define

$$\omega_\epsilon(k)^2 := \sum_r V_r e^{i\epsilon k\cdot r}.$$

For a stable scalar model, this symbol is non-negative, with the zero-mode policy stated separately in the massless case. The helper parameter $m$ is a nonnegative mass label; the generator statements below use only the stricter case $\omega_\epsilon(k) > 0$ on the chosen patch.

## 25.3 Nearest-Neighbour Klein-Gordon Example

The sourced lattice scalar-field example has, in the notation of this shard,

$$V_0 = m^2 + \frac{2d}{\epsilon^2}, \qquad V_{\pm e_a} = -\frac{1}{\epsilon^2}.$$

Therefore

$$\begin{aligned}\omega_{\epsilon,m}(k)^2 &= m^2 + \frac{2d}{\epsilon^2} - \frac{1}{\epsilon^2}\sum_{a=1}^{d}(e^{i\epsilon k_a} + e^{-i\epsilon k_a}) \\ &= m^2 + \frac{2}{\epsilon^2}\sum_{a=1}^{d}(1 - \cos(\epsilon k_a)).\end{aligned}$$

For fixed physical $k$ and $\epsilon \to 0$,

$$\omega_{\epsilon,m}(k)^2 = m^2 + |k|^2 + O(\epsilon^2|k|^4).$$

## 25.4 Momentum-Space Position

After diagonalization, this shard uses the flat local momentum space $L^2(\mathcal{U}, dk)$. The one-particle Hamiltonian is multiplication by $\omega(k)$, with $\omega(k) > 0$ on $\mathcal{U}$, and

$$X_a = i\partial_{k_a}$$

on a common smooth core. The time-zero boost candidate corresponding to the energy first moment is the flat-measure symmetrized operator

$$K_a = \frac{1}{2}(X_a\omega + \omega X_a).$$

The symmetrization is the one-particle version of taking a self-adjoint first moment of the energy density. If the one-particle scalar product is changed, the adjoint and hence the boost formula must be rederived; no unitary equivalence with the OAR weighted space or Schottenloher mass-shell realization is asserted here.

## 25.5 Boost-Time Residual

For a test wavefunction $\psi$,

$$\begin{aligned}[K_a, \omega]\psi &= \frac{1}{2}(X_a\omega^2 - \omega X_a\omega)\psi \\ &= i\,\omega\,(\partial_a\omega)\psi.\end{aligned}$$

Thus

$$i[\omega, K_a] = \omega\,\partial_a\omega = \frac{1}{2}\partial_a\omega(k)^2.$$

This is the algebraic output of the boost-time commutator. With the vector-field-to-Stone sign map of CONVENTIONS.md (k), the continuum Poincare target is $i[H, K_a] = P_a$. Taking $P_a = k_a$, one needs

$$\frac{1}{2}\partial_a\omega(k)^2 = k_a.$$

Integrating over $k$ gives

$$\omega(k)^2 = m^2 + |k|^2$$

on the connected low-energy patch. With speed $c$, the right side is $c^2k_a$, giving $\omega(k)^2 = m^2 + c^2|k|^2$.

## 25.6 Lattice Residual

For the nearest-neighbour lattice Klein-Gordon symbol,

$$\frac{1}{2}\partial_a\omega_{\epsilon,m}(k)^2 = \frac{\sin(\epsilon k_a)}{\epsilon}.$$

Therefore the boost-time commutator has the correct continuum limit:

$$\frac{\sin(\epsilon k_a)}{\epsilon} = k_a + O(\epsilon^2 k_a^3).$$

The Julia tests check this in $d = 1, 2, 3$, both from the closed formula and by evaluating the coefficient symbol $\sum_r V_r e^{i\epsilon k\cdot r}$.

## 25.7 Rotation Residual

For the one-particle rotation candidate

$$J_{ab} = X_a P_b - X_b P_a, \qquad P_a = k_a,$$

one obtains

$$i[\omega, J_{ab}] = k_b \partial_a \omega - k_a \partial_b \omega.$$

Thus rotations commute with the Hamiltonian exactly when the dispersion is radial on the momentum region under consideration. For a hypercubic lattice this is not exact over the full Brillouin zone, but the Klein-Gordon expansion is isotropic to quadratic order at $k = 0$.

## 25.8 Boundary

This shard is a one-particle symbol calculation. The many-body generators are second quantizations plus vacuum-energy and domain policies. Real-space locality of $K_a = \sum_x x_a h_x$, boundary terms, and convergence of the second-quantized operators are deferred to later shards.

# 26 Gaussian Boson Diagonalization and the One-Particle Symbol

## 26.1 Status, Sources, and Dependencies

**Status: sourced template plus flat local coefficient derivation.** The standard lattice free-scalar Hamiltonian, Fock representation, continuum one-particle space, and second-quantized Hamiltonian are sourced. The extension from nearest-neighbour coefficients to an arbitrary real translation-invariant scalar symbol is a local algebraic calculation on the flat $L^2(dk)$ symbol space fixed in CONVENTIONS.md (j).

Dependencies: CA-23–CA-24 and CONVENTIONS.md (a), (f), (j).

## 26.2 Scalar Coefficient Tier

Fix $d \in \{1, 2, 3\}$. The first Gaussian tier is

$$H = \frac{1}{2}\sum_x p_x^2 + \frac{1}{2}\sum_{x,r} q_x V_r q_{x+r}, \qquad V_r = V_{-r} \in \mathbb{R}.$$

On an infinite translation-invariant lattice, or on a periodic lattice before taking volume limits, the Fourier transform sends the potential kernel to

$$\omega_\epsilon(k)^2 = \sum_r V_r e^{i\epsilon k\cdot r}.$$

The reality condition $V_r = V_{-r}$ makes this symbol real:

$$\omega_\epsilon(k)^2 = V_0 + \sum_{r>0} 2V_r \cos(\epsilon k \cdot r).$$

The stability condition for this tier is

$$\omega_\epsilon(k)^2 \geq 0$$

on the Brillouin zone. The `mass` parameter is a nonnegative label; in the nearest-neighbour Klein-Gordon family, $m > 0$ gives the positive-dispersion case used by the current generator statements. For a general coefficient symbol the required condition remains $\omega_\epsilon(k) > 0$ on the chosen patch. When $m = 0$, zero modes are coefficient data only until a zero-mode convention excludes or otherwise controls them.

## 26.3 Mode Decoupling

The Hamilton equations are a local derivation from the canonical commutators:

$$\dot{q}_x = p_x, \qquad \dot{p}_x = -\sum_r V_r q_{x+r}.$$

Fourier transforming gives

$$\dot{\widehat{q}}(k) = \widehat{p}(k), \qquad \dot{\widehat{p}}(k) = -\omega_\epsilon(k)^2 \widehat{q}(k),$$

and hence

$$\ddot{\widehat{q}}(k) + \omega_\epsilon(k)^2 \widehat{q}(k) = 0.$$

Thus each momentum mode is an oscillator with frequency $\omega_\epsilon(k)$. This is the algebraic reason the compiler can reduce the scalar Gaussian tier to multiplication by a dispersion symbol.

## 26.4 Complex Structure and Fock Space

For $\omega > 0$ on the chosen momentum patch, the oscillator complex structure is

$$(q, p) \longmapsto (-\omega p, \omega^{-1} q).$$

The sourced free-scalar construction uses this pattern with $\gamma_m(k) = \sqrt{|k|^2 + m^2}$ in the continuum and with $\gamma_m^{(N)}(k)$ on the lattice. It also realizes the Weyl algebra in a bosonic Fock representation by creation and annihilation operators. The source scalar product is the OAR one on real Cauchy data, with $\gamma_m^{-1/2}\widehat{q} + i\gamma_m^{1/2}\widehat{p}$; the text then realizes the representation on a mass-independent flat $L^2$ Fock space. This shard uses only the resulting flat symbol bookkeeping and does not prove that the boost formula of CA-24/CA-26 intertwines the OAR or mass-shell representation.

The compiler-level output for the positive-dispersion scalar tier is therefore:

| | |
|---|---|
| $\mathfrak{h}_\epsilon$ | flat one-particle momentum space on the chosen patch or grid, |
| $h_\epsilon$ | multiplication by $\omega_\epsilon(k)$, |
| $\mathcal{F}_+(\mathfrak{h}_\epsilon)$ | symmetric Fock completion, |
| $H_\epsilon$ | $d\Gamma(h_\epsilon) + E_{\epsilon,0}$ if a vacuum-energy policy is named. |

The additive constant $E_{\epsilon,0}$ is not part of the commutator algebra for the generators considered here, but it is part of a Hamiltonian-density normal-ordering policy and must be recorded before using ground-state energies as numerical claims.

## 26.5 What Is Not Yet Included

This shard does not define the full bosonic BdG tier. A general quadratic Hamiltonian in creation and annihilation operators may include pairing terms, and then the correct diagonalization is a symplectic/Bogoliubov problem with positivity, implementability, and domain conditions. Those conventions are not fixed here.

The current compiler rule is deliberately narrower:

$$\text{scalar } (q, p) \text{ tier} \quad \Rightarrow \quad \omega_\epsilon(k)^2 \text{ symbol} \quad \Rightarrow \quad h_\epsilon = \omega_\epsilon(k).$$

BdG inputs must fail closed or route to a later convention until that tier is written.

## 26.6 Compiler Consequence

For the positive-dispersion scalar Gaussian block, later candidate generators can be constructed first on the one-particle smooth momentum core and then second-quantized. The coefficient data enter only through $\omega_\epsilon(k)^2$ and its derivatives, while the continuum proof still requires the CA-15 payload: scaling maps, state convergence, domains, and observable covariance.

# 27 One-Particle Generator Algebra for Gaussian Bosons

## 27.1 Status, Sources, and Dependencies

**Status: flat local derivation with limited Julia checks.** The Poincare target, Stone convention, and free-boson Fock realization are sourced. The commutator identities in this shard are local differential-operator derivations on a flat $L^2(dk)$ smooth momentum-space core. The Julia suite currently checks coefficient/symbol consequences: Klein-Gordon symbols, boost-time symbols, sampled rotation-Hamiltonian residuals, sampled boost-boost residual coefficient data, low-energy Hessian examples, finite periodic massive spectra, the massless doubler coefficient-rejection witness, and the Gaussian tolerance policy. It does not check the full generator bracket table, finite coordinate commutators, or any second-quantized identity. The smooth-core derivation is not yet a sourced OAR or mass-shell Poincare representation theorem.

Dependencies: CA-11, CA-24–CA-25, and CONVENTIONS.md (h), (j), (k).

## 27.2 Momentum Core

There are two separate settings.

**Smooth patch.** Work on a connected smooth momentum patch $\mathcal{U}$ where $\omega(k) > 0$, with the flat measure $dk$. Let

$$\mathcal{D}_0 = C_c^\infty(\mathcal{U}) \subset L^2(\mathcal{U}, dk)$$

and define

$$X_a = i\partial_{k_a}, \qquad h = \omega(k), \qquad p_a = k_a.$$

The one-particle Hamiltonian and momenta are multiplication operators:

$$H^{(1)} = h, \qquad P_a^{(1)} = p_a.$$

The rotation and time-zero boost candidates are

$$J_{ab}^{(1)} = X_a P_b - X_b P_a, \qquad K_a^{(1)} = \frac{1}{2}(X_a h + h X_a).$$

These are written in the self-adjoint sign convention of CONVENTIONS.md (k); in particular $J_{ab}$ is the Gaussian rotation sign, not the raw lowered-index CA-11 vector field $M_{ab}$. All commutators below are identities on $\mathcal{D}_0$; promoting them to self-adjoint generator statements is a separate domain theorem. Promoting this flat formula to the OAR one-particle Hilbert space or Schottenloher's mass-shell/free-boson Poincare representation additionally requires a sourced or locally derived unitary intertwiner.

**Finite periodic grids.** On a finite periodic Gaussian lattice, the dual grid is used for multiplication symbols and Fourier eigenvalue checks. The current finite suite does not place the differential operator $X_a = i\partial_{k_a}$ on that grid and does not claim an exact finite $[X_a, p_b] = i\delta_{ab}$ commutator. Periodic first-moment or momentum-coordinate generator tests require an explicit branch, sawtooth, Fourier-interpolation, or comparable periodic-coordinate convention before they can be stated. Thus the exact differential commutators below are smooth-patch algebra; the finite periodic checks remain symbol/eigenvalue checks.

## 27.3 Exact Translation and Rotation Brackets

On the smooth core, because $h$ and $p_a$ are multiplication operators,

$$i[h, p_a] = 0.$$

The canonical differential identity is the smooth-patch identity

$$[X_a, p_b] = i\delta_{ab}.$$

Therefore

$$i[P_c, J_{ab}] = \delta_{ac}P_b - \delta_{bc}P_a.$$

This is independent of the dispersion. It is kinematical on the smooth patch: it comes from the chosen momentum coordinate and the differential representation of rotations, in the rotation sign convention of CONVENTIONS.md (k).

### 27.4 Boost-Momentum Bracket

For the boost candidate,

$$\begin{aligned}[P_b, K_a] &= \frac{1}{2}\big([P_b, X_a]h + h[P_b, X_a]\big) \\ &= -i\delta_{ab}h.\end{aligned}$$

Hence

$$i[P_b, K_a] = \delta_{ab}H.$$

This is the boost-momentum line of the self-adjoint target table in CONVENTIONS.md (k). On the smooth patch, this part of the Poincare algebra is exact for every positive scalar dispersion because it only uses $K_a$ as the symmetrized first moment of energy.

### 27.5 Boost-Time Residual

CA-24 derived

$$i[H, K_a] = \frac{1}{2}\partial_a\omega(k)^2.$$

The continuum target with speed $c = 1$ wants $P_a = k_a$. The residual is

$$R_a(k) := \frac{1}{2}\partial_a\omega(k)^2 - k_a.$$

For speed $c$, replace $k_a$ by $c^2k_a$. This is the first algebraic filter on Hamiltonian coefficients. Julia now exposes it both as the boost-time symbol $B_a = \frac{1}{2}\partial_a\omega^2$ and as the residual vector $R_a = B_a - c^2k_a$, still only at coefficient-symbol level.

### 27.6 Rotation-Hamiltonian Residual

The Hamiltonian commutes with rotations iff the dispersion is radial on the patch:

$$i[h, J_{ab}] = k_b\partial_a\omega - k_a\partial_b\omega.$$

Equivalently,

$$k_b\partial_a\omega^2 - k_a\partial_b\omega^2 = 0$$

where $\omega > 0$. Hypercubic lattice symbols usually fail this globally on the Brillouin zone but may satisfy it to leading order at a low-energy minimum. The Julia helper records the sampled antisymmetric matrix

$$2(k_bB_a - k_aB_b), \qquad B_a = \frac{1}{2}\partial_a\omega^2,$$

so the factor of two follows the boost-time convention instead of being re-derived independently.

## 27.7 Boost-Boost Residual

Write

$$B_a(k) := \frac{1}{2}\partial_a \omega(k)^2, \qquad R_a(k) := B_a(k) - k_a.$$

Equivalently $B_a = h\partial_a h$. Since

$$K_a = i\left(h\partial_a + \frac{1}{2}\partial_a h\right),$$

the commutator of the first-order differential parts gives

$$i[K_a, K_b] = J_{ab} + R_b X_a - R_a X_b.$$

The leading $+J_{ab}$ is the sign dictated by CONVENTIONS.md (k): CA-11 has $[M_{0a}, M_{0b}] = -M_{ab}$, while the self-adjoint rotation here is $J_{ab} = -A_{M_{ab}}$. Here the possible divergence term vanishes because $R$ is a gradient residual: $\partial_a R_b = \partial_b R_a$. Thus the boost-boost bracket closes to the rotation bracket whenever the boost-time residual $R_a$ vanishes on the patch. The Julia helper records only the sampled coefficient tensor for the first-order residual $R_b X_a - R_a X_b$: its $(a, b, c)$ entry is the coefficient of $X_c$ for the ordered boost pair $(a, b)$. It is not a finite-grid coordinate commutator and does not check the boost-boost operator bracket. The remaining caveat is analytic, not algebraic: all of these are unbounded operators and still require a common self-adjointness/core theorem before this becomes a continuum representation statement.

## 27.8 Second Quantization

Let

$$\mathcal{F}_{\mathrm{alg}}(\mathcal{D}_0) := \bigoplus_{n=0}^{\mathrm{fin}} \mathcal{D}_0^{\otimes_s n} \subset \mathcal{F}_+\big(L^2(\mathcal{U}, dk)\big)$$

be the algebraic finite-particle core. The formal many-body candidates are

$$H = d\Gamma(h), \quad P_a = d\Gamma(p_a), \quad J_{ab} = d\Gamma(J_{ab}^{(1)}), \quad K_a = d\Gamma(K_a^{(1)}),$$

with a separate vacuum-energy convention if absolute Hamiltonian values are used. The formal core identity is

$$[d\Gamma(A), d\Gamma(B)]\psi = d\Gamma([A, B])\psi, \qquad \psi \in \mathcal{F}_{\mathrm{alg}}(\mathcal{D}_0),$$

whenever the one-particle commutator identity is already established on $\mathcal{D}_0$ and the displayed products preserve that algebraic domain. This is a local $d\Gamma$ bookkeeping statement on the algebraic finite-particle core, not a Julia-checked invariant and not a self-adjoint many-body representation theorem. The current tests do not lift residuals to Gaussian Fock space; domain/self-adjointness for the unbounded $d\Gamma$ generators and the sourced OAR/mass-shell bridge remain open proof obligations.

# 28 Lorentz Residual Conditions on Gaussian Coefficients

## 28.1 Status, Sources, and Dependencies

**Status: checked local coefficient calculus.** The sourced massive free scalar model supplies the target dispersion. The general coefficient residuals are local Taylor calculations from $\omega_\epsilon(k)^2 = \sum_r V_r e^{i\epsilon k\cdot r}$. The examples in this shard are checked in Julia; the massless doubler is only a coefficient-level rejection witness, not a positive-dispersion generator example.

Dependencies: CA-24–CA-26 and CONVENTIONS.md (j).

## 28.2 Gradient Residual

For a scalar coefficient family,

$$\omega_\epsilon(k)^2 = \sum_r V_r e^{i\epsilon k\cdot r}.$$

The boost-time residual from CA-26 is

$$R_a(k) = \frac{1}{2}\partial_a\omega_\epsilon(k)^2 - c^2 k_a.$$

Explicitly,

$$\frac{1}{2}\partial_a\omega_\epsilon(k)^2 = \frac{i\epsilon}{2}\sum_r r_a V_r e^{i\epsilon k\cdot r}.$$

When $V_r = V_{-r}$, this is real and can be written as a sine sum:

$$\frac{1}{2}\partial_a\omega_\epsilon(k)^2 = -\epsilon\sum_{r>0} r_a V_r \sin(\epsilon k\cdot r),$$

where $r > 0$ denotes any fixed choice of one representative from each $\{r, -r\}$ pair.

## 28.3 Low-Energy Hessian Condition

At a symmetric minimum $k = 0$, the first derivative vanishes. The Hessian is

$$\partial_a\partial_b\omega_\epsilon(0)^2 = -\epsilon^2\sum_r r_a r_b V_r.$$

Therefore the quadratic Lorentz-speed condition is

$$-\frac{\epsilon^2}{2}\sum_r r_a r_b V_r = c^2\delta_{ab}.$$

This is the first finite-dimensional coefficient test. In compiler form:

$$\mathsf{HessResidual}_{ab} := -\frac{\epsilon^2}{2}\sum_r r_a r_b V_r - c^2\delta_{ab}.$$

It must vanish at the chosen low-energy point before the model can be routed to a Klein-Gordon continuum target.

It is not a positivity or patch certificate. The Hessian only sees the Taylor coefficient at the chosen point; a finite-grid symbol can still be negative elsewhere or have additional low-energy minima.

### 28.4 Nearest-Neighbour Klein-Gordon

For

$$V_0 = m^2 + \frac{2d}{\epsilon^2}, \qquad V_{\pm e_a} = -\frac{1}{\epsilon^2},$$

one obtains

$$-\frac{\epsilon^2}{2}\sum_r r_a r_b V_r = \delta_{ab}.$$

The gradient residual is the stronger finite-$\epsilon$ expression

$$R_a(k) = \frac{\sin(\epsilon k_a)}{\epsilon} - k_a,$$

which vanishes as $O(\epsilon^2 k_a^3)$ for fixed physical momentum.

### 28.5 Anisotropic Counterexample

Let the nearest-neighbour coefficients carry different speeds $c_a$:

$$\omega_\epsilon(k)^2 = m^2 + \frac{2}{\epsilon^2}\sum_a c_a^2(1 - \cos(\epsilon k_a)).$$

Then

$$-\frac{\epsilon^2}{2}\sum_r r_a r_b V_r = c_a^2 \delta_{ab}.$$

Unless all $c_a$ agree with the target speed, the boost-time residual cannot match a single Lorentz cone. The Julia test suite checks this as a deliberate failure case, not as an approximate success.

### 28.6 Hessian-Passing Instability Witness

In $d = 1$, at $\epsilon = 1$, consider the scalar coefficients

$$V_0 = 0, \qquad V_{\pm 1} = 1, \qquad V_{\pm 2} = -\frac{1}{2}.$$

The symbol is

$$\omega(k)^2 = 2\cos k - \cos(2k).$$

Its low-energy Hessian data pass the speed-one condition:

$$-\frac{1}{2}\sum_r r^2 V_r = 1.$$

However, the $L = 4$ periodic sample has values

$$\omega(0)^2 = 1, \quad \omega(\pi/2)^2 = 1, \quad \omega(\pi)^2 = -3, \quad \omega(3\pi/2)^2 = 1.$$

The checked minimum-data helper records the negative minimum $-3$, with uncentered label 2 and momentum $\pi$, plus centered label $-2$ and momentum $-\pi$. Enabling its nonnegative finite-grid gate makes the helper raise an error. Thus the Hessian residual can vanish while the finite periodic symbol is unstable.

## 28.7 Doubler Counterexample

The Hessian condition is necessary but not sufficient. The symbol

$$\omega_\epsilon(k)^2 = m^2 + \frac{1}{\epsilon^2}\sum_a \sin^2(\epsilon k_a)$$

has the correct quadratic Hessian at $k = 0$ when $c = 1$. The checked example sets $m = 0$, so it is massless and has zero modes; by CONVENTIONS.md (j), this is outside the current positive-dispersion generator theorem. Its role is coefficient-level rejection: on even periodic grids it has multiple minima at Brillouin-zone points with $\sin(\epsilon k_a) = 0$. Therefore the compiler also needs a *single-patch* or *sector-selection* condition before a continuum relativistic scalar target is attached.

The current tests record this in $d = 1, 2, 3$ on the $L_a = 4$ periodic grids: the Hessian residual still vanishes at speed 1, while `symbol_minimum_data` finds $2^d$ zero minima with each centered label coordinate in $\{0, -2\}$. This is a finite-grid witness of extra patches, not a continuum species-count theorem.

## 28.8 Compiler Rule

For the scalar Gaussian tier, a Lorentz-target compile attempt must emit at least:

| | |
|---|---|
| mass term | $\omega(0)^2$, |
| speed matrix | $-\frac{\epsilon^2}{2}\sum_r r_a r_b V_r$, |
| boost residual | $R_a(k)$, |
| rotation residual | $k_b\partial_a\omega^2 - k_a\partial_b\omega^2$, |
| boost-boost residual | $R_b X_a - R_a X_b$, |
| patch data | finite-grid minimum value, labels, momenta, and count. |

Only the mass and speed entries are finite Taylor data. The rotation and boost-boost entries are now sampled smooth-symbol residuals in Julia, with the boost-boost entry represented as first-order $X_c$-coefficient data rather than as a finite-grid operator commutator. The residual and patch entries are what prevent an anisotropic or doubler model from being mislabeled as a single Klein-Gordon field.

# 29 Numerical Verification Suite for Gaussian Boson Examples

## 29.1 Status, Sources, and Dependencies

**Status: checked Julia shard.** This shard records the numerical invariants now implemented for the scalar Gaussian tier. The checks are not continuum proofs; they are finite algebraic witnesses that the coefficient compiler is computing the intended symbols and residuals. The massless doubler check is a coefficient-level rejection test, not a Fock-generator check. The finite periodic checks below do not assert an exact differential $[X_a, P_b] = i\delta_{ab}$ commutator.

The current numerical surface is deliberately narrow: it checks scalar coefficient structure, finite-periodic symbol values, finite stiffness spectra, assigned Fourier eigenvectors, periodic minimum data, and the implemented coefficient residuals/Hessians, including sampled rotation-Hamiltonian and boost-boost residual data on smooth-symbol momenta. It also checks centered periodic momentum labels for branch-aware low-energy windows, a loud finite-grid nonnegativity gate when requested, and test-local central finite-difference boost-time derivative oracles. It does not test finite boost generators, finite rotation generators, coordinate-generator commutators, or full generator closure.

Dependencies: CA-24–CA-27, INDEX.md, and CONVENTIONS.md (j).

## 29.2 Why These Tests

The numerical layer is not allowed to assert that a plot "looks relativistic". Each check must compare two independently constructed objects:

$$\text{real-space coefficient data} \quad \longleftrightarrow \quad \text{momentum-space symbol data.}$$

For Gaussian systems this is unusually powerful because the finite periodic stiffness matrix is exactly diagonalized by Fourier modes.

The boost-time derivative oracle follows the same rule. The production boost-time helper is compared to $\frac{1}{2}\nabla_k \omega(k)^2$ obtained by central finite differences of the separate scalar-symbol evaluator. Those tests use generic symmetric finite-range coefficient sets in $d = 1, 2, 3$, with off-axis terms in the higher-dimensional cases and sample momenta away from high-symmetry points.

The rotation and boost-boost residual tests are also smooth-symbol checks. They sample coefficient formulas at named momenta and compare residual sizes or tensor entries; they do not introduce $X_a = i\partial_{k_a}$ as a finite periodic matrix.

## 29.3 Numerical Tolerance Policy

CONVENTIONS.md (j) fixes the tolerance names used by this shard. With the `GAUSSIAN_` prefix and `_ATOL`/`_RTOL` suffix understood, the default pairs are

| | |
|---|---|
| `SYMBOL_IMAG` | $(10^{-10}, 10^{-10})$, |
| `SYMBOL_VALUE` | $(10^{-10}, 10^{-10})$, |
| `EIGENVALUE` | $(10^{-10}, 10^{-10})$, |
| `MINIMUM_COUNT` | $(10^{-10}, 10^{-10})$, |
| `SMALL_SPACING_RESIDUAL` | $(5 \cdot 10^{-4}, 10^{-10})$. |

The symbol-imaginary validator compares the complex symbol with its real part after the dimension and spacing preconditions are checked. The small-spacing pair is a named finite-sample tolerance, not a continuum error bound. The finite-difference derivative oracle is test-local and uses $(2 \cdot 10^{-8}, 2 \cdot 10^{-8})$; it is not part of the production tolerance surface.

### 29.4 Structural Coefficient Validation

The scalar Gaussian helpers now fail closed by default on coefficient data outside CONVENTIONS.md (j). The validator requires nonempty data, integer offsets, a common dimension $d \in \{1, 2, 3\}$, real coefficients, and aggregate paired equality $V_r = V_{-r}$ after duplicate offsets are summed. Negative tests cover a missing partner, unequal paired aggregates, complex coefficients, dimension mismatch, and non-integer offsets.

### 29.5 Periodic Stiffness Matrix

For finite periodic sizes $L_1, \ldots, L_d$, define the stiffness matrix

$$V_{xy} = \sum_r V_r \, \mathbf{1}_{y=x+r \bmod L}.$$

For the Fourier vector

$$\varphi_k(x) = |\Lambda|^{-1/2} e^{i\epsilon k \cdot x},$$

one has the exact identity

$$(V\varphi_k)(x) = \left( \sum_r V_r e^{i\epsilon k \cdot r} \right) \varphi_k(x) = \omega_\epsilon(k)^2 \varphi_k(x).$$

Therefore each dual label $n$ gives a tested eigenvector with its assigned symbol value, and the sorted eigenvalues of the finite stiffness matrix must equal the sorted symbol values on the periodic Brillouin grid. These finite Fourier invariants are checked in Julia for $d = 1, 2, 3$, including mixed odd/even sizes with every $L_a > 2R$ and an off-axis symmetric pair $(1, 1), (-1, -1)$. They are not finite periodic first-moment or momentum-coordinate commutator tests; those require a branch, sawtooth, Fourier-interpolation, or other coordinate convention before they can become numerical claims.

The eigenvector suite also includes a one-sided shift sentinel to pin the implementation direction $y = x + r$. That sentinel is not a scalar Gaussian Hamiltonian claim. The strict default rejects it, and the test reaches this low-level matrix path only through an explicit non-validating keyword.

The eigenvalue oracle deliberately keeps the uncentered grid helper, because the tested symbol spectra are branch-insensitive. The separate centered grid helper returns the centered integer dual label and its physical representative

$$k_a = \frac{2\pi n_a}{\epsilon L_a}.$$

For even $L = 2r$, the branch is $n_a \in \{-r, \ldots, 0, \ldots, r - 1\}$, matching the centered dual lattice in the OAR source. Thus, at $L = 4$ and $\epsilon = 1$, the uncentered representative $3\pi/2$ is recorded on the centered branch as $n = -1$ and $k = -\pi/2$. This is only a momentum-label helper; it is not a periodic position-coordinate or generator construction.

### 29.6 Implemented Example Systems

**Nearest-neighbour Klein-Gordon.** The tests check the coefficient symbol against

$$m^2 + \frac{2}{\epsilon^2} \sum_a (1 - \cos(\epsilon k_a)),$$

the boost-time symbol against $\sin(\epsilon k_a)/\epsilon$, the residual vector against $\sin(\epsilon k_a)/\epsilon - k_a$, and the low-energy Hessian residual against zero. The sampled rotation and boost-boost residual helpers are

small on fixed low-energy momenta as $\epsilon$ is made small. The boost-time helper is also checked on non-Klein-Gordon finite-range symbols by the independent finite-difference oracle above.

**Anisotropic cones.** The tests build coefficients with speeds $(c_1, c_2, c_3)$. They verify that the Hessian is

$$2\,\mathrm{diag}(c_1^2, c_2^2, c_3^2)$$

and that the Lorentz residual is nonzero when the target speed is 1. A separate sampled residual test checks that anisotropy also gives nonzero rotation-Hamiltonian and boost-boost residual data at a generic smooth momentum.

**Hessian-passing unstable scalar.** The tests build the one-dimensional coefficient witness

$$V_0 = 0, \qquad V_{\pm 1} = 1, \qquad V_{\pm 2} = -\frac{1}{2}.$$

Its speed-one Hessian residual vanishes, but finite periodic sampling at $L = 4$ has minimum $-3$. The minimum-data helper records one minimum at location 3, uncentered label 2, centered label $-2$, uncentered momentum $\pi$, and centered momentum $-\pi$. Requesting a nonnegative finite-grid symbol raises an error, so this example cannot pass as a stable scalar symbol.

**Doubler symbol.** The tests build

$$\omega_\epsilon(k)^2 = m^2 + \epsilon^{-2} \sum_a \sin^2(\epsilon k_a)$$

with $m = 0$. In $d = 1, 2, 3$, the speed-one Hessian residual vanishes on the chosen patch, but the $L_a = 4$ periodic grid has $2^d$ zero minima with centered label coordinates in $\{0, -2\}$. This catches the case where the Hessian at one minimum is correct but the compiler would need a multi-species, sector-selection, or zero-mode policy output. It is deliberately outside the positive-dispersion generator statements. The J6 symbol-residual test also samples a doubler momentum where the rotation residual is globally nonzero and an extra zero where $R_a$ is nonzero relative to the selected $k = 0$ branch.

### 29.7 Current Julia Surface

The checked helper layer is:

| | |
|---|---|
| `kg_nearest_neighbor_coefficients` | standard scalar KG coefficients, |
| `anisotropic_kg_nearest_neighbor_coefficients` | anisotropic coefficient witness, |
| `doubler_quadratic_coefficients` | coefficient-level doubler witness, |
| `validate_scalar_coefficients` | structural scalar-coefficient validator, |
| `scalar_quadratic_symbol` | $\sum_r V_r e^{i\epsilon k\cdot r}$, |
| `boost_time_symbol_from_coefficients` | $\frac{1}{2}\nabla\omega^2$, |
| `boost_time_residual_from_coefficients` | $\frac{1}{2}\nabla\omega^2 - c^2 k$, |
| `rotation_hamiltonian_residual_from_coefficients` | $k_b\partial_a\omega^2 - k_a\partial_b\omega^2$, |
| `boost_boost_residual_coefficients_from_coefficients` | coefficients of $R_bX_a - R_aX_b$, |
| `nearest_neighbor_integrated_energy` | 1D integrated A4 current symbol, |
| `current_symbol_low_energy_hessian_from_coefficients` | $\partial_a\partial_b\omega^2(0)$, |
| `lorentz_hessian_residual` | $\frac{1}{2}\mathrm{Hess}(\omega^2)(0) - c^2 I$, |
| `periodic_stiffness_matrix` | finite periodic real-space matrix, |
| `periodic_momentum_grid` | uncentered spectrum grid, |
| `centered_periodic_momentum_grid` | centered labels and momenta, |
| `periodic_fourier_vector` | Fourier vector for a dual label, |
| `periodic_symbol_values` | finite Brillouin-grid eigenvalue oracle, |
| `symbol_minimum_data` | finite-grid minimum value, labels, momenta, and count, |
| `count_periodic_symbol_minima` | finite-grid minima counter. |

This list is a coverage boundary, not a generator-algebra interface: no helper currently constructs a finite coordinate generator, a finite boost, a finite rotation, or their commutator table.

### 29.8 Next Numerical Milestones

Proposed next suite additions, none claimed as achieved here, are:

1. systematic sampled residual scans on shrinking low-energy windows,
2. comparison of open-boundary first moments with periodic symbol tests.

Any finite periodic coordinate-generator version of the first-moment comparison requires a branch or Fourier-interpolation convention before it can be promoted beyond a local experiment.

# 30 Gaussian Boson Real-Space Energy Density

## 30.1 Status, Sources, and Dependencies

**Status: convention, not a theorem.** This shard fixes how the scalar Gaussian Hamiltonian is split into site-attached real-space energy densities. It preserves the total Hamiltonian used in the sourced nearest-neighbour scalar model and in CA-24–CA-28, but it does not claim a continuum tensor object, a conservation law, or covariance.

Dependencies: CA-23–CA-28 and CONVENTIONS.md (j), (l).

## 30.2 Cell Volume and Notation

Let the hypercubic lattice spacing be $\epsilon$, so one spatial cell has volume $\epsilon^d$. In this shard $h_x$ is the physical energy density attached to the cell at $x$:

$$H_\epsilon = \epsilon^d \sum_x h_x.$$

The integrated site energy is

$$e_x := \epsilon^d h_x, \qquad H_\epsilon = \sum_x e_x.$$

This distinction matters because the generic first-moment notation of CONVENTIONS.md (h) writes $H = \sum_x h_x$; for the Gaussian block, the object to insert there is $e_x$, not the density $h_x$.

For fields $\Phi_x, \Pi_x$ and a real symmetric translation-invariant kernel $V_r = V_{-r}$, use

$$H_\epsilon = \epsilon^d \left( \frac{1}{2} \sum_x \Pi_x^2 + \frac{1}{2} \sum_{x,r} \Phi_x V_r \Phi_{x+r} \right).$$

For the nearest-neighbour Klein-Gordon source, the renormalized coefficients are

$$V_0 = m^2 + \frac{2d}{\epsilon^2}, \qquad V_{\pm e_a} = -\frac{1}{\epsilon^2},$$

which give the dispersion recorded in CA-24.

## 30.3 Equal-Endpoint Bond Split

The chosen local density is

$$h_x = \frac{1}{2}\Pi_x^2 + \frac{1}{2} V_0 \Phi_x^2 + \frac{1}{4} \sum_{r \neq 0} V_r \big( \Phi_x \Phi_{x+r} + \Phi_{x-r} \Phi_x \big).$$

Each nonzero displacement bond is assigned half to each endpoint cell. Summing over $x$ gives

$$\sum_x h_x = \frac{1}{2} \sum_x \Pi_x^2 + \frac{1}{2} \sum_{x,r} \Phi_x V_r \Phi_{x+r},$$

so this convention reproduces the total scalar Hamiltonian above after the cell-volume factor is restored.

The $q, p$ notation in CA-24–CA-28 absorbs the cell volume into the coefficient bookkeeping. In that notation the same displayed formula defines $e_x$, and the physical density is $h_x = e_x/\epsilon^d$.

## 30.4 Boundary Policy

The default exact identities are infinite-lattice identities with summable tails, or finite periodic total-energy identities. On a finite open region, the density is obtained from the finite Hamiltonian actually named: include only bonds present in that Hamiltonian, and share each included bond equally between its included endpoints. No exterior ghost sites, omitted half-bonds, or boundary counterterms are implicit.

For periodic finite lattices, wrap-around bonds are included in the total energy split. This does not create a periodic position convention. Any periodic first moment still needs the branch, sawtooth, or interpolation policy required by CONVENTIONS.md (j).

## 30.5 Vacuum-Energy Policy

The density above is bare. No normal-ordering or vacuum subtraction is implicit. Once a state $\omega$ is named, the expectation-subtracted version may be recorded as

$$:h_x:_\omega := h_x - \omega(h_x)\mathbf{1}.$$

This changes $H_\epsilon$ by a scalar and therefore drops out of commutators, but it matters for absolute energies, expectation values, and Fourier modes of quadratic densities. The local subtraction source is the free-fermion Koo-Saleur setting, where the scaling-limit representation relies on subtracting vacuum or ground-state expectation values; this shard only records the analogous policy needed before such quantities are used in the scalar Gaussian block.

## 30.6 Improvement Policy

The default convention has no divergence improvement:

$$h'_x = h_x$$

unless a later shard explicitly names an edge field $b$, a discrete divergence $(\nabla_\epsilon \cdot b)_x$, and a boundary convention. Such a rewrite

$$h'_x = h_x + (\nabla_\epsilon \cdot b)_x$$

may leave the total energy unchanged on a periodic lattice or for decaying fields, but it is not harmless for position-weighted generators. A discrete summation-by-parts calculation shifts first moments by an edge-field term plus boundary contributions. For example, in one dimension with

$$(\nabla_\epsilon b)_n = \frac{b_{n+1/2} - b_{n-1/2}}{\epsilon}, \qquad x_n = \epsilon n,$$

one has

$$\epsilon \sum_n x_n (\nabla_\epsilon b)_n = -\epsilon \sum_n b_{n+1/2} + \text{boundary terms}.$$

Therefore improvements are separate data, not gauge choices to be silently modded out.

## 30.7 Compiler Consequence

A scalar Gaussian real-space generator proposal must now carry the tuple

$$\big(\epsilon^d,\ h_x,\ e_x = \epsilon^d h_x,\ \text{boundary policy},\ \text{subtraction policy},\ \text{improvement policy}\big).$$

The total symbol $\omega(k)^2$ remains sufficient for CA-24–CA-28, but it is not sufficient once a construction uses position weights or local quadratic modes. Those later constructions must cite this density convention or perform an explicit convention sweep.

# 31 One-Dimensional Gaussian Boson Stress-Energy Candidates

## 31.1 Status, Sources, and Dependencies

**Status: proposal with checked $T_{01}$ seed.** This shard fixes a candidate dictionary for the finite open one-dimensional scalar Gaussian chain. Only the energy-current entry has a checked local witness. No convergence theorem, no continuum stress tensor, and no Virasoro representation is claimed.

Dependencies: CA-12, CA-23–CA-29, CONVENTIONS.md (j), (k), (l), and the A4 finite open-chain current tests.

## 31.2 Continuum Motivation and Boundary

The continuum notation $T_{\mu\nu}$ is used here as a target shape, not as an imported lattice theorem. Weinberg supplies the QFT motivation that spacetime translations lead to conserved currents grouped as an energy-momentum tensor and that translation generators are spatial integrals of time components. The same registered OCR also records two caveats that matter for the lattice dictionary: the canonical raised tensor need not be symmetric, while the Belinfante tensor is a modified conserved tensor with the same integrated translation generators and symmetry. The OCR around formulas is noisy, so this shard uses those anchors only for motivation and caveats, not for exact component signs. The self-adjoint commutator signs for any later Poincare comparison are the project convention in CONVENTIONS.md (k), not raw OCR import.

Schottenloher's Axiom 5 is narrower but sharper for the CFT endpoint: in a two-dimensional conformal field theory there are four fields $T_{\mu\nu}$, $\mu, \nu \in \{0, 1\}$, with symmetry and conservation. The later Luescher–Mack theorem gives tracelessness and two Virasoro mode families under the full Axioms 1–5. We do not use those CFT conclusions to define the lattice $T_{10}$ or $T_{11}$. They are acceptance targets for a future continuum proof.

## 31.3 Finite Open Chain Setup

Work first with a finite open chain of $N$ cells and spacing $\epsilon$. The CA-29 density convention distinguishes the physical density $h_j$ from the integrated cell energy

$$e_j = \epsilon h_j, \qquad H_\epsilon = \sum_{j=1}^{N} e_j.$$

Finite commutators use $e_j$. This avoids inserting an accidental factor of $\epsilon$ into a discrete continuity identity.

For the nearest-neighbour scalar Gaussian chain, the A4 helper implements the CA-29 equal-endpoint split of the finite Hamiltonian: kinetic and onsite terms belong to the site, and each present bond $q_j q_{j+1}$ is split equally between its two endpoints. Open-chain means no exterior ghost cells, no missing half-bonds, and no boundary counterterms are implicit.

## 31.4 Candidate Dictionary

**$T_{00}^{\mathrm{lat}}$.** The lattice energy-density entry is the CA-29 density:

$$T_{00}^{\mathrm{lat}}(j) := h_j.$$

For exact finite open-chain algebra, the cell-integrated version is

$$E_j := e_j = \epsilon h_j.$$

Thus $T_{00}^{\mathrm{lat}}$ is the physical density label, while $E_j$ is the object that appears in the tested quadratic commutators.

$T_{01}^{\mathrm{lat}}$**.** The energy-current or energy-flux seed lives on open-chain edges:

$$J_{j+1/2} := i[e_j, e_{j+1}], \qquad 1 \le j < N.$$

Set endpoint currents to zero,

$$J_{1/2} = J_{N+1/2} = 0.$$

The checked finite continuity identity is

$$i[H_\epsilon, e_j] = J_{j-1/2} - J_{j+1/2}, \qquad 1 \le j \le N.$$

This is the only stress-energy component in this shard with a Julia witness. It is the lattice candidate corresponding to $T_{01}$ in the cell-integrated continuity law

$$\frac{d}{dt} \int_{\text{cell } j} T_{00}\, dx = T_{01}(x_{j-1/2}) - T_{01}(x_{j+1/2}).$$

No continuum normalization or convergence of $J_{j+1/2}$ is claimed.

$T_{10}^{\mathrm{lat}}$**.** Do not identify $T_{10}^{\mathrm{lat}}$ with $T_{01}^{\mathrm{lat}}$ at this stage. Define it instead as the future integrated momentum-density comparison target:

$$M_j \quad \text{with} \quad P^{\mathrm{lat}} = \sum_j M_j.$$

The required next witness is that the chosen $M_j$, or at least its total $P^{\mathrm{lat}}$, matches the Gaussian one-particle flat-symbol momentum target after the appropriate $\mathrm{d}\Gamma(k)$ comparison is made. That is the S3 task, not an assumption in this shard.

$T_{11}^{\mathrm{lat}}$**.** Define $T_{11}^{\mathrm{lat}}$ operationally as the future current of the chosen momentum density. Once $M_j$ is fixed, the candidate stress component must come with edge terms $S_{j+1/2}$ and a checked equation

$$i[H_\epsilon, M_j] = S_{j-1/2} - S_{j+1/2}$$

with an explicit boundary convention. Until that witness exists, there is no lattice $T_{11}$ formula. In particular, $T_{11}^{\mathrm{lat}}$ is not defined as $-T_{00}^{\mathrm{lat}}$, not derived from tracelessness, and not inferred from CFT holomorphic factorization.

## 31.5 Boundary, Subtraction, and Improvement

This candidate dictionary is open-boundary first. Endpoint currents are part of the statement, not small errors:

$$i[H_\epsilon, e_1] = -J_{3/2}, \qquad i[H_\epsilon, e_N] = J_{N-1/2}.$$

Periodic total-energy density splits are allowed by CA-29, but periodic first moments and periodic coordinate-dependent generator tests remain blocked until CONVENTIONS.md (j) fixes a coordinate branch, sawtooth, Fourier interpolation, or equivalent policy.

The default density is bare. A state-dependent subtraction $:h_j:_\omega = h_j - \omega(h_j)\mathbf{1}$ may later be named; it changes the Hamiltonian by a scalar and does not change commutators, but it is part of any claim about absolute energy or density modes. The default improvement is zero. Any later replacement $h'_j = h_j + (\nabla_\epsilon b)_j$ must name the edge field, discrete divergence, and boundary convention before first moments or currents are recomputed.

### 31.6 Compiler Payload and Acceptance

A compiler output that cites this shard must carry:

| | |
|---|---|
| density convention | $h_j,\ e_j = \epsilon h_j,$ CA-29 split, |
| boundary policy | finite open chain, $J_{1/2} = J_{N+1/2} = 0$, |
| subtraction policy | bare unless a state $\omega$ is named, |
| improvement policy | $b = 0$ unless a discrete divergence is named, |
| checked $T_{01}$ witness | $i[H_\epsilon, e_j] = J_{j-1/2} - J_{j+1/2}$. |

The acceptance boundary is equally explicit. The checked witnesses are the finite open-chain continuity residual in `src/GaussianBosonCurrents.jl` and `test/runtests.jl`, plus the CA-31 translation-invariant current-symbol equality for the nearest-neighbour Klein-Gordon orientation. Still pending are the finite open-chain wave-packet comparison with the flat $\mathrm{d}\Gamma(k)$ momentum target; the $T_{11}$ witness consisting of a chosen momentum density $M_j$ and checked current $S_{j+1/2}$; and the continuum witness: scaling maps, domains, state sequence, and covariance convergence. Without those witnesses, the dictionary remains a proposal with checked $T_{01}$ evidence, not a stress-tensor theorem.

# 32 Gaussian Current-Symbol Equivalence in One Dimension

## 32.1 Status, Sources, and Dependencies

**Status: checked translation-invariant symbol bridge.** This shard adds the first bridge between the finite real-space current convention and the momentum-symbol layer. It is deliberately one-dimensional and nearest-neighbour. The checked statement is the equality of the translation-invariant integrated-current symbol with the CA-26 boost-time symbol for the scalar Klein-Gordon coefficients. No finite open-chain momentum operator, no finite periodic first moment, no $d\Gamma(k)$ theorem, and no continuum convergence statement is claimed.

Dependencies: CA-24, CA-26, CA-29–CA-30, and CONVENTIONS.md (j), (k), (l).

## 32.2 Real-Space Current Orientation

The finite open-chain A4 witness fixes the local current on an open bond by

$$J_{j+1/2} = i[e_j, e_{j+1}]$$

with endpoint currents zero. For a nearest-neighbour scalar chain with bond coefficient $b$, the checked closed form is

$$J_{j+1/2} = \frac{b}{2}\big(q_{j+1}p_j - q_j p_{j+1}\big).$$

This sign is not adjustable in this shard: it is the CONVENTIONS.md (l) orientation already used in the open-chain continuity test.

To compare with the boost-time symbol, pass from the local edge current to the first-moment integrated current density. In uniform spacing $\epsilon$, the nearest-neighbour first-moment coefficient contributes the spacing factor $\epsilon$. The translation-invariant integrated current therefore has symbol

$$\mathcal{J}_\epsilon(k) = -\epsilon b \sin(\epsilon k).$$

This is a Fourier-symbol identity for the infinite or translation-invariant nearest-neighbour expression. It does not introduce a periodic coordinate branch and does not construct an open finite-chain momentum operator.

## 32.3 Klein-Gordon Equality

For the one-dimensional nearest-neighbour lattice Klein-Gordon coefficients,

$$b = -\epsilon^{-2}, \qquad \omega_{\epsilon,m}(k)^2 = m^2 + 2\epsilon^{-2}(1 - \cos(\epsilon k)).$$

The integrated-current symbol becomes

$$\mathcal{J}_\epsilon(k) = \frac{\sin(\epsilon k)}{\epsilon}.$$

The CA-26 boost-time symbol is

$$B_\epsilon(k) = \frac{1}{2}\partial_k \omega_{\epsilon,m}(k)^2 = \frac{\sin(\epsilon k)}{\epsilon}.$$

Thus, at this checked symbol level,

$$\mathcal{J}_\epsilon(k) = B_\epsilon(k).$$

The mass drops out because the current and the derivative of the onsite mass term are both insensitive to $m^2$.

### 32.4 Julia Witness

The helper

`nearest_neighbor_integrated_energy_current_symbol`

implements

$$-\epsilon b \sin(\epsilon k)$$

for one-dimensional data only. The Klein-Gordon wrapper inserts $b = -\epsilon^{-2}$. The tests compare that wrapper against

`kg_lattice_boost_time_symbol`

for several momenta and spacings. They also include a sign sentinel: for positive $k$ and the Klein-Gordon negative bond, the current symbol is positive and agrees with the boost-time target; reversing the bond sign gives a negative symbol and fails the target comparison. A small-spacing check compares the current symbol with the centered continuum momentum representative $k$ near zero using the existing Gaussian small-spacing tolerance.

### 32.5 Open Obligations

This shard closes only the first symbol-level bridge. The following remain pending:

1. a finite open-chain wave-packet comparison with boundary residuals,
2. an identification with the one-particle $\mathrm{d}\Gamma(k)$ momentum target,
3. a periodic first-moment construction, which still needs a branch policy,
4. a scaling or continuum convergence theorem.

Any compiler output citing this shard must therefore say "current-symbol equality" rather than "finite-chain momentum" or "continuum stress tensor".

# 33 Higher-Dimensional Gaussian Cell-Current Proposal

## 33.1 Status, Sources, and Dependencies

**Status: proposal and compiler-input specification.** This shard states the support data and local equations a higher-dimensional Gaussian stress-energy candidate must carry in $d = 2, 3$. It is not a theorem, does not define a global orientation convention, and makes no convergence claim.

Dependencies: CA-13, CA-23, CA-29–CA-31, and CONVENTIONS.md (j), (k), (l).

## 33.2 Compiler Input: Oriented Support Data

For each finite or infinite hypercubic support in $d = 2$ or $d = 3$, the compiler input must include an oriented incidence structure:

$$\mathcal{C}, \qquad \mathcal{F}, \qquad \sigma(c, f) \in \{-1, 0, +1\}.$$

Here $\mathcal{C}$ is the set of energy cells, attached to sites $x_c \in \epsilon\mathbb{Z}^d$. The set $\mathcal{F}$ contains oriented codimension-one faces. In $d = 2$, these faces are oriented edges between neighbouring square cells. In $d = 3$, they are oriented plaquettes between neighbouring cube cells. If an interior face $f$ is oriented from its minus cell $c_-(f)$ to its plus cell $c_+(f)$, set

$$\sigma(c_-(f), f) = +1, \qquad \sigma(c_+(f), f) = -1,$$

and set $\sigma(c, f) = 0$ for all other cells. This sign convention is local input for this proposal, not a repository-wide orientation convention.

Boundary faces are separate data. For an open box, $\mathcal{F}_\partial$ contains oriented exterior faces with only one adjacent cell; their incidence sign is recorded as $\sigma_\partial(c, f) = +1$ for outward flux. Setting a boundary flux to zero is then a named boundary condition, not an omitted term. For a periodic support, exterior faces are absent, but any first moment or coordinate-weighted test still needs the branch or Fourier policy required by CONVENTIONS.md (j).

## 33.3 Energy Density and Energy Flux

The energy component starts from CA-29:

$$T_{00}^{\mathrm{lat}}(c) := h_c, \qquad e_c := \epsilon^d h_c, \qquad H_\epsilon = \sum_{c \in \mathcal{C}} e_c.$$

All finite commutator identities use the integrated cell energy $e_c$.

An energy-current candidate is face data

$$J_0(f) \quad (f \in \mathcal{F}),$$

with optional boundary data $J_0^\partial(f)$ on $f \in \mathcal{F}_\partial$. If $f$ has normal direction $a$, then $J_0(f)$ is the proposed lattice $T_{0a}$ flux through that oriented face. The required energy-continuity witness is the cell equation

$$i[H_\epsilon, e_c] + \sum_{f \in \mathcal{F}} \sigma(c, f) J_0(f) + \sum_{f \in \mathcal{F}_\partial} \sigma_\partial(c, f) J_0^\partial(f) = 0.$$

For an open no-leak boundary this specializes to $J_0^\partial(f) = 0$. For a driven or truncated-region diagnostic, the same displayed equation records the boundary residual explicitly. This shard does not propose a universal formula for $J_0(f)$; a later numerical suite must derive or solve for it from the chosen Gaussian density and then test the equation above.

### 33.4 Momentum Densities and Stress Currents

The momentum-density components are additional witnesses:

$$M_a(c) \quad (a = 1, \dots, d), \qquad P_a^{\mathrm{lat}} = \sum_c M_a(c).$$

They are not inferred from the energy flux $J_0(f)$. The reason is already visible in CA-30: $T_{10}$ was kept as a momentum-density target, not as a consequence of $T_{01}$. In higher dimensions this separation is mandatory because rotations require $M_a$ before angular momentum even has a candidate density.

For each chosen momentum density $M_a(c)$, the stress components are its currents. If $f$ has normal direction $b$, write

$$S_a(f) \equiv T_{ab}^{\mathrm{lat}}(f).$$

The required momentum-continuity witness is

$$i[H_\epsilon, M_a(c)] + \sum_{f \in \mathcal{F}} \sigma(c, f) S_a(f) + \sum_{f \in \mathcal{F}_\partial} \sigma_\partial(c, f) S_a^\partial(f) = 0$$

for every $a$ and every cell $c$. Until $M_a$ and $S_a$ are supplied with this witness, there is no lattice $T_{a0}$ or $T_{ab}$ claim.

### 33.5 Boost and Symbol Diagnostics

The boost first moment can be formed from the integrated energy cells:

$$K_a^{\mathrm{lat}} := \sum_c x_c^a e_c.$$

Using the energy-continuity equation, $i[H_\epsilon, K_a^{\mathrm{lat}}]$ reduces by discrete summation by parts to the integrated $a$-normal energy flux plus boundary terms. The higher-dimensional analogue of CA-31 is then a test, not an assumption:

$$\text{integrated } T_{0a} \text{ symbol} \quad \stackrel{?}{=} \quad \frac{1}{2} \partial_a \omega(k)^2.$$

Only after this equality is checked in the relevant local or symbol regime may it be compared with the flat one-particle momentum target $k_a$ and the residuals of CA-26–CA-27. Passing the symbol equality would still not prove a continuum stress tensor.

### 33.6 Rotation and Antisymmetric-Stress Residual

For $a < b$, a rotation candidate requires the momentum-density witnesses:

$$J_{ab}^{\mathrm{lat}} := \sum_c \big( x_c^a M_b(c) - x_c^b M_a(c) \big).$$

This formula is unavailable if only $T_{00}$ and $T_{0a}$ have been supplied. Applying the momentum-continuity equation gives a discrete summation-by-parts diagnostic of the form

$$i[H_\epsilon, J_{ab}^{\mathrm{lat}}] = \epsilon \sum_{f \perp a} S_b(f) - \epsilon \sum_{f \perp b} S_a(f) + \text{boundary terms},$$

with the displayed sign tied to the incidence convention above. In continuum language this is the antisymmetric-stress obstruction: rotation conservation targets $T_{ab} = T_{ba}$, after boundary terms and any spin/improvement data have been named. Weinberg's registered anchors motivate that a symmetric Belinfante tensor can carry the same integrated translations, but the canonical tensor need not be symmetric. Therefore stress symmetry is a target or witness, not an assumption baked into the lattice proposal.

### 33.7 Trace, Improvement, and Boundary Alternatives

Trace diagnostics are also target-dependent. A compiler may report the diagonal stress combination built from $h_c$ and the $S_a(f)$, but it must name the continuum target and metric/sign convention before calling it a trace residual. No massless-CFT tracelessness test, massive Klein-Gordon trace test, or improvement formula is imported here.

The CA-29 improvement policy remains active:

$$h'_c = h_c + (\nabla_\epsilon \cdot b)_c.$$

Any such change must name the edge or face field $b$, the discrete divergence, and the boundary convention. It must also recompute $J_0$, $M_a$, $S_a$, and the first-moment boundary terms; improvements are not silent gauge moves for boosts or rotations.

Two boundary regimes are admissible compiler inputs:

| | |
|---|---|
| open box | use explicit $\mathcal{F}_\partial$ terms and test endpoint/face residuals, |
| periodic box | use no boundary faces, but fix a coordinate branch, sawtooth, or Fourier-interpolation policy before any first-moment test. |

The existing Gaussian numerical layer is periodic-symbol first. The next stress-energy numerical suite, S5 in the action plan, should therefore start with open-boundary local continuity tests or with a separately sourced periodic coordinate construction, and should include anisotropic/doubler failure witnesses at the current-density level.

# 34 First Gaussian Stress-Energy Numerical Suite

## 34.1 Status, Sources, and Dependencies

**Status: checked local-current numerical suite.** This shard records the first stress-energy-specific numerical tests. The checked surface is narrow: finite open-chain $T_{01}$ continuity for the CA-29/CA-30 density-current choice, and a one-dimensional nearest-neighbour current-symbol slope residual. It is not a finite boost construction, not a $\mathrm{d}\Gamma(k)$ theorem, not a periodic first-moment test, and not a continuum stress-tensor claim.

Dependencies: CA-29–CA-32 and CONVENTIONS.md (j), (k), (l).

## 34.2 Finite Open-Chain Continuity

The first suite strengthens the A4 local-current witness instead of replacing it. For each tested open chain, the Julia layer constructs the CA-29 equal-endpoint site energies $e_j$, the total Hamiltonian

$$H = \sum_{j=1}^{N} e_j,$$

and the edge currents

$$J_{j+1/2} = i[e_j, e_{j+1}], \qquad 1 \le j < N.$$

The residual helper checks

$$i[H, e_j] - (J_{j-1/2} - J_{j+1/2}) = 0$$

with endpoint currents $J_{1/2} = J_{N+1/2} = 0$.

The test cases now cover several chain sizes and onsite/bond choices:

$$(N, a, b) \in \{(2, 0.75, -0.5), (3, 2.0, 1.25), (5, 7.25, -1.5), (7, 0.1, -0.875)\}.$$

For each case the suite asserts:

1. $\sum_j e_j = H$,
2. $H_{qq}$ is the expected tridiagonal matrix with diagonal $a$ and bond $b$,
3. $H_{pp} = I$,
4. $J_{j+1/2}$ equals the closed form $\frac{b}{2}(q_{j+1}p_j - q_j p_{j+1})$,
5. $J_{j+1/2}$ is independent of the onsite coefficient,
6. all continuity residual matrices vanish,
7. $i[H, e_1] = -J_{3/2}, \quad i[H, e_j] = J_{j-1/2} - J_{j+1/2}, \quad i[H, e_N] = J_{N-1/2}$.

The explicit endpoint tests are the important stress-energy addition: boundary signs are part of the local $T_{01}$ claim, not discarded finite-size errors. The suite also checks that a one-site chain cannot produce a bond current and that an out-of-range open bond index raises an error.

## 34.3 Current-Symbol Slope Residual

The second suite is the smallest symbol-level current-density failure witness currently justified by CA-31. For the one-dimensional nearest-neighbour symbol

$$\mathcal{J}_\epsilon(k) = -\epsilon b \sin(\epsilon k),$$

the small-$k$ slope is

$$\mathcal{J}_\epsilon'(0) = -\epsilon^2 b.$$

The new helper returns the speed residual

$$\rho_J(b, \epsilon, c) = -\epsilon^2 b - c^2.$$

It validates positive spacing, positive target speed, and real bond data before returning a number.

For the speed-$c$ nearest-neighbour Klein-Gordon bond

$$b = -c^2\epsilon^{-2},$$

the residual is zero. The tests assert this for several spacings and speeds. They also assert failure witnesses:

$$b = -\epsilon^{-2} \quad \text{fails when } c \neq 1,$$

wrong bond magnitude gives the expected nonzero residual, and reversing the bond sign gives

$$\rho_J(c^2\epsilon^{-2}, \epsilon, c) = -2c^2.$$

These are current-density symbol failures for the CA-31 orientation. They are not Hessian-only failures and they do not use a periodic coordinate branch.

### 34.4 What This Falsifies

The suite falsifies three classes of local-current mistakes:

1. incorrect open-boundary signs in $i[H, e_j] = J_{j-1/2} - J_{j+1/2}$,
2. wrong orientation in $J_{j+1/2} = i[e_j, e_{j+1}]$,
3. nearest-neighbour current symbols whose low-energy slope does not match the selected $c^2$.

The onsite-independence checks also catch an accidental insertion of the mass or onsite term into the nearest-neighbour current.

### 34.5 Open Obligations

The suite deliberately leaves the following undone:

1. finite open-chain wave packets and comparison with $\mathrm{d}\Gamma(k)$,
2. periodic first moments or periodic coordinate generators,
3. higher-dimensional current-density formulas and incidence-sign tests,
4. next-nearest-neighbour or doubler current-density witnesses,
5. momentum-density $T_{10}$ and stress-current $T_{11}$ continuity tests,
6. scaling maps, state choices, domains, and continuum convergence.

The doubler evidence remains the CA-28 coefficient/minimum-data rejection. It should not be described as a current-density failure until a current formula for that non-nearest-neighbour symbol is derived and tested.

# 35 Qubit Nearest-Neighbour Symmetry Quest

## 35.1 Status, Sources, and Dependencies

**Status: programme shard with checked seed.** This shard scopes the next lattice-symmetry quest for qubit systems. CA-35 supplies the first checked finite algebraic obstruction. CA-36 and CA-37 state proposal-level diagnostics for $2+1$-dimensional Poincare tests and $1+1$-dimensional Witt/Virasoro tests.

Dependencies: CA-10–CA-17, CONVENTIONS.md (h), (k), (m), and the review orchestration in `reviews/2026-05-31_qubit_nn_symmetry/`.

## 35.2 Input Tier

The first input class is a translation-invariant qubit nearest-neighbour density. In the convention of CONVENTIONS.md (m),

$$h = \sum_{\alpha,\beta=0}^{3} h_{\alpha\beta}\, \sigma^{\alpha} \otimes \sigma^{\beta}, \qquad h_{\alpha\beta} \in \mathbb{R},$$

with $\sigma^0 = I$ and $\sigma^1, \sigma^2, \sigma^3 = X, Y, Z$. On a one-dimensional open chain the density $h_j$ acts on sites $(j, j+1)$, so

$$H = \sum_j h_j.$$

In two spatial dimensions the same coefficient format may be used separately on horizontal and vertical edges, but the edge orientation, midpoint, boundary, and bulk-projection conventions are not fixed by this shard.

## 35.3 Question

The question is not whether a finite qubit lattice has exact Poincare or Virasoro symmetry. CA-15 already forbids that shortcut. The question is whether the finite algebra forced by the candidate generator ansatz gives cheap rejection tests:

$$\begin{gathered}\text{Pauli coefficient data} \longrightarrow \text{finite commutator residuals} \\ \longrightarrow \text{rule out impossible symmetry candidates early.}\end{gathered}$$

The first such test is the adjacent-density current

$$J(h) := i[h_{12}, h_{23}],$$

because CA-12 shows that the first energy moment

$$K = \sum_j x_j h_j$$

obeys $i[H, K]$ equal to the sum of adjacent commutators in the translation-invariant bulk. If this current vanishes, the first-moment boost ansatz produces zero spatial momentum.

## 35.4 Investigation Plan

**One-dimensional Poincare filters.** Compute $J(h)$ directly from $h_{\alpha\beta}$. Reject the density as a nontrivial Poincare candidate for the CA-12 first-moment route if the bulk current vanishes while the energy density is non-scalar. Then test stronger finite residuals such as $i[P, K] - H$, but only after boundary, scaling, and normalization policies are named.

**Two-dimensional Poincare filters.** Use horizontal and vertical edge densities $h^x, h^y$. Define ramp moments from edge midpoints and compute $P_x = i[H, K_x]$, $P_y = i[H, K_y]$ on open patches. Proposed finite residuals include current nontriviality, $i[H, P_a]$, $i[P_x, P_y]$, isotropy under $x \leftrightarrow y$, and later rotation closure after a local momentum-density split is fixed.

**Witt/Virasoro filters.** On a periodic one-dimensional chain, form Fourier modes of the density and of the adjacent current. Koo–Saleur supplies the sourced pattern: Hamiltonian-density modes plus commutator corrections. The first rejection test is again current collapse; central terms and finite-size non-closure are not early failure tests.

## 35.5 Immediate Rule-Outs

The first checked obstruction is deliberately modest. Under the symmetric nearest-neighbour split of a genuine on-site qubit Hamiltonian, $J(h) = 0$. Thus a fully local single-site Hamiltonian cannot provide nonzero bulk momentum through the CA-12 first-moment boost ansatz. The same obstruction also catches commuting classical nearest-neighbour densities, such as pure diagonal $ZZ$ terms without noncommuting fields.

Nonzero $J(h)$ is not sufficient. It is only the first algebraic gate. A density that passes still needs the CA-15 evidence payload: local support policy, state sequence, renormalisation maps, convergence topology, low-energy sector or core, and observable covariance.

# 36 One-Dimensional Qubit Boost-Current Obstructions

## 36.1 Status, Sources, and Dependencies

**Status: checked local derivation.** This shard turns the CA-12 nearest-neighbour identity into explicit polynomial conditions on the two-qubit Pauli coefficient matrix. The obstruction is checked in Julia.

Dependencies: CA-12, CA-34, CONVENTIONS.md (h), (k), (m).

## 36.2 Adjacent Current in Pauli Coefficients

Let

$$h_j = \sum_{\alpha,\beta=0}^{3} h_{\alpha\beta}\, \sigma_j^{\alpha} \sigma_{j+1}^{\beta}.$$

The only noncommuting overlap between $h_j$ and $h_{j+1}$ is at site $j+1$. With

$$[\sigma^b, \sigma^c] = 2i \sum_{e=1}^{3} \epsilon_{bce} \sigma^e \qquad (b, c \in \{1,2,3\}),$$

one obtains

$$i[h_j, h_{j+1}] = -2 \sum_{\alpha,\delta=0}^{3} \sum_{e=1}^{3} \left( \sum_{b,c=1}^{3} h_{\alpha b} h_{c\delta} \epsilon_{bce} \right) \sigma_j^{\alpha} \sigma_{j+1}^{e} \sigma_{j+2}^{\delta}.$$

Equivalently, define right-leg and left-leg vectors

$$R_\alpha = (h_{\alpha 1}, h_{\alpha 2}, h_{\alpha 3}), \qquad L_\delta = (h_{1\delta}, h_{2\delta}, h_{3\delta}).$$

Then the current coefficients are the cross products

$$i[h_j, h_{j+1}] = -2 \sum_{\alpha,\delta=0}^{3} \sum_{e=1}^{3} (R_\alpha \times L_\delta)_e \sigma_j^{\alpha} \sigma_{j+1}^{e} \sigma_{j+2}^{\delta}.$$

Thus the simple current-collapse condition is

$$R_\alpha \times L_\delta = 0 \qquad \text{for every } \alpha, \delta.$$

When this holds, every adjacent-current contribution to the CA-12 bulk momentum candidate vanishes.

## 36.3 Fully On-Site Terms

For a genuine translation-invariant on-site density

$$A = \sum_{\alpha=0}^{3} a_\alpha \sigma^\alpha,$$

CONVENTIONS.md (m) embeds it into the two-site bond tier by the symmetric split

$$h_{\text{onsite}} = \frac{1}{2}(A \otimes I + I \otimes A).$$

Then the right-leg and left-leg vectors are all parallel in the only places where they overlap, so

$$i[(h_{\text{onsite}})_{12}, (h_{\text{onsite}})_{23}] = 0.$$

The Julia test checks this directly for a generic real linear combination of $I, X, Y, Z$.

Consequently the first-moment boost ansatz gives

$$i[H, K] = 0$$

in the translation-invariant bulk. If the target is a nontrivial Poincare representation with $i[H, K] = P$ and $i[P, K] = H$, this first-moment route cannot realise it unless the energy itself is trivial after the relevant compression or subtraction.

### 36.4 Other Immediate Rejections

The same current-collapse test catches commuting nearest-neighbour density families. A pure diagonal classical interaction such as $ZZ$ has

$$[h_j, h_{j+1}] = 0$$

because the overlapping Pauli factor is the same diagonal matrix. Such a model may still be a well-defined lattice Hamiltonian, but it fails the first nontrivial bulk-symmetry diagnostic: the boost-current candidate supplies no spatial momentum.

By contrast, the tested transverse-Ising-style density with a $ZZ$ bond and a symmetric $X$-field split has nonzero adjacent current. This does not prove Poincare invariance; it only passes the first obstruction.

### 36.5 The Next Residual

The next necessary Poincare relation is translation conservation:

$$i[H, P] = 0.$$

For the local candidate $P = \sum_j C_j$, $C_j = i[h_j, h_{j+1}]$, the minimal bulk density after the Jacobi cancellation of outer-overlap terms is

$$A_j = i[h_j + h_{j+1}, C_j].$$

The full infinite-chain equation is not necessarily $A_j = 0$, but

$$A_j = u_j - u_{j+1}$$

for a finite density $u$. CA-39 writes this and the next boost relation $i[P, K] = v^2 H$ as explicit Pauli-coefficient equations.

This residual is useful for detecting density-split artifacts. An asymmetric encoding of an on-site Hamiltonian,

$$h = U \otimes I + I \otimes V,$$

can create the fake current

$$i[h_{12}, h_{23}] = I \otimes i[V, U] \otimes I.$$

But the checked sentinel with $U = X$, $V = Z$ has nonzero

$$i[h_{12} + h_{23}, i[h_{12}, h_{23}]],$$

so it fails translation conservation even though the adjacent current is nonzero. The CA-39 boost equation then tests the remaining relation after the allowed density-divergence and scalar-energy subtraction have been fixed.

### 36.6 Boundary

The obstruction is conditional on the density convention. An asymmetric encoding of an on-site Hamiltonian as $A \otimes I + I \otimes B$ can create spurious adjacent commutators when $A$ and $B$ fail to commute. Such currents are density-split artifacts unless a separate convention justifies the split, and the conservation residual above is the first backstop against them. Passing the obstruction is also far weaker than CA-15 acceptance: boundary residuals, scaling maps, low-energy sectors, and observable covariance remain unproved.

# 37 Two-Dimensional Qubit Plaquette Symmetry Diagnostics

## 37.1 Status, Sources, and Dependencies

**Status: proposal diagnostics.** This shard gives a finite algebraic test plan for $2+1$-dimensional qubit lattices. No two-dimensional current or rotation convention is fixed yet, so every condition below is a proposed compiler filter.

Dependencies: CA-13, CA-15, CA-16, CA-32, CONVENTIONS.md (h), (k), (m).

## 37.2 Edge-Density Input

On an open square patch, use oriented horizontal and vertical edges:

$$h^a_{r,r+\hat{a}} = \sum_{\alpha,\beta=0}^{3} c^a_{\alpha\beta}\, \sigma^\alpha_r \sigma^\beta_{r+\hat{a}}, \qquad a \in \{x, y\}.$$

Let $m_e$ be the midpoint of edge $e$, and write

$$H = \sum_e h_e, \qquad K_a = \sum_e m^a_e h_e.$$

The ramp-current candidate is

$$P_a := i[H, K_a] = \sum_{\{e,f\}} (m^a_f - m^a_e) i[h_e, h_f],$$

where only overlapping edge pairs contribute. This is the direct higher-dimensional analogue of CA-13's scalar-ramp formula.

## 37.3 Patch Diagnostics

The smallest useful checks are local:

1. three-site lines for straight $x$ and $y$ currents,
2. $2 \times 2$ plaquettes for corner commutators and $C_4$ covariance,
3. $3 \times 3$ open patches for $P_x, P_y$ and $i[H, P_a]$,
4. $4 \times 4$ or larger patches for bulk-projected $i[P_x, P_y]$.

Boundary-supported residuals should be projected away or recorded separately. The diagnostic object is the interior Pauli-string residual, not raw equality on the whole open patch.

## 37.4 Proposed Rejection Filters

**Current nontriviality.** Both $P_x$ and $P_y$ should be nonzero in the bulk for a nontrivial $2+1$-dimensional Poincare candidate.

**Translation conservation.** The bulk residual $i[H, P_a]$ should vanish or be controlled in the intended finite algebra. A nonzero interior Pauli-string residual is an early rejection witness for exact finite bulk closure.

**Translation commutativity.** The bulk part of $i[P_x, P_y]$ should vanish or be a named boundary/irrelevant term before the model is promoted.

**Boost-translation bracket.** After normalisation, the residual

$$i[P_b, K_a] - \delta_{ab} H$$

is the direct $2+1$-dimensional Poincare diagnostic. This is strong and must be treated as a finite filter, not a theorem.

**Isotropy.** For scalar qubits with no internal spatial action, a microscopic $90^\circ$ rotation should relate horizontal and vertical edge data after endpoint reversal. Obvious decoupled stacks, such as $c^y = 0$, fail this filter.

### 37.5 Rotation Is Not Free

A rotation candidate needs a local momentum-density split

$$P_a = \sum_v p_a(v)$$

before one can write

$$J_{xy} = \sum_v (x_v p_y(v) - y_v p_x(v)).$$

CA-13 already marks this as extra data. A plausible first convention is to assign each edge-pair current to the shared vertex, but that convention must be written and tested before rotation closure is claimed. Only then should one check

$$i[H, J_{xy}], \qquad i[P_x, J_{xy}] - P_y, \qquad i[P_y, J_{xy}] + P_x.$$

### 37.6 Rule-Out Classes

The proposal immediately targets four classes:

1. fully on-site Hamiltonians: $P_x = P_y = 0$,
2. commuting classical edge densities: all ramp currents vanish,
3. decoupled stacks: one momentum component vanishes,
4. anisotropic edge data: fail the scalar-qubit $C_4$ filter.

None of these rejections is a continuum theorem. They are finite algebraic filters for the specific candidate-generator route.

# 38 Qubit Nearest-Neighbour Witt/Virasoro Diagnostics

## 38.1 Status, Sources, and Dependencies

**Status: proposal with local derivation targets.** This shard is a finite-scale diagnostic proposal, not a continuum Virasoro claim.

Dependencies: CA-12, CA-14–CA-16, CA-34–CA-35, CONVENTIONS.md (m). CFT normalisations in CONVENTIONS.md (d) remain unfixed, so no central-charge extraction convention is available here.

## 38.2 Fourier Density and Current Modes

Let a periodic qubit chain of length $N$ have

$$H = \sum_j h_j, \qquad h_j = \sum_{\alpha,\beta=0}^{3} h_{\alpha\beta}\sigma_j^\alpha \sigma_{j+1}^\beta.$$

For $q_n = 2\pi n/N$, define

$$H_n = \sum_j e^{iq_n j} h_j, \qquad J_n = \sum_j e^{iq_n j} j_j, \qquad j_j = i[h_j, h_{j+1}].$$

The nearest-neighbour commutator identity is

$$i[H_m, H_n] = (e^{iq_n} - e^{iq_m}) J_{m+n}.$$

This is the finite Pauli-chain analogue of the Koo–Saleur pattern in which Hamiltonian-density modes and commutator corrections approximate chiral stress-tensor modes.

## 38.3 First Failure Witness

CA-35 gives the coefficient condition

$$R_\alpha \times L_\delta = 0 \qquad \text{for all } \alpha, \delta,$$

which implies $j_j = 0$ and hence $J_n = 0$ for every Fourier mode. In that case the density modes cannot acquire the Koo–Saleur commutator correction needed to split left-moving and right-moving candidates. Fully on-site Hamiltonians under the symmetric split fail here, as do commuting classical densities.

## 38.4 Chiral-Rank Diagnostic

A later finite diagnostic may define tentative chiral candidates of the form

$$\ell_n = a_N{:}H_n{:} + \lambda_n J_n + \mu_N \delta_{n0} I, \qquad \bar{\ell}_{-n} = a_N{:}H_n{:} - \lambda_n J_n + \mu_N \delta_{n0} I.$$

Before testing brackets, one should check that $H_n$ and $J_n$ are linearly independent on the chosen low-energy sector $P_N$. A rank-collapse test such as

$$\det \mathrm{Gram}_{P_N}(H_n, J_n) \neq 0$$

is an early finite witness that the two proposed chiral directions have not collapsed to one.

## 38.5 Compressed Witt Residuals

The raw full Hilbert-space commutator is the wrong first test. Koo–Saleur already warns that finite-size quantities do not close as Virasoro generators and that commutators interact subtly with the scaling limit. The diagnostic should therefore choose a low-energy projection $P_N$, an intermediate projection $Q_N$, and test residuals such as

$$P_N\big(\ell_m Q_N \ell_n - \ell_n Q_N \ell_m - (m-n)\ell_{m+n}\big)P_N$$

for $m+n \neq 0$, together with the barred and cross-chiral versions.

The $m+n=0$ brackets should not be the first rejection test: central terms require a Virasoro normalization, vacuum or ground-state subtraction, and a scaling state sequence. These are not fixed for arbitrary qubit chains.

## 38.6 Boundary

This shard gives finite algebraic diagnostics only. Passing them does not produce a Virasoro representation. A continuum claim still needs the full CA-15 payload plus Koo–Saleur-style scaling representation data: state sequence, subtraction policy, mode normalization, low-energy sector, operator topology, and convergence theorem.

# 39 Qubit Local-Algebra Equation Framework

## 39.1 Status, Sources, and Dependencies

**Status: checked local framework.** This shard supplies the algebraic language used by CA-39–CA-44. It does not assert that a solution has continuum Poincare or Virasoro symmetry.

Dependencies: CA-11, CA-12, CA-15, CA-34–CA-35, CONVENTIONS.md (m), (n).

## 39.2 Pauli Word Algebra

Use zero-based Pauli labels $a \in \{0,1,2,3\}$, with $\sigma^0 = I$. For a finite interval, a Pauli word is

$$P_{\alpha_0 \dots \alpha_{\ell-1}} = \sigma^{\alpha_0} \otimes \cdots \otimes \sigma^{\alpha_{\ell-1}}.$$

The local product is encoded by constants $M_{ab}^c$:

$$\sigma^a \sigma^b = \sum_{c=0}^{3} M_{ab}^c \sigma^c,$$

where

$$M_{0b}^c = \delta_{bc}, \qquad M_{a0}^c = \delta_{ac},$$

and, for $a, b \geq 1$,

$$M_{ab}^0 = \delta_{ab}, \qquad M_{ab}^c = i\epsilon_{abc} \quad (c \geq 1).$$

Thus for two $\ell$-site coefficient tensors $A, B$,

$$(AB)_\gamma = \sum_{\alpha,\beta} A_\alpha B_\beta \prod_{r=0}^{\ell-1} M_{\alpha_r \beta_r}^{\gamma_r}.$$

The commutator coefficient is therefore the finite expression

$$(i[A,B])_\gamma = i \sum_{\alpha,\beta} A_\alpha B_\beta \left( \prod_r M_{\alpha_r \beta_r}^{\gamma_r} - \prod_r M_{\beta_r \alpha_r}^{\gamma_r} \right).$$

When two local densities have different supports, first embed both tensors into their common finite window by padding with Pauli label 0, then apply this formula.

## 39.3 Bulk Equality Modulo Divergence

Let $\tau$ be the unit shift of the infinite chain. A finite density $R$ represents a zero bulk formal sum if it is a coboundary:

$$R = u - \tau(u).$$

If $u$ has $n$ Pauli indices, the coefficient form fixed in CONVENTIONS.md (n) is

$$(Du)_{a_0 \dots a_n} = u_{a_0 \dots a_{n-1}} \delta_{a_n 0} - \delta_{a_0 0} u_{a_1 \dots a_n}. \tag{38.1}$$

Consequently a local residual $R(h)$ passes the bulk equation precisely when there exists a finite tensor $u$ such that

$$R_{a_0 \dots a_n}(h) - (Du)_{a_0 \dots a_n} = 0 \qquad \text{for every Pauli word } a_0 \dots a_n. \tag{38.2}$$

The scalar coefficient of $u$ lies in the kernel of $D$, so it is a gauge variable. A coefficientwise condition $R(h) = 0$ is allowed as a stronger filter, but it is not the default infinite-chain equation.

### 39.4 What "Necessary" Means Here

Every equation in the next shards is necessary only for the named generator route: the fixed Pauli density, the fixed first-moment boost ansatz, and the fixed coboundary quotient. Solving these equations is not sufficient for continuum symmetry. Failing one of them is still useful: it excludes the Hamiltonian from this route before any low-energy sector, scaling map, or state-space analysis is attempted.

# 40 One-Dimensional Qubit Poincare Necessary Equations

## 40.1 Status, Sources, and Dependencies

**Status: local derivation, partly checked.** The current, coboundary, and residual tensors are implemented in Julia. The equations are necessary for the fixed CA-12 first-moment route, not sufficient for continuum Poincare symmetry.

Dependencies: CA-11, CA-12, CA-35, CA-38, CONVENTIONS.md (k), (m), (n).

## 40.2 Input and Current

Let

$$h_j = \sum_{\alpha,\beta=0}^{3} h_{\alpha\beta}\sigma_j^{\alpha}\sigma_{j+1}^{\beta}, \qquad h_{\alpha\beta} \in \mathbb{R}.$$

The first-moment route fixes

$$H = \sum_j h_j, \qquad K = \sum_j j h_j, \qquad P = i[H,K] = \sum_j p_j, \qquad p_j = i[h_j, h_{j+1}].$$

The current coefficients are

$$p_{aed} = -2 \sum_{b,c=1}^{3} h_{ab} h_{cd} \epsilon_{bce}, \qquad e \in \{1,2,3\}, \tag{39.1}$$

with $p_{a0d} = 0$. Equivalently the right-leg vector $(h_{a1}, h_{a2}, h_{a3})$ must cross the left-leg vector $(h_{1d}, h_{2d}, h_{3d})$. If

$$p_{aed} = 0 \quad \text{for all } a, e, d, \tag{39.2}$$

then the candidate bulk momentum is $P = 0$. For a nontrivial speed $v^2 \neq 0$, the later equation $i[P,K] = v^2 H$ can then hold only if the Hamiltonian density is scalar-trivial in the coboundary quotient. The symmetric fully on-site split in CONVENTIONS.md (m) is checked to have $p = 0$; hence fully local non-scalar Hamiltonians are ruled out for this route.

## 40.3 Translation Conservation

Let

$$A_j = i[h_j + h_{j+1}, p_j].$$

The outer-overlap terms in $i[H,P]$ cancel in the formal sum by the Jacobi identity, so translation conservation is

$$A = Du \tag{39.3}$$

for some two-site tensor $u_{rs}$. In Pauli coefficients,

$$\begin{aligned} A_{rst} = i \sum_{\alpha,\beta,a,b} h_{\alpha\beta} p_{abt} \left( M^r_{\alpha a} M^s_{\beta b} - M^r_{a\alpha} M^s_{b\beta} \right) \\ + i \sum_{\alpha,\beta,b,c} h_{\alpha\beta} p_{rbc} \left( M^s_{\alpha b} M^t_{\beta c} - M^s_{b\alpha} M^t_{c\beta} \right). \end{aligned}$$

Thus the explicit degree-three equations are

$$A_{rst} - u_{rs}\delta_{t0} + \delta_{r0} u_{st} = 0 \qquad (r,s,t = 0,\ldots,3). \tag{39.4}$$

## 40.4 Boost-Translation Relation

Assume (39.3) has been witnessed by a chosen $u$. The remaining bulk representative of $i[P,K]$ is

$$B_j - u_j, \qquad B_j = i[p_j, h_{j+1}] + 2i[p_j, h_{j+2}].$$

The four-site coefficients of $B$ are

$$\begin{aligned} B_{rstu} = i\delta_{u0} \sum_{b,c,\alpha,\beta} p_{rbc} h_{\alpha\beta} \left( M^s_{b\alpha} M^t_{c\beta} - M^s_{\alpha b} M^t_{\beta c} \right) \\ + 2i \sum_{c,\alpha} p_{rsc} h_{\alpha u} \left( M^t_{c\alpha} - M^t_{\alpha c} \right). \end{aligned}$$

Allow an explicit scalar energy-origin subtraction

$$\bar{h}_{rs} = h_{rs} - e\,\delta_{r0}\delta_{s0}.$$

The necessary speed-$v$ boost equation $i[P,K] = v^2 H$ is that there exists a three-site tensor $w$ such that

$$\begin{aligned} 0 = B_{rstu} - u_{rs}\delta_{t0}\delta_{u0} - v^2 \bar{h}_{rs}\delta_{t0}\delta_{u0} \\ - w_{rst}\delta_{u0} + \delta_{r0} w_{stu} \qquad (r,s,t,u = 0,\ldots,3). \end{aligned} \tag{39.5}$$

Equations (39.1), (39.4), and (39.5), together with real variables $u, w, e, v^2$, are the advertised explicit finite polynomial conditions on $h_{\alpha\beta}$ for the one-dimensional first-moment Poincare route.

## 40.5 Boundary

The scalar $h_{00}$ cannot be detected by commutators; it must be handled by the explicit $e$ variable or by a state-dependent normal ordering. The coboundary witnesses are also not unique. These are convention choices, not loopholes: changing the density by a divergence changes $K = \sum_j j h_j$, so it changes the equations.

# 41 Computer Algebra for One-Dimensional Qubit Residuals

## 41.1 Status, Sources, and Dependencies

**Status: checked.** This shard documents the finite algebra engine behind CA-35 and CA-39.
Dependencies: CA-35, CA-38, CA-39, CONVENTIONS.md (m), (n).

## 41.2 Operator and Coefficient Roundtrip

The file `src/QubitPauliLattice.jl` implements

$$\sum_{\alpha_0,\ldots,\alpha_{n-1}=0}^{3} c_{\alpha_0\ldots\alpha_{n-1}} \sigma^{\alpha_0} \otimes \cdots \otimes \sigma^{\alpha_{n-1}}$$

for $n = 1, 2, 3$ and for general $n$. The inverse map uses

$$c_\alpha = 2^{-n} \operatorname{Tr}(P_\alpha A).$$

The tests assert coefficient roundtrips for two-site densities and for the five-site residual representatives. The code rejects arrays whose axes are not all length 4, and rejects non-self-adjoint matrix inputs.

## 41.3 Checked Residual Helpers

The current helper returns

$$p = i[h_{12}, h_{23}]$$

and the direct coefficient helper returns exactly (39.1). The conservation helper

$$A = i[h_{12} + h_{23}, p]$$

is the minimal three-site density for $i[H, P]$ after the Jacobi cancellation of outer-overlap terms.

The file `src/QubitPauliResiduals.jl` adds

$$(Du)_{a_0\ldots a_n} = u_{a_0\ldots a_{n-1}}\delta_{a_n 0} - \delta_{a_0 0} u_{a_1\ldots a_n}$$

and

$$B = i[p_{123}, h_{23}] + 2i[p_{123}, h_{34}].$$

The test suite checks that the coboundary coefficients reconstruct the operator identity

$$Du = u_{12} - u_{23}$$

on a three-site window, including the fact that the scalar $u_{00}I$ is in the kernel.

## 41.4 Sentinels

The checked examples are deliberately small:

- A generic real one-site density embedded by the symmetric split has $p = 0$, $A = 0$, and $B = 0$, while the embedded Hamiltonian density itself is nonzero. This is the concrete fully-local exclusion witness for CA-39.

- A pure classical $ZZ$ density also has $p = 0$.
- A transverse-Ising-style density has nonzero $p$ and nonzero boost residual density $B$. This only means it survives the first obstruction.
- An asymmetric fake on-site split creates a nonzero current, but the conservation sentinel is larger than the current and catches the density-split artifact.

### 41.5 Current Gap

The code does not yet solve the polynomial system (39.1), (39.4), (39.5), and does not build moment matrices. Its role is narrower: it makes every displayed coefficient tensor executable and gives mutation-friendly invariants before an SDP exporter is added.

# 42 Two-Dimensional Qubit Poincare Equation Schema

## 42.1 Status, Sources, and Dependencies

**Status: proposal-level local schema.** This is a finite algebraic target for a future checker. It is not yet a canonical stress tensor or a continuum $2+1$-dimensional Poincare theorem.

Dependencies: CA-13, CA-36, CA-38, CONVENTIONS.md (o).

## 42.2 Edge Densities and Vertex Currents

On $\mathbb{Z}^2$, write positively oriented edge densities

$$h_r^a = \sum_{\alpha,\beta=0}^{3} c_{\alpha\beta}^a \, \sigma_r^\alpha \sigma_{r+\hat{a}}^\beta, \qquad a \in \{x, y\}.$$

At a vertex $v$, order incident edges as $W, S, E, N$, with midpoint offsets

$$m_W = (-1/2, 0), \quad m_S = (0, -1/2), \quad m_E = (1/2, 0), \quad m_N = (0, 1/2).$$

Use shared-leg coefficient tensors

$$q_{\lambda\mu}^W = c_{\lambda\mu}^x, \quad q_{\lambda\mu}^E = c_{\mu\lambda}^x, \quad q_{\lambda\mu}^S = c_{\lambda\mu}^y, \quad q_{\lambda\mu}^N = c_{\mu\lambda}^y.$$

For ordered incident pairs $A < B$ in $W < S < E < N$, the local commutator coefficient on outer leg $A$, center $v$, outer leg $B$ is

$$J_{\lambda\kappa\rho}^{AB} = -2 \sum_{\mu,\nu=1}^{3} q_{\lambda\mu}^A q_{\rho\nu}^B \epsilon_{\mu\nu\kappa}. \tag{41.1}$$

The ramp-generated momentum densities are

$$p_a(v) = \sum_{A<B} (m_B^a - m_A^a) J^{AB}(v). \tag{41.2}$$

For $AB = WS, WE, WN, SE, SN, EN$, the $x$-weights are

$$1/2,\ 1,\ 1/2,\ 1/2,\ 0,\ -1/2,$$

and the $y$-weights are

$$-1/2,\ 0,\ 1/2,\ 1/2,\ 1,\ 1/2.$$

## 42.3 Divergence-Relaxed Residuals

Every residual below is a finite Pauli tensor $R$ on a named patch. The strong filter is $R = 0$. The default bulk filter is the existence of finite-support tensors $U_x, U_y$ with

$$R = (1 - \tau_{-\hat{x}})U_x + (1 - \tau_{-\hat{y}})U_y. \tag{41.3}$$

Coefficientwise, this is an explicit linear equation in the auxiliary divergence tensors and a polynomial equation in $c^x, c^y$, using the Pauli product constants of CA-38.

## 42.4 Necessary Bulk Equations

Momentum conservation is

$$C_a := \sum_{e:\operatorname{supp} h_e \cap \operatorname{supp} p_a(0) \neq \varnothing} i[h_e, p_a(0)] = \nabla \cdot U^{(C_a)}, \qquad a = x, y. \tag{41.4}$$

The commuting-translation equation is

$$T_{xy} := \sum_{r:\operatorname{supp} p_x(0) \cap \operatorname{supp} p_y(r) \neq \varnothing} i[p_x(0), p_y(r)] = \nabla \cdot U^{(T)}. \tag{41.5}$$

Let $e_0 = h_0^x + h_0^y$ denote the chosen local energy cell. With $m_e^a$ the $a$-coordinate of the edge midpoint,

$$B_{ab} := \sum_{e:\operatorname{supp} h_e \cap \operatorname{supp} p_b(0) \neq \varnothing} m_e^a \, i[p_b(0), h_e] - \delta_{ab} v_a^2 e_0 = \nabla \cdot U^{(B_{ab})}. \tag{41.6}$$

This is the $i[P_b, K_a] = \delta_{ab} v_a^2 H$ equation after the coordinate weighted conservation term is removed by (41.4).

If one defines the rotation candidate

$$J = \sum_v (v_x p_y(v) - v_y p_x(v)),$$

then the translation-rotation residual is

$$R_c = \sum_r \left( r_x \, i[p_c(0), p_y(r)] - r_y \, i[p_c(0), p_x(r)] \right) - (\delta_{cx} p_y(0) - \delta_{cy} p_x(0)) = \nabla \cdot U^{(R_c)}. \tag{41.7}$$

This last equation is quartic in $c^x, c^y$ before the quadratic subtraction. It depends on the chosen momentum-density split, so it is less canonical than the boost equations.

## 42.5 Implementation Boundary

CA-41 supplies the finite equations a checker should implement. The next implementation step is to enumerate the patch supports in (41.4)–(41.7), embed every edge tensor into the common patch, and emit Pauli coefficients plus the divergence auxiliaries. Until that exists, the $2+1$-dimensional claims remain proposal-level.

# 43 Qubit Witt/Virasoro Necessary Equations

## 43.1 Status, Sources, and Dependencies

**Status: exact finite algebra plus proposal residuals.** The mode identities are finite local algebra. Their use as Virasoro diagnostics is a necessary filter only.

Dependencies: CA-14, CA-35, CA-37–CA-39, CONVENTIONS.md (d), (m), (n).

## 43.2 Exact Density-Mode Algebra

On a periodic chain of length $N$, set $z_n = \exp(2\pi i n/N)$ and

$$H_n = \sum_j z_n^j h_j, \qquad C_j = i[h_j, h_{j+1}], \qquad J_n = \sum_j z_n^j C_j.$$

Then $H_n^* = H_{-n}$, $J_n^* = J_{-n}$, and the nearest-neighbour identity is

$$[H_m, H_n] = -i(z_n - z_m) J_{m+n}. \tag{42.1}$$

This is the exact lattice antecedent of the Koo-Saleur $H(f), P(f)$ commutator. In Pauli coefficients $C_j$ is the tensor $p$ from (39.1).

Define higher-current densities

$$D_{r,j} = i[h_{j+r}, C_j] \quad (r = 0, 1, 2), \qquad F_j^{(r)} = i[C_j, C_{j+r}] \quad (r = 1, 2).$$

Using the Jacobi cancellation of the $r = -1$ term,

$$i[H_m, J_n] = \sum_j z_{m+n}^j \left( D_{0,j} + z_m D_{1,j} + (z_m^2 - z_n) D_{2,j} \right), \tag{42.2}$$

and

$$i[J_m, J_n] = \sum_j z_{m+n}^j \left( (z_n - z_m) F_j^{(1)} + (z_n^2 - z_m^2) F_j^{(2)} \right). \tag{42.3}$$

The $D$-coefficients are degree three polynomials in $h_{\alpha\beta}$; the $F$-coefficients are degree four.

## 43.3 Chiral Residuals

A Koo-Saleur-style chiral ansatz has the form

$$\ell_n = A_n {:}H_n{:}_\omega + B_n {:}J_n{:}_\omega + \mu \delta_{n0} I,$$
$$\bar{\ell}_n = A_n {:}H_n{:}_\omega - B_n {:}J_n{:}_\omega + \mu \delta_{n0} I,$$

with $A_{-n} = \overline{A_n}$ and $B_{-n} = \overline{B_n}$. The necessary Virasoro residuals for a chosen finite mode set are

$$R_{mn}^{++} = [\ell_m, \ell_n] - (m - n)\ell_{m+n} - \frac{c}{12} m(m^2 - 1)\delta_{m+n,0} I, \tag{42.4}$$

$$R_{mn}^{--} = [\bar{\ell}_m, \bar{\ell}_n] - (m - n)\bar{\ell}_{m+n} - \frac{c}{12} m(m^2 - 1)\delta_{m+n,0} I, \tag{42.5}$$

and

$$R_{mn}^{+-} = [\ell_m, \bar{\ell}_n]. \tag{42.6}$$

For example,

$$[\ell_m, \ell_n] = -iA_m A_n (z_n - z_m) J_{m+n} - iA_m B_n \mathcal{D}_{mn} + iB_m A_n \mathcal{D}_{nm} - iB_m B_n \mathcal{F}_{mn},$$

where $\mathcal{D}_{mn} = i[H_m, J_n]$ and $\mathcal{F}_{mn} = i[J_m, J_n]$ are given by (42.2)–(42.3).

### 43.4 Necessary Filter

At finite size, the Koo-Saleur source explicitly warns that the lattice quantities need not close. Therefore the safe condition is not "finite Virasoro symmetry". The safe condition is: for the chosen ansatz and mode set, every Pauli coefficient of $R^{++}, R^{--}, R^{+-}$ outside the named span

$$\mathrm{span}\{I, H_{m+n}, J_{m+n}\}$$

must either vanish, be a registered finite-size correction, or be annihilated in the chosen low-energy/state restriction. CA-43–CA-44 turn the last option into moment constraints.

# 44 Vacuum Moment Constraints for Qubit Residuals

## 44.1 Status, Sources, and Dependencies

**Status: proposal-level SDP interface backed by sourced state/GNS architecture.** The moment equations are finite algebra. The existence of a continuum vacuum or positive-energy representation is not claimed.

Dependencies: CA-15, CA-38–CA-42, CONVENTIONS.md (n).

## 44.2 Finite Moment Data

Let $P_s$ be a finite Pauli word on $\mathbb{Z}$, padded by identities outside its support. A candidate translation-invariant global state $\omega$ has moments

$$y_s = \omega(P_s), \qquad y_\emptyset = 1, \qquad y_{\tau^n s} = y_s. \tag{43.1}$$

For a finite word set $\mathcal{W}$, positivity is the moment-matrix constraint

$$M_{\mathcal{W}}(y)_{u,v} = \omega(P_u^* P_v) = \sum_s \mu_{u,v}^s y_s \succeq 0, \tag{43.2}$$

where the constants $\mu_{u,v}^s$ are computed from the Pauli product table of CA-38. Hermiticity gives

$$M_{\mathcal{W}}(y)_{v,u} = \overline{M_{\mathcal{W}}(y)_{u,v}}.$$

The SDP may be represented over real variables by splitting real and imaginary parts.

## 44.3 Zero Expectations

The infinite sums $H, P, K$ are not finite observables. The finite data are their local densities and residuals. After choosing the scalar subtraction $\bar{h} = h - eI$, a vacuum-normalized test may impose

$$\omega(\bar{h}_j) = 0, \qquad \omega(p_j) = 0, \tag{43.3}$$

and, for the residuals of CA-39,

$$\omega(A_j - Du_j) = 0, \qquad \omega(B_j - u_j - v^2 \bar{h}_j - Dw_j) = 0. \tag{43.4}$$

These are only the level-zero moment consequences. They are usually too weak to exclude much.

## 44.4 GNS Relation Constraints

The stronger finite restriction says that a residual vanishes on the vacuum sector as a relation in the GNS representation. For each chosen residual density $R(h)$ and probe words $X, Y$,

$$\omega(X^* R(h) Y) = 0. \tag{43.5}$$

For fixed $h$, fixed $v^2, e, u, w$, and fixed residual coefficients, (43.5) is linear in the moments $y_s$. The same rule applies to the finite Witt/Virasoro residual densities $r_{mn}^{++}, r_{mn}^{--}, r_{mn}^{+-}$ from CA-42:

$$\omega(X^* r_{mn}^{\pm\pm}(h) Y) = 0, \qquad \omega(X^* r_{mn}^{+-}(h) Y) = 0. \tag{43.6}$$

These equations are the promised finite data extracted from the demand that a global state $\omega$ make the generators and relations vanish in expectation on a finite probe space.

### 44.5 Highest-Weight Caveat

If a chiral vacuum sector is named, one may also impose finite analogues of

$$\ell_n\Omega = 0, \quad \bar{\ell}_n\Omega = 0 \qquad (n > 0),$$

for example

$$\omega(\ell_{-n}\ell_n) = 0. \tag{43.7}$$

But central charge, normal ordering, and finite-volume mode normalisation are still convention-sensitive. Until CONVENTIONS.md (d) is fixed for this block, such constraints are optional diagnostics rather than canonical Virasoro vacuum equations.

# 45 SDP Exclusion Hierarchy for Qubit Hamiltonians

## 45.1 Status, Sources, and Dependencies

**Status: rigorous proposal.** The hierarchy is a finite-state consequence of the sourced state/GNS framework and the local residual derivations. No solver is implemented yet.

Dependencies: CA-38–CA-43, CONVENTIONS.md (n).

## 45.2 Level Data

Fix a Hamiltonian coefficient matrix $h_{\alpha\beta}$. Also fix, or scan externally, the scalar subtraction $e$, speed $v^2$, and any coboundary witnesses $u, w$ used in CA-39. This is important: if those witnesses are unknown variables multiplied by unknown moments, the problem is no longer an SDP without an additional lifting relaxation.

At level $N$, choose finite sets

$$\mathcal{W}_N = \{\text{Pauli words supported in } [-N, N]\},$$

$$\mathcal{A}_N = \{\text{probe words whose residual products fit the window}\},$$

and residual densities

$$\mathcal{R}_N(h) = \{A - Du,\ B - u - v^2\bar{h} - Dw\}$$

plus any selected finite Witt/Virasoro residuals from CA-42.

## 45.3 SDP Constraints

The variables are finitely many moments $y_s$. The constraints are

$$y_\emptyset = 1, \qquad y_{\tau s} = y_s, \tag{44.1}$$

$$M_{\mathcal{W}_N}(y) \succeq 0, \tag{44.2}$$

and optional stationarity constraints for local probes:

$$\omega(\delta_H(X)) = 0, \qquad \omega(\delta_P(X)) = 0, \qquad X \in \mathcal{A}_N. \tag{44.3}$$

The generator-relation constraints are

$$\omega(X^* R Y) = 0, \qquad R \in \mathcal{R}_N(h), \quad X, Y \in \mathcal{W}_N, \tag{44.4}$$

whenever the Pauli product $X^*RY$ fits inside the moment support. Each entry in (44.1)–(44.4) is affine in $y$ for fixed residual coefficients, so this is a semidefinite feasibility problem.

## 45.4 Exclusion Statement

If a translation-invariant global state $\omega$ exists whose GNS representation carries the chosen local residual relations, then every finite restriction of $\omega$ satisfies every level $N$. Therefore:

$$\text{SDP level } N \text{ infeasible} \quad \Longrightarrow \quad h_{\alpha\beta} \text{ is excluded for this generator route.} \tag{44.5}$$

The hierarchy is monotone: increasing $N$ adds moment variables and relation tests, and cannot turn a previously infeasible level into evidence of symmetry.

### 45.5 Non-Implications

Feasibility at all checked levels does not prove that a global state exists, does not prove uniqueness or clustering, does not prove positive energy, and does not prove a continuum Poincare or Virasoro representation. It only means that the tested finite restrictions have not yet found a contradiction.

When $h_{\alpha\beta}$ is also a variable, the residual coefficients are polynomial in $h$. Then the feasibility problem becomes a polynomial moment problem over both Hamiltonian and state variables. That is a separate relaxation; the first implementation should instead take $h$ as input and return either an SDP infeasibility certificate or "not excluded at this level".

# 46 Roadmap for Qubit Symmetry Exclusion

## 46.1 Status, Sources, and Dependencies

**Status: implementation plan.** This closes the current qubit necessary-equation expansion and names the next build steps.

Dependencies: CA-34–CA-44, CONVENTIONS.md (m)–(o).

## 46.2 What We Have Now

The one-dimensional first-moment route now has explicit necessary equations on the Pauli coefficient matrix $h_{\alpha\beta}$:

$$p_{aed} = -2 \sum_{b,c=1}^{3} h_{ab} h_{cd} \epsilon_{bce},$$

$$A(h) = Du, \qquad B(h) - u - v^2(h - eI) = Dw.$$

These equations already rule out fully local symmetric on-site Hamiltonians and commuting classical densities for any nontrivial Poincare route of this form, because their generated momentum current is zero.

The two-dimensional shard gives a square-lattice edge schema, not yet code. The Witt/Virasoro shard gives exact finite mode commutators and residuals, but keeps Koo-Saleur's finite-nonclosure warning active. The vacuum/SDP shards turn "there exists a global state $\omega$" into finite positive moment data and linear relation constraints.

## 46.3 Implementation Steps

1. Add symbolic coefficient builders for $A(h)$, $B(h)$, and coboundary equations, returning sparse polynomial tensors rather than dense matrices.

2. Add a solver-facing algebraic filter: for a fixed $h$, check current collapse; optionally solve the linear coboundary witnesses $u, w$ after $p, A, B$ are computed.

3. Build a Pauli-word moment enumerator for intervals $[-N, N]$, including product constants, translation-invariance canonicalisation, and Hermitian realification.

4. Export the fixed-$h$ feasibility problem (44.1)–(44.4) through a local SDP interface. Record every run under `runs/` with the command line, $h$, level $N$, parameters $e, v^2, u, w$, and solver status.

5. Add mutation tests: perturb a known zero-current density to make the current nonzero; perturb the coboundary sign to make the coboundary reconstruction test fail; perturb a residual coefficient and require a small SDP level to change.

## 46.4 Acceptance for an Exclusion Claim

A future statement "$h$ is excluded at level $N$" must point to:

- the exact Pauli coefficient matrix $h_{\alpha\beta}$ and density convention;
- the algebraic residual tensors and any witnesses $e, v^2, u, w$;

- the finite word sets $\mathcal{W}_N, \mathcal{A}_N$;
- the SDP solver transcript or certificate in a run bundle;
- a worklog entry explaining why the infeasible constraints are necessary for the named route.

### 46.5 Open Gaps

The central charge convention remains unfixed. The two-dimensional edge schema has no implementation. The SDP hierarchy is fixed-$h$; scanning continuous families of $h_{\alpha\beta}$ requires a separate polynomial relaxation. None of the filters replaces CA-15's scaling-limit obligations. They are designed to cheaply reject impossible candidates before the expensive continuum work begins.

# 47 Qubit SDP Implementation Contract

## 47.1 Status, Sources, and Dependencies

**Status: checked implementation contract.** This shard records the Julia layer that implements the fixed-$h$ hierarchy of CA-43–CA-45.

Dependencies: CA-38–CA-45, CONVENTIONS.md (n).

## 47.2 Package Split

The implementation is deliberately layered.

- `QubitPauliWords.jl`: exact positioned Pauli words, exact multiplication phases, translation canonicalisation, and coefficient-term extraction.
- `QubitPoincareWitnesses.jl`: linear algebraic witnesses for $A = Du$ and $B - u - \lambda h + \mu I = Dw$, used before invoking an SDP.
- `QubitMomentSDP.jl`: finite moment bases, moment matrices, relation constraints $\omega(X^* RY) = 0$, JuMP model construction, and Mosek-backed solve helpers.
- `QubitHamiltonianScreening.jl`: deterministic sentinel Hamiltonians and current-collapse filters.

The fixed-$h$ boundary from CA-44 is enforced in the API: an SDP instance takes residual terms as fixed coefficients. If $h$, $u$, $w$, and $\lambda$ are all variables, the problem is no longer this SDP; it becomes a separate polynomial/lifted hierarchy.

## 47.3 Verdicts

The solver helper returns one of three finite-level statuses:

`:excluded`, `:not_excluded_at_level`, `:solver_unknown`.

Only the first is a mathematical rejection, and only for the named level and named residual constraints. The second is not evidence for symmetry; it means that this finite relaxation did not find a contradiction.

An exclusion claim must cite the coefficient matrix $h_{\alpha\beta}$, the residuals, the moment/probe word sets, the solver status, and a run bundle or test that reproduces the result.

# 48 Pauli Word Moment Basis for Infinite Qubit Chains

## 48.1 Status, Sources, and Dependencies

**Status: checked.** The word algebra is implemented without dense matrices and checked against exact Pauli identities.

Dependencies: CA-38, CA-43, CA-46.

## 48.2 Positioned Words

A word is stored as sorted non-identity labels on integer sites. Identities are not stored, but gaps are preserved:

$$X_0 Z_2 \quad \leftrightarrow \quad \texttt{sites=(0,2), labels=(1,3).}$$

The empty word is the identity. Multiplication is sitewise with

$$XY = iZ, \quad YZ = iX, \quad ZX = iY,$$

and the reversed products carrying $-i$. The implementation returns

$$P_u P_v = \phi(u, v) P_{u \cdot v}, \qquad \phi(u, v) \in \{1, -1, i, -i\}.$$

The tests assert $XY = iZ$, $YX = -iZ$, $X^2 = I$, and commutation of disjoint words.

## 48.3 Translation Quotient

The SDP row/column words are not quotiented by translation. They are actual local probes in a finite window. Translation invariance enters only when a moment variable is named:

$$y_s = \omega(P_s) = \omega(P_{\tau^n s}).$$

The canonical representative shifts the first non-identity site to 0. Thus $X_{-3}Z_{-1}$ and $X_0 Z_2$ share the same moment variable.

Because each phase-free Pauli word is self-adjoint, each canonical moment variable is real. Complex numbers enter only as multiplication phases.

# 49 Moment Matrices and Positivity Constraints

## 49.1 Status, Sources, and Dependencies

**Status: checked.** Moment matrices are constructed by `build_qubit_sdp_model`.

Dependencies: CA-43–CA-47.

## 49.2 Affine Moment Matrix

For a finite actual probe set $\mathcal{W}$, define

$$M_{u,v} = \omega(P_u^* P_v) = \phi(u,v) y_{\mathrm{can}(u \cdot v)}.$$

The identity moment is the constant 1. All other canonical words appearing in products or relation constraints receive real JuMP variables.

Write $M = A + iB$, with $A, B$ real affine matrices. Positivity of the complex Hermitian moment matrix is imposed by the real symmetric cone

$$\begin{pmatrix} A & -B \\ B & A \end{pmatrix} \succeq 0. \tag{48.1}$$

For the one-site probe set $\{I, X, Y, Z\}$, the implementation creates three real moment variables and an $8 \times 8$ realified PSD block. This exact dimension is checked in the Julia suite.

## 49.3 Tiny Feasibility Tests

The zero-residual one-site moment problem solves with Mosek as

`:not_excluded_at_level`.

The artificial relation $\omega(I) = 0$ solves as `:excluded`. These are smoke tests for the solver path, not physical Hamiltonian claims.

# 50 Compiling Poincare Residuals into Moment Equations

## 50.1 Status, Sources, and Dependencies

**Status: checked local implementation.** Dense matrix residual oracles from CA-40 are converted into coefficient terms and then into moment equations.

Dependencies: CA-39–CA-40, CA-43–CA-48, CONVENTIONS.md (n).

## 50.2 Algebraic Witnesses

For fixed $h$, the first algebraic gate solves

$$A(h) = Du$$

as a real least-squares linear system after fixing the scalar gauge $u_{00} = 0$. Passing this gate supplies a concrete $u$.

The next gate solves the linearized boost equation

$$B(h) - u - \lambda h + \mu I = Dw, \tag{49.1}$$

again with the scalar gauge $w_{000} = 0$. Here $\lambda = v^2$ and $\mu = v^2 e$. A positive $\lambda$ is the nontrivial-speed branch; a solution with $\lambda = 0$ is only the trivial current-collapsed branch.

## 50.3 Moment Relations

Once residual terms $R = \sum_r c_r P_r$ are fixed, each pair of probe words gives

$$\omega(X^* R Y) = \sum_r c_r \, \phi(X, r, Y) \, y_{\mathrm{can}(XrY)} = 0. \tag{49.2}$$

The implementation splits (49.2) into real and imaginary affine constraints. This is the finite SDP condition promised by CA-43.

## 50.4 Important Boundary

Witness solving is an algebraic necessary filter. The SDP layer does not optimize over $h$, $u$, $w$, or $\lambda$. It checks whether a fixed residual system is compatible with a finite positive moment restriction.

# 51 SDP Levels and the Mosek Backend

## 51.1 Status, Sources, and Dependencies

**Status: checked implementation.** The solver path is exercised by tests and by a recorded smoke run.

Dependencies: CA-44–CA-49.

## 51.2 Level Data

An implemented level consists of:

$$\mathcal{W}_{\mathrm{psd}}, \qquad \mathcal{W}_{\mathrm{rel}}, \qquad \mathcal{R}.$$

The first set is the row/column basis for the PSD moment matrix. The second set supplies $X, Y$ in $\omega(X^* R Y) = 0$. The third is a list of fixed residual coefficient terms.

The convenience constructor uses full Pauli-word windows:

$$\mathcal{W} = \{P_s : \operatorname{supp} s \subseteq [a, a + L - 1]\}.$$

Optional weight truncation is an implementation diagnostic, not the canonical hierarchy.

## 51.3 Solver Status

The feasibility problem has no objective. The status map is:

| MOI termination status | hierarchy verdict |
|---|---|
| `OPTIMAL` | `:not_excluded_at_level` |
| `INFEASIBLE`, `INFEASIBLE_OR_UNBOUNDED` | `:excluded` |
| anything else | `:solver_unknown`. |

This map is intentionally conservative.

## 51.4 Smoke Run

The run bundle

`runs/2026-05-31-qubit-sdp-smoke/`

records three tiny Mosek solves: zero residual, identity-zero relation, and a forced $ZZ = 0$ relation. The first is not excluded at the level; the latter two are excluded. The $ZZ$ case is artificial: it forces $\omega(ZZ \cdot ZZ) = \omega(I) = 0$.

# 52 Sentinel Hamiltonians for the SDP Hierarchy

## 52.1 Status, Sources, and Dependencies

**Status: checked sentinels.** These examples test the implementation and do not assert physical criticality.

Dependencies: CA-35, CA-39–CA-50.

## 52.2 Matrices

The package records five $4 \times 4$ Pauli coefficient matrices:

- a symmetric fully onsite split of $0.25I + 0.7X - 0.3Y + 0.5Z$;
- a commuting classical $ZZ$ density;
- a transverse-Ising-style density $-ZZ - 0.35(XI + IX)$;
- a generic currentful density;
- the fake asymmetric onsite split $XI + IZ$.

The deterministic current filter gives

$$\text{onsite}, ZZ : \texttt{:current_collapsed},$$

and

$$\text{transverse-Ising}, \text{generic}, \text{fake} : \texttt{:currentful}.$$

## 52.3 Gate Outcomes

The onsite and $ZZ$ examples are excluded before SDP for the nontrivial first-moment route because $p = 0$. Their boost-witness solve has only the trivial $\lambda = 0$ branch.

The generic and fake examples have nonzero current but fail $A = Du$. This is the second algebraic gate catching more than current collapse.

The transverse-Ising-style example has nonzero current and passes $A = Du$, but the implemented linear boost witness does not solve (49.1) with the tested data. It is therefore a useful near-candidate sentinel: it reaches a deeper gate before failing.

# 53 Candidate Hamiltonian Selection Roadmap

## 53.1 Status, Sources, and Dependencies

**Status: roadmap.** This shard connects the implementation to the north star: identifying Hamiltonians worth expensive continuum analysis.

Dependencies: CA-15, CA-42–CA-51.

## 53.2 Screening Order

For a fixed nearest-neighbour qubit density $h_{\alpha\beta}$, use:

1. current collapse $p = 0$;
2. conservation witness $A = Du$;
3. boost witness $B - u - \lambda h + \mu I = Dw$ with $\lambda > 0$;
4. finite moment SDP feasibility for the chosen residuals;
5. only then, low-energy/scaling and Koo-Saleur/Virasoro tests.

The point is to reject impossible candidates cheaply. A Hamiltonian surviving all finite checks is not accepted as symmetric; it is merely promoted to the next, more expensive analysis.

## 53.3 Next Implementation Steps

The immediate next step is a run format for candidate scans: input $h_{\alpha\beta}$, current status, witness data, level data, solver status, and a headline verdict. Continuous families require external parameter scans or a later polynomial hierarchy. Named physical model families need registered sources before the report can claim criticality, universality class, or continuum CFT content.

# 54 Qubit Candidate Scan Contract

## 54.1 Status, Sources, and Dependencies

**Status: checked implementation.** The scan layer packages the CA-46–CA-52 fixed-$h$ gates into reproducible candidate runs.

Dependencies: CA-39, CA-46–CA-52, CONVENTIONS.md (m)–(q).

## 54.2 Contract

Each scan point is a real $4 \times 4$ Pauli coefficient matrix $h_{\alpha\beta}$. The gate order is:

$$p = i[h_j, h_{j+1}], \qquad A = Du, \qquad B - u - \lambda h + \mu I = Dw,$$

followed, only when requested, by finite moment-SDP checks.

The serialized row records the family, parameters, coefficient matrix, current norm, witness residual norms, speed data, terminal gate, and scoped verdict. The scope field records the fixed first-moment route.

## 54.3 Verdict Boundary

`excluded_current_collapsed` and `excluded_no_conservation_witness` are early algebraic failures for the fixed current route. `excluded_no_boost_witness` means the exact local boost-witness equation failed; it does not reject other lattice stress-tensor constructions such as Koo–Saleur/free-fermion routes. This distinction is essential for sourced critical transverse Ising.

# 55 Pauli Coefficients for Scan Families

## 55.1 Status, Sources, and Dependencies

**Status: checked implementation convention.** The family constructors are implemented in the Julia backend.

Dependencies: CA-34–CA-53, CONVENTIONS.md (m), (q).

## 55.2 Implemented Maps

The scan uses $(I, X, Y, Z)$ row/column order. A one-site field $f_a\sigma_j^a$ is represented on bonds by

$$h_{a0} = h_{0a} = f_a/2.$$

Thus the self-dual transverse-Ising sample is

$$h = -ZZ - \frac{1}{2}(XI + IX),$$

which is the bond-density version of the sourced transverse-Ising form.

The XXZ constructor uses

$$h = J(XX + YY + \Delta ZZ),$$

and Heisenberg is the $\Delta = 1$ specialization. The transverse-XY constructor uses $J_xXX + J_yYY + h_zZ$ with the same symmetric field split.

## 55.3 Synthetic Families

XYZ, compass, DM-style, field-deformed Heisenberg, and deterministic dense generic matrices are stress tests. They exercise the gates but carry no criticality or universality-class claim in this lab book.

# 56 Locally Sourced Spin-Chain Inputs

## 56.1 Status, Sources, and Dependencies

**Status: source audit.** This shard records which family names the first scan may treat as physical rather than synthetic.

Dependencies: CA-14, CA-52–CA-54.

## 56.2 Allowed Physical Labels

The defect-gauging source gives an effective qubit Hamiltonian proportional to $\sum_i (I + Z_i Z_{i+1} + X_i)$, identifies it as transverse Ising, and states that it gives a critical theory. The Ising anyon-chain source independently identifies the localized chain with critical TFIM and the Ising CFT.

The lattice-fermion source supports transverse XY as a Jordan–Wigner image and also records TFIM/Koo–Saleur convergence on the even observable subalgebra. The XXZ source gives a Pauli-chain Hamiltonian and the gapless/free-boson continuum anchors for $-1 < \Delta \leq 1$, with Heisenberg as a special point.

## 56.3 Excluded From Physical Labeling

The first scan does not make physical claims for XYZ, DM, compass, clock, Potts, or Ashkin–Teller. Potts is qutrit in the local source. The sourced Ising zigzag/Ashkin–Teller relation uses a three-site term, outside the present nearest-neighbour two-qubit input class.

# 57 Synthetic and Stress-Test Hamiltonian Grids

## 57.1 Status, Sources, and Dependencies

**Status: checked implementation input.** These grids are generated by the candidate-scan sample constructor.

Dependencies: CA-53–CA-55, CONVENTIONS.md (q).

## 57.2 Grids

The synthetic block includes XYZ triples on $\{-1, 0, 1\}^3 \setminus \{0\}$, XY anisotropy, compass $XX/ZZ$, DM-style antisymmetric $XY - YX$, Heisenberg with one field component, and 24 deterministic dense generic matrices. The generic matrices are generated by fixed sine/cosine formulas in the code rather than by an ambient random number generator.

## 57.3 Purpose

These samples answer engineering questions: Does the conservation gate reject generic dense inputs? Do structured bilinear families reach the boost gate? Do current-collapsed one-axis cases remain caught? They do not answer whether XYZ or DM chains are critical.

# 58 Current, Conservation, and Boost Witness Gates

## 58.1 Status, Sources, and Dependencies

**Status: checked scan gates.** The implementation calls the CA-40 and CA-49 helpers pointwise. Dependencies: CA-39–CA-40, CA-49, CA-53–CA-56.

## 58.2 Gates

The current gate checks whether $p = i[h_j, h_{j+1}]$ collapses. The conservation gate solves $A = Du$. The boost gate solves

$$B - u - \lambda h + \mu I = Dw$$

with $w_{000} = 0$ and requires nonzero positive speed for the nontrivial operator route.

## 58.3 Interpretation

The first full scan has no survivor of the exact boost gate. The self-dual TFIM point, which is locally sourced as critical, passes current and conservation but fails the exact boost witness. Therefore this gate is best read as a test for an exact UV first-moment construction, not as a general criterion for CFT emergence.

# 59 Optional SDP Levels for Scan Survivors

## 59.1 Status, Sources, and Dependencies

**Status: implemented option, unused by the first broad run.** The scan API can call `solve_qubit_sdp` after a nontrivial boost witness.

Dependencies: CA-44, CA-50, CA-53–CA-57.

## 59.2 Level Data

An SDP row must record the PSD window, relation window, moment count, relation constraint count, optimizer, and solver status. The status vocabulary remains

`:excluded`, `:not_excluded_at_level`, `:solver_unknown`.

## 59.3 First-Run Boundary

The 99-point run has no nontrivial exact boost witness, so it does not invoke Mosek per point. This keeps the run honest: adding SDP rows before fixing a residual witness would mix the fixed-$h$ SDP with a polynomial variable-witness problem.

# 60 Reproducible Scan Bundle Format

## 60.1 Status, Sources, and Dependencies

**Status: checked run bundle.** The bundle is

`runs/2026-05-31-qubit-candidate-scan/`.

Dependencies: CA-53–CA-58 and AGENTS.md Law 3.

## 60.2 Files

`inputs.toml` records the Pauli basis, indexing rule, family inventory, source-kind inventory, and sample count. `summary.toml` records verdict counts and notable sourced points. `results.toml` records one row per sample, including coefficients, parameters, terminal gate, residual norms, and scope.

## 60.3 Acceptance

The run command prints the same headline counts that the TOML stores: 99 samples, 9 current-collapsed, 27 conservation failures, and 63 exact boost-witness failures.

# 61 First Candidate Scan Results

## 61.1 Status, Sources, and Dependencies

**Status: checked run result.** This shard reports only the run bundle named in CA-59.

Dependencies: CA-53–CA-59.

## 61.2 Headline Table

The 99 samples split as:

| verdict | count |
|---|---|
| `excluded_current_collapsed` | 9 |
| `excluded_no_conservation_witness` | 27 |
| `excluded_no_boost_witness` | 63 |

No point reaches `not_excluded_algebraic`.

## 61.3 Notable Points

The self-dual TFIM sample

$$-ZZ - \frac{1}{2}(XI + IX)$$

is currentful and has a conservation witness, but its exact boost residual norm is nonzero in the run. The Heisenberg point and the $XX$ point behave the same way: they reach the boost gate and fail it.

Because TFIM and XXZ have local source anchors as critical or gapless models, the conclusion is about the route: the exact CA-39 boost witness is too strict as a universal candidate filter.

# 62 From Scan Failures to Scaling Work

## 62.1 Status, Sources, and Dependencies

**Status: roadmap from checked failures.** This is the handoff after CA-60.
Dependencies: CA-14, CA-15, CA-42, CA-52–CA-60.

## 62.2 Queue

Do not retry the same exact boost-witness gate expecting TFIM to pass. The next candidate path should use the sourced Koo–Saleur/free-fermion/OAR evidence chain for TFIM and transverse XY, and the Koo–Saleur/XXZ literature for XXZ. Those routes use low-energy or scaling-limit stress tensors, not an exact finite-support UV boost density satisfying CA-39.

## 62.3 SDP Upgrade

The state-level hierarchy should next learn how to quotient or relax the unknown $Dw$ term without turning the problem into an uncontrolled polynomial relaxation. Until then, the implemented SDP layer is still useful for fixed residual systems and artificial contradictions, but the broad scan has shown where the exact-local route stops.

# 63 The Categorical Borchers–Uhlmann Pipeline: a Programme Refinement

## 63.1 Status, Sources, and Dependencies

**Status: proposal/question.** This shard records a programme refinement, not a theorem. Its central proposal is that the CA-02/CA-03 indefinite-particle grammar should be reinterpreted first as an observable $*$-algebra of local words, and only later as a Hilbert space after a state and GNS quotient have been chosen.

Dependencies: CA-01, CA-02, CA-03, CA-09, CA-43, CA-44, CA-57, CA-60, CA-61; CONVENTIONS.md (q); companion shards CA-63, CA-64, CA-65, CA-66 (this block).

## 63.2 Refined North Star

**Proposal.** The 2026-07-05 refinement of the north star is a recipe, as functorial as possible, which starts from a unitary fusion category or modular tensor category $\mathcal{C}$, together with any conformal data required by the target CFT, and produces a family of microscopic lattice models whose scaling limit realises a chiral or full CFT whose defects and excitations recover the input category.

**Proposal.** The key reinterpretation is algebra-first. CA-03 builds

$$\mathfrak{F}(H) = \widehat{\bigoplus}_{n\geq 0} H^{\widehat{\otimes} n}$$

as a Hilbert completion and explicitly identifies the underlying tensor algebra $T(H) = \bigoplus_n H^{\otimes n}$. In this shard, the tensor algebra is not primarily the pre-Hilbert finite-particle subspace. It is the proposed free local-word $*$-algebra from which observables are generated. The local anchor for this style is the free unital $*$-algebra of generators $\Phi(f)$ over test functions, quotiented by a two-sided $*$-ideal of relations; the registered source builds exactly this object (it does not use the name Borchers–Uhlmann, which we keep as informal terminology).

**Proposal.** Candidate lattice models emerge only after three extra choices: a reference state $\omega$, approximate discrete Virasoro or Koo–Saleur-type residual relations, and quotienting by the state-null relations. In symbols, the intended order is

$$\text{local-word } *\text{-algebra} \quad \xrightarrow{\omega,\mathcal{R}} \quad \mathfrak{N}_{\omega,\mathcal{R}} \quad \xrightarrow{\text{GNS}} \quad (\mathcal{H}_\omega, \pi_\omega, \Omega_\omega),$$

not "choose a Hilbert space first, then search for relations".

## 63.3 Five Stages

**Stage 0: input data. Status: prior-shard question.** The input is a unitary fusion category or modular tensor category, plus conformal/OPE data when the target CFT needs more than $\mathcal{C}$. CA-01 states this as the global programme map. CA-09 gives the first checked kinematic example: Fibonacci fusion admissibility produces a sector projector, but full anyons still need coherent $F/R$-data.

**Stage 1: free algebra of anyonic words. Status: proposal.** Replace the Hilbert-first reading of CA-03 by an observable-algebra reading: generate formal local anyonic words from the categorical variable-$N$ sectors

$$\bigoplus_{N,c,\text{configs}} \operatorname{Mor}(X_c, X_{a_1} \otimes \cdots \otimes X_{a_N}).$$

The precise algebra generators, involution, locality grading, and completion policy are fixed in CA-65 and CA-66 (companion shards, this block).

**Stage 2: categorical relations. Status: mixed checked/proposal.** Fusion admissibility is already a sector-projector relation in CA-09. Additional categorical relations should include defect/MPO relations. Hollands supplies the finite-size anchor: the bipartite anyonic-chain MPO algebra is isomorphic to the CFT defect algebra at finite size, while the persistence of that algebra in the scaling limit is explicitly left as an analytic question.

**Stage 3: reference state and relaxed residuals. Status: proposal, currently exclusion-mode.** The state must be chosen before a Hilbert space is claimed. CA-43 and CA-44 provide the present finite architecture: translation-invariant moments, positivity, and residual relation constraints of the form

$$\omega(X^* R Y) = 0.$$

In the GNS reading, these are the finite probe shadows of the null-relation condition. CA-63 (companion shard, this block) is the construction-mode state existence route; CA-64 (companion shard, this block) is the relaxation route.

**Stage 4: GNS quotient and lattice model. Status: sourced scaffold plus proposal.** Once $(\mathcal{A}, \omega)$ and a chosen residual set $\mathcal{R}$ are fixed, the candidate Hilbert space and operator representation are produced by the GNS quotient. The local OAR sources support this ordering: build scaling maps and an inductive-limit algebra, construct compatible/projective limit states, then perform GNS to get Hilbert-space representations and compatible representation systems.

**Stage 5: continuum limit and fidelity. Status: precedent plus question.** The continuum-limit technology is anchored by OAR and the free-fermion Koo–Saleur theorem: lattice Virasoro approximants and correlation functions converge in that sourced free-fermion setting. The anyonic precedent is Stottmeister's braiding-RG note, where braiding defines a real-space RG for anyonic chains and the Ising-chain example reaches the Ising CFT vacuum. The general shape of the open problem is now also locally sourced: Zini–Wang define scaling limits of quantum theories as double colimits and conjecture that every unitary minimal model VOA arises as a strong scaling limit with approximating lattice Virasoro observables. The categorical fidelity target remains a question:

$$\text{DHR/defect content of the limit} \simeq \mathcal{C}?$$

Hollands makes this a precise open problem by proving a finite algebraic defect correspondence and asking whether it persists under scaling limits.

### 63.4 Two Load-Bearing Constraints

**Constraint 1: the Jones no-go changes the order.** The locally registered no-go theorem states that the block-spin/Temperley–Lieb semicontinuous Hilbert space does not support a chiral CFT whose Hamiltonian generates circle translation as a continuum limit of lattice rotations; in the main theorem, triadic-rotation-rigid vectors in the triadic limit of an annular TL representation are necessarily zero. The OAR discussion corroborates the programme reading: nontrivial Thompson implementers are not implementable in the presence of nonzero central charge, and the Koo–Saleur strategy succeeds in those situations. The consequence is load-bearing: do not expect a fixed inductive-limit Hilbert space with dyadic rotations to carry the continuum geometry first. The

state comes first; geometry and symmetry are recovered through state-dependent scaling limits and generator convergence.

**Constraint 2: exact UV relations are too strict.** CA-57 defines the current, conservation, and exact boost-witness gates. CA-60 reports the checked 99-point scan: no point reaches `not_excluded_algebraic`, and the locally sourced self-dual TFIM point fails the exact boost witness. CA-61 and CONVENTIONS.md (q) record the route-scoped reading: this rejects the exact first-moment route, not the Ising scaling limit. Therefore Stage 3 must use quantified state-level or GNS-norm residuals, not exact coefficient identities, as the default relation language.

### 63.5 Functoriality and Fidelity Assessment

**Proposal.** Stages 1 and 2 should be functorial in principle: free algebras and imposed categorical relations are natural constructions once the generator and convention data are fixed. This is not yet a theorem; CA-65 records the intended functor on pairs of a category and a site object.

**Proposal with sourced scaffold.** GNS should be treated functorially on compatible $(\mathcal{A}, \omega)$-pairs. The local source support is not an abstract category theorem; it is the OAR architecture in which projective nets of states induce compatible nets of GNS representations and Hilbert-space connecting maps.

**Question.** The state assignment is the non-functorial crux. The realistic target is canonicity by a universal property: define a compact or closed convex set of states satisfying the relaxed residual constraints, then prove non-emptiness, uniqueness or controlled faces, and fidelity of the resulting limit. The strongest intended theorem would say that this state set exists and its continuum DHR/defect category recovers the input $\mathcal{C}$.

### 63.6 Boundary

This shard does not fix the categorical local-word algebra; the $F/R$-symbol gauge, object indexing, bracketing, and locality/completion conventions are fixed in CA-65/CA-66, not here.

It does not prove existence of the reference state. CA-43/CA-44 are exclusion-mode finite constraints; CA-63 is the companion construction-mode shard.

It does not fix the relaxation convention. The exact-relation scan failure only says exact UV boost witnesses are too strict for the current route; CA-64 must define the norm, scale, tolerance, and limiting quantifiers.

It does not prove a new continuum theorem. The only continuum precedents cited here are the sourced free-scalar/OAR, free-fermion Koo–Saleur, and Ising braiding-RG examples; the Zini–Wang minimal-model statement is cited as a conjecture, not a result.

It does not claim that SDP feasibility or finite residual smallness implies symmetry. CA-44 already records that feasibility does not prove a global state, positive energy, or a continuum Poincare/Virasoro representation.

# 64 All-Level Moment Feasibility Gives a Qubit Reference State

## 64.1 Status, Sources, and Dependencies

**Status: derived compactness theorem, conditional on standard C*-algebra facts.** This upgrades CA-44 only for the exact monotone full-window hierarchy. It does not apply to optional weight truncations, arbitrary implemented diagnostic levels, solver-unknown statuses, or nonmonotone residual lists.

Dependencies: CA-43, CA-44, CA-62, CONVENTIONS.md (n), CONVENTIONS.md (p).

## 64.2 Canonical Full-Window Levels

Let

$$I_N = [-N, N], \qquad \mathcal{A}_N = \bigotimes_{j \in I_N} M_2(\mathbb{C}),$$

and let $\mathcal{W}_N$ be the set of all Pauli words supported in $I_N$. For this shard, a level means the full-window problem:

$$y_\emptyset = 1, \qquad y_s = y_{\tau^k s} \tag{63.1}$$

whenever both words fit in the level window,

$$M_N(y)_{u,v} = \omega(P_u^* P_v) \succeq 0, \qquad u, v \in \mathcal{W}_N, \tag{63.2}$$

and every fixed residual relation

$$\omega(X^* R Y) = 0 \tag{63.3}$$

whose product support fits in $I_N$. The residual family is fixed in advance, or is enlarged monotonically. This monotonicity is part of the theorem; without it the compactness argument below has no nesting premise.

## 64.3 Finite Moments Are Finite States

Given a feasible $y$ at level $N$, define a linear functional on $\mathcal{A}_N$ by

$$\omega_N(P_s) = y_s$$

and extend linearly in the Pauli-word basis. If $X = \sum_{u \in \mathcal{W}_N} c_u P_u$, then

$$\omega_N(X^* X) = \sum_{u,v} \overline{c_u} c_v \, \omega_N(P_u^* P_v) = c^* M_N(y) c \geq 0. \tag{63.4}$$

Since every positive element in a finite matrix algebra has a square root, (63.4) makes $\omega_N$ positive. The normalization $y_\emptyset = 1$ makes it unital, hence a state on $\mathcal{A}_N$.

Equivalently, with $d_N = 2^{|I_N|}$,

$$\rho_N = d_N^{-1} \sum_{s \in \mathcal{W}_N} y_s P_s \tag{63.5}$$

satisfies $\omega_N(A) = \mathrm{Tr}(\rho_N A)$, the density-matrix presentation of the finite state. Thus $M_N(y) \succeq 0$ if and only if $\rho_N \succeq 0$.

## 64.4 Compactness

Let $\mathcal{A}$ be the quasi-local spin C*-algebra, the norm closure of the local algebras. The normalized product trace outside $I_N$ gives a unital completely positive conditional expectation

$$E_N : \mathcal{A} \to \mathcal{A}_N.$$

Therefore $\omega_N \circ E_N$ is a global state and agrees with $\omega_N$ on $\mathcal{A}_N$. This proves nonemptiness of the level set below whenever the level moment problem is feasible.

Let $K_N \subset S(\mathcal{A})$ be the set of global states whose restriction to $\mathcal{A}_N$ satisfies the affine level-$N$ translation and residual constraints. Positivity is automatic for elements of the state space. Each constraint is a finite linear equation in evaluations $\varphi \mapsto \varphi(A)$, hence $K_N$ is weak-* closed. Since $S(\mathcal{A})$ is weak-* compact, each $K_N$ is compact.

The full-window levels are nested. Every level-$N$ translation equality and every level-$N$ residual test also appears at level $N+1$. Hence

$$K_{N+1} \subseteq K_N. \tag{63.6}$$

If every full-window level is feasible, the $K_N$ are nested nonempty compact sets, so

$$\bigcap_{N \geq 0} K_N \neq \emptyset. \tag{63.7}$$

Choose $\omega$ in this intersection.

For any Pauli word $s$ and any integer shift $k$, choose $M$ so large that the intermediate translates from $s$ to $\tau^k s$ fit in $I_M$. Since $\omega \in K_M$, (63.1) gives

$$\omega(P_s) = \omega(P_{\tau^k s}).$$

By linearity, density of the local Pauli algebra, and continuity of the shift automorphism, $\omega \circ \tau^k = \omega$ on all of $\mathcal{A}$.

## 64.5 Residual Relations in the GNS Representation

Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega)$ be the GNS representation. Fix one residual density $R$. For local Pauli words $X, Y$, choose $N$ so that $X^*RY$ fits the level window. Because $\omega \in K_N$,

$$\langle \pi_\omega(X)\Omega_\omega, \pi_\omega(R)\pi_\omega(Y)\Omega_\omega \rangle = \omega(X^*RY) = 0. \tag{63.8}$$

The vectors $\pi_\omega(A)\Omega_\omega$, with $A$ local, are dense in the GNS Hilbert space. Since $\pi_\omega(R)$ is bounded, (63.8) on this dense subspace implies

$$\pi_\omega(R) = 0. \tag{63.9}$$

Translation-invariant states implement the lattice shift by a GNS unitary, so the translated residual densities vanish as well.

## 64.6 Exact Alternative

Conversely, if a translation-invariant state $\omega$ satisfies all residual relations, then $y_s = \omega(P_s)$ is feasible at every full-window level. Indeed, $y_\emptyset = 1$, translation invariance gives (63.1), positivity of $\omega$ gives (63.2), and the residual hypothesis gives (63.3).

Thus, for fixed exact residual data, there is no third mathematical possibility:

$$\text{some full-window level is infeasible} \quad \text{or} \quad \text{a global reference state exists.} \tag{63.10}$$

This is an exact compactness statement, not a finite-time algorithm.

## 64.7 Non-Implications

The state in (63.7) need not be unique. It need not be a ground state, a positive-energy state, a clustering state, or a scaling-limit state. The argument does not construct a continuum limit and does not produce a projective-unitary Poincare or Virasoro representation. It only proves that the imposed finite residual relations have a genuine C*-algebraic state model.

## 64.8 Boundary

Finite feasibility at checked levels still proves no existence theorem. The compactness result requires exact feasibility at every monotone full-window level. Numerically, exclusion remains semi-decidable by finding a certified infeasible finite level; existence is not certified by any finite prefix of feasible solver runs. The finite witnesses for the proof ingredients (density-matrix reconstruction, extension moment preservation, level nesting, and a positivity mutation) are checked in the Julia testset "qubit full-window moment compactness witnesses".

# 65 Quantified Relaxations of the Qubit Symmetry Gates

## 65.1 Status, Sources, and Dependencies

**Status: proposal with checked finite pieces.** The definitions below are finite Pauli-coefficient linear algebra plus fixed-residual moment SDPs. This shard answers the CA-61 obligation to relax or quotient the boost relation instead of requiring an exact local witness, and it supplies the Stage-3 relation language of CA-62.

Dependencies: CA-39, CA-42, CA-43, CA-44, CA-57, CA-60, CA-61, CA-62, CONVENTIONS.md (m), (n), (p), (q).

## 65.2 Coefficient Norm and Gauge

Let $V_L$ be the Pauli coefficient space of $L$-site densities, with

$$\langle r, s\rangle_\tau = \sum_a \overline{r_a} s_a.$$

This is the normalized tracial Hilbert–Schmidt inner product. Padding by trailing identities is denoted $\iota_{L,N} : V_L \to V_N$.

The coboundary map of CONVENTIONS.md (n) is

$$(Du)_{a_0\cdots a_L} = u_{a_0\cdots a_{L-1}}\delta_{a_L 0} - \delta_{a_0 0} u_{a_1\cdots a_L}.$$

Residuals are distances to affine subspaces in this norm. Scalar witness components are gauge directions; they are fixed to zero for reported witnesses, but the distance itself is gauge-invariant.

## 65.3 Algebraic Profiles

The current gate remains hard for the nonzero-speed first-moment route:

$$\|p\|_\tau = 0 \quad\Longrightarrow\quad \text{fail this route.}$$

For $A = i[h_j + h_{j+1}, p_j]$, define the conservation profile

$$\epsilon_c(L; h) = \operatorname{dist}\big(\iota_{3,n} A, \iota_{L+1,n} \operatorname{im} D_L\big), \qquad n = \max(3, L+1).$$

The reported dimensionless value is

$$\widehat{\epsilon}_c(L; h) = \epsilon_c(L; h) / \max(\|A\|_\tau, \varepsilon_{\text{mach}}).$$

For candidates with exact conservation at support $L$, define the exact-conservation boost profile

$$\epsilon_b^0(L; h, \lambda, e) = \min_{u,w:\, D_L u = A} \|\iota B - \iota u - \lambda\, \iota(h - eI) - D_L w\|_\tau,$$

with $B = i[p_j, h_{j+1}] + 2i[p_j, h_{j+2}]$ and $\lambda = v^2$. Letting $u$ float freely in the boost residual is wrong: it can absorb boost error while no longer witnessing the conservation coboundary. For candidates without exact conservation, use the joint profile

$$\epsilon_{cb}^2 = \min_{u,w} \|\widehat{A - Du}\|_\tau^2 + \|B - u - \widehat{\lambda(h - eI)} - Dw\|_\tau^2.$$

The raw profiles are nonincreasing in $L$, because a support-$L$ witness is also a support-$L+1$ witness after padding by an identity. The speed $\lambda$ is optimized only by external scan or bounded quadratic programming, with $\mu = \lambda e$ as the linear scalar variable; the sign of the optimizing $\lambda$ is a recorded diagnostic, not an assumption.

## 65.4 State-Level Residuals

For a fixed residual density $R$, the GNS quantity is

$$\|\pi_\omega(R)\Omega_\omega\|^2 = \omega(R^*R).$$

At moment level $N$, define

$$v_N(R) = \min_y \omega_y(R^*R)$$

subject to the CA-43–CA-44 moment PSD, translation-invariance, normalization, and optional stationarity constraints. For fixed $R$, this is an SDP because $\omega_y(R^*R)$ is affine in $y$. Normalize by

$$\nu_N(R) = v_N(R)/\tau(R^*R),$$

when $\tau(R^*R) > 0$; the tracial value $\tau(R^*R) = \sum_s |r_s|^2$ is computable exactly from Pauli coefficients. A dual (certified) lower bound $\nu_N(R) \geq \delta > 0$ is an exclusion certificate for the relaxed route; $\nu_N(R)$ small is survival, never symmetry evidence.

Do not optimize $u, w, \lambda, e$ jointly with the moments in this SDP tier. For $R(a) = R_0 + \sum_i a_i R_i$, the expression $\omega(R(a)^*R(a))$ is quadratic in $a$ with coefficients affine in $y$, hence bilinear jointly. That joint minimization is not a plain SDP. Alternation gives upper-bound candidates; certified lower bounds require fixed witnesses or a bounded polynomial/sum-of-squares lift recorded as a new convention before use.

## 65.5 Scaling Reading

A continuum claim is not a fixed-$N$ claim. A scaling statement must name: state sequence, renormalisation maps, vacuum subtraction, and operator topology, as required by CA-14 and CA-15. The target form is that normalized GNS residuals vanish on a growing long-wavelength window:

$$\sup_{|m|,|n|\leq K_N} \nu_N(r^{\sigma\sigma'}_{mn}) \to 0, \qquad K_N/N \to 0,$$

applied to the CA-42 Witt/Virasoro residual densities $r^{++}_{mn}, r^{--}_{mn}, r^{+-}_{mn}$. Modes near the Brillouin scale are not expected to close first; long-wavelength modes are the diagnostic window.

## 65.6 Sentinel Expectations

These are predictions to be checked by the relaxed re-scan, not results. The self-dual TFIM point is currentful, has an exact conservation witness, and fails the exact boost gate with recorded residual norm 1.8257418583505538; the relaxed verdict should therefore be survival into a GNS/scaling queue rather than terminal exclusion. Fully on-site and classical commuting densities still fail the hard current gate. Generic dense samples must show actual profile decay to survive; a single small value at one support is not a pass.

## 65.7 Boundary

No continuum theorem is claimed. The profiles and SDP objectives are finite diagnostics; converting decay of $\nu_N$ into a continuum symmetry statement requires the scaling payload of CA-15 and remains open. The longer-support TFIM profile values computed during design are not quoted here because their run bundle is not yet registered; only the recorded scan artifact is cited.

# 66 Anyonic Word Algebra from a Site Object

## 66.1 Status, Sources, and Dependencies

**Status: proposal shard from the adopted W4 design.** The finite local algebras are source-local C*-algebra objects; the free presentation and functoriality clauses are the adopted interface and still need implementation tests.

Dependencies: CA-02, CA-03, CA-09, CA-62, CA-63, CA-64, and CONVENTIONS.md (a), (b) (fixed with this block), (c), (f), (g), and (r).

## 66.2 Site Object First

The input to the word algebra is a pair $(\mathcal{C}, O)$, not the category $\mathcal{C}$ alone. Here $\mathcal{C}$ is a unitary fusion category and $O \in \mathcal{C}$ is the one-site object. The default mobile hard-core choice is

$$O = \mathbf{1} \oplus X,$$

where $X$ is the allowed non-vacuum species object. For Fibonacci, $X = \tau$; for an all-species hard-core site, $X = \bigoplus_{a \neq \mathbf{1}} X_a$.

For a finite open interval $I$ with $|I| = L$, fix the left-associated tensor power from CONVENTIONS.md (c),(r):

$$O_I := (((O \otimes O) \otimes O) \cdots) \otimes O.$$

The local anyonic observable algebra is

$$\mathcal{A}_I(\mathcal{C}, O) := \mathrm{End}_{\mathcal{C}}(O_I), \qquad f^* := f^\dagger.$$

Because endomorphism algebras in a unitary category are finite-dimensional unitary *-algebras, each $\mathcal{A}_I$ is a finite-dimensional C*-algebra.

Choose representatives $\{X_c\}_{c \in \mathrm{Irr}(\mathcal{C})}$, with $\mathbf{1}$ first by CONVENTIONS.md (a) as fixed in this block, and define

$$m_c(L) := \dim_{\mathbb{C}} \mathrm{Hom}_{\mathcal{C}}(X_c, O_I).$$

After choosing orthonormal bases of these Hom spaces,

$$\mathcal{A}_I(\mathcal{C}, O) \cong \bigoplus_{c \in \mathrm{Irr}(\mathcal{C})} M_{m_c(L)}(\mathbb{C}).$$

This matrix form is a presentation. The abstract algebra $\mathrm{End}_{\mathcal{C}}(O_I)$ is basis-free; fusion-tree bases and $F$-symbols enter only when we write matrices.

## 66.3 Placed Matrix Blocks

Let $w, w' \in \mathrm{Hom}_{\mathcal{C}}(X_c, O_I)$ be basis isometries in the same charge block and let $f \in \mathrm{End}_{\mathcal{C}}(X_c)$. The corresponding placed matrix block is

$$E_I(w', f, w) := w' \circ f \circ w^\dagger \in \mathcal{A}_I(\mathcal{C}, O).$$

For basis vectors $u, v, w, z$ in compatible charge blocks,

$$E_I(u, g, v)\, E_I(w, f, z) = \delta_{v,w}\, E_I(u, g \circ f, z),$$

and

$$E_I(w', f, w)^* = E_I(w, f^\dagger, w').$$

For simple $X_c$, $f$ is a scalar multiple of $\mathrm{id}_{X_c}$, so these are ordinary matrix units inside the $c$-block. This is the categorical replacement for positioned Pauli words in CA-47: rows and columns are not bare strings of letters, but charge-resolved fusion-tree basis vectors.

## 66.4 Inclusions, Shift, and AF Limit

For $I \subset J$, the local inclusion is obtained by tensoring with identities on the added site factors, using the fixed bracketing convention:

$$f \longmapsto \mathrm{id}_O^{\otimes \ell} \otimes f \otimes \mathrm{id}_O^{\otimes r} .$$

On a one-sided chain the shift is

$$\alpha(f) = \mathrm{id}_O \otimes f.$$

The quasi-local observable algebra is the inductive-limit C*-algebra

$$\mathcal{A}(\mathcal{C}, O) := \overline{\bigcup_I \mathcal{A}_I(\mathcal{C}, O)}.$$

It is AF by construction: it is the norm closure of an increasing union of finite-dimensional C*-algebras.

The Bratteli diagram is the fusion graph of $O$. If $O_I \cong \bigoplus_c m_c(L) X_c$, then

$$m_d(L+1) = \sum_c m_c(L)\, N_{c,O}^d, \qquad N_{c,O}^d := \dim \mathrm{Hom}_{\mathcal{C}}(X_d, X_c \otimes O).$$

Sanity check: for $\mathcal{C} = \mathrm{Hilb}_{\mathrm{fd}}$ and $O = \mathbb{C} \oplus \mathbb{C}^d$, this gives

$$\mathcal{A}_L \cong M_{(d+1)^L}(\mathbb{C}),$$

and the inductive limit is the ordinary UHF spin-chain algebra with a $(d{+}1)$-dimensional maybe-spin site.

## 66.5 The BU-Style Free Pre-Layer

The adopted free layer is the tensor *-algebra

$$\mathcal{BU}_0(\mathcal{C}, O) := T(V), \qquad V := \bigoplus_{j \in \mathbb{Z}} \bigoplus_{a \neq \mathbf{1}} \mathrm{Hom}_{\mathcal{C}}(X_a, O)_j.$$

The subscript $j$ records the lattice site. The involution is the formal dual/adjoint operation fixed by CONVENTIONS.md (b),(r); in a fusion-tree matrix presentation the unitary gauge is mandatory so that $*$ is entrywise conjugate transpose.

The intended presentation map is

$$\pi : \mathcal{BU}_0(\mathcal{C}, O) \twoheadrightarrow \mathcal{A}(\mathcal{C}, O),$$

with

$$J_{\mathrm{fus}} := \ker \pi.$$

Thus

$$\mathcal{BU}_0(\mathcal{C}, O)/J_{\mathrm{fus}} \cong \mathcal{A}(\mathcal{C}, O).$$

The ideal $J_{\mathrm{fus}}$ is kinematic. It contains the hard-core site relations, fusion admissibility, $F$-coherence, and the matrix-block relations above. This mirrors the AQFT construction in which one starts from a free unital *-algebra and quotients by a two-sided *-ideal of relations.

The load-bearing split is therefore

$$\mathcal{BU}_0 \xrightarrow{\pi \text{ (kinematic fusion)}} \mathcal{A}(\mathcal{C}, O) \xrightarrow{\text{state/GNS}} H_\omega.$$

Fusion is not deferred to the state. The GNS null space may impose dynamical or scaling relations, but category fidelity lives already in $J_{\mathrm{fus}}$.

## 66.6 Defect Algebra Is a Different Layer

Hollands' double triangle algebra is a finite-dimensional vector space built from Hom spaces such as $\mathrm{Hom}_M(cd, \bar{a}b)$, with its own vertical product. That algebra is not the local observable algebra $\mathcal{A}_I(\mathcal{C}, O)$. In this programme it is the deferred defect/MPO/symmetry layer represented on, or acting by endomorphisms of, the observable tower. A later shard must specify that representation and its compatibility with the shift.

## 66.7 Functoriality Target

For a unitary monoidal dagger functor $F : \mathcal{C} \to \mathcal{C}'$ and a chosen unitary $F(O) \cong O'$, the expected finite-level map is

$$\mathrm{End}_{\mathcal{C}}(O^{\otimes L}) \longrightarrow \mathrm{End}_{\mathcal{C}'}((O')^{\otimes L}),$$

given by applying $F$ and conjugating by the chosen site-unitary tensor power. The coherence data make this pseudofunctorial rather than strictly functorial on the nose. The free construction $\mathcal{BU}_0$ is natural at the generator level, and $\pi$ should be natural once $J_{\mathrm{fus}}$ is implemented.

## 66.8 Lamport Dependency Skeleton

$D65.1$ : Input pair $(\mathcal{C}, O)$, $\quad O = \mathbf{1} \oplus X$ by default.
$D65.2$ : $O_I = O^{\otimes |I|}$ with fixed left bracketing.
$D65.3$ : $\mathcal{A}_I(\mathcal{C}, O) = \mathrm{End}_{\mathcal{C}}(O_I), \quad f^* = f^\dagger$.
$D65.4$ : $m_c(L) = \dim \mathrm{Hom}_{\mathcal{C}}(X_c, O^{\otimes L}), \quad \mathcal{A}_L \cong \bigoplus_c M_{m_c(L)}(\mathbb{C})$.
$D65.5$ : $E_I(w', f, w) = w' f w^\dagger$.
$D65.6$ : $\mathcal{BU}_0 = T\Big(\bigoplus_{j, a \neq \mathbf{1}} \mathrm{Hom}(X_a, O)_j\Big)$.
$D65.7$ : $J_{\mathrm{fus}} = \ker(\mathcal{BU}_0 \twoheadrightarrow \mathcal{A})$.
$T65.1$ : $E(u, g, v)E(w, f, z) = \delta_{v,w} E(u, g \circ f, z)$.
$T65.2$ : $E(w', f, w)^* = E(w, f^\dagger, w')$.
$T65.3$ : $m_d(L+1) = \sum_c m_c(L) N_{c,O}^d$.
$T65.4$ : $\mathcal{A}(\mathcal{C}, O) = \overline{\bigcup_I \mathcal{A}_I}$ is AF.

## 66.9 Boundary

This shard does not construct the dynamical vacuum, prove a scaling limit, or derive Virasoro generators. Approximate Virasoro relations becoming exact in a continuum limit remain a programme conjecture. The categorical Markov trace and its naturality under spherical/unitary functors are not used here because the local source chain for canonical spherical structure is still open. The exact generator-level implementation of $\mathcal{BU}_0$, the endpoint-closing rule for charged words, and the double-triangle representation on $\mathcal{A}$ are implementation tasks, not completed results. “‘

# 67 Anyonic States and Variable-$N$ Fock Space

## 67.1 Status, Sources, and Dependencies

**Status: proposal shard from the adopted W4 design.** The finite carrier and Fibonacci arithmetic are local derivations from sourced fusion rules; no Julia invariant test has been added yet.

Dependencies: CA-02, CA-03, CA-06, CA-09, CA-62, CA-63, CA-64, CA-65, and CONVENTIONS.md (a), (b) (fixed with this block), (c), (f), (g), and (r).

## 67.2 The Finite Hard-Core Carrier

Fix the site object $O = \mathbf{1} \oplus X$ from CA-65. For $L$ open sites, define

$$H_L(\mathcal{C}, O) := \bigoplus_{c \in \mathrm{Irr}(\mathcal{C})} \mathrm{Hom}_{\mathcal{C}}(X_c, O^{\otimes L}).$$

The inner product on each Hom space is the categorical one: if $s, t$ have irreducible source $X_c$, then $s^\dagger t$ is a scalar multiple of $\mathrm{id}_{X_c}$. The direct sum over charges is orthogonal.

Expanding the maybe-object gives the variable-particle grading. For one species $X$,

$$O^{\otimes L} = (\mathbf{1} \oplus X)^{\otimes L} \cong \bigoplus_{S \subseteq \{1,\dots,L\}} X^{\otimes S},$$

where $S$ records the occupied sites and the tensor factors are kept in increasing site order. Hence

$$H_L(\mathcal{C}, O) \cong \bigoplus_{N=0}^{L} \bigoplus_{|S|=N} \bigoplus_{c \in \mathrm{Irr}(\mathcal{C})} \mathrm{Hom}_{\mathcal{C}}(X_c, X^{\otimes S}).$$

This is the sourced variable-$N$ target from `literature/SYNTHESIS.md`. The hard-core condition is built into the site object: each site contributes either $\mathbf{1}$ or one non-vacuum summand, never two particles at the same site.

For multiple species, take $X = \bigoplus_{a \neq \mathbf{1}} X_a$. The same formula refines by a species labelling of the occupied set $S$.

## 67.3 Action of the Word Algebra

The finite algebra

$$\mathcal{A}_L(\mathcal{C}, O) = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$$

acts on every charge sector by postcomposition:

$$T \cdot w := T \circ w, \qquad T \in \mathcal{A}_L, \quad w \in \mathrm{Hom}_{\mathcal{C}}(X_c, O^{\otimes L}).$$

Thus $H_L$ is the direct sum of the defining modules for the matrix blocks

$$\mathcal{A}_L \cong \bigoplus_c M_{m_c(L)}(\mathbb{C}), \qquad m_c(L) = \dim \mathrm{Hom}(X_c, O^{\otimes L}).$$

This is the finite carrier on which the kinematic quotient from CA-65 acts.

A charge-diagonal finite state can be written after choosing one minimal projection $e_c$ in each nonzero charge block. Let

$$p := \bigoplus_c e_c.$$

Using the ordinary finite matrix trace in the chosen block presentation, set

$$\omega_{L,p}(T) := \frac{\mathrm{Tr}(pTp)}{\mathrm{Tr}(p)}.$$

The corresponding GNS quotient is intended to recover one cyclic column from each charge block, hence a space of dimension $\sum_c m_c(L) = \dim H_L$. The choice of $e_c$ is not canonical when $m_c(L) > 1$; coherent choices in $L$ are an open point in this construction.

## 67.4 Birth and Death Operators

Let $a$ be a simple object with conjugate $\bar{a}$, and choose the raw cup

$$v_a : \mathbf{1} \to a \otimes \bar{a}.$$

The raw normalization is

$$v_a^\dagger v_a = d_a \, \mathrm{id}_{\mathbf{1}} .$$

At adjacent sites $j, j+1$, define the local birth map

$$b^a_{j,j+1} := (\iota_a \otimes \iota_{\bar{a}}) \circ v_a \circ (\pi_{\mathbf{1}} \otimes \pi_{\mathbf{1}}),$$

where $\iota_a : a \to O$ is the site summand inclusion and $\pi_{\mathbf{1}} : O \to \mathbf{1}$ is the vacuum summand projection. Its adjoint

$$d^a_{j,j+1} := (b^a_{j,j+1})^\dagger$$

is the death map. These maps change the occupation number by $N \leftrightarrow N \pm 2$, and annihilate configurations not matching the two-vacuum or $a, \bar{a}$ local pattern.

The default convention is raw cups with explicit $d_a$ bookkeeping:

$$(b^a_{j,j+1})^\dagger b^a_{j,j+1} = d_a \, P_{\mathbf{11}}.$$

The normalized cup $d_a^{-1/2} v_a$ would make the birth map a partial isometry on the two-vacuum corner, but that is not the default convention.

## 67.5 Hollands' Dense Chain as a Corner

Hollands' chain Hilbert space $V^L$ is built from sequences of intertwiners, and the corresponding operators are relative commutants. In the present tower it is recovered by two compressions. First take the fixed-$N = L$ corner that drops the vacuum summand at every site:

$$O = \mathbf{1} \oplus \tau \quad \rightsquigarrow \quad P_\tau O^{\otimes L} \cong \tau^{\otimes L}.$$

Then take the total-charge-$\mathbf{1}$ corner:

$$e_{\mathbf{1}} P_\tau H_L \cong \mathrm{Hom}_{\mathcal{C}}(\mathbf{1}, \tau^{\otimes L}).$$

Hollands' Jones projections are local elements in this compressed corner. The extra Q-system/subfactor data used in the dense-chain source are not supplied by a bare UFC input and remain deferred.

## 67.6 Fibonacci Test Targets

For Fibonacci, take

$$\mathrm{Irr}(\mathrm{Fib}) = \{\mathbf{1}, \tau\}, \qquad \tau \otimes \tau = \mathbf{1} \oplus \tau, \qquad O = \mathbf{1} \oplus \tau.$$

With $F_0 = 0$, $F_1 = 1$, the multiplicities are

$$(m_{\mathbf{1}}(L), m_\tau(L)) = (F_{2L-1}, F_{2L}) \qquad (L \geq 1).$$

Therefore

$$\dim H_L = m_{\mathbf{1}}(L) + m_\tau(L) = F_{2L+1},$$

and

$$\dim \mathcal{A}_L = m_{\mathbf{1}}(L)^2 + m_\tau(L)^2 = F_{4L-1}.$$

The first arithmetic targets are

$$\dim H_2 = 5, \quad \dim H_3 = 13, \quad \dim H_4 = 34,$$

and

$$\dim \mathcal{A}_2 = 13, \quad \dim \mathcal{A}_3 = 89, \quad \dim \mathcal{A}_4 = 610.$$

For the dense-chain corner $O = \tau$, the same recurrence reduces to CA-09's $(a_n, b_n)$ path-count sequence.

For the raw Fibonacci cup

$$v : \mathbf{1} \to \tau \otimes \tau,$$

Penneys gives

$$v^\dagger v = \phi \, \mathrm{id}_{\mathbf{1}}, \qquad \phi = \frac{1+\sqrt{5}}{2}.$$

Thus the local cup-cap invariant for the raw birth/death pair is $\phi$, not 1. The unitary Fibonacci recoupling matrix is

$$F_\tau^{\tau\tau\tau} = \begin{pmatrix} \phi^{-1} & \phi^{-1/2} \\ \phi^{-1/2} & -\phi^{-1} \end{pmatrix} = (F_\tau^{\tau\tau\tau})^\dagger .$$

For this nontrivial block the matrix is also involutory, but the adopted convention is still the unitary gauge so that matrix presentations use the ordinary adjoint.

## 67.7 Lamport Dependency Skeleton

$D66.1$ : $H_L(\mathcal{C}, O) = \bigoplus_c \mathrm{Hom}(X_c, O^{\otimes L})$.
$D66.2$ : $O = \mathbf{1} \oplus X \Rightarrow H_L \cong \bigoplus_{N,S,c} \mathrm{Hom}(X_c, X^{\otimes S})$.
$D66.3$ : $\mathcal{A}_L = \mathrm{End}(O^{\otimes L})$ acts by $T \cdot w = T \circ w$.
$D66.4$ : $p = \bigoplus_c e_c$, $\omega_{L,p}(T) = \mathrm{Tr}(pTp)/\mathrm{Tr}(p)$.
$D66.5$ : $b^a_{j,j+1} = (\iota_a \otimes \iota_{\bar{a}}) v_a (\pi_{\mathbf{1}} \otimes \pi_{\mathbf{1}})$, $d = b^\dagger$.
$T66.1$ : $(b^a)^\dagger b^a = d_a P_{\mathbf{11}}$ for raw cups.
$T66.2$ : $O = \mathbf{1} \oplus \tau$ : $(m_{\mathbf{1}}(L), m_\tau(L)) = (F_{2L-1}, F_{2L})$.
$T66.3$ : $\dim H_L = F_{2L+1}$, $\dim \mathcal{A}_L = F_{4L-1}$.
$T66.4$ : Hollands fixed-$N$, charge-$\mathbf{1}$ chain is a corner of $H_L$.

### 67.8 Boundary

This shard does not prove existence or uniqueness of a dynamical vacuum. The finite state $\omega_{L,p}$ is a useful charge-diagonal target, not a canonical vacuum. The categorical Markov trace is the expected natural "infinite-temperature" state only after the spherical/pivotal convention is locally sourced. The residual choice of $e_c$ for $m_c > 1$ is open.

The general statement "unitary UFC implies canonical spherical structure and positive quantum dimensions" is still a source gap for this block. The Fibonacci self-duality sign $\kappa_\tau = +1$ is also not locally sourced. Approximate Virasoro relations becoming exact in the scaling-limit GNS null space remains a programme conjecture, not a result of this shard.

# 68 Refinement-Map Requirements and Obstructions

## 68.1 Status, Sources, and Dependencies

**Status: source-local synthesis plus derived checklist.** Every requirement and failure mode below is anchored in a registered local source; the checklist assembly and the dense-chain diagnosis are local readings.

Dependencies: CA-14, CA-15, CA-62, CA-65, CA-66; CONVENTIONS.md (r).

## 68.2 Strict OAR Requirements

The sourced strict architecture asks of a refinement family $\alpha^N_{N'} : \mathcal{A}_N \to \mathcal{A}_{N'}$:

1. unital and completely positive (states pull back to states);
2. a *-morphism, injective, hence isometric;
3. the semigroup property $\alpha^{N'}_{N''} \circ \alpha^N_{N'} = \alpha^N_{N''}$;
4. bounded, scale-independent support growth (locality);
5. smearing: a lattice field maps to a smeared continuum field, not a point-like one.

On the state side, a scaling limit is a projectively consistent family $\omega^{(N)} = \omega^{(N')} \circ \alpha^N_{N'}$; on the dynamics side, the refined evolutions must form Cauchy sequences in the scaling-limit GNS representation, and the source stresses that success depends sensitively on the choice of scaling maps.

## 68.3 Fallback and Soft Relaxations

Two sourced fallbacks weaken the strict demands without abandoning the continuum target. First, when the *-morphism property or regularity fails (block-spin, point-localization), the algebra-level limit is lost but the vacuum can still be recovered through Wightman two-point functions; continuum local operators are then only indirectly accessible by reconstruction.

Second, the soft-inductive-system formalism replaces exact structure by asymptotic structure: connecting maps need only be contractions with asymptotic transitivity; operations converge iff they *asymptotically intertwine* the connecting maps,

$$\lim_{n \gg m} \left\| (\tilde{\jmath}_{nm} T_m - T_n \jmath_{nm}) x_m \right\| = 0,$$

and C*-structure survives under an asymptotic multiplicativity condition. For symmetry recovery the same source is explicit: Thompson-group implementers fail for nonzero central charge, and the successful strategy is Koo–Saleur approximation of conformal symmetries. **Consequence [derived]:** for a CFT target with $c \neq 0$, "compatible with the Virasoro action" can only mean *asymptotic* intertwining of normal-ordered Koo–Saleur generators on a suitable core — there is no exact-implementer formulation to aim for.

## 68.4 The No-Go Mechanism as a Checklist

The Jones no-go concerns the semicontinuous limit built from a single scale-invariant tree-local refinement element. The registered source identifies the root cause: the underlying dangling binary tree does not feel the circle topology, and the pulled-back smallest rotation is a long product of local elements controlled by a norm-contracting renormalisation (squaring) map, whose iterates vanish — forcing rotation matrix elements weakly to zero. Kliesch–Koenig extend this: discontinuity has

*full measure* among tree isometries, and they isolate an easily verified necessary condition for a continuous limit.

**Checklist [derived reading].** A refinement family risks the no-go mechanism when:

| | |
|---|---|
| NG1 | it is generated by a single scale-invariant tree-local element, decoupled from the circle; |
| NG2 | its induced squaring/renormalisation map is norm-contracting on the relevant subspace; |
| NG3 | it embeds point-like, without low-pass smearing; |
| NG4 | it operates in the large-loop-parameter regime where the contraction estimate bites; |
| NG5 | it demands exact unitary implementers of the discrete rotation/conformal action on a fixed limit space. |

Any candidate family must either avoid each mode or verify the Kliesch–Koenig necessary condition explicitly. The wavelet precedent shows what success looks like: translation-covariant smearing recovers continuous translations, while Lorentz/conformal covariance is separate further work.

## 68.5 Where Fixed Anyon Number Enters the Braiding RG

The braiding-RG refinement for anyonic chains is sourced as: attach new through-strings to a string-box diagram and braid their ends into position, *increasing the local density of $\sigma$-anyons*; the basis carries a $\sigma$-anyon at every site. **Diagnosis [derived].** Three fixed-$N$ commitments follow: (i) the braid needs a strand at every site it acts on, so the map is undefined on empty sites; (ii) the identification of local operators with Temperley–Lieb generators presupposes exactly one anyon per site; (iii) the refinement doubles the strand count deterministically. For the CA-65/CA-66 site object $O = \mathbf{1} \oplus X$, occupation is a quantum variable, so a refinement must be defined on the whole word space including empty sites — the braiding-RG step does not port. Conversely, dense chains have no refinable vacuum: every refined site must carry an anyon, so any fine-graining changes the physical content. The variable-$N$ tower is exactly what removes this obstruction, at the price of the new proof obligations recorded in CA-68.

## 68.6 Boundary

This shard proves no convergence theorem and constructs no refinement; it fixes the requirement set and the failure modes. The Kliesch–Koenig necessary condition is registered but its verification for any variable-$N$ family is future work. The Koo–Saleur compatibility target is stated precisely in CA-68; nothing here claims it holds.

# 69 Refinement Maps for the Variable-$N$ Anyonic Word Algebra

## 69.1 Status, Sources, and Dependencies

**Status: proposal plus local derivations.** The vacuum-insertion map is an exact finite-level construction in the CA-65/CA-66 tower. Its Virasoro compatibility is an open scaling problem, not a theorem.

Dependencies: CA-62, CA-63, CA-64, CA-65, CA-66, CA-67, and CONVENTIONS.md (a), (b), (c), and (r).

## 69.2 Vacuum Insertion

**Definition [choice].** Fix $O = \mathbf{1} \oplus X$. Let $\iota_{\mathbf{1}} : \mathbf{1} \to O$ be the empty-site summand inclusion. The canonical site-doubling refinement is

$$V_L : O^{\otimes L} \longrightarrow O^{\otimes 2L}, \qquad \text{old site } j \mapsto 2j-1, \quad \text{fine site } 2j \mapsto \mathbf{1}.$$

Equivalently, in each two-site fine cell the local factor is $\mathrm{id}_O \otimes \iota_{\mathbf{1}} : O \to O \otimes O$, with the fixed left-associated bracketing of CONVENTIONS.md (r).

**Lemma [derived].** $V_L$ is an isometry: $V_L^\dagger V_L = \mathrm{id}_{O^{\otimes L}}$. Each inserted factor contributes $\iota_{\mathbf{1}}^\dagger \iota_{\mathbf{1}} = \mathrm{id}_{\mathbf{1}}$, and tensor products of isometries are isometries.

**Lemma [derived].** The induced map on charge sectors

$$J_{L,c} : \mathrm{Hom}(X_c, O^{\otimes L}) \to \mathrm{Hom}(X_c, O^{\otimes 2L}), \qquad w \mapsto V_L \circ w,$$

is isometric and preserves the total charge $c$. It is strictly local: the cellwise factors act on disjoint fine two-site cells.

## 69.3 Algebra Map

**Observation [derived].** Let $P_L = V_L V_L^\dagger$. The map

$$\theta_L(T) = V_L T V_L^\dagger, \qquad T \in \mathcal{A}_L := \mathrm{End}_{\mathcal{C}}(O^{\otimes L}),$$

is a *-isomorphism from $\mathcal{A}_L$ onto the corner $P_L \mathcal{A}_{2L} P_L$. It is not a unital embedding into $\mathcal{A}_{2L}$, because $\theta_L(1) = P_L \neq 1$.

**Boundary [open].** Strict OAR, as locally sourced, wants unital injective *-morphisms of algebras. Thus $\theta_L$ is not yet a strict OAR scaling map on the full $\mathcal{A}_{2L}$. A unital extension needs complement data that this shard does not canonically supply; the honest alternatives are a unital completely positive completion inside a *soft* C*-system with asymptotic multiplicativity (CA-67), or the state/GNS corner formalism. *Amendment (2026-10-22, round 2):* a local unital extension for pointed $O$ with a relative half-braiding on $\langle O \rangle$ is derived in CA-90 (Theorem 90.1).

## 69.4 State Consistency

**Exact corner consistency [derived].** If a state $\omega_{2L}$ on $\mathcal{A}_{2L}$ satisfies $\omega_{2L}(P_L) = 1$, then $\omega_L := \omega_{2L} \circ \theta_L$ is a state on $\mathcal{A}_L$. For the CA-66 charge-diagonal states, exact consistency is obtained by choosing fine minimal projections $e_c^{(2L)} = V_L e_c^{(L)} V_L^\dagger$.

**Dynamical-vacuum condition [open].** A physical scaling vacuum should satisfy exact or asymptotic GNS consistency $J_L \Omega_L \approx \Omega_{2L}$. Bare vacuum insertion supplies a candidate reference subspace, not a dynamical-vacuum theorem.

## 69.5 Density Reading

**Observation [derived].** A word with $N$ occupied sites refines to a word with the same $N$ occupied sites on $2L$ fine sites: the site-filling fraction halves while the *physical* particle number at fixed physical volume is unchanged as the lattice spacing halves. Dense chains cannot do this: every refined site must carry the chain anyon, which is the CA-67 diagnosis of why the braiding-RG route does not port.

**Boundary [open].** This reading fits fixed-particle and finite physical-density sectors. Whether a *critical* vacuum sector is reachable depends on the dressed state and coupling flow below.

## 69.6 Virasoro Compatibility Target

**Target [open].** Let $:L^{(L)}_{k,\pm}(g_L):$ denote normal-ordered Koo–Saleur approximants at renormalized coupling $g_L$ (vacuum subtraction per CA-14). A refinement family $W_L$ should satisfy, on a low-energy GNS core and a growing long-wavelength window,

$$\sup_{|k|\le K_L} \big\| :L^{(2L)}_{k,\pm}(g_{2L}): W_L\psi - W_L :L^{(L)}_{k,\pm}(g_L):\psi \big\| \longrightarrow 0, \qquad K_L/L \to 0.$$

This is the CA-67 asymptotic-intertwining condition specialised to Koo–Saleur modes, the anyonic analogue of the sourced free-fermion convergence theorem. Per CA-67 there is no exact-implementer formulation to aim for when $c \neq 0$; asymptotic intertwining in a soft system is the target statement.

**Proposal [open].** Bare $V_L$ is not expected to satisfy the target for a fixed local Hamiltonian. Use a dressed family

$$W_L = U_L V_L,$$

with $U_L \in \mathcal{A}_{2L}$ a bounded-depth, charge-preserving local circuit, scale- and coupling-dependent. $W_L$ remains local, isometric, and charge-preserving; the role of $U_L$ is to create the short-range entanglement that bare empty-site insertion lacks, playing the part of the wavelet low-pass filter (and of the braiding disentangler in the dense setting) so that NG1–NG3 of CA-67 are avoided.

## 69.7 Dead Prior Art: the Deficit/Pair-Creation Precedent

The archived predecessor project (dead, not canonical; re-derive before use) validated two fine-graining isometries numerically for Fibonacci/Ising at small size under D4 wavelets: a *normalised product map* (spread each anyon through a one-particle wavelet isometry, column-normalise the diagonal Gram matrix) and an experimental *number-changing isometry* $V_0 + V_2$ in which the isometry deficit $I - V_0^\dagger V_0$ of the raw fixed-$N$ wavelet map is filled by a pair-creation channel into the $N{+}2$ sector. **Reading [derived].** The deficit phenomenon is structural evidence for the present programme: fixed-$N$ wavelet refinement fails isometry exactly where indefinite anyon number can repair it, using the CA-66 birth maps. The archived construction was never reconciled with any algebraic framework, never connected to Virasoro data, and its existence conditions were not proved; those are re-derivation targets for this block, not inherited results.

## 69.8 Fibonacci $H_2 \to H_4$ Test

**Data [derived].** For Fibonacci $O = \mathbf{1} \oplus \tau$, $(m_{\mathbf{1}}(2), m_\tau(2)) = (2, 3)$ and $(m_{\mathbf{1}}(4), m_\tau(4)) = (13, 21)$.

**Matrix target [derived].** In the occupied-subset path basis, $V_2$ sends $S \subseteq \{1,2\}$ to $2S - 1 \subseteq \{1,2,3,4\}$: the $\mathbf{1}$-block maps $\emptyset \mapsto \emptyset$, $\{1,2\} \mapsto \{1,3\}$; the $\tau$-block maps $\{1\} \mapsto \{1\}$, $\{2\} \mapsto \{3\}$, $\{1,2\} \mapsto \{1,3\}$. The charge-block matrices are $13\times 2$ and $21\times 3$ zero-one isometries. Julia invariants:

$V_c^\dagger V_c = I$ on both charge blocks, $P = VV^\dagger$ idempotent self-adjoint, occupation covariance $Vn_j = n_{2j-1}V$ and $n_{2j}V = 0$.

**Birth covariance [derived].** For the CA-66 adjacent birth $b^a_{j,j+1}$,

$$V_L b^a_{j,j+1} = \widetilde{b}^a_{2j-1,2j+1} V_L,$$

where $\widetilde{b}$ creates the pair on fine sites $2j-1, 2j+1$ and keeps fine site $2j$ in the vacuum. The naive adjacent-fine-site covariance is false for this insertion convention; a mutation test must pin this.

## 69.9 Comparison and Boundary

The braiding RG refines dense chains by increasing string density through braiding; the present family refines a vacuum-capable word algebra by adding empty sites. This shard does not claim to evade all rotation obstructions: it avoids the specific fixed TL semicontinuous Hilbert space of the no-go, and any future discrete-rotation action on this tower must be checked against the CA-67 checklist, including the registered Kliesch–Koenig necessary condition. The expected algebraic home of the Fibonacci word tower — a dilute Temperley–Lieb-type diagram algebra — and any Koo–Saleur construction for *dilute* loop models are open research targets; the dilute-algebra sources are registered but not yet used. No continuum limit, Virasoro representation, or category-fidelity theorem is proved here. The output is a local, isometric, charge-preserving refinement candidate, its dressing proposal, and its proof obligations.

# 70 The Dilute Temperley–Lieb Algebra and the Anyonic Word Algebra

## 70.1 Status, Sources, and Dependencies

**Status: source-local registration plus derived finite arithmetic.** The algebra data are registered from the local sources; the evaluation map is a proposal; the non-surjectivity obstruction and the small-$L$ dimension matches are checked local arithmetic.

Dependencies: CA-65, CA-66, CA-68; CONVENTIONS.md (b), (r).

## 70.2 The Dilute Algebra

The dilute Temperley–Lieb algebra $\mathrm{dTL}_n(\beta)$ is spanned by dilute $n$-diagrams: $n$ points on each vertical side, an even number of which may be *vacancies*, all remaining points joined pairwise by non-intersecting strings. The product is concatenation, with a factor $\beta$ per closed loop and zero whenever a floating string fails to close. A generating set is $\{a_i, a_i^t, b_i, b_i^t, e_j, x_j\}$, with $e_i + x_i = \mathrm{id}$ and the dense Temperley–Lieb generator recovered as $u_i = b_i^t b_i$. Two facts are load-bearing for this block:

*Parity.* The vacancy numbers on the two sides of a diagram always share parity, and the even/odd spans are two-sided ideals: $\mathrm{dTL}_n = \mathrm{edTL}_n \oplus \mathrm{odTL}_n$. Equivalently, a dilute diagram never changes the occupied-site number $N$ by an odd amount: strings pair endpoints, so occupied counts on the two sides are congruent mod 2.

*Dense corners.* For any subset $A$ of positions, the idempotent $\pi_A = \prod_{i\in A,\, j\notin A} x_j e_i$ satisfies $\pi_A\, \mathrm{dTL}_n\, \pi_A \cong \mathrm{TL}_{|A|}$: the fully occupied corner is the ordinary dense algebra.

The dimension is a Motzkin number:

$$\dim \mathrm{dTL}_n = \sum_{k=0}^{n} \binom{2n}{2k} \dim \mathrm{TL}_k = M_{2n},$$

so $\dim \mathrm{dTL}_n = 1, 2, 9, 51, 323$ for $n = 0, \ldots, 4$. For $q$ not a root of unity ($\beta = q + q^{-1}$) the algebra is semisimple.

## 70.3 The Evaluation Map

**Proposal.** Let $\mathcal{C}$ be a unitary fusion category with self-dual simple $X$ of quantum dimension $d$, and $O = \mathbf{1} \oplus X$ the CA-65 site object. Define

$$\rho_L : \mathrm{dTL}_L(d) \longrightarrow \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$$

by reading a vacancy as the $\mathbf{1}$-summand, an occupied point as the $X$-summand, a through-string as the identity on $X$ transported between its endpoint sites, and cups/caps as the raw coevaluation/evaluation of CONVENTIONS.md (r) (Fibonacci: $v^\dagger v = \varphi\, \mathrm{id}_{\mathbf{1}}$, so closed loops evaluate to $d = \varphi$, matching the diagrammatic $\beta$). The Frobenius–Schur flag of (r) is a recorded obligation before $\rho_L$ is promoted beyond proposal status for general self-dual $X$.

## 70.4 The Obstruction: No Surjection for Fibonacci

**Checked arithmetic.** For Fibonacci, CA-66 gives $\dim \mathrm{End}(O^{\otimes L}) = F_{4L-1}$ (checked in the Julia testset "fibonacci anyonic word algebra and vacuum insertion"), while $\dim \mathrm{dTL}_L = M_{2L}$:

| $L$ | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| $\dim \mathrm{dTL}_L = M_{2L}$ | 2 | 9 | 51 | 323 |
| $\dim \mathrm{End}(O^{\otimes L}) = F_{4L-1}$ | 2 | 13 | 89 | 610 |

Since $M_{2L} < F_{4L-1}$ for $L \geq 2$, no map from $\mathrm{dTL}_L$ can be surjective, and passing to any quotient (e.g. by a Jones–Wenzl-type ideal) only shrinks the image. The dense intuition $\mathrm{TL}_L(\varphi) \twoheadrightarrow \mathrm{End}(\tau^{\otimes L})$ does *not* extend to the dilute pair.

**Derived mechanism.** The failure is exactly the parity ideal structure: dilute diagrams preserve occupied-number parity, while Fibonacci has $N_{\tau\tau}^{\tau} = 1$, so the word algebra contains fusion morphisms changing $N$ by one (the trivalent vertex $\tau \to \tau \otimes \tau$). At $L = 2$ the missing $13 - 9 = 4$ dimensions are precisely the matrix units linking the $N = 1$ and $N = 2$ states of the charge-$\tau$ block.

**Derived dimension coincidence.** Splitting each CA-66 charge block by occupied-number parity, the parity-preserving subalgebra of $\mathrm{End}(O^{\otimes L})$ has dimension

$$L = 2: \ (2^2) + (2^2 + 1^2) = 9, \qquad L = 3: \ (4^2 + 1^2) + (5^2 + 3^2) = 51,$$

matching $M_{2L}$ exactly in both cases. **Conjecture (checked at the dimension level for $L \leq 3$; open beyond):** for Fibonacci, $\rho_L$ is an isomorphism onto the parity-even–preserving subalgebra of $\mathrm{End}(O^{\otimes L})$ for small $L$, with a nontrivial kernel expected only once dense corners reach the first negligible Jones–Wenzl projector ($\mathrm{TL}_4$ at $\beta = \varphi$), i.e. from $L = 4$. At $L = 4$ the counts 323 versus the parity-preserving dimension must be computed, not assumed; this is a CA-71-tier numerical obligation.

## 70.5 Consequences for the Programme

Two tiers separate cleanly. *Tier 1 (dilute TL proper):* Hamiltonians built from the dTL generators — occupancy projectors, dense exchange $u_i$, hopping $a_i + a_i^t$, pair creation/annihilation $b_i + b_i^t$ — act on the word algebra through $\rho_L$ and conserve $N$-parity. The sourced anyonic t-J models are of this shape, and the dilute Koo–Saleur programme (CA-70) lives here. *Tier 2 (beyond dTL):* the full word algebra needs the fusion vertex; densities using $N$-parity-violating terms leave the dilute algebra and require category-specific data. This is the precise algebraic sense in which the variable-$N$ word algebra is strictly richer than the dilute diagram algebra.

## 70.6 Boundary

No Hamiltonian, criticality, or Virasoro statement is made here. The evaluation map is a proposal pending the Frobenius–Schur convention; its injectivity on $\mathrm{dTL}_L(\varphi)$ for $L \leq 3$ is a dimension-level argument only and must be confirmed by an explicit rank computation (CA-71). The affine/periodic dilute algebra needed by a periodic Koo–Saleur setup is not defined in the registered sources and remains an acquisition/derivation target.

# 71 The Dilute Koo–Saleur Ansatz

## 71.1 Status, Sources, and Dependencies

**Status: proposal.** To our knowledge (2026-07-05 survey, recorded in the review bundle) no Koo–Saleur construction exists for any dilute loop model or the dilute Temperley–Lieb algebra; this shard fixes the candidate generators and their proof obligations. Nothing here is a convergence claim.

Dependencies: CA-14, CA-42, CA-64, CA-69; CONVENTIONS.md (r).

## 71.2 What the Dense Template Requires

The sourced dense-TL recipe takes: (i) a critical Hamiltonian $H_N = \sum_j h_j$ with density built from algebra generators, normalized so the sound velocity is one; (ii) the ground-state energy density $e_\infty$ subtracted termwise; (iii) a momentum density from the nearest-neighbour commutator of Hamiltonian densities; (iv) periodic boundary conditions; and (v) the $c/24$ zero-mode shift. The generators are Fourier modes of the subtracted density plus the current correction. The known frontier is recorded in the same source: even in the dense case, the limit of commutators agrees with the commutator of limits only up to a modified central term, and parts of that statement remain conjectural.

## 71.3 The Dilute Ansatz

**Proposal.** Let $h_j$ be a self-adjoint, translation-invariant two-site density in the dilute algebra: a real linear combination of occupancy projectors, the dense exchange $u_j = b_j^t b_j$, hopping $a_j + a_j^t$, and pair creation/annihilation $b_j + b_j^t$ (all dilute diagrams, hence occupied-parity-preserving by CA-69). On the periodic chain — the algebraic setting being the registered periodic dilute Temperley–Lieb algebra — define

$$E_n^{(N)} = \frac{N}{2\pi v} \sum_{j=1}^{N} e^{inj2\pi/N} \, (h_j - e_\infty), \qquad p_j = i[h_j, h_{j+1}],$$

$$P_n^{(N)} = \frac{N}{2\pi} \sum_{j=1}^{N} e^{in(j+\frac{1}{2})2\pi/N} \, p_j,$$

$$L_n^{(N)} = \tfrac{1}{2}\big(E_n^{(N)} + \tfrac{1}{v} P_n^{(N)}\big) + \tfrac{c}{24}\,\delta_{n,0}, \qquad \bar{L}_{-n}^{(N)} = \tfrac{1}{2}\big(E_n^{(N)} - \tfrac{1}{v} P_n^{(N)}\big) + \tfrac{c}{24}\,\delta_{n,0},$$

where $v$ and $e_\infty$ are the model's sound velocity and ground-state energy density, and $c$ its central charge. This is the direct dilute transcription of the sourced dense formula; every ingredient except the algebra membership of $h_j$ is inherited. The candidate modes live in the even (parity-preserving) dilute ideal, consistently with CA-69.

## 71.4 Two Sourced Critical Anchors

*Integrable anchor (dilute O(n)/Izergin–Korepin).* The registered sources give exact branch central charges, $c = 1 - \frac{3\phi^2}{\pi(\pi-2\lambda)}$ (branches 1, 2) and $c = \frac{3}{2} - \frac{3\phi^2}{2\pi\lambda}$ (branches 3, 4), and the periodic dilute algebra carries the loop transfer matrices. The explicit quantum-chain Hamiltonians (the Hamiltonian limits of these transfer matrices, with their $v$ and $e_\infty$) are in a journal-only source flagged above; until it is registered, the integrable anchor supports regime selection but not a shard-ready density. A recorded warning: at least one regime of the $A_2^{(2)}$ model has a *non-compact*

continuum limit (registered source VernierJacobsenSaleur2014), so regime selection is part of the input data, not a detail.

*Anyonic anchor (itinerant t-J).* The locally sourced anyonic t-J models are dilute in exactly the CA-69 sense: anyon hopping plus a dense projector exchange term, with hard-core occupancy — an $h_j$ in the image of the dilute generators under the CA-69 evaluation map, acting on the CA-66 Fibonacci sectors. Their central charges are numerically identified ($c = 7/10$ or $4/5$ for Fibonacci, $1/2$ for Ising; entanglement gives $1 + c$ with the charge sector). This anchor is shard-ready on the input side but its criticality is numerical, not Bethe-exact.

## 71.5 Proof Obligations

1. $e_\infty,\ v,\ c$ must be sourced (integrable anchor) or estimated with recorded finite-size protocols (anyonic anchor);
2. the central-term/limit gap is inherited from the dense case and remains conjectural there; no dilute claim may be stronger;
3. periodic dilute algebra: the CA-69 evaluation map must be extended to the periodic setting before $L_n^{(N)}$ acts on word sectors;
4. gaplessness alone is insufficient: a dilute-looking density can be gapless with dynamical exponent $z \neq 1$ (no local source yet; registered as an external caution for CA-71 acceptance tests);
5. diagnostics run through CA-64: normalized GNS residuals $\nu_N$ of the dilute Witt relations on long-wavelength windows.

## 71.6 Validation Plan (CA-71 Tier)

Exact diagonalization in the Fibonacci $O = \mathbf{1} \oplus \tau$ charge sectors (bases already implemented and tested for CA-65–CA-68): build the dense projector from the sourced unitary $F$-matrix, hopping across vacancies, and raw-cup birth/death; estimate $e_\infty$, $v$, and $c$ from finite-size spectra; construct $L_n^{(N)}$ and test low-energy commutator closure and the CA-64 residuals. Acceptance: dilute relations hold exactly; sourced or estimated $(e_\infty, v, c)$ reproduce the anchor values; current modes do not collapse; residuals decrease on the named window. Failure witnesses: current collapse, wrong central charge, residuals bounded away from zero, or non-conformal ($z \neq 1$) finite-size scaling.

## 71.7 Boundary

No convergence theorem, no continuum Virasoro representation, and no claim that the ansatz closes even at the level of the dense case's conjectures. The integrable anchor's Hamiltonian is a registered acquisition gap. The anyonic anchor's criticality is numerical input, not a theorem. The extension of the CA-69 evaluation map to the periodic dilute algebra is future work.

# 72 The Refinement Placement Category and Exact Composition

## 72.1 Status, Sources, and Dependencies

**Status: local derivations plus sourced structure theory.** The placement definitions, isometry, charge preservation, and the exact composition theorem are finite-level derivations checked in Julia. The forest-functor identification and the Ore dichotomy are sourced. No dressing, no state selection, no dynamics, and no continuum statement appears here.

Dependencies: CA-65, CA-66, CA-67, CA-68; CONVENTIONS (a), (b), (c), (r), (s), (t). This shard executes item W1.1 of `reviews/2026-07-06_pipeline_consistency_scoping/PLAN.md`.

## 72.2 Placement Injections and Bare Refinements

**Definition [choice].** For $k \le l$, a *placement* is an order-preserving injection

$$\varphi : \{1, \dots, k\} \hookrightarrow \{1, \dots, l\},$$

recorded as its strictly increasing slot list $(\varphi(1), \dots, \varphi(k))$. Placements compose as functions; the identity placement is $k = l$, $\varphi = \mathrm{id}$. We write Pl for the category with objects $\mathbb{Z}_{>0}$ and morphisms the placements.

**Definition [choice].** Fix the site object $O = \mathbf{1} \oplus X$ of CONVENTIONS (r) and let $\iota_{\mathbf{1}} : \mathbf{1} \to O$ be the empty-site summand inclusion. The *bare refinement* attached to a placement $\varphi : [k] \to [l]$ is

$$V_\varphi : O^{\otimes k} \longrightarrow O^{\otimes l}, \qquad \text{coarse site } i \mapsto \text{fine slot } \varphi(i), \quad \text{slots } s \notin \operatorname{im}\varphi \mapsto \iota_{\mathbf{1}}(\mathbf{1}),$$

with the left-associated bracketing of CONVENTIONS (c)/(r). CA-68's dyadic vacuum insertion is the placement $\varphi_{\mathrm{dyad}}(j) = 2j - 1$.

**Lemma [derived].** $V_\varphi$ is an isometry, $V_\varphi^\dagger V_\varphi = \mathrm{id}_{O^{\otimes k}}$: every inserted factor contributes $\iota_{\mathbf{1}}^\dagger \iota_{\mathbf{1}} = \mathrm{id}_{\mathbf{1}}$, and tensor products of isometries are isometries (the CA-68 argument, verbatim, for an arbitrary slot pattern).

**Lemma [derived].** On charge sectors, $w \mapsto V_\varphi \circ w$ maps $\mathrm{Hom}(X_c, O^{\otimes k})$ isometrically into $\mathrm{Hom}(X_c, O^{\otimes l})$ for every charge $c$; the total charge is preserved because composing with $V_\varphi$ does not touch the source object $X_c$.

**Observation [derived].** In the occupied-set/fusion-path basis of CONVENTIONS (r) and `src/AnyonicWordAlgebra.jl`, $V_\varphi$ acts by relabelling: the occupied set $S$ maps to $\varphi(S)$, and the cumulative-charge path repeats the running charge at every vacuum slot. This is exact, with no F-move corrections, because every associator moved past an inserted factor carries the unit label, and unit-label F-symbols are identities in the unitary gauge (CONVENTIONS (b)). Consequently the matrix of $V_\varphi$ on each charge block is a zero-one matrix whose columns are distinct standard basis vectors.

## 72.3 Exact Composition

**Theorem [derived].** For composable placements $\varphi : [k] \to [l]$ and $\psi : [l] \to [m]$,

$$V_\psi \circ V_\varphi = V_{\psi \circ \varphi}.$$

*Proof sketch.* Both sides place coarse site $i$ at fine slot $\psi(\varphi(i))$ and the vacuum summand everywhere else: a slot of $[m]$ is outside $\operatorname{im}(\psi \circ \varphi)$ iff it is outside $\operatorname{im}\psi$ or is the $\psi$-image of a slot outside $\operatorname{im}\varphi$, and in either case it carries $\iota_{\mathbf{1}}(\mathbf{1})$. All bracketing moves involve unit labels only, so the two zero-one matrices agree column by column. □

**Corollary [derived].** The map $\varphi \mapsto V_\varphi$ is a functor Pl $\to$ Hilb (isometries), and $\theta_\varphi(T) = V_\varphi T V_\varphi^\dagger$ satisfies $\theta_\psi \circ \theta_\varphi = \theta_{\psi\circ\varphi}$ exactly: each $\theta_\varphi$ is a $*$-isomorphism of $\mathcal{A}_k$ onto the corner $P_\varphi \mathcal{A}_l P_\varphi$, $P_\varphi = V_\varphi V_\varphi^\dagger$ (the CA-68 corner statement, now closed under composition).

**Corollary [derived].** The dyadic tower composes on the nose: $V^{(2L\to 4L)}_{\varphi_{\mathrm{dyad}}} \circ V^{(L\to 2L)}_{\varphi_{\mathrm{dyad}}} = V^{(L\to 4L)}_{\varphi}$ with $\varphi(j) = 4j - 3$. This settles the semigroup question left open in the 2026-07-06 consistency map (seam S2 at finite level): the reindexing is the composite placement itself.

## 72.4 Cellwise Refinement and Cell-Shift Compatibility

**Definition [choice].** For a chain of $M$ cells of $k$ sites, the uniform cellwise refinement applies one fixed cell placement $\varphi : [k] \to [l]$ in every cell; the result is the whole-chain placement $\Phi_M(\varphi) : [Mk] \to [Ml]$, cell $m$ occupying fine slots $(m-1)l + 1, \ldots, ml$.

**Lemma [derived].** On its open-chain domain, $V_{\Phi_M(\varphi)}$ intertwines whole-cell shifts: shifting the coarse chain by one cell ($k$ sites) corresponds to shifting the fine chain by one cell ($l$ sites). No exact fine counterpart of a *single-site* coarse shift exists when $l/k \notin \mathbb{Z}$, and none is used: cross-scale position-weighted comparisons go through the cell-shift subgroup only (CONVENTIONS (t)).

## 72.5 The Forest-Functor Reading

**Sourced identification.** The registered Brothier–Stottmeister construction assigns to any covariant functor $\Phi$ from the category of binary forests to a concrete category the inductive limit $X_\Phi = \varinjlim_t X_t$ over the directed set of trees, together with the *Jones action* of Thompson's group $F$ on $X_\Phi$; a *tensor* functor is characterized by the single seed $R = \Phi(Y) \in \mathrm{Hom}(\Phi(1), \Phi(1)\otimes\Phi(1))$, any pair $(A, R)$ conversely defines one, and for tensor functors the $F$-action always extends to Thompson's $V$; the targets of record are Hilbert spaces with isometries and C*-algebras with injective $*$-morphisms.

**Observation [derived].** The vacuum caret

$$R = \mathrm{id}_O \otimes \iota_{\mathbf{1}} : O \longrightarrow O \otimes O$$

is such a seed, and the forest generator $f_{j,n}$ maps to $\mathrm{id}^{\otimes j-1} \otimes R \otimes \mathrm{id}^{\otimes n-j}$ = "insert one vacuum slot after site $j$". Since every placement is a composite of single vacuum insertions, the placement functor of this shard is exactly the tensor functor generated by the vacuum caret; distinct forests mapping to the same placement record insertion histories, so Pl is the image of the forest category under this functor. CA-68's refinement tower therefore carries, for free and at the level of sourced theorems, an inductive-limit space and a Thompson-$F$ (indeed $V$) action. **Boundary [open]:** whether that action is continuous or physically relevant is precisely the no-go territory of CA-67 (NG1–NG5, Kliesch–Koenig); nothing here claims continuity, and the Kliesch–Koenig applicability question for this charge-graded carrier is deliberately deferred to the W2.4 audit.

## 72.6 The Ore Dichotomy and Fallback Homes

**Sourced dichotomy.** Fraction groups exist for a category of refinement diagrams iff it is left-cancellative and satisfies Ore's property (morphisms with a common target admit a common extension). The registered forest-skein source proves: the free forest-skein category over a colour set $S$ is always left-cancellative; one colour recovers Thompson's $F$; with more than two colours Ore's property *always fails*, and skein relations are the mechanism that can restore it.

**Consequence [derived reading].** Homogeneous refinement schedules (one caret type, e.g. pure vacuum insertion) retain a Thompson-like fraction-group symmetry. Mixed-radix or multi-caret

schedules — the kind that Wave 2's content-creating dressings and filling-targeted $k \to l$ schedules will need — generically lose the group and retain only the refinement *groupoid*. The two sourced fallback homes are: divisibility-directed inductive systems over scales ordered by divisibility (the AF/hyperfinite pattern over matrix building blocks), and soft inductive systems over an arbitrary directed set, where exact transitivity is relaxed to asymptotic transitivity — the recorded destination for dressed families $W = UV$ (CA-67, CA-68).

### 72.7 Julia Invariants

`src/RefinementPlacements.jl` implements placements (validated strictly increasing slot lists), composition, the charge-block matrices of $V_\varphi$ in the CA-66 basis, and uniform cellwise placements. The testset "refinement placement category (CA-71)" pins: the bridge $V_{\varphi_{\mathrm{dyad}}} =$ `vacuum_insertion_matrix` at $L = 1, 2, 3$; isometry on both charge blocks for all placements with $k \le 3$, $l \le 5$; exact functoriality $V_\psi V_\varphi = V_{\psi \circ \varphi}$ over an exhaustive composable sweep and the dyadic $2 \to 4 \to 8$ chain against the direct $j \mapsto 4j - 3$ placement; occupation covariance $V_\varphi n_i = n_{\varphi(i)} V_\varphi$ and $n_s V_\varphi = 0$ for $s \notin \operatorname{im} \varphi$; and the Fibonacci multiplicity table $(m_{\mathbf{1}}, m_\tau)(L) = (F_{2L-1}, F_{2L})$ at the sizes used. Each invariant is mutation-proved per Rule 6 (composition off-by-one, wrong path repetition, unmapped occupied set), with the mutation-to-failing-test record kept at the head of the testset.

### 72.8 Boundary

This shard defines no dressing $U$, selects no state, and makes no dynamical or continuum claim; the asymptotic cocycle condition for dressed families and the filling-flow constructions are Wave-2 targets (W2.1). The cell-shift lemma is an open-chain statement; periodic chains await the CONVENTIONS (r) periodic entry. The Thompson action inherited from the forest functor is an algebraic fact about the finite tower and its inductive limit; its continuity is not claimed and is exactly what the Jones and Kliesch–Koenig no-gos constrain (CA-67). Nothing here defines Koo–Saleur modes; the categorical residual set and the scale ledger of CONVENTIONS (s) are consumed by the following shards (W1.2, W1.3).

# 73 GNS Descent and the Corner Calculus of Placement Refinements

## 73.1 Status, Sources, and Dependencies

**Status: local derivations with checked finite shadows, plus two recorded interface contracts.** Every lemma below is an elementary finite-dimensional derivation, checked in Julia. The two items that are *not* theorems here — the residual-compatibility contract and the BU-level square — are stated as proposals with their consuming shards named. This shard settles the residual-free part of seam S1 of the 2026-07-06 consistency map; the residual part waits for the categorical residual set (W1.3).

Dependencies: CA-63, CA-65, CA-66, CA-67, CA-68, CA-71; CONVENTIONS (r), (s), (t). Executes item W1.2 of `reviews/2026-07-06_pipeline_consistency_scoping/PLAN.md`.

## 73.2 Setting

Fix a placement $\varphi : [k] \to [l]$ with bare refinement $V_\varphi$, corner projection $P_\varphi = V_\varphi V_\varphi^\dagger$, and corner morphism $\theta_\varphi(T) = V_\varphi T V_\varphi^\dagger$ (CA-71). States on $\mathcal{A}_l = \bigoplus_c M_{m_c(l)}(\mathbb{C})$ are block density pairs $\omega_l = (w_c, \rho_c)$, $\omega_l(T) = \sum_c w_c \operatorname{tr}(\rho_c T_c)$ (CA-66). Throughout, $\Omega_\omega$ denotes the GNS cyclic vector and $[T]_\omega$ the GNS class of $T$.

## 73.3 The Corner Calculus

**Lemma 72.1 [derived].** $\theta_\varphi$ is an injective $*$-morphism onto the corner $P_\varphi \mathcal{A}_l P_\varphi$, and

$$\theta_\varphi(S)^\dagger \theta_\varphi(T) = \theta_\varphi(S^\dagger T), \qquad \theta_\varphi(S)\,(1 - P_\varphi) = 0 = (1 - P_\varphi)\,\theta_\varphi(S).$$

*Proof.* $V^\dagger V = 1$ gives $VS^\dagger V^\dagger V T V^\dagger = VS^\dagger T V^\dagger$; and $VSV^\dagger(1 - VV^\dagger) = VS(V^\dagger - V^\dagger) = 0$. □

**Lemma 72.2 [derived] (pullback functionals).** $\omega_l \circ \theta_\varphi$ is a positive linear functional of total mass $\omega_l(P_\varphi)$; it is a state iff $\omega_l(P_\varphi) = 1$ (the CA-68 exact corner case). For $\omega_l(P_\varphi) > 0$ the *normalized pullback*

$$\omega_k := \frac{\omega_l \circ \theta_\varphi}{\omega_l(P_\varphi)}$$

is a state on $\mathcal{A}_k$. **Boundary:** translation invariance of $\omega_l$ does *not* imply translation invariance of $\omega_k$ in any sense finer than the cell-shift subgroup of CONVENTIONS (t); no invariance is claimed here.

## 73.4 The GNS Descent Lemma

**Theorem 72.3 [derived].** Let $\omega_l(P_\varphi) > 0$ and let $\omega_k$ be the normalized pullback. Then

$$J\,[T]_{\omega_k} := \omega_l(P_\varphi)^{-1/2}\,[\theta_\varphi(T)]_{\omega_l}$$

is a well-defined isometry $\mathcal{H}_{\omega_k} \to \mathcal{H}_{\omega_l}$ satisfying, for all $S, T \in \mathcal{A}_k$,

$$J\,\pi_{\omega_k}(S) = \pi_{\omega_l}(\theta_\varphi(S))\,J, \qquad \left\| J\Omega_{\omega_k} - \Omega_{\omega_l} \right\|^2 = 2\Big(1 - \sqrt{\omega_l(P_\varphi)}\Big).$$

In particular $J\Omega_{\omega_k} = \Omega_{\omega_l}$ iff $\omega_l(P_\varphi) = 1$.

*Proof.* Isometry and well-definedness: $\langle J[S], J[T]\rangle = \omega_l(P)^{-1}\omega_l(\theta(S)^\dagger\theta(T)) = \omega_l(P)^{-1}\omega_l(\theta(S^\dagger T)) = \omega_k(S^\dagger T)$ by Lemma 72.1, so $\omega_k$-null classes map to $\omega_l$-null classes. Intertwining: $J[ST] = \omega_l(P)^{-1/2}[\theta(S)\theta(T)] =$

$\pi_{\omega_l}(\theta(S))J[T]$. Vacuum defect: $\langle J\Omega_k, \Omega_l\rangle = \omega_l(P)^{-1/2}\,\omega_l(P^\dagger 1) = \sqrt{\omega_l(P)}$, hence $\|J\Omega_k - \Omega_l\|^2 = 1 + 1 - 2\sqrt{\omega_l(P)}$. □

**Reading.** Every placement refinement descends to the GNS level *unconditionally*: multiplicativity survives the corner because $V^\dagger V = 1$. What is conditional is only whether the refinement is *pointed* (vacuum-preserving), and that is measured by the single scalar $1 - \omega_l(P_\varphi)$ — the same scalar that controls the completion defect below. This closes the residual-free part of seam S1.

### 73.5 Residual Descent: the Contract for the Categorical Residual Set

**Lemma 72.4 [derived].** In the setting of Theorem 72.3, for any $R' \in \mathcal{A}_k$ and $T \in \mathcal{A}_k$,

$$\big\|\pi_{\omega_k}(R')\,[T]_{\omega_k}\big\| = \big\|\pi_{\omega_l}(\theta_\varphi(R'))\,J[T]_{\omega_k}\big\|.$$

Hence $\pi_{\omega_k}(R') = 0$ iff $\pi_{\omega_l}(\theta_\varphi(R'))$ vanishes on $\operatorname{ran} J$; in particular $\pi_{\omega_l}(\theta_\varphi(R')) = 0 \Rightarrow \pi_{\omega_k}(R') = 0$: coarse relations are inherited from fine ones through $\theta_\varphi$.

**Contract [proposal, consumed by W1.3].** A scale family of residual sets $\{\mathcal{R}_k\}$ is *descent-compatible* for $(\omega_l, \varphi)$ when $\pi_{\omega_l}(\theta_\varphi(R'))\,J = 0$ for every $R' \in \mathcal{R}_k$ — e.g. when $\theta_\varphi(\mathcal{R}_k)$ lies in the $\pi_{\omega_l}$-null structure generated by $\mathcal{R}_l$. Once W1.3 defines the categorical residual densities, this is the equation their scale coherence must satisfy; nothing about it is claimed here.

**Lemma 72.5 [derived] (independent families).** If instead $\omega_k$ is *given* (a projective family, not the pullback), then $J[T]_{\omega_k} := [\theta_\varphi(T)]_{\omega_l}$ is well-defined and bounded iff the Gram domination $\omega_l(\theta(T)^\dagger\theta(T)) \le C\,\omega_k(T^\dagger T)$ holds for some $C$; the pullback case is $C = \omega_l(P_\varphi)$ with equality. This is the finite Gram condition to check when comparing two independently constructed state families.

### 73.6 The Unital CP Completion and Its Exact Defect

Strict OAR wants unital injective $*$-morphisms; $\theta_\varphi$ is not unital. For a state $\chi$ on $\mathcal{A}_k$ define the *$\chi$-completion*

$$\Phi_\chi(T) := \theta_\varphi(T) + \chi(T)\,(1 - P_\varphi).$$

**Lemma 72.6 [derived].** $\Phi_\chi$ is unital and completely positive (sum of a $*$-morphism and the CP map $T \mapsto \chi(T)(1 - P_\varphi)$), with state pullback the convex mixture

$$\omega_l \circ \Phi_\chi = \omega_l(P_\varphi)\,\omega_k + \big(1 - \omega_l(P_\varphi)\big)\,\chi.$$

**Theorem 72.7 [derived] (exact multiplicativity defect).** For all $S, T \in \mathcal{A}_k$,

$$\Phi_\chi(ST) - \Phi_\chi(S)\Phi_\chi(T) = \big(\chi(ST) - \chi(S)\chi(T)\big)\,(1 - P_\varphi),$$

and consequently

$$\big\|\Phi_\chi(ST) - \Phi_\chi(S)\Phi_\chi(T)\big\|_{\omega_l} = \big|\chi(ST) - \chi(S)\chi(T)\big|\,\sqrt{\omega_l(1 - P_\varphi)}.$$

*Proof.* Expand $\Phi_\chi(S)\Phi_\chi(T)$; the cross terms vanish by Lemma 72.1, leaving $\theta(ST) + \chi(S)\chi(T)(1 - P)$; subtract from $\Phi_\chi(ST) = \theta(ST) + \chi(ST)(1 - P)$. The seminorm follows from $(1 - P)^\dagger(1 - P) = 1 - P$. □

**Corollary 72.8 [derived reading].** On the matrix-block algebras $\mathcal{A}_k$ no state is multiplicative on all of $\mathcal{A}_k$ (unless a block is one-dimensional and $\chi$ is concentrated there), so the completion family $\{\Phi_\chi\}$ satisfies the soft-C*-system asymptotic homomorphism property (the sourced eq. (29) condition) *along a scale sequence precisely when* $\omega_l(1 - P_\varphi) \to 0$ in the relevant GNS seminorms. The

corner scalar of Theorem 72.3 is thus also the exact modulus of softness of the unital completion: strict corner ($\omega(P) = 1$) gives an exactly pointed descent, and $\omega(P) \to 1$ gives an asymptotically multiplicative unital CP system. Whether a physically relevant (critical) fine-state family can achieve $\omega_l(P) \to 1$ for dressed placements is exactly the W2.1 dressed-exact-corner question; nothing is claimed here.

### 73.7 The BU-Level Square

**Proposal [contract, blocked on W1.5].** On the free word layer $\mathcal{BU}_0$ (CA-65), a placement acts on generators by slot relabelling, $\widehat{R}_\varphi(v_{a,j}) := v_{a,\varphi(j)}$, extended freely. The required compatibility square is

$$\pi_l \circ \widehat{R}_\varphi = \theta_\varphi \circ \pi_k \quad \text{on local words,} \qquad \widehat{R}_\varphi(J_{\mathrm{fus},k}) \subseteq J_{\mathrm{fus},l},$$

so that refinement is defined already at the presentation level and descends through the kinematic quotient. Its verification needs the endpoint-closing rule (the CA-65 generator-to-block bridge, scheduled as W1.5); until then the square is an interface contract, not a result.

### 73.8 Julia Invariants

`src/GnsCornerCalculus.jl` implements block states, corner weights $\omega_l(P_\varphi)$, normalized pullbacks, explicit finite GNS data (Gram matrices, left actions, cyclic vectors), the descent intertwiner $J$, and the completion $\Phi_\chi$. The testset "gns descent and corner calculus (CA-72)" pins: the corner-calculus identities of Lemma 72.1 exactly on integer matrices; the CA-68 corner regression ($\omega_l(P_\varphi) = 1$ and pullback recovery for fine states built on the corner — a regression by construction, not a probe); isometry $J^\dagger G_l J = G_k$, the intertwining relation, and the vacuum-defect scalar $2(1 - \sqrt{\omega_l(P_\varphi)})$ for faithful fine states; the residual-descent norm identity of Lemma 72.4; the exact defect identity of Theorem 72.7 with its GNS-norm equality and the convex-mixture pullback of Lemma 72.6; and one exact rational pin, $\omega_l(P_\varphi) = 5/34$ for the carrier-trace state $\mathrm{Tr}_{H_4} / \dim H_4$ under the dyadic placement (corner ranks $2 + 3$ against $\dim H_4 = 13 + 21 = 34$; the implementing agent caught and corrected the orchestrator's wrong draft pin here — recorded in the worklog). Each invariant is mutation-proved per Rule 6 (dropped $J$-normalization, $P$ in place of $1 - P$ in the completion, $V^\dagger V$ in place of $VV^\dagger$), with the record at the head of the testset.

### 73.9 Boundary

This shard proves nothing about dynamics, translation invariance, criticality, dressings, or continuum limits; it fixes the exact finite-$L$ calculus through which any of those must pass. The completion state $\chi$ is a non-canonical choice, and no canonical choice is proposed. The residual-descent contract and the BU-level square are named interfaces for W1.3 and W1.5. Whether $\omega_l(P) \to 1$ is achievable on physical state families (and with which dressed placements) is the W2.1 question; the Kliesch–Koenig applicability audit remains W2.4. Seam S1 is closed here only in its residual-free part, as the plan demands.

**Erratum / qualification (2026-10-22).** Corollary 72.8's reading is not supported as an operator-norm claim. Theorem 72.7 gives the completion defect $(\chi(ST) - \chi(S)\chi(T))(1-P)$, whose operator norm is $|\chi(ST) - \chi(S)\chi(T)|$ whenever $P \neq 1$, whatever $\omega_l(1-P)$; the sourced eq. (29) is an operator-norm condition. Smallness of $\omega_l(1-P)$ controls only the GNS seminorm, so $\omega(P) \to 1$ does not give a soft C*-system. The phrase "exact modulus of softness" should be read as "GNS-seminorm defect". Found independently in D1 (T5b), D3 (§1) and D6 (N1); stated as Correction 76.A of CA-76. Theorem 72.7 itself is unaffected.

# 74 The Categorical Residual Set on the Anyonic Word Tower

## 74.1 Status, Sources, and Dependencies

**Status: derived port plus one new lemma and one definition.** The residual calculus below is the CA-38/CA-39/CA-42 qubit framework transported to the anyonic word tower; the transport lemma and the coherent residual family are new local derivations. The densities are the CA-70 Tier-1 proposal. Everything is stated on the *open* chain per the Wave-1 commitment of the 2026-07-06 plan; periodic mode calculus is gated behind decision D1. Nothing here claims that residual annihilation implies symmetry (CA-44/CA-64 semantics are preserved verbatim).

Dependencies: CA-38, CA-39, CA-42, CA-43, CA-63, CA-64, CA-69, CA-70, CA-71, CA-72; CONVENTIONS (n), (r), (s), (t). Executes item W1.3 of `reviews/2026-07-06_pipeline_consistency_scoping/`

## 74.2 Setting

Fix $(\mathcal{C}, O)$ with $O = \mathbf{1} \oplus X$ per CONVENTIONS (r), the open-chain tower $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$, and a self-adjoint two-site density $h \in \mathcal{A}_2$ placed at bonds by the shift, $h_j \in \mathcal{A}_L$ for $1 \le j \le L-1$, so that $[h_j, h_k] = 0$ for $|j-k| \ge 2$ (disjoint supports). The Tier-1 dilute density of CA-70 is the default instance: a real combination of occupancy projectors, the dense exchange $u_j = b_j^t b_j$, hopping $a_j + a_j^t$, and pair terms $b_j + b_j^t$, all parity-even by CA-69. The qubit block is the $\mathcal{C} = \mathrm{Hilb}$, $O = \mathbb{C}^2$ degenerate case, which is what makes the port below a specialization-in-reverse rather than an analogy.

## 74.3 The Ported Local Residual Calculus

**Definition [port].** For a local density $u$ supported on $m$ sites at position $j$, the coboundary is $(Du)_j = u_j - u_{j+1}$. The local residual densities are, verbatim from CONVENTIONS (n) but now as elements of $\mathcal{A}_L$:

$$p_j = i[h_j, h_{j+1}], \qquad A_j = i[h_j + h_{j+1},\, p_j], \qquad B_j = i[p_j, h_{j+1}] + 2i[p_j, h_{j+2}],$$

with the conservation and boost residual equations

$$R_j^{\mathrm{cons}}(h;u) = A_j - (Du)_j, \qquad R_j^{\mathrm{boost}}(h;u,w,\lambda,e) = B_j - u_j - \lambda\,\bar{h}_j - (Dw)_j, \qquad \bar{h} = h - e\,\mathbf{1},$$

where $\lambda$ is the unconstrained multiplier of CONVENTIONS (s) (not a squared velocity) and the scalar parts of $u, w$ are gauge, exactly as in CA-38. **Derived [port]:** nothing in the CA-38/CA-39 derivations uses the Pauli basis — only bilinearity, locality, and the shift — so every identity holds in $\mathcal{A}_L$; the Pauli *coefficient* equations of CA-38 are recovered by expanding in the matrix-unit or diagram basis of the charge blocks $\bigoplus_c M_{m_c(L)}(\mathbb{C})$.

## 74.4 Open-Window Mode Calculus

**Definition [choice].** Fix the phase base $z_n = e^{2\pi i n/L}$ on the length-$L$ open chain (CONVENTIONS (t): index phases at own scale). For a window $W = [j_0, j_1] \subset [1, L-1]$ define $H_n^{(W)} = \sum_{j \in W} z_n^j h_j$ and $J_n^{(W)} = \sum_{j \in W} z_n^j p_j$.

**Lemma [derived].** The exact periodic identity (42.1) ports to the open window with an edge remainder:

$$[H_m^{(W)}, H_n^{(W)}] = -i(z_n - z_m)\, J_{m+n}^{(W)} + \partial_{mn}^{(W)},$$

where $\partial^{(W)}_{mn}$ is supported in the two edge cells of $W$ (the telescoping of CA-42's derivation fails only where the window ends). The same holds for the ported (42.2)–(42.3) with their $D$- and $F$-density expansions. The Julia suite asserts the support statement directly; it is the strong form of "bulk identity modulo boundary". Periodic modes — the setting of CA-70's $L^{(N)}_n$ and of the registered periodic dilute algebra — are deliberately *not* defined here (decision D1; CA-70 obligation 3).

### 74.5 Witt/Virasoro Residuals with Two Central Constants

**Definition [port + CONVENTIONS (s)].** For a state $\omega$ let $:X:_\omega = X - \omega(X)\mathbf{1}$ termwise on densities. The chiral ansatz on a window is

$$\ell_n = A_n{:}H^{(W)}_n{:}_\omega + B_n{:}J^{(W)}_n{:}_\omega + \mu\,\delta_{n0}\mathbf{1}, \qquad \bar\ell_n = A_n{:}H^{(W)}_n{:}_\omega - B_n{:}J^{(W)}_n{:}_\omega + \mu\,\delta_{n0}\mathbf{1},$$

and the closure residuals carry the *two-constant* central term of CONVENTIONS (s):

$$r^{++}_{mn} = [\ell_m,\ell_n] - (m-n)\ell_{m+n} - \tfrac{\delta_{m+n,0}}{12}\big(m^3 c^*_L - m\,c_L\big)\mathbf{1},$$

with $r^{--}_{mn}$, $r^{+-}_{mn} = [\ell_m,\bar\ell_n]$ analogous. Setting $c^*_L = c_L$ is a measured outcome or theorem target, never an input. The estimator pair $(c_L(m), c^*_L(m))$ is defined by solving $\omega([\ell_m,\ell_{-m}] - 2m\ell_0) = \frac{1}{12}(m^3c^*_L - mc_L)$ at two mode values, relative to a declared $(\omega, W)$.

### 74.6 The Categorical Residual Set

**Definition.** The scale-$L$ residual family of the pair ($h$, window data) is

$$\mathcal{R}_L(h) = \big\{R^{\mathrm{cons}}_j,\ R^{\mathrm{boost}}_j\big\}_j \cup \big\{r^{\sigma\sigma'}_{mn}\big\}_{|m|,|n|\le K},$$

as elements of $\mathcal{A}_L$. This is the object whose absence was seam S3 of the 2026-07-06 consistency map: the moment/GNS constraints $\omega(X^\dagger RY) = 0$ of CA-43/CA-63, and the relaxed profiles and $\nu$-objectives of CA-64, are now well-posed on the categorical tower by substitution. **Boundary:** CA-63's existence theorem is proved for the qubit UHF algebra; its transport to the AF tower is W1.6's categorical corollary, not a claim of this shard.

### 74.7 Exact Transport and the Stretched Dichotomy

**Lemma 73.1 [derived] (exact transport).** Let $\theta_\varphi$ be the corner morphism of a placement (CA-71, CA-72). Since $\theta_\varphi$ is a $*$-morphism (Lemma 72.1), for every noncommutative $*$-polynomial $F$,

$$\theta_\varphi\big(F(h_{j_1},\dots,h_{j_r})\big) = F\big(\theta_\varphi(h_{j_1}),\dots,\theta_\varphi(h_{j_r})\big).$$

Hence the *entire* residual calculus transports exactly: the $\theta_\varphi$-image of every element of $\mathcal{R}_k(h)$ is the same residual built from the *stretched densities* $\theta_\varphi(h_j)$, which for the dyadic placement are supported on fine sites $\{2j-1, 2j, 2j+1\}$ with the inserted site pinned to vacuum (the CA-68 stretched-covariance phenomenon at the density level).

**Observation [derived] (the dichotomy).** The stretched density is *not* the fine density: $\theta_\varphi(h_j) \neq h^{(l)}_{j'}$ for any placement of the same $h$ at fine scale. Consequently the CA-72 descent-compatibility contract *fails* for the naive pairing $(\mathcal{R}_k(h), \mathcal{R}_l(h))$: annihilating the fine residuals does not annihilate the $\theta$-transported coarse ones. This is the precise residual-level content of "bare vacuum insertion is naive".

**Definition + Proposition 73.2 [derived] (coherent family).** The *stretched-augmented* residual family assigns to the fine scale

$$\mathcal{R}_l^{\mathrm{coh}} := \mathcal{R}_l(h) \ \cup \ \bigcup_{\varphi} \theta_\varphi\big(\mathcal{R}_k(h)\big)$$

over the placements $\varphi$ of the chosen refinement schedule. If a state $\omega_l$ annihilates $\mathcal{R}_l^{\mathrm{coh}}$ in the CA-63 sense ($\pi_{\omega_l}(R) = 0$ for all $R \in \mathcal{R}_l^{\mathrm{coh}}$), then the normalized pullback $\omega_k$ annihilates $\mathcal{R}_k(h)$, by Lemma 72.4 applied to $\pi_{\omega_l}(\theta_\varphi(R')) = 0$. Descent-compatibility thus holds *by construction* for the coherent family. **Price [recorded]:** the fine moment problem gains the stretched constraints — a strictly stronger feasibility demand on $\omega_l$. **Alternative [recorded, W2.1]:** a dressing $U$ may instead be chosen so that $\mathrm{Ad}_U \circ \theta_\varphi$ carries stretched residuals asymptotically into the null structure of $\mathcal{R}_l(h)$; quantifying that is precisely the dressed-refinement task, now stated as an equation rather than a slogan.

## 74.8 Julia Invariants

`src/FibonacciLocalOperators.jl` implements the two-site local-operator embedding on the CA-66 path basis (the single nontrivial Fibonacci $F$-matrix anchored to the registered Trebst source, unit-label moves trivial per CONVENTIONS (b)) and the Tier-1 generators (occupancies, hop, raw-cup pair terms with $u^2 = \varphi u$, dense $e_j$). `src/CategoricalResiduals.jl` implements the ported residual densities, coboundary distances (pinv/nullspace per the CA-64 lesson), the open-window mode identity with its edge-support assertion, and the transport tests of Lemma 73.1 against the CA-71/CA-72 machinery. The testset "categorical residual set (CA-73)" pins: disjoint-support commutation; the occupancy cross-check against the independent `occupation_number_matrix`; $e_j^2 = \varphi e_j$ on the full space and the braid relation $e_j e_{j\pm1} e_j = e_j$ *on the dense corner only* (on the full dilute space $e_1e_2e_1 = 0$, because $u_j = b_j^t b_j$ annihilates states with an empty neighbour — the checked computational shadow of CA-69's corner statement $\pi_A \,\mathrm{dTL}\, \pi_A \cong \mathrm{TL}$); source-anchored dilute relations; parity commutation of every Tier-1 generator (the shadow of CA-69's parity finding); the edge-support lemma in its sharp form (the bulk telescopes and exactly one top-edge current survives); a checked physics value — the pure-hopping density's conservation residual is a coboundary to $10^{-14}$ (hopping conserves momentum) — and the dilute-image dimension data at $L = 2, 3$ (pinned to CA-69's $9, 51$) together with the computed $L = 4$ values, independently re-run by the orchestrator — recorded here as *data for CA-74*, whose interpretation belongs to that shard, not this one.

## 74.9 Boundary

Tier-2 (parity-odd, fusion-vertex) densities are not defined here; they are the S4/W3.1 item. Periodic modes and the periodic evaluation map remain gated behind D1 (CA-70 obligation 3; interchiral caveat recorded in the plan). No estimation protocol for $(e_\infty, v_L, c_L, c_L^*)$ is fixed here (W2.2). Residual annihilation, exact or relaxed, is never symmetry evidence (CA-44, CA-64). The transport lemma is exact but says nothing about convergence; the coherent family makes descent hold by fiat at the cost of stronger constraints, and whether any physical state family satisfies them is precisely the open state-existence frontier (W1.6, W2.1).

**Erratum / qualification (2026-10-22).** Two corrections to the support reading of Lemma 73.1 (the transport identity itself stands). (i) Since $\theta_\varphi(1) = P_\varphi$ pins *every* inserted site, $\theta_\varphi(h_j)$ carries vacancy projectors on inserted sites far from $j$; the statement "supported on $\{2j-1, 2j, 2j+1\}$ with the inserted site pinned" holds only for a local representative $S_j$ with $\theta_\varphi(h_j) = S_j P_\varphi$, a different element of the fine algebra. (ii) Polynomials with constant terms transport with the corner unit:

$\theta_\varphi(F(h;1)) = F(\theta_\varphi(h); P_\varphi)$, so central terms and energy shifts go to multiples of $P_\varphi$, not of 1. Both are Corrections 76.B and 76.C of CA-76 (also CA-77 qualification (d), CA-79 Lemma 79.6).

# 75 The Jones–Wenzl Kernel Decision for the Dilute Evaluation Map

## 75.1 Status, Sources, and Dependencies

**Status: derived, with one numerical input.** This shard interprets the CA-73 decision data and settles the CA-69 conjecture at $L = 4$. The combinatorics and the kernel identification are local derivations from registered sources; the single essential numerical input is the computed rank $\operatorname{rank}\rho_4 = 322$ (CA-73, re-run here). Executes item W1.4 of `reviews/2026-07-06_pipeline_consistency_scoping` orchestration record in `reviews/2026-07-06_ca74_jw_kernel_decision/`.

Dependencies: CA-66, CA-69, CA-71, CA-72, CA-73; CONVENTIONS (b), (r).

## 75.2 Setting and the Decision Data

Fix Fibonacci $(\mathcal{C}, O)$, $O = \mathbf{1} \oplus \tau$, per CONVENTIONS (r), the tower $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$, and the CA-69 evaluation map $\rho_L : \mathrm{dTL}_L(\varphi) \to \mathcal{A}_L$. CA-73 computed (and this session re-ran) the decision data: the image of the Tier-1 dilute generators has dimension 9, 51, 322 at $L = 2, 3, 4$, equal at every $L$ to the dimension of the parity-preserving subalgebra, and at $L = 4$ exactly one short of $\dim \mathrm{dTL}_4 = M_8 = 323$. This shard answers the two questions those numbers pose: *why* 322, and *which* element spans the kernel.

## 75.3 Parity-Refined Multiplicities

**Derived.** Write $m_c^{\pm}(L)$ for the number of CA-66 basis paths of $O^{\otimes L}$ with total charge $c \in \{\mathbf{1}, \tau\}$ and even (+) or odd (−) occupied number $N$. Appending a site keeps $N$ (vacancy summand) or increments it and fuses the charge ($\tau$ summand), so

$$m_{\mathbf{1}}^{\pm}(L+1) = m_{\mathbf{1}}^{\pm}(L) + m_{\tau}^{\mp}(L), \qquad m_{\tau}^{\pm}(L+1) = m_{\tau}^{\pm}(L) + m_{\mathbf{1}}^{\mp}(L) + m_{\tau}^{\mp}(L),$$

with seed $(m_{\mathbf{1}}^{+}, m_{\mathbf{1}}^{-}, m_{\tau}^{+}, m_{\tau}^{-})(1) = (1, 0, 0, 1)$. The alternating sums $M_c^{-}(L) = m_c^{+}(L) - m_c^{-}(L)$ obey the Fibonacci recursion generated by $I - T$, $T = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right)$, giving $M_{\mathbf{1}}^{-}(L) = F_{L+1}$, $M_{\tau}^{-}(L) = -F_L$ and the closed forms

$$m_{\mathbf{1}}^{\pm}(L) = \tfrac{1}{2}\big(F_{2L-1} \pm F_{L+1}\big), \qquad m_{\tau}^{\pm}(L) = \tfrac{1}{2}\big(F_{2L} \mp F_L\big),$$

consistent with $m_{\mathbf{1}} = F_{2L-1}$, $m_{\tau} = F_{2L}$ (CA-66, T66.2). Charge $\mathbf{1}$ is even-heavy (the all-vacuum state and the $\tau \otimes \tau \to \mathbf{1}$ channel), charge $\tau$ odd-heavy (single-$\tau$ states); the asymmetry $\sim \varphi^L$ is subdominant to $m_c \sim \varphi^{2L}$.

| $L$ | $m_{\mathbf{1}}^{+}$ | $m_{\mathbf{1}}^{-}$ | $m_{\tau}^{+}$ | $m_{\tau}^{-}$ |
|---|---|---|---|---|
| 1 | 1 | 0 | 0 | 1 |
| 2 | 2 | 0 | 1 | 2 |
| 3 | 4 | 1 | 3 | 5 |
| 4 | 9 | 4 | 9 | 12 |
| 5 | 21 | 13 | 25 | 30 |
| 6 | 51 | 38 | 68 | 76 |

The $L = 4$ row is the observed split $13 = 9 + 4$, $21 = 9 + 12$. The parity-*preserving* subalgebra $\bigoplus_c \big(M_{m_c^{+}} \oplus M_{m_c^{-}}\big)$ has dimension

$$P(L) = \sum_c \big((m_c^{+})^2 + (m_c^{-})^2\big) = 9,\ 51,\ 322,\ 2135,\ 14445 \quad (L = 2, \ldots, 6),$$

with complement check $\dim \mathcal{A}_L - P(L) = \sum_c 2m_c^+ m_c^-$ ($610 - 322 = 288$ at $L = 4$). This derives the 322: the CA-73 datum says $\rho_4$ is *onto* the parity-preserving subalgebra, the maximum permitted by the parity obstruction of CA-69.

### 75.4 The Kernel Theorem

**Convention bridge (Rule 3, recorded).** SRC-TL-JONES works with $\mathrm{TL}_n(q)$, loop parameter $-(q + q^{-1})$, and $\ell = |q^2|$. Our diagram algebra has loop value $\beta = +\varphi = 2\cos(\pi/5)$ (CONVENTIONS (r); $e_j^2 = \varphi e_j$ checked in CA-73), i.e. $q = e^{i\pi/5}$, $\ell = 5$; quantum integers $[1], \ldots, [5] = 1, \varphi, \varphi, 1, 0$. The loop sign is a gauge choice ($\kappa_\tau = +1$ is sourced) and touches none of the dimension, trace, or radical statements used below; the identification is pinned by the source's own $\ell = 5$ example, whose module dimensions are exactly our path counts. In its notation the first Jones–Wenzl idempotent at the root is $E_{\ell-1} = E_4 \in \mathrm{TL}_4$; we write $p_4$.

**Theorem 74.1 [derived; one numerical input].** Let $\iota : \mathrm{TL}_4(\varphi) \cong \pi_A \, \mathrm{dTL}_4 \, \pi_A$ be the fully-occupied corner isomorphism ($A = \{1, 2, 3, 4\}$). Then

$$\ker \rho_4 = \mathrm{span}\{\iota(p_4)\},$$

and $\rho_4$ maps $\mathrm{dTL}_4(\varphi)$ onto the parity-preserving subalgebra of $\mathcal{A}_4$. *Proof.* (i) $p_4$ *exists and is negligible:* at $\ell = 5$ the trace form on $\mathrm{TL}_4$ is degenerate with one-dimensional radical generated by the unique idempotent $p_4$ satisfying $e_i p_4 = p_4 e_i = 0$ ($i = 1, 2, 3$), and $\mathrm{tr}_4(p_4) = 0$ [sourced :63–69, :135–141, :818]. (ii) $\iota(p_4) \neq 0$*:* $p_4$ has coefficient 1 on the identity diagram [sourced :792], diagrams are a basis, and $\iota$ is injective [sourced corner iso]. (iii) $\rho_4(\iota(p_4)) = 0$*:* the corner restriction $\rho_4 \circ \iota$ is the dense evaluation $\mathrm{TL}_4(\varphi) \to \mathrm{End}_{\mathcal{C}}(\tau^{\otimes 4})$ (CA-69), whose kernel is exactly the radical $\mathrm{span}\{p_4\}$: the quotient $Q_4(5) \cong \mathrm{End}(\Delta_q(1)^{\otimes 4})$ has dimension $2^2 + 3^2 = 13 = C_4 - 1$ [sourced :74–80; ledger derived]. Independently: $\rho_4 \circ \iota$ is a unital $*$-representation in the unitary gauge, so $\rho_4(\iota(p_4))$ is an orthogonal projection whose quantum trace is $\mathrm{tr}_4(p_4) = 0$; positivity of quantum dimensions (CONVENTIONS (r)) forces it to vanish. (iv) *Rank–nullity:* $\dim \mathrm{dTL}_4 = M_8 = 323$ [sourced CA-69] and $\mathrm{rank}\, \rho_4 = 322$ [numerical input, CA-73 datum, re-run this session], so the kernel is one-dimensional and (ii)+(iii) identify it. Since $322 = P(4)$ and the image lies in the parity-preserving subalgebra (CA-69), the map is onto it. □

Every ingredient except $\mathrm{rank}\, \rho_4 = 322$ is analytical and locally sourced; the numerical rank is the surjectivity half, and the Julia suite additionally confirms $\rho_4(\iota(p_4)) = 0$ to machine precision.

### 75.5 The Upgraded Conjecture

**Conjecture 74.2 [proposal; refines CA-69].** For all $L \geq 1$, $\rho_L$ maps $\mathrm{dTL}_L(\varphi)$ onto the parity-preserving subalgebra of $\mathcal{A}_L$, with kernel the two-sided ideal generated by the corner Jones–Wenzl embeddings, $\ker \rho_L = \langle \iota_A(p_4) : A \subseteq [L], |A| = 4 \rangle$. Exact for $L \leq 3$ (zero kernel) and $L = 4$ (Theorem 74.1; a two-sided ideal inside a one-dimensional kernel equals it). Under the conjecture,

$$\dim \ker \rho_L = M_{2L} - P(L) = 0,\ 0,\ 1,\ \mathbf{53},\ \mathbf{1066} \quad (L = 2, \ldots, 6);$$

because image $\subseteq$ parity-preserving is *proved*, these are unconditionally rigorous *lower bounds* on $\dim \ker \rho_L$ — the open half at $L \geq 5$ is surjectivity alone. For comparison, the dense kernels $\ker(\mathrm{TL}_n(\varphi) \to \mathrm{End}(\tau^{\otimes n}))$ have dimensions $C_n - F_{2n-1} = 1, 8, 43$ ($n = 4, 5, 6$); the dilute kernel is strictly larger because $\mathrm{dTL}_L$ glues binomially many dense corners [sourced :2314–2318, the dilute-radical decomposition]. A within-budget $L = 5$ check must be combinatorial — either exhibiting dilute words spanning all $P(5) = 2135$ parity-blocks, or computing $\dim\langle \iota_A(p_4) \rangle = 53$ through the dilute cellular structure; the 2188-dimensional rank computation is out of budget (AGENTS.md platform note).

### 75.6 Julia Invariants

`src/JonesWenzlKernel.jl` builds the corner unit $\Pi_A$, the corner-compressed dense generators $E_j$, and the Wenzl recursion $p_{r+1} = p_r - ([r]/[r+1])\, p_r E_r p_r$ on the CA-66 path basis, carrying the formal word expansion alongside the matrices. Because the local extraction of the recursion truncates its diagram factor (source note above), the testset "jones-wenzl kernel decision (CA-74)" pins the *defining* properties instead of the formula: idempotency, self-adjointness, two-sided annihilation $E_j p_k = p_k E_j = 0$ for $j < k$, and unit-word coefficient exactly 1. It further pins: $\operatorname{tr} \Pi_A = (2, 3)$ on the two charge blocks; $e_j^2 = \varphi e_j$ and the braid relation in the corner; $[1..5] = (1, \varphi, \varphi, 1, 0)$ with $[5] = 0$ at $10^{-14}$; survival $\|\rho(\iota(p_2))\| = \|\rho(\iota(p_3))\| = 1$; **the decision** $\|\rho(\iota(p_4))\|_{\mathrm{op}} = 2.3 \times 10^{-15} \leq 10^{-12}$; the CA-73 cross-checks `dilute_image_dimension`$(4) = 322 =$ `parity_even_dimension`$(4)$ (reused, not forked); and the full parity table above for $L = 2..5$ by pure path counting. 113 assertions, standalone green, re-run independently by the orchestrator; three mutations (Wenzl coefficient, recursion $\beta$, corner unit) each RED (14/25/8 failures), reverted.

### 75.7 Gram-Rank Deferral and Boundary

The Gram-rank kernel dictionary (descendant Gram matrices, JW $\leftrightarrow$ null-vector reading) is W2.3 and stays behind the endpoint-closing ledger W1.5/T1: Gram data need a canonical fusion-tree ONB in each fixed-$S$ sector, which no shard yet fixes; CA-74's statement is rank-only and basis-independent, which is why it could proceed (PLAN.md W1.4). Not claimed here: any continuum or Koo–Saleur statement; a closed form for the dilute ideal $\langle \iota_A(p_4) \rangle$ at general $L$; surjectivity beyond $L = 4$. Recorded gaps (acquisition queue): a source with the explicit algebraic Wenzl recursion (e.g. Kauffman–Lins or Morrison arXiv:1503.00384) and the categorical negligible-morphism formalism (e.g. EGNO Ch. 8 / Goodman–Wenzl); one line of the ILZ extraction (`:666`, "$\dim Q_8 = 233$") contradicts the paper's own $\ell = 5$ formulas (610) and is treated as an extraction artifact — only $\dim Q_4(5) = 13$ is used, triple-checked.

# 76 Integration Map: One Model-Independent Pipeline

## 76.1 Status, Sources, and Dependencies

**Status: map.** This shard makes *no new mathematical claim*. Every row cites the shard or note that carries the statement and its own status label; the two "unwritten links" are **Questions**; the acceptance contract is a **Proposal** that only reorders existing requirements. Everything is stated for a unitary fusion category $\mathcal{C}$, or a modular one when the declared target needs braiding, together with *declared extra data* $\mathsf{E}$ (site object, boundary module category, central-charge lift, chiral/full choice, fidelity target; CONVENTIONS (u), (ac)). Models occur only in the closing examples table (AGENTS.md Rule 16).

Dependencies: CA-01, CA-14, CA-15, CA-62, CA-65–CA-68, CA-71–CA-73, CA-81; CA-86 and CA-87 (merged 2026-10-22); CA-76–CA-80 (reviewed exploration content, 2026-10-22); `reconstruction/` LC-1–LC-5, LC-9, LC-10, LC-12 (LC-6, LC-11, LC-13 and CA-82 in the examples table only); CONVENTIONS (r), (s), (u), (v), (x), (ac)–(af).

## 76.2 The Pipeline as Typed Layers

Each layer is a typed object or map depending on $(\mathcal{C}, \mathsf{E})$ and a cutoff index $n$. "General" means stated for arbitrary $\mathcal{C}$ in the cited source; "missing" lists what no current source supplies.

| Layer (general object) | Supplied by | Already general | Missing |
|---|---|---|---|
| (1) Carrier: site object $O \in \mathcal{C}$; $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$; number filtration for $O = \mathbf{1} \oplus X$ | CA-65, CA-66; CA-87 Lemma 87.1 | AF tower, Bratteli rule $m_d(L{+}1) = \sum_c m_c(L) N_{c,O}^d$, carrier $\bigoplus_c \mathrm{Hom}(X_c, O^{\otimes L})$, $N$-grading (proposal-status shards; finite algebra elementary) | Criterion for when a site object is genuinely new: Lemma 87.1 blocks $O \cong Y \otimes \bar{Y}$; no general classification of non-blocked $O$ |
| (2) Dynamics: $h \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ self-adjoint; tile decomposition; Trotter history sum | CA-86; LC-4 §1 | History sum $= \mathrm{Ev}_{\mathcal{C}}(\Gamma) \times \prod_p w(p)$ for any $\mathcal{C}$ (Lemma 86.1, exact gate form Prop. 78.2; Cor. 86.2); Prop. 86.3 only under its commutativity hypothesis: the brick sweep is the noncommuting row-to-row transfer, and the ground-state claim needs nondegeneracy (CA-78 erratum) | A selection principle for $h$ (criticality, branch); CA-78 gives rejection tests only; in-repo integrable lines only for Temperley–Lieb-type input |
| (3) Refinement: isometries $V_\varphi$, corner maps $\theta_\varphi$, GNS descent | CA-68, CA-71, CA-72, CA-73 | Exact composition; unconditional descent with defect $2(1 - \sqrt{\omega(P_\varphi)})$; exact completion defect; residual transport (all derived) | Unital $*$-maps (strict OAR); dressing $W = UV$ with $\omega(P) \to 1$ on a critical family; NG1–NG5 check |
| (4) Continuum state and generators | CA-81; `wilson_triangle.md`, `virasoro_routes.md` | WT-1 for any coherent unital C*-tower; VR-1 under A1–A4 (local derivations) | Any instance of VR-1 (referee note :155–161); joint use of one state system |
| (5) Lattice symmetry: KS modes from the density | CA-14; LC-4 §2; `PROOF.md` | Prescription determined by $h_n$, with $e_n, b_n, c_n, c_n^*$ declared (CONVENTIONS (s), (af)) | Convergence beyond free fields; the no-wrap lesson for central terms generalised |
| (6) Fidelity: sewing and four-port equations, obstruction complexes, declared target | LC-1–LC-5, LC-9, LC-10, LC-12 §5; `realisation_and_fidelity.md` | C0–C7 for any $\mathcal{C}$ and module category; $\mathrm{HH}^1$, CE, $\mathrm{H}^3$, mixed complex, cokernel obstructions; $S, T$ minimum target; character-growth threshold sharp at $c > 1$ (Prop. 79.9, Lemma 80.12; conditional) | Lattice avatar of the sewing system; complete carriers; extended-algebra content of $V_{\mathbf{1}}$ |

**Notes on the rows.** (1) Lemma 87.1 is a *blocking criterion*: if a simple $Y$ has $Y \otimes \bar{Y} \cong \mathbf{1} \oplus X$, the dilute tower is a dense $Y$-tower with paired sites, so layers (2)–(6) inherit nothing new from dilution. (4) WT-1 needs unital injective $*$-maps; CA-72 supplies a non-unital $*$-morphism or a unital non-multiplicative map (Link (b) below). VR-1 has no verified instance: the accepted free-fermion estimate holds on fixed momentum cores, not as a uniform graph-norm bound. (5) The only sourced convergence theorem is free-fermion (CA-14 boundary); the fermion benchmark records that periodic momentum wrap invalidates its central-term recipe. (6) LC-12 §5, under an explicit modular-character hypothesis, forces infinitely many Virasoro-primary families in the vacuum carrier for the instance treated there; the general-$c$ form is now sharp at $c > 1$ (CA-79 Prop. 79.9, CA-80 Lemma 80.12, both conditional).

### 76.3 Unwritten Links

**Link (a) [Question]: the history sum as lattice avatar of the cell equations.** Fix $(\mathcal{C}, O, h)$. For a brick-wall rectangle $P$ with $\ell$ sites and $M$ rows, the Trotter product $Z_P^{\mathrm{lat}} := U(\delta\tau)^M \in \mathrm{End}_{\mathcal{C}}(O^{\otimes\ell})$ is, to first order per plaquette, the CA-86 history sum. LC-1 asks for cell maps $Z_P : D_{\mathrm{in}} \to D_{\mathrm{out}} \otimes L_P$ obeying C0–C7. Open:

(a1) *Typing.* Which module category and channel spaces $M_{ab}^i$ of LC-1 §1 correspond to the open-chain carriers $\mathrm{Hom}_{\mathcal{C}}(X_c, O^{\otimes\ell})$ used as lattice boundary data?

(a2) *Sewing (C2, C3).* Vertical stacking of rectangles multiplies Trotter products; spatial gluing is not a tensor product, because plaquettes straddle the seam. Is the seam correction a bounded-support insertion whose limit is LC-1's anomaly-line identification? (*Amendment 2026-10-22:* the seam *operator* $K_s = s^\dagger\,\mathrm{rot}$ exists for a declared seam pairing $s$ and decides one-tile spatial reflection positivity, CA-92 Thm. 92.3 [cond. on $s$, (RH)]; this sewing question stays open.)

(a3) *Ward (C4, C7).* Corollary 86.2 says isotopy fails at finite cutoff through the geometric weights. Do derivatives of $Z_P^{\mathrm{lat}}$ in the anisotropy parameter produce insertions equal to the layer-(5) KS densities up to residuals that vanish under layers (3)–(4)?

(a4) *Limit.* Does C0 hold at every cutoff (since $\mathrm{Ev}_{\mathcal{C}}$ obeys the $F$-moves), and do refined lattice cells converge to a solution of C1–C6?

**Link (b) [Question]: the CA-72 defect scalar as input to WT-1's A3.** WT-1 A3 asks for summable $d_{k,n} \geq \|\omega_{n+1} \circ \alpha_{n+1}^k - \omega_n \circ \alpha_n^k\|$. CA-72 gives, for each placement $\varphi_n$, the scalar $\varepsilon_n := 1 - \omega_{n+1}(P_{\varphi_n})$ (written $q_n$ in CA-76), which fixes the vacuum defect of the descended isometry (Theorem 72.3) and the GNS-seminorm defect of the unital completion (Corollary 72.8); the operator-norm completion defect is $|\chi(ST) - \chi(S)\chi(T)|$ whenever $P \neq 1$, whatever $\varepsilon_n$ (CA-76 Correction 76.A). Open:

(b1) *Typing.* A1 of WT-1 demands coherent unital injective $*$-maps; $\theta_\varphi$ is not unital and $\Phi_\chi$ is not multiplicative. Does a soft variant of WT-1 hold for coherent-up-to-defect UCP towers whose multiplicativity defect is the Theorem 72.7 expression? *Status 2026-10-22:* a strict unital tower exists globally by scalar augmentation (Prop. 76.2, derived, nonlocal), and a unital reducing extension of vacuum insertion exists for every $\mathcal{C}$ (Prop. 76.3, Cor. 76.4). *Amendment 2026-10-22:* a *local* unital lift is derived for pointed $O$ with a relative half-braiding on $\langle O\rangle$ (CA-90 Thm. 90.1; braided $\mathcal{C}$ suffices); (b1) is closed there, open at bare $\mathsf{L0}$.

(b2) *Split.* Is $d_{k,n}$ bounded by an explicit function of $\{\varepsilon_j\}_{j\geq n}$ plus the mismatch between $\omega_n$ and the normalized pullback of $\omega_{n+1}$ (Lemma 72.5 Gram domination)? *Answered in the strict case* (Prop. 76.1, derived): $d_{k,n} \leq p_n r_n + 2\varepsilon_n$ for intertwining unital lifts, $r_n$ that mismatch; no bound by $\varepsilon_n$ alone exists.

(b3) *Attainability.* Corollary 72.8 needs only $\varepsilon_n \to 0$; WT-1 needs summability. Does any dressed family $W = UV$ (CA-68) achieve $\sum_n \varepsilon_n < \infty$ for ground states of a fixed layer-(2) Hamiltonian? *Open*; sharpened as global fidelity versus local lift (Question 76.12), with a conditional gap route (Thm. 76.6).

### 76.4 One Acceptance Contract

**Proposal.** A *lattice tower* for $(\mathcal{C}, \mathsf{E})$ is accepted only with the following data and proofs. Each item names the approach supplying a construction [C] and the one supplying the proof obligation [P]; "none" means no current source constructs it.

1. *Declared input.* $(\mathcal{C}, \mathsf{E})$, with the fidelity target (modular data up to unit-preserving relabelling, or braided equivalence, or a defect target). [C] CA-65 pair $(\mathcal{C}, O)$; [P] CONVENTIONS (u), `realisation_and_fidelity.md`:23–53.

2. *Local algebras.* Finite-dimensional C*-algebras $\mathcal{A}_n$ with declared support structure and translation. [C] CA-65; [P] CA-15 item 1, LC-4 §1.

3. *Local dynamics.* Self-adjoint finite-range density $h_n$, cutoff-dependent couplings allowed. [C] CA-86 tiles, LC-4 §1 transfer factors; [P] LC-1 C7, CONVENTIONS (ac).

4. *Refinement.* Coherent unital injective $*$-maps of bounded support growth with smearing, or a declared soft fallback. [C] CA-68, CA-71, CA-72 completion; [P] CA-67 strict list and NG1–NG5, LC-4 §3.

5. *States.* Microscopic states $\omega_n$ with summable row drift and a computable tail. [C] CA-66 states, LC-4 §3 ground-state tier; [P] WT-1 A2–A3, WT-2.

6. *GNS refinement.* $J_k[a] = [\alpha_k(a)]$, pointed or with quantified vacuum defect. [C] CA-72 Theorem 72.3; [P] WT-1 step $\langle 1 \rangle 5$, LC-4 §3.

7. *Lattice symmetry.* Standard KS modes of $h_n$ with declared $e_n, b_n$, velocity and the two central constants. [C] CA-14, LC-4 §2; [P] CA-15 items 2 and 6, CONVENTIONS (s), (af).

8. *Residual descent.* A residual family compatible with refinement and controlled on observable words, not only on the vacuum. [C] CA-73 coherent family; [P] CA-72 contract, VR-2.

9. *Generator convergence.* Graph-norm certificate on the *same* GNS system as item 6. [C] VR-1 (no instance); [P] LC-4 §4, CA-68 compatibility target, CA-81.

10. *Covariance.* Observable covariance of the limiting generators on local algebras. [C] none; [P] CA-15 scaling and covariance checks.

11. *Analytic completion.* Positive energy, essential self-adjointness, integration to a projective unitary representation, local-net axioms. [C] none; [P] LC-4 §5, VR-1 boundary.

12. *Local conformal consistency.* The stress of the limit is an internal field generating coordinate changes, matching item 7. [C] LC-1, LC-9 (equations only); [P] LC-1 C4–C7, LC-W.

13. *Fidelity.* Identification of the limit's sector data with the declared target, including exclusion of unintended sectors. [C] LC-9, LC-10 (equations only); [P] `realisation_and_fidelity.md`:46–62, LC-4 §5, LC-12 §5 constraint.

### 76.5 Status of the Six Explorations

Reviewed paragraphs: `explorations/2026-10-22-astra/D*/INTEGRATION.md` §(a); the content now lives in CA-76–CA-80 with per-statement grades. Every headline was graded *needs reformulation* (§(d) there).

**D1 carrier theory.** CA-77: unital vs corner towers, occupation grading and parity criterion, decidable dual-pair blocking (generalises Lemma 87.1), constant Morita class, configuration Fock carrier [derived]; corner corrections in CA-76. D1-C [conjecture; conditional-ok, framing overstated]: category analytically inert, tuned contact term, $z = 2$.

**D2 selection principle.** CA-78: coefficient compiler, exact gate ODE, blocking leakage, cubic cokernel test (rejection only), YBE vs brick sweep (converse as repaired) [derived]; interface Lemmas 78.9–78.11 [conditional-ok]; erratum to Prop. 86.3. D2-C [conjecture]: sufficiency not selection, category-blind, no verified instance.

**D3 refinement meets dynamics.** CA-76: drift bounds and scalar augmentation (76.1–76.2) and polar tangent (76.9) [derived]; gap route 76.6 [conditional-ok]. D3-C (76.10) [conjecture, unverifiable]: a UV locality lemma that cannot by itself give summable critical drift.

**D4 Koo–Saleur beyond free fields.** CA-79: tile modes, refinement defect, exact-covariance no-go [derived]; shell certificate 79.2 and $c > 1$ threshold 79.9 [conditional]; D4-X stays in the note [conditional, sketch]. D4-C [conjecture; overstated as a statement about $\mathcal{C}$]: any spin chain fits the carrier, the input is the full CFT, site growth is unconstrained.

**D5 fidelity and lattice sewing.** CA-80: exact sewing along tile legs, temporal reflection positivity, Ward product rule, scalar falsifier [derived]; two-sided certificate 80.9 [conditional, index limit step open]. D5-C [conjecture, unverifiable]: contains the open UMTC realisation problem and nearly duplicates D6-C.

**D6 functor and obstruction theory.** No shard. P1, P2, N1 [confirmed] are in CA-76 (76.3, 76.1, 76.A); braided local lift R1 [proposal, 76.5]. Typed source/target and contract K0–K12, adding K8 dynamics, K10 typed defects, K12 effectivity [proposal; conditional-ok], P3, P4 [confirmed], N2, N3 remain in the note. D6-C [conjecture, unverifiable]: contains the realisation problem; its chiral-limit clause conflicts with D2/D4/D5.

### 76.6 Decisions Taken (2026-10-22): the Layering Principle

Tobias resolved the three open decisions by one rule (AGENTS.md, Rule 17): state each result at the most general layer at which it is true and name the extra data the next layer needs. (1) *Chirality* (PLAN.md D1): chiral and full two-chiral targets are both in scope; each statement declares its setting (D6 chiral; D2, D4, D5 full; CA-80). (2) *Input data*: layers of increasing data (fusion, braided, modular, enriched $\mathsf{E}$); the data a statement needs is part of the statement. (3) *Targets*: a ladder, limit exists $\to$ conformal $\to$ faithful to a declared theory; a chain $h \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ is an instance at the rung its certificate reaches, otherwise a candidate. The six headline conjectures are restated on this grid in CA-88.

### 76.7 Examples (Instances Only)

| Instance | Calibration datum it supplies to the general map |
|---|---|
| Fibonacci | Lemma 87.1 applies ($\tau\otimes\tau \cong \mathbf{1}\oplus\tau$): dilute tower is blocked dense; LC-12 §5 genus-one constraint. |
| Ising | Lemma 87.1 applies with $X = \psi$; CA-82 gives a fermionic compiler with a sourced local limit (calibration, not the general route). |
| Semion | LC-6 associator obstruction for $A = \mathbf{1} \oplus s$; LC-11/LC-13 conditional four-port solution. |
| Haagerup | hypothesis of Lemma 87.1 fails for $X = \rho$ (CA-87 proposal): first non-blocked carrier test case. |

### 76.8 Boundary

This shard proves nothing and promotes no status. Rows inherit the labels of their sources; the links and the contract are questions and proposals. CA-86 and CA-87 are merged (2026-10-22); the D1–D6 entries carry the review grades of their sources and promote nothing.

# 77 Corner-to-Wilson Bridge: Refinement Meets Dynamics

## 77.1 Status, Sources, and Dependencies

**Status: mixed; every statement carries its own label.** Derived: Corrections 76.A–C, Propositions 76.1–76.3, Corollary 76.4, Proposition 76.9, Negative 76.11. Conditional: Theorem 76.6, Corollary 76.7, Remark 76.8. Proposal: 76.5 (superseded by CA-90). Conjecture: 76.10. Question: 76.12. All content is lifted from reviewed in-repo exploration records; their review grades are carried, not upgraded. **Abbreviations:** "D*i* NOTE/REVIEW/INTEGRATION" are the files `NOTE.md`, `REVIEW.md`, `INTEGRATION.md` in `explorations/2026-10-22-astra/D1-carrier-theory/`, `.../D3-refinement-meets-dynamics/`, `.../D4-koo-saleur-beyond-free/`, `.../D6-functor-and-obstruction-the` "D3 checks", "D6 checks" are their `checks/` bundles. Reproducibility: the exact finite checks live in those bundles; porting them to `scripts/` with a run bundle is open. No Julia or LaTeX run was made for this shard. Dependencies: CA-68, CA-71, CA-72, CA-73, CA-75 (Link (b)), CA-84; WT-1; CONVENTIONS (h), (r), (t), (af), (ak), (am), (an).

## 77.2 Setting and Notation

**Setting [CONVENTIONS (r), (t); notation local to this shard].** $\mathcal{C}$ is unitary fusion, $O = \mathbf{1} \oplus X$ with a chosen isometric unit inclusion, $N_n = 2^n N_0$ sites on an *open* ordered chain, $\mathcal{A}_n = \mathrm{End}_{\mathcal{C}}(O^{\otimes N_n})$ acting faithfully on $\mathcal{H}_n = \bigoplus_c \mathrm{Hom}_{\mathcal{C}}(c, O^{\otimes N_n})$. $V_n$ is the CA-71 dyadic vacuum insertion, $W_n = U_n V_n$ a dressing ($U_n$ unitary in $\mathcal{A}_{n+1}$), $P_n = W_n W_n^\dagger$, $\theta_n = W_n(\cdot)W_n^\dagger$, $W_n^k = W_{n-1}\cdots W_k$, $P_n^k = W_n^k W_n^{k\dagger}$, $\theta_n^k = \mathrm{Ad}\, W_n^k$. States $\omega_n$ on $\mathcal{A}_n$ are chosen independently; $p_n = \omega_{n+1}(P_n)$, $q_n = 1 - p_n$, and for $p_n > 0$ the normalized pullback $\tau_n = \omega_{n+1}\theta_n/p_n$ has mismatch $e_n = \|\tau_n - \omega_n\|$. State distances are full dual norms (at most 2); $\delta_n^2 = 2(1-\sqrt{p_n})$ is the Theorem 72.3 defect. $\alpha_n : \mathcal{A}_n \to \mathcal{A}_{n+1}$ always denotes a genuine unital injective $*$-map, $\alpha_n^k = \alpha_{n-1}\cdots\alpha_k$; $\Phi_\chi$ is the CA-72 UCP completion; $d_{k,n}$ is as in WT-1 A3. **Signs.** $H = \sum_Z h_Z$ with Hermitian interval energy densities, as in (h), (af); a CA-86 Hamiltonian $H = -\sum_j h_j$ of (ak) enters as $h_Z = -h_j^{(\mathrm{ak})}$. The (ak) cup-cap tile $e_j$ is not used, and the mismatch $e_n$ is unrelated to the density subtraction of (af). **Clash.** CA-75 Link (b) writes $\varepsilon_n$ for what is $q_n$ here; below $\epsilon_n$ is the vacuum residual of Theorem 76.6.

## 77.3 Corrections to CA-72 and CA-73

**Correction 76.A [derived; found independently by D1, D3, D6; confirmed].** Put $\mathrm{cov}_\chi(S,T) = \chi(ST) - \chi(S)\chi(T)$. If $P \neq 1$, the Theorem 72.7 defect $\mathrm{cov}_\chi(S,T)(1-P)$ has operator norm $|\mathrm{cov}_\chi(S,T)|$, *whatever* $\omega(1-P)$ is, since $\|1-P\| = 1$. Witness (plain multi-matrix algebras): $\theta : M_2 \to M_4$ into the first block, $\chi = \mathrm{Tr}\,/2$, $S = T = \mathrm{diag}(1,-1)$; the defect is $1-P$, of norm 1, while its GNS seminorm vanishes when $\omega(P) = 1$ (D3 checks `output.txt`:5). Source eq. (29) is an operator-norm asymptotic homomorphism property for composites. Hence Corollary 72.8's reading, that the completion family satisfies eq. (29) precisely when $\omega_l(1-P) \to 0$ in GNS seminorms, is **not supported as an operator-norm claim**; the GNS-seminorm identity of Theorem 72.7 stands. CA-75:145–147 ("modulus of softness") inherits this qualification. A soft-C*-system statement for any completion family needs its own operator-norm and composition argument.

**Correction 76.B [derived; D1, D3, D4 concur; confirmed].** $\theta_n(1) = P_n$ pins *every* inserted site, so $\theta_n(a_I)$ for $a_I$ supported on an interval $I$ contains vacancy projectors on inserted sites far from $I$. The support statement of Lemma 73.1 (CA-73:169–172, stretched densities on $\{2j-1, 2j, 2j+1\}$) holds for a *local representative*, which agrees with $\theta_n(h_j)$ on $\mathrm{Ran}\, W_n$ but is a different element of $\mathcal{A}_{n+1}$. General witness (D3 REVIEW:34): an (ak) hop $h^+$ moving $X$ onto

a distant inserted site does not commute with that site's vacancy pin. This identifies explicit projectors; it infers no nonlocality from fusion constraints.

**Correction 76.C [derived; confirmed].** Polynomials with constants transport with corner unit: $\theta(F(h;1)) = F(\theta(h);P)$, not $F(\theta(h);1)$; constants in residuals (central terms, energy shifts) transport to multiples of $P$. Centering: with $m = \tau(h)$, the transported coarse centering $\theta(h) - mP$ and the fine centering $\theta(h) - pm1$ differ by $m(p1 - P)$, whose fine GNS seminorm is exactly $|m|\sqrt{pq}$.

## 77.4 Corner Weight to Wilson Drift

**Proposition 76.1 [derived; D6 P2 = D3-1, found independently; confirmed].** ASSUME $W$ is an isometry implementing $\theta : \mathcal{A} \to \mathcal{B}$ between finite multi-matrix algebras, $P = WW^\dagger$, $\sigma$ a state on $\mathcal{B}$, $p = \sigma(P) > 0$, $q = 1 - p$, $\tau = \sigma\theta/p$, $\rho$ a state on $\mathcal{A}$. PROVE

(i) $\|\sigma\theta - \rho\| \le p\|\tau - \rho\| + q$, (ii) $\|\sigma\Phi_\chi - \rho\| \le p\|\tau - \rho\| + 2q$,
(iii) if $\alpha$ is a unital $*$-map with $\alpha(a)W = Wa\ \forall a:\ \|\sigma\alpha - \rho\| \le p\|\tau - \rho\| + 2q$,
(iv) if each $\alpha_n$ intertwines $W_n:\ d_{k,n} \le p_n e_n + 2q_n$, $q_n = \delta_n^2 - \frac{1}{4}\delta_n^4$, $2q_n = (1 + \sqrt{p_n})\,\delta_n^2 \le 2\delta_n^2$.

⟨1⟩1. $\sigma\theta = p\tau$, so $\sigma\theta - \rho = p(\tau - \rho) - q\rho$; the triangle inequality gives (i).

⟨1⟩2. Lemma 72.6: $\sigma\Phi_\chi = p\tau + q\chi$; subtract $\rho = p\rho + q\rho$, use $\|\chi - \rho\| \le 2$.

⟨1⟩3. Intertwining for $a$ and $a^*$ gives $P\alpha(a) = WaW^\dagger = \alpha(a)P$, so $\alpha(a) = \theta(a) + (1-P)\alpha(a)(1-P)$ and $\sigma\alpha = p\tau + q\eta$, $\eta$ a state if $q > 0$; conclude as in ⟨1⟩2.

⟨1⟩4. $\omega_{n+1}\alpha_{n+1}^k - \omega_n\alpha_n^k = (\omega_{n+1}\alpha_n - \omega_n)\alpha_n^k$, and pullback along a unital $*$-map is contractive; apply ⟨1⟩3 with $\sigma = \omega_{n+1}$, $\rho = \omega_n$. The identities follow from $\sqrt{p_n} = 1 - \delta_n^2/2$. □

The bound is attained ($p = 9/25$, drift $2q = 32/25$: D6 checks `raw_output.txt`:5). No function of $q_n$ alone bounds $d_{k,n}$: even $p_n = 1$ leaves $e_n$ free. **Caution [D6 REVIEW:155–161].** Since $P_n$ pins every inserted site, $\sum_n q_n < \infty$ is a *global* fidelity demand of exact-RG strength on an extensive chain.

**Proposition 76.2 [derived; D3-1 (2.2)–(2.3), generalising CA-84; confirmed].** For any isometries $W_n$, $\hat{\mathcal{A}}_n = \mathcal{A}_n \oplus \mathbb{C}$, $\hat{\alpha}_n(a,z) = (\theta_n(a) + z(1 - P_n), z)$, $\hat{\omega}_n(a,z) = \omega_n(a)$ define a strict unital injective $*$-tower (cross terms vanish; composites obey $\theta_m^n(1 - P_n^k) + (1 - P_m^n) = 1 - P_m^k$) with exact adjacent drift $\hat{D}_n = \|p_n\tau_n - \omega_n\| + q_n \in [\,2q_n,\ p_n e_n + 2q_n\,]$ (the dual norm of a direct sum is the sum). If $\omega_n(P_n^k) \to 0$ for each fixed $k$, the limit on row $k$ is the character $(a,z) \mapsto z$ (*scalar collapse*). **Boundary:** $\hat{\alpha}_n$ is generally nonlocal through $P_n$; for it, as for a bare corner, every local approximant $b_B$ of $\theta(a)$ has $\|\theta(a) - b_B\| \ge \|a\|/2$ (D3 REVIEW:60–61).

**Proposition 76.3 [derived; D6 P1; confirmed].** ASSUME $\mathcal{A} = \bigoplus_a M_{d_a}$, $\mathcal{B} = \bigoplus_b M_{e_b}$, $\theta$ an injective corner $*$-map with $\theta(1) = P$ in which block $a$ occurs in block $b$ with multiplicity $r_{ba}$. PROVE: a unital $*$-map $\alpha$ with $P\alpha(x)P = \theta(x)$, $[P, \alpha(x)] = 0$ exists iff there are integers $q_{ba} \ge 0$ with $e_b - \sum_a r_{ba}d_a = \sum_a q_{ba}d_a$ for all $b$.

⟨1⟩1. Diagonal matrix units of $M_d$ map to equal-rank projections, so every representation has dimension in $d\,\mathbb{Z}_{\ge 0}$; restricting $\alpha$ to $(1 - P)\mathcal{B}(1 - P)$ blockwise gives the condition.

⟨1⟩2. Conversely, $q_{ba}$ defining copies on the complement give $\beta$ with $\beta(1) = 1 - P$; set $\alpha = \theta + \beta$. □ By ⟨1⟩3 of 76.1 every $\alpha$ intertwining $W$ is such an extension; block ranks, hence the condition, are invariant under unitary dressing.

**Corollary 76.4 [derived; reviewer remark R2, D6 REVIEW:142–151].** For vacuum insertion $\mathcal{A}_L \to \mathcal{A}_{2L}$ the condition of 76.3 holds for *every* $\mathcal{C}$. *Proof.* Block $c$ maps to block $c$ with $r_{cc} = 1$; with $m_c(L) = \dim \mathrm{Hom}(c, O^{\otimes L})$ and $O^{\otimes 2L} \cong O^{\otimes L} \otimes O^{\otimes L}$ the complement is $m_c(2L) - m_c(L) = \sum_a (N^c_{a,O^{\otimes L}} - \delta_{ac}) m_a(L)$, with nonnegative coefficients because $\mathbf{1} \subset O^{\otimes L}$. □ So

the arithmetic never obstructs vacuum insertion; the open requirement is **locality** (and coherence) of the unital lift.

**Proposal 76.5 [proposal; reviewer R1; superseded by CA-90 Theorem 90.1, derived, non-pointed check included].** For braided $\mathcal{C}$ let $B : O^{\otimes L} \otimes O^{\otimes L} \to O^{\otimes 2L}$ interleave the new strands across the old with one fixed crossing type, and $\alpha(T) = B(T \otimes \mathrm{id})B^\dagger$. Proposed: $\alpha$ is a unital injective $*$-map with $\alpha(T)V_L = V_L T$ (braiding with $\mathbf{1}$ is trivial), it maps a coarse two-site density on $j, j+1$ into fine sites $\{2j-1, 2j, 2j+1\}$ by naturality, and $\mathrm{Ad}_U \circ \alpha$ intertwines $W = UV$. A pointed braided instance passes at $L = 2, 3$ with a range-2 mutation rejected (D6 checks `review_braided_lift_output.txt`:2–4) but cannot discriminate braiding phases; a non-pointed check with sourced $F, R$ is owed (D6 INTEGRATION:117–123). For non-braided $\mathcal{C}$ the construction needs a half-braiding on $O$ (a lift to $Z(\mathcal{C})$): open. *Amendment (2026-10-22, round 2):* established in CA-90 (Theorem 90.1, derived): a pointed relative half-braiding on $\langle O\rangle$ suffices, no lift to $Z(\mathcal{C})$; one-gap lifts are obstructed when $[O]$ is not central (Proposition 90.2).

## 77.5 Dynamics: Vacuum Residual and Gap

**Theorem 76.6 [conditional; D3-4; proof checked, grade conditional-ok].** Assume $K_n \geq 0$ ($K_n = H_n - E_n$, $E_n$ a certified ground energy) with unit $\Omega_n$, $K_n\Omega_n = 0$, $\omega_n = \langle \Omega_n, \cdot\, \Omega_n\rangle$; a unique gapped fine vacuum in the declared comparison carrier, $K_{n+1} \geq \Delta_{n+1}(1 - |\Omega_{n+1}\rangle\langle\Omega_{n+1}|)$; a residual $\|(K_{n+1}W_n - \lambda_n W_n K_n)\Omega_n\| \leq \epsilon_n$, $\lambda_n > 0$; and either coherent $\alpha_n$ intertwining $W_n$ or the tower of 76.2. Prove $d_{k,n} \leq \min(2, 2\epsilon_n/\Delta_{n+1})$ and $q_n \leq \min(1, (\epsilon_n/\Delta_{n+1})^2)$ on every row; hence $\sum_n \epsilon_n/\Delta_{n+1} < \infty$ gives WT-1 A3.

⟨1⟩1. Put $\psi = W_n\Omega_n$, $Q = 1 - |\Omega_{n+1}\rangle\langle\Omega_{n+1}|$. As $K_n\Omega_n = 0$ the residual is $K_{n+1}\psi = K_{n+1}Q\psi$, so $\|Q\psi\| \leq \epsilon_n/\Delta_{n+1}$.

⟨1⟩2. Two pure states with overlap modulus $u$ are at distance $2\sqrt{1-u^2} = 2\|Q\psi\|$.

⟨1⟩3. The trial state $\sigma_n = \langle\psi, \cdot\,\psi\rangle$ obeys $\sigma_n\alpha_n = \omega_n$ (and $\hat\sigma_n\hat\alpha_n = \hat\omega_n$ for $\hat\sigma_n(b,z) = \sigma_n(b)$, as $\psi \in \mathrm{Ran}\, P_n$); pullback is contractive.

⟨1⟩4. $\psi \in \mathrm{Ran}\, P_n$ gives $|\langle\Omega_{n+1}, \psi\rangle|^2 \leq p_n$, so $q_n \leq \|Q\psi\|^2$. □

Energy variant: $E_n^{\mathrm{tr}} = \langle\psi, K_{n+1}\psi\rangle$ gives $d_{k,n} \leq 2\sqrt{E_n^{\mathrm{tr}}/\Delta_{n+1}}$, $q_n \leq E_n^{\mathrm{tr}}/\Delta_{n+1}$; the overlap inequality is source-local (`latticefermions.tex`:487–491). Degenerate ground spaces need a selected sector; zero variance does not select the ground state (D3 checks `output.txt`:13).

**Corollary 76.7 [conditional; reviewer non-collapse corollary, D3 REVIEW:65–68].** Under 76.6 let $s_k = \sqrt{2}\sum_{j\geq k} \epsilon_j/\Delta_{j+1} \leq \sqrt{2}$. Then $\omega_n(P_n^k) \geq (1 - s_k^2/2)^2$ for all $n > k$, so scalar collapse (76.2) cannot occur, and D1-T5b's sufficient condition $\sum_n \sqrt{1-p_n} < \infty$ (D1 NOTE:383–385) holds.

⟨1⟩1. Fix phases with $c = \langle\Omega_{j+1}, W_j\Omega_j\rangle \geq 0$; then $\|W_j\Omega_j - \Omega_{j+1}\|^2 = 2 - 2c \leq 2(1-c^2) \leq 2(\epsilon_j/\Delta_{j+1})^2$.

⟨1⟩2. Telescoping through isometries, $\|W_n^k\Omega_k - \Omega_n\| \leq s_k$; so $\omega_n(P_n^k) \geq (\mathrm{Re}\langle W_n^k\Omega_k, \Omega_n\rangle)^2 \geq (1 - s_k^2/2)^2$.

⟨1⟩3. $\sqrt{q_n} \leq \epsilon_n/\Delta_{n+1}$ by 76.6. □

**Remark 76.8 [derived arithmetic on conditional inputs; D3 (4.3), (5.1)].** If $\Delta_n \geq c\, a_n^z$ and $\epsilon_n \leq C a_n^{z+y}$, $y > 0$, $a_n = 2^{-n}$, then $d_{k,n} \leq C' 2^{-yn}$, a geometric tail. A circuit-conjugacy comparison with at most $cN_{n+1}$ mismatch terms of norm $\eta_n \leq C a_n^u$ gives $\epsilon_n \leq cN_{n+1}\eta_n$ plus an energy-origin cost $2\zeta_n$; summability needs $u > 1 + z$. Dyadic spacing does not sum $O(1)$ or $O(1/n)$ per-step errors.

## 77.6 Polar Heat Refinement

**Proposition 76.9 [derived; D3-5; confirmed, also on a constrained multiplicity-two carrier].** Assume finite $L$, $H = \sum_Z h_Z$ Hermitian categorical interval terms on the fine open chain, $V = V_L$, $P = VV^\dagger$, $P_Z$ the product of vacancy projectors on inserted sites in $Z$, $G(t) = V^\dagger e^{-2tH} V$ and $W(t) = e^{-tH} V\, G(t)^{-1/2}$. Prove $W(0) = V$, $W(t)^\dagger W(t) = 1$, $\dot W(0) = -(1-P)HV = \sum_Z [P_Z, h_Z]\, V$ with each $[P_Z, h_Z]$ anti-Hermitian, supported in $Z$, of norm $\le 2\|h_Z\|$, and $\frac{d}{dt} W^\dagger H W|_0 = -2V^\dagger H(1-P)HV \le 0$.

⟨1⟩1. $e^{-tH}$ is invertible, so $G > 0$ and $W^\dagger W = G^{-1/2} G G^{-1/2} = 1$ in every charge block.

⟨1⟩2. $\dot G(0) = -2V^\dagger HV$ and $\frac{d}{dt} G^{-1/2}|_0 = V^\dagger HV$, so $\dot W(0) = -HV + PHV$.

⟨1⟩3. $h_Z$ acts trivially on inserted sites outside $Z$: $P h_Z V = P_Z h_Z V$ and $P_Z V = V$, hence $[P_Z, h_Z]V = -(1-P)h_Z V$. No braiding and no Hilbert tensor factorisation is used.

⟨1⟩4. Differentiating $W^\dagger H W$, both terms equal $-X^\dagger X$, $X = (1-P)HV$. □

Supplements [derived; confirmed]: $V^\dagger h_Z (1-P) h_Y V = 0$ unless $Z, Y$ share an inserted site; a local tangent $D = \sum_I x_I V$ with $V^\dagger D + D^\dagger V = 0$ extends to anti-Hermitian $a_I = x_I P_I - P_I x_I^\dagger - \frac{1}{2} P_I (x_I - x_I^\dagger) P_I$ supported in $I$, $\|a_I\| \le 3\|x_I\|$, $\sum_I a_I V = D$ (D3 (6.3)). For $t > 0$, $\dot W = -(1-P_t)HW + W K_{\text{frame}}$ with $P_t = WW^\dagger$ not a product of local projectors and $K_{\text{frame}}$ generally nonzero (D3 (6.2)): this is the all-orders difficulty.

**Conjecture 76.10 [conjecture D3-C; review: well posed, no counterexample found, unverifiable].** Assume $\mathcal{C}$ computably presented unitary fusion, $O = \mathbf{1} \oplus X$, dyadic $V_L$, $H_L = \sum_Z h_{L,Z}$ with $\operatorname{diam} Z \le R$, $\sup_x \sum_{Z \ni x} \|h_{L,Z}\| \le J$; no gap, uniqueness or translation invariance. Conjecture: computable $t_0, \mu, C_0, D_0$ depending only on $(\mathcal{C}, O, R, J)$ exist such that, uniformly in $L$ and $0 \le t \le t_0$: (1) $W_L(t) = U_L(t) V_L$, $\dot U_L = A_L U_L$, $A_L = \sum_I a_{L,I}$ anti-Hermitian and charge-preserving with $\sup_x \sum_{I \ni x} |I| e^{\mu \operatorname{diam} I} \|a_{L,I}\| \le C_0 J$; (2) $W_L^\dagger H_L W_L$ has a coarse interval decomposition with a like bound, scalars kept local; (3) $U_L(t)$ has a charge-preserving circuit of depth $\le D_0$, gate range $\le \min(2L, \lceil C_0 \log((2L+1)/\varepsilon) \rceil)$ and norm error $\varepsilon$. **Reviewer verdict (D3 REVIEW:54, INTEGRATION:76–84).** 76.9 is the first-order step; an all-orders proof needs a quasi-local replacement for $P_t$; short-imaginary-time analyticity or cluster expansion is the plausible route, but no such source is local, so nothing is asserted. D3-C is a *UV locality lemma*: by 76.11 a fixed-$t$ near-identity dressing cannot by itself drive $q_n$ or $\epsilon_n/\Delta_{n+1}$ to zero summably for a critical vacuum. First attackable lemma: a locally summable generator for the $t^2$ coefficient, via (6.3) and pair locality.

**Negative 76.11 [derived; confirmed].** (a) If $[H, P] = 0$, then $e^{-tH} V = V e^{-tV^\dagger HV}$ and $W(t) = V$ for all $t$: heat filtering inside the retained carrier cancels in the polar normalisation. (b) Normalised imaginary-time filtering of a vector with ground overlap $b > 0$ leaves excited weight $\le e^{-2t\Delta}(1-b^2)/b^2$; a fixed short time does not beat a closing gap. (c) (D3-6) A fixed UV shell gap with vanishing shell coupling does not select the IR vacuum: retain $\operatorname{span}\{e_1, e_2\}$ in $H(t) = \left(\begin{smallmatrix} 0 & 0 & 0 \\ 0 & t^2/2 & t \\ 0 & t & 1 \end{smallmatrix}\right)$, $0 < t < 1$. The shell gap is 1 and the coupling $t \to 0$, yet the lower $2 \times 2$ block has determinant $-t^2/2 < 0$, so the unique ground state is orthogonal to the retained ground vector $e_1$ (D3 checks `output.txt`:12). The missing condition is control of the induced metric and self-energy (D3 (7.2)–(7.3)) relative to the infrared scale. (d) (D4-V, D4 NOTE:465–482) Bare dyadic insertion compresses a nearest-neighbour fine energy to a sum of one-site coarse morphisms; if the coarse energy lies outside one-site sums plus scalars, the KS zero modes of (af) cannot be exactly covariant under bare insertion.

## 77.7 The Dichotomy: CA-75 Link (b3)

**Question 76.12 [question; reviewer reading, D3 REVIEW:93–101].** Every drift route above is of one of two kinds. *(G) Global fidelity:* 76.1, 76.2, 76.6 control $d_{k,n}$ through $q_n$ or $\epsilon_n/\Delta_{n+1}$, which see the global $P_n$ and carry an $O(N_n)$ volume factor; with $\Delta_n \gtrsim a_n^z$, summability needs local density mismatch $o(a_n^{1+z})$ (76.8). At fixed local accuracy the source *expects* zero overlap at infinite volume (`latticefermions.tex`:478–479), so dressing accuracy must improve with $n$ [expectation, not proved]. *(L) Local unital lift:* the global-fidelity-free route (D3-3, conditional-ok, D3 NOTE:226–260) bounds $d_{k,n}$ by local witness energies, a full-ball compression constant and a localisation error, but needs a local unital lift (76.2's boundary excludes the augmentation) and state-adapted witnesses (natural ones are expected to be vacuous at criticality, D3 REVIEW:62). **Link (b) status:** (b2) answered by 76.1, with no bound by $q_n$ alone; (b1) answered globally by 76.2, locally by CA-90 Theorem 90.1 (L0+half-braiding); **(b3) open**: does a declared dressing family achieve (G) at rate $o(a_n^{1+z})$, or does a local unital lift with state-adapted witnesses achieve (L)? No route here is presently non-vacuous for a critical family.

**Interface [open].** All statements use open chains with declared endpoints, while LC-4's KS tier is periodic (`reconstruction/04_cutoff_oar.md`:22–24, (af)). LC-4 can consume them only via an open-chain KS tier or a periodic version of this shard.

**What this does not claim.** No critical vacuum, gap, coupling selection, local unital lift (braided or not), proof of D3-C, KS convergence, projective-unitary symmetry or sector fidelity is constructed. Corrections 76.A–C qualify CA-72 and CA-73 without editing them. 76.5 is a proposal here; the local lift is derived in CA-90.

| Instance (calibration only) | Calibrates | Boundary |
|---|---|---|
| $M_2 \to M_4$ corner (no category) | 76.A norm 1 at zero GNS seminorm; 76.1 sharp at 32/25 | linear algebra only |
| $\mathcal{C} = \mathrm{Vec}$, $O = \mathbf{1} \oplus \mathbf{1}$ | global pins, factor 2, polar tangent (D3 checks) | site flip exists only here |
| pointed $\mathbb{Z}/3$, $O = \mathbf{1} \oplus g \oplus g$ | 76.9 on a constrained, non-self-dual, multiplicity-two carrier | first order only |
| pointed braided instance | 76.5: unital, extends $\theta$, range 3, $L = 2, 3$ | blind to braiding phases |
| free-fermion Wilson example | fixed-row computable dyadic tail (D3 NOTE:745) | nonlocal; not uniform in row |

# 78 Pointed Carriers: Occupation Grading, Dual-Pair Blocking, and Corner Qualifications

## 78.1 Status, Sources, and Dependencies

**Status: derived, with negative statements and one conjecture.** The numbered statements are local derivations of `explorations/2026-10-22-astra/D1-carrier-theory/NOTE.md`, graded *confirmed* by the sceptical review `REVIEW.md` in that directory (REVIEW.md:161–177) unless labelled otherwise; Proposition 77.2 is correct but near-tautological (REVIEW.md:165). The collision-sewing statement is **conjectural** (conditional-ok, overstated framing; REVIEW.md:163), not promoted. Finite evidence: `checks/carrier_checks.py` (4 groups GREEN, 4 mutations RED; REVIEW.md:21–24) and `checks/review_contact_check.py`. General $\mathcal{C}$ throughout; models only in the closing table (AGENTS.md Rule 16).

Dependencies: CA-65, CA-66, CA-68, CA-69, CA-71–CA-73, CA-75, CA-81, CA-87 (Lemma 87.1); CONVENTIONS (a), (r), (t), (u); the pointed-object amendment (am) of (r), stated below.

## 78.2 Setting and the Two Towers

Let $(O,\eta)$ be a *pointed* object: $\eta : \mathbf{1} \to O$, $\eta^\dagger\eta = 1$, $p_0 = \eta\eta^\dagger$, and $\iota : X \to O$ an orthogonal splitting of $p_1 = 1 - p_0$, so $O \cong \mathbf{1} \oplus X$ (Penneys:219–237). $O$ is *connected* if $\mathrm{Hom}(\mathbf{1}, X) = 0$. Put $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$, $H_{L,c} = \mathrm{Hom}(c, O^{\otimes L})$, $\alpha_L(a) = a \otimes \mathrm{id}_O$; for an order-preserving injection $\varphi : [k] \to [l]$ (CONVENTIONS (t)) let $V_\varphi : O^{\otimes k} \to O^{\otimes l}$ insert $\eta$ outside the image, $\theta_\varphi(a) = V_\varphi a V_\varphi^\dagger$, $P_\varphi = V_\varphi V_\varphi^\dagger$. $d_O$ is the positive spherical dimension (Penneys:1015–1019; EGNO:1700–1706), $\tau_L = \mathrm{Tr}_{\mathcal{C}}/d_O^L$. **Convention (amends (r); CONVENTIONS (am), 2026-10-22).** The site is the pointed pair $(O,\eta)$; $X$ may contain unit summands, and (r)'s species object is the connected case. Configuration spaces are ordered, with no $N!$ and fibres $\mathrm{Hom}(c, X^{\otimes N})$ (NOTE.md:75–79, :86–90; REVIEW.md:51).

**Proposition 77.1 [derived] (D1-T1, NOTE.md:120–151).** $\alpha$ is a unital injective $*$-map; $V$ is a coherent isometric placement functor, $V_\psi V_\varphi = V_{\psi\varphi}$; $\theta_\varphi$ is an injective $*$-map onto the corner $P_\varphi \mathcal{A}_l P_\varphi$ with $\theta_\varphi(1) = P_\varphi$. The unital tower has the fusion-graph incidence $m_d(L{+}1) = \sum_c m_c(L) N_{c,O}^d$ (CA-65); the corner tower has block maps $a_c \mapsto J_c a_c J_c^\dagger$ and not that incidence. A pointed unitary $u$ ($u\eta = \eta'$) intertwines both towers via $\mathrm{Ad}\, u^{\otimes L}$. *Proof sketch.* Compressing $a \otimes \mathrm{id}_O$ with $\mathrm{id} \otimes \eta$ returns $a$; $V_\varphi^\dagger V_\varphi = 1$ by cancelling each $\eta^\dagger\eta$, and composites by coherence (Penneys:394–398); $B \mapsto V_\varphi^\dagger B V_\varphi$ inverts $\theta_\varphi$ on the corner.

**Proposition 77.2 [derived; near-tautological] (D1-T2, NOTE.md:153–182).** All $\mathcal{A}_L$, $\theta_\varphi$, concatenation and the partial isometries $V_\varphi V_\psi^\dagger$ reconstruct the Cauchy completion (Penneys:358–362) of the subcategory generated by $O$, hence $\mathcal{C}$ if $O$ tensor-generates it, up to pointed unitary monoidal equivalence. Concatenation already carries the monoidal structure (REVIEW.md:113–116); the content is the negative lesson that algebras plus $\theta$ do not suffice. Braiding and twist must be marked separately.

**Proposition 77.3 [negative] (D1-N1, NOTE.md:184–197).** Along any nested placement schedule with lengths $\to \infty$, $\varinjlim(\mathcal{A}_L, \theta) = \bigoplus_c K(H_{\infty,c})$ over reachable charges: the unmarked corner limit retains only block Hilbert dimensions.

## 78.3 Occupation Grading and Parity

Let $Q_{L,n}$ sum the occupation-pattern projections with $n$ factors $p_1$, and $Z_{L,n} = \bigoplus_{|S|=n} X^{\otimes n}$.

**Proposition 77.4 [derived] (D1-T3, NOTE.md:208–247; counts re-derived at REVIEW.md:123–125).**

1. $\mathcal{A}_L = \bigoplus_{m,n} \mathrm{Hom}(Z_{L,n}, Z_{L,m})$; $\mathcal{A}_L^{(k)} = \bigoplus_{m-n=k} Q_{L,m}\mathcal{A}_L Q_{L,n}$ has $\mathcal{A}_L^{(k)}\mathcal{A}_L^{(l)} \subseteq \mathcal{A}_L^{(k+l)}$, $(\mathcal{A}_L^{(k)})^\dagger = \mathcal{A}_L^{(-k)}$. Each block is an $(\mathrm{End}\, Z_m, \mathrm{End}\, Z_n)$-bimodule, in charge $c$ of size $\binom{L}{m} \dim \mathrm{Hom}(c, X^{\otimes m}) \times \binom{L}{n} \dim \mathrm{Hom}(c, X^{\otimes n})$; $\hat{N}_l V_\varphi = V_\varphi \hat{N}_k$.

2. With $s = \dim \mathrm{Hom}(\mathbf{1}, X^{\otimes 2})$, $t = \dim \mathrm{Hom}(X, X^{\otimes 2})$, $z = \dim \mathrm{Hom}(\mathbf{1}, X)$: $\dim \mathcal{A}_2^{(+2)} = s$, $\dim \mathcal{A}_2^{(+1)} = 2t + 2z$, so the number-changing part of $\mathcal{A}_2$ has dimension $2s + 4t + 4z$, i.e. $2s + 4t$ for connected $O$. The number-preserving part is not an ideal and its complement is only a bimodule.

3. $\mathcal{A}_L^{(k)} = 0$ for all odd $k$ and all $L$ (parity preservation) iff $\mathrm{Hom}(X^{\otimes m}, X^{\otimes n}) = 0$ for all $m, n \geq 0$ with $m - n$ odd.

⟨1⟩**1.** Distribute $O^{\otimes L}$ over $\mathbf{1} \oplus X$ and insert $1 = \sum_n Q_{L,n}$ on both sides (Penneys:219). For two sites $Z_0 = \mathbf{1}$, $Z_1 = X \oplus X$, $Z_2 = X^{\otimes 2}$, giving item 2.

⟨1⟩**2.** Necessity in 3: a nonzero $f : X^{\otimes n} \to X^{\otimes m}$ sits in the occupied corner of $\mathcal{A}_{\max(m,n)}$ as an odd-degree element. Sufficiency: every block of ⟨1⟩1 is such a Hom space. QED.

Testing only $t = 0$ does not establish the all-scale condition; the CA-69 parity-even diagram subalgebra can be proper for this structural reason (NOTE.md:243–247).

## 78.4 Dual-Pair Blocking Is Decidable

**Theorem 77.5 [derived] (D1-T4, NOTE.md:251–287; REVIEW.md:117–122).** Write $[O] = \sum_c o_c[c]$ ($Y$ is a blocking root as in CA-87). There is $Y \in \mathcal{C}$, not necessarily simple, with $O \cong Y \otimes \bar{Y}$ iff the integer system

$$o_c = \sum_{a,b} y_a y_b N_{a,\bar{b}}^c \quad (c \in \mathrm{Irr}\,\mathcal{C}), \qquad \sum_a y_a^2 = o_{\mathbf{1}}, \qquad y_a \in \mathbb{Z}_{\geq 0}$$

is solvable. The search is finite. If it succeeds, the unitary $u : O \to Y \otimes \bar{Y}$ can be chosen with $u\eta = d_Y^{-1/2}\mathrm{coev}_Y$, so the vacancy projector becomes the normalised cup projection $up_0u^\dagger = d_Y^{-1}\mathrm{coev}_Y\mathrm{coev}_Y^\dagger$. **Assume** the fusion coefficients and dagger. **Prove** criterion, finiteness, transport.

⟨1⟩**1.** For $Y = \bigoplus_a y_a a$, the multiplicity of $c$ in $Y \otimes \bar{Y}$ is the first sum. By distributivity.

⟨1⟩**2.** $N_{a,\bar{b}}^{\mathbf{1}} = \delta_{ab}$ gives the second equation; hence $y_a \leq \lfloor \sqrt{o_{\mathbf{1}}} \rfloor$ and exhaustive search terminates, with exact integer arithmetic. By Penneys:427–430, :484–487.

⟨1⟩**3.** Equal multiplicities give an isomorphism; polar decomposition makes it unitary. By semisimplicity, Penneys:233.

⟨1⟩**4.** $U(o_{\mathbf{1}})$ acts transitively on unit vectors of $\mathrm{Hom}(\mathbf{1}, Y\bar{Y})$, so $u$ may be corrected on the unit-isotypic block to send $\eta$ to $d_Y^{-1/2}\mathrm{coev}_Y$, a unit vector since $\mathrm{coev}_Y^\dagger\mathrm{coev}_Y = d_Y$ (CONVENTIONS (r)). QED.

**Remarks [derived].** (i) For connected $O$, $o_{\mathbf{1}} = 1$ forces $Y$ simple, the hypothesis of Lemma 87.1; Theorem 77.5 strictly generalises it by the converse, decidability, non-simple $Y$ with multiplicities, and transport of $\eta$. Self-duality of $O$ and $\operatorname{FPdim} O = (\operatorname{FPdim} Y)^2$ are necessary only. (ii) After blocking, range $k$ on $O$-sites is range $2k$ on the alternating $Y, \bar{Y}$ chain, vacancy insertion is insertion of a normalised cup pair, and occupation counts pairs outside that channel. (iii) Failure excludes only dual-pair blocking inside $\mathcal{C}$, not other words $Y_1 \otimes \cdots \otimes Y_k$, enlarging $\mathcal{C}$, module-category dualities or equal universality classes; success selects no Hamiltonian (NOTE.md:281–287).

**Question (REVIEW.md:99–108; unchecked).** Over alternating $O\bar{O}$ words the projections $d_O^{-1}\operatorname{coev}\operatorname{coev}^\dagger$ match the cup Jones projections of Hollands:1240–1271, and after blocking $p_0$ is one; is the unital tower a basic construction?

**Proposition 77.6 [negative; strengthened] (D1-N2, NOTE.md:289–313; REVIEW.md:109–112).** A bare $O$ is not an algebra object (FRS:2272–2284). $O \otimes \bar{O}$ carries the matrix algebra structure of FRS:2331–2336, and by FRS:2913–2920 all $U \otimes U^\vee$, including $U = \mathbf{1}$, are Morita equivalent [source-local; the authors defer the proof]. Hence the Morita class of $O \otimes \bar{O}$ is *the same for every nonzero* $O$ and carries no information about the carrier. Independently [derived], $O$ and $O \oplus O$ have equivalent module categories but $\dim \operatorname{End}$ differing by four.

## 78.5 Vacuum Insertion, States, and Qualifications of CA-71–CA-73

**Proposition 77.7 [derived] (D1-T5, D1-T5b, NOTE.md:317–336, :377–395).**

1. For every $T$, $f \mapsto V_\varphi f$ is a bijection $\operatorname{Hom}(T, O^{\otimes k}) \to \{f : T \to O^{\otimes l} : P_\varphi f = f\}$, natural in $T$.

2. For a state $\omega$ with $p = \omega(P_\varphi) > 0$, the normalised corner pullback has GNS vector $\pi(P_\varphi)\Omega/\sqrt{p}$, at squared distance $2(1 - \sqrt{p})$ from $\Omega$, and $1 - p \le 2(1 - \sqrt{p}) \le 2(1 - p)$. $\sum_n \sqrt{1 - p_n} < \infty$ suffices for convergence of transported vacuum vectors.

3. $\tau_l(P_\varphi) = d_O^{k-l}$, invariant under unitary dressing; trace states cannot approach weight one under extensive insertion when $d_O > 1$.

4. The CA-72 unital completion has multiplicativity defect $\operatorname{cov}_\chi(S, T)(1 - P_\varphi)$, of operator norm $|\operatorname{cov}_\chi(S, T)|$ whenever $P_\varphi \neq 1$, independently of $\omega(P_\varphi)$.

*Proofs.* (2) norm one, overlap $\sqrt{p}$; (3) the categorical trace of an isometry's range projection is its domain's dimension; (4) multiply out (REVIEW.md:126).

**Qualifications [derived; REVIEW.md:127–130, :173].** (a) CA-72 Corollary 72.8 reads GNS-seminorm smallness as the soft-system condition, but eq. (29) of 2306.16063:568–575 is an operator-norm condition; by 77.7(4) the seminorm is not that norm. (b) The right caret $\mathrm{id}_O \otimes \iota_{\mathbf{1}}$ never inserts before the first strand, so it does not generate placements with $\varphi(1) > 1$; the CA-71:167–169 generation sentence is false as stated. (c) $\bigoplus_c \operatorname{Hom}(c, O^{\otimes L})$ is not $\Phi(1)^{\otimes L}$ in Hilb, so the Thompson-$V$ extension of B–S:522–541 is unavailable (CA-71:173–174 "indeed $V$" unsupported); the $F$-action of B–S:500–511 survives. (d) $\theta_\varphi$ pins *every* inserted site and sends constants to $P_\varphi$: $\theta(F(a; 1)) = F(\theta(a); P_\varphi)$, qualifying CA-73:159–172. (e) The isometry deficit read at CA-68:160–162 arises for overlapping smearing under hard-core exclusion; box cells have none (Theorem 77.8).

## 78.6 The Configuration Carrier

For a cell split into children of lengths $\ell, \ell'$ let $R : O \to O^{\otimes 2}$, $R\eta = \eta \otimes \eta$, $R\iota = \sqrt{\ell/(\ell+\ell')}\,(\iota \otimes \eta) + \sqrt{\ell'/(\ell+\ell')}\,(\eta \otimes \iota)$, i.e. $R = U(\mathrm{id}_O \otimes \eta)$ with a real rotation $U$ on $X\eta \oplus \eta X$. With $\Delta_N = \{0 < x_1 < \cdots < x_N < 1\}$, set $\mathcal{F}_{\mathcal{C},X} = \widehat{\bigoplus}_{c,N} L^2\big(\Delta_N; \operatorname{Hom}(c, X^{\otimes N})\big)$.

**Theorem 77.8 [derived] (D1-T6, NOTE.md:399–455; REVIEW.md:131–135).** Along nested partitions of $(0,1)$ with mesh $\to 0$, the $R$ are exact charge- and number-preserving isometric refinements, coherent under all parenthesisations (amplitudes telescope to $\sqrt{|J|/|I|}$), and the box maps $K_\pi(t \text{ at } j_1 < \cdots < j_N) = \prod_r |I_{j_r}|^{-1/2} \mathbf{1}_{I_{j_r}}(x_r)\, t$ extend to a unitary from the Hilbert inductive limit onto $\mathcal{F}_{\mathcal{C},X}$. For adjacent intervals, $\mathcal{F}_c(I \cup J) \cong \widehat{\bigoplus}_{a,b} \mathcal{F}_a(I) \otimes \mathcal{F}_b(J) \otimes \mathrm{Hom}(c, a \otimes b)$: fusion fibres are retained, not replaced by onsite factors; no braiding is used. **Caveat:** $\mathrm{Ad}\, R$ is again a corner map; this is a Hilbert carrier theorem, not a unital observable refinement.

**Proposition 77.9 [negative] (D1-N4, NOTE.md:464–473).** Bare dyadic insertion has an atomic limit: $\ell^2$ of a countable position set with fusion fibres; moving a vector to a nearby point changes it by norm $\sqrt{2}$, so the geometric translations are not strongly continuous.

## 78.7 Conjecture, with Caveat

D1-C (NOTE.md:9–62) conjectures that explicit nearest-neighbour $H_\varepsilon$, built from $D_{\mathrm{hop}} = \pi_{X0} - \pi_{0X}$ and declared reaction morphisms $b_\alpha : X \to X \otimes X$, $v_\beta : \mathbf{1} \to X \otimes X$ (NOTE.md:35–41), converge in projected-resolvent sense on $\mathcal{F}_{\mathcal{C},X}$ to a sector-coupling collision form, under declared closedness and core hypotheses. **Status: conjectural; framing overstated** (REVIEW.md:163). The reviewer found (REVIEW.md:78–91) that $D_{\mathrm{hop}}^\dagger D_{\mathrm{hop}}$ equals standard hard-core hopping plus a contact attraction $-2\varepsilon^{-2} n_j n_{j+1}$: it engineers Neumann collisions (two-particle $E_0 = 0$, $E_1 = 9.8673$ at $L = 60$), whereas standard hopping gives Dirichlet collisions ($E_0 = 9.8673$; $\pi^2 = 9.8696$) (`checks/review_contact_output.txt`:4–7), under which all traces vanish and the number-changing coupling disappears. The limit is thus non-universal, sees $\mathcal{C}$ only through finite fibres and two linear maps (REVIEW.md:92–98), and is a nonrelativistic ($z = 2$) benchmark with no conformal, Wilson or fidelity content. **Amendment (2026-10-22, round 2).** D1-C is proved for the tuned family, without its closedness and core hypotheses, in CA-94 (Theorem 94.6, Lemmas 94.3–94.5); the limit changes $\hat{N}$ iff $\sum_\alpha \lambda_\alpha b_\alpha \neq 0$ or $\sum_\beta \kappa_\beta v_\beta \neq 0$ (Corollary 94.7). The caveat stands: the tuning is necessary (Proposition 94.1), and the limit is an off-ladder $z$=2 resolvent limit, not R1.

**Open problem [Question] (INTEGRATION.md:103–106, :133–136).** On $\mathcal{F}_{\mathcal{C},X}$, construct coherent local *unital* refinements of a number-changing field algebra (a categorical second quantisation), compatible with one state family satisfying WT-1 (CA-81). Corner maps do not solve this.

## 78.8 What This Does Not Claim

No dynamics, critical point, unital OAR map, continuum CFT or fidelity statement is made. Occupation $\hat{N}$ is microscopic, not a CFT particle number; no argument identifies the empty configuration with a conformal vacuum (NOTE.md:475–479). The Jones-tower identity is a Question. *Amendment 2026-10-22:* D1-T7 and the D1-C analytic gap are closed for the tuned family in CA-94 (Lemma 94.5, Theorem 94.6), with obstructions 94.1, 94.9, 94.11; that is a $z$=2 resolvent limit, not a CFT.

## 78.9 Examples (calibration only)

| Instance | Calibration | Local evidence |
|---|---|---|
| Fibonacci, $O = \mathbf{1} \oplus \tau$ | connected; $\tau \otimes \tau = \mathbf{1} \oplus \tau$ solves Theorem 77.5 with $Y = \tau$ | Penneys:610–631; CA-87:71–75 |
| Haagerup $H_3$, $X = \rho$ | connected, so $Y$ simple; dimension test fails: no dual-pair blocking | CA-87:78–81 |
| $\mathrm{Hilb}_{\mathrm{fd}}$, $O = \mathbb{C}^2, \mathbb{C}^3$ | both matrix objects Morita trivial, End differ; 2 not a square, so no root | FRS:2331–2336, :2913–2920 |
| $\mathrm{Vec}_G$, $O = \bigoplus_g g$, $\lvert G\rvert = r$ | $m_g(L) = r^{L-1}$: $C_4$, $C_2 \times C_2$ have identical unmarked towers | Penneys:277–288, :522 |

# 79 Finite Selection Equations for a Categorical Nearest-Neighbour Hamiltonian

## 79.1 Status, Sources, and Dependencies

**Status: mixed, labelled per statement** (derived 78.1–78.8, with exact SymPy checks; conditional 78.9–78.11; conjectural 78.12; proposals 78.13 and the ranking). Origin: D2 in `explorations/2026-10-22-astra/D2-selection-principle/`: N = `NOTE.md`, reviewed in R = `REVIEW.md`, I = `INTEGRATION.md`; review grades carried unchanged. No Julia or continuum run; porting the checks is open (I:67–68). Dependencies: CA-65, CA-72, CA-75 row (2), CA-81 (WT-1, VR-1), CA-86, CA-87; CONVENTIONS (a), (b), (r), (af), (ak), (al), (ao).

## 79.2 Setting and the Coefficient Compiler

$\mathcal{C}$ is unitary fusion with positive spherical trace $\mathrm{tr}_{\mathcal{C}}$; $O \neq 0$ is the site object of (r); $B_2 = \mathrm{End}_{\mathcal{C}}(O^{\otimes 2})$, $B_3 = \mathrm{End}_{\mathcal{C}}(O^{\otimes 3})$. As in (ak), $H = -\sum_j h_j$, $h = h^\dagger \in B_2$; the physical bond energy fed to the KS tier (af) is $k_j := -h_j$ (N:58–61). $e_j$ denotes only the cup-cap tile of (ak). A spectral $R(u)$ is a regular operator on $O \otimes O$ with additive parameter, distinct from a braiding (N:66–70).

**Proposition 78.1 [derived; compiler].** For a basis $\{D_\alpha\}$ of $B_2$ and $h = \sum_\alpha c_\alpha D_\alpha$, the Gram matrix $\mathcal{G}_{\alpha\beta} = \mathrm{tr}_{\mathcal{C}}(D_\alpha^\dagger D_\beta)$ is positive definite, and $\tau^\alpha(X) = \sum_\beta (\mathcal{G}^{-1})_{\alpha\beta} \mathrm{tr}_{\mathcal{C}}(D_\beta^\dagger X)$ satisfy $X = \sum_\alpha \tau^\alpha(X) D_\alpha$. Constants: $M^\gamma_{\alpha\beta} = \tau^\gamma(D_\alpha D_\beta)$ and coordinates $\sigma^\mu$ (same construction on $B_3$) of words in $A_\alpha = D_\alpha \otimes 1$, $B_\alpha = F^\dagger(1 \otimes D_\alpha)F$, $F$ the associator. **Proof.** $B_2 \cong \bigoplus_a M_{m_a}(\mathbb{C})$ and $\mathrm{tr}_{\mathcal{C}}$ is $d_a \,\mathrm{Tr}$ on block $a$, $d_a > 0$ (Penneys :1010–1019): faithful and positive. □ No self-duality, multiplicity-freeness or FS sign is used (contrast AFM :175–184).

**Proposition 78.2 [derived; exact gate ODE, upgrading Lemma 86.1].** Write $e^{th} = \sum_\alpha w_\alpha(t) D_\alpha$. Then

$$w_\alpha(0) = \tau^\alpha(1), \qquad \dot{w}_\gamma = \textstyle\sum_{\alpha\beta} c_\alpha w_\beta M^\gamma_{\alpha\beta}, \tag{78.1}$$

and every finite ordered product of gates $e^{\delta\tau h_j}$, in particular $U(\delta\tau)^M$ of CA-86, equals *exactly* $\sum_\Gamma \prod_p w_{\alpha(p)}(\delta\tau)\,\mathrm{Ev}_{\mathcal{C}}(\Gamma)$, one basis tile per plaquette. **Proof.** $\frac{d}{dt}e^{th} = he^{th}$; expand, apply $\tau^\gamma$. Substitute per plaquette; multilinearity and composition in $\mathcal{A}_L$ give $\mathrm{Ev}_{\mathcal{C}}$ (CA-86:89–92). □ Lemma 86.1's remainder disappears; only the Trotter comparison remains (R:101–103).

## 79.3 Blocking, Leakage, and Metric Response

Contracting a fixed $2 \times 2$ patch $P$ gives a degree-four map $b_\beta(w) = \sum C^\beta_{\alpha_1\ldots\alpha_4} w_{\alpha_1}\cdots w_{\alpha_4}$, $C^\beta = \tau^\beta_{\mathrm{new}}(\mathrm{Ev}_{\mathcal{C}}(P[D_{\alpha_1},\ldots,D_{\alpha_4}]))$, into tiles with $O^{\otimes 2}$ boundaries, not back into $B_2$. With a retained isometry $I$, $\pi = I^\dagger$: $\mathcal{B}_P = Ib + L$, $L = (1 - I\pi)\mathcal{B}_P$; a closed fixed point solves $b(w^*) = \lambda w^*$, $\ell(w^*) = 1$, $L(w^*) = 0$ (N:140–199).

**Lemma 78.3 [derived; D2-L1, confirmed].** Assume finite categorical spaces, fixed $P$, the isometries above and $V : B' \to B \otimes B$ isometric. Prove (i) $I\pi$ is the orthogonal projection onto retained tiles; (ii) if $X = V(V^\dagger X V)V^\dagger$ for every boundary endomorphism $X$, $V$ is unitary; (iii) lossless retention of the same nonzero $O$ on every boundary forces $d_O = 1$.
⟨1⟩1. $I^\dagger I = 1$ gives (i). ⟨1⟩2. $X = 1$ gives $VV^\dagger = 1$; with $V^\dagger V = 1$, (ii). ⟨1⟩3. Then $O \otimes O \cong O$, $d_O^2 = d_O$, $d_O > 0$. ⟨1⟩4. QED.
So for $d_O \neq 1$ no fixed-carrier compression of *all* boundary endomorphisms is lossless: blocking leaks. Closure of special tensors is not excluded; $\mathrm{Ev}_{\mathcal{C}}$ is compatible with blocking, the weights flow.

**Lemma 78.4 [negative, derived; D2-L2, confirmed].** Assume an unnormalised generating functional $W(g)$, smooth and local in the metric, unchanged by every compactly supported metric

variation for all fixed probes; stress is its metric derivative. PROVE all separated stress insertions vanish. ⟨1⟩1. Differentiate once: zero. ⟨1⟩2. Differentiate at disjoint supports; locality excludes contact terms. ⟨1⟩3. QED.
The target is Euclidean covariance with nontrivial metric response, not Levin–Wen isotopy (N:239–245). The Levin–Wen principle selects a topological fixed point, so CA-86:57 is no selection precedent for a critical chain (R:166–170).

### 79.4 Additive Yang–Baxter as a Coefficient System

For $R(u) = \sum_\alpha r_\alpha(u) D_\alpha$, additive YBE (precedent 1408.1282:159)

$$R_{12}(u)R_{23}(u{+}v)R_{12}(v) = R_{23}(v)R_{12}(u{+}v)R_{23}(u) \tag{78.2}$$

reads $\sum[\sigma^\mu(A_\alpha B_\beta A_\gamma) r_\alpha(u) r_\beta(u{+}v) r_\gamma(v) - \sigma^\mu(B_\alpha A_\beta B_\gamma) r_\alpha(v) r_\beta(u{+}v) r_\gamma(u)] = 0$ for all $\mu$: cubic *functional* equations, finite only after a fixed ansatz (N:253–267).

**Theorem 78.5 [derived and checked; D2-L3, confirmed; cubic cokernel test].** ASSUME $R$ analytic near 0, $R(0) = 1$, $R'(0) = h$, (78.2) for small $u, v$. PROVE there is $K \in B_2$ with

$$[h_{12} + h_{23}, [h_{12}, h_{23}]] = K_{23} - K_{12}. \tag{78.3}$$

⟨1⟩1. Write $R = 1{+}uh{+}u^2a{+}u^3b{+}\cdots$, $A = h_{12}$, $B = h_{23}$. The $uv$ coefficient is $2(a_{23}{-}a_{12}){+}A^2{-}B^2$, so $q_{12} = q_{23}$ for $q = 2a - h^2$.
⟨1⟩2. $q_{12} = q_{23}$ forces $q$ scalar. ⟨2⟩1. Left partial trace over factor 1, right over factor 3: $d_O q = \mathrm{tr}_1(q) \otimes 1 = 1 \otimes \mathrm{tr}_2(q)$. ⟨2⟩2. The right partial trace of $x \otimes 1 = 1 \otimes y$ gives $d_O x = \mathrm{tr}(y)1$; so $x$, hence $q$, is scalar. ⟨2⟩3. QED.
⟨1⟩3. Multiplying $R$ by the scalar $1 - \kappa u^2/2$ ($q = \kappa 1$) preserves (78.2) and $h$, and makes $a = h^2/2$.
⟨1⟩4. With $r_3 = R'''(0)$, the $u^2v$ coefficient is $(K_{23} - K_{12} - [A{+}B, [A, B]])/2$, $K = r_3 - h^3$ (free words, `exact_checks.py`:69–81). ⟨1⟩5. QED.
Only pivotal structure and $d_O \neq 0$ are used (R:107–112). With $\mathcal{D} : K \mapsto K_{23} - K_{12}$, projecting (78.3) onto $\mathrm{coker}\,\mathcal{D}$ gives finitely many cubic equations in $c$ with constants from Prop. 78.1. This *rejection test* is necessary for analytic additive YBE, neither sufficient (higher jets) nor a criticality test. The review identifies (78.3) as the classical Reshetikhin condition; **local source missing**, so no attribution is made.

**Lemma 78.6 [derived as algebra; D2-L4, confirmed].** ASSUME $e_i^2 = d\,e_i$, $e_1e_2e_1 = e_1$, $e_2e_1e_2 = e_2$, $X_i(u) = 1 + x(u)e_i$. PROVE the residual $X_1(u)X_2(u{+}v)X_1(v) - X_2(v)X_1(u{+}v)X_2(u)$ equals $(x + y + dxy + xyz - z)(e_1 - e_2)$, where $x = x(u)$, $y = x(v)$, $z = x(u{+}v)$.
⟨1⟩1. The left side is $1 + (x{+}y{+}dxy{+}xyz)e_1 + ze_2 + xz\,e_1e_2 + yz\,e_2e_1$. ⟨1⟩2. The right side swaps the $e_1, e_2$ coefficients only; subtract. QED.
For $d = 2\cos\lambda$, $\sin\lambda \neq 0$, $x(u) = \sin u/\sin(\lambda - u)$ solves it (`exact_checks.py`:51–58; precedent 1608.04927:453–458). *Applicability* (R:113–117): a TL pair in $B_2$ needs a cup/cap on $O \otimes O$, i.e. a self-dual summand $X$ of $O$, whose FS indicator $\kappa_X$ of (r) fixes the sign relating $d$ to $d_X$; the lemma does not supply the representation, and exhausts $B_2$ only if $B_2 = \mathrm{span}\{1, e\}$. On-site solutions $\exp(u(p \otimes 1 + 1 \otimes p))$ lie outside $\mathrm{span}\{1, e\}$ when $\mathrm{End}_{\mathcal{C}}(O) \neq \mathbb{C}$: more solutions need not mean better dynamics (N:327–334).

**Proposition 78.7 [negative, derived and checked; D2-L5 with repaired converse].** (i) With $e_1, e_2$ as above and $\mathsf{B}(x) = (1 + xe_1)(1 + xe_2)$, $[\mathsf{B}(x), \mathsf{B}(y)] = xy(y - x)(e_2 - e_1)$: local YBE gates do not make an open brick sweep commute. (ii) If $\mathrm{End}_{\mathcal{C}}(O) \neq \mathbb{C}$, take a nontrivial projection $p$, $h = p \otimes 1 + 1 \otimes p$, $g = p \otimes p$, $X(u) = \exp(uh + u^2 g)$. All gates commute, so every brick or diagonal product commutes for all $u$; yet no real-analytic $\varphi \not\equiv 0$, $\varphi(0) = 0$, makes $\exp(\varphi h + \varphi^2 g)$

satisfy (78.2).
⟨1⟩1. (i): $\mathsf{B}(x) = 1 + x(e_1{+}e_2) + x^2 e_1 e_2$ and $[e_1{+}e_2, e_1 e_2] = e_2 - e_1$ (the $d\,e_1 e_2$ terms cancel); expand. RED mutation: `run.txt`:20–29.
⟨1⟩2. (ii): one-site projections commute, so both sides are exponentials of commuting self-adjoint sums differing by $(a{+}b{-}c)(h_{12} - h_{23}) + (a^2{+}b^2{-}c^2)(g_{12} - g_{23})$, $a = \varphi(u)$, $b = \varphi(v)$, $c = \varphi(u{+}v)$.
⟨1⟩3. The differences are linearly independent and exp is injective on commuting self-adjoints, so $a + b = c$, $a^2 + b^2 = c^2$, hence $ab = 0$; at $u = v$, $\varphi \equiv 0$ (`review_checks.py`:21–31). ⟨1⟩4. QED.
The original witness $\exp((u{+}u^2)\,p \otimes p)$ was a reparametrisation artefact (graded *overstated*, R:124–134); unused. Simple $O$: converse open.

**Erratum to CA-86 Proposition 86.3.** (a) Prop. 86.3 (CA-86:112–116) calls the brick sweep $T(u) = \prod_{\text{odd}} X_j \prod_{\text{even}} X_j$ diagonal-to-diagonal and assumes it commutes "as the sourced $\hat{\tau}_D$ does". It is Koo–Saleur's *row-to-row* $\hat{\tau}$ (9312156.tex:603), which on the homogeneous self-dual line does *not* commute (:716–718); the commuting family is the sequential diagonal product $\hat{\tau}_D = \prod_j X_{2j} X_{2j+1}$ (:725). The attribution is wrong; by Prop. 78.7(i) YBE gates do not repair it. The conditional implication (commuting family $\Rightarrow [H, T(u)] = 0$) stands, and applied to the sourced commuting $\hat{\tau}_D$ (:729–738) the Taylor argument of CA-86:124–126 gives $[\sum_j e_j, \hat{\tau}_D(u)] = 0$. (b) "The ground state is a common eigenvector" (CA-86:120–122) needs a nondegenerate ground state: $[H, T] = 0$ only preserves eigenspaces. CA-86:146, :237–239 and CA-75 row (2) (:73) inherit both qualifications; whether a Trotter product equals a commuting transfer is open.

**Proposition 78.8 [derived and checked; braided linear ansatz].** Assume $\mathcal{C}$ braided, $\mathsf{b} = c_{O,O}$, $b = \mathsf{b}_{12}$, $b' = \mathsf{b}_{23}$, $R(u) = 1 + x(u)\mathsf{b}$. Prove the YBE residual is $(x{+}y{-}z)(b - b') + xy(b^2 - b'^2)$; if it vanishes where $xy \neq 0$, then $\mathsf{b}^2 - \alpha\mathsf{b} = \beta 1$; conversely that relation reduces (78.2) to $z = x + y + \alpha xy$.
⟨1⟩1. Expand; the cubic words cancel by $bb'b = b'bb'$ (`exact_checks.py`:84–89). ⟨1⟩2. Vanishing gives $b^2 - \alpha b = b'^2 - \alpha b'$, $\alpha = -(x{+}y{-}z)/xy$; apply ⟨1⟩2 of Theorem 78.5. ⟨1⟩3. Substitute. QED.
A unitary braid with more than two eigenvalues has no *linear* spectralisation. This limits the ansatz only: a local source Baxterises through BMW with a three-eigenvalue braid, nonlinearly in $G, G^{-1}$ (1408.1282:217–235). Unbraided $\mathcal{C}$ supplies no $\mathsf{b}$. **Remark [derived].** $R \equiv 1$ solves (78.2) with zero dynamics, so YBE implies neither criticality, nor a nonzero central charge, nor fidelity (N:391–397).

## 79.5 Defect Symmetry

Sliding a defect $a$ past $D_\alpha$ gives residuals $Q_{a,\alpha}$ and equations $\sum_\alpha c_\alpha Q_{a,\alpha} = 0$. If crossings are natural for *every* endomorphism of $O$ with coherent closure, $Q \equiv 0$: no equation on $c$ (N:470–485); only a larger non-categorical ansatz is cut down. Defect commutation is weaker than YBE (AFM :58–60).

**Lemma 78.9 [conditional; D2-L7, conditional-ok].** Assume a common limiting Hilbert system; defect and junction operators and adjoints converge strongly, uniformly bounded; finite products obey the categorical relations; junction reflected-norm Grams $\geq \eta 1$, $\eta > 0$. Prove the relations survive and independent junctions stay independent. ⟨1⟩1. $A_n B_n v - ABv = A_n(B_n - B)v + (A_n - A)Bv \to 0$. ⟨1⟩2. Induct over finite products and adjoints. ⟨1⟩3. The Gram bound keeps combinations nonzero. ⟨1⟩4. QED.
Faithful, *not full*: no fullness, distinct simple images or sector completeness. Symmetry survives in gapped phases and beside neutral spectators: it constrains the search space, fixes no critical point.

## 79.6 Interfaces to WT-1 and VR-1

**Lemma 78.10 [conditional; D2-L8, conditional-ok].** ASSUME $F, G$ positive nonzero functionals on a unital C*-algebra, $z = F(1)$, $w = G(1)$. PROVE $\|F/z - G/w\| \le 2\|F - G\|/\min(z, w)$. ⟨1⟩1. Split as $(F - G)/z + G(1/z - 1/w)$. ⟨1⟩2. $\|G\| = w$, $|z - w| \le \|F - G\|$. ⟨1⟩3. Triangle; QED.
If $d_{n+1} \le \rho d_n + \ell_n$, $0 \le \rho < 1$, $\sum \ell_n < \infty$, and unnormalised row functionals obey relative bounds $\le C_k d_n$, then $\sum_n d_n \le (d_0 + \sum \ell_n)/(1 - \rho)$ and the normalised drift is $\le 2C_k d_n$: WT-1 A3, given unital injective $*$-maps (A1). With CA-72 pullbacks of mass $1 - \varepsilon_n$ it feeds CA-75 Link (b2) given (b1) (R:171–175).

**Lemma 78.11 [conditional; D2-L10, conditional-ok].** ASSUME finite Hilbert spaces, output/input controls $K \ge 1$, a residual $M$, $s \in \mathbb{N}$, $0 \le \alpha < 1$, $\|K^s M e^{-tK}\| \le b\, t^{-s-\alpha}$ for $0 < t \le 1$. PROVE $\|K^s M K^{-(s+1)}\| \le C_{s,\alpha} b$, $C_{s,\alpha} = \frac{1}{s!}\big(\frac{1}{1-\alpha} + \int_1^\infty t^s e^{1-t} dt\big)$.
⟨1⟩1. $K^{-(s+1)} = \frac{1}{s!}\int_0^\infty t^s e^{-tK} dt$. ⟨1⟩2. On $(0, 1]$ the integral is $\le b/(s!(1 - \alpha))$. ⟨1⟩3. For $t \ge 1$, $e^{-tK} = e^{-K} e^{-(t-1)K}$ bounds the integrand by $b\, t^s e^{1-t}/s!$. ⟨1⟩4. QED.
This is the shape of VR-1 A2 (two losses: A4). **Counterexample [negative, derived]:** two-sided nuclear collar bounds do not suffice. For $E_n = \mathrm{diag}(0, \dots, n)$, $X_n = e^{\sqrt{n}}(|n\rangle\langle 0| + |0\rangle\langle n|)$: $\mathrm{Tr}\, e^{-tE_n} \le (1 - e^{-t})^{-1}$ and $\|e^{-tE_n} X_n e^{-tE_n}\|_1 = 2e^{\sqrt{n} - tn} \le 2e^{1/(4t)}$ uniformly in $n$, yet $\|X_n(1 + E_n)^{-p}|0\rangle\| = e^{\sqrt{n}}$ for every $p$.

## 79.7 The Sufficiency Conjecture and Its Attackable Core

**Conjecture 78.12 [conjectural; D2-C].** Let $O$ tensor-generate $\mathcal{C}$ and $h$ carry a geometric stress certificate (N:606–693): C1 analytic local gate families whose metric derivatives have time and mixed responses $k = -h$ and $i[k_j, k_{j+1}]$ of (af), one positive velocity; C2 a blocking hierarchy in a fixed Banach space, $d_{n+1} \le \rho d_n + \ell_n$, leakage kept; C3 separating positive boundary probes, $\sup_n \mathrm{Tr}\, e^{-tE_n} < \infty$, and the one-sided bounds of Lemma 78.11 for metric insertions; C4 discrete Ward contour identities with contact term and summable residuals; C5 coherent unital injective local $*$-refinements in one GNS system; C6 a nonzero separated connected stress two-point limit. Then the scaling limit is a local unitary positive-energy 2D conformal net whose stress is the limit of the standard KS modes of $k$ on a common core, generating projective unitary $\mathrm{Diff}(S^1)$ actions, with $c_{\mathrm{conf}}$ the Ward contact coefficient.

**Reviewer verdict: needs reformulation.** Sufficiency, not selection; category-blind (for $\mathcal{C} =$ Hilb, $O = \mathbb{C}^n$ every qudit chain is in scope); no instance of C1–C6 verified; the Ward-to-KS, domain, locality and integration steps are open (R:76–97, :198–202; N:695–701). Chiral-sector equality with $\mathcal{C}$ is not part of it. **Target 78.13 [proposal; Ward-to-KS transfer lemma].** Under C1 and C3, the discrete contour identity of C4 with summable residuals implies VR-1 A3/A4 for the standard KS modes of $k$ on the same GNS core (I:110–115). First test: C3 on the in-repo free-fermion benchmark (I:117–121). *Amendment (2026-10-22, round 2): split by CA-89.* A4 transfers from geometric modes given A2 (Theorem 89.1(i)); A3 needs differentiated, coordinate-resolved matching (89.1(ii)); the clause "A3 for every gate-intertwining refinement" is **false** when $c^* \ne 0$ (Proposition 89.7); $K = 1 + (H - E_0)/v$ violates A2 (89.3); open core: the operator lift W2 (Lemma 89.L).

**Ranked principles [proposal; N:705–723].** (1) Geometric stress with controlled critical RG (C1–C6): strongest sufficiency candidate; (2) non-null holomorphic endpoint/stress equations with contact normalisation: best local selector (a finite rank-drop problem); (3) regular spectral YBE with a specified transfer and branch, starting from (78.3): neither necessary nor sufficient; (4) isotropy or normalised blocking alone: filters; (5) topological symmetry alone: no selector. Output is a selected subset of coefficient space with all auxiliary choices, never a canonical $h$.

**What this does not claim.** No selection theorem and no selected coupling for any $\mathcal{C}$; no

existence of a certificate; no attribution of (78.3) without a local source; the endpoint equations (N:399–466) and Gram amplification D2-L9 are only pointed to. The examples below calibrate; no proof uses them.

| Instance | Calibration (source) |
|---|---|
| Dense TL / ABF $\mathcal{A}_{k+1}$ | two sign regimes, two central charges; lattice category $\neq$ primary category (AFM :619–623) |
| Dilute $A_L$, branches 1–2 | branch and seam are extra data beyond the loop weight (ZB 9611156.tex:430–455) |
| Tambara–Yamagami input | lattice category not even a subcategory of the CFT's MTC (AFM :775) |
| Hilb, $O = \mathbb{C}^2$ | witness for Prop. 78.7(ii) (`review_checks.py`:5–7) |
| CGS loop $O(n)$, $n = 1$ | lattice stress proved only at $n = 1$ (StressTensor.tex:529–532) |

# 80 Koo–Saleur Tiles: Spectral-Shell Certificate and Refinement Defect

## 80.1 Status, Sources, and Dependencies

**Status: mixed, labelled per statement.** Lemmas 79.1, 79.6, 79.7 and Theorem 79.8 are **derived** (finite algebra; graded "confirmed" by the in-repo review). Theorem 79.2, Propositions 79.3, 79.4 and 79.9 are **conditional** (graded "conditional-ok"; hypotheses listed). Example 79.5 is **negative**. Position 79.10 is a derived argument under stated assumptions; §80.6 is **conjectural** and a **proposal**. No chain, free or interacting, is shown here to satisfy P0–P3. Everything holds for a unitary fusion $\mathcal{C}$ with all multiplicity indices and no self-duality, braiding or Temperley–Lieb relation, except Proposition 79.9 and §80.6, which need a declared modular target. Reproducibility: `explorations/2026-10-22-astra/D4-koo-saleur-beyond-free/checks/run_checks.py` (Python; 20 exact equality checks, five rejected mutations, qubit sites only, i.e. $\mathcal{C} = \mathrm{Vec}$; the identities checked are formal).

Dependencies: CA-14, CA-67, CA-68, CA-71, CA-73, CA-75 rows (4)–(6), CA-81, CA-86; LC-2, LC-12; CONVENTIONS (r), (s), (t), (v), (af), (ak), (al), (ap).

## 80.2 The KS Prescription Is a Tile Prescription

**Setting.** $O \in \mathcal{C}$ a site object, $h_{N,j} \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ self-adjoint, $H_N^{\mathrm{raw}} = -\sum_j h_{N,j}$ (CONVENTIONS (ak)), a given velocity $v_N > 0$, and the velocity-one density $k_j := -h_{N,j}/v_N$, which is the $h$ of (af). Declared extra data: a periodic representation on $N \geq 4$ sites with a unitary translation $U_N$, $U_N^N = 1$, $U_N k_j U_N^* = k_{j+1}$, and disjoint-support commutativity. The forward current is $\mathcal{J}_j := i[k_j, k_{j+1}]$ (three-site support). With $z = e^{2\pi i/N}$, $a_N = N/4\pi$, a scalar density subtraction $\varepsilon_N$ (the $e_n$ of (af), renamed to avoid the cup-cap tile $e_j$ of (ak)) and a real zero-mode shift $b_N$:

$$T_{N,n} = a_N \sum_j z^{nj}(k_j - \varepsilon_N - \mathcal{J}_j) + b_N \delta_{n,0}, \qquad \bar{T}_{N,n} = a_N \sum_j z^{-nj}(k_j - \varepsilon_N + \mathcal{J}_j) + b_N \delta_{n,0}. \quad (1)$$

Expanding $h_j = \sum_\alpha c_\alpha D_\alpha^{(j)}$ in a dagger-stable tile basis (CA-86) gives the forward chiral tile $-\sum_\alpha \frac{c_\alpha}{v_N} D_\alpha^{(j)} - \varepsilon_N - \frac{i}{v_N^2} \sum_{\alpha,\beta} c_\alpha c_\beta [D_\alpha^{(j)}, D_\beta^{(j+1)}]$: both powers $1/v_N$, $1/v_N^2$ matter. The categorical evaluation fixes recoupling; the coefficients fix dynamics. No isotropy, velocity or Ward identity follows.

**Lemma 79.1 (D4-T) [derived].** $i[\sum_l k_l, k_j] = \mathcal{J}_{j-1} - \mathcal{J}_j$, $T_{N,n}^* = T_{N,-n}$, and $[\sum_j z^{mj} k_j, \sum_j z^{nj} k_j] = -i(z^n - z^m) \sum_j z^{(m+n)j} \mathcal{J}_j$. **Proof.** Only neighbouring densities fail to commute (interchange law); the neighbouring pair carries coefficient $z^{mj} z^{n(j+1)} - z^{m(j+1)} z^{nj} = z^{(m+n)j}(z^n - z^m)$; $k_j, \mathcal{J}_j$ are self-adjoint. □ If $\varepsilon_N = E_0(N)/(N v_N)$ then $T_{N,0} + \bar{T}_{N,0} = \frac{N}{2\pi v_N}(H_N^{\mathrm{raw}} - E_0(N)) + 2b_N$; replacing $T_n \mapsto T_n + d\,\delta_{n,0}$ maps $c \mapsto c + 24d$ and fixes $c^*$ [derived], so a wrong cubic anomaly is never repaired by an energy origin (cf. GS:3224–3229). *Amendment (2026-10-22):* with $K = 1 + (H_N^{\mathrm{raw}} - E_0)/v_N$ this identity makes A2 fail at $n = 0$ whenever the spectral width of $K - 1$ stays bounded below (CA-89, Proposition 89.3); this is one reason the regularity control of Theorem 79.2 is an auxiliary $Q^p$.

## 80.3 A Spectral-Shell Certificate for VR-1

**Setting.** A tower of finite Hilbert spaces $H_r$ ($N = N_r \to \infty$), isometries $J_r$ (the GNS maps of the proposed refinement), modes $T_{r,n}$ of (1), and an *auxiliary* grading $Q_r \geq 1$. Physical energy

$E_N = \frac{N}{2\pi v_N}(H_N^{\mathrm{raw}} - E_0)$ almost never intertwines exactly with a refinement, so interpretation $(E_N)$ and regularity $(Q_r)$ are separated; their uniform comparison is a further obligation. Shells $E_{r,a} = 1_{[2^a,2^{a+1})}(Q_r)$, cutoffs $P_r(\Lambda) = 1_{[1,\Lambda]}(Q_r)$; constants below are independent of $r$; each statement applies to both chiralities and to the mixed residuals $[T_{r,m}, \bar{T}_{r,n}]$. *Amendment (2026-10-22):* under equal ledgers the mixed residuals reduce to the zero-mode commutator $[T_{r,0}, \bar{T}_{r,0}]$ (CA-91, Theorem 91.5); a complex-linear parity supplies equal ledgers (Theorem 91.1).

**Property P [hypothesis; not synthesised from $\mathcal{C}$]. P0.** $Q_{r+1}J_r = J_rQ_r$, $T_{r,n}^* = T_{r,-n}$. **P1.** For every $d > 0$: $\|E_{r,a}T_{r,n}E_{r,b}\| \le C_{n,d}\,2^b 2^{-d|a-b|}$, full shell blocks with all multiplicities. **P2.** With $z_{mn} = \delta_{m+n,0}(m^3c^* - mc)/12$ and $R_{r,m,n} = [T_{r,m}, T_{r,n}] - (m-n)T_{r,m+n} - z_{mn}$: $\|Q_r^t R_{r,m,n} P_r(\Lambda)\| \to 0$ for each fixed $\Lambda$ and real $t \ge 0$. **P3.** $\Delta_{r,n} = T_{r+1,n}J_r - J_rT_{r,n}$ obeys $\|Q_{r+1}^t \Delta_{r,n} P_r(\Lambda)\| \le A_{n,t}2^{-\alpha_{n,t}r}\Lambda^{\gamma_{n,t}}$, $\alpha_{n,t} > 0$, for all $\Lambda \ge 1$ (full output, low input).

P1 is the uniform lattice analogue of the Carpi–Weiner energy bound $\|L_n v\| \le r(1+|n|^{3/2})\|(1+L_0)v\|$ (CW:727–747), with $Q$ in the role of $1 + L_0$ and off-diagonal shell decay added [source-local comparison]. The only local evidence on an interacting chain is numerical and fixed-window: on a nonintegrable chain the KS modes' low-window overlaps outside the CFT selection rule were fitted to decay as $1/N^2$ for $N = 13, \dots, 22$ (MV:115, :567, :573) [source-local; not a shell bound].

**Theorem 79.2 (D4-P) [conditional]. ASSUME:** P0–P3; a fixed integer $p > 1$. **PROVE:** $K_r = Q_r^p$ satisfies VR-1 A1–A4 (`virasoro_routes.md`:55–80); hence modes and all finite words converge on $\bigcap_t \mathrm{Dom}(Q^t)$. Genuine Virasoro needs in addition $c^* = c$.

⟨1⟩1. The $(a,b)$ block of $Q^tT_nQ^{-t-1}$ is $\le 2^tC_{n,d}2^{-(d-t)|a-b|}$ $(d > t)$; a Schur test on row/column sums bounds the operator norm by $M_{n,t} = 2^tC_{n,d}\frac{1+2^{-(d-t)}}{1-2^{-(d-t)}}$, with no shell-dimension factor.

⟨1⟩2. A1 by functional calculus from P0. A2 at $t = ps$: $K^sT_nK^{-s-1} = (Q^{ps}T_nQ^{-ps-1})Q^{-(p-1)}$.

⟨1⟩3. Inserting $Q^{-t-1}Q^{t+1}$: $\|Q^tR_{mn}Q^{-t-2}\| \le B_{mn,t} := M_{m,t}M_{n,t+1} + M_{n,t}M_{m,t+1} + |m-n|M_{m+n,t} + |z_{mn}|$.

⟨1⟩4. Splitting the input at $\Lambda$, the $K$-weighted residual is $\le \|Q^tRP(\Lambda)\| + B_{mn,t}\Lambda^{-2(p-1)}$ at $t = ps$; take $\Lambda$ large, then $r$ large (P2): A4.

⟨1⟩5. Exact $Q$-transport and ⟨1⟩1 give $\|Q_{r+1}^t\Delta_{r,n}Q_r^{-t-1}\| \le 2M_{n,t}$; the same split gives $e_{r,n,s} \le A2^{-\alpha r}\Lambda^\gamma + 2M_{n,ps}\Lambda^{-(p-1)}$ (P3 constants at $t = ps$); $\Lambda = 2^{\theta r}$, $\theta\gamma < \alpha$, makes it summable: A3.

⟨1⟩6. QED by ⟨1⟩1–⟨1⟩5 and VR-1. The powers $ps$ are cofinal, so the smooth domain is unchanged.

The spare power $p > 1$ is exactly the "stronger auxiliary regularity weight" the referee requires before a fixed-core estimate may be advertised as a graph-norm certificate (referee:155–161; REVIEW:51).

**Proposition 79.3 (D4-B) [conditional]. ASSUME:** P1 and $\|E_{r,a}R_{r,m,n}E_{r,b}\| \to 0$ for each fixed $(a,b)$. **PROVE:** P2. ⟨1⟩1. Products obey an order-two envelope: summing two P1 bounds over the intermediate shell with exponent $D > d + 2$ gives $C_d 2^{2b}2^{-d|a-b|}$. ⟨1⟩2. For fixed $\Lambda$ only finitely many $b$ occur; the geometric output tail is uniformly small. ⟨1⟩3. The finite rectangle tends to zero. QED.

**Proposition 79.4 (D4-I) [conditional]. ASSUME:** $\|\mathrm{ad}_{Q_r}^k(T_{r,\pm n})E_{r,b}\| \le A_{n,k}2^{(k+1)b}$ for all $k \ge 0$, uniformly in $r$. **PROVE:** P1. ⟨1⟩1. For $a \ge b + 2$ the shells are separated by $g \ge 2^{a-1}$, and $Y \mapsto \int_0^\infty e^{-sQ_a}Ye^{sQ_b}ds$ inverts $X \mapsto Q_aX - XQ_b$ with norm $\le 1/g$. ⟨1⟩2. Applying it $k$ times gives $2^kA_{n,k}2^b2^{-k(a-b)}$; adjoints give $a < b$; take $k > d$. QED. Lieb–Robinson bounds give no such *relative* estimate by themselves: the rescaling in $E_N$ is extensive.

**Example 79.5 [negative; spare power is needed].** On orthonormal pairs $a_j, b_j$ with $Q = 2^j$ on the $j$-th pair, $T_{r,1} = 2^r|a_r\rangle\langle b_r|$, $T_{r,-1} = T_{r,1}^*$, other modes 0, $c = c^* = 0$: P1 holds and all fixed blocks vanish eventually, but $R_{r,1,-1} = 2^{2r}(|a_r\rangle\langle a_r| - |b_r\rangle\langle b_r|)$ has $\|Q^t R Q^{-t-2}\| = 1$ while $\|K^s R K^{-s-2}\| = 2^{-2(p-1)r}$. (It isolates A4; it does not satisfy A3.)

## 80.4 KS Modes under Categorical Refinement

**Setting.** Any isometry $V : H_N \to H_M$ (a CA-71 placement is one instance), $P = VV^*$, $P^\perp = 1-P$, fine densities $k'_a$, compression $B_a = V^* k'_a V$, leakage $F_a = P^\perp k'_a V$.

**Lemma 79.6 (D4-R) [derived].** $V^*\mathcal{J}'_a V = i([B_a, B_{a+1}] + F_a^* F_{a+1} - F_{a+1}^* F_a)$; $\Delta_n := T'_{M,n}V - VT_{N,n} = P^\perp T'_{M,n}V + V(V^*T'_{M,n}V - T_{N,n})$ with orthogonal ranges; $V^*T'_{M,n}V = a_M \sum_a z_M^{na}(B_a - \varepsilon_M - V^*\mathcal{J}'_a V) + b_M\delta_{n,0}$. If local stretched representatives $S_j$ satisfy $S_j V = Vk_j$, then $\Delta_n = \{a_M \sum_a z_M^{na}(k'_a - \varepsilon_M - \mathcal{J}'_a) - a_N \sum_j z_N^{nj}(S_j - \varepsilon_N - i[S_j, S_{j+1}])\}V + (b_M - b_N)\delta_{n,0}V$: fine-versus-stretched tiles, Fourier phase/scale/velocity, scalar origin. **Proof.** Insert $1 = P + P^\perp$ between $k'_a$ and $k'_{a+1}$; use $V^*V = 1$. □ Compressed currents are not currents of compressed densities; the $F^*F$ terms are the missing part. One inserted vacancy shifts the phase of every later site, so a local density error does not give a local KS error; $O(N_r^{-\epsilon})$ is summable on geometric $N_r$, $O(1/N)$ on $N \to N+1$ is not. *Support remark [derived]:* the corner image is $\theta(k_j) = Vk_jV^* = S_jP$, which pins *every* inserted slot; CA-73:169–171 should be read through the local representative $S_j$ (REVIEW:50).

**Lemma 79.7 (D4-V) [derived; negative]. ASSUME:** fine energy $\sum_a k'_a$ nearest-neighbour; $V$ the bare dyadic insertion of CA-68 (one $\iota_{\mathbf{1}}$ after each site, no dressing). **PROVE:** $V^*(\sum_a k'_a)V$ is a sum of one-site morphisms in $\mathrm{End}_{\mathcal{C}}(O)$. ⟨1⟩1. Each fine bond meets one coarse factor and one $\iota_{\mathbf{1}}$; its compression is $(\mathrm{id} \otimes \iota_{\mathbf{1}}^\dagger)k'(\mathrm{id} \otimes \iota_{\mathbf{1}})$ or the mirror. ⟨1⟩2. Sum over bonds. QED. *Consequence:* exact covariance of both $T_0$ and $\bar{T}_0$ would make $V^*(T'_0 + \bar{T}'_0)V = T_0 + \bar{T}_0$, i.e. coarse $H_N^{\mathrm{raw}}$ a one-site sum plus scalar; whenever it is not, exact covariance fails for this $V$. Leakage in Lemma 79.6 can still mediate two-site terms; dressings change the premise.

**Theorem 79.8 (D4-N) [derived; negative]. ASSUME:** nonzero finite $H_r$, A1, A4, adjoint pairing, and $T_{r+1,n}J_r = J_rT_{r,n}$ for all $n$ on all of $H_r$. **PROVE:** $c^* = 0$, $T_{r,n} = 0$ for $n \neq 0$, $T_{r,0} = \frac{c}{24}1$. ⟨1⟩1. Covariance transports $R_{r,m,n}$ isometrically to every later stage, where A4 sends it to zero on each vector: $R_{r,m,n} = 0$. ⟨1⟩2. With $D = \dim H_r$, traces of the $(1,-1)$, $(2,-2)$ brackets give $0 = 2\,\mathrm{Tr}\,T_0 + (c^* - c)D/12$ and $0 = 4\,\mathrm{Tr}\,T_0 + (8c^* - 2c)D/12$; so $c^*D/2 = 0$. ⟨1⟩3. $[T_0, T_n] = -nT_n$ kills $T_n$ beyond the spectral diameter of $T_0$; the bracket of $T_a, T_{k-a}$ then gives $(2a-k)T_k = 0$. ⟨1⟩4. The $(1,-1)$ bracket gives $T_0 = \frac{c}{24}$. QED. *Prior art:* trace cyclicity against a finite-$N$ central term (GS:2489); finite-matrix central terms are coboundaries (LC-2 Prop. 5); no exact implementer for $c \neq 0$ (CA-67:104–109, NG5). New here: ⟨1⟩1 and the zero-representation conclusion. Hence A3 must be approximate; exact $Q$-covariance, exact categorical relations and eventual equality on fixed low cores are not excluded.

## 80.5 Extended Fields Are Declared Data

**Proposition 79.9 (D4-G) [conditional]. ASSUME:** finitely many convergent unitary sector characters $\chi_a(\tau) = \mathrm{Tr}_{V_a}\, q^{L_0 - c/24}$, nonnegative, transforming by the declared modular $S$ with $S_{\mathbf{1}a} > 0$; weight-zero vacuum; $c > 1$. **PROVE:** $V_{\mathbf{1}}$ is not a finite sum of Virasoro highest-weight modules. ⟨1⟩1. $M$ modules would give $\chi_{\mathbf{1}}(i\beta/2\pi) \le Me^{\beta c/24}\prod_n(1-e^{-\beta n})^{-1}$ (PBW). ⟨1⟩2. $\log\prod \le \sum_k [k(e^{\beta k}-1)]^{-1} \le \pi^2/6\beta$. ⟨1⟩3. $S$ and the positive vacuum row give $\chi_{\mathbf{1}}(i\beta/2\pi) \ge S_{\mathbf{1}\mathbf{1}}e^{\pi^2 c/6\beta}$. ⟨1⟩4. As $\beta \to 0$ this beats ⟨1⟩2 when $c > 1$. QED. The threshold is sharp for this argument. LC-12

§5's bound $\sum k^{-2} < 2$ needed $c > 12/\pi^2$ and was applied at one $c$ (D5 REVIEW:105–106); D5-H reaches the same $c > 1$. This answers the general-$c$ question of CA-75:108–111.

**Position 79.10 [derived argument; assumptions as stated]. ASSUME** the hypotheses of Proposition 79.9 for the declared target. Then (i) Virasoro modes alone, with finitely many families per sector label, cannot carry $V_{\mathbf{1}}$: additional strong generators and their singular OPE coefficients are part of the declared target data, not output of $\mathcal{C}$. (ii) Any $D$ commuting with every $h_j$ (closed topological defects do, in the self-dual multiplicity-free setting of AFM-T:186–212, AFM-F:175–184) commutes with every $T_n, \bar{T}_n$; if $D = W_n$ with $[T_0, W_n] = -nW_n$, then $nW_n = 0$. So defect operators and their polynomials are zero modes, never nonzero chiral generators. (iii) Endpoint/junction fields, higher conserved densities and microscopic currents are *proposals* for avatars; their weights, OPEs and positivity are extra equations. Transport of such avatars under VR-1-type bounds with finite losses (D4-X) remains a conditional proof sketch in the note.

## 80.6 Headline Conjecture and First Attackable Lemma

**Conjecture (D4-C) [conjectural; review verdict: overstated as a statement about $\mathcal{C}$].** From an effective positive rational sewing completion $\mathcal{S}$ of a UMTC $\mathcal{C}$ one can effectively build finite categorical chains, nearest-neighbour tile Hamiltonians, local refinements and states whose standard KS modes satisfy P, which satisfy WT-1, and whose limit is unitarily the diagonal theory of $\mathcal{S}$ with full fidelity to $\mathcal{C}$. Verdict (REVIEW:167–192): the carrier admits unit multiplicities and growing site objects, hence *any* spin chain; the input $\mathcal{S}$ is the full CFT; site growth is unconstrained relative to locality. It is an inverse-discretisation (lattice-realisability) conjecture strengthening ZW Conj. 4.3 and Rmk. 9 (ZW:871–895) and the route of `virasoro_routes.md`:41, which the note does not cite. **Proposal:** restate with a pointed carrier $O = \mathbf{1} \oplus X$ and bounded support growth, separating the KS-certificate part from fidelity (REVIEW:231–235).

**First attackable lemma [proposal].** Prove P0–P3 for the sourced quasi-free calibration (`cft_machine/benchmarks/fermion/PROOF.md`:12–76; its source-modified modes) with $Q = 1 +$ normal-ordered one-particle |momentum|: zero padding transports $Q$ exactly and a mode shifts it by at most $|n|$ (finite band, P1); the fixed-core defect $O(M^{-2}|r|^3)$ on fixed cores (PROOF.md:50–64) suggests P3 with $\gamma \approx 3$. Through Theorem 79.2 this would be the first VR-1 instance (CA-75 row (4)) and answer the referee boundary. The next general lemma is Proposition 79.4's relative commutator bound for an interacting density.

**What this does not claim.** No instance of P, free or interacting; no selection of $h$, $v_N$, $\varepsilon_N$, $b_N$ from $\mathcal{C}$; no periodic categorical representation or unital OAR map; no positivity, self-adjointness, integration or local-net identification; no fidelity. The no-wrap filter of the note is not adopted.

| Instance | Role (calibration only) |
|---|---|
| Source-modified free fermion | Fixed-core KS errors (PROOF.md:50–65); candidate first P instance; not verified. |
| Ising / nonintegrable ANNNI | MV:115, :551–575: fixed-window KS selection rules, fitted $1/N^2$ leakage; numerical. |
| Fibonacci, $c = 14/5$ | LC-12 §5: single-$c$ case of Proposition 79.9. |
| Vec, qubit sites | D4 check bundle: formal identities of Lemmas 79.1, 79.6, 79.7, Example 79.5. |

# 81 Exact Lattice Cells, Temporal Reflection Positivity, and a Two-Sided Fidelity Certificate

## 81.1 Status, Sources, and Dependencies

**Status: mixed; every statement carries its own label.** Lemmas 80.1–80.6 are *derived*, finite, and hold for every unitary fusion $\mathcal{C}$ (no braiding used). Lemma 80.7 is *conditional*; Proposition 80.8 and Lemma 80.10 are *derived, negative*. Theorem 80.9 is *conditional*; its finite-dimensional gap is closed along route (ii) in CA-93 (amendment 2026-10-22; the review graded the original "overstated"). Lemmas 80.11–80.12 are *conditional*; Conjecture 80.C is *conjectural*; the chiral-versus-full target is an *open decision*. This shard answers part of CA-75 row (6) and Link (a) and the "exclusion of unintended sectors" in CA-75 contract item 13. Reproducibility: exact SymPy checks in the exploration directory (`checks/check_cells.py`: six invariant groups pass, five mutations rejected; `checks/review_pp_constant.py`); a Julia port is an open item.

Dependencies: CA-66, CA-75, CA-82 (instance only), CA-86 (Lemma 86.1); LC-4, LC-9 (Lemma FP-D), LC-12 §5; CONVENTIONS (r), (af), (ak), and the entry (al) (bond energy $k_j$; see the boundary paragraph).

## 81.2 Setting and Notation

$\mathcal{C}$ is unitary fusion with site object $O$ (CONVENTIONS (r)). Following (ak), $H = -\sum_j h_j$, $h_j \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$, $h_j = \sum_\alpha c_\alpha D_\alpha^{(j)}$ in the elementary tiles $D_\alpha$; $e_j = b_j d_j$ is *only* the cup-cap tile. The *exact tile* is $\mathcal{T}_j(t) := e^{t h_j} = \sum_\alpha w_\alpha(t) D_\alpha^{(j)}$; its coefficients $w_\alpha(t)$ differ from $c_\alpha$ (CA-86's first-order weight is $w_\alpha = \delta\tau\, c_\alpha$). The positive bond energy is $k_j := -h_j$, so $H = \sum_j k_j$, with current $J_j := i[k_j, k_{j+1}]$; $k_j$ is the density called $h_{(n,j)}$ in (af) and LC-4. A cut $\gamma$ carrying ordered object $W_\gamma$ has Hilbert space $\mathcal{H}_\gamma = \bigoplus_{a \in \mathrm{Irr}\,\mathcal{C}} \mathrm{Hom}_{\mathcal{C}}(a, W_\gamma)$, all charges and multiplicities kept; this is not a tensor factorisation across a fusion constraint. Hermitian forms are linear in the second slot. A finite patch $P$ is a fixed oriented brick graph with input/output cuts, and

$$Z_P = \sum_{\Gamma \text{ on } P} \Big( \prod_p w_{p,\alpha_p} \Big) \, \mathrm{Ev}_{\mathcal{C}}(\Gamma),$$

internal labels summed and external labels free; with exact tiles there is no $O(\delta\tau^2)$ remainder (contrast Lemma 86.1).

## 81.3 Exact Cells and Sewing

**Lemma 80.1 [derived; D5-S, REVIEW "confirmed"]. Assume** finite patches $P, Q$ sharing a cut $\gamma$ that runs *along tile legs* (a zigzag through no plaquette) with compatible orientation, a basis $\{u_A\}$ of $\mathcal{H}_\gamma$ with Gram matrix $\mathbf{G}_{AB} = \langle u_A, u_B \rangle$, and synthesis map $U$. **Prove** $Z_{Q \circ P} = Z_Q\big(\sum_{AB} |u_A\rangle (\mathbf{G}^{-1})_{AB} \langle u_B|\big) Z_P = Z_Q Z_P$, and dual block bases transform by $F^{-T}$. $\langle 1 \rangle 1$. A history on the sewn graph is a pair of histories with a common interface label; weights multiply and evaluations compose. BY the definition of $Z$. $\langle 1 \rangle 2$. $\mathbf{G} = U^\dagger U$ gives $U \mathbf{G}^{-1} U^\dagger = 1_{\mathcal{H}_\gamma}$; summing the common labels is this insertion. BY $\langle 1 \rangle 1$. $\langle 1 \rangle 3$. With tree kets $t_L = F t_R$, a dual basis $b_L = \tilde{U} b_R$ satisfies $\tilde{U} F^T = 1$, so $\tilde{U} = F^{-T} = \overline{F}$ in unitary gauge. This is LC-9 Lemma FP-D (the continuum statement) restated at a lattice cut. QED. Regroupings of one weighted graph agree; an $F$-move changes basis but does not identify two different weighted brick graphs.

## 81.4 Temporal Reflection Positivity

**Lemma 80.2 [derived; D5-P, "confirmed"]. Assume** a temporal reflection cut whose quotient $\mathcal{H}_\gamma$ is spanned by half-patch insertions $x_r$, and seam operator $\Sigma = \sum_\alpha w_\alpha D_\alpha$. **Prove** all-insertion reflection positivity holds iff $\Sigma \geq 0$, iff each charge block $\Sigma^a = [w_{a,\mu\nu}] \geq 0$ in multiplicity matrix units. ⟨1⟩1. The reflected gluing is $Q_{rs} = \langle x_r, \Sigma x_s\rangle$, BY Lemma 80.1 and the dagger reflection. ⟨1⟩2. $\sum \bar{z}_r z_s Q_{rs} = \langle x, \Sigma x\rangle$ with $x = \sum z_s x_s$ ranging over $\mathcal{H}_\gamma$ by spanning. ⟨1⟩3. $\mathrm{End}_{\mathcal{C}}(W) = \bigoplus_a \mathrm{End}(H_a)$ faithfully, so positivity is blockwise; categorical weights $d_a > 0$ do not change it. QED.

**Corollary 80.3 [derived; coefficient constraints].** (i) If $\mathcal{T}(t)$ is reflection positive for all $t \in [0, \varepsilon)$ (spanning as in Lemma 80.2), then $h = \lim_{t\downarrow 0}(\mathcal{T}(t) - 1)/t$ is a limit of self-adjoint operators, so $h = h^\dagger$. (ii) If $h = h^\dagger$, $e^{th} > 0$ for every real $t$. Writing $D_\alpha^\dagger = \sum_\beta J_{\beta\alpha} D_\beta$, the condition is $c_\beta = \sum_\alpha J_{\beta\alpha}\overline{c_\alpha}$ (in matrix units $c_{a,\mu\nu} = \overline{c_{a,\nu\mu}}$). **No sign condition $c_\alpha \geq 0$ is necessary**, and positive coefficients are not sufficient; for $h = h^\dagger$ a linearised tile $1+th$ is positive iff $\lambda_{\min}(h^a) \geq -1/t$ on every block ($t > 0$).

**Lemma 80.4 [derived; D5-R and REVIEW note (e)]. Assume** $h_j = h_j^\dagger$, mutually commuting terms within the odd and within the even layer, $A = \prod_{\mathrm{odd}} e^{th_j}$, $B = \prod_{\mathrm{even}} e^{th_j}$, $t$ real. **Prove** $R = A^{1/2}BA^{1/2} > 0$, $R'(0) = \sum_j h_j = -H$, and $R = A^{-1/2}(AB)A^{1/2}$; hence CA-86's row $U = AB$ is similar to a positive operator and $\mathrm{Tr}(AB)^m = \mathrm{Tr}\, R^m > 0$. ⟨1⟩1. $A, B > 0$: commuting positive factors. ⟨1⟩2. $R = (B^{1/2}A^{1/2})^\dagger(B^{1/2}A^{1/2})$ is invertible, so $R > 0$. ⟨1⟩3. Product rule at $t = 0$: two half odd layers give one odd sum. ⟨1⟩4. Similarity is $A^{-1/2}ABA^{1/2} = A^{1/2}BA^{1/2}$. QED. At fixed volume $R^M \to e^{-\beta H}$ as the step shrinks; no uniform-in-volume rate is claimed. Range-$r$ densities need an $r$-layer palindromic schedule (REVIEW §5), routine but not written.

**Lemma 80.5 [derived; D5-G, energy-resolved positivity]. Assume** a finite collar generator $E = E^\dagger \geq 0$ with spectral projections $P_\lambda$ and $X$ mapping external states to the cut. **Prove** $Q(t) = X^\dagger e^{-tE} X = \sum_\lambda e^{-t\lambda} X^\dagger P_\lambda X$ has $(-1)^k Q^{(k)}(t) \geq 0$, and every multi-time kernel $[X_r^\dagger e^{-(t_r+t_s)E} X_s]_{rs} \geq 0$. ⟨1⟩1. Each coefficient is $(P_\lambda X)^\dagger(P_\lambda X)$. ⟨1⟩2. $\lambda^k e^{-t\lambda} \geq 0$. ⟨1⟩3. The kernel's form is $\|\sum_s e^{-t_s E} X_s u_s\|^2$. QED. Pointwise positivity does not suffice: $Q(t) = 1 - \frac{1}{2}e^{-t} > 0$ has a negative energy-one coefficient, and its two-time determinant is $-\frac{1}{2}(e^{-t_1} - e^{-t_2})^2 < 0$.

**Lemma 80.6 [derived finite identities; D5-W].** For differentiable $h(s)$, $\partial_s e^{th(s)} = \int_0^t e^{(t-u)h}(\partial_s h) e^{uh}\, du$ (variation of constants), and for nearest-neighbour $k_j$ with disjoint-support commutativity $i[H, k_j] = J_{j-1} - J_j$ (this restates (af) and LC-4:39). For a proposed connection with residual $W_X = \nabla_X Z$, $W_X(Z_m \cdots Z_1) = \sum_k Z_m \cdots Z_{k+1} W_X(Z_k) Z_{k-1} \cdots Z_1$ and $F_{\mathrm{out}}(X,Y)Z - ZF_{\mathrm{in}}(X,Y) = \nabla_X W_Y - \nabla_Y W_X - W_{[X,Y]}$ (anomaly line in "out"). For contractions $\|W_X(Z_m \cdots Z_1)\| \leq \sum_k \|W_X(Z_k)\|$: over a growing patch the *total* residual must vanish; small per-tile error is insufficient. Conformality, the flux and improvement equations of LC-4, and scalar curvature are *not* derived.

**Coverage of CA-75 Link (a).** (a2) is answered only for cuts along tile legs. *Not covered:* straight spatial seams through plaquettes (the case Link (a2) actually poses) and the anomaly-line limit. *Spatial* reflection positivity, whose crossing-kernel test changes with the leg reshuffling (NOTE.md:231–239), is treated for one crossing tile and a declared seam pairing in CA-92 (Theorems 92.3, 92.5; amendment 2026-10-22). (a3) gains only the total-residual bookkeeping; (a1) and (a4) are untouched.

## 81.5 Defect Operators Do Not Certify Fidelity

**Lemma 80.7 [conditional; D5-D, "conditional-ok"]. Assume** isometric WT-compatible embeddings $J_n$, uniformly bounded defects $D_{a,n}$ with $D_{\mathbf{1},n} = 1$, $D_{a,n}D_{b,n} = \sum_c N_{ab}^c D_{c,n}$, $D_{a,n}^\dagger = D_{\bar{a},n}$, and summable $\|D_{a,n+1}J_n - J_n D_{a,n}\| \leq \epsilon_{a,n}$ (likewise for adjoints). **Prove** strong limits $D_a$ exist

and obey the same relations; they commute with the limiting dynamics if the bounded evolutions and their commutators converge strongly. *Sketch:* telescoping with tail $\|v\| \sum_{k\geq n} \epsilon_{a,k}$ on the dense union, extension by uniform bounds, and strong convergence of products of uniformly bounded strongly convergent sequences (NOTE.md:356–363).

**Proposition 80.8 [derived, negative; scalar falsifier].** For every unitary fusion $\mathcal{C}$, $D_a := d_a 1$ satisfies every hypothesis-level relation of Lemma 80.7 (by $d_a d_b = \sum_c N_{ab}^c d_c$ and $d_{\bar{a}} = d_a$), commutes with every $H$, and carries no charge separation; for $|\operatorname{Irr}\mathcal{C}| > 1$ it is a nonfaithful representation of the fusion algebra. Hence convergent fusion operators, fusion relations and commutation with $H$ cannot certify fidelity for any $\mathcal{C}$. Actual sectors need localised, locally normal, transportable endomorphisms with convergent junctions; Gram lower bounds give only injectivity, not fullness (NOTE.md:374–381).

## 81.6 The Two-Sided Certificate

Fix four cyclically ordered intervals covering $S^1$ up to endpoints; $\mathcal{N} = \mathcal{A}(I_1) \vee \mathcal{A}(I_3) \subset \mathcal{M} = (\mathcal{A}(I_2) \vee \mathcal{A}(I_4))'$, and $\mu_{\mathcal{A}} = [\mathcal{M} : \mathcal{N}]$, the minimal Pimsner–Popa index (KLM:430–443). Let $\mathsf{B}$ be the declared chiral UMTC, $\operatorname{Dim}\mathsf{B} = \sum_i d_i^2$.

**Theorem 80.9 [conditional; D5-F in the review's repaired form; (F3′) from finite data: CA-93]. Assume** (F1) the OAR limit is a completely rational conformal net $\mathcal{A}$ (irreducible, strongly additive, split, with modular PCT; none of this follows from WT-1/VR-1); (F2) lattice endpoints give a fully faithful unitary braided tensor functor $\Phi : \mathsf{B} \to \mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$ with $d(\Phi(i)) = d_i$, preserving twists, duality and orientation; (F3′) there is a normal conditional expectation $E : \mathcal{M} \to \mathcal{N}$ with $E(x) \geq (\operatorname{Dim}\mathsf{B})^{-1} x$ for $x \in \mathcal{M}_+$. **Prove** $\Phi$ is a braided equivalence onto the complete finite-sector category and $\mu_{\mathcal{A}} = \operatorname{Dim}\mathsf{B}$. ⟨1⟩1. $\mu_{\mathcal{A}} \leq [\mathcal{M} : \mathcal{N}]_E \leq \operatorname{Dim}\mathsf{B}$ BY (F3′) and the PP definition. ⟨1⟩2. BY (F1) and finiteness, $\mu_{\mathcal{A}} = \sum_\rho d(\rho)^2$ over all irreducible sectors, and every sector is a finite direct sum (KLM:1719–1729, :1899–1900). ⟨1⟩3. The image of $\Phi$ contributes exactly $\operatorname{Dim}\mathsf{B}$: simples stay simple ($\mathrm{End} = \mathbb{C}$), distinct simples stay inequivalent, dimensions are preserved, BY (F2). ⟨1⟩4. An extra simple adds $d(\rho)^2 > 0$, contradicting ⟨1⟩1–⟨1⟩3; so $\Phi$ is essentially surjective, hence a braided equivalence. QED. Normalised $S$ and twists then match by ribbon traces; the character $T$ needs the declared central-charge lift as well.

**The gap, explicitly.** The note's checkable hypothesis was finite-dimensional: $\mathcal{N}_k \subset \mathcal{M}_k$ exhausting $\mathcal{N}, \mathcal{M}$ with $E_k \geq (\operatorname{Dim}\mathsf{B} + \varepsilon_k)^{-1}\mathrm{id}$, passed to the limit by KLM's monotone exhaustion (Prop. p3(a), KLM:507–527). KLM proves the increasing case "after taking commutants" (KLM:528–530), which uses commutant-invariance of the index. That fails for Pimsner–Popa constants of finite-dimensional inclusions: $\mathbb{C}1 \subset M_n$ has index $n$, while its commutant inclusion $1 \otimes M_n \subset M_n \otimes M_n$ on $\mathbb{C}^n \otimes \mathbb{C}^n$ has index $n^2$ (`checks/review_pp_constant.py`, exact for $n = 2, 3$). The finite form also assumes neither compatible $E_k$ nor a preserved state, so no normal limit expectation is built. *Theorem 80.9 therefore holds only under (F3′) or factor approximants; a finite-algebra positive-map test is currently not a sufficient check.* No counterexample to the conclusion is known; the gap is in the route. *Repair routes* (any one suffices; REVIEW.md:157–163): (i) (F3) for infinite-factor approximants; (ii) compatible $E_k$ preserving a faithful normal state, yielding a normal limit $E \geq \lambda\,\mathrm{id}$; (iii) a complete PP bound plus a sourced commutant lemma (source not local: Law 1 gap). By Lemma 80.10 the state in (ii) cannot be the vacuum. *Amendment (2026-10-22, round 2):* route (ii) is closed in CA-93 (Lemma 93.2, derived): a common normal state, faithful on $\mathcal{N}$ and preserved by every $E_k$, replaces compatibility; no complete positivity or commutant is used; routes (i), (iii) are unnecessary. So a finite positive-map test *is* sufficient given bounded exhaustion of both algebras and such a (non-vacuum) state; no instance is known (CA-93). Exhaustion of $\mathcal{M}$ itself is the separate conditional repair estimate D5-A (NOTE.md:458–504), valid with probes exactly in

$\mathcal{A}(I_2) \vee \mathcal{A}(I_4)$; its hypothesis nearly restates its conclusion (REVIEW.md:167–171).

**Lemma 80.10 [derived, negative; D5-V]. Assume** $\mathcal{N} \subset \mathcal{M}$, $\Omega$ cyclic for $\mathcal{N}$ and separating for $\mathcal{M}$, and an $\mathcal{N}$-bimodular expectation $E$ preserving $\omega = \langle \Omega, \cdot\, \Omega\rangle$. **Prove** $\mathcal{M} = \mathcal{N}$. $\langle 1\rangle 1$. $\langle n\Omega, E(x)\Omega\rangle = \omega(E(n^*x)) = \langle n\Omega, x\Omega\rangle$. $\langle 1\rangle 2$. Cyclicity gives $E(x)\Omega = x\Omega$; separation gives $E(x) = x$. QED. By Reeh–Schlieder the hypotheses hold for the four-interval inclusion, so an annular expectation preserves the vacuum only if $\mu_{\mathcal{A}} = 1$.

**Lemma 80.11 [conditional; D5-M, valid for $\mathsf{B} = \mathcal{C}$ only]. Assume** resolved modular covariance with multiplicities $M_{ij} \in \mathbb{Z}_{\geq 0}$, $M_{\mathbf{11}} = 1$ (M1); trace-norm convergence of lattice traces including defect insertions (M2); defects forming a unital $*$-representation of $\mathsf{B}$'s fusion algebra with Verlinde $S$ (M3). **Prove** $MS = SM$, $MT = TM$; defect eigencharacters are $\lambda_i(a) = S_{ai}/S_{\mathbf{1}i}$ via projections $p_i = S_{\mathbf{1}i} \sum_a \overline{S_{ai}} D_a$; every $i$ occurs iff the representation is faithful. Multiplicities are *not* fixed (proof NOTE.md:545–561; self-adjointness of $p_i$ uses $S_{\bar{a}i} = \overline{S_{ai}}$, $S_{\mathbf{1}i} > 0$). For $\mathsf{B} = Z(\mathcal{C})$ or fusion-only $\mathcal{C}$ a tube-algebra version is needed and is not written.

**Lemma 80.12 [conditional; D5-H, sharp].** Under $c > 1$, nonnegative weights, a weight-zero vacuum, positive characters with $S$-covariance and $S_{\mathbf{1}i} > 0$, the vacuum carrier is not a finite sum of Virasoro highest-weight families: PBW gives $\log \chi_{\mathbf{1}}(i/t) \leq \pi t/12 + O(1)$, against $\geq \pi c t/12 + \log S_{\mathbf{11}}$ (NOTE.md:586–599). It concurs with D4-G and sharpens LC-12 §5 ($c > 12/\pi^2$ there) to $c > 1$.

**Why selection is indispensable.** (F2) with $\mathsf{B} = \mathcal{C}$ is a strong demand on $(O, h)$: the category defining a lattice model and the category of the continuum primaries typically differ (AFM-J:123–128, AFM-S:775). A selection principle (CA-78; D2) must supply it; nothing in this shard does.

**Conjecture 80.C [conjectural; D5-C].** Every computably presented UMTC admits, with some compatible declared enrichment, a local lattice family whose OAR/KS limit satisfies (F1)–(F3′) with $\mathsf{B} = \mathcal{C}$. *Review verdict* (REVIEW.md:26–28, :182–211; INTEGRATION.md:115–117): it contains the open UMTC realisation problem (Bischoff2018:364–372) plus the whole analytic layer; it is degenerate for $\mathcal{C} = \mathrm{Vec}$ unless $c > 0$ is required; its uniformity clause is trivialised by proof search; it nearly duplicates D6-C. It is not a pipeline step.

**Open decision: chiral versus full target (PLAN.md D1).** D6's headline asks a 1D tower for a *chiral* limit net, while D2, D4 and D5 target full two-chiral theories with a tested chiral subnet (D6 REVIEW.md:20–24, :74–100). No local source settles whether a 1D lattice can have a chiral OAR limit; Theorem 80.9 is stated for whichever chiral subnet the declared target names, and neither periodic boundary conditions nor commutation with $H$ chooses it.

## 81.7 What This Does Not Claim

No critical Hamiltonian, convergent cell family, chiral or full OAR limit, endpoint functor, or index bound is constructed for any $\mathcal{C}$. Spatial reflection positivity, straight spatial seams, OS reconstruction, conformality of the Ward residual, and an instance of the finite-dimensional route to $\mu_{\mathcal{A}} \leq \operatorname{Dim} \mathsf{B}$ (CA-93) remain open; CA-92 covers one-tile spatial RP only. Matching $S, T$, fusion dimensions or closed-loop spectra is not fidelity.

| Instance | Calibration datum for the general statements |
|---|---|
| Arbitrary $\mathcal{C}$, $D_a = d_a 1$ | Proposition 80.8: fusion identities survive with no charge separation. |
| $\mathbb{C}1 \subset M_n$, $n = 2, 3$ | PP index $n$ versus commutant $n^2$: the gap in Theorem 80.9. |
| Ising (CA-82) | Sector completeness imported from a classified net; Theorem 80.9 is the general mechanism it bypasses. |
| Semion (LC-13) | Lemma 80.5: graded positivity rejects a pointwise-positive kernel. |
| Fibonacci (LC-12) | Lemma 80.12 generalises its $c = 14/5$ character-growth argument. |

# 82 The CFT Machine and Its Quantitative OAR Contract

## 82.1 Sources, Dependencies, and Status

This shard records Tobias's 2026-09-22 formulation of the programme and the new workspace `cft_machine/`. It does not assert a universal construction. Dependencies: CA-62, CA-67, CA-72, CA-73; CONVENTIONS (u)–(y).

## 82.2 The Actual Construction Target

**Programme.** Discover an algorithm, or functor, from categorical data to concrete lattice systems and microscopic Virasoro generators, whose limit through operator-algebraic renormalisation (OAR) and Wilson's triangle is a rigorous CFT with the input modular data. Indeterminate-number anyon spaces are one mechanism; they do not restrict the search.

A successful output names the microscopic algebras and Hamiltonians, refinement maps, renormalized states, symmetry generators, limiting local CFT, and modular-data comparison. Any additional conformal input must be explicit. Matching a finite spectrum, producing limiting commutators, or checking a finite moment problem does not supply the complete output.

The new workspace investigates a source-controlled fermion seed, a graph-norm convergence criterion, and equivariant fixed-point constructions alongside the anyonic tower. The source-grounded comparison and scoped claims are in `cft_machine/research/`.

## 82.3 A Quantitative Wilson-State Theorem

**Theorem WT-1 [local derivation]. Assume** unital C*-algebras $A_k$ and coherent unital injective $*$-maps $\alpha_n^k : A_k \to A_n$ for $n \geq k$. Let $\omega_n$ be microscopic states. Suppose for each fixed $k$ there are nonnegative $d_{k,n}$ with

$$\|\omega_{n+1} \circ \alpha_{n+1}^k - \omega_n \circ \alpha_n^k\| \leq d_{k,n}, \qquad \sum_{n \geq k} d_{k,n} < \infty.$$

The norm is the dual operator norm. Set $t_k(n) = \sum_{j \geq n} d_{k,j}$. **Prove** there is a unique state $\omega$ on the inductive algebra whose restrictions are the horizontal row limits, and

$$\|\omega \circ i_k - \omega_n \circ \alpha_n^k\| \leq t_k(n).$$

⟨1⟩1. Every row is norm Cauchy. BY telescoping adjacent differences and summability; for $m > n$ the difference is bounded by $\sum_{j=n}^{m-1} d_{k,j}$.

⟨1⟩2. Its limit $\omega_k$ is a state with the stated tail bound. BY completeness of the dual, closure of positivity and normalization under norm limits, and ⟨1⟩1.

⟨1⟩3. The limiting family is compatible: $\omega_l \circ \alpha_l^k = \omega_k$. BY coherence, the analogous equality holds for every microscopic scale $n \geq l$; take row limits.

⟨1⟩4. Define $\omega(i_k(a)) = \omega_k(a)$ on the dense algebraic union. It is well-defined, bounded, positive and unital. BY ⟨1⟩2–⟨1⟩3; continuous extension gives existence and density gives uniqueness.

⟨1⟩5. The GNS maps $[a]_k \mapsto [\alpha_l^k(a)]_l$ are compatible isometries preserving cyclic vectors. BY multiplicativity, unitality, and $\omega_l(\alpha_l^k(a^*a)) = \omega_k(a^*a)$.

⟨1⟩6. QED BY ⟨1⟩2–⟨1⟩5.

**Effective corollary WT-2.** If expectations and refinements are computable and the tail has a computable modulus, an observable $a \in A_k$ with a supplied computable bound $B \geq \|a\|$ can be evaluated in the limit to error $\eta$: choose $n$ with $t_k(n)B < \eta/2$ and compute its row expectation to error $\eta/2$. The modulus is substantive analytic input; finitely sampled decay does not provide it.

## 82.4 The Generator Contract and Its Boundary

**Conditional local theorem VR-1.** The companion proof in `cft_machine/research/virasoro_routes.md` constructs limiting modes and all their finite products under four explicit hypotheses: compatible isometric refinements with a positive auxiliary control operator, uniform bounds between successive graph norms, summable refinement defects in those norms, and vanishing weighted closure residuals. Its proof uses complete graph spaces and a direct telescoping identity for products.

The finite identity embeds as a projection $P_N$, with $P_N \to 1$ on the common smooth domain. A corner embedding is not identified with a unital observable map. Both central coefficients of CONVENTIONS (s) remain explicit. Adjoint pairing on the domain does not prove essential self-adjointness.

These two contracts must use the *same* state and representation system. WT-1 alone constructs no dynamics; VR-1 alone constructs no local OAR algebras. Positive energy, integration of smeared modes, local-net axioms, rationality, and modular fidelity need separate proofs or carefully verified applications of existing theorems.

**Numerical scope.** The fermion benchmark separately checks a generator-refinement square and the pullbacks of finite Dirac ground-state covariances. Its compatible limiting chiral sea is not substituted for these nontrivial Wilson rows. The corresponding source and run paths are listed in `INDEX.md` and the benchmark README.

# 83 A Supported Ising Modular-Data Compiler

## 83.1 Sources, Dependencies, and Scope

**Status: explicit supported construction using existing continuum theorems.** The code and structured derivations live in `cft_machine/compiler/`; the source-controlled numerical backend is `cft_machine/benchmarks/fermion/`. This is not a constructor for arbitrary categories. Dependencies: CA-81; CONVENTIONS (u), (w), (y), (z).

## 83.2 Exact Input and Concrete Output

In order $(1, \sigma, \psi)$ the supported seed has

$$S = \frac{1}{2}\begin{pmatrix} 1 & \sqrt{2} & 1 \\ \sqrt{2} & 0 & -\sqrt{2} \\ 1 & -\sqrt{2} & 1 \end{pmatrix}, \qquad \Theta = \mathrm{diag}(1, e^{\pi i/8}, -1), \qquad c = \frac{1}{2}.$$

The compiler compares $S$, twists, fusion multiplicities and the selected real $c$ exactly, allowing unit-preserving relabelling. Rational polynomial arithmetic modulo $z^8 + 1$ represents $z = e^{\pi i/8}$ and $\sqrt{2} = z^2 - z^6$. Character $T$ retains its separate vacuum phase $e^{-2\pi i c/24}$; it is not coerced into this smaller cyclotomic field. Explicit factor lists declare products; no general factorization algorithm or reconstruction of arbitrary $F/R$ data is claimed.

At even size $M$, take antiperiodic momenta $r = -M/2 + 1/2, \ldots, M/2 - 1/2$ and $\epsilon_M = 2\pi/M$. The finite microscopic mode matrix is the source-modified weighted shift

$$A_{M,n} e_r = \cos^2(\epsilon_M n/4) \frac{\sin(\epsilon_M (r - n/2))}{\epsilon_M} e_{r-n}$$

when both labels are in the window, and zero otherwise. The normal-ordered self-dual quadratic $Q_M(A_{M,n})$ includes the factor $1/2$ and the vacuum scalar; the recipe emits its actual CAR terms. The $n = 0$ sine Hamiltonian is nearest-neighbour in the real spatial basis, with an antiperiodic seam. Its positive-momentum occupation energies are positive, so the declared sea is its finite ground state. This chiral Hamiltonian is distinct from the auxiliary two-component Dirac benchmark.

The algebra map refines CAR generators by the isometric zero-padding map, $\Psi(f) \mapsto \Psi(Jf)$. It is unital and preserves conjugation and parity. A one-particle corner matrix $JAJ^*$ is not substituted for this CAR algebra map.

## 83.3 The Local Observable Theory

The selected finite observable algebra is the even self-dual CAR algebra. For $k$ factors take independent parity fixed points $(\mathbb{Z}_2)^k$. Total parity alone would retain odd-times-odd fields and give a different observable extension. Refinements are equivariant and the sea states are compatible. Thus these initial chiral states give constant Wilson rows; CA-81's nonconstant Dirac rows are a separate example of the same formalism.

The local reconstruction uses norm limits of even products of smoothly sampled, interval-supported real antiperiodic test sections. Its represented weak closure is the output interval algebra. This prescription defines the sampled-generator local net; it does not assert equality with a maximal construction using every arbitrary bounded scaling sequence.

**Local convergence argument [structured derivation]. Assume** a smooth real test section $f$ supported inside a fixed interval chart, sampled as $v_M(x) = \sqrt{\epsilon_M} f(x)$. **Prove** its sharp Fourier embedding converges to $f$ in $L^2(dx)$.

⟨1⟩1. Every fixed Fourier coefficient converges. BY the Riemann sum for the coefficient in the orthonormal basis $e^{irx}/\sqrt{2\pi}$.

⟨1⟩2. The norms converge to $\|f\|_2$. BY unitarity of the discrete Fourier transform and the Riemann sum $\epsilon_M \sum_x |f(x)|^2 \to \int |f|^2$.

⟨1⟩3. Weak convergence and convergence of norms give strong convergence. BY density of finite Fourier sums, ⟨1⟩1, boundedness from ⟨1⟩2, and expansion of the squared Hilbert-space difference.

⟨1⟩4. QED BY ⟨1⟩3.

CAR norm continuity then gives convergence of every fixed even field product. Density of supported smooth tests identifies the generating local even CAR algebra. The NS vacuum representation and the sourced even-net identity identify its vacuum interval net with $\mathrm{Vir}_{1/2}$. The universal global circle observable algebra is not identified with a plain even CAR algebra; the source explicitly distinguishes them.

### 83.4 Symmetry and Modular Fidelity

The source's modified-KS theorem supplies the strong finite-energy limit and convergence of its unitary implementers in the Majorana representation. The finite Schwinger computation separately checks the self-dual central coefficient $(n^3 - n)/24$; it is not a central term of a finite matrix commutator without normal ordering.

The complete-net sector theorem at minimal-model parameter $m = 3$ gives three sectors, with weights $0, 1/16, 1/2$, Ising fusion and the displayed twists and $S$. This supplies fidelity of the identified local net rather than guessing its sectors from finite spectra.

For products, the compiler retains each factor's stress family and their placed sum. The sum alone is not the full tensor-product net. The sourced tensor-sector completeness result and factorwise braiding give product modular data and additive central charge. The output explicitly records the relation between input labels and canonical product labels.

This construction establishes a supported branch. General category inputs, other Ising chiralities, arbitrary central-charge lifts, and general tensor functors remain outside its implemented domain. The broader machine must expand that domain with additional constructions and proofs.

# 84 Coset Lattice Stress Generators with Exact Core Stabilization

## 84.1 Sources, Dependencies, and Scope

**Status: source-based recipe with a local stabilization proof and exact finite witnesses.** This constructs stress generators inside an ambient fermion theory. It does not yet construct the local coset lattice theory itself. Dependencies: CA-81, CA-82; CONVENTIONS (z).

## 84.2 An Explicit Microscopic Polynomial

Take $k+1$ two-flavor complex fermion copies, $k \geq 1$, with half-integer momenta in the finite window and the filled negative sea. Let $t^a = \sigma_a/2$ and use the no-wrap one-particle shifts

$$j^a_{M,p}(e_r \otimes v) = e_{r-p} \otimes t^a v$$

when both labels are in the window, and zero otherwise. Normal-ordered complex second quantization gives finite quadratic currents $J^{u,a}_{M,p}$. Let $I$ sum the first $k$ copies, $J$ denote the last, and $D = I + J$. For a level-$l$ family $X$ define

$$\mathsf{S}^{[l]}_{M,n,R}(X) = \frac{1}{l+2} \sum_{a=1}^{3} \sum_{\substack{|p|\leq R \\ |n-p|\leq R}} :X^a_{M,n-p} X^a_{M,p}:,$$

$$C_{M,n,R} = \mathsf{S}^{[k]}(I) + \mathsf{S}^{[1]}(J) - \mathsf{S}^{[k+1]}(D).$$

The normal ordering orders current modes, keeping their contraction terms when the displayed products are expanded. Thus $C$ is an explicit CAR polynomial of degree at most four, not an unspecified continuum operator.

The infinite current cocycle has coefficient $p\delta_{ab}/2$ at level one: exactly $p$ Fermi crossings contribute $\mathrm{Tr}(t^a t^b) = \delta_{ab}/2$. Independent copies add levels. The sourced GKO result gives

$$c = \frac{3k}{k+2} + 1 - \frac{3(k+1)}{k+3} = 1 - \frac{6}{(k+2)(k+3)}.$$

This fixes the normalization before any finite computation.

## 84.3 Exact Stabilization on Free-Energy Cores

**Theorem CR-2 [local derivation]. Assume** the no-wrap finite currents above, compatible sea GNS embeddings, and auxiliary excitation energy

$$E_M = \sum_{u,f,r>0} r\, n_{u,f,r} + \sum_{u,f,r<0} |r|(1 - n_{u,f,r}).$$

Let $P_e$ project onto auxiliary energy at most $e$. **Prove** the embedded $C_{M,n,R}$ agrees with the continuum coset mode $C_n$ on $P_e$ whenever

$$R > e + |n|, \qquad M/2 - 1/2 > e + 2R.$$

⟨1⟩1. $[E_M, J_{M,p}] = -pJ_{M,p}$. By each nonzero bilinear moving occupation from $r$ to $r-p$; the scalar subtraction has $p = 0$. Consequently $J_{M,p}P_e = 0$ for $p > e$ by nonnegativity of $E_M$. The same holds in the infinite sea.

⟨1⟩2. Every omitted normally ordered current pair annihilates $P_e$. BY its indices summing to $n$: if either absolute value exceeds $R$, one index is positive and greater than $e$. Normal ordering places that mode on the right; use ⟨1⟩1.

⟨1⟩3. Retained products agree with continuum products in the safe window. BY deviations from the sea lying within $|r| \le e$, one current action involving only $|r| \le e + |p|$, and two retained actions involving only $|r| \le e + 2R$.

⟨1⟩4. The finite retained sum equals the continuum sum on $P_e$. QED BY ⟨1⟩2–⟨1⟩3.

The choice $R_M = \lfloor M/8 \rfloor$ eventually meets both conditions for every fixed $e, n$. A fixed word of modes uses the enlarged core $e + \sum_j |n_j|$ and therefore also stabilizes exactly. The symmetric two-index cutoff preserves finite adjoints; same-sign pairs must be grouped before taking adjoints, and zero-index pairs use the exact same-component commutation of $J_0^a$ with $J_p^a$.

**Vacuum corollary CR-3.** For $n \ge 1$, the smaller conditions $R \ge n$ and $M/2 - 1/2 > n$ suffice for $C_{M,-n,R}\Omega = C_{-n}\Omega$. Indeed, positive and zero currents annihilate the sea, so only two negative indices of total magnitude $n$ contribute; all involved orbitals have $|r| \le n$.

## 84.4 Exact Witnesses and the Remaining Interface

The sparse-CAR producer evaluates the actual polynomials with exact complex rational amplitudes and canonical fermion signs. No many-body matrix is allocated. At $M = 8, 16$ and $R = 3, 4$, it gives

| $k$ | $c$ | $\|C_{-2}\Omega\|^2$ | $\|C_{-3}\Omega\|^2$ |
|---|---|---|---|
| 1 | 1/2 | 1/4 | 1 |
| 2 | 7/10 | 7/20 | 7/5 |

The independent oracle is $c(n^3 - n)/12$, not a fitted central charge. The run distinguishes the general CR-2 window from the smaller CR-3 vacuum window. CAR signs, level addition, adjoint pairings and refined descendants are checked separately; coefficient and embedding-sign mutations are rejected.

What remains is a coherent *local coset observable* construction. The ambient fermion tower retains additional sectors. A stress difference does not remove them, and finite commutants or vacuum-cyclic compressions have not been shown to commute with refinement. Global finite-scale positivity and microscopic ground states of the quartic Hamiltonian are also unproved. The result solves a generator and cutoff-control problem; it does not yet enlarge the completed modular-data compiler beyond CA-82.

# 85 Coset Positivity and a Scalar-Augmented OAR Tower

## 85.1 Sources, Dependencies, and Scope

**Status: local derivations and exact finite witnesses.** Dependencies: CA-83, CH-1–CH-3 in `cft_machine/research/coset_hamiltonian.md`, CO-1–CO-3 in `cft_machine/research/coset_observable_system.md`; CONVENTIONS (aa).

## 85.2 The Raw Finite Zero Mode Is Not Positive

For $k = 1$, fill flavor 1 at every momentum in both copies and leave flavor 2 empty. Let $v$ be this normalized polarized occupation vector. For even $M$ and $0 \le R < M$, the raw coset polynomial of CA-83 obeys

$$\langle v, C_{M,0,R} v \rangle = -\frac{(M-R)(M-R-1)}{12}.$$

**Proof CH-1 [structured calculation].**

⟨1⟩1. In each copy, $\sum_a \|J_0^a v\|^2 = M^2/4 + M/2$: the third current has eigenvalue $M/2$, and each of the other two flips one of $M$ flavors with amplitude magnitude $1/2$. The two-copy diagonal current has squared norm $M^2 + M$.

⟨1⟩2. The zero-mode Sugawara difference is $2(M^2/4 + M/2)/3 - (M^2 + M)/4 = -M(M-1)/12$.

⟨1⟩3. For $1 \le p \le R$, each copy has $M - p$ retained flavor-flip transitions. Counting the two normally ordered terms gives a coset contribution $(M-p)/6$.

⟨1⟩4. Sum ⟨1⟩3 and add ⟨1⟩2. QED by finite summation.

Thus the sea, whose raw energy is zero, is not a ground state when $R < M - 1$. No diagonalization is needed. Exact checks give

| $M$ | $R = 3$ | $R = 4$ |
|---|---|---|
| 8 | $-5/3$ | $-1$ |
| 16 | $-13$ | $-11$ |

The witness has auxiliary energy $M^2/2$, so it does not contradict stabilization on a fixed low-energy subspace as $M$ grows.

## 85.3 Computable Vacuum-Descendant Spaces

**Construction CH-2.** Let $V_b$ be the span of the vacuum and words $C_{-n_1} \cdots C_{-n_t}\Omega$ with $n_j \ge 1$ and $\sum_j n_j \le b$. Construct them using the finite CAR polynomials at

$$R_b = 2b + 1, \qquad M_b = 16(b+1).$$

There are finitely many words. At every intermediate grade at most $b$, CR-2's conditions hold: $R_b > 2b$ and $M_b/2 - 1/2 = 8b + 15/2 > b + 2R_b$. Thus the finite vectors agree exactly with their continuum representatives.

Exact Gram elimination removes linear dependencies. If the retained column vectors form $U_b$ and $G_b = U_b^* U_b$, then $P_b = U_b G_b^{-1} U_b^*$ is their orthogonal projector. Refinement identifies the same words across scales, so the isometries $J_{Bb} : V_b \to V_B$ compose exactly. On these spaces both $E$ and $C_0$ act by total grade. The grade-zero space is the vacuum line.

The implemented preview through grade three uses $(\Omega, C_{-2}\Omega, C_{-3}\Omega)$ when present, with exact Gram diagonal $(1, c/2, 2c)$ and grades $(0, 2, 3)$. It uses the smaller CR-3 vacuum windows at $M = 8, 16$, not a large computation of the general schedule.

**Positive Hamiltonians CH-3.** The ambient operator

$$H_b = P_b E P_b + (b+1)(1 - P_b)$$

is positive and has unique vacuum ground state: its first block has nonnegative grades and one zero vector, while its other block is strictly positive. This is a different Hamiltonian from the raw quartic zero mode. Its global projection currently has no proved local fusion-space realization.

## 85.4 An Independent Scalar Gives Genuine Unital OAR

Define $A_b = B(V_b) \oplus \mathbb{C}$ and, for $B \geq b$,

$$\alpha_{Bb}(a, \lambda) = \big(J_{Bb} a J_{Bb}^* + \lambda(1 - J_{Bb} J_{Bb}^*), \lambda\big).$$

The scalar is independent of $a$; replacing it with a state expectation would generally preserve only complete positivity, not multiplication.

**Proof CO-1.**

⟨1⟩**1.** The two first-coordinate summands have orthogonal supports. Multiplication and adjoints are preserved, and $(1, 1)$ maps to $(1, 1)$.

⟨1⟩**2.** Compressing by $J_{Bb}^*$ recovers $a$; the second coordinate recovers $\lambda$. The map is injective.

⟨1⟩**3.** The complementary projection splits into the newly added shell and the transported previous shell. Substitution proves $\alpha_{DB}\alpha_{Bb} = \alpha_{Db}$.

⟨1⟩**4.** The states $\omega_b(a, \lambda) = \langle \Omega, a\Omega \rangle$ are compatible. Their Wilson rows are constant. The norm inductive limit is $K(\mathcal{H}_{\text{vac}}) + \mathbb{C}1$, and the vacuum GNS is $\mathcal{H}_{\text{vac}} = \overline{\bigcup_b V_b}$, since finite-rank operators act cyclically on $\Omega$.

⟨1⟩**5.** QED by ⟨1⟩1–⟨1⟩4.

The finite Hamiltonian $(E|_{V_b}, b+1)$ is positive. Its dynamics intertwine exactly: the old subspace and the new shell are energy-reducing, and the embedded coarse algebra is scalar on the new shell. The Hamiltonians themselves need not intertwine.

## 85.5 What the Construction Does Not Identify

This is a new constrained observable prescription. It is generally not the restriction of the original unital CAR refinement, which retains all new-mode multiplicities. The new map instead inserts the selected sea when it transports the matrix block.

Likewise the ambient penalized resolvents tend to $P(E + i)^{-1}P$, where $P$ projects onto the limiting vacuum module. This has a kernel on the discarded ambient sectors and is not a self-adjoint resolvent on that entire ambient Hilbert space. The new vacuum GNS representation is essential.

Global constraints can be intrinsic fusion constraints. Their presence is not an obstruction to locality. What remains unproved is the identification of these descendant spaces with categorical fusion spaces, together with a local realization of the Hamiltonian and refinement operations on them.

# 86 Smooth Coset Symmetry Limits and Local Reconstruction

## 86.1 Sources and Scope

Dependencies: CA-83, CA-84; SC-1/SC-2 in `cft_machine/research/smeared_coset_limits.md`; CO-2/CO-3 in `cft_machine/research/coset_observable_system.md`; CONVENTIONS (ab).

## 86.2 Three Cutoffs and an Ambient Operator Theorem

For a real smooth function $f$ on the circle, set $\widehat{f}_n = (2\pi)^{-1}\int f(x)e^{-inx}\,dx$ and

$$\|f\|_{3/2} = \sum_n |\widehat{f}_n|(1+|n|^{3/2}).$$

The microscopic quartic operator is

$$T_M(f) = \sum_{|n|\le K_M} \widehat{f}_n C_{M,n,R_M}, \qquad K_M = \lfloor M/16\rfloor, \quad R_M = \lfloor M/8\rfloor.$$

Its lift $A_M(f)$ uses the unital CAR refinement, not a one-particle or many-body corner extension. Each finite lift is bounded self-adjoint; its norm is not assumed uniformly bounded in $M$.

**Common-core lemma SC-1.** The algebraic finite-free-energy space decomposes orthogonally into unitary highest-weight coset modules. At the lowest unused free grade, positive modes annihilate; diagonalize the Hermitian $C_0$ there and generate descendants. Adjoint pairing preserves orthogonal complements. Finite free-grade multiplicity makes this procedure exhaust the space. The positive norm identity $\|C_{-1}v\|^2 = 2h\|v\|^2$ gives nonnegative lowest weights.

The sourced Carpi–Weiner bound is uniform in lowest weight at fixed $c$:

$$\|T(f)\xi\| \le r_c\|f\|_{3/2}\|(1+C_0)\xi\|.$$

It therefore applies to the module sum. The self-adjoint closures and their resolvents form direct sums, and the finite-free-energy vectors are a common core. This modulewise argument does not declare the whole ambient sector sum to be a single projective vacuum representation: its rotation phases can differ.

**Theorem SC-2 [structured proof]. Prove** $A_M(f)$ converges to $T(f)$ on that core, in strong resolvent sense, and through its one-parameter unitaries.

⟨1⟩**1.** On $P_e$, every retained microscopic mode agrees exactly with its continuum mode once $R_M > e+K_M$ and $M/2-1/2 > e+2R_M$. The displayed schedules meet both conditions eventually; $M \ge 32(e+1)$ suffices.

⟨1⟩**2.** Hence the remaining error is only a continuum Fourier tail, bounded by

$$r_c\|(1+C_0)\xi\| \sum_{|n|>K_M} |\widehat{f}_n|(1+|n|^{3/2}) \longrightarrow 0.$$

⟨1⟩**3.** For core vectors,

$$(A_M - i)^{-1}(T-i)\xi - \xi = (A_M-i)^{-1}(T-A_M)\xi \longrightarrow 0.$$

The resolvent norms are at most one, and $(T-i)$ maps a graph core densely into the Hilbert space. Extend the limit by density; similarly at $-i$.

⟨1⟩4. The self-adjoint strong-resolvent theorem gives strong convergence of $e^{itA_M}$, uniformly on compact $t$ intervals. Finite unitary products converge by telescoping and unit norm.

⟨1⟩5. QED by ⟨1⟩1–⟨1⟩4.

This is an ambient microscopic symmetry limit. It does not by itself remove the extra current and charge sectors or prove positivity of the raw finite Hamiltonian, which CA-84 disproves.

## 86.3 Local Fields in the Descendant OAR Representation

In the vacuum module of CA-84, grade selection gives the exactly finite compression

$$T_b(f) = P_b T(f) P_b = \sum_{|n|\le b} \widehat{f}_n P_b C_n P_b.$$

The CAR construction computes every retained matrix element using the safe schedule of CH-2. A computable smearing supplies only finitely many requested Fourier coefficients at each stage; no large many-body exponentiation was performed in the numerical work.

The algebra element $(e^{itT_b(f)}, 1)$ is represented as identity off $V_b$. Its strong limit is $e^{itT(f)}$: the zero-extended finite compressions converge on the essential core, and the preceding resolvent argument applies. The complementary scalar matters; a bare corner unitary would omit identity on the remaining Hilbert space.

Let $\mathcal{B}(I)$ be the sup-norm C*-algebra of sequences generated by these bounded unitary approximants with real smooth $f$ supported in $I$. The strong-star limit map is a $*$-homomorphism on this generated algebra. Its kernel is a closed ideal, and the quotient identifies with the C*-algebra generated by the limiting stress unitaries. Vacuum GNS weak closures give the sourced minimal-model Virasoro net and its complete sector data.

This represented field construction is additional to the genuine strict OAR norm limit $K(\mathcal{H}_{\mathrm{vac}}) + \mathbb{C}1$. Strong-star-null sequences are not generally norm-null, so the quotient is not relabelled as the norm-soft inductive limit of the source.

## 86.4 The Local Fusion-Space Question

The finite descendant matrices currently do not retain spatial intervals: finitely many Fourier moments of functions supported in any nonempty interval can be independently specified. Nor can refinement preserve every same-smearing compression exactly, since a smearing can have all low moments zero and higher moments nonzero.

These facts concern this specific encoding, not global fusion constraints in general. The next task is to identify a categorical fusion-space basis and local interactions/refinements that realize the required matrices and limits. No such identification or bounded-range implementation is claimed in the present work.

# 87 String-Net Histories as Trotter Expansions: the Spacetime Dictionary

## 87.1 Status, Sources, and Dependencies

**Status: derived.** Lemma 86.1 and Proposition 86.4 are checked local derivations; the face weights, branch data and central charges they use are source-local; the final subsection is a proposal. Reproducibility: the trigonometric identities of Proposition 86.4 were evaluated numerically for $L = 2, \dots, 5$ on both branches (worklog chunk 016); a Julia port is an open item.

Dependencies: CA-65, CA-66, CA-69, CA-74; CONVENTIONS.md (a), (b), (r), (ak).

## 87.2 The Levin–Wen Rules and What They Fix

**Source-local.** Levin and Wen specify a fixed-point wave function $\Phi$ on string-net configurations by four local linear relations: (1) topological invariance, (2) a closed type-$i$ loop contributes $d_i$, (3) a bubble on a string is $\delta_{ij}$, (4) the $F$-move. Rules (2)–(3) are motivated by scale invariance of an RG fixed point, rule (1) by the expectation that topological phases have topologically invariant fixed points, and rule (4) is the simplest completing constraint. The plaquette coefficients $a_s = d_s / \sum_i d_i^2$ are not chosen: they are the unique choice with a smooth continuum limit. The lesson carried forward is that a *principle* (scale invariance of the fixed point) fixes the model.

## 87.3 The Trotter Dictionary

**Setting (CONVENTIONS (r), (ak)).** $\mathcal{C}$ unitary fusion, $O = \mathbf{1} \oplus X$, $A_L = \mathrm{End}_{\mathcal{C}}(O^{\otimes L})$. A nearest-neighbour Hamiltonian is $H = -\sum_j h_j$ with $h_j \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ acting on sites $j, j+1$. Fix a basis of *elementary tiles* $D_\alpha$ of $\mathrm{End}_{\mathcal{C}}(O \otimes O)$ consisting of morphism diagrams (vacancy projectors, through-strings, cup, cap, hops, and for non-loop categories the fusion vertices), and expand $h_j = \sum_\alpha c_\alpha D_\alpha^{(j)}$ with $c_\alpha \in \mathbb{C}$. Brick-wall Trotter with step $\delta\tau$:

$$U(\delta\tau) = \prod_{j \text{ odd}} e^{\delta\tau h_j} \prod_{j \text{ even}} e^{\delta\tau h_j}, \qquad e^{-\beta H} = U(\delta\tau)^M + O(\beta\,\delta\tau) \quad (\beta = M\delta\tau)$$

on a finite chain, by the Lie–Trotter formula. A *history* $\Gamma$ assigns to each plaquette of the $M$-row brick lattice either the identity tile or one elementary tile $D_{\alpha(p)}$. Composing the tiles row by row gives a morphism $\mathrm{Ev}_{\mathcal{C}}(\Gamma) \in A_L$, the $\mathcal{C}$-evaluation of the spacetime string-net drawn by $\Gamma$.

**Lemma 86.1 [derived; tile expansion].** To first order per plaquette,

$$U(\delta\tau)^M = \sum_\Gamma \Big( \prod_{p:\, \alpha(p) \neq \mathrm{id}} \delta\tau\, c_{\alpha(p)} \Big)\, \mathrm{Ev}_{\mathcal{C}}(\Gamma) \;+\; R,$$

where $R$ collects every term carrying a second-order factor $O(\delta\tau^2 \|h_j\|^2)$ on at least one plaquette. **Proof.** $e^{\delta\tau h_j} = 1 + \delta\tau h_j + O(\delta\tau^2\|h_j\|^2)$ per plaquette. Expanding $\prod_p (1 + \delta\tau h_{j(p)})$ and each $h_j$ in tiles gives exactly the sum over histories with one tile per plaquette; composition of the tiles in the brick order is the composition in $A_L$, which is $\mathrm{Ev}_{\mathcal{C}}(\Gamma)$. The remainder collects all terms with a second-order factor on at least one plaquette. □

**Corollary 86.2 [the Levin–Wen rules are automatic].** $\mathrm{Ev}_{\mathcal{C}}(\Gamma)$ obeys rules (2), (3), (4): a closed $X$-loop evaluates to $d_X$ by the raw cup normalisation $v^\dagger v = d_X\, \mathrm{id}_{\mathbf{1}}$ of (r); a bubble between simple labels is $\delta$ by Schur; recoupling is the $F$-move. Rule (1), isotopy, holds for $\mathrm{Ev}_{\mathcal{C}}$ alone (planar

isotopy of morphism diagrams in a pivotal category) and fails for the full amplitude, which carries the geometric factor $\prod_p \delta\tau\, c_{\alpha(p)}$ depending on how many plaquettes each string traverses. Hence

$$\text{amplitude}(\Gamma) = \underbrace{\mathrm{Ev}_{\mathcal{C}}(\Gamma)}_{\text{topological}} \times \underbrace{\textstyle\prod_p w(p)}_{\text{geometric}}, \qquad w(\mathrm{id}) = 1,\ w(\alpha) = \delta\tau\, c_\alpha.$$

This is the spacetime reading made precise: rules (2)–(4) intact, rule (1) replaced by anisotropic tile weights (identity tiles weigh 1, all others $O(\delta\tau)$); isotropy can only emerge in the continuum.

**Proposition 86.3 [derived; exactness on an integrable line].** Suppose the tiles carry weights $\rho_\alpha(u)$ forming a face operator $X_j(u) = \sum_\alpha \rho_\alpha(u) D_\alpha^{(j)}$ with $X_j(0) = N(0)\cdot 1$, whose diagonal-to-diagonal (brick-wall) transfer matrices $T(u) = \prod_{j\ \mathrm{odd}} X_j(u) \prod_{j\ \mathrm{even}} X_j(u)$ commute for all $u$, as the sourced $\hat\tau_D$ does. Then $h_j = X_j'(0)/N(0) - (N'(0)/N(0))\,1$ satisfies $1 + \delta\tau h_j = X_j(\delta\tau)/N(\delta\tau) + O(\delta\tau^2)$, the Trotter product is $T(\delta\tau)^M$ up to normalisation, and $H$ commutes with $T(u)$ for every $u$. In particular the ground state of $H$ is a common eigenvector of the transfer matrices at every $u$, including the isotropic point.

**Proof.** The first two claims are Taylor expansion. For the third, the Taylor coefficients of a commuting one-parameter family commute with every member of the family, and $H$ is the first coefficient of $-\log T(u)$. □ On an integrable line no perturbative matching of rates is needed: $u$ interpolates exactly between the Hamiltonian limit and the isotropic point.

## 87.4 Dense Calibration

Take $X$ with $d = d_X$ and $H = -\sum_j e_j$, $e_j = b_j d_j$ the cup-cap generator of CA-66 ($e_j^2 = d\, e_j$). Because $e_j^2 \propto e_j$,

$$e^{\beta e_j} = 1 + w(\beta)\, e_j, \qquad w(\beta) = \frac{e^{\beta d} - 1}{d} = \beta + O(\beta^2),$$

so the brick-wall expansion is *exact* with two tiles per plaquette and

$$\langle \mathrm{out}|U^M|\mathrm{in}\rangle = \sum_\Gamma w^{\#\{e\text{-tiles}\}}\, d^{\#\{\text{closed loops}\}}\, \langle \mathrm{out}|\Gamma_{\mathrm{open}}|\mathrm{in}\rangle.$$

This is the sourced diagonal Temperley–Lieb transfer matrix at small spectral parameter: with $d = \sqrt{Q} = 2\cos\gamma$, $\log(Q^{-L/2}\epsilon_y^L \hat\tau_D(u)) = (u/\sin\gamma)\sum_j e_j + O(u^2)$, a commuting family, so Proposition 86.3 applies. For $d = \varphi$, $\gamma = \pi/5$, the antiferromagnetic sign is the golden chain, critical with $c = 7/10$ by the exact RSOS map. The dense chain has no continuous parameter at all: the category fixes $\gamma$ and the remaining choice is a sign.

## 87.5 Dilute Calibration

**Tiles (CONVENTIONS (ak)).** The dilute face operator is a linear combination of nine tiles. With in-nodes on the left and bottom edges (sites $j$, $j+1$) and out-nodes on the top and right edges, the tiles are, in the source's numbering: 1: both vacant, $P_{00}$; 2, 3: one through-string, $P_{X0}, P_{0X}$; 8: two through-strings, $P_{XX}$; 9: cap-then-cup, $e_j$; 4, 5: cap $d_j$ and cup $b_j$; 6, 7: hops $h_j^{\pm}$. The sourced weights, normalised so that $\rho_{1,2,3,8}(0) = 1$ and $\rho_{4,\ldots,7,9}(0) = 0$, are

$$\rho_1 = 1 + \tfrac{\sin u\,\sin(3\lambda - u)}{\sin 2\lambda \sin 3\lambda}, \quad \rho_{2,3} = \tfrac{\sin(3\lambda-u)}{\sin 3\lambda}, \quad \rho_{4,5} = \tfrac{\sin u}{\sin 3\lambda},$$
$$\rho_{6,7} = \tfrac{\sin u\,\sin(3\lambda-u)}{\sin 2\lambda \sin 3\lambda}, \quad \rho_8 = \tfrac{\sin(2\lambda-u)\sin(3\lambda-u)}{\sin 2\lambda \sin 3\lambda}, \quad \rho_9 = -\tfrac{\sin u\,\sin(\lambda-u)}{\sin 2\lambda \sin 3\lambda},$$

with loop fugacity $n = -2\cos 4\lambda$. The $u = 0$ values confirm the tile assignment: $X_j(0) = P_{00}+P_{X0}+P_{0X}+P_{XX} = 1$, matching the sourced factorisation into identity-arc triangles. The Hamiltonian limit $H_{\rm dil} = -\sum_j X_j'(0)$ is

$$H_{\rm dil} = \sum_j \Big[ -\tfrac{P_{00}}{\sin 2\lambda} + \cot 3\lambda\,(P_{X0}+P_{0X}) + \tfrac{\sin 5\lambda}{\sin 2\lambda \sin 3\lambda} P_{XX} - \tfrac{b_j+d_j}{\sin 3\lambda} - \tfrac{h_j^+ + h_j^-}{\sin 2\lambda} + \tfrac{\sin\lambda}{\sin 2\lambda \sin 3\lambda}\, e_j \Big].$$

Every coupling is a function of $\lambda$ alone. For a Temperley–Lieb type category with $d = 2\cos(\pi/(L+1))$ the equation $n = d$ has the two solutions $\lambda = \frac{\pi}{4}\frac{L}{L+1}$ (branch 1) and $\lambda = \frac{\pi}{4}\frac{L+2}{L+1}$ (branch 2), both critical, with $c = 1 - 6/(h(h-1))$, $h = L+2$ resp. $h = L+1$. Fibonacci is $L = 4$: branch 1 gives $c = 4/5$, branch 2 gives $c = 7/10$.

**Proposition 86.4 [derived; honeycomb reduction at $u = \lambda$].** At $u = \lambda$ the nine-tile model is the honeycomb $O(n)$ loop model with edge fugacity $x = 1/(2\cos\lambda)$. On branch 1 this is Nienhuis's critical point $x_c(n) = 1/\sqrt{2+\sqrt{2-n}}$; on branch 2 it is the second-branch value $1/\sqrt{2-\sqrt{2-n}}$.

**Proof.** Split each square tile along the diagonal from the bottom-left to the top-right corner into two trivalent vertices, $A$ with edges $\{\text{top}, \text{left}, \text{diag}\}$ and $B$ with $\{\text{bottom}, \text{right}, \text{diag}\}$. Give each occupied half-edge of a tile weight $x^{1/2}$ and the internal diagonal weight $x$. Then the tiles weigh: $1 \mapsto 1$; $2, 3 \mapsto x$ (an arc inside $A$ or $B$); $4, 5, 6, 7 \mapsto x^2$ (an arc through the diagonal); $8 \mapsto x^2$; $9 \mapsto 0$ (it would use the diagonal twice). At $u = \lambda$: $\rho_9 = 0$; $\rho_4 = \rho_5 = \rho_6 = \rho_7 = \rho_8 = \sin\lambda/\sin 3\lambda$; $\rho_{2,3} = \sin 2\lambda/\sin 3\lambda$; $\rho_1 = 1 + \sin\lambda/\sin 3\lambda = 2\sin 2\lambda\cos\lambda/\sin 3\lambda$. Hence $\rho_2/\rho_1 = 1/(2\cos\lambda) =: x$ and $\rho_4/\rho_1 = \sin\lambda/(4\sin\lambda\cos^2\lambda) = x^2$, consistent. Finally $x^{-2} = 4\cos^2\lambda = 2 + 2\cos 2\lambda$ and $\sqrt{2-n} = \sqrt{2+2\cos 4\lambda} = 2|\cos 2\lambda|$, so $x = x_c$ when $\cos 2\lambda \geq 0$ (branch 1, $\lambda < \pi/4$) and $x = 1/\sqrt{2-\sqrt{2-n}}$ when $\cos 2\lambda < 0$ (branch 2). □

**Checked arithmetic (Fibonacci, $L = 4$).**

| branch | $\lambda$ | $x = 1/(2\cos\lambda)$ | Nienhuis value | $c$ | couplings $(-\rho_1', -\rho_2', -\rho_4', -\rho_6', -\rho_8', -\rho_9')$ |
|---|---|---|---|---|---|
| 1 | $\pi/5$ | 0.618034 | $x_c = 0.618034$ | 4/5 | $(-1.0515,\ -0.3249,\ -1.0515,\ -1.0515,\ 0,\ +0.6498$ |
| 2 | $3\pi/10$ | 0.850651 | $\tilde{x}_c = 0.850651$ | 7/10 | $(-1.0515,\ -3.0777,\ -3.2361,\ -1.0515,\ -3.4026,\ +2.7$ |

The same identities hold for $L = 2, 3, 5$ (worklog 016). Note the sign of the $e_j$ term: the integrable dilute chain has a *repulsive* cup-cap term on both branches, opposite to the dense golden chain, and on branch 1 the $P_{XX}$ coupling vanishes because $\sin 5\lambda = \sin\pi = 0$.

## 87.6 What the Calibration Says About the Principle

For Temperley–Lieb type input the machine closes with no continuous choice. The category fixes $n = d$, hence $\lambda$ up to a branch; the couplings of $H_{\rm dil}$ are functions of $\lambda$; the Hamiltonian, the isotropic point $u = 3\lambda/2$ and the honeycomb point $u = \lambda$ are one commuting family; the limit is critical with a sourced $c$. Isotropy is restored along the family, not imposed. The geometric parameter is the anisotropy $u$, not a coupling. The residual freedom is the branch: two critical points for $u > 0$, two more with an Ising factor for $u < 0$. Whether a principle selects among branches is open.

## 87.7 Beyond Loops: the Fibonacci Net Gas (Proposal)

For Fibonacci on $O = \mathbf{1} \oplus \tau$, $O \otimes O = 2\cdot\mathbf{1} \oplus 3\cdot\tau$, so $\mathrm{End}(O\otimes O) = M_2 \oplus M_3$ has dimension 13, while $\mathrm{dTL}_2(\varphi)$ has dimension 9 (CA-69). The four missing dimensions are the vertex tiles $\tau\otimes\tau \leftrightarrow \tau$ with one vacancy, which CA-74 proves lie outside the dilute loop algebra. The general reflection-symmetric nearest-neighbour Hamiltonian therefore has, beyond the loop couplings above, a vertex coupling $v$:

$$h = \big[\mathrm{diag}(\epsilon_{00}, \epsilon_1) + w(b+d)\big]_{\mathbf{1}\text{-block}} \oplus \big[\mathrm{diag}(\mu, \mu, \epsilon_\tau) + t(h^+ + h^-) + v(\nu + \nu^\dagger)\big]_{\tau\text{-block}},$$

with $\nu : |\tau\tau \to \tau\rangle \mapsto |\tau\mathbf{1}\rangle + |\mathbf{1}\tau\rangle$ built from the vertex morphism of $\mathcal{C}$ (normalisation to be fixed in CONVENTIONS before use). Lemma 86.1 turns any such $h$ into a *trivalent net gas* in spacetime with per-tile weights $\delta\tau \cdot$ (coupling) and the $\mathcal{C}$-evaluation of the net. Two corners are calibrated on the quantum side: $\mu \to -\infty$ freezes every site to $\tau$ and recovers the dense chain ($c = 7/10$ or $4/5$ by sign); the $v = 0$ hyperplane is the parity-preserving dilute loop sector of CA-74, which contains $H_{\mathrm{dil}}$. No integrable line is known once $v \neq 0$: the sourced statement is that beyond Temperley–Lieb, integrability requires further fine-tuning. Proposition 86.3 is therefore unavailable, and the comparison of rates is genuinely first-order: the open problem is to find the locus in $(\epsilon, \mu, w, t, v)$ at which the net gas is critical. The first analytical tool to point at it is a lattice holomorphicity condition on a $Z(\mathcal{C})$-spin defect endpoint, which is the next shard.

**Erratum / qualification (2026-10-22).** (a) The brick sweep $T(u)$ of Proposition 86.3 is Koo–Saleur's *row-to-row* $\hat{\tau}$, which on the homogeneous self-dual line does *not* commute; the commuting family is the sequential diagonal product $\hat{\tau}_D = \prod_j X_{2j} X_{2j+1}$. The conditional implication (commuting family $\Rightarrow [H, T(u)] = 0$) stands and applies to $\hat{\tau}_D$; the attribution "as $\hat{\tau}_D$ does" does not. (b) "The ground state is a common eigenvector" needs a nondegenerate ground state. Both qualifications reach the dense calibration and "What the Calibration Says About the Principle". Details: CA-78, Erratum to Proposition 86.3.

# 88 The Net Case: the Dilute Fibonacci Chain Is the Golden Chain Blocked

## 88.1 Status, Sources, and Dependencies

**Status: derived and checked.** Lemma 87.1 is a categorical identity; Theorem 87.2 is derived by hand and checked by the reproducible run `runs/2026-09-23-net-tl-representation/` (script `scripts/python/net_tl_representation_check.py`, INDEX.md); Proposition 87.3 assembles three sourced identities; the last subsection is a proposal.

Dependencies: CA-65, CA-66, CA-69, CA-74, CA-86; CONVENTIONS.md (b), (r), (aj), (ak).

## 88.2 Blocking: When Dilute Is Dense in Disguise

**Lemma 87.1 [derived].** Let $\mathcal{C}$ be a unitary fusion category and $O = \mathbf{1} \oplus X$ the site object of (r). If there is a simple $Y$ with $Y \otimes \bar{Y} \cong \mathbf{1} \oplus X$, then for every $L$

$$O^{\otimes L} \;\cong\; (Y \otimes \bar{Y})^{\otimes L}, \qquad \mathrm{End}_{\mathcal{C}}(O^{\otimes L}) \;\cong\; \mathrm{End}_{\mathcal{C}}\left((Y\bar{Y})^{\otimes L}\right),$$

and a nearest-neighbour operator on $O$-sites $j, j+1$ is a range-four operator on the alternating $Y\bar{Y}$ chain, aligned to pairs. For self-dual $Y$ the right-hand side is the dense $Y$-chain on $2L$ sites.

**Proof.** Tensor the isomorphism $O \cong Y \otimes \bar{Y}$ $L$ times; End is functorial; a morphism on $O \otimes O \cong (Y\bar{Y})^{\otimes 2}$ has support on four consecutive $Y$-sites. □

**Instances.** Fibonacci: $\tau \otimes \tau = \mathbf{1} \oplus \tau$, so the dilute $\mathbf{1} \oplus \tau$ chain of $L$ sites is the golden chain of $2L$ sites. The dimensions agree: CA-66 gives $\dim \mathcal{H}_L = F_{2L+1}$ for the dilute chain and $\dim \mathrm{End}(O^{\otimes L}) = F_{4L-1}$, which are the fusion-path count and $\sum_c m_c^2$ of $\tau^{\otimes 2L}$. Ising: $\sigma \otimes \sigma = \mathbf{1} \oplus \psi$. $SU(2)_k$: $\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1$, so the dilute spin-1 chain is the dense spin-$\frac{1}{2}$ chain. Haagerup $H_3$ with $X = \rho$: the sourced rule $\rho^2 = \mathbf{1} + (\mathbf{1} + a + a^2)\rho$ gives $d_\rho^2 = 1 + 3d_\rho$, so $d_\rho = (3 + \sqrt{13})/2$ and the invertibles have dimension 1; no simple $Y$ has $d_Y^2 = 1 + d_\rho = (5 + \sqrt{13})/2$, so no such $Y$ exists.

**Consequence for CA-69/CA-74.** The correct two-site algebra of the dilute Fibonacci chain is $\mathrm{End}(\tau^{\otimes 4})$, the Fibonacci quotient of $\mathrm{TL}_4(\varphi)$, of dimension 13; the dilute algebra $\mathrm{dTL}_2(\varphi)$ of dimension 9 is its parity-preserving subalgebra. The "vertex tile outside dTL" of CA-74 is a Temperley–Lieb generator acting across a pair boundary. For Fibonacci there is no net gas beyond loops: every trivalent history is a loop history of the blocked chain.

## 88.3 The Temperley–Lieb Generators in Tile Form

Write $\mathrm{TL}_{2L}(\varphi)$, $e_i^2 = \varphi e_i$, $e_i e_{i\pm1} e_i = e_i$, on the dilute chain as

$$e_{2j-1} = \varphi\, P_0^{(j)} \;(j = 1, \ldots, L), \qquad e_{2j} = F_{j,j+1} \;(j = 1, \ldots, L-1),$$

with $P_0^{(j)}$ the vacancy projector of site $j$ (the pair $\tau_{2j-1}\tau_{2j}$ fused to $\mathbf{1}$) and $F \in \mathrm{End}(O \otimes O)$ unknown. In the pair basis $\{|\mathbf{11}\rangle, |\tau\tau; \mathbf{1}\rangle\} \oplus \{|\mathbf{1}\tau\rangle, |\tau\mathbf{1}\rangle, |\tau\tau; \tau\rangle\}$ of (aj) the relations $e_{2j\pm1}e_{2j}e_{2j\pm1} = e_{2j\pm1}$ force the vacancy-diagonal entries $\langle \mathbf{11}|F|\mathbf{11}\rangle = \langle \tau\mathbf{1}|F|\tau\mathbf{1}\rangle = \langle \mathbf{1}\tau|F|\mathbf{1}\tau\rangle = \varphi^{-1}$; $F^2 = \varphi F$ with reflection symmetry then forces rank one in each block.

**Theorem 87.2 [derived; checked].** Up to the sign of the vertex entry, the unique reflection-symmetric solution is

$$F = \varphi\, |u\rangle\langle u| \;\oplus\; \varphi\, |w\rangle\langle w|, \qquad u = (\varphi^{-1},\ \varphi^{-1/2}), \qquad w = (\varphi^{-1},\ \varphi^{-1},\ \pm\varphi^{-3/2}),$$

with the cup sign + tied to the $F$-gauge of (b). All Temperley–Lieb relations, including $[F_{j,j+1}, F_{j+1,j+2}] = 0$, which involves the $F$-move, hold to machine precision for $L = 3, \ldots, 6$; the opposite cup sign fails

at $7.6\times10^{-1}$; a 1% mutation of the vertex entry fails at $3\times10^{-3}$. Moreover $H=-\sum_i e_i$ on the dilute chain of $L$ sites and the golden chain $-\varphi\sum_j P_{\mathbf{1}}^{(j,j+1)}$ on $2L$ sites have identical dimension $F_{2L+1}$ and identical spectra:

| $N=2L-1$ | dim | $E_0$ | first gaps |
|---|---|---|---|
| 5 | 13 | $-6.5141330104$ | 0.746587, 1.891335, 2.463767 |
| 9 | 89 | $-11.4453171319$ | 0.483796, 1.270064, 1.624217 |
| 15 | 1597 | $-18.8535471512$ | 0.316696, 0.841769, 1.066473 |

in both representations, digit for digit (run bundle). This is Lemma 87.1 made explicit: $F_{j,j+1}=\varphi P_{\mathbf{1}}^{(2j,2j+1)}$ rewritten in the pair basis. In tile language every one of the six couplings of CA-86 is on:

$$F=\tfrac{1}{\varphi}P_{00}+\tfrac{1}{\varphi}(b+d)+P_{\mathbf{1}}^{\tau\tau}\ \oplus\ \tfrac{1}{\varphi}(P_{0\tau}+P_{\tau 0})+\tfrac{1}{\varphi}(h^{+}+h^{-})+\varphi^{-7/4}(\nu^{R}+\nu^{L}+\text{h.c.})+\varphi^{-2}P_{\tau}^{\tau\tau},$$

with the raw cup $b$ of (r) and the (aj)-normalised vertex tiles.

**Colour reading.** $u_1^2=\varphi^{-2}=1/Q$, $u_2^2=\varphi^{-1}=(Q-1)/Q$, $w_1^2=1/Q$, $w_3^2=\varphi^{-3}=(Q-2)/Q$ with $Q=\varphi+1$: these are the probabilities that a Potts spin flipped uniformly over $Q$ colours stays, creates a wall pair, hops a wall, or splits a wall into two. $F$ is the flip generator of the $Q=\varphi+1$ Potts chain in domain-wall variables, and $H$ is that chain at its self-dual point. The gap ratios at $N=15$, 2.658 and 3.368, sit near 8/3 and 10/3, the ratios $(3/5+1)/(3/5)$ and $2/(3/5)$ expected from the weights 3/5 and the level-two vacuum descendant of the sourced golden-chain spectrum.

## 88.4 The Isotropic Net Gas Is the $Q=\varphi+1$ Potts Model

**Proposition 87.3 [derived from sourced identities].** With the vertex normalisation (aj), the isotopy-invariant evaluation of a planar Fibonacci net $N$ with $V_3$ trivalent vertices is

$$\langle N\rangle=\varphi^{-2}\,\varphi^{\frac{3}{4}V_3(N)}\,\chi_{\widehat{N}}(\varphi+1),$$

$\chi_{\widehat{N}}$ the chromatic polynomial of the dual graph. Hence the isotropic net gas with edge fugacity $x$ and vertex fugacity $y=\varphi^{-3/4}$ is

$$\sum_N x^{E(N)}y^{V_3(N)}\langle N\rangle=\varphi^{-2}\,Z_{\text{Potts}}(Q=\varphi+1,\ K=x),$$

the low-temperature expansion of the $Q=\varphi+1$ Potts model with domain walls on the net lattice and spins on its dual.

**Proof.** The sourced formula reads $\langle G\rangle=\varphi^{-5}\varphi^{\frac{3}{2}F(G)}\chi_{\widehat{G}}(\varphi+1)$ for a connected net, $F(G)$ its number of faces. Euler with $3V_3=2E$ gives $F=2+\frac{1}{2}V_3$, so $\langle G\rangle=\varphi^{-2}\varphi^{\frac{3}{4}V_3}\chi_{\widehat{G}}$. Checks: a loop ($V_3=0$, $\chi=Q(Q-1)=\varphi^3$) gives $\varphi$; a theta ($V_3=2$, $\chi=Q(Q-1)(Q-2)=\varphi^2$) gives $\varphi^{3/2}$, the theta value of (aj). For $k$ components $\langle N\rangle$ is multiplicative while $\widehat{N}$ is a $k$-fold one-point union, so $\chi_{\widehat{N}}=\prod_i\chi_{\widehat{N}_i}/Q^{k-1}$; the factors $\varphi^{-2k}$ and $Q^{k-1}=\varphi^{2(k-1)}$ combine to $\varphi^{-2}$. The sourced low-temperature expansion $Z_{\text{Potts}}=\sum_N K^{E(N)}\chi_{\widehat{N}}(Q)$ then matches term by term at $y=\varphi^{-3/4}$. □

Since $Q=\varphi+1<4$, the model has a critical point, in the universality class of completely packed loops with weight $\sqrt{Q}=\varphi$, i.e. the dense branch of CA-86 with $c=7/10$. The sourced condition $e^{\beta_c J}-1=\sqrt{Q}$ is the square-lattice self-dual value; the triangular-lattice value for honeycomb domain walls needs a local source (open). Two remarks. Chromatic polynomials at non-integer $Q$

take both signs, so the net-gas weights are not positive: positivity lives in the fusion-path basis of the blocked chain, not in the net basis, exactly as the strange-correlator source warns. And the critical net gas is, by Lemma 87.1, the golden chain again, with the Potts critical point as its self-dual point.

## 88.5 What This Does to the Principle

Every calibrated critical point of the Fibonacci $\mathbf{1} \oplus \tau$ chain is now a point of the pair-aligned range-four golden chain family:

| point | identity | $c$ |
|---|---|---|
| dense, antiferromagnetic | golden chain | 7/10 |
| dense, ferromagnetic | $Z_3$ parafermion (sourced) | 4/5 |
| dilute $A_2^{(2)}$ branch 1 (CA-86) | range-four integrable point | 4/5 |
| dilute $A_2^{(2)}$ branch 2 (CA-86) | range-four integrable point | 7/10 |
| Potts / isotropic net gas | golden chain, self-dual | 7/10 |
| hard-hexagon strange correlator (sourced) | tau-projected string-net | 4/5 |

Two universality classes, both minimal models with Fibonacci defect lines. For Fibonacci the trivalent vertex adds nothing that Temperley–Lieb does not already contain, and Intuition 3's "forced" creation and annihilation is the statement that pairs of $\tau$ fuse to the vacuum. The net-gas question is therefore not void in general but void for Fibonacci.

**Proposal.** The genuine net case is a category whose site object is not a product: Haagerup with $O = \mathbf{1} \oplus \rho$, for which $\mathrm{End}(O \otimes O)$ is not a Temperley–Lieb quotient and the vertex tiles $\rho \otimes \rho \to \rho$, $\alpha\rho$, $\alpha^2\rho$ carry three independent couplings. The blocked-chain trick is unavailable, no integrable line is known, and the isotropic net evaluation is not a chromatic polynomial. This is the concrete form of the acid test: the first computation is the Haagerup analogue of Theorem 87.2, namely whether any tile combination in $\mathrm{End}(O \otimes O)$ generates a known algebra with a critical Hamiltonian, and the second is the lattice holomorphicity condition of the sweep. Open items: Julia port of the run bundle; a local source for the triangular-lattice Potts critical coupling; and whether the range-four golden-chain family has any critical point outside $\{7/10, 4/5\}$.

# 89 The Layered Target Grid: Setting, Input Data, and Target Rungs

## 89.1 Status, Sources, and Dependencies

**Status: definitions and placement [proposal]; nothing is proved or promoted.** This shard implements the layering principle (AGENTS.md, Rule 17): every statement names its chirality setting, its input-data layer and the target rung it reaches. Definitions 88.1–88.3 are organisational conventions; Remark 88.4 is a trivial derived monotonicity; every placement inherits the grade of its source shard or review and changes none. The six restatements of §89.4 are *proposals* built from the reviewers' reformulations. The "CA-88" IDs proposed in the D1, D2, D4, D5 and D6 integration packages are superseded by this shard. *Round 2 (2026-10-22):* CA-89–CA-94 are placed with the tags verified in their reviews; verdicts in §89.5. No placement upgrades a review grade.

Dependencies: CA-68, CA-71–CA-74, CA-75 (layering paragraph), CA-76–CA-82, CA-86, CA-87, CA-89–CA-94; LC-1–LC-13; CONVENTIONS (r), (u), (af), (ak), (aq).

## 89.2 Three Axes

Throughout, $\mathcal{C}$ is a unitary fusion category. Microscopic choices (site object $O$, tiles $h \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$, boundary and seam, refinements, states; D5's E5, NOTE.md:124–127) are *declarations at every layer*, not a data layer: they are part of a construction, not of its input.

**Definition 88.1 [proposal; setting axis S].** $\mathsf{S}_{\mathrm{ch}}$: an *open* chain with declared endpoints, targeting a *chiral* conformal net $\mathcal{A}$ (one stress family). Fidelity means: the sector category $\mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$ is braided equivalent to a declared chiral UMTC $\mathsf{B}$ ($\mathsf{B} = \mathcal{C}$ for modular $\mathcal{C}$), as in Theorem 80.9. $\mathsf{S}_{\mathrm{full}}$: a *periodic* chain targeting a full two-dimensional local net with chiral subnets $\mathcal{A}_L, \mathcal{A}_R$, two stress families ($\mathrm{Vir} \times \overline{\mathrm{Vir}}$) and a declared gluing (extension; for the diagonal case the Cardy invariant $M = 1$, D5 P6). Fidelity means: the lattice $\mathcal{C}$-symmetry, i.e. the periodic defect operators that commute with the transfer matrix and fuse by $N_{ab}^{c}$ (AFM-D:163–212), converges into the full theory's symmetry (defect) data, and the declared identifications hold: $\mathrm{Rep}(\mathcal{A}_L) \simeq \mathcal{C}$ for modular $\mathcal{C}$ (D6 REVIEW:94–98), with two-chiral bulk data $\mathcal{C} \boxtimes \overline{\mathcal{C}} \simeq Z(\mathcal{C})$ (Bischoff:661–664), the sector category of the doubled chiral net or the centre of the defect category $\mathcal{C}$, *not* the DHR category of the full diagonal extension, which is trivial (Corollary 93.9, Proposition 93.10); for fusion-only $\mathcal{C}$ the braided target is $Z(\mathcal{C})$, with $\mathcal{C}$ recovered as defects/solitons (Bischoff:384–397). *Warning (PLAN.md:244–247):* the periodic lattice algebra's scaling limit is the interchiral algebra, not $\mathrm{Vir} \times \overline{\mathrm{Vir}}$; the open/periodic choice is part of the setting, not a boundary detail. *Source gap (Law 1):* no local source settles whether a 1D local tower can have a chiral limit with $c \neq 0$ ($c_L \neq c_R$); the nearest is 1903.12334:76 (D6 REVIEW:86–90). *Delineation (CA-91; full; category-free;* R2 *[cond.]):* a complex-*linear* unitary parity commuting with the control, with both standard KS families VR-1-certified on one system, forces $c_L = c_R$, $c_L^* = c_R^*$ (Theorem 91.1) and excludes an absent right stress (Corollary 91.10); an antiunitary (dagger-type) reflection preserves each chirality and constrains nothing (Remark 91.4). Parity-odd perturbations exist iff the total current is nonzero: never on the fixed carrier $O = \mathbf{1}$ (Negative 91.12), always after $O \otimes (\mathbf{1} \oplus \mathbf{1})$ (91.13) [L0, finite]. The general lattice-chirality question remains a source gap, so $\mathsf{S}_{\mathrm{ch}}$ is neither excluded nor established; it is a declared tag.

**Definition 88.2 [proposal; data ladder L].** L0 *fusion*: $\mathrm{Irr}\,\mathcal{C}$, $N_{ab}^{c}$, $F$, dagger, pivotal/spherical structure, $d_a > 0$. L1 *braided*: in addition $R$ obeying the hexagons. L2 *modular*: $R$ nondegenerate (Bischoff:655–658); hence $S$, twists $\theta_i$, and $c$ mod 8 (EG:334). L3 *enriched* E: a real lift $c$ ($c_L, c_R$ in $\mathsf{S}_{\mathrm{full}}$), weights $h_i$ with $e^{2\pi i h_i} = \theta_i$, and characters or a finite presentation of the vacuum chiral algebra (D5 E2–E3, NOTE.md:107–118; D6 E.energy/E.fields, NOTE.md:14–17); in $\mathsf{S}_{\mathrm{full}}$ also the

sewing data E4 (NOTE.md:119–123). L4 *declared target theory*: an effective sewing completion $\mathcal{S}$ (D4-C's input, D4 NOTE.md:11–21) or a sourced rigorous net (CA-82). The ladder is not fully linear: for fusion-only $\mathcal{C}$, L0 already determines the modular $Z(\mathcal{C})$ (Bischoff:384–386), which is why $\mathsf{S}_{\text{full}}$ reaches modular-type targets from fusion input, whereas L3 is never determined by L2: $c$ only mod 8, and tensoring with a holomorphic net preserves the category (Bischoff:374–381; `realisation_and_fidelity.md`:39–44).

**Definition 88.3 [proposal; target rungs R].** Minimal data and current certificate source per rung:

| Rung | Content | Min. data | Certificate in repo |
|---|---|---|---|
| $\mathsf{R0}^c$ | local isometric/corner refinement tower | L0 | CA-68, CA-71 [derived] |
| $\mathsf{R0}^u$ | local *unital* injective $*$-refinements | L0+half-braiding on $\langle O\rangle$ (so L1) | 76.2 unital, nonlocal; 76.4 all $\mathcal{C}$; CA-90 local [derived] |
| R1 | a Wilson-triangle limit state | none | WT-1 (CA-81); drift via 76.1, 76.6, 90.W [cond.]; 90.V trivial vacuum only |
| R2 | Virasoro / extended-mode rep. on the same GNS system | none (Vir); L3 (extended) | VR-1 (CA-81); 79.2 via P0–P3 [cond.]; 89.1 transfer interfaces, Virasoro form forced by consistency [cond.]; 91.1, 91.5 two families [cond.]; 78.13 [proposal, split by CA-89] |
| R3 | local conformal theory (positive energy, integration, net axioms) in the declared setting | none | D2-C's conclusion; ingredients 80.2–80.5, 92.3/92.5 (spatial RP, finite), 78.9–78.11; no categorical instance |
| R4 | faithful to declared $(\mathcal{C}, \mathsf{E})$ | L0+L4 (sectors); L1+L4 (braided); L3 ($T$, characters) | 80.9 + 93.8 [cond. on R3 plus lattice certificates 93.11/93.12]; 93.2 closes the PP gap; 80.8: fusion operators never reach R4 |
| R5 | coherent/functorial over enrichment-preserving equivalences | L3 | D6-C clause; none |

R4 is itself layered: defect fidelity ($\mathcal{C}$ acts faithfully by defects; L0, D2 §5, Lemma 80.7) is necessary, never sufficient (Prop. 80.8); sector fidelity as a tensor equivalence needs only L0+L4 and Theorem 93.8's (F1), ($\text{F2}_0$), (F3′); braided sector fidelity needs L1+L4, modularity being a conclusion (CA-93); matching $T$ and characters needs L3 (80.9 closing remark); identification with one net needs L4. $S, T$ agreement alone is the *minimum* target, not categorical fidelity (`realisation_and_fidelity.md`:46–56). $\mathsf{R0}^u$ at bare L0: needs only a relative half-braiding on $\langle O\rangle$; one-gap lifts are obstructed when $[O]$ is not central in $K(\langle O\rangle)$ (90.2); reached at bare L0 for the CA-96 families, open for general $\mathcal{C}$ (96.10). A seam pairing (CA-92) shows the same split: it exists iff $O \otimes \bar{O}$, $\bar{O} \otimes O$ have equal multiplicities (92.2), so for self-dual $O$ at L0; expected at L1 (ribbon identity not locally sourced); it can fail for non-self-dual $O$ in non-braided $\mathcal{C}$ ($\text{Vec}_{S_3}$).

**Remark 88.4 [derived; monotonicity].** A statement proved at L$k$ holds verbatim at every L$k' \geq$ L$k$ by forgetting data. A rung certificate implies the lower rungs only if all are certified on the *same* refinement, state and representation system (CA-81, "same state and representation system"); separately obtained R1 and R2 certificates do not compose.

## 89.3 Existing Results on the Grid

Grades are those of the source shard; "$\to$ R$j$" marks a finite or conditional ingredient that feeds rung $j$ without reaching it; "any" means the statement is setting-agnostic.

| Result | S | L | R | Comment |
|---|---|---|---|---|
| CA-71–CA-74, CA-68 | open | L0 | R0$^c$ | July tower: corner maps, residual calculus, JW kernel; nonunital |
| WT-1, VR-1 (CA-81) | any | none | R1; R2 | category-free contracts; must share one system |
| 76.1–76.4, 76.9 | open | L0 | → R1 | drift from corner weight; unital extension; local first tangent |
| 76.6–76.7 | open | L0 | → R1 | gap route [cond.]; 76.5 superseded by CA-90 |
| 76.10 (D3-C); 76.12 | open | L0 | → R0, R1 | UV locality [conj.]; dichotomy (G)/(L) [open] |
| CA-77 | any | L0 | R0$^c$ | pointed carrier, grading, dual-pair blocking, Fock carrier |
| CA-78 | any (D2-C full) | L0 (L1 for 78.8) | → R2, R3 | finite selection equations; interface lemmas [cond.]; D2-C |
| CA-79 | full | L0; 79.9 L3 | → R2; → R4 | shell certificate 79.2 [cond.]; no-go 79.8; $c > 1$ growth |
| CA-80 | any; 80.9 ch | L0; 80.9 L2/L3 | → R3; → R4 | sewing, reflection positivity [derived]; 80.9 assumes (F1)= R3 |
| CA-82 | ch | L4 | R4 | Ising calibration; R4 by cited net theorems, reads no $F, R$ |
| CA-83–CA-85 | ch | L4 | R2–R3 | coset calibrations; scalar-augmented, no local fusion-space encoding |
| CA-86, CA-87 | any | L0 | finite | Trotter dictionary; blocking lemma; no tower, no limit |
| CA-89 | full (per family) | none | → R2 | transfer 89.1(i), derivative A3 89.1(ii), cocycle 89.1(iii) [cond.]; 89.3, 89.7 [negative] |
| CA-90 | any | L0+$t$ (L1) | R0$^u$ | local unital lift [derived]; 90.2 one-gap no-go [negative] |
| CA-91 | full | none; L0 for 91.11–91.15 | R2 [cond.]; → R0 | parity ledger balance 91.1; mixed brackets ⇔ ZC (91.5); delineation 91.10; finite parity-odd ingredients |
| CA-92 | full (ch: twisted refl.) | L0 + decl. $s$ | finite, → R3 | spatial seam kernel, Schur criterion [derived, cond. on $s$]; finite OS [derived]; $F \circ \dagger$ crossing, perpendicular commutator ill-typed [negative] |
| CA-93 | any; 93.8 ch; 93.9 full | none; 93.8 L0/L1+L4; 93.9 L2+L4 | → R4; R4 given R3 | normal limit expectation [derived]; repaired two-sided fidelity; doubled net vs LR extension; EP/AN [cond.] |
| CA-94 | any (open) | L0 + decl. | R0$^c$ + $z$=2 | D1-C proved for tuned family [derived]; N1–N3 [negative]; fixed-$h$ E6 false; $\mathcal{C}$ not inert: $G_r$ bounds range-$r$ number change (94.11) |
| LC-1–LC-3, LC-5–LC-13 | ch | L2–L3 | → R3, R4 | continuum equations only; declared lifts (LC-8, LC-11) |
| LC-4 | full | L0 | → R1, R2 | periodic KS tier; both chiral families |

## 89.4 The Six Headline Conjectures, Restated

Each entry gives the tags (S; L; rung reached / rung claimed) of the headline *as written*, then a restatement [proposal] at the most general layer where it is plausibly true, the first attackable lemma, and the next layer's demand. All six were graded "needs reformulation" (INTEGRATION.md §(d)).

**D1-C** (open; L0 + declared reaction morphisms and contact tuning; carrier (77.8) plus an off-ladder $z$=2 resolvent limit reached (CA-94) / a $z = 2$ resolvent limit, off the conformal ladder). *Restated (*L0*, any,* R0$^u$ + R1*):* on the pointed configuration carrier of 77.8, for any unitary fusion

$\mathcal{C}$ and $O = \mathbf{1} \oplus X$, there are coherent local unital refinements of a number-changing field algebra compatible with one state family satisfying WT-1 (D1 INTEGRATION:102–105; CA-77 open problem). The collision-sewing version is now **derived** for the tuned family (CA-94, Theorem 94.6), and the tuning is necessary: a fixed pure-hop density has no full-Fock limit after any scalar subtraction (Proposition 94.1); hard-core hopping gives Dirichlet collisions and decoupled sectors (CA-77 §Conjecture). The former first lemma (collision-trace bound for $N \leq 2$, INTEGRATION:107–111) is superseded by Lemma 94.5 for all $N$. *First lemma (restated target, open):* a unital local lift of the dressed refinement $R = U(\mathrm{id} \otimes \eta)$ (CA-77:209–212) commuting with $\hat{N}$ (E6 INTEGRATION:87–89; CA-90 gives the bare-insertion version). *Next layer:* R2 needs KS/Ward data D1 does not supply; R4 needs braided/ribbon marking of the tower (L1/L2; INTEGRATION (e)7, (e)9).

**D2-C** (full; L0 + declared certificate C1–C6; none reached / R3). Split in two. *(i) Ward-to-KS transfer lemma (category-free, full, → R2):* the operator lift W2 of the discrete contour identity, plus A2 for an admissible control and differentiated, coordinate-resolved matching, implies VR-1 A3/A4 for the standard KS modes of $k$ on the same GNS core (Theorem 89.1); A3 for arbitrary gate-intertwining refinements is false when $c^* \neq 0$ (Proposition 89.7). *(ii) Selection proper:* which $h \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ carries a certificate; at L0 it is category-blind ($\mathcal{C} = \mathrm{Vec}$ admits every qudit chain, D2 REVIEW:79) and remains a question. *First lemma:* the probe-word Ward-to-residual Lemma 89.L, tested first on the in-repo free-fermion benchmark with $K = 1 + E_N$ (D2 INTEGRATION:117–121). *Next layer:* chiral-sector equality with $\mathcal{C}$ is not in D2-C (NOTE.md:40–45); it is R4 at L2, plus a $c$ lift at L3 (INTEGRATION (e)13).

**D3-C** (open; L0, pointed $O$; first order reached by 76.9 / uniform locality, → R0). *Restated (L0, open, → R0):* a UV locality lemma: the polar heat refinement has a quasi-local generator uniformly in $L$ for $0 \leq t \leq t_0$ (Conjecture 76.10 as stated). The IR question is separate and is the dichotomy 76.12: (G) global fidelity at rate $o(a_n^{1+z})$, or (L) a local unital lift with state-adapted witnesses. *First lemma:* a locally summable generator for the $t^2$ coefficient via (6.3) and pair locality (D3 INTEGRATION:84). *Next layer:* R1 needs a selected fine vacuum and gap (76.6) or $\mathsf{R0}^u$ locally (CA-90, available at L1; D3-3 applicable, its witness hypotheses remain).

**D4-C** (full, diagonal; L4; → R2 by 79.2 [cond.] / R4). *Restated, relative (L4, full, R4):* given a modular $\mathcal{C}$ and a supplied effective completion $\mathcal{S}$, a pointed categorical carrier with bounded support growth carries tile Hamiltonians whose standard KS modes satisfy P and whose limit is the diagonal theory of $\mathcal{S}$, with the KS part separated from fidelity (D4 INTEGRATION:98–101). This is an inverse-discretisation conjecture strengthening Zini–Wang Conj. 4.3 and Rmk. 9, which are chiral ($\mathsf{S}_{\mathrm{ch}}$, L4: a UMM VOA; approximants of $L_m$) (ZW:871–895). *Attackable core (L0, any, → R2):* the shell certificate P0–P3 of Theorem 79.2. *First lemma:* P0–P3 for the sourced quasi-free calibration (INTEGRATION:103–111). *Next layer:* complete sector correspondence and braided equivalence with $\mathcal{C}$ (INTEGRATION (e)12).

**D5-C** (full; L2 + selected E1–E7; none / R4 uniformly). *Restated (L2; setting of whichever chiral subnet is declared; R4 given R3):* two-sided fidelity, a fully faithful endpoint functor plus an index bound $\leq \dim \mathsf{B}$ imply complete braided fidelity (Theorem 80.9); its finite-dimensional gap ($n$ vs $n^2$) is closed along route (ii) by Lemma 93.2, and the data layer lowers to L0+L4 / L1+L4 (Theorem 93.8). The universal-procedure clause is dropped (it contains Bischoff:364–373 and is trivialised by proof search, D5 REVIEW:210, :251); for $\mathcal{C} = \mathrm{Vec}$ a nontrivial target needs $c > 0$, an L3 datum (REVIEW:182–185). *First lemma* (the earlier compatible-CP lemma is superseded by Lemma 93.2): the annular approximant lemma of CA-93, then EP/AN (Lemmas 93.11–93.12). *Next layer:* $T$ and characters at L3; (F1) is R3 and is assumed, not supplied.

**D6-C** (chiral; L2 input, selects L3; none / R5). *Restated* as three parts: (a) *enrichment synthesis* L2 → L3: select E with a realising completely rational net; open and containing the realisation problem (Bischoff:364–373); (b) the relative D4-C above (L4 → R4); (c) *coherent selection* (R5) along

enrichment-preserving braided equivalences, with the automorphism-lifting hypothesis explicit (D6 REVIEW:121–125). The chiral clause is replaced by the declared-setting tag: in $\mathsf{S}_{\mathrm{full}}$ the target is a full net whose left chiral subnet realises $\mathcal{C}$ (REVIEW:94–98). *First lemma (*$\mathsf{L}1$*, open,* $\to \mathsf{R}0^u, \mathsf{R}1$*):* the braided vacancy tower with lifts $\mathrm{Ad}_{U_n} \circ B_n(\cdot \otimes \mathrm{id})B_n^\dagger$ is a strict local OAR tower with drift $d_{k,n} \le p_n e_n + 2(1-p_n)$ (D6 INTEGRATION:110–115; Proposal 76.5 + 76.1), checked first on a non-pointed braided instance with sourced $F, R$; proved for $U_n = 1$ (CA-90 90.1, 90.W), local dressing only for supplied circuits (90.D). *Next layer:* (a) itself.

### 89.5 Round-2 Verdicts (2026-10-22)

Grades are the Opus reviewers' (REVIEW.md in `explorations/2026-10-22-astra-r2/`), carried unchanged; tags as verified there.

- E1 → CA-90: **PROOF-COMPLETE** (braided headline; non-braided converse weaker, sharpened by R-1; E1 REVIEW:12–19); (any; $\mathsf{L}0$+relative half-braiding; $\mathsf{R}0^u$). Next: dyadic transport-extraction, Target 90.T.
- E5 → CA-93: **PROOF-COMPLETE** (E5 REVIEW:10); Lemma 93.2 (any; category-free; $\to \mathsf{R}4$), Theorem 93.8 (ch; $\mathsf{L}0/\mathsf{L}1$+$\mathsf{L}4$; $\mathsf{R}4$ given $\mathsf{R}3$). Next: the annular approximant lemma (no instance known).
- E6 → CA-94: **COUNTEREXAMPLE** (verified) to E6 as posed, with a proof-complete positive Theorem 94.6 (E6 REVIEW:9–12); (any; $\mathsf{L}0$+decl.; $\mathsf{R}0^c$+$z$=2). Next: contact trichotomy, Target 94.16.
- E2 → CA-91: **WEAKER-RESULT-PROVED** (E2 REVIEW:11–16); (full; category-free; $\mathsf{R}2$ [cond.]). Next: zero-mode (current–momentum) lemma.
- E3 → CA-92: **WEAKER-RESULT-PROVED** (E3 REVIEW:10); (full, ch only with twisted reflection; $\mathsf{L}0$+decl. $s$; finite, $\to \mathsf{R}3$). Next: reflection-coherence lemma.
- E4 → CA-89: **WEAKER-RESULT-PROVED** (E4 REVIEW:8); (full per family; category-free; $\to \mathsf{R}2$). Next: probe-word Ward-to-residual Lemma 89.L.

No cell moved up a rung except $\mathsf{R}0^u$ (reached at $\mathsf{L}0$+half-braiding, so at $\mathsf{L}1$); every other round-2 result is an ingredient ($\to$), conditional, finite, off-ladder or negative. *Round-3 verdicts (2026-10-25; Astra reviewers, REVIEW.md in* `explorations/2026-10-25-astra-r3/`*, carried unchanged; worklog 022).* F1 → CA-96 **COUNTEREXAMPLE** to the brief (blocked transport proof-complete; Target 90.T superseded); F5 → CA-99 **COUNTEREXAMPLE** (exact-family $N \le 2$ classification proof-complete; Target 94.16 refuted and replaced); F2 → CA-100, F3 → CA-98, F4 → CA-95, F6 → CA-97 **WEAKER-RESULT-PROVED**. No cell moved up a rung; $\mathsf{R}0^u$ is now reached at bare $\mathsf{L}0$ for the group-graded families of CA-96 (single-site centrality is neither necessary nor sufficient).

### 89.6 How to Read a Statement under the Layering Principle

Every statement carries three tags: ($\mathsf{S}$; $\mathsf{L}$; $\mathsf{R}$), with $\mathsf{S} \in \{\mathrm{ch}, \mathrm{full}, \mathrm{any}\}$, $\mathsf{L} \in \{\mathrm{none}, \mathsf{L}0, \ldots, \mathsf{L}4\}$ and $\mathsf{R}$ the rung it *reaches*; a conjecture also states the rung it *claims*. Microscopic declarations ($O$, $h$, boundary, refinements, states) are listed beside the tags. A statement that silently fixes a layer (one chirality, one enrichment, one target) is incomplete until it names the layer and the data (AGENTS.md:113–115). Two working rules follow: state the result at the lowest $\mathsf{L}$ and broadest $\mathsf{S}$ at which its proof runs, and record the extra data the next rung needs as part of the statement, as the restatements above do.

## 89.7 What This Does Not Claim

No rung is reached here. The grid does not decide chirality: $\mathsf{S}_{\mathrm{ch}}$ versus $\mathsf{S}_{\mathrm{full}}$ is a declared tag pending the source gap above. The full-setting fidelity notion (defect data plus $\mathcal{C} \boxtimes \overline{\mathcal{C}} \simeq Z(\mathcal{C})$ bulk identification) is a definition, not a theorem about any lattice limit. The minimal-data column of Definition 88.3 records current evidence; $\mathsf{R0}^u$ at bare $\mathsf{L0}$ for general $\mathcal{C}$ is open (Target 96.10; 90.T superseded).

## 89.8 Examples (Instances Only)

| Instance | Known placements on the grid |
|---|---|
| Fibonacci | $\mathsf{L2}$ available; tower CA-71–CA-74 at (open; $\mathsf{L0}$; $\mathsf{R0}^c$); CA-87 finite; LC-12 §5 obstruction at $\mathsf{L3}$ ($c = 14/5$ lift). |
| Ising | CA-82 at (ch; $\mathsf{L4}$; $\mathsf{R4}$ by cited theorems); the only instance at $\mathsf{R4}$, and it reads no $F, R$. |
| Semion | $\mathsf{L2}$ (pointed modular); LC-11/LC-13 conditional four-port at $\mathsf{L3}$ (declared $R = i$); no rung reached. |
| Haagerup | $\mathsf{L0}$ only (not braided); $\mathsf{S}_{\mathrm{full}}$ via $Z(\mathcal{C})$ is its natural modular target (Bischoff:384–390, open); no rung. |
| Vec | At $\mathsf{L0}$ every qudit chain is in scope (D2 REVIEW:79); at $\mathsf{R4}$ nontrivial only with $c > 0$ at $\mathsf{L3}$. |
| $\mathrm{Vec}_{S_3}$ | $\mathsf{L0}$, not braided: one-gap unital lifts obstructed (90.2(b)); no seam pairing for $O = \mathbf{1} \oplus (12) \oplus (123)$ (92.2). |

# 90 Ward-to-Koo–Saleur Transfer: Graph-Norm Interfaces, Derivative Matching and the Contact Cocycle

## 90.1 Status, Sources, and Dependencies

**Status: mixed, labelled per statement; review grade WEAKER-RESULT-PROVED.** Theorem 89.1(i)–(iii) is *conditional* (proved under its hypotheses); Lemma 89.2 and Proposition 89.8 are *derived*; Lemma 89.5 is *conditional* but carries almost no transfer content (REVIEW F1); Propositions 89.3, 89.4, 89.6, 89.7 are *negative, derived*; 89.3 and the strong form of 89.7 are the reviewer's. The round-2 conjecture E4 is *not proved*; its "any refinement" A3 clause is *false* whenever $c^* \neq 0$ (89.7). Lemma 89.L is a *proposal*. **Layer tags (aq):** 89.1–89.7 are *full, one chiral family at a time; category-free;* $\to$ R2 (89.4, 89.6 setting *any*; 89.8 setting *both*). No categorical datum enters any proof: they apply verbatim to every unitary fusion $\mathcal{C}$ with a declared periodic realisation. The periodic translation $U_r$ used in 89.1(iii) and 89.7 is *declared extra data*: for non-braided $\mathcal{C}$ with flux sectors it is a half-braiding ($Z(\mathcal{C})$) datum, not a fusion consequence (REVIEW:206). Nothing reaches R2: no instance of any hypothesis set is verified. Reproducibility: exact SymPy checks (`checks/run_checks.py`, five groups, five mutations rejected) and reviewer checks R1–R4 (`checks/review/review_checks.py`, all pass) in the exploration directory; they verify displayed finite identities and calibrations, not W1/W2 or any certificate.

Dependencies: CA-78 (Conjecture 78.12, Lemma 78.11, Target 78.13), CA-79 ((1), Lemma 79.1, Theorem 79.2, Lemma 79.6), CA-81 (VR-1), CA-88 (grid, D2-C); LC-W; CONVENTIONS (s), (v), (ad), (ak), (ap), (aq).

## 90.2 Setting and the Conjecture as Posed

A tower $(H_r, J_r, K_r)$, $N = N_r \to \infty$, satisfies VR-1 A1: finite Hilbert spaces, isometries $J_r$, $K_r \geq 1$, $K_{r+1}J_r = J_rK_r$. The modes $T_{r,n}$ are the standard KS modes (1) of (ap), $T_{r,n} = \frac{N}{4\pi}\sum_j z^{nj}(k_j - \varepsilon_r - \mathcal{J}_j) + b_r\delta_{n0}$, $k_j = -h_{r,j}/v_r$, $\mathcal{J}_j = i[k_j, k_{j+1}]$, with no change of prescription. Graph norms (v): $|X|_{s,d}^{t\leftarrow r} := \|K_t^s X K_r^{-s-d}\|$ (scale superscripts omitted within one scale). VR-1 A2: $|T_{r,n}|_{s,1} \leq C_{n,s}$; A3: $e_{r,n,s} := |T_{r+1,n}J_r - J_rT_{r,n}|_{s,1}$ summable in $r$; A4: $|[T_m, T_n] - (m-n)T_{m+n} - z_{mn}1|_{s,2} \to 0$, $z_{mn} = \delta_{m+n,0}(m^3c^* - mc)/12$, constants kept separate as in (s). *Geometric modes* $S_{r,m} = N^{-1}\sum_j f_m(j)Q_{r,j} + b_r\delta_{m0}$, $f_m(j) = e^{2\pi imj/N}$, are built from boundary metric insertions $Q_{r,j}$ on the positive quotient; $D_{r,n} := T_{r,n} - S_{r,n}$, $d_{r,n,s} := |D_{r,n}|_{s,1}$. A *periodic translation* is a unitary $U_r$, $U_r^N = 1$, commuting with $K_r$ and the time gates, with $U_rT_{r,n}U_r^* = e^{-2\pi in/N}T_{r,n}$ (declared; see Status).

**Conjecture E4 [conjectural as posed; full, L0, R2; brief:66–77].** If the gate metric derivative equals $h$ minus a continuity current with residual summable in the graph norm of $K_N = 1 + (H_N - E_0)/v$ (C1), and discrete Ward equations hold with convergent contact coefficients (C3), then (a) the KS modes satisfy A4 with $z_{mn}$ fixed by the contacts, (b) they satisfy A3 for *any* refinement that intertwines the gates up to a summable error, and (c) the central charge is the Ward contact coefficient. *Verdict (REVIEW:226–240).* (a) is not proved: its whole content sits in W2 (Lemma 89.5, Remark F1). (b) is **false** whenever $c^* \neq 0$ (Proposition 89.7), and real gate errors cannot give A3 (89.6). (c) holds only for $c^* = 12\kappa_3$; $c$ also records the declared zero-mode shift $b$ (89.1(iii)). Its literal control $K_N$ violates A2 (89.3).

**Theorem 89.1 [conditional; full, one chiral family at a time; category-free;** $\to$ **R2; INTEGRATION:9–34].** Let $(H_r, J_r, K_r)$ satisfy A1, with $\min\operatorname{spec} K_r = 1$, and a periodic translation. (i) *Transfer.* If $S_{r,n}$ satisfy A2 ($|S_{r,n}|_{s,1} \leq M_{n,s}$) and A4, and $d_{r,n,s} \to 0$, then $T$ satisfies A2 and A4 with the same central term, with bound (2). (ii) *Refinement.* If holomorphic metric

families $B_{r,n}(\eta)$ with $B'_{r,n}(0) = S_{r,n}$ obey (3), $J_r$ and $K_r$ are independent of $\eta$, and $d$ is summable, then $T$ satisfies A3 with bound (4). (iii) *Contact cocycle.* If $T$ satisfies A2 and A4-type residuals with convergent scalars $z^{(r)}_{mn} \to z_{mn}$ vanish in $|\cdot|_{s,2}$, then $z_{mn} = \delta_{m+n,0}(m^3c^* - mc)/12$ with $c^* = 2(a_2 - 2a_1)$, $c = 2a_2 - 16a_1$, $a_m := z_{m,-m}$. If the scalar contacts are first- and third-derivative Ward contacts $\kappa_1, \kappa_3$ (cylinder chart) with zero-mode shift $b$, then $c^* = 12\kappa_3$, $c = 24b - 12\kappa_1$.

### 90.3 Graph-Norm Calculus and the Choice of Control

**Lemma 89.2 [derived; any; category-free].** $|XY|_{s,d+e} \le |X|_{s,d}|Y|_{s+d,e}$ and $|X|_{s,d+1} \le |X|_{s,d}$. **Proof.** Insert $K^{-s-d}K^{s+d}$ between $X$ and $Y$; $\|K^{-1}\| \le 1$. □ (NOTE:50–69.)

**Proposition 89.3 (R-A2) [negative, derived; reviewer; full; category-free; → R2].** **Assume** the KS modes with $\varepsilon_N = E_0/(Nv_N)$, $K_N = 1 + (H_N - E_0)/v_N$, $b_N$ bounded, and spectral width $w_N := \max\operatorname{spec} K_N - 1 \ge w > 0$ for large $N$. **Prove** A2 fails at $(n, s) = (0, 0)$. ⟨1⟩1. $T_0 + \bar{T}_0 = N(K - 1)/(2\pi) + 2b$ (Lemma 79.1). ⟨1⟩2. On the top eigenvector, $\max(|T_0|_{0,1}, |\bar{T}_0|_{0,1}) \ge \frac{1}{2}|Nw_N/(2\pi) + 2b|/(1 + w_N) \to \infty$. QED (REVIEW:162–172). So E4's literal control is not admissible; every A2-dependent statement here needs another $K$, e.g. the conformal $1 + E_N$ or CA-79's $Q^p$, $p > 1$ (Theorem 79.2). This sharpens NOTE:71–82, which called A2 for this $K$ only "very strong".

### 90.4 Transfer of A2 and A4: Theorem 89.1(i)

**Proof of (i) (E4-T; NOTE:208–243, REVIEW:100–106).** ⟨1⟩1. A2 for $T$ with $C_{n,s} = M_{n,s} + \sup_r d_{r,n,s}$, BY the triangle inequality. ⟨1⟩2. With $R^S, R^T$ the A4 residuals, exactly $R^T_{mn} = R^S_{mn} + [D_m, S_n] + [S_m, D_n] + [D_m, D_n] - (m - n)D_{m+n}$. ⟨1⟩3. Lemma 89.2 on each product, with consecutive graph orders:

$$\begin{aligned} q^T_{r,m,n,s} \le q^S_{r,m,n,s} &+ d_{r,m,s}M_{n,s+1} + M_{n,s}d_{r,m,s+1} + M_{m,s}d_{r,n,s+1} + d_{r,n,s}M_{m,s+1} \\ &+ d_{r,m,s}d_{r,n,s+1} + d_{r,n,s}d_{r,m,s+1} + |m - n|\, d_{r,m+n,s} \longrightarrow 0. \end{aligned} \tag{2}$$

⟨1⟩4. QED; the barred family is identical. The two-chiral conclusion additionally needs zero mixed contact with vanishing two-loss error; the intended supplier is CA-91 Theorem 91.5, which needs *equal ledgers* (e.g. from a linear parity, Theorem 91.1) and the zero-mode condition ZC (E2 INTEGRATION:107–112; this corrects INTEGRATION:94–99).

**Proposition 89.4 (E4-N1) [negative, derived; any; category-free].** Summable insertion error does not bound commutators without a product bound: on $\mathbb{C}^2$, $K = 1$, $S_r = 2^r \operatorname{diag}(1, 0)$, $D_r = 2^{-r}\sigma_x$ give $\sum_r |D_r|_{s,1} < \infty$ but $|[S_r, D_r]|_{s,2} = 1$ (NOTE:245–258). This is why A2 enters (i); special cancellation may replace it only as a proved product bound.

**Lemma 89.5 (E4-W, demoted) [conditional; full; category-free; → R2; NOTE:84–204].** **Assume** W0: $N \to \infty$, $S_{r,m}$ as above, $D_N F(j) = (F(j+1) - F(j-1))/(2i\ell_N)$, $\ell_N = 2\pi/N$, $d_N(m) = \sin(m\ell_N)/\ell_N$. W1: on all boundary vectors of the positive quotient, $[S_{r,m}, Q_{r,j}] = f_m D_N Q_{r,j} + 2(D_N f_m)Q_{r,j} + [\kappa_{3,r}D^3_N f_m + \kappa_{1,r}D_N f_m](j)1 + X_{r,m,j} + W_{r,m,j}$ (exchange error $X$, Ward error $W$). W2: $E_{r,m,n} := N^{-1}\sum_j f_n(j)(X + W)_{r,m,j}$ has $|E_{r,m,n}|_{s,2} \le w_{r,m,n,s} \to 0$; $\kappa_{i,r} \to \kappa_i$, $b_r \to b$. W3: A2 for $S$. **Prove** $S$ satisfies A4 with $z_{mn} = \delta_{m+n,0}\{\kappa_3 m^3 + (\kappa_1 - 2b)m\}$. ⟨1⟩1. Periodic summation by parts ($D_N f_m = d_N(m) f_m$) gives, for $N > |m + n|$, $[S_m, S_n] = \beta_N S_{m+n} + z^{(r)}_{mn}1 + E_{r,m,n}$, $\beta_N = 2d_N(m) - d_N(m+n)$, $z^{(r)}_{mn} = \delta_{m+n,0}[\kappa_{3,r}d_N(m)^3 + (\kappa_{1,r} - 2b_r)d_N(m)]$. ⟨1⟩2. $|\sin x - x| \le |x|^3/6$ gives $|\beta_N - (m - n)| \le \ell_N^2(2|m|^3 + |m + n|^3)/6 =: a_{N,m,n}$. ⟨1⟩3. $q^S_{r,m,n,s} \le w_{r,m,n,s} + a_{N,m,n}M_{m+n,s} + |z^{(r)}_{mn} - z_{mn}| \to 0$, BY W2, W3, $\|K^{-2}\| \le 1$. QED. At finite $N$, $\beta_N$ is not antisymmetric ($\beta_N(m, m) \ne 0$); $X + W$ absorbs it.

**Remark F1 [derived; reviewer; REVIEW:15–22, :86–98].** W1 does not constrain $X, W$: they are defined by it. Hence $E_{r,m,n} \equiv [S_m, S_n] - \beta_N S_{m+n} - z^{(r)}_{mn}1$ identically (check R1, deviation $4 \cdot 10^{-15}$), and W2 *is* A4 with $\beta_N$ in place of $m - n$. Lemma 89.5 is the elementary implication "$\beta_N$-A4 $\Rightarrow$ A4"; the contour-to-commutator problem sits entirely inside W2. The local sufficient condition $N^{-1}\sum_j(|X_j|_{s,2} + |W_j|_{s,2}) \to 0$ would require each local commutator to carry the scalar $\kappa_3 d_N^3 f_m(j)$, while the KS central term is a state-level effect of the assembled sum: this local route is **unverified and plausibly never satisfied**, and is not advertised.

## 90.5 A3 from Differentiated Matching: Theorem 89.1(ii)

Hypothesis of (ii): for each $r, n$, $B_{r,n}$, $B_{r+1,n}$ are holomorphic on a neighbourhood of the same disk $|\eta| \le \rho_{r,n}$ (Fourier factors, cell counts, velocity, anomaly line and zero-mode origin included), and

$$F_{r,n}(\eta) := B_{r+1,n}(\eta)J_r - J_r B_{r,n}(\eta), \qquad \sup_{|\eta|=\rho_{r,n}} |F_{r,n}(\eta)|_{s,1} \le u_{r,n,s}, \qquad \sum_r u_{r,n,s}/\rho_{r,n} < \infty. \quad (3)$$

**Proof of (ii) (E4-A; NOTE:262–306, REVIEW:112–119).** $\langle 1\rangle 1$. $F'(0) = (2\pi i)^{-1}\oint F(\eta)\eta^{-2}d\eta$ for the finite matrix $F$; the fixed weights pass through the integral, so $|S_{r+1,n}J_r - J_r S_{r,n}|_{s,1} \le u_{r,n,s}/\rho_{r,n}$. $\langle 1\rangle 2$. Exact $K$ transport and isometry (functional calculus on the reducing range of $J_r$) give $|D_{r+1,n}J_r|_{s,1} \le d_{r+1,n,s}$, $|J_r D_{r,n}|_{s,1} \le d_{r,n,s}$. $\langle 1\rangle 3$. Adding,

$$e_{r,n,s} \le u_{r,n,s}/\rho_{r,n} + d_{r+1,n,s} + d_{r,n,s}, \quad (4)$$

summable. QED. A2 is not used. A directly certified summable bound on $F'(0)$, or D2-L10 collars for this derivative (Lemma 78.11), may replace (3). If $J$ depends on $\eta$, the connection terms $B_{r+1,n}(0)J_r'(0) - J_r'(0)B_{r,n}(0)$ must be kept; CA-79 Lemma 79.6 (compression/leakage) is their concrete source. A1 plus (i) and (ii) let VR-1 apply unchanged.

**Proposition 89.6 (E4-N2) [negative, derived; any; category-free].** Uniformly close, positive, entire real gate families need not have close metric derivatives: for an orthogonal projection $A \ne 0$, $\epsilon_r = 2^{-r-2}$, $G_r(t,\eta) = \exp[-t(1+\epsilon_r \sin(\eta/\epsilon_r^2))A]$, $G = e^{-tA}$, one has $\sup_{0\le t\le 1,\eta\in\mathbb{R}} \|G_r - G\| \le \epsilon_r$ but $\|\partial_\eta(G_r - G)(1,0)\| = e^{-1}/\epsilon_r$ (NOTE:308–334). The complex growth $\sim e^{\rho/\epsilon^2}$ breaks (3): the Cauchy bound is a genuine added hypothesis. It is an operator family, not a C1 certificate.

**Proposition 89.7 (E4-N3 strengthened to R-N3′) [negative, conditional reductio; reviewer; full; category-free; → R2; NOTE:336–365, REVIEW:174–197].** **Assume** A1, $N_r \to \infty$, a periodic translation, $\min\operatorname{spec} K_r = 1$, and that $T$ satisfies A3 for $J$ and A4 with $(c^*, c)$. **Prove** either $c^* = 0$, or A3 fails for $J_r' := U_{r+1}^{k_r} J_r$, $k_r$ even, $|k_r - N_{r+1}/2| \le 1$; $J'$ has the same gate-intertwining errors as $J$. $\langle 1\rangle 1$. $\Theta := U_{r+1}^{k_r}$ gives $\Theta T_n = \lambda_r T_n \Theta$, $\lambda_r \to -1$ for odd $n$; with $\Delta, \Delta'$ the defects of $J, J'$, $\Delta' = \Theta(\lambda^{-1}\Delta + (\lambda^{-1} - 1)JT_{r,n})$, and $\Theta J$ is a graph isometry, so $|\Delta'|_{s,1} \ge |\lambda^{-1} - 1|\,|T_{r,n}|_{s,1} - |\Delta|_{s,1}$. $\langle 1\rangle 2$. Case A: $\limsup_r |T_{r,n}|_{s,1} > 0$ for some odd $n$, some $s$: A3 fails for $J'$. $\langle 1\rangle 3$. Case B: $|T_{r,n}|_{s,1} \to 0$ for all odd $n$, all $s$. Then $|T_n T_{-n}|_{s,2} \le |T_n|_{s,1}|T_{-n}|_{s+1,1} \to 0$, so A4 gives $T_0 + z_{n,-n}/(2n) \to 0$ in $|\cdot|_{s,2}$ for $n = 1, 3$; the difference is a scalar and $\|K^{-2}\| = 1$, so $z_{1,-1}/2 = z_{3,-3}/6$, i.e. $c^* = 0$ (check R4). QED. A2 is not used. Hence E4(b) holds only vacuously or with $c^* = 0$; A3 needs *coordinate-resolved* differentiated matching as in (ii). An even $k_r$ preserves a brick-wall sweep (REVIEW:132–135).

## 90.6 The Contact Cocycle: Theorem 89.1(iii)

**Proof of (iii) (E4-C; NOTE:412–493, REVIEW:144–158).** Let $R_{r,m,n}$ be the residuals with the limiting scalars, norms $q_{r,m,n,s} \to 0$. $\langle 1\rangle 1$. $z$ is antisymmetric, diagonal and real. $\langle 2\rangle 1$. Add

$(m,n)$, $(n,m)$: operator parts cancel, $\|K^{-2}\| = 1$. $\langle 2\rangle 2$. For $m+n \neq 0$, $N > |m+n|$, averaging over $U_r^j$ kills the operator part by its Fourier character, is contractive ($[U_r, K_r] = 0$), and leaves the scalar. $\langle 2\rangle 3$. For $(m,-m)$ the operator part is self-adjoint; test on a $K$-eigenvector of eigenvalue 1. $\langle 1\rangle 2$. Finite Jacobi. By A2 and Lemma 89.2, $|[T_l, R_{mn}]|_{s,3} \leq C_{l,s} q_{r,m,n,s+1} + q_{r,m,n,s} C_{l,s+2} \to 0$. The $T_{l+m+n}$ coefficients cancel algebraically; the scalar $(m-n)z_{l,m+n} + (n-l)z_{m,n+l} + (l-m)z_{n,l+m}$ has vanishing $|\cdot|_{s,3}$ norm, and $\|K^{-3}\| = 1$ forces it to vanish. $\langle 1\rangle 3$. The triple $(m, 1, -m-1)$ gives $(m-1)a_{m+1} = (m+2)a_m - (2m+1)a_1$ $(m \geq 2)$, solved by $a_m = (a_2 - 2a_1)(m^3 - m)/6 + a_1 m = (m^3 c^* - mc)/12$; oddness covers $m \leq 0$. This is the diagonal recurrence of Schottenloher (2507–2634) with the linear coboundary kept. $\langle 1\rangle 4$. The $\kappa$ dictionary: with Lemma 89.5's contacts $a_m = \kappa_3 m^3 + (\kappa_1 - 2b)m$, so $c^* = 12\kappa_3$, $c = 24b - 12\kappa_1$. QED. The anchor $\min \operatorname{spec} K = 1$ is essential: without it a scalar identity can vanish in weighted norm. Only $c^*$ is a Ward output; $c = c^*$ needs $b = (c^* + 12\kappa_1)/24$, an input or a further equation (REVIEW:210), consistent with Lemma 79.1 ($c \mapsto c + 24d$, $c^*$ fixed).

**Proposition 89.8 (E4-P) [derived, formal; both; category-free; → R2; NOTE:369–410].** In the plane convention (ad), if the scalar part of the ordered contact is $\sum_{k\geq 1} a_k(w)(z-w)^{-k}1$ (finitely many terms), then with $f_m = z^{m+1}$, $g_n = w^{n+1}$, $z_{mn} = \operatorname{Res}_w g_n(w) \sum_{k\geq 1} \frac{a_k(w)}{(k-1)!} \partial_w^{k-1} f_m(w)$. For D2 (19) ($a_4 = c_W/2$): $z_{mn} = \delta_{m+n,0} c_W (m+1)m(m-1)/12$; the nonscalar residue is $(m-n)L_{m+n}$, and pole coefficients $A_2, A_1$ would give $(A_2 - A_1)(m+1) - A_1(n+1)$, so the Ward values $A_2 = 2, A_1 = 1$ are essential; a shift $T_n = L_n + b\delta_{n0}$ gives $c^* = c_W$, $c = c_W + 24b$. Plane-to-cylinder formally gives $\kappa_1 = 0$, $\kappa_3 = c_W/12$, consistent with 89.1(iii) (REVIEW:137–142). Formal residue calculus only: the lattice chart-change estimate is open.

## 90.7 The First Gap and the Next Lemma

**First gap [open].** Turn the correlation-level contour identity D2 (19) into the operator two-loss estimate W2. By Remark F1 this is A4 up to $\beta_N$: the conjecture's entire A4 content sits here. LC-W proves only the propagation law, not a bound on its accumulated sum (`reconstruction/01_cells.md`:164–195). Further open lemmas (NOTE:522–547): uniform Ward errors after Gram conditioning (D2-L9); A2 for an admissible control; the differentiated matching (3) with $K$ transport; chart change and both chiralities.

**Lemma 89.L (probe-word Ward-to-residual) [proposal; full; category-free; → R2; INTEGRATION:120–136]. Assume** D2's contour identity (19) holds with residuals $R_n(f, y, A)$ for every probe word $A$ in a family whose reflected boundary vectors are dense in each graph space, and the residuals and collar insertions obey the one-sided two-loss bound of D2-L10 (Lemma 78.11) with constants tending to zero. **Prove** W2: $|[S_m, S_n] - \beta_N S_{m+n} - z_{mn}^{(r)}|_{s,2} \to 0$. All probe words, not the vacuum alone, are needed: vacuum-null is not operator-null (VR-2, virasoro_routes:179–192). *First test:* the in-repo free-fermion benchmark (`cft_machine/benchmarks/fermion/`; D2 INTEGRATION:117–121) with $K = 1 + E_N$, at the laptop budget.

## 90.8 What This Does Not Claim

No instance of any hypothesis set of Theorem 89.1 is constructed or verified for any $\mathcal{C}$; nothing reaches R2. No discrete Ward identity is shown to imply A4 (that is W2). No $c = c^*$, no positive energy, smooth smearing, mode-number uniformity, self-adjointness, integration or net (VR-1 exclusions, virasoro_routes:151–177). The Carpi–Weiner continuum bound is not lattice A2. Proposition 89.8 is formal, not contour convergence. No categorical selection: the statements are category-free, which is their scope, not a fidelity claim.

| Instance | Calibration datum for the general statements |
|---|---|
| Any unitary fusion $\mathcal{C}$, declared periodic realisation | 89.1, 89.5: no fusion identity enters; translation declared (half-braiding datum if non-braided). |
| Vec, $O = \mathbf{1}$ | All modes scalar or zero; 89.1(iii) gives $c = c^* = 0$; consistent (REVIEW:203). |
| Vec, $O = \mathbb{C}^2$, Heisenberg density | 89.3: $\max \lvert T_0 K^{-1} \rvert = 0.849, 2.072, 3.119$ at $L = 4, 6, 8$, linear growth (check R2). |
| Unit-multiplicity $2 \times 2$ block of any $\mathcal{C}$ | 89.4, 89.6 witnesses; not conformal certificates. |
| CGS $n = 1$ point | Stress convergence proved in source (StressTensor:527–532); **not verified** to satisfy W2 or (3). |

# 91 Local Unital Lifts from Relative Half-Braidings

## 91.1 Status, Sources, and Dependencies

**Status: mixed; every statement carries its own label.** Derived: Lemma 90.0, Theorem 90.1, Negative 90.N, Corollary 90.W, Proposition 90.V, Proposition 90.2 (part (b) is the reviewer's Proposition R-1), Proposition 90.3, Corollary 90.D (an implication from a *supplied* circuit). Proposal: Target 90.T. Open: arbitrary local dyadic lifts at L0, WT-1 A3 for any non-trivial state family, the periodic seam, parity covariance. Review grades (carried, not upgraded): braided headline **PROOF-COMPLETE**; non-braided converse **WEAKER-RESULT-PROVED**, sharpened by R-1 (E1 REVIEW:12–21). **Abbreviations:** "E1 NOTE/REVIEW/INTEGRATION" are `NOTE.md`, `REVIEW.md`, `INTEGRATION.md` in `explorations/2026-10-22-astra-r2/E1-local-unital-lift/`, whose `checks/` hold Python/NumPy Float64 calibrations (commands: `checks/README.md`:8–13, REVIEW:180–183) that calibrate and do not prove; a port to `scripts/` with a run bundle is open. No Julia or LaTeX run was made. Proofs are structured, not proof-assistant. This shard **closes CA-75 Link (b1)** and **supersedes CA-76 Proposal 76.5** for braided $\mathcal{C}$ and for L0 with a declared pointed relative half-braiding, and answers D3 open problem 1 at that layer. Dependencies: CA-68, CA-71, CA-75, CA-76, CA-88; WT-1; CONVENTIONS (r), (t), (x), (am), (an), (aq).

**Layer tags [(aq)].** Setting is any: no target theory enters; the microscopic declaration is an *open* chain with declared endpoints (REVIEW:114–119). "+H" means Definition 90.H's datum.

| Statement | S | L | R |
|---|---|---|---|
| Lemma 90.0, Theorem 90.1, Neg. 90.N | any | L0+H (in particular L1) | R0$^u$ |
| Corollary 90.W | any | L0+H, declared states | $\to$ R1 (conditional on summability) |
| Proposition 90.V | any | L0+H, constructed states | R1 (vacuum product only); negative |
| Proposition 90.2 (a); (b) | any | L0 | $\to$ R0$^u$; negative for one-gap lifts |
| Proposition 90.3, Corollary 90.D | any | L0+H (90.D: + a supplied circuit) | R0$^u$ implementation |
| Target 90.T | any | L0 | $\to$ R0$^u$ [proposal] |

## 91.2 Setting and the Transport Datum

$\mathcal{C}$ is unitary fusion and $(O,\eta)$ is pointed as in (am): $\eta^\dagger\eta = 1$, $p_0 = \eta\eta^\dagger$, $p_1 = 1 - p_0$, $O \cong \mathbf{1} \oplus X$; $X$ may contain unit summands and need not be simple, self-dual or multiplicity-free. $W_M = O^{\otimes M}$ (left-associated, (r)), $\mathcal{A}_M = \mathrm{End}_{\mathcal{C}}(W_M)$ acting on $\bigoplus_c \mathrm{Hom}(c, W_M)$, and $\mathcal{D} = \langle O\rangle$ is the *full* fusion subcategory generated by $O$, so every $W_M$ and every endomorphism of it lies in $\mathcal{D}$. **Support** means membership: $a \in \mathcal{A}_{[j,k]} := 1_{W_{j-1}} \otimes \mathrm{End}(W_{k-j+1}) \otimes 1_{W_{M-k}}$ (single-site algebras need not generate $\mathcal{A}_M$). **Degree**: with $N_M = \sum_j p_1^{(j)}$, $a$ has degree $q$ iff $[N_M, a] = qa$; this is the (am) grading since the $Q_{M,n}$ are the spectral projections of $N_M$ (REVIEW:49–53). $V_L$ is the CA-68 dyadic insertion (placement $\varphi(j) = 2j-1$, (t)), $\theta_L = \mathrm{Ad}\, V_L$ its corner map.

**Definition 90.H [definition; L0 plus a declared datum; NOTE:19–26].** A *pointed unitary relative half-braiding* of $(O,\eta)$ on $\mathcal{D}$ is a family of unitaries $t_Y : Y \otimes O \to O \otimes Y$, $Y \in \mathcal{D}$, with (associators and unitors suppressed; Penneys:319, :395–398; FRS:806–819)

$$t_{\mathbf{1}} = 1_O, \quad t_Z(f\otimes 1_O) = (1_O \otimes f)t_Y \ (f : Y \to Z), \quad t_{Y\otimes Z} = (t_Y \otimes 1_Z)(1_Y \otimes t_Z), \quad t_Y(1_Y \otimes \eta) = \eta \otimes 1_Y. \tag{H}$$

$z_Y = t_Y^{-1}$ is a half-braiding of $O$ relative to $\mathcal{D}$ (1704.04221:298–307); unitarity and pointedness are imposed, not inferred from that source.

**Remark 90.R [derived; NOTE:16, :160, :283–284; REVIEW:49–53, :79–83].** (a) A unitary braiding $c$ on $\mathcal{C}$, or only on $\mathcal{D}$, gives $t_Y = c_{Y,O}$ satisfying (H) (FRS:921–924, :1023–1125); the $\eta$ identity uses $c_{Y,\mathbf{1}} = \mathrm{id}$, standard but not displayed at a cited local line (*Law 1 gap*, REVIEW:31). The opposite crossing gives a different valid tower; no mixture is made. (b) A lift of $O$ to $Z(\mathcal{C})$ with a central pointed inclusion restricts to (H) but is strictly stronger ($\mathrm{Vec}_G$ row of the table). (c) Centrality of the chosen $\eta$ is needed: a half-braiding on $O$ with non-central point breaks intertwining ($\mathrm{Vec}_{\mathbb{Z}/2}$ row). (d) Conversely a central $\eta$ splits $O = \mathbf{1} \oplus X$ in $Z(\mathcal{D})$, as $p_0, p_1$ are then central.

## 91.3 The One-Gap Lift

**Lemma 90.0 [derived; E1.1, NOTE:30–68; REVIEW:55–61, every step valid].** Assume (H), $0 \le s \le M$, $b_{M,s} = 1_{W_s} \otimes t_{W_{M-s}} : W_M \otimes O \to W_{M+1}$ (new site after old site $s$), $\beta_{M,s}(a) = b(a \otimes 1_O)b^\dagger$, and $v_{M,s}$ the $\eta$-insertion in that gap. Prove: $\beta$ is a unital injective $*$-homomorphism, $\beta(a)v = va$, and for $a \in \mathcal{A}_{[j,k]}$

$$\beta(a) \in \mathcal{A}_{[j,k]} \ (s \ge k), \qquad \mathcal{A}_{[j+1,k+1]} \ (s < j), \qquad \mathcal{A}_{[j,k+1]} \ (j \le s < k). \tag{90.1}$$

$\langle 1\rangle 1$. $b$ is unitary; tensor interchange and dagger (Penneys:319) give a unital $*$-map; compressing by $1 \otimes \eta$ returns $a$, so it is injective. $\langle 1\rangle 2$. The $\eta$ identity on the suffix gives $b(1 \otimes \eta) = v$, so $\beta(a)v = b(1 \otimes \eta)a = va$. $\langle 1\rangle 3$. $s \ge k$: $b$ acts only after site $k$ and on the ancilla; interchange. $\langle 1\rangle 4$. $s < j$: $a = 1_{W_s} \otimes a_{\mathrm{tail}}$ with $a_{\mathrm{tail}}$ an endomorphism of the whole suffix; naturality gives $t(a_{\mathrm{tail}} \otimes 1)t^\dagger = 1_O \otimes a_{\mathrm{tail}}$. No generation by one-site operators is used; this is where (H) on the *full* $\mathcal{D}$ is needed. $\langle 1\rangle 5$. $j \le s < k$: with $A = W_{j-1}$, $J = W_{s-j+1}$, $R = W_{k-s}$, $C' = W_{M-k}$, the tensor identity gives $b = (1_{AJ} \otimes t_R \otimes 1_{C'})(1_{AJR} \otimes t_{C'})$; the second factor commutes past $a$, leaving $1_A \otimes [(1_J \otimes t_R)(a_{JR} \otimes 1_O)(1_J \otimes t_R^\dagger)] \otimes 1_{C'} \in \mathcal{A}_{[j,k+1]}$. □ Only (H) is used: no self-duality, multiplicity-freeness, ribbon structure or TL quotient (REVIEW:61).

## 91.4 The Dyadic Lift

**Theorem 90.1 [derived; E1.2 (braided) and E1.5 (L0+H), NOTE:72–117, :151–158; REVIEW:63–70, :79–83; PROOF-COMPLETE].** Assume Definition 90.H. Let $\alpha_L = \beta_{M_L,s_L} \circ \cdots \circ \beta_{M_1,s_1}$ with $M_i = L + i - 1$, $s_i = 2i - 1$; equivalently, with $B_L : W_L \otimes W_L \to W_{2L}$ interleaving $\mathrm{old}_1..\mathrm{old}_L, \mathrm{new}_1..\mathrm{new}_L$ into $\mathrm{old}_1, \mathrm{new}_1, \ldots, \mathrm{old}_L, \mathrm{new}_L$ by a crossing $t_O$ of $\mathrm{old}_j$ with $\mathrm{new}_i$ exactly when $i < j$,

$$\alpha_L(a) = B_L(a \otimes 1_{W_L})B_L^\dagger. \tag{90.2}$$

Prove: (i) $\alpha_L$ is a unital injective $*$-homomorphism with $\alpha_L(a)V_L = V_L a$; (ii) $\alpha_L(\mathcal{A}_{[j,k]}) \subset \mathcal{A}_{[2j-1,2k-1]}$, uniformly in $L$, $n$ and $\mathcal{C}$, so $r = 0$ in the brief's convention, and $\alpha_L(f^{(j)}) = f^{(2j-1)}$ for $f \in \mathrm{End}(O)$; (iii) $[N_{2L}, \alpha_L(a)] = \alpha_L([N_L, a])$; (iv) for a unitary (braided, at L1) strong monoidal equivalence with unitary $u : F(O) \to O'$ carrying $(F(\eta), F(t))$ to $(\eta', t')$, $F_{2L}\alpha_L = \alpha'_L F_L$; (v) for $L_n = 2^n L_0$ the composites $\alpha_n^k = \alpha_{n-1} \cdots \alpha_k$ form a coherent directed system of unital injective $*$-maps (WT-1 A1).

$\langle 1\rangle 1$. Before step $i$, $\mathrm{old}_i$ sits at $2i - 1$ and the suffix is $\mathrm{old}_{i+1..L}$; so step $i$ inserts right after $\mathrm{old}_i$, and expanding suffix crossings by the tensor identity gives (90.2).

$\langle 1\rangle 2$. (i): Lemma 90.0 for each $\beta$; the one-gap isometries compose to $V_L$ (CA-71:102–113, (t)).

$\langle 1\rangle 3$. (ii): by (90.1), insertions $i < j$ shift both endpoints, $j \le i < k$ only the right one, $i \ge k$ neither; hence the length-$(2(k - j + 1) - 1)$ interval

$$[\, j + (j-1),\ k + (j-1) + (k-j)\,] = [2j - 1, 2k - 1]. \tag{90.3}$$

⟨1⟩4. (iii): naturality gives $t_O(p_e \otimes 1) = (1 \otimes p_e)t_O$; the $\eta$ identity and unitarity give $t_Y(1_Y \otimes p_0)t_Y^\dagger = p_0 \otimes 1_Y$ (the one place pointedness enters). Summing strand by strand,

$$N_{2L}B_L = B_L(N_L \otimes 1 + 1 \otimes N_L), \qquad [N_{2L}, \alpha_L(a)] = \alpha_L([N_L, a]). \tag{90.4}$$

⟨1⟩5. (iv): $F$ and the tensorators (Penneys:325–348) carry the finite composite $B_L$ to $B'_L$. *Law 1 gap:* braided and unitary monoidal functors are not defined at a cited local line (REVIEW:30). Compatibility with composition of equivalences is asserted (NOTE:113), not proved, and not used (REVIEW:68).

⟨1⟩6. (v): cocycle by associativity of composition; WT-1 A1 and the norm inductive limit (2010.11121:164–176). Path independence across placement forests is not asserted (A1 does not need it). □

*Consequences [derived].* For $h \in \mathcal{A}_{[j,j+1]}$, $\alpha_L(h) \in \mathcal{A}_{[2j-1,2j+1]}$ and $\alpha_L(h)P_L = \theta_L(h)$: a *homomorphic* local representative replacing the local representative of Correction 76.B (INTEGRATION:80–82). The lift transports constants with unit 1, the corner with unit $P_L$; these are different observables (NOTE:263). $\alpha_L$ is not reflection-equivariant (placement $2j-1$, one crossing type; the mirror is the opposite-crossing lift with even placement; INTEGRATION:69–71).

**Negative 90.N [derived, negative; NOTE:119; REVIEW:70].** By (90.3), $\alpha_L(N_L) = \sum_j p_1^{(2j-1)}$, so $N_{2L} - \alpha_L(N_L) = \sum_j p_1^{(2j)}$, of norm exactly $L$ when $X \neq 0$. Reading (iii) as a uniformly bounded *absolute-number* difference is false; what holds is the (am) endomorphism-degree statement (90.4).

## 91.5 Wilson Consequences

**Corollary 90.W [derived; E1.3, NOTE:123–134; REVIEW:72–73].** Assume Theorem 90.1, states $\omega_n$ on $\mathcal{A}_{L_n}$, and the (an) quantities with $W_n = V_n$. Prove $d_{k,n} \leq p_n e_n + 2(1-p_n)$ for $k \leq n$ ($p_n e_n := 0$ at $p_n = 0$). ⟨1⟩1. Theorem 90.1(i) is the intertwining hypothesis of Proposition 76.1(iii)–(iv) (CA-76:90–108). ⟨1⟩2. At $p_n = 0$, $\tau_n, e_n$ are undefined and the bound 2 is the trivial one; at $p_n = 1$ the bound is $e_n$, not 0. □ Summable bounds would give WT-1 A3 (`wilson_triangle.md`:18–107; computable tails for WT-2, :109–129); none is supplied here.

**Proposition 90.V [derived; E1.4, NOTE:136–147; REVIEW:75–78].** (a) $e_M = \eta^{\otimes M}$ and $\omega_M^{\mathrm{vac}}(a) = e_M^\dagger a e_M$ satisfy $e_{2M} = V_M e_M$, so $\omega_{2M}^{\mathrm{vac}}\alpha_M = \omega_M^{\mathrm{vac}}$: $p_n = 1$, $e_n = 0$, zero drift, and a WT-1 limit state exists. This R1 instance is the *trivial vacuum product state*; no criticality is claimed. (b) *Negative.* If $X \neq 0$, take a simple $x$, an isometry $u : x \to X$, and $f_M = (\iota u) \otimes \eta^{\otimes(M-1)}$; $\omega_M^{\mathrm{occ}}(a) = f_M^\dagger a f_M$ is also compatible. Alternating $\omega_n$ between the two families gives $p_n = 1$ at every step, yet on $2p_0^{(1)} - 1$ adjacent pullbacks alternate $\pm 1$: drift exactly 2, $e_n = 2$, no limit. So $p_n = 1$ alone does not give A3.

## 91.6 Non-Braided Input: Finite Equations and the One-Gap Obstruction

Theorem 90.1 already runs at L0 with Definition 90.H; neither a braiding of $\mathcal{C}$ nor a lift of $O$ to $Z(\mathcal{C})$ is needed (Remark 90.R). The question is when the datum exists, and whether anything else can work.

**Proposition 90.2 [derived; (a) E1.6, NOTE:162–189, REVIEW:84–86; (b) reviewer Proposition R-1, REVIEW:148–168].** (a) Let $s_i$ run over the simples of $\mathcal{D}$ (unit included) and $f_{ij;k,\mu} : s_k \to s_i \otimes s_j$ be complete orthonormal isometric embeddings, all multiplicities kept. Pointed unitary relative half-braidings are exactly the solutions of the finite system (associators on both sides)

$$z_{\mathbf{1}} = 1_O, \quad z_i(\eta \otimes 1_i) = 1_i \otimes \eta, \quad (f_{ij;k,\mu} \otimes 1_O)z_k = (1_i \otimes z_j)(z_i \otimes 1_j)(1_O \otimes f_{ij;k,\mu}), \quad z_i \text{ unitary.} \tag{90.5}$$

A necessary integer condition is

$$\Delta_{x,c} := \dim \mathrm{Hom}(c, O \otimes x) - \dim \mathrm{Hom}(c, x \otimes O) = 0 \quad \forall\, x, c \text{ simple}, \quad \text{i.e. } [O][x] = [x][O] \text{ in } K(\mathcal{D}). \tag{90.6}$$

$\langle 1\rangle 1$. Necessity: naturality, tensor identity, unitarity, centrality of $\eta$; a unitary is an isomorphism in each charge block, so $\Delta \neq 0$ excludes (90.5) (cf. ENO:1221). $\langle 1\rangle 2$. Sufficiency: $z_Y = \sum_{i,\mu}(v_{i,\mu} \otimes 1_O) z_i (1_O \otimes v_{i,\mu}^\dagger)$ is unitary and basis-independent, natural by Schur, tensorial by summing (90.5) against $f^\dagger$. □ (90.6) is only necessary; the associator-dependent equations (90.5) remain. (b) Assume $M \geq 2$, $1 \leq s \leq M-1$, $J = W_s$, $R = W_{M-s}$, and *any* unital $*$-homomorphism $\beta : \mathcal{A}_M \to \mathcal{A}_{M+1}$ with $\beta(a) v_{M,s} = v_{M,s} a$, $\mathcal{A}_{[j,k]} \to \mathcal{A}_{[j,k]}$ for $k \leq s$ and $\mathcal{A}_{[j,k]} \to \mathcal{A}_{[j+1,k+1]}$ for $j > s$ (straddling intervals unconstrained). Prove $\Delta_{x,c} = 0$ for every simple $c$ and every simple $x$ that is a summand of both $J$ and $R$. $\langle 1\rangle 1$. $\beta(a \otimes 1_R) = a \otimes 1_O \otimes 1_R$ and $\beta(1_J \otimes b) = 1_J \otimes 1_O \otimes b$, by the pattern, intertwining and $v^\dagger v = 1$. $\langle 1\rangle 2$. On $S = \mathrm{End}(J) \otimes \mathrm{End}(R)$, block $(i,j)$ sits in block $c$ of $\mathrm{End}(J \otimes R)$ with multiplicity $N_{ij}^c$ and in block $c'$ of $\mathrm{End}(J \otimes O \otimes R)$ with multiplicity $T_{ij}^{c'} = \dim \mathrm{Hom}(c', s_i \otimes O \otimes s_j)$. $\langle 1\rangle 3$. With $\Lambda$ the inclusion matrix of $\beta$, $\sum_c N_{ij}^c \Lambda_{cc'} = T_{ij}^{c'}$ (uniqueness of decomposition of finite-dimensional representations). $\langle 1\rangle 4$. $i = \mathbf{1}$ (as $\mathbf{1} \subset O \subset J$) gives $\Lambda_{xc'} = \dim \mathrm{Hom}(c', O \otimes x)$; $j = \mathbf{1}$ gives $\Lambda_{xc'} = \dim \mathrm{Hom}(c', x \otimes O)$. □ So (90.6) is necessary for *every* one-gap lift with the support pattern (90.1), not only for the half-braiding construction, once $x$ occurs in $W_s$ and $W_{M-s}$. **Not decided:** dyadic lifts with slack $r$ or another internal placement, where the left/right images are only lifts with unknown inclusion matrices (REVIEW:165–168).

**Boundary [open; NOTE:191, :269].** (i)–(iii) constrain algebra maps, not crossings; $z$ is not reconstructed from an arbitrary $\alpha$. Existence or a complete obstruction for arbitrary local dyadic unital lifts at $\mathsf{L0}$ is open.

## 91.7 Circuit Depth and Dressing

**Proposition 90.3 [derived; E1.7, NOTE:195–214; REVIEW:91–102].** Assume Theorem 90.1, $X \neq 0$, the fine $2L$-site chain and the contiguous input $a \mapsto a \otimes 1_{W_L}$. Prove: $B_L$ is a nearest-neighbour circuit of $t_O$ crossings of depth $L-1$ with $L(L-1)/2$ gates, and any range-$w$, depth-$D$ circuit $U$ with $U(a \otimes 1)U^\dagger = \alpha_L(a)$ obeys $D(w-1) \geq L-1$ (90.7) $\langle 1\rangle 1$. Layer $t = 1, \ldots, L-1$: crossings on $(p, p+1)$, $p = L - t + 2i - 1$, $i = 1, \ldots, t$, exchanging $\mathrm{old}_{L+i-t}$ with $\mathrm{new}_i$; disjoint and adjacent by induction on $t$. $\langle 1\rangle 2$. Each pair $i < j$ crosses once (at $t = L + i - j$) and the final word is interleaved. A shuffle of two increasing sequences is 321-avoiding, so all its reduced words are commutation-equivalent: no braid (Yang–Baxter) move, which a relative half-braiding need not supply among old strands, is used, and tensor interchange identifies the schedule with $B_L$ (the combinatorial lemma made explicit at REVIEW:92–97). $\langle 1\rangle 3$. One layer enlarges an interval support by at most $w-1$ at each end (interchange). $\langle 1\rangle 4$. $p_0^{(L)} \otimes 1$ has support $\{L\}$ and image $p_0^{(2L-1)}$ (Theorem 90.1(ii)). A $p_0$ outside an interval is not in that interval algebra: compressing the other sites to $\eta$, interval operators give equal scalars on $\eta$ and on an isometric simple summand of $X$ at the omitted site, $p_0$ gives 1 and 0. □ So no fixed depth and range implement the family in this geometry, while $\alpha_L$ itself has zero support growth. Distributed-ancilla geometries change the input convention and are not compared (NOTE:216).

**Corollary 90.D [derived implication; NOTE:218–224; REVIEW:104].** For a *supplied* output circuit $U$ of depth $D$ and range $w$, $W = UV_L$ and $\alpha^U = \mathrm{Ad}_U \circ \alpha_L$ satisfy $\alpha^U(a)W = Wa$, and an $m$-site input has output support length $\leq 2m - 1 + 2D(w-1)$: $r = 2D(w-1)$. Degree preservation needs $[U, N_{2L}] = 0$ in addition. $U$ is not constructed; a quasi-local $U$ needs its own tail estimate (D3's dressing lemma stays open).

## 91.8 Integration Status and the Next Lemma

CA-75 Link (b1) is closed at (any; L0 + H; R0$^u$); without the datum at L0 it is open. Proposal 76.5 is superseded by Theorem 90.1, including the owed non-pointed check with sourced $F, R$ (Fibonacci row). D3-3's local-witness route (Question 76.12 (L)) becomes applicable; its witness hypotheses remain. Corrections 76.A, 76.C and Negative 76.11(d) are untouched. Theorem 90.1 is open-chain only: periodic seam coherence of $B_L$ (needed by $\mathsf{S}_{\text{full}}$, LC-4) is open (NOTE:273).

**Target 90.T [proposal; dyadic transport-extraction; INTEGRATION:85–91].** Let $\alpha_L : \mathcal{A}_L \to \mathcal{A}_{2L}$ be unital $*$-maps with $\alpha_L(a)V_L = V_L a$ and $\alpha_L(\mathcal{A}_{[j,k]}) \subset \mathcal{A}_{[2j-1-r,\, 2k-1+r]}$ for a fixed $r$ and all $L$. Show that the Bratteli argument of 90.2(b) survives the slack (block $r$ sites at the cut, track the inclusion matrices of the left and right partial lifts), forcing $[O][x] = [x][O]$ in $K(\mathcal{D})$ for every simple $x$ of $\mathcal{D}$; or exhibit a local dyadic lift for the $\mathrm{Vec}_{S_3}$ row. Either outcome settles R0$^u$ at L0 up to the associator equations (90.5).

## 91.9 What This Does Not Claim

No WT-1 A3 for any non-trivial state family; R1 is reached only for the vacuum product state. No critical state, gap, KS convergence, or rung $\geq$ R2; no periodic or parity-covariant lift; no decision for arbitrary local dyadic lifts at L0, and (90.6) is not claimed sufficient. Naturality (iv) concerns the tower construction, not R5. The depth bound concerns contiguous ancillas only. The finite checks are Float64 calibrations; the "not braided" status of $\mathrm{Vec}^{\omega}_{\mathbb{Z}_3}$ is a finite-grid search, not a proof (REVIEW:189–190).

## 91.10 Examples (Instances Only)

| Instance | Calibration datum for the general statements |
|---|---|
| any $\mathcal{C}$, $X = 0$; Vec, $O = \mathbb{C}^d$ | $\alpha_L = \mathrm{id}$ (excluded from (90.7)); tensor shuffle with $t =$ flip (REVIEW:144). |
| Fibonacci, $O = \mathbf{1} \oplus \tau$ | Sourced $F, R$ (`RSWfinal3.tex`:2379–2407), $\leq 4$ sites: all matrix-unit tests pass; phase-free crossing fails block naturality, error 0.786 (`output.txt`:4–6, `red_phase_free.txt`:12). |
| $\mathrm{Vec}_{\mathbb{Z}_3}$, $O = \mathbf{1} \oplus g$ | Non-self-dual $X$: passes $L = 2, 3$; the $g$–$g$ phase is tested only at $L = 4$ ($\mathbb{Z}_n$ needs $n$ old sites) (`review_pointed_lift_output.txt`:2–3). |
| semion, $O = \mathbf{1} \oplus 2g$; $\mathbf{1} \oplus \mathbf{1}' \oplus g$ | Multiplicity, nontrivial associator, unit summand in $X$: pass; phase-free crossing fails support (:4–6). |
| $\mathrm{Vec}^{\omega}_{\mathbb{Z}_3}$, $O = \mathbf{1} \oplus g$ | Non-braided (grid search): three pointed half-braidings; Theorem 90.1 checks pass at $L = 4$; also $O = \mathbf{1} \oplus g \oplus g^2$ (:7–11). |
| $\mathrm{Vec}_G$, $O = \mathbf{1} \oplus g$, $g$ non-central | $\langle g \rangle$ cyclic with symmetric braiding, so 90.1 applies; $[O][h] - [h][O] = [gh] - [hg] \neq 0$: no lift to $Z(\mathrm{Vec}_G)$ (NOTE:283). |
| $\mathrm{Vec}_{\mathbb{Z}/2}$, non-central point | Half-braiding on $O$ with $\eta$ not central: $\alpha(h)V = -Vh$ (NOTE:284; `review_pointed_lift_output.txt`:12). |
| $\mathrm{Vec}_{S_3}$, $O = \mathbf{1} \oplus g \oplus h$, $gh \neq hg$ | $\Delta \neq 0$ on $\langle O \rangle = S_3$ (four clashing entries at $M = 2, s = 1$): no one-gap lift (90.2(b)); dyadic lifts undecided (Target 90.T) (`review_one_gap_obstruction_output.txt`:1–4). |

# 92 Parity, Mixed Koo–Saleur Brackets, and the Chiral/Full Delineation

## 92.1 Status, Sources, and Dependencies

**Status: conditional R2 implications, derived negatives and finite ingredients; every statement carries its own label.** Lifted from the round-2 note E2, graded WEAKER-RESULT-PROVED with *no invalid step* (REVIEW.md:11–16, :147–149). Theorems 91.1, 91.5, 91.6, Propositions 91.2, 91.7, 91.14 and Corollary 91.10 are implications from hypotheses (two VR-1 certificates on one system, a linear parity, ZC/MW/PC) that no categorical chain is shown to satisfy: *conditional*. 91.3, 91.8, 91.12 are *derived, negative*; 91.4 is *derived* (reviewer); 91.9 is a reviewer's numerical observation; 91.11, 91.13, 91.15 are *derived*, finite. The brief's conjecture is *not* proved as worded (REVIEW.md:214–246): part (b) needs the added zero-mode hypothesis, parity transport needs PC, and part (c) holds only in the restricted form 91.10. **Layer tags (CONVENTIONS (aq)).** 91.1–91.10 and 91.14: (full; category-free; R2 [cond.]); 91.10 is the delineation, speaking about chiral targets from the full setting; 91.3 and 91.8 are category-free abstract counterexamples testing the R2 interface. 91.11–91.13 and 91.15: (any; L0; $\to$ R0, finite). The note's own "L0" tags (NOTE.md:762–769) are superseded by the reviewer's retag (REVIEW.md:43–44, :156–166). Reproducibility: `checks/check_e2.py` (exact SymPy; `raw_output.txt`:1–13, identities pass, six mutations rejected) and `checks/review_e2_checks.py` (reviewer, numpy, a $\mathrm{Vec}_{\mathbb{Z}_3}$ non-self-dual instance; `review_raw_output.txt`:1–10), both in the E2 directory. No Julia port.

Dependencies: CA-79 ((1), Lemma 79.1, Theorems 79.2, 79.8), CA-81 (VR-1), CA-82 (instance), CA-88; CONVENTIONS (m), (s), (v), (af), (al), (ap), (aq).

## 92.2 Two Corrections Recorded First

**(1) The reflection must be complex-linear** [derived; reviewer, REVIEW.md:179–187]. The brief's parity is "site reflection composed with the pivotal/dagger structure" (brief:67–68). Dagger is conjugate-linear (Penneys:145), and so is $f \mapsto (f^{\vee})^{\dagger}$ (NOTE.md:641–653). Remark 91.4 shows that such a reflection *preserves* each chirality and gives no $c_L = c_R$ constraint; every result below takes a complex-linear unitary $P_r$ as data. E3's reflection-positivity reflection is conjugate-linear (E3 NOTE.md:355–357), so RP reflection and chirality-exchanging parity are distinct data (INTEGRATION.md:102–106). **(2) Equal ledgers are a hypothesis.** The note's abstract (NOTE.md:8–10) derives mixed commutation from locality, two VR-1 certificates and current conservation and *omits* equal anomaly ledgers, which E2-Z assumes (NOTE.md:295; REVIEW.md:37–42, (M1)). Here every statement that uses them lists them; a linear parity supplies them (Theorem 91.1). Dropping them is open: when the mixed brackets are scalars, the 2-cocycle condition forces the defect to vanish, but the general case is unproved (REVIEW.md:208–212).

## 92.3 Setting

A tower $(H_r, J_r, K_r)$ as in VR-1 A1, $N = N_r \to \infty$; on $H_r$ a unitary translation $U_r$, $U_r^N = 1$, and self-adjoint velocity-one densities $k_j$ with $U_r k_j U_r^* = k_{j+1}$, $k_j$ supported on sites $j, j+1$, and $[k_j, k_l] = 0$ unless $j, l$ are equal or adjacent (cyclically). In a categorical instance $k_j = -h_{N,j}/v_N \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$ as in (ap); no categorical datum is used before §92.7. *Range qualification:* for a range-$R$ density the adjacent current is not the conserved-current prescription; range-$R$ is open (NOTE.md:33–41).

With $z$, $a_N = N/4\pi$, $\mathcal{J}_j$, $\varepsilon_N$, $b_N$ of (ap), rewrite (1) as

$$E_n = a_N \sum_j z^{nj}(k_j - \varepsilon_N) + b_N \delta_{n,0}, \quad B_n = a_N \sum_j z^{nj} \mathcal{J}_j, \quad L_n := T_{r,n} = E_n - B_n, \quad S_n := \bar{T}_{r,-n} = E_n + B_n, \tag{5}$$

$D_r := L_0 - S_0 = -2B_0$, total current $\mathcal{Q}_N := \sum_j \mathcal{J}_j$ (not the grading $Q_r$ of (ap)). Ledgers: $z^L_{mn} = \delta_{m+n,0}(m^3 c_L^* - m c_L)/12$, $z^R_{mn}$ likewise, with residuals $F^L_{mn} = [L_m, L_n] - (m-n)L_{m+n} - z^L_{mn}$, $F^R_{mn}$ the same for $\bar{T}$ (CONVENTIONS (s)). Graph norms $\|A\|_{(s,d)} = \|K^s A K^{-s-d}\|$ (CONVENTIONS (v)) obey $\|AB\|_{(s,d+e)} \le \|A\|_{(s,d)} \|B\|_{(s+d,e)}$. "Both families VR-1" means A1 for the common system and A2–A4 for each family with its own constants $C^{L,R}_{n,s}$, $q^{L,R}$ and ledger (VR-1:55–80).

### 92.4 Parity: Anomaly Balance without Convergence

**Theorem 91.1 (E2-P) [conditional; full; category-free; R2].** **ASSUME** both families VR-1 on one system, and complex-linear unitaries $P_r$ with $P_r k_j P_r^* = k_{-j}$, $[P_r, K_r] = 0$. **PROVE** (a) both families and all mixed words converge on one invariant domain $D$; (b) $c_L = c_R$ and $c_L^* = c_R^*$. ⟨1⟩1. VR-1's domain is built from $(J, K)$ only, and its word telescoping uses A2, A3 one letter at a time; label letters by (chirality, $n$). BY VR-1:115–137 (REVIEW.md:51–52). ⟨1⟩2. $P\mathcal{J}_j P^* = -\mathcal{J}_{-j-1}$, $PE_nP^* = E_{-n}$, hence exactly

$$P_r L_n P_r^* - \bar{T}_{r,n} = (z^{-n} - 1)B_{-n}, \qquad \|\cdot\|_{(s,1)} \le \pi|n|(C^R_{n,s} + C^L_{-n,s})/N. \tag{6}$$

BY linearity of $P_r$, the substitution $l = -j-1$, and A2 (NOTE.md:116–129). ⟨1⟩3. Conjugate $F^L_{mn}$ by $P_r$, replace each $PL_jP^*$ by (6) and subtract $F^R_{mn}$: $\|(z^R_{mn} - z^L_{mn})1\|_{(s,2)} \le q^L + q^R + O_{m,n,s}(N^{-1})$, uniformly in $r$. ⟨1⟩4. Do not assume a lower bound on $K_r^{-2}$: transport a fixed coarse vector $v \neq 0$ isometrically; A1 keeps $\|v\|$, $\|K^2 v\|$ constant, so $z^L_{mn} = z^R_{mn}$. At $(1,-1), (2,-2)$: $\Delta c^* - \Delta c = 0$, $8\Delta c^* - 2\Delta c = 0$, so $\Delta c = \Delta c^* = 0$. QED (NOTE.md:131–150).

**Proposition 91.2 (E2-P, transport) [conditional; full; category-free; R2].** If in addition $P_r^2 = 1$ and $P_{r+1} J_r = J_r P_r$ (PC), then $Pi_r v := i_r P_r v$ extends to a unitary involution preserving $D$ with $PT_nP^* = \bar{T}_n$ on $D$. BY PC (well-defined), $[P_r, K_r] = 0$ (isometric on every $H_s$), and (6) in the limit (NOTE.md:152–158). Only this statement uses PC.

**Negative 91.3 (E2-P0) [derived; negative].** For any $\mathcal{C}$, $O = \mathbf{1} \oplus \mathbf{1}$, embeddings appending $|0\rangle$, $K = 1$, zero modes and $P_N$ the site reversal satisfy every hypothesis of 91.1, but $P_N$ sends the embedded first-site excitation to mutually orthogonal vectors: no strong limit, on any subsequence (NOTE.md:163–176). This refutes convergence of the *specified* $P_r$ only; here the identity intertwines.

**Remark 91.4 (antiunitary reflection) [derived; reviewer, REVIEW.md:179–187].** For antiunitary $\Theta$ with $\Theta k_j \Theta^* = k_{-j}$: $\Theta \mathcal{J}_j \Theta^* = +\mathcal{J}_{-j-1}$, $\Theta E_n \Theta^* = E_n$ and $\Theta T_{r,n} \Theta^* = E_n - z^n B_n = T_{r,n} + O(|n|/N)$ in $(s,1)$. $\Theta$ preserves each family and constrains nothing. Checked exactly at $N = 5, 6$ (`review_raw_output.txt`:7–8).

### 92.5 Mixed Brackets Reduce to One Zero-Mode Commutator

**Identity (E2-M ⟨1⟩1) [derived].** With $C_{mn} = [L_m, S_n]$, $A_{mn} = C_{mn} - C_{nm}$, $\kappa_{mn} = -ia_N(z^n - z^m)$ (Lemma 79.1) and *equal ledgers*,

$$A_{mn} = (4\kappa_{mn} - 2(n-m))B_{m+n} - F^L_{mn} - F^R_{-m,-n}, \qquad |4\kappa_{mn} - 2(n-m)| \le 2\pi(m^2+n^2)/N, \tag{7}$$

so $\|A_{mn}\|_{(s,2)} \to 0$. The central terms cancel *only* by equal ledgers (REVIEW.md:71–76).

**Theorem 91.5 (E2-Z) [conditional; full; category-free; R2].** **ASSUME** both families VR-1 on one system with *equal ledgers* (e.g. by 91.1). **PROVE** on $D$: all $[T_m, \bar{T}_n] = 0$ iff $[T_0, \bar{T}_0] = 0$. A sufficient finite certificate is

$$\|[T_{r,0}, \bar{T}_{r,0}]\|_{(s,2)} = \left\| \tfrac{N^2}{8\pi^2} [\textstyle\sum_j k_j, \mathcal{Q}_N] \right\|_{(s,2)} \to 0 \quad \forall s \qquad \text{(ZC)}, \tag{8}$$

since $[L_0, S_0] = 2[E_0, B_0] = 2a_N^2[\sum_j k_j, \mathcal{Q}_N]$ (Z0); ZC is sufficient, not claimed necessary. ⟨1⟩1. In the limit, (7) gives $C_{mn} = C_{nm}$; $C_{00} = 0$ by hypothesis. ⟨1⟩2. Jacobi and the cyclic sum give $C_{mn} = X_{m+n} := C_{m+n,0}$ (off-diagonal: $k = 0$; diagonal: the triple $(p, n-p, n)$, $p \notin \{0, n, n/2\}$); so $X_0 = 0$. ⟨1⟩3. Using $X_0 = 0$ explicitly: $[L_p, X_q] = -qX_{p+q}$, $[S_p, X_q] = qX_{p+q}$. ⟨1⟩4. $[[L_1, S_{-1}], X_q] = -q(q-1)X_q + q(q+1)X_q = 2qX_q$, while $[L_1, S_{-1}] = X_0 = 0$; so $X_q = 0$. QED (NOTE.md:305–349; REVIEW.md:89–101). Exact conservation of the total energy current implies ZC; locality alone does not (NOTE.md:351–357). Another sufficient condition: the current–momentum identification (12) of 91.7 with $p_r$ a spectral function of $U_r$, since then $[p_r, \bar{T}_{r,0}] = 0$ (NOTE.md:358–363).

**Theorem 91.6 (E2-M) [conditional; full; category-free; R2].** **ASSUME** 91.5's hypotheses and $\|[D_r, \bar{T}_{r,n}] - n\bar{T}_{r,n}\|_{(s,2)} \le \mu_{r,n,s} \to 0$ (MW). **PROVE** explicit $b_{r,m,n,s} \to 0$, computable from finitely many A2, A4, MW constants, with $\|[T_{r,m}, \bar{T}_{r,n}]\|_{(s,4)} \le b_{r,m,n,s}$ (graph loss four). ⟨1⟩1 (7). ⟨1⟩2 $W_n = [L_0, S_n] = ([D_r, S_n] + nS_n) + F^R_{0,-n}$, small in $(s,2)$. ⟨1⟩3 $(n-m)C_{mn} = [L_0, A_{mn}] + nA_{mn} - V_{mn} + V_{nm}$: off-diagonal loss three. ⟨1⟩4 with $a = |n| + 1$, $b = n - a$: $(a-b)C_{nn} = [L_a, C_{bn}] - [L_b, C_{an}] - [F^L_{ab}, S_n]$: loss four. QED; the bounds (7)–(11) are at NOTE.md:199–290. If all $q, \mu = O(N^{-\gamma})$, $b = O(N^{-\min(1,\gamma)})$. *Reviewer remark* (REVIEW.md:84–87): in the limit MW is *equivalent* to $[T_0, \bar{T}_n] = 0$ for all $n$; 91.6's content is the finite-stage rate and the upgrade from that row to all $(m, n)$.

**Proposition 91.7 (E2-S) [conditional; full; category-free; → R2].** **ASSUME** 91.6 without MW, and self-adjoint $p_r$ with $U_r = e^{2\pi i p_r/N}$ (lifts of the phases; extra data) and

$$\|D_r - p_r\|_{(s,1)} \le \delta_{r,s} \to 0 \quad (12), \qquad \|[p_r, \bar{T}_{r,n}] - n\bar{T}_{r,n}\|_{(s,2)} \le \nu_{r,n,s} \to 0 \quad (13).$$

**PROVE** MW with $\mu \le \nu + \delta_{r,s} C^R_{n,s+1} + C^R_{n,s}\delta_{r,s+1}$. BY $U\bar{T}_n U^* = z^n \bar{T}_n$ (blocks $q \to q'$ need $q' - q - n \in N\mathbb{Z}$) and the product rule (NOTE.md:367–394). A sufficient full-output form of (13) is $\|\bar{T}_{r,n} F_{r,n}\|_{(s,2)} = O(N^{-1-\epsilon})$ for the projection $F_{r,n}$ onto $p$-eigenspaces within $|n|$ of the branch ends, with $[p, K] = 0$ (NOTE.md:396–409). *The substantive gap is (12)*; a spectral fit "energy ≈ |momentum|" does not prove it, and finite range alone leaves an $O(N^3)$ bound on a mixed sum (NOTE.md:411–420).

**Negative 91.8 (E2-V) [derived; negative; category-free, not a KS chain].** On $\ell^2(\mathbb{Z})$ put $A_n e_k = (k - n/2)e_{k-n}$, $Ke_k = (1 + |k|)^p e_k$ ($p > 1$ integer), $H_r = \text{span}\{e_k : |k| \le 2^r\}$, $T_{r,n} = \bar{T}_{r,n} = \pi_r A_n \pi_r$, $P_r = 1$. Both families satisfy A1–A4 with $c = c^* = 0$, coherent exchange and linear zero-mode spectrum, yet $[T_1, \bar{T}_{-1}]e_1 = 2e_1$ (NOTE.md:422–462; A4 residual $\sim M^{2-2p}$ re-run, `review_raw_output.txt`:9–10). Scope: refutes inference from A1–A4 alone; the copies lack opposite spatial momenta and (7) fails, so it is *not* a counterexample to 91.5's open converse.

**Remark 91.9 (ZC is a selection condition at finite $N$) [numerical; reviewer].** $\|[H, \mathcal{Q}]\| = 0$ for an XXZ-type density but 3.1 ($N = 5$), 4.7 ($N = 6$) after a swap-symmetric term $\beta(n_j + n_{j+1})$ (`review_raw_output.txt`:3–6; REVIEW.md:191–196): exact ZC depends on the density split, not on $H$. The graph-norm limit is not settled.

## 92.6 The Delineation

**Corollary 91.10 (E2-C) [conditional; full, delineating chiral targets; category-free; R2].** **ASSUME** 91.1's hypotheses, and *absent right stress*: $\bar{T}_n = t\,\delta_{n,0}1$ on $D$. **PROVE** $T_n = t\,\delta_{n,0}1$,

$c_L^* = c_R^* = 0$, $c_L = c_R = 24t$; in the normalisation $c = c^*$, $t = 0$. PC is not needed. $\langle 1\rangle 1$. Barred $(1,-1),(2,-2)$: $0 = 2t + (c_R^* - c_R)/12$, $0 = 4t + (8c_R^* - 2c_R)/12$; so $c_R^* = 0$, $c_R = 24t$. $\langle 1\rangle 2$. 91.1(b) transfers the ledger. $\langle 1\rangle 3$. VR-1 gives $\|\bar{T}_{r,n} - t\delta_{n,0}1\|_{(s,1)} \to 0$; conjugate by $P_r$ and use (6). QED (NOTE.md:478–500; REVIEW.md:119–124). The valid contrapositive is a *disjunction* (NOTE.md:507–516): a nonzero purely chiral target from a periodic chain must drop a linear microscopic parity, *or* the common two-family certificate (refinement, state or retained carrier), *or* the standard modes or prescribed limit. "Chiral targets require parity-breaking densities" is *not* proved. A chiral subnet of a larger limit is not absent right stress (NOTE.md:503–505).

## 92.7 Finite Ingredients for Parity Breaking

Here $\mathcal{C}$ is unitary fusion, $k_j \in \mathrm{End}_{\mathcal{C}}(O \otimes O)$, $P_N$ a declared linear reflection.

**Proposition 91.11 (E2-D) [derived; any; L0; → R0].** $H_N(\lambda) = \sum_j k_j - \lambda \mathcal{Q}_N$, $\lambda \in \mathbb{R}$, is a self-adjoint range-three $\mathcal{C}$-morphism Hamiltonian with $P_N H_N(\lambda) P_N^* = H_N(-\lambda)$; it breaks $P_N$ iff $\lambda \mathcal{Q}_N \neq 0$. BY $P\mathcal{J}_j P^* = -\mathcal{J}_{-j-1}$ (91.1 $\langle 1\rangle 2$), so $P\mathcal{Q}_N P^* = -\mathcal{Q}_N$ (NOTE.md:529–551). For any local self-adjoint $a$, the translated sum of $a - \rho(a)$ works if that *sum* is nonzero.

**Negative 91.12 (E2-D0) [derived; negative; any; L0].** No perturbation exists on every fixed carrier: for $O = \mathbf{1}$, $\mathrm{End}_{\mathcal{C}}(\mathbf{1}) = \mathbb{C}$ (Penneys:442), every density is scalar and $\mathcal{Q}_N = 0$.

**Proposition 91.13 (E2-A) [derived; any; L0; → R0].** On $O' = O \otimes (\mathbf{1} \oplus \mathbf{1})$, $N \geq 4$, with Pauli flags (CONVENTIONS (m)) and the *bond-centred* site reflection $x \mapsto 1 - x$ (the convention making $Pk_jP^* = k_{-j}$ possible; REVIEW.md:132–134), $g_j = X_jX_{j+1} + Z_jZ_{j+1}$ gives $i[g_j, g_{j+1}] = 2(X_jY_{j+1}Z_{j+2} - Z_jY_{j+1}X_{j+2})$, a nonzero sum of $2N$ Hilbert–Schmidt-orthogonal words; 91.11 applies. It changes $O$, adds degrees of freedom and supplies *no* chiral limit (NOTE.md:564–601).

**Proposition 91.14 (E2-T) [conditional; full; category-free; R2 in the fixed representation].** Without changing states or modes, $2a_N(H_N(\lambda) - N\varepsilon_N) + 2b_N = (L_0 + S_0) + \lambda D_r \to (1+\lambda)T_0 + (1-\lambda)\bar{T}_0$ on $D$; at $\lambda = 1$ the operators $\bar{T}_n$ and their ledger remain: a tilt is not deletion (NOTE.md:603–619). No general "discard one Fermi edge" functor is established; the sharp Fourier filter onto $M$ modes has $\Pi_{xy} = (1 - z^{M(x-y)})/[N(1 - z^{x-y})]$, nonzero at every odd separation for $M = N/2$ (NOTE.md:621–634).

**Lemma 91.15 (E2-R, conjugate-dual mirror) [derived; any; L0 + unitary spherical duality; → R0].** On $V_N = \bigoplus_c \mathrm{Hom}_{\mathcal{C}}(c, O^{\otimes N})$, $f \mapsto (f^\vee)^\dagger$ is conjugate-linear and changes $O \to O^\vee$, $c \to c^\vee$ (Penneys:145, :682–698). With $W_N$ the conjugate Hilbert space of $\bigoplus_c \mathrm{Hom}(c^\vee, (O^{\otimes N})^\vee)$, $R_N f = \overline{(f^\vee)^\dagger}$ is a linear unitary, $\mu_N(a) = \overline{(a^\vee)^\dagger}$ a support-reversing $*$-representation, and on $V_N \otimes W_N$ the swap $P$ with $d_j = k_j \otimes 1 + 1 \otimes R_N k_{-j} R_N^*$ satisfies $Pd_jP^* = d_{-j}$ and PC for mirrored refinements. No self-duality or braiding is used (Penneys:917–935, :992–999; NOTE.md:665–698). It doubles the theory; it supplies no state or mode estimate.

## 92.8 What This Says about Rule 17's Chirality Layer

The results *delineate*; they do not *restrict*. The layering principle keeps chiral and full targets both in scope (AGENTS.md:100–103); 91.10 says exactly which hypotheses a chiral target obtained from a periodic chain must drop, and nothing more. $\mathsf{S}_{\mathrm{ch}}$ remains a declared tag: a chiral net may arise as a subnet of a full limit (D6 REVIEW.md:94–98), from a parity-breaking construction, or with modified modes (CA-82's sine chain is reflection-odd and uses modified modes, so it evades 91.10 through the disjunction; INTEGRATION.md:118–120). The Definition 88.1 source gap is narrowed in the restricted parity-paired standard-KS sense only; the general lattice-chirality question remains a source gap.

**Next lemma (zero-mode, current–momentum) [conjectural target].** Under 91.1's hypotheses with $U_r$ covariant for both families and $[U_r, K_r] = 0$, prove $\|\frac{N_r^2}{8\pi^2}[\sum_j k_j, \mathcal{Q}_r]\|_{(s,2)} \to 0$ for every $s$, via (12) for a spectral logarithm $p_r$ of $U_r$; *or* exhibit a nearest-neighbour chain (any $\mathcal{C}$, even Vec) with both VR-1 certificates and $[T_0, \bar{T}_0] \neq 0$ in the limit (INTEGRATION.md:125–131; REVIEW.md:248–252). Either settles the brief's part (b). *Further open* (NOTE.md:794–817): PC from a concrete OAR construction; the range-$R$ current; a category-uniform chiral extraction; R3/R4 completion.

## 92.9 What This Does Not Claim

That ZC follows from the other hypotheses; that PC follows from finite symmetries; that chiral targets require parity-breaking densities; any categorical edge-discard functor; any categorical chain meeting either VR-1 certificate; that a Virasoro *algebra* action on $D$ is the projective-unitary symmetry of the north star (NOTE.md:771–774); anything at R3–R5.

| Instance | Calibration datum for the general statements |
|---|---|
| Any $\mathcal{C}$, $O = \mathbf{1}$ | 91.12: all local endomorphisms scalar; no fixed-carrier parity breaking. |
| Any $\mathcal{C}$, flags $O \otimes (\mathbf{1} \oplus \mathbf{1})$ | Exact (6), (7), 91.13 at $N = 4$; current HS norm$^2$ = 512 (`raw_output.txt`:2). |
| $\mathrm{Vec}_{\mathbb{Z}_3}$, $O = \mathbf{1} \oplus g$ | Non-self-dual: (6), (7), Z0, $P\mathcal{Q}P^* = -\mathcal{Q}$ exact at $N = 5, 6$; site reversal is linear (REVIEW.md:171–174). |
| Witt shifts (91.8) | Separate certificates, mixed $(1, -1)e_1 = 2e_1$; not a KS chain. |
| Ising compiler (CA-82) | Chiral by a reflection-odd sine $H$ with modified modes: evades 91.10 via the disjunction. |

# 93 Spatial Seams, Reflection Pairings and a Finite Osterwalder–Schrader Lemma

## 93.1 Status, Sources, and Dependencies

**Status: mixed; every statement carries its own label.** Source grade: WEAKER-RESULT-PROVED (REVIEW.md:10–23). Theorems 92.3 and 92.5 are *derived, conditional* on a declared seam pairing $s$ and on the reflected-half hypothesis (RH), which the review found *assumed through the definition of the pairing, not derived* (REVIEW.md:116–120). Lemma 92.2 is *derived*; Proposition 92.4 is *conditional*; Propositions 92.6 and 92.9 are *derived, negative*; Remark 92.7 is *derived* in Vec and *conjectural* in general; Lemmas 92.8 and 92.10 are *derived* and category-free; the reflection-coherence lemma is a *proposal*. This shard addresses the spatial-RP gap stated at CA-80:137–140. It does *not* answer CA-75 Link (a2), which asks for a spatial *sewing* identity and an anomaly-line limit (REVIEW.md:51). Reproducibility: exact SymPy/NumPy scripts `checks/check_seams.py`, `checks/check_categories.py` (with recorded mutations) and the reviewer's independent `checks/review_checks.py` (19 PASS); no Julia port. NOTE, REVIEW, INTEGRATION below are the files in `explorations/2026-10-22-astra-r2/E3-reflection-positivity/`.

**Layer tags (CONVENTIONS (aq)).** 92.3–92.5: ($\mathsf{S}_{\text{full}}$ for the untwisted reflection; chiral only with a declared twisted reflection; $\mathsf{L0}$ plus the microscopic declaration $s$; finite, $\to \mathsf{R3}$). A seam pairing is a declaration, not a data layer (CA-88:49–51); the note's tags "L3 over L0" and "R0" are superseded (REVIEW.md:236–254; INTEGRATION.md:4–6, :119–122). 92.6, 92.9: same setting, $\mathsf{L0}$, finite. 92.8, 92.10: (any; category-free; finite, 92.8 $\to \mathsf{R3}$).

Dependencies: CA-78 (78.1, 78.2, 78.7, erratum), CA-80 (80.1–80.4, :137–140), CA-86 (86.1), CA-88 (88.2–88.3); CONVENTIONS (b), (r), (ak), (al), (am), (ao), (aq).

## 93.2 Setting and the Seam Pairing

$\mathcal{C}$ is unitary fusion, $O \neq 0$ a site object (CONVENTIONS (r); (am) for $O = \mathbf{1} \oplus \mathbf{1}$), with dual $(\overline{O}, \mathrm{ev}_O : \overline{O} \otimes O \to \mathbf{1}, \mathrm{coev}_O : \mathbf{1} \to O \otimes \overline{O})$ from the unitary spherical structure, raw cups of squared norm $d_O$ (CONVENTIONS (b); Penneys:992–1030). Tiles, $h$ and $H = -\sum h_j$ follow (ak), (al). The *linear history* tile $1 + th$ (CA-86 Lemma 86.1) and the *exact* tile $e^{th}$ (CA-78 Prop. 78.2) are kept apart. Put $W_+ = O \otimes \overline{O}$, $W_- = \overline{O} \otimes O$, with cut spaces $\mathcal{H}(W) = \bigoplus_a \mathrm{Hom}(a, W)$ as in CA-80. *Spatial RP* means positivity over all linear combinations of half-patch insertions, not of one scalar partition sum (NOTE.md:52–55).

**Definition 92.1 [declaration; $\mathsf{L0}$].** A *seam pairing* is a unitary $s : W_+ \to W_-$ with $s\,\mathrm{coev}_O = \mathrm{ev}_O^\dagger$, declared with a rule pairing all non-seam tiles with their reflected adjoints. The rule is antilinear, involutive and compatible with dagger and sewing; the reverse seam uses $s^\dagger$ (NOTE.md:37–47). It is fixed before $h$. *Gauge (REVIEW.md:122–132):* $s \circ W$ with $W$ unitary in $\mathrm{End}_{\mathcal{C}}(W_+)$ and $W\mathrm{coev}_O = \mathrm{coev}_O$ is equally admissible and changes the kernel to $W^\dagger K_s$. On non-vacuum blocks and on $\mathrm{coev}_O^\perp$ $W$ is free, so polar decomposition makes those blocks PSD for *some* $s$: single-tile RP is a property of $(\mathcal{C}, O, h, s)$.

**Lemma 92.2 [derived; existence of $s$].** A unitary $s$ with $s\,\mathrm{coev}_O = \mathrm{ev}_O^\dagger$ exists iff $\dim \mathrm{Hom}(a, W_+) = \dim \mathrm{Hom}(a, W_-)$ for all $a \in \mathrm{Irr}\,\mathcal{C}$ (INTEGRATION.md:17–18); in particular when $O \cong \overline{O}$. $\langle 1\rangle 1$. Necessity: a unitary isomorphism preserves multiplicities (NOTE.md:216–218). $\langle 1\rangle 2$. Sufficiency: matching orthonormal bases of each $\mathrm{Hom}(a, W_\pm)$ give a unitary $V$; $V\mathrm{coev}_O$ and $\mathrm{ev}_O^\dagger$ lie in $\mathrm{Hom}(\mathbf{1}, W_-)$ with equal norm $\sqrt{d_O}$, so a unitary $U$ on that block maps one to the other; set $s = (U \oplus 1)V$. $\langle 1\rangle 3$. If $\psi : O \to \overline{O}$ is an isomorphism, $\psi \otimes \psi^{-1} : W_+ \to W_-$ is one, so the multiplicities agree. QED. $s$ alone does not supply the whole reflection rule (NOTE.md:45–47). At $\mathsf{L1}$ the identity

$\mathrm{ev}^\dagger = c_{O,\overline{O}}(\theta_O \otimes 1)\mathrm{coev}$ would supply $s$; it is a reviewer remark with *no local source* (Law 1 gap, REVIEW.md:241–245) and is not used. Self-duality need not exist (Penneys:917).

### 93.3 The Bent Kernel and Single-Tile Spatial RP

With associators inserted for the named bracketings, and $v_{a,\mu} : a \to W_+$ isometric inclusions,

$$\mathrm{rot}(T) = (1_{\overline{O}} \otimes 1_O \otimes \mathrm{coev}_O^\dagger)(1_{\overline{O}} \otimes T \otimes 1_{\overline{O}})(\mathrm{ev}_O^\dagger \otimes 1_O \otimes 1_{\overline{O}}), \quad K_s(T) = s^\dagger \,\mathrm{rot}(T), \quad K^a_{\mu\nu} = d_a^{-1}\,\mathrm{tr}_{\mathcal{C}}\left(v^\dagger_{a,\mu} K_s(T) v_{a,\nu}\right). \tag{92.1}$$

*Evaluation prescription* (NOTE.md:144–173): expand $O, \overline{O}$ into simples keeping every copy and vertex index; place a tile on a non-adjacent slot as $F^\dagger_{\mathrm{path}}(1 \otimes D_\alpha \otimes 1)F_{\mathrm{path}}$ (CA-78:44–53); insert the oriented cups in their actual slots; multiply by $s^\dagger$, not by an unprovided braiding; divide Penneys-normalised vertices by $(d_i d_j/d_k)^{1/4}$ (Penneys:530–568). Each entry is then a finite sum of products of $F$-symbols, pivotal cups and entries of $s$, with all multiplicities; in arbitrary cut bases use the inverse Gram of CA-80:70–81.

**(RH) [hypothesis, not derived].** After bending, the reflected half-patch, read through $s$, evaluates to the adjoint of the half-amplitude.

**Theorem 92.3 [derived, conditional on $s$ and (RH); E3-S]. Assume** Definition 92.1, one crossing tile $T = \sum_\alpha w_\alpha D_\alpha$, half-insertions spanning $\mathcal{H}(W_+)$, and (RH). **Prove** spatial RP iff every block $K^a = [K^a_{\mu\nu}]$ of (92.1) is PSD. ⟨1⟩1. rot is type-correct and complex-linear into $\mathrm{Hom}(W_+, W_-)$ (intermediate word $\overline{O} \otimes O \otimes O \otimes \overline{O}$); it is not an endomorphism. BY Penneys:398–430. ⟨1⟩2. $v^\dagger X v \in \mathrm{End}(a) = \mathbb{C}$ and $\mathrm{tr}_{\mathcal{C}} = d_a \cdot$scalar, so $K^a$ is the $a$-block of $K_s(T)$. BY Penneys:1010–1019. ⟨1⟩3. BY (RH), all remaining cuts run along tile legs and CA-80 Lemma 80.1 gives the mixed form $q(x, y) = \langle x, K_s(T)y\rangle$. This is where (RH) enters; the review: E3-S is "close to a definition plus Lemma 80.2; the real content is the explicit oriented formula" (REVIEW.md:116–120). ⟨1⟩4. CA-80 Lemma 80.2 with spanning; $d_a > 0$ is harmless. QED. The Hermiticity equations $K^a_{\mu\nu} = \overline{K^a_{\nu\mu}}$ are *spatial*, not the temporal dagger pairing of Cor. 80.3. Theorem 92.3 is the typed form of D5's crossing-kernel test (D5 NOTE.md:231–239; REVIEW.md:57).

**Proposition 92.4 [conditional; multi-tile sufficiency]. Assume** several crossing tiles whose ports and pairings tensor and reassociate compatibly, disjoint along the seam, all other tiles in reflected pairs, any extra seam insertion positive and placed inside the reflected factorisation. **Prove** $K_{s,i} \geq 0$ for all $i$ implies all-patch spatial RP. ⟨1⟩1. $K_{s,i} = B_i^\dagger B_i$ blockwise, so $\bigotimes_i K_{s,i} = (\bigotimes B_i)^\dagger(\bigotimes B_i) \geq 0$. ⟨1⟩2. Reassociation is unitary; charge restriction and half-patch attachment give a compression $V^\dagger(\bigotimes K_{s,i})V \geq 0$. QED (NOTE.md:201–214). Necessity holds only if the admitted patches include single-tile full-span tests. Compatibility is not automatic for non-braided $\mathcal{C}$ (planar order of the seam word) and may be unsatisfiable (REVIEW.md:133–141). On a translation-invariant brick lattice "bulk tiles in reflected pairs" is $\Theta_x(h) = h$.

### 93.4 The Linear History: an Exact Schur Criterion

Let $u = \mathrm{coev}_O$, $e = u/\sqrt{d_O}$, $P = |e\rangle\langle e|$, $Q = 1 - P$, $A = K_s(h)$, $a = \langle e, Ae\rangle$, $b = QAe$, $C = QAQ|_{Q\mathcal{H}}$, $C^+$ the inverse on $\mathrm{Ran}\, C$.

**Theorem 92.5 [derived, conditional on $s$, (RH), one crossing tile with full span; E3-L]. Prove** (i) $K_s(1) = uu^\dagger = d_O P$; (ii) $1 + th$ is spatially RP for all $0 < t < \varepsilon$, some $\varepsilon > 0$, iff

$$A = A^\dagger, \qquad C \geq 0, \qquad b \in \mathrm{Ran}\, C; \tag{92.2}$$

(iii) at a given $t > 0$ iff (92.2) and $d_O + t(a - b^\dagger C^+ b) \geq 0$, i.e. for $t \leq t_* = d_O/(b^\dagger C^+ b - a)$ when the denominator is positive, and for all $t$ otherwise. ⟨1⟩1. At $T = 1$ the interchange law cancels the

associators: $\mathrm{rot}(1) = \mathrm{ev}_O^\dagger \mathrm{coev}_O^\dagger$; unitarity of $s$ gives $s^\dagger \mathrm{ev}_O^\dagger = \mathrm{coev}_O$. Hence

$$K_s(1+th) = \begin{pmatrix} d_O + ta & tb^\dagger \\ tb & tC \end{pmatrix}. \tag{92.3}$$

Checked with nontrivial 3-cocycles, including non-self-dual $O$ (`review_output.txt`:2–4). ⟨1⟩2. Positivity at any $t > 0$ forces $A = A^\dagger$ and $C \geq 0$; for $z \in \ker C$ the term linear in $\lambda$ on $\lambda e + z$ forces $b^\dagger z = 0$, so $b \in (\ker C)^\perp = \mathrm{Ran}\, C$. ⟨1⟩3. Under (92.2), $\langle(\alpha, y), K_s(1+th)(\alpha, y)\rangle = t\|C^{1/2}(y + C^+ b\alpha)\|^2 + (d_O + ta - tb^\dagger C^+ b)|\alpha|^2$; $y = -C^+ b\alpha$ gives necessity. ⟨1⟩4. $d_O > 0$ and finite dimension give (ii) from (iii). QED (NOTE.md:233–272). $\varepsilon$ depends only on $(d_O, a, b, C)$, so for identical crossing tiles it is *uniform in chain length $N$ and row number $M$* (REVIEW.md:159–162). The criterion is not $K_s(h) \geq 0$ (Prop. 92.6(iii)). Checks: 300 random pencils and a range failure (`review_output.txt`:31–32; `seams_output.txt`:6). *Exact gates [derived remark].* For $e^{th}$ the criterion is only pointwise, $K^a(w(t)) \geq 0$ with the CA-78 ODE; RP near $t = 0$ implies $A = A^\dagger$ and $QAQ \geq 0$, but the range condition is not inferable: $\left[\begin{smallmatrix} 1 & t \\ t & t^2 \end{smallmatrix}\right] \geq 0$ has derivative $\left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$, whose pencil fails it (an illustration, not a categorical gate; NOTE.md:274–282).

## 93.5 Negative Results and Pairing Dependence

**Proposition 92.6 [derived, negative; E3-N, flip pairing on a unit-multiplicity corner].** In any $\mathcal{C}$ take $O = \mathbf{1} \oplus \mathbf{1}$ and the flip pairing. Then $K(T)_{(ik),(jl)} = T_{(il),(kj)}$ and $K(A \otimes B) = |\mathrm{vec}A\rangle\langle \mathrm{vec}B^\dagger|$ (N1), with $Z = \mathrm{diag}(1, -1)$: (i) $h = -Z \otimes Z = h^\dagger$, $1 + th > 0$ for $0 < t < 1$, yet $\langle \mathrm{vec}Z, K(1+th)\mathrm{vec}Z\rangle = -4t$; the exact gate gives $-4 \sinh t$. Self-adjointness is not sufficient. (ii) $h = E_{01} \otimes E_{10} \neq h^\dagger$, $h^2 = 0$, and $K(e^{th}) = K(1+th) = |\mathrm{vec}1\rangle\langle \mathrm{vec}1| + t|\mathrm{vec}E_{01}\rangle\langle \mathrm{vec}E_{01}| \geq 0$. Self-adjointness is not necessary. (iii) $h = -1$: $K(h) \not\geq 0$, yet $K(1+th) = (1-t)|\mathrm{vec}1\rangle\langle \mathrm{vec}1| \geq 0$ for $t \leq 1$ and $K(e^{th}) \geq 0$. *Proof.* (N1) by hand from (92.1) (REVIEW.md:84–86) and on all 16 matrix units (`seams_output.txt`:2–5). □ *Reviewer additions* (REVIEW.md:87–107): (a) the witnesses *depend on $s$*: with $s' = \mathrm{flip} \circ (X \otimes \overline{X})$, $X$ the global on-site flip commuting with $h$, $K_{s'}(1 - tZ \otimes Z) \geq 0$ (`review_output.txt`:35); (i) refutes only the literal untwisted $\Theta_x$. (b) The witness transfers to the connected site $O = \mathbf{1} \oplus g$ in $\mathrm{Vec}^\omega_{\mathbb{Z}_2}$, $\mathrm{Vec}^\omega_{\mathbb{Z}_3}$, $Z = P_{\mathbf{1}} - P_g$: eigenvalue $-2t$, $\omega$-independent (`review_output.txt`:29–30). For general $O = \mathbf{1} \oplus X$ it needs $s$ natural for $\mathrm{End}(O)$: a remark, not proved.

**Remark 92.7 [derived in** Vec **with the flip; conjectural in general].** By (N1), $K_{\mathrm{flip}}(h)$ is Hermitian iff $h_{(il),(kj)} = \overline{h_{(jk),(li)}}$ for all indices, i.e. $h = Ph^\dagger P$ with $P$ the swap (`review_output.txt`:33). So untwisted spatial RP needs a parity-invariant $h$; the general-$\mathcal{C}$ form "$K_s(h)$ Hermitian iff $\Theta_x(h) = h$" is part (iii) of the coherence proposal below. Hence the setting tag "full; chiral only with a twisted reflection" (REVIEW.md:248–254; E2 link: INTEGRATION.md:100–105).

## 93.6 A Finite Osterwalder–Schrader Lemma (Category-Free)

**Lemma 92.8 [derived; E3-OS]. Assume** a finite space $V$, a shift $\tau$, a nonzero PSD form $q_0$, and $q_1(f, g) = q_0(f, \tau g)$ with $0 \leq q_1 \leq q_0$. **Prove** $T[f] = [\tau f]$ is a well-defined positive contraction on $\mathcal{H} = V/\mathrm{rad}\, q_0$; if $T$ is injective, $E = -a^{-1} \log T \geq 0$ and $T^n = e^{-naE}$; if $\tau 1 = 1$, $q_0(1, 1) = 1$, then $\Omega = [1]$ has zero energy; multi-time kernels are positive. ⟨1⟩1. For PSD $q$, $q(z, z) = 0 \Rightarrow q(f, z) = 0$ (vary $\lambda$ in $z + \lambda f$). ⟨1⟩2. $z$ null $\Rightarrow q_1(z, z) = 0 \Rightarrow q_0(f, \tau z) = 0$ $\forall f$: $T$ is well defined. ⟨1⟩3. $\langle [f], T[g]\rangle = q_1(f, g)$, so $T = T^\dagger$, $0 \leq T \leq 1$. ⟨1⟩4. Diagonalise; eigenvalues in $(0, 1]$ under injectivity. ⟨1⟩5. $\sum_{ij} \langle v_i, T^{n_i + n_j} v_j\rangle = \|\sum_i T^{n_i} v_i\|^2$. QED (NOTE.md:286–317). *Strictness* (NOTE.md:333–341): injectivity is needed for a Hamiltonian on the whole quotient ($K(1)$ above has rank one), and the lower bound is needed ($q_0 = 1$, $\tau = -1$). *What it does not give:* Theorem 92.3 supplies

$q_0$; the bound $q_1 \le q_0$, shift closure and compatibility of the two forms are unproved for any strip. No covariance, locality, field regularity or continuum. A reflection excludes positive momentum: $JPJ^\dagger = -P$, $P \ge 0$ force $P = 0$; positive energy and signed momentum (Schottenloher:4113–4117, :4150–4172), and covariance is a separate axiom (:4616–4642).

### 93.7 Two Directions: What Fails as Posed and What Survives

**Proposition 92.9 [derived, negative; brief E3(b), E3(c) as posed].** (i) The temporal and spatial transfers act on cut spaces of different objects and dimensions; a perpendicular commutator is undefined until a common carrier is identified (NOTE.md:366–376). (ii) rot is complex-linear and "$F$ composed with dagger" conjugate-linear, so they differ on any nonzero complex space (literal reading). Robustly, no single-tile similarity exists: the identity tile has full temporal rank and rank-one spatial $K(1)$. *Not refuted:* a relation between whole-lattice row and column transfers, e.g. $\operatorname{Tr} T_t^M = \operatorname{Tr} T_x^L$ on a torus, true for any tensor network without an $F$-move; E3(b) is false only in its literal clause (REVIEW.md:174–184). (iii) On one cut, YBE gates need not give commuting brick sweeps (CA-78 Prop. 78.7(i)) and commuting gates need not satisfy YBE under any analytic reparametrisation (78.7(ii)); for the perpendicular pair these are analogies (REVIEW.md:197–199).

**Lemma 92.10 [derived under its hypotheses; E3-Y, E3-D, E3-T].** (Y) **Assume** one carrier $\mathcal{H}$, auxiliary $A, B$, $L_{aj}(u)$ trivial on $B$, $L_{bj}(v)$ trivial on $A$, $R$ invertible on $A \otimes B$, $[L_{aj}, L_{bk}] = 0$ $(j \ne k)$, $RL_{aj}L_{bj} = L_{bj}L_{aj}R$, and a linear closure with $\mathrm{Cl}(M_aM_b) = t(u)t(v)$, $\mathrm{Cl}(M_bM_a) = t(v)t(u)$, $\mathrm{Cl}(R^{-1}XR) = \mathrm{Cl}(X)$. **Prove** $[t(u), t(v)] = 0$. ⟨1⟩1. Regroup $M_aM_b = \mathsf{A}_N \cdots \mathsf{A}_1$, $\mathsf{A}_j = L_{aj}L_{bj}$. ⟨1⟩2. $R\mathsf{A}_j = \mathsf{B}_j R$ for $j = N, \ldots, 1$. ⟨1⟩3. Apply Cl. QED. (D) With defects $D_j = R\mathsf{A}_j - \mathsf{B}_j R$, $[t(u), t(v)] = \mathrm{Cl}\big(R^{-1} \sum_k \mathsf{B}_N \cdots \mathsf{B}_{k+1} D_k \mathsf{A}_{k-1} \cdots \mathsf{A}_1\big)$; $N$ terms, no uniform bound, no converse. (T) On a common carrier two transfers starting at 1 commute to first order, and

$$[e^{\delta A}e^{\delta B}, e^{\delta B}e^{\delta A}] = -\delta^3[A+B,[A,B]] + O(\delta^4), \qquad A = \textstyle\sum_{\mathrm{odd}} h_j,\ B = \sum_{\mathrm{even}} h_j \tag{92.4}$$

(NOTE.md:378–473; `seams_output.txt`:7; `review_output.txt`:41). Identifying the two directional transfers with $t(u), t(v)$ of one closed family is *unproved* for general $\mathcal{C}$; (T) answers the brief's first-order request with zero (REVIEW.md:200–204).

### 93.8 The First Gap: Reflection Coherence

(RH) for a multi-tile brick seam is *assumed via the definition of the pairing, not derived*; $s$ is pinned only on $\mathrm{coev}_O$ and is gauge beyond it (REVIEW.md:289–299). **Reflection-coherence lemma [proposal; full; $\mathsf{L}0$ + a declared unitary self-duality $\psi : O \to \overline{O}$, or $\mathsf{L}1$; finite].** Build $\Theta_x$ from the antilinear anti-monoidal functor $X \mapsto (X^\vee)^\dagger$ (Penneys:682–708, :992–1003) composed with $\psi$ (or a braiding, or E2's mirror carrier). Prove (i) $\Theta_x$ is antiunitary on every cut space, compatible with Lemma 80.1 sewing, $\Theta_x^2$ trivial on observables; (ii) the induced $s$ satisfies $s\,\mathrm{coev}_O = \mathrm{ev}_O^\dagger$, Frobenius–Schur phases tracked; (iii) $K_s(h)$ Hermitian iff $\Theta_x(h) = h$ (INTEGRATION.md:124–136). With Prop. 92.4 this would give spatial RP of the whole CA-86 linear history for self-dual sites, a statement about $(\mathcal{C}, O, h)$, not $(\mathcal{C}, O, h, s)$. An FS phase is absorbed in the vacuum entry of $s$; $\Theta_x^2 = -1$ on pseudo-real sites is unexamined (REVIEW.md:270). Further open lemmas (NOTE.md:566–591): exact-gate, strip-translation, crossing-and-closure, continuum. *Layer observation:* a coherent $s$ is available for self-dual $O$ at $\mathsf{L}0$ with declared $\psi$, expected at $\mathsf{L}1$ (unsourced), and can fail for non-self-dual $O$ in non-braided $\mathcal{C}$: the $\mathsf{L}0/\mathsf{L}1$ split of $\mathsf{R}0^u$ (CA-88:104).

## 93.9 What This Does Not Claim

No whole-history iff for spatial RP for general $(\mathcal{C}, O, h)$; no coherent $s$ for the whole brick grammar; no spatial sewing identity across a plaquette seam and no anomaly-line limit (CA-75 Link (a2) stays open; only the seam operator exists); no crossing relation between directional transfers; no lattice OS axioms beyond Lemma 92.8; no positive momentum, Euclidean covariance or continuum. Nothing reaches R1–R5. Theorem 92.5 is for the linear history only; exact gates are characterised pointwise; multi-tile necessity is not claimed.

## 93.10 Examples (Instances Only)

| Instance | Calibration datum for the general statements |
|---|---|
| Any $\mathcal{C}$, $O = \mathbf{1} \oplus \mathbf{1}$ | Prop. 92.6: flip pairing; $-Z \otimes Z$ gives $-4t$; $E_{01} \otimes E_{10}$ positive; $h = -1$ (`seams_output.txt`:2–5). |
| $\mathrm{Vec}^{\omega}_{\mathbb{Z}_2}$, $\mathrm{Vec}^{\omega}_{\mathbb{Z}_3}$, $O = \mathbf{1} \oplus g$ | $K_s(1) = uu^{\dagger}$ with cocycles; witness eigenvalue $-2t$; $\omega$ changes $K_s(T)$ beyond the left gauge (`review_output.txt`:2–30). |
| $\mathrm{Vec}_{\mathbb{Z}_3}$, $O = \mathbf{1} \oplus g$, $\bar{g} = g^2$ | $T = 1 + \mathrm{Swap}/2$, graded flip $s$: blocks $\left[\begin{smallmatrix} 3/2 & 1 \\ 1 & 3/2 \end{smallmatrix}\right]$, [1/2], [1/2] (`categories_output.txt`:2–4). |
| $\mathrm{Rep}(A_4)$, real $V$, $N^V_{VV} = 2$ | $T = 1 + \mathrm{Swap}/2$, $s = \mathrm{Swap}$: blocks $7/2, 1/2, 1/2, \frac{1}{2}I_2$; $K$ via the fibre functor, so (92.1) is not exercised entry by entry; the multiplicity block is scalar (weak test; REVIEW.md:217–222). |
| $\mathrm{Vec}_{S_3}$, $O = \mathbf{1} \oplus (12) \oplus (123)$ | $W_+ \ncong W_-$: no $s$ (Lemma 92.2); self-dual $\mathbf{1} \oplus (12) \oplus (13) \oplus (23)$ has matching ports (`review_output.txt`:37–40). |

*Caveat (REVIEW.md:209–216):* both positive calibrations use a *symmetric braiding* as $s$ (the flip in abelian $\mathrm{Vec}_{\mathbb{Z}_3}$, Swap in $\mathrm{Rep}(A_4)$): a hidden L1 ingredient of the examples, not of the theorems. No positive instance has a non-braided $\mathcal{C}$ with a non-self-dual site.

# 94 Normal Limit Expectations and the Repaired Two-Sided Fidelity Certificate

## 94.1 Status, Sources, and Dependencies

**Status: mixed; every statement carries its own label.** Lemmas 93.1–93.2, Corollary 93.3 and Proposition 93.4 are *derived* (pure operator algebra); Examples 93.5–93.6 are *derived, negative*; Example 93.7 is *derived* and was added by the review; Theorem 93.8 and Corollary 93.9 are *derived implications* whose hypotheses are not supplied for any $\mathcal{C}$; Proposition 93.10 is *derived, negative*; Lemmas 93.11–93.13 are *conditional*; the bare minimal-constant variant is *conjectural*; the annular approximant lemma is a *proposal*. Reviewer grade: PROOF-COMPLETE under the stated hypotheses (REVIEW.md:10–21, :207–217). Ordinary structured proofs, not proof-assistant checked. Reproducibility: exact SymPy witnesses in `explorations/2026-10-22-astra-r2/E5-two-sided-fidelity/checks/` (`check_expectations.py`: 4 invariant groups, 5 mutations rejected; `review_check.py`: X1, X2, calibration); finite algebra calibrations only. Unqualified `NOTE.md`, `REVIEW.md`, `INTEGRATION.md` below are the E5 files of that directory. *Layer tags* (CONVENTIONS (aq)), (S; L; R): 93.1–93.7 (any; category-free; → R4); 93.8 (chiral; L0+L4 for a tensor equivalence, L1+L4 for a braided one; R4 sector sub-rung, given R3); 93.9–93.10 (full, with a named doubled chiral net; L2+L4; R4 for that net); 93.11–93.13 (chiral, or an identified chiral subnet; L2+L4; conditional R3 → R4).

Dependencies: CA-80 (Theorem 80.9, Proposition 80.8, Lemma 80.10, "The gap, explicitly"), CA-75 item 13, CA-88 (Definitions 88.1–88.3); CONVENTIONS (u), (al), (aq). This shard closes route (ii) of CA-80's gap.

## 94.2 Scoped Notation

All inclusions are unital, in $\mathcal{B}(\mathcal{H})$. For a normal conditional expectation $E : \mathcal{M} \to \mathcal{N}$ the *cost* is $K(E) := \inf\{\beta \geq 1 : E(x) \geq \beta^{-1}x \ \forall x \in \mathcal{M}_+\} = [\mathcal{M} : \mathcal{N}]_E$ (KLM:430–443; the infimum is attained when finite), and $q(\mathcal{N} \subset \mathcal{M}) := \inf_E K(E)$. For the four-interval inclusion of (al), $\mu_{\mathcal{A}} \leq K(E)$ for every $E$; KLM's minimum runs over all expectations, which does not affect this inequality (REVIEW.md:55–58). A *bounded exhaustion* of $\mathcal{M}$ by increasing $\mathcal{M}_k \subset \mathcal{M}$ means: every $x \in \mathcal{M}$ is the strong* limit of a norm-bounded sequence $x_k \in \mathcal{M}_k$; it is a hypothesis, not a density theorem (NOTE.md:58–63). $\psi$ is an *auxiliary* normal state, never the physical vacuum (Lemma 80.10); $\varphi := \psi|_{\mathcal{N}}$. $D := \operatorname{Dim}\mathcal{C} = \sum_i d_i^2$.

## 94.3 The Normal Limit Expectation

**Lemma 93.1 [derived; E5-G; any; category-free; → R4] (positive-graph passage). Assume** a normal expectation $F : \mathcal{M} \to \mathcal{N}$, normal expectations $E_k : \mathcal{M}_k \to \mathcal{N}_k$ inside $\mathcal{N} \subset \mathcal{M}$, $b = \liminf_k K(E_k) < \infty$ realised along $k_j$, and for each $x \in \mathcal{M}_+$ bounded positive $x_j \in \mathcal{M}_{k_j}$ with $x_j \to x$ and $E_{k_j}(x_j) \to F(x)$ ultraweakly. **Prove** $K(F) \leq b$. ⟨1⟩1. $E_{k_j}(x_j) - K(E_{k_j})^{-1}x_j \geq 0$, BY definition of $K$. ⟨1⟩2. Pairing with normal positive functionals and passing to the limit gives $F(x) - b^{-1}x \geq 0$, BY boundedness and closedness of the positive cone. QED (NOTE.md:84–90).

**Lemma 93.2 [derived; E5-L; any; category-free; → R4] (normal limit expectation). Assume** (i) increasing $\mathcal{N}_k \subset \mathcal{M}_k$ with $\mathcal{N}_k \subset \mathcal{N}$, $\mathcal{M}_k \subset \mathcal{M}$, bounded exhaustions of both $\mathcal{N}$ and $\mathcal{M}$; (ii) normal conditional expectations $E_k : \mathcal{M}_k \to \mathcal{N}_k$, *not* assumed compatible; (iii) a normal state $\psi$ on $\mathcal{M}$, faithful on $\mathcal{N}$, with $\psi \circ E_k = \psi|_{\mathcal{M}_k}$ for all $k$; (iv) $b := \liminf_k K(E_k) < \infty$. **Prove** there is a unique $\psi$-preserving normal conditional expectation $E : \mathcal{M} \to \mathcal{N}$; it satisfies $E(x) \geq b^{-1}x$ on $\mathcal{M}_+$ (so $q(\mathcal{N} \subset \mathcal{M}) \leq b$), and $E$ and $\psi$ are faithful. (i)–(iii) alone already construct $E$. ⟨1⟩1.

For $n \in \mathcal{N}_r$, $k \geq r$, $a \in \mathcal{M}_k$, BY bimodularity and (iii):

$$\psi\big(n^* E_k(a)\big) = \psi\big(E_k(n^* a)\big) = \psi(n^* a). \tag{9}$$

$\langle 1\rangle 2$. For bounded $a_k \to x$ strong$^*$, every ultraweak cluster point $y \in \mathcal{N}$ of $E_k(a_k)$ obeys

$$\varphi(n^* y) = \psi(n^* x) \qquad \forall n \in \mathcal{N}, \tag{10}$$

BY (9), normality of $\psi$, compactness of the ball and bounded exhaustion of $\mathcal{N}$. If $y, z$ both satisfy (10), take $n = y - z$: $\varphi((y-z)^*(y-z)) = 0$, so $y = z$ BY faithfulness of $\varphi$. Define $E(x) := y$; it is independent of the approximants, and every bounded approximating sequence has $E_k(a_k) \to E(x)$ ultraweakly. $\langle 1\rangle 3$. $E$ is linear, unital, completely positive and fixes $\mathcal{N}$: approximate the entries of $X^{1/2}$ for positive $X \in M_m(\mathcal{M})$ and take positive ultraweak limits (fixed matrix size). $\langle 1\rangle 4$. $\psi \circ E = \psi$ ($n = 1$ in (10)). *Normality:* if $x_\alpha \uparrow x$, then $y := \sup_\alpha E(x_\alpha) \leq E(x)$ and $\varphi(E(x) - y) = \psi(x) - \lim_\alpha \psi(x_\alpha) = 0$, so $y = E(x)$. This is the step finite-stage normality cannot supply (Example 93.5). $\langle 1\rangle 5$. Bimodularity passes from $\mathcal{N}_r$ to $\mathcal{N}$ BY $\langle 1\rangle 2$ and normality. $\langle 1\rangle 6$. Uniqueness: any $\psi$-preserving normal expectation satisfies (10); apply $\langle 1\rangle 2$. $\langle 1\rangle 7$. Bounded positive approximants along a subsequence realising $b$ satisfy the hypotheses of Lemma 93.1 BY $\langle 1\rangle 2$; hence $E(x) \geq b^{-1} x$. $\langle 1\rangle 8$. $E(x) = 0$, $x \geq 0$ forces $x = 0$ BY $\langle 1\rangle 7$; $\psi(x) = 0$ forces $E(x) = 0$. QED (full proof NOTE.md:116–187; every step graded valid, REVIEW.md:67–86). *Remarks.* Neither commutants (KLM:507–556) nor complete positivity of $E_k - \lambda\,\mathrm{id}$ is used; type-I or nonfactor approximants cause no difficulty. Increasing $\mathcal{M}_k$ is not used in $\langle 1\rangle 1$ (REVIEW.md:69).

**Corollary 93.3 [derived; E5-J; any; category-free].** Under Lemma 93.2, in the GNS representation of $\psi$ with vector $\Omega_\psi$ (not the vacuum), the projections onto $\overline{\mathcal{M}_k \Omega_\psi}$ converge strongly to 1, the Jones projections $e_k$ onto $\overline{\mathcal{N}_k \Omega_\psi}$ converge strongly to $e$ onto $\overline{\mathcal{N} \Omega_\psi}$, and $E_k(x_k) \to E(x)$ strong$^*$ for bounded $x_k \to x$ strong$^*$. Jones convergence is a *consequence*, not a substitute for normality or exhaustion (NOTE.md:195–227; REVIEW.md:88–93).

**Proposition 93.4 [derived; E5-N; any; category-free] (compatible case, exact criterion). Assume** (i), (ii) of Lemma 93.2 with $E_l|_{\mathcal{M}_k} = E_k$ ($l \geq k$), $E_0$ the induced map on $\bigcup_k \mathcal{M}_k$, and a faithful normal state $\varphi$ on $\mathcal{N}$ (exists on a separable Hilbert space, REVIEW.md:100–101). **Prove** $E_0$ extends to a normal expectation $\mathcal{M} \to \mathcal{N}$ iff $\varphi \circ E_0$ extends to a normal positive functional on $\mathcal{M}$; and if $b < \infty$, then $K(E) = \sup_k K(E_k) = \lim_k K(E_k) = b$. $\langle 1\rangle 1$. Necessity: $\psi := \varphi \circ E$. $\langle 1\rangle 2$. Sufficiency: the extension $\psi$ restricts to $\varphi$ on $\mathcal{N}$ and is preserved by every $E_k$; apply Lemma 93.2; constant sequences show $E$ extends $E_0$. $\langle 1\rangle 3$. Restriction gives $K(E_k) \uparrow$ and $\sup_k K(E_k) \leq K(E)$; Lemma 93.2 gives $\leq b$. QED (NOTE.md:242–252). *Selection remark.* If only minimal stage constants $q_k = q(\mathcal{N}_k \subset \mathcal{M}_k)$ are known, a selection certificate (O) is needed: expectations with $K(E_{k_j}) \leq q_{k_j} + \delta_j$, $\delta_j \to 0$, along a subsequence realising $\liminf q_k$ (NOTE.md:260–269). The brief's literal minimal-constant form without (O) is *conjectural*: neither proved nor refuted (REVIEW.md:16–17, :212).

## 94.4 Sharpness of the Hypotheses

**Example 93.5 [derived, negative; E5-X1, verified].** Let $D_0, D_1 \subset \{0,1\}^{\mathbb{N}}$ be the eventually-0 and eventually-1 sequences, $\mathcal{H} = \ell^2(D_0 \sqcup D_1)$, $\mathcal{M}_k$ the functions of the first $k$ bits on each copy independently, $\mathcal{N}_k$ the same function on both copies, $E_k(f_0, f_1) = \big(\frac{f_0+f_1}{2}, \frac{f_0+f_1}{2}\big)$. The $E_k$ are compatible and normal with $K(E_k) = q_k = 2$, and both exhaust $\mathcal{N} = \mathcal{M} = \ell^\infty(D_0 \sqcup D_1)$ boundedly; yet $E_0(P_0) = \frac{1}{2} \neq P_0$ for $P_0 = (1,0)$, so *no* expectation extends $E_0$. The failure is of normality: $(0, f_k) \downarrow 0$ while $E_k(0, f_k) \downarrow p_d/2 \neq 0$ (NOTE.md:273–317; REVIEW.md:105–114). It refutes extension of the maps, not index semicontinuity (the limit has $q = 1 \leq 2$).

**Example 93.6 [derived, negative; E5-X2].** $\mathcal{N} = \mathbb{C}1 \subset \mathcal{M} = M_m(\mathbb{C})$, $m \geq 2$, on $L^2(\mathcal{M}, \tau)$, with $\mathcal{N}_k = \mathcal{M}_k = \mathbb{C}1$: the stage Jones projections *equal* $e_{\mathcal{N}}$ and the stage cost is 1, but $q(\mathcal{N} \subset \mathcal{M}) = m$. Jones convergence without ambient exhaustion underestimates the index (NOTE.md:319–334). The review notes X2 drops a hypothesis the brief already contained; no counterexample to the index inequality under full exhaustion without a normal state is known; the review sketches why none exists in the abelian case, and its noncommutative attempts failed (REVIEW.md:118–123, :176–184; open problem NOTE.md:613–614).

**Example 93.7 [derived; added by the review] (the bound is attained).** For the hyperfinite factor, $R \otimes 1 \subset R \otimes M_n$ with $\mathcal{N}_k = M_{2^k} \otimes 1 \subset \mathcal{M}_k = M_{2^k} \otimes M_n$, $E_k = \mathrm{id} \otimes \mathrm{tr}$ and $\psi$ the trace: the hypotheses of Lemma 93.2 and Proposition 93.4 hold and $K(E_k) = n \min(2^k, n) \uparrow n^2 = [R \otimes M_n : R \otimes 1]$ (REVIEW.md:168–171; `checks/review_raw_output.txt`:6). Lemma 93.2 is not vacuous for genuine subfactors.

## 94.5 The Two-Sided Fidelity Theorem at its Lowest Layer

Let $\mathcal{N} = \mathcal{A}(I_1) \vee \mathcal{A}(I_3) \subset \mathcal{M} = (\mathcal{A}(I_2) \vee \mathcal{A}(I_4))'$ be the four-interval inclusion of (al).

**Theorem 93.8 [derived implication; E5-F with layer lowered by REVIEW §4; chiral; L0+L4 / L1+L4; R4 given R3]. Assume** (F1) $\mathcal{A}$ is a completely rational conformal net on a separable Hilbert space (this is R3, not supplied here); (F2$_0$) $\mathcal{C}$ is unitary fusion and $\Phi : \mathcal{C} \to \mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$ is a fully faithful unitary tensor functor; (F3$''$) the hypotheses of Lemma 93.2 (or of Proposition 93.4) hold for the four-interval inclusion with $b \leq D$. **Prove** (a) $\mu_{\mathcal{A}} = D$, $\Phi$ is a unitary tensor equivalence onto $\mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$, and $\mathcal{A}$ has no further irreducible sectors; (b) if $\mathcal{C}$ is braided and $\Phi$ braided (L1), $\Phi$ is a braided equivalence and $\mathcal{C}$ is *modular*. $\langle 1\rangle 1$. $\mu_{\mathcal{A}} \leq K(E) \leq b \leq D$, BY Lemma 93.2 and KLM:430–443. $\langle 1\rangle 2$. $\mu_{\mathcal{A}} = \sum_\rho d(\rho)^2$ over all irreducible finite sectors, BY KLM:1719–1729 (c21), whose modular PCT hypothesis is automatic for conformal nets (KLM:782–792); every sector is a direct sum of finite-dimensional ones (KLM:1899–1900). $\langle 1\rangle 3$. The image of $\Phi$ contributes exactly $D$: simples stay simple and pairwise inequivalent (full faithfulness), and $d(\Phi(i)) = \mathrm{FPdim}(i) = d_i$ *automatically*, since statistical dimension is the Frobenius–Perron dimension (B:620–626; REVIEW.md:197–199). No dimension-preservation hypothesis is needed. $\langle 1\rangle 4$. An irreducible sector outside the image adds $d(\rho)^2 > 0$, contradicting $\langle 1\rangle 1$–$\langle 1\rangle 3$; $\Phi$ is essentially surjective; the inverse is chosen unitary (NOTE.md:376–377). No braiding is used (REVIEW.md:200–201). $\langle 1\rangle 5$. At L1 a braided equivalence results; $\mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$ is nondegenerate BY KLM:1861–1863 (c23), hence so is $\mathcal{C}$. Twists and ribbon traces for the canonical positive spherical structure are preserved (B:617–645; REVIEW.md:128–131). QED. *Scope.* For a non-braidable $\mathcal{C}$ (F2$_0$) is unsatisfiable (the image would inherit a braiding), so 93.8 is vacuous there (REVIEW.md:165–167). $T$, characters and a central-charge lift remain L3 data (CONVENTIONS (u)). The chiral tag must be read with E2's caveat: a parity-paired KS limit carries both stress families, so a chiral target needs parity breaking or is the left chiral subnet of a full limit (INTEGRATION.md:97–100).

## 94.6 The Full-Setting Corollary and a False Reading

**Corollary 93.9 [derived; E5-D; full, named doubled chiral net; L2+L4; R4 for that net]. Assume** Theorem 93.8 at L2 ($\mathcal{C}$ modular). **Prove** KLM's net $\mathcal{A} \otimes \mathcal{A}^{\mathrm{o}}$ has $\mathrm{DHR}_{\mathrm{fin}}(\mathcal{A} \otimes \mathcal{A}^{\mathrm{o}}) \simeq \mathcal{C} \boxtimes \mathcal{C}^{\mathrm{rev}} \simeq Z(\mathcal{C})$, of dimension $D^2$. $\langle 1\rangle 1$. Irreducible sectors are exactly $\rho_i \otimes \rho_j^{\mathrm{o}}$ (KLM:969–972, :1518–1559, l17, c18); intertwiners and exchanges factor, the right copy with reversed braiding $\varepsilon^{\mathrm{o}} = j(\varepsilon)^*$ (KLM:1089–1092; B:643–645). $\langle 1\rangle 2$. $Z(\mathcal{C}) \simeq \mathcal{C} \boxtimes \mathcal{C}^{\mathrm{rev}}$ iff modular (B:660–664); $\sum_{ij} (d_i d_j)^2 = D^2$. QED (NOTE.md:407–419). *Which net.* KLM's c18 concerns their *one-dimensional* net $\mathcal{A} \otimes \mathcal{A}^{\mathrm{o}}$ (time-zero picture); the corresponding statement for the 2D light-ray product $\mathcal{A} \otimes \bar{\mathcal{A}}$ on $\mathbb{R} \times \mathbb{R}$ is

not locally sourced (REVIEW.md:138–140): a Law 1 gap, recorded, not filled.

**Proposition 93.10 [derived, negative].** For modular $\mathcal{C} \neq \mathrm{Vec}$ realised as in 93.8, the canonical (diagonal) Longo–Rehren extension $\mathcal{B}_2 \supset \mathcal{A} \otimes \mathcal{A}^{\mathrm{o}}$ (KLM:1086–1098) has *trivial* finite DHR category: by KLM:1578–1587 (p18) and :1861–1879 (c23, $\mu_{\mathcal{B}} = 1$) it has no nontrivial bounded-localised finite-index endomorphism, and double-cone sectors restrict to time-zero interval sectors by cofinality (NOTE.md:425–440). Since $(a, \mathbf{1})$ is a nonunit simple of $Z(\mathcal{C})$, *"*DHR *of the full diagonal extension is $Z(\mathcal{C})$" is false as stated* (NOTE.md:441–444; REVIEW.md:20–21). Correct readings: $Z(\mathcal{C})$ is the sector category of the doubled chiral net, or the centre of the diagonal theory's defect category, which is $\mathcal{C}$ itself (FRS:641–646; NOTE.md:446–450); a general full extension declares its own bimodule category (B:3323–3337).

## 94.7 What a Lattice Construction Must Supply

**Lemma 93.11 [conditional; E5-EP; chiral; L2+L4; →(F2), → R4 given R3].** Finite charged endpoint operators, junction matrices for bases of $\mathrm{Hom}_{\mathcal{C}}(a, b)$, tensorators and conjugate pairs, with bounded product/adjoint limits that are unital, multiplicative, locally normal, localised and transportable; identity errors $\to 0$ (both $U_k^* U_k \to 1$ and $U_k U_k^* \to 1$); limiting conjugate pairs proved *standard*; and a fullness certificate (bounded coefficient approximation of every limiting intertwiner), imply the functor of (F2) (NOTE.md:458–492). The review notes the fullness certificate nearly restates fullness (REVIEW.md:151–153).

**Lemma 93.12 [conditional; E5-AN; chiral; L2+L4; →(F3″)].** Represented finite inclusions $\mathcal{N}_k \subset \mathcal{M}_k$ with $\mathcal{N}_k \subset \mathcal{N}$, $\mathcal{M}_k \subset \mathcal{M}$, bounded exhaustions of *both*, matrix expectations with $E_k(x) \geq b_k^{-1} x$ and $\liminf b_k \leq D$, and a common normal non-vacuum state (or criterion 93.4), give (F3″) and Corollary 93.3 (NOTE.md:494–522). The literal inclusions must come from the inductive-limit embedding: an extra obligation (REVIEW.md:154–156). No commutant and scaling limit may be interchanged.

**Lemma 93.13 [conditional; E5-ALL; chiral; L2+L4; R3 → R4].** If one lattice/OAR system has an identified completely rational conformal limit net and 93.11, 93.12 hold in that representation, Theorem 93.8 applies (NOTE.md:524–532). It derives no R1–R3 and no convergence to $\mathcal{B}_2$.

**Non-vacuity caveat.** No instance of Lemma 93.2's hypotheses is known for a type $\mathrm{III}_1$ four-interval inclusion. That inclusion is irreducible with finite index, so its expectation is unique and $K(E) = \mu_{\mathcal{A}}$ (REVIEW.md:56–58); certifying costs must therefore have $\liminf K(E_k) = \mu_{\mathcal{A}}$ *exactly* (and lim, in the compatible case 93.4): a sharp lattice demand (REVIEW.md:173–175). Without an instance, 93.8 is a correct implication with no known application (INTEGRATION.md:121–123).

**Next lemma [proposal; any; category-free, then L0+L4] (annular approximant lemma).** For a sourced completely rational net, construct increasing finite-dimensional $\mathcal{N}_k \subset \mathcal{M}_k$ inside the four-interval inclusion with bounded exhaustion of both, a non-vacuum normal state preserved by the $E_k$, and $K(E_k) \to \mu_{\mathcal{A}}$; equivalently, show the minimal-cost expectations of some exhausting family preserve one normal state (INTEGRATION.md:114–123).

**Effect on CA-80.** This closes route (ii) of CA-80's "The gap, explicitly" (CA-80:198–202), with a common normal state replacing compatibility and without complete positivity; routes (i) and (iii) are unnecessary. Theorem 80.9 is now conditional on (F1) only, given the 93.2 certificate. Proposition 80.8 and Lemma 80.10 stay in force.

## 94.8 What This Does Not Claim

No lattice family, annular approximant, common normal state, endpoint functor or completely rational limit net is constructed for any $\mathcal{C}$; (F1) is R3 and assumed. The minimal-constant variant

without (O), index semicontinuity without a normal state, and non-vacuity for conformal-net inclusions are not established (REVIEW.md:211–215). Nothing identifies $T$, characters, $c$ or OPE data, gives R5, or shows lattice convergence to the Longo–Rehren full extension.

| Instance | Calibration datum for the general statements |
|---|---|
| $\mathcal{C} = \mathrm{Vec}$ | $D = 1$: 93.8 gives $\mu_{\mathcal{A}} = 1$, a holomorphic net; consistent, degenerate (REVIEW.md:160–162). |
| Non-braidable $\mathcal{C}$ (e.g. Haagerup) | ($\mathrm{F2}_0$) unsatisfiable; 93.8 vacuous; full setting via $Z(\mathcal{C})$ instead. |
| Any modular $\mathcal{C} \neq \mathrm{Vec}$ | Doubled net $Z(\mathcal{C})$, $D^2$; diagonal LR extension: Vec (93.9, 93.10). |
| Two-copy cylinders (X1) | Stage cost 2, compatible, full exhaustion; no normal extension (93.5). |
| $\mathbb{C}1 \subset M_m$ (X2) | Equal Jones projections, stage cost 1, true cost $m$ (93.6); commutant $n$ vs $n^2$ is CA-80's gap. |
| $R \otimes 1 \subset R \otimes M_n$ | Stage costs $n \min(2^k, n) \uparrow n^2$: bound attained (93.7). |

# 95 Number-Changing Collision Dynamics on the Pointed Carrier: a Tuned Continuum Theorem and Three Obstructions

## 95.1 Status, Sources, and Dependencies

**Status: derived (positive and negative); each statement carries its own label.** NOTE, REVIEW, INTEGRATION are the files in `explorations/2026-10-22-astra-r2/E6-number-changing-continuum/`; "D1 NOTE" etc. those in `explorations/2026-10-22-astra/D1-carrier-theory/`. The review grades the brief's E6 (`briefs/E6-number-changing-continuum.md`:69–97) **COUNTEREXAMPLE (verified)**, with a **proof-complete replacement** and no invalid step (REVIEW:9–29): E6(a), (b) fail by N1 (94.1); E6(c), as unrestricted surjectivity, fails by N3 (94.11), which does not settle (c) without $(G_r)$ or for a local target class (REVIEW:21–24); the positive theorem is for a *different, $\varepsilon$-dependent* family (REVIEW:26–29). Global $H^2$ regularity (94.8) and general range $r$ (94.13) are *conditional*; 94.16 is *conjectural*. Finite evidence (`checks/`: `check_e6.py`, five groups GREEN, three mutations RED, `raw_output.txt`:3–14; `review_N1_N3.py`; `review_tuned_convergence.py`) supports finite algebra and counting steps only, not the theorem (NOTE:776–778).

**Layer tags (CONVENTIONS (aq)) of the main results 94.6, 94.7, 94.10.** Setting *any* (open interval $(0,1)$, ordered configurations, nonrelativistic kinetic energy; no chiral or full target selected). Data $\mathsf{L0}$ plus declared ($b_\alpha, v_\beta, \lambda_\alpha, \kappa_\beta, \mu$, contact tuning); $\mathsf{L1}$–$\mathsf{L4}$ unused. Rung $\mathsf{R0}^c$ (carrier, Theorem 77.8) plus an *off-ladder $z$=2 resolvent limit*; **not** $\mathsf{R1}$, which in CA-88 is a Wilson-triangle limit state (Definition 88.3), none being constructed (NOTE:807–811; REVIEW:218–223). *Per statement:* all of 94.1–94.15 have setting any (open interval) and data $\mathsf{L0}$ plus the declared data they name (94.3 uses only a finite-dimensional fibre). 94.6–94.8, 94.10 reach the off-ladder limit; 94.3–94.5 and 94.13 feed it ($\to$ that limit); 94.1, 94.9, 94.11, 94.14 are negative at this layer; 94.12 is $\mathsf{L0}$ naturality, not $\mathsf{R5}$; 94.16 is conjectural at the same tags.

**Source gap (Law 1).** No local source treats Mosco convergence, extension classification, boundary triples, point interactions or interior-boundary-condition Hamiltonians (NOTE:58–66; REVIEW:58–61); all analytic lemmas are proved inline. Prior art on this operator class may exist; a novelty or comparison claim needs a source first (REVIEW:239–242). Dependencies: CA-77 (77.4, 77.8, Conjecture with Caveat), CA-88 (88.3, D1-C entry), D1-T7; CONVENTIONS (a), (b), (r), (t), (u), (ak), (al), (am), (aq), and the shard-local conventions of §95.2 (proposed for CONVENTIONS).

## 95.2 Setting and Shard-Local Conventions

$\mathcal{C}$ is unitary fusion; $(O,\eta)$ is pointed, $O \cong \mathbf{1} \oplus X$ (CONVENTIONS (am)), not necessarily connected. Fibres $E_{c,n} = \mathrm{Hom}(c, X^{\otimes n})$ carry the isometry inner product (Penneys:528–538); $\mathcal{F} = \widehat{\bigoplus}_{c,n} L^2(\Delta_n; E_{c,n})$, no $n!$; $P_n$ is the occupation projection; $K_\varepsilon$ are the box maps of Theorem 77.8 on the uniform mesh $\varepsilon = 1/L$. $\pi_{00}, \pi_{X0}, \pi_{0X}, \pi_{XX}$ are the coisometries of $O^{\otimes 2}$ onto its summands, $P_{XX} = \pi_{XX}^\dagger \pi_{XX}$. Energies: $H = \sum_{j=1}^{L-1} k_j$, $k = -h^{(\mathrm{ak})}$ as in (al). $T_{n,i}$ is the collision trace on $\{x_i = x_{i+1}\}$ with collapsed measure $dy = dS/\sqrt{2}$; $D_{n,i} = \partial_i - \partial_{i+1}$ is its outward flux. $B_{b,n,i} : E_{c,n} \to E_{c,n+1}$ inserts $b \in \mathrm{Hom}(X, X^{\otimes 2})$ at position $i$; $V_{v,n,i} : E_{c,n} \to E_{c,n+2}$ inserts $v \in \mathrm{Hom}(\mathbf{1}, X^{\otimes 2})$ in gap $i$, on $\Gamma_{n,i} = \{y_1 < \cdots < y_i < t < y_{i+1} < \dots\}$. Coherent unitors/associators are implicit; no braiding is used. "Extension of the minimal Laplacian" means containing $A_{\min} = \bigoplus(-\Delta + \mu n)$ on finite-sector $C_c^\infty$ vectors, vacuum included (NOTE:18–56).

## 95.3 Fixed Densities Fail, and the Corrected Scaling

**Proposition 94.1 [negative; E6-N1] (NOTE:74–121; REVIEW:69–83). Assume** any $\mathcal{C}$, any $X \neq 0$, the pure-hop density $k = -\pi_{X0}^\dagger \pi_{0X} - \pi_{0X}^\dagger \pi_{X0}$, $H_\varepsilon = \varepsilon^{-2} \sum_j k_j$, real scalars $s_\varepsilon$. **Prove** no self-adjoint $H$ on $\mathcal{F}$ has $K_\varepsilon (H_\varepsilon - s_\varepsilon - i)^{-1} K_\varepsilon^* \to (H-i)^{-1}$ strongly.
$\langle 1\rangle 1$. $\Omega$ has energy 0 at every scale: projected resolvent $a_\varepsilon \Omega$, $a_\varepsilon = (-s_\varepsilon - i)^{-1}$. $\langle 1\rangle 2$. A resolvent is injective, so $a_\varepsilon \to a \neq 0$ and $s_\varepsilon$ tends to a finite real number. $\langle 1\rangle 3$. For unit $e \in E_{c,1}$, $0 \neq f \in C_c^\infty(0,1)$, $u_\varepsilon(j) = \sqrt{\varepsilon} f((j - \frac{1}{2})\varepsilon) e$: $K_\varepsilon u_\varepsilon \to fe$, $H_\varepsilon u_\varepsilon = -2\varepsilon^{-2} u_\varepsilon + w_\varepsilon$, $\sup_\varepsilon \|w_\varepsilon\| \leq \sup |f''|$. $\langle 1\rangle 4$. With $d_\varepsilon = -2\varepsilon^{-2} - s_\varepsilon - i$, $R_\varepsilon u_\varepsilon = d_\varepsilon^{-1}(u_\varepsilon - R_\varepsilon w_\varepsilon) \to 0$ ($|d_\varepsilon| \to \infty$, $\|R_\varepsilon\| \leq 1$). $\langle 1\rangle 5$. Hence $(H-i)^{-1}(fe) = 0$, contradicting injectivity. QED. The real-resolvent wording fails too ($|1 - (\lambda + i)t|^2 \geq (1+\lambda^2)^{-1}$; NOTE:123–131). *Reviewer additions:* N1 survives scaling the diagonal part by $\varepsilon^{-2}$ too, the mismatch being a number-proportional chemical potential, never a scalar (REVIEW:79–80); a generalised (pseudo-)resolvent limit exists only for an operator on the vacuum line, so N1 refutes exactly the claimed self-adjoint $H_h$ on $\mathcal{F}$ (REVIEW:81–83). $k = 0$ refutes a mandatory nonzero Laplacian in E6(b) (NOTE:121).

**The tuned family (D1-C; D1 NOTE:35–41).** Fix finitely many $b_\alpha \in \mathrm{Hom}(X, X^{\otimes 2})$, $v_\beta \in \mathrm{Hom}(\mathbf{1}, X^{\otimes 2})$, weights $\lambda_\alpha, \kappa_\beta \geq 0$, and $\mu > 0$;

$$H_\varepsilon = \sum_j \Big[ \varepsilon^{-2} D_h^\dagger D_h + \sum_\alpha \lambda_\alpha \varepsilon^{-1} D_{b_\alpha}^\dagger D_{b_\alpha} + \sum_\beta \kappa_\beta \varepsilon^{-1} D_{v_\beta}^\dagger D_{v_\beta} \Big]_j + \mu \hat{N},$$

$D_h = \pi_{X0} - \pi_{0X}$, $D_b = \pi_{XX} - \sqrt{\varepsilon}\, b\, \pi_{X0}$, $D_v = \pi_{XX} - \varepsilon\, v\, \pi_{00}$. $D_h^\dagger D_h$ is hard-core hopping plus the contact attraction $-2\varepsilon^{-2} n_j n_{j+1}$ (D1 REVIEW:78–87): CA-77's caveat (77:240–246), kept deliberately.

**Proposition 94.2 [derived; E6-P1, scale identification] (NOTE:150–173; REVIEW:85–88).** Expanding the squares,

$$\varepsilon^{-1} D_b^\dagger D_b = \varepsilon^{-1} P_{XX} - \varepsilon^{-1/2}(\pi_{XX}^\dagger b \pi_{X0} + \text{h.c.}) + \pi_{X0}^\dagger b^\dagger b \pi_{X0}, \quad \varepsilon^{-1} D_v^\dagger D_v = \varepsilon^{-1} P_{XX} - (\pi_{XX}^\dagger v \pi_{00} + \text{h.c.}) + \varepsilon\, \pi_{00}^\dagger v^\dagger v \pi_{00},$$

so the finite collision-form scalings are hop $\varepsilon^{-2}$, contact (upper trace-square) $\varepsilon^{-1}$, fusion cross term $\varepsilon^{-1/2}$, pair cross term $\varepsilon^0$. $\langle 1\rangle 1$. With $n$ spectators a $1 \to 2$ birth has amplitudes $\varepsilon^{(n+1)/2}, \varepsilon^{(n+2)/2}$ on $\sim \varepsilon^{-(n+1)}$ grid points: a cross coefficient $g$ contributes $g\sqrt{\varepsilon}$. $\langle 1\rangle 2$. A $0 \to 2$ birth contributes $g$. $\langle 1\rangle 3$. Hence $g_b = \varepsilon^{-1/2}$, $g_v = 1$, by the D1-T7 Riemann sums (D1 NOTE:498–508). QED. The brief's uniform $\varepsilon^{-1}$ on fixed $b, v$ tiles gives cross sizes $\varepsilon^{-1/2}, \varepsilon^{-1}$, inconsistent even for D1-C; in a mutation run the vacuum energy grows as $223, 933, 3813$ at $L = 12, 24, 48$ (`checks/review_tuned_convergence_output.txt`:6).

## 95.4 The Closed Collision Form

On its maximal finite-energy domain (NOTE:175–188),

$$q(\psi) = \sum_{c,n} \int_{\Delta_n} \Big( \sum_i \|\partial_i \psi_n\|^2 + \mu n \|\psi_n\|^2 \Big) + \sum_\alpha \lambda_\alpha \sum_{n \geq 1, i} \|T_{n+1,i} \psi_{n+1} - B_{b_\alpha, n, i} \psi_n\|^2 + \sum_\beta \kappa_\beta \sum_{n \geq 0, i} \|T_{n+2,i+1} \psi_{n+2} - V_{v_\beta, n, i} \psi_n$$

**Lemma 94.3 [derived; E6-L1] (NOTE:192–229; REVIEW:90–93).** For a finite-dimensional fibre, functions smooth on $\overline{\Delta_n}$ are dense in $H^1(\Delta_n)$, and face traces are continuous $H^1 \to L^2$: $\|f|_F\|^2 \leq C_n(d^{-1}\|f\|^2 + d\|\nabla f\|^2)$, $0 < d \leq \frac{1}{2}$. Proof: dilation about an interior point, mollification, integration along the inward pyramid (Jacobian $\propto (1-t)^{n-1}$). Constants depend on $n$; used sectorwise only.

**Lemma 94.4 [derived; E6-L2] (NOTE:231–289; REVIEW:95–103). Assume** the tuned data, $\mu > 0$. **Prove** $q$ is densely defined, nonnegative and closed, finite-particle vectors smooth on closed chambers form a form core, and a unique nonnegative self-adjoint $H$ has $q(\varphi,\psi) = \langle \varphi, H\psi\rangle$. ⟨1⟩1. $V_0 = D(q_0)$ is complete. ⟨1⟩2. $q = q_0 + \|T\psi - S\psi\|_G^2$ with $\|S\psi\|_G^2 \le B\sum_n n\|\psi_n\|^2 + V\|\psi\|^2$, $B = \sum \lambda_\alpha \|b_\alpha\|^2$, $V = \sum \kappa_\beta v_\beta^\dagger v_\beta$ (D1 NOTE:534–543; $\|\mathrm{id}\otimes b\otimes \mathrm{id}\| \le \|b\|$ as $\otimes$ is a linear †-functor, Penneys:319, REVIEW:63–65); so $S$ is bounded on $V_0$ since $\mu > 0$. ⟨1⟩3. $T : V_0 \to G$ is closed by sectorwise trace continuity (94.3). ⟨1⟩4. The form norm is equivalent to $\|\psi\|_0^2 + \|T\psi\|_G^2$; $q$ is closed. ⟨1⟩5. Truncation at particle number $M$ converges in this norm, keeping the lower sources into sectors $M{+}1, M{+}2$: the core. ⟨1⟩6–⟨1⟩7. The variational resolvent $R_z$ gives $D(H) = \mathrm{Ran}\, R_z$; $w = R_z(g + zw)$ gives $H^* = H$. QED. $\mu > 0$ is needed (REVIEW:103).

**Lemma 94.5 [derived; E6-L3, collision-trace Mosco lower bound] (NOTE:293–398; REVIEW:105–127). Assume** $K_\varepsilon u_\varepsilon \rightharpoonup \psi$, $\sup_\varepsilon \langle u_\varepsilon, H_\varepsilon u_\varepsilon\rangle < \infty$. **Prove** $q(\psi) \le \liminf_\varepsilon \langle u_\varepsilon, H_\varepsilon u_\varepsilon\rangle$. ⟨1⟩1. With $f_\varepsilon = \varepsilon^{-n/2} u_\varepsilon$ the hopping energy is exactly $E_{\varepsilon,n} = \varepsilon^{n-2}\sum_{\text{edges}} \|f_\varepsilon(j') - f_\varepsilon(j)\|^2$; fibres move by unitors only. ⟨1⟩2. Summation by parts on $\Omega_d \Subset \Delta_n$ gives $\psi_n \in H^1$, $\int \|\nabla\psi_n\|^2 \le \liminf E_{\varepsilon,n}$. ⟨1⟩3. On a separated patch $U$ of a collision face, the normal collar mean $M_{\varepsilon,h}$ over $m = \lfloor h/\varepsilon\rfloor$ cells obeys $\|A_\varepsilon - M_{\varepsilon,h}\|_{L^2(U)}^2 \le hE_{\varepsilon,n}$ (telescoping gives the sharper $(m/2)\sum\|\Delta_s\|^2$, REVIEW:112–113) and $\|M_{\varepsilon,h}\|^2 \le (m\varepsilon)^{-1}\|K_\varepsilon u_{\varepsilon,n}\|^2$; $M_{\varepsilon,h} \rightharpoonup A_h\psi$ and $\|T\psi - A_h\psi\|_{L^2(U)}^2 \le (h/3)\|\partial_{i+1}\psi\|_{\text{collar}}^2$, so $A_\varepsilon \rightharpoonup T\psi$. ⟨1⟩4. On $U$ the discrete lower sources are exactly $B(K_\varepsilon u_{\varepsilon,n-1})$ and the pulled-back $V(K_\varepsilon u_{\varepsilon,n-2})$, weakly convergent; the face weight is $\varepsilon^{-1}\varepsilon^n = \varepsilon^{n-1}$. ⟨1⟩5. Weak lower semicontinuity on finite collections of terms, each counted once, then monotone exhaustion of $\Omega_d$, patches, sectors and reactions. QED. No constant is $N$-uniform: each depends on a finite collection $(n, d, U, h)$, and the supremum is taken after the liminf (REVIEW:126–127).

## 95.5 The Tuned Continuum Theorem

**Theorem 94.6 [derived; E6-T1] (NOTE:402–446; REVIEW:129–137). Assume** any $\mathcal{C}$, any $X$, the tuned family. **Prove** $K_\varepsilon(H_\varepsilon + z)^{-1}K_\varepsilon^* f \to (H+z)^{-1}f$ strongly for all $z > 0$, $f \in \mathcal{F}$, along all $\varepsilon = 1/L$ (REVIEW:205–206), with $H$ from 94.4: no subsequence, scalar subtraction, imported trace/core hypothesis or $N$-uniform estimate.
⟨1⟩1. Smooth finite-particle vectors have strongly convergent sampling sequences with energies $\to q$ (D1-T7, D1 NOTE:483–516; re-verified, REVIEW:130–132). ⟨1⟩2. The core of 94.4 and a diagonal choice of mesh thresholds give every $\psi \in D(q)$ a recovery sequence. ⟨1⟩3. The minimiser $u_\varepsilon$ of $F_\varepsilon(u) = \langle u, H_\varepsilon u\rangle + z\|u\|^2 - 2\,\mathrm{Re}\langle K_\varepsilon^* f, u\rangle$ has $\|u_\varepsilon\| \le \|f\|/z$, energy $\le \|f\|^2/z$. ⟨1⟩4. By 94.5 and ⟨1⟩2 every weak cluster point minimises $F(u) = q(u) + z\|u\|^2 - 2\,\mathrm{Re}\langle f, u\rangle$, whose unique minimiser is $(H+z)^{-1}f$. ⟨1⟩5. For a recovery sequence $v_\varepsilon$, $z\|u_\varepsilon - v_\varepsilon\|^2 \le F_\varepsilon(v_\varepsilon) - F_\varepsilon(u_\varepsilon) \to 0$. QED. $K_\varepsilon K_\varepsilon^* \to I$ is not used; $\langle K_\varepsilon^* f, v\rangle = \langle f, K_\varepsilon v\rangle$ suffices (REVIEW:136–137). This closes D1 INTEGRATION (e)2–(e)3, the closedness/core hypotheses and the liminf gap (D1 INTEGRATION:128–130).

**Corollary 94.7 [derived; number-change criterion] (NOTE:448–461; REVIEW:138–140).** With $\beta = \sum_\alpha \lambda_\alpha b_\alpha$, $\nu = \sum_\beta \kappa_\beta v_\beta$: $H$ fails to commute with some $P_k$ iff $\beta \ne 0$ or $\nu \ne 0$. ⟨1⟩1. Only $\beta, \nu$ are off-diagonal in $n$ (94.10); if both vanish $q$ splits. ⟨1⟩2. Otherwise semisimplicity gives smooth tests with nonzero mixed form. ⟨1⟩3. Reduction by $P_k$ passes to the form, $\mathrm{Ran}(H+1)^{-1}$ being form-dense (94.11 ⟨2⟩1–⟨2⟩2). QED. Individual maps can cancel: equal weights on $b$ and $-b$ give $\beta = 0$.

**Proposition 94.8 [derived on regular vectors; global $H^2$ conditional; E6-P2] (NOTE:465–493; REVIEW:142–144).** $D(H)$ is the variational domain of 94.4. For vectors regular enough to integrate by parts, $(H\psi)_n = -\Delta\psi_n + \mu n\psi_n - \sum_\alpha \lambda_\alpha \sum_i B_{b_\alpha}^* R_{b_\alpha} - \sum_\beta \kappa_\beta \sum_i \int_{\mathrm{gap}_i} V_{v_\beta}^* R_{v_\beta}\, dt$, and on collision face $i$: $D_{n,i}\psi_n + \sum_\alpha \lambda_\alpha(T_{n,i}\psi_n - B_{b_\alpha}\psi_{n-1}) + \sum_\beta \kappa_\beta(T_{n,i}\psi_n - V_{v_\beta}\psi_{n-2}) = 0$; outer faces

are Neumann. $\partial_\nu = (\partial_i - \partial_{i+1})/\sqrt{2}$ and $dS = \sqrt{2}\,dy$ leave no residual $\sqrt{2}$. No global $H^2$ theorem is asserted (regularity at intersecting faces is open, NOTE:832).

**Proposition 94.9 [negative; E6-N2] (NOTE:495–518; REVIEW:146–148).** If an active reaction map is nonzero, $H \not\supseteq A_{\min}$. ⟨1⟩1. Semisimplicity gives a one-particle $C_c^\infty$ vector $\psi$ with $B\psi \neq 0$, or $\psi = \Omega$ for $v \neq 0$ (Penneys:195–219). ⟨1⟩2. Its upper sectors and traces vanish, so $q(\psi) = \langle \psi, A_{\min}\psi\rangle + \|S\psi\|^2 > \langle \psi, A_{\min}\psi\rangle$, contradicting $\psi \in D(H)$. QED. *Reviewer strengthening:* such $\psi$ is not in $D(H)$ at all, the mixed form being a boundary functional of the upper sector (REVIEW:147–148). Lower-sector *interior* values feed upper-sector collision conditions: the limit is of interior-boundary type on a stratified configuration space, not a Lagrangian datum of independent chamber Laplacians.

**Theorem 94.10 [derived; E6-T2, coefficient image] (NOTE:522–598; REVIEW:150–156).** For arbitrary finite reaction lists put $a_b = \sum \lambda_\alpha$, $Q_b = \sum \lambda_\alpha b_\alpha^\dagger b_\alpha \in \operatorname{End} X$, $a_v = \sum \kappa_\beta$, $\gamma = \sum \kappa_\beta v_\beta^\dagger v_\beta$, $a = a_b + a_v$. Then $q = q_0 + a\sum \|T\psi\|^2 - 2\operatorname{Re}\sum\langle T\psi_{n+1}, B_\beta\psi_n\rangle - 2\operatorname{Re}\sum\langle T\psi_{n+2}, V_\nu\psi_n\rangle + \sum\langle\psi_n, (Q_b)_i\psi_n\rangle + \gamma\|\psi\|^2$ depends only on $(a, \beta, \nu, Q_b, \gamma)$. With $M = \operatorname{Hom}(X, X^{\otimes 2})$, $W = \operatorname{Hom}(\mathbf{1}, X^{\otimes 2})$, a tuple is attained iff some split $a = a_b + a_v$, $a_b, a_v \geq 0$, satisfies

$$a_b > 0:\ \beta \in M,\ Q_b - \beta^\dagger\beta/a_b \in G_M; \qquad a_v > 0:\ \nu \in W,\ \gamma - \nu^\dagger\nu/a_v \in G_W; \qquad a_b = 0 \Rightarrow \beta = Q_b = 0;\ \ a_v = 0 \Rightarrow \nu$$

$G_M = \{\sum d_j^\dagger d_j : d_j \in M\}$. Writing $X = \bigoplus_c c^{m_c}$, $X^{\otimes 2} = \bigoplus_c c^{n_c}$, $G_M$ is the cone of PSD blocks of $\operatorname{End} X$ supported on charges with $n_c > 0$; $G_W = \mathbb{R}_{\geq 0}$ if $W \neq 0$, else $\{0\}$. ⟨1⟩1. Expand the squares; both reaction types range once over every collision face. ⟨1⟩2. Necessity: $Q_b - \beta^\dagger\beta/a_b = \sum \lambda_\alpha (b_\alpha - \beta/a_b)^\dagger (b_\alpha - \beta/a_b)$. ⟨1⟩3. Reconstruction: if $Q_b - \beta^\dagger\beta/a_b = \sum_{j\leq k} d_j^\dagger d_j$, the $2k$ maps $\beta/a_b \pm \sqrt{k/a_b}\,d_j$ of weight $a_b/2k$ have the right zeroth, first and second moments; likewise for $v$. ⟨1⟩4. Row-Gram decomposition gives the blocks (Penneys:195–237, 470–478, 528–530). QED. *Reviewer addition:* the identities hold exactly at finite $\varepsilon$ (REVIEW:156). The test is finitely many PSD and zero-block conditions, decidable under an effective exact encoding (NOTE:546–549). T2 classifies this family only, not arbitrary cutoff-dependent densities.

## 95.6 Obstructions, Naturality, and Range

**Proposition 94.11 [negative; E6-N3, grading obstruction] (NOTE:602–687; REVIEW:158–168). Assume** $(G_r)$: $\operatorname{Hom}(X^{\otimes p}, X^{\otimes q}) = 0$ for $0 \leq p \neq q \leq r$, and some $c$ occurs in both $X^{\otimes m}$, $X^{\otimes n}$, distinct $m, n \geq 2$. **Prove** (i) every strong projected-resolvent limit of range-$r$ categorical densities, under any scaling or scalar subtraction, commutes with every $P_k$; (ii) some charge-preserving self-adjoint $H_* \supseteq A_{\min}$ on $\mathcal{F}$ does not.
⟨1⟩1. Degree-$(q-p)$ tiles lie in $\operatorname{Hom}(Z_p, Z_q) = 0$ (Proposition 77.4, 77:93–117). ⟨1⟩2. Commutation with the bounded $P_k$ survives strong limits. ⟨1⟩3–⟨1⟩4. With face functionals $\ell_s(\psi_s) = \int \rho_s \langle e_s, T\psi_s\rangle$ $(s = m, n)$, $H^1$-bounded by 94.3, $q_* = \sum \|\nabla\psi\|^2 + |\ell_m - \ell_n|^2$ is closed. ⟨1⟩5. $H_* \supseteq A_{\min}$. ⟨1⟩6. Tests $u_m \pm u_n$ get added energies 0, 4: $q_*$ does not split, and reduction would pass to the form (⟨2⟩1: $\operatorname{Ran}(H_* + 1)^{-1}$ is form-dense; ⟨2⟩2: $P_k$ would split $q_*$). QED. *Reviewer additions:* N3 also holds for the reading "induced by a morphism, natural in $c$", since $e_m e_n^*$ is induced by $\iota_m \iota_n^* : X^{\otimes n} \to X^{\otimes m}$, which exists as $n > r$ (REVIEW:164–166); the Lagrangian graph remark (NOTE:645–656) is *formal*, an $L^2 \times L^2$ trace pairing not being a boundary triple on polytopes (REVIEW:167–168, :234–235). The countertarget is nonlocal along faces and couples faces of different dimension: N3 refutes unrestricted surjectivity only, not a local stratified target class (NOTE:682–687). It is the $\mathbb{Z}$-graded analogue of the parity statement 77.4(3); its instance is in the examples table only.

**Proposition 94.12 [derived; E6-P3, naturality at the nonrelativistic layer] (NOTE:691–711; REVIEW:170–171).** A unitary tensor equivalence transporting $(O, \eta, X)$, reactions and

couplings induces unitaries with $K'_\varepsilon W_\varepsilon = W K_\varepsilon$, $H'_\varepsilon W_\varepsilon = W_\varepsilon H_\varepsilon$, $H' = WHW^*$ (unitary †-functors preserve isometry inner products; tensorators are unitary). Not R5.

**Proposition 94.13 [derived estimate; general range $r$ conditional; E6-P4] (NOTE:721–742; REVIEW:173).** If an adjacent-pair face sample is replaced by one reached through $\le r$ allowed moves, edge multiplicity $\le C_{n,r}$, then $\|A_\varepsilon - A'_\varepsilon\|^2_{L^2(U)} \le C_{n,r}\,\varepsilon E_{\varepsilon,n} \to 0$. T1 extends to a positive range-$r$ family *provided* it controls nearest-neighbour graph energy, has only codimension-one reaction traces, bounded lower sources, and smooth recovery with a closed core; this is a conditional extension, not a claim about every range-$r$ density.

**Proposition 94.14 [negative; higher-codimension $H^1$ trace obstruction] (NOTE:744–761; REVIEW:175–176).** A tile creating three coincident particles addresses a codimension-two stratum. With normal radius $R$ and cutoff $\log(h/R)/\log(h/\delta)$, the normal gradient energy is at most a fixed multiple of $\int_\delta^h [R\log(h/\delta)]^{-2} R\,dR = 1/\log(h/\delta) \to 0$ while the stratum value stays fixed; no continuous $H^1$-to-triple-trace map exists. This defeats 94.5's trace hypothesis for such tiles, not renormalised constructions.

**Remark 94.15 [derived; where $\mathcal{C}$ enters] (NOTE:713–719; REVIEW:224–228).** The category enters through the fibres $E_{c,n}$, the admissible Hom spaces $M, W$, the coherent insertions (which may contain $F$-matrices in fusion-tree bases), the support cones of 94.10 (charges with $n_c > 0$) and the obstruction $(G_r)$. Every analytic estimate is *category-inert*: finite-dimensional norms, bounded maps and the exclusion grid only. No braiding, twist, modular data or central charge enters; L0 does not select $b, v, \mu$ or the contact tuning (D1 INTEGRATION (e)4 stands); no sector category is recovered. 94.11 sharpens D1 REVIEW R2 ("inert category", D1 REVIEW:92–98): fusion rules restrict the *attainable* range-$r$ dynamics. *Scope omission (REVIEW:208–211):* for non-connected $O$, degree-$\pm1$ tiles create a single particle from a vacancy; these bulk sources lie outside the D1 family and T1, and a fixed-density classification must include them.

## 95.7 Next Lemma: Contact Trichotomy

**Target 94.16 [conjectural; any (open interval); L0 + declared $t, c_\varepsilon$; R0$^c$ + off-ladder $z$=2] (INTEGRATION:104–117).** Let $h \in \mathrm{End}_{\mathcal{C}}(O\otimes O)$ be fixed, self-adjoint, with number-conserving block $t$·hop plus diagonal, scaled by $\varepsilon^{-2}$, with counterterm $\varepsilon^{-2}\epsilon_{\mathrm{bot}}\hat{N}$ ($\epsilon_{\mathrm{bot}}$ the one-particle band bottom), D1's outer-endpoint terms, and two-particle contact coefficient $c_\varepsilon \varepsilon^{-2} n_j n_{j+1}$. Conjecture: the projected resolvents on $P_{\le 2}\mathcal{F}$ converge iff $c_\varepsilon = -2t + \kappa\varepsilon + o(\varepsilon)$, giving Robin collisions with parameter $\kappa$ plus D1-type reaction coupling; if $(c_\varepsilon + 2t)/\varepsilon \to +\infty$, collisions are Dirichlet and sectors decouple; if $(c_\varepsilon + 2t)/\varepsilon \to -\infty$, a bound pair dives to $-\infty$ and there is no limit. The tuned family is the instance $t = 1$, $c_\varepsilon = -2 + a\varepsilon$, $a$ as in 94.10. Reading "above/below" at scale $\varepsilon$ sharpens INTEGRATION:113–115. It turns N1 and open problem 1 (NOTE:830) into a classification, testable with a modified `checks/review_tuned_convergence.py`.

**Effect on other shards.** D1-C (CA-77, Conjecture with Caveat) is proved for the tuned family by 94.6, without its closedness/core hypotheses; the caveat stands, since the tuning is necessary (94.1). CA-88's D1-C collision-sewing version is derived, its first lemma ($N \le 2$) superseded by 94.5 for all $N$; the restated R0$^u$ + R1 target stays open (INTEGRATION:61–79).

## 95.8 What This Does Not Claim

No limit for arbitrary or fixed densities, no classification of all self-adjoint or local stratified extensions, no global $H^2$ regularity, no renormalised higher-codimension reactions, no range-$r$ theorem beyond 94.13's hypotheses. The result is a $z$=2 nonrelativistic resolvent limit: no local unital refinement (R0$^u$), Wilson state (R1), modes, conformality, reflection-positive history sum, Ward/KS

data or fidelity (NOTE:821–834). $\hat{N}$ is not a CFT particle number.

## 95.9 Examples (Instances Only)

| Instance | Calibration or obstruction datum | Evidence |
|---|---|---|
| Any $\mathcal{C}$, $X = 0$ | $\mathcal{F}$ is the vacuum line; $H = 0$; consistent with 94.10. | NOTE:842; REVIEW:201 |
| Any $\mathcal{C}$, $X \neq 0$, pure hop | No scalar subtraction gives a full-Fock limit (94.1); the one-particle projected resolvent falls $\sim \varepsilon^2$ while the vacuum response stays fixed. | `review_N1_N3_output.txt`:1–4; `raw_output.txt`:10–13 |
| Vec, $X = \mathbf{1}$ (scalar fibres) | Tuned $N \leq 2$ compression, $\lambda = \kappa = \mu = 1$, $b = 0.7$, $v = 1.3$, $L \leq 96$: low eigenvalues Cauchy (max. differences $0.265, 0.143, 0.074$); $\langle\Omega\|H\|\Omega\rangle \to \gamma = 1.69$; $N$=2 weight $\approx 0.052$ retained. | `review_tuned_convergence_output.txt`:1–7 |
| $\mathrm{Hilb}_{\mathrm{fd}}(\mathbb{Z}_d)$, $X$ a generating grade, $d > r$ | $X^0, \ldots, X^r$ distinct simples, so $(G_r)$ holds; $X^{\otimes 2}$, $X^{\otimes(d+2)}$ share a simple; 94.11 applies (nearest-neighbour: $d = 3$, $r = 2$); $d = 3$ fails $G_3$. | Penneys:321; `review_N1_N3_output.txt`:5–8 |
| Any $\mathcal{C}$ with $M \neq 0$, weights on $b$ and $-b$ | Nonzero tiles, $\beta = 0$; with $\nu = 0$ the limit conserves $\hat{N}$ (94.7). | NOTE:845 |
| Non-self-dual $X$, $W = 0$ | $\nu = \gamma = 0$; pure contact weight from zero maps allowed (94.10). | REVIEW:196–197 |

# 96 Probe-Word Ward Lifting: Uniform Bounds, the Geometric Gap, and a False Converse

## 96.1 Status, Sources, and Dependencies

**Status: WEAKER-RESULT-PROVED (REVIEW:1–42).** Statements 95.1, 95.6, 95.8, 95.10, 95.11 are *derived*; 95.2, 95.3, 95.7 are *conditional*; 95.4, 95.5, 95.9 are *negative*; 95.12 is *conjectural*. The analytic implication is proved; G1–G2 are unverified geometric hypotheses. **CA-88 placement (INTEGRATION:44–65):** only annotations change in the R2 certificate column, CA-89 row and D2-C first-lemma pointer (optional CA-78/79 pointers); none becomes a general R2 certificate. $\mathsf{R0}^c, \mathsf{R0}^u, \mathsf{R1}, \mathsf{R3}$–$\mathsf{R5}$, WT-1/VR-1 and D2-C selection remain unchanged. The counterexample has only the trivial zero representation; the AP calibration does not move the open/chiral cell. NOTE, REVIEW and INTEGRATION mean the files in `explorations/2026-10-25-astra-r3/F4-ward-to-residual/`; check paths below are relative to that directory. Dependencies: CA-78 (78.1–78.2, 78.11, 78.13), CA-79 (prescribed modes), CA-81 (VR-1), CA-88 (grid and same-system rule), CA-89 (89.1, 89.5, 89.L); CONVENTIONS (v), (w), (al), (ao)–(ar). Supplementary notation is declared below; the proposed registration is in `SHARD_META.md`. No new categorical data are inferred.

## 96.2 Physical Probe Bounds and the Conditional Geometric Interface

**Lemma 95.1 [derived; any; category-free; → R2] (NOTE:49–111; REVIEW:51–58). Assume** finite positive quotients $H_r$, $K_r \geq 1$, dense linear spans $D_r$ of reflected probes; $\rho_r(u,v) = \langle u, R_r v\rangle$ descends through both null spaces. Pairings are linear in the second slot (al). Write $|R|_{s,2} = \|K^s R K^{-s-2}\|$ as in (ar). For each integer $s \geq 0$, fix $0 \leq \alpha_s < 1$ independently of $r$. Require, uniformly for *all linear combinations* $u, v \in D_r$,

$$|\rho_r(K_r^s u, e^{-tK_r} v)| \leq b_{r,s} t^{-s-1-\alpha_s} \|u\|\|v\|, \qquad 0 < t \leq 1. \tag{11}$$

**Prove** $|R_r|_{s,2} \leq C_{s,\alpha_s} b_{r,s}$, where $C_{s,\alpha} = ((s+1)!(1-\alpha))^{-1} + \sum_{j=0}^{s+1} 1/j!$.

⟨1⟩1. Finite dimensionality gives $D_r = H_r$; the dual norm identity turns (11) into $\|K_r^s R_r e^{-tK_r}\| \leq b_{r,s} t^{-s-1-\alpha_s}$.

⟨1⟩2. Spectral diagonalisation and integration by parts give $K^{-s-2} = ((s+1)!)^{-1} \int_0^\infty t^{s+1} e^{-tK}\, dt$. On $(0,1]$ the bound integrates to $b/((s+1)!(1-\alpha))$; on $[1,\infty)$, factor $e^{-tK} = e^{-K} e^{-(t-1)K}$, giving $b \int_1^\infty t^{s+1} e^{1-t} dt/(s+1)! = b \sum_{j=0}^{s+1} 1/j!$. Triangle inequality proves the claim.

⟨1⟩3. For an independent probe basis with Gram $G$ and *form* matrix $B$, physical unit vectors have coordinates $G^{-1/2}x$; the form norm is $\|G^{-1/2} B G^{-1/2}\|$. Quotient null coordinates first. This is not D2-L9's operator-coordinate transformation. No ambient-state faithfulness is needed.

**Theorem 95.2 [conditional; full per family; category-free; → R2] (NOTE:115–166, 175–183; REVIEW:59–64). Assume** 95.1's quotient, $N = N_r \to \infty$, geometric modes $S_{r,m} = N^{-1} \sum_j f_m(j) Q_{r,j} + b_r \delta_{m0}$, $f_m(j) = e^{2\pi i m j/N}$, as in (ar). Put $\ell_N = 2\pi/N$, $d_N(m) = \sin(m\ell_N)/\ell_N$, $\beta_N = 2d_N(m) - d_N(m+n)$, $N > |m+n|$. Declare a finite sewn-diagram sequence $F_0, \ldots, F_L$ on the *same* quotient, with: **G1** $F_L - F_0 = \langle u, [S_m, S_n] v\rangle + \rho_{\mathrm{end}}(u,v)$, representing both orderings; **G2** each step is its Ward contact plus residual, and the transported contact sum is $\langle u, (\beta_N S_{m+n} + z_{mn}^{(r)} I) v\rangle + \rho_{\mathrm{chart}}(u,v)$, where

$$z_{mn}^{(r)} = \delta_{m+n,0} \{\kappa_{3,r} d_N(m)^3 + (\kappa_{1,r} - 2b_r) d_N(m)\}.$$

**G3** all errors obey (11) with the same $\alpha_s$, after quotient, port transport, Fourier factors and insertion counts; $B_{r,m,n,s} := b_{\mathrm{end}} + b_{\mathrm{chart}} + \sum_i |a_i| b_i \to 0$. A directly proved smaller bound on the

assembled form may replace this absolute sum. **Prove** W2: $|\mathcal{R}_{r,m,n}|_{s,2} \le C_{s,\alpha_s} B_{r,m,n,s} \to 0$, with $\mathcal{R}_{r,m,n} = [S_{r,m}, S_{r,n}] - \beta_N S_{r,m+n} - z^{(r)}_{mn} I$.

$\langle 1\rangle 1$. Sum successive diagram differences: internal diagrams cancel. G1–G2 give $\langle u, \mathcal{R}_{r,m,n} v\rangle = \sum_i a_i \rho_i + \rho_{\text{chart}} - \rho_{\text{end}}$.

$\langle 1\rangle 2$. G3 and the triangle inequality give (11) with constant $B$. Apply 95.1. This identifies the actual residual; it does not derive G1 or G2 from density or D2's plane identity.

**Proposition 95.3 [conditional; full per family; category-free; → R2] (NOTE:168–183; REVIEW:63–64,131–139).** In addition to 95.2, assume A2 for $S$: $|S_{r,n}|_{s,1} \le M_{n,s}$, real $\kappa_{i,r} \to \kappa_i$, $b_r \to b$, and $|T_{r,n} - S_{r,n}|_{s,1} \to 0$ for the *prescribed* KS modes at all required orders. Then $T$ has A4, $c^* = 12\kappa_3$, $c = 24b - 12\kappa_1$. BY Lemma 89.5, the correction is bounded by $C_{s,\alpha_s} B + \ell_N^2(2|m|^3 + |m+n|^3) M_{m+n,s}/6 + |z^{(r)}_{mn} - z_{mn}| \to 0$; Theorem 89.1(i) transfers to $T$. A3 and mixed brackets remain separate; certificates must share CA-88's state/refinement system.

## 96.3 Two Obstructions and the Precise Analytic Converse

**Negative 95.4 [negative; any; category-free; → R2] (NOTE:185–201; REVIEW:65–67).** Fixed-word decay cannot replace (11): take $H_r = \operatorname{span}(e_0, \dots, e_r)$, $K_r = I$, $\Omega = e_0$, inclusion refinements and $R_r = |e_r\rangle\langle e_r|$. Finite matrix-unit probes acting on $\Omega$ span all finite-support vectors. Each fixed probe pair has eventually zero residual, also after heat collaring; yet $R_r e_r = e_r$, so $|R_r|_{s,2} = 1$. This refutes only word-dependent constants, not a uniform bound.

**Negative 95.5 [negative; full, both KS families; category-free; trivial R2] (NOTE:205–340; REVIEW:68–82).** A1–A4, including dense cyclic probes, do not imply vanishing subcritical collar constants for the same residual. **Construct** $N_r = 8 \cdot 2^r$, $a_r = e^{N_r^2}$, $\lambda_r = a_r^2 N_r^3$, $H_r = (\mathbb{C}|0\rangle \oplus \mathbb{C}^2)^{\otimes N_r}$, with active projection $P$. On the active block $X = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, $Y = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix}\right)$, $Z = \operatorname{diag}(1,-1)$; all vanish on $|0\rangle$. Set $k_j = a_r(P_j P_{j+1} + X_j Z_{j+1})$, $h_j = -k_j$, $v_r = 1$, $\varepsilon_r = b_r = 0$. With $z = e^{2\pi i/N}$, use exactly $T_n = \frac{N}{4\pi} \sum_j z^{nj}(k_j - \mathcal{J}_j)$, $\bar{T}_n = \frac{N}{4\pi} \sum_j z^{-nj}(k_j + \mathcal{J}_j)$, $\mathcal{J}_j = i[k_j, k_{j+1}]$. Let $U$ shift one site; $J_r$ inserts vacancies at odd *zero-based* fine sites, retaining old sites at even ones.

$\langle 1\rangle 1$. The active density is $a(I + X \otimes Z) \ge 0$, with eigenvalues $0, 2a$; other occupation blocks vanish. The vacancy vacuum has energy zero and is cyclic for product matrix units, giving dense probes.

$\langle 1\rangle 2$. Set $K_0 = 1$ on the vacuum and $\lambda_0$ off it. At the next scale let $F_+ = \operatorname{Ran} J_r$, $F_- = U_{r+1} \operatorname{Ran} J_r$: their intersection is the vacuum line and their nonvacuum parts are orthogonal. Transport $K_r$ by $J_r$ on $F_+$, by $U_{r+1} J_r$ on $F_-$; set $K_{r+1} = \lambda_{r+1} I$ on $(F_+ + F_-)^\perp$. The prescriptions agree at the vacuum. Since $U_{r+1}^2 J_r = J_r U_r$, induction gives $K \ge 1$, $\min\operatorname{spec} K = 1$, $[K, U] = 0$, $K_{r+1} J_r = J_r K_r$.

$\langle 1\rangle 3$. Every density kills $F_+ + F_-$; self-adjointness makes the complement invariant. All mode support therefore has $K_r = \lambda_r I$. Direct multiplication gives $\mathcal{J}_j = -2a^2 X_j Y_{j+1} Z_{j+2}$ and $\sup_n \|T_n\|, \sup_n \|\bar{T}_n\| \le L_N := N^2 a^2/\pi$. Thus adjoint pairing holds, A2 is bounded by $1/(\pi N)$, and, with $B_{mn} = 2|m| + |m+n|$,

$$|[T_m, T_n] - \beta_N T_{m+n}|_{s,2} \le \frac{2L_N^2 + B_{mn} L_N}{\lambda_N^2} \le \frac{2}{\pi^2 N^2} + \frac{B_{mn}}{\pi a_N^2 N^4} \longrightarrow 0.$$

Replacing $\beta_N$ by $m - n$ proves A4 with $c = c^* = 0$, also for the barred family; the mixed bracket is at most $2L_N^2/\lambda_N^2$. Since $T_{r+1,n} J_r = 0$, the A3 defect is $-J_r T_{r,n}$, bounded by $1/(\pi N_r)$, summable in $r$. Every limiting mode is zero.

$\langle 1\rangle 4$. On the all-active block, the coefficient of $X_j Y_{j+1} Y_{j+2} Y_{j+3} Z_{j+4}$ in $E = [T_1, T_{-1}] - \beta_N T_0$ is $c_N = N^2 a_N^4 \sin(4\pi/N)/\pi^2$. Only distance-two current pairs contribute: their commutator is

$8ia^4XYYYZ$, multiplied by $(N/(4\pi))^2(z^{-2}-z^2)$. Normalized Pauli trace gives $\|E\| \geq |c_N| \geq 8Na_N^4/\pi^2$.

$\langle 1\rangle 5$. Fix $s \geq 0$, $0 \leq \alpha < 1$, $p = s+1+\alpha$. For *sufficiently large* $N$, $t = p/\lambda_N \leq 1$; a proposed collar bound forces

$$b_N \geq p^p e^{-p}|c_N|\lambda_N^{-1-\alpha} \geq \frac{8p^p e^{-p}}{\pi^2} e^{2(1-\alpha)N^2} N^{-2-3\alpha} \longrightarrow \infty.$$

This control is nonphysical and stress collapses: physical-energy comparison, noncollapse and the full D2 certificate are not refuted.

**Proposition 95.6 [derived; any; category-free; → R2] (NOTE:344–365, 383–388; REVIEW:83–84, 86).** For finite $R_r$, $K_r \geq 1$, $q_{r,s} = |R_r|_{s,2} \to 0$, W2 is equivalent to the uniform assembled pairing bound

$$|\langle u, R_r v\rangle| \leq q_{r,s}\|K_r^{-s}u\|\|K_r^{s+2}v\|, \qquad \|K_r^s R_r e^{-tK_r}\| \leq A_{s+2}q_{r,s}t^{-s-2}, \quad A_p = (p/e)^p.$$

$\langle 1\rangle 1$. Factor the pairing through $K^s R K^{-s-2}$; uniform duality proves the converse.

$\langle 1\rangle 2$. Factor the heat operator similarly and maximize $x^p e^{-x}$ at $x = p$. The heat exponent is critical ($\alpha = 1$); its gamma-integral majorant diverges logarithmically at zero. Finite fixed Fourier sums obey $|\sum f_m g_n R_{r,m,n}|_{s,2} \leq \sum |f_m g_n| q_{r,m,n,s} \to 0$; growing support, shrinking contours and spatial Ward identities do not follow.

**Lemma 95.7 [conditional; any; category-free; → R2] (NOTE:367–381; REVIEW:85).** Assume additionally $\|K_r^s R_r e^{-tK_r}\| \leq B_s t^{-s-1-\alpha_0}$, $0 < t \leq 1$, $0 \leq \alpha_0 < 1$, $B_s$ independent of $r$. For $\alpha_0 < \alpha < 1$, $\theta = (\alpha-\alpha_0)/(1-\alpha_0)$, the collar constant $(A_{s+2}q_{r,s})^\theta B_s^{1-\theta} \to 0$ works with exponent $s+1+\alpha$. BY taking the minimum of this uniform envelope and 95.6's critical bound, then $\min(x,y) \leq x^\theta y^{1-\theta}$. The extra regularity repairs the analytic converse only; G1–G2 remain separate.

## 96.4 Spectral Calibration and the First Categorical Input

**Proposition 95.8 [derived; AP chiral CAR; category-free; → R2] (NOTE:390–504; REVIEW:87–94, 153–159).** Here *AP chiral CAR* means one chiral family on an antiperiodic circle, not an open-chain realization. For even $M \geq 8$, $\epsilon = 2\pi/M$, $r = -M/2+1/2, \ldots, M/2-1/2$, use (w)'s $A_n e_r = \cos^2(\epsilon n/4)\sin(\epsilon(r-n/2))e_{r-n}/\epsilon$, deleting transitions outside the window. Let $L_n = :d\Gamma(A_n):$ in the complex sea $1_{r<0}$, $Q = \sum_{\text{exc}}|r|$, $K = 1+Q$. For $n = 1,2,3$, put $R_{n,-n} = [L_n, L_{-n}] - 2nL_0 - (n^3-n)I/12$, $C_n = [A_n, A_{-n}] - 2nA_0$, $\delta_n = c_S(A_n, A_{-n}) - (n^3-n)/12$. Then, for every integer $s \geq 0$,

$$|R_{n,-n}|_{s,2} \leq U_n(\epsilon) := C_n^{\mathrm{bd}}\epsilon + B_n^{\mathrm{bd}}\epsilon^2, \qquad C_n^{\mathrm{bd}} = \pi(n + 7n^3/24) + \frac{(n-1)^2}{\pi(4.5-n)^2},$$

$$B_n^{\mathrm{bd}} = \sum_{r=1/2,3/2,\ldots,n-1/2} \left[\frac{(r-n/2)^4}{3} + \frac{n^2(r-n/2)^2}{8}\right], \quad (B_1^{\mathrm{bd}}, B_2^{\mathrm{bd}}, B_3^{\mathrm{bd}}) = (0, 7/24, 35/12),$$

$$\|K^s R_{n,-n} e^{-tK}\| \leq \sqrt{A_{s+2}U_n(\epsilon)H_n A_{s+1}}\, t^{-s-3/2}, \quad H_n = 4n + (n-1)^2/2 + B_n^{\mathrm{bd}}\pi^2/16, \quad 0 < t \leq 1.$$

$\langle 1\rangle 1$. The sourced CAR identity gives $R = :d\Gamma(C_n): + \delta_n I$; $C_n$ is diagonal, so $[R, Q] = 0$. At interior labels, $C_n(r) = 2n\sin(\epsilon r)[\cos(\epsilon r)\cos^4(\epsilon n/4)\operatorname{sinc}(\epsilon n) - 1]/\epsilon$, where $\operatorname{sinc}(x) = \sin(x)/x$, continuously extended at zero.

$\langle 1\rangle 2$. Elementary sine bounds give $|C_n(r)| \leq n\epsilon^2|r|^3 + (7n^3/12)\epsilon^2|r|$ in the interior. A deleted edge square is at most $(n-1)^2/4$; its label satisfies $\epsilon r^2 \geq \pi(4.5-n)^2/4$. Hence $|C_n(r)| \leq C_n^{\mathrm{bd}}\epsilon r^2$

everywhere. The $n$ Fermi-crossing terms give $|\delta_n| \le B_n^{\mathrm{bd}}\epsilon^2$; summing excitation contributions and using $\sum_{\mathrm{exc}} r^2 \le Q^2$ proves the full finite-Fock graph bound, independently of $s$.

$\langle 1\rangle 3$. Also $|C_n(r)| \le [4n + (n-1)^2/2]|r|$, so $|R| \le H_n K$. This supplies 95.7 with $\alpha_0 = 0$, $B_s = H_n A_{s+1}$; choose $\alpha = 1/2$. The all-size majorants are $O(M^{-1})$ and $O(M^{-1/2})$, only for these three diagonal pairs.

**Negative 95.9 [negative; AP chiral CAR; category-free; → R2] (NOTE:506–524; REVIEW:95–97).** Replace $Q$ by $E_{\mathrm{chiral}} = :d\Gamma(A_0):$, the excitation sum $\sum_{\mathrm{exc}} |\sin(\epsilon r)|/\epsilon$. Then $\liminf_M |R_{1,-1}|_{s,2;K=1+E_{\mathrm{chiral}}} \ge 1$.

$\langle 1\rangle 1$. Add a particle at $r_+ = M/2 - 1/2$ and a hole at $r_- = -r_+$: a neutral even unit vector with energy $2e_M$, $e_M = \sin(\epsilon/2)/\epsilon$. The two absent outward shifts give residual $-2u_M^2 - 4e_M$, $u_M = \cos^2(\epsilon/4)\sin(\epsilon)/\epsilon$; the $n = 1$ Schwinger term is zero.

$\langle 1\rangle 2$. Diagonality gives the lower bound $(2u_M^2 + 4e_M)/(1 + 2e_M)^2 \to 1$. This moving ultraviolet vector does not refute the source's fixed-core strong convergence. The third control $E_D = \sum_{\mathrm{exc}} 2|\sin(\epsilon r/2)|/\epsilon$ on this sea is only a dispersion proxy, not the finite two-component Dirac GNS system (NOTE:412–428; REVIEW:114–118).

**Proposition 95.10 [derived; AP chiral CAR; category-free; → R2] (NOTE:568–586; REVIEW:112–127).** With $a_n = c_S(A_n, A_{-n})$, $a_1 = 0$ and the inherited finite crossing sum gives $a_n \to (n^3 - n)/12$. Thus the two-mode estimators $c_M^* = 2(a_2 - 2a_1)$, $c_M = 2a_2 - 16a_1$ both tend to 1; at finite $M$ they need not fit $a_3$. The separate self-dual prescription halves the cocycle, giving $c^* = c = 1/2$ (source Eq. (28), (w)); it supplies no Majorana full-Fock graph estimate. These are spectral cocycles, not measured $\kappa_1, \kappa_3$ or a lattice chart-change theorem.

**Proposition 95.11 [derived; any locally, full with declared closure; L0; → R2] (NOTE:595–627; REVIEW:98).** For any unitary fusion $\mathcal{C}$, pointed $O$, complete dagger-stable basis $\{D_\alpha\}$ of $\mathrm{End}_{\mathcal{C}}(O \otimes O)$, $h = \sum_\alpha c_\alpha D_\alpha = h^\dagger$, the insertion to match is

$$k_j - \varepsilon_N I - \mathcal{J}_j = -\sum_\alpha \frac{c_\alpha}{v_N} D_\alpha^{(j)} - \varepsilon_N I - \frac{i}{v_N^2} \sum_{\alpha,\beta} c_\alpha c_\beta [D_\alpha^{(j)}, D_\beta^{(j+1)}].$$

BY CA-79's fixed prescription $k = -h/v_N$, $\mathcal{J}_j = i[k_j, k_{j+1}]$. CA-78 supplies positive Grams, associator recoupling and exact gate coefficients. Retain all charge/multiplicity matrix units, including available vacancy, hop, cup/cap and split/merge terms; no universal nine-tile basis or Ward-forcing subset follows. The next layer needs metric gate families, positive boundary quotient, coordinate/port transports and G1–G3; periodic closure/translation is declared (possibly relative half-braiding data). No $h$, conformal target or fidelity is selected.

## 96.5 Effect on other shards

*Amendment (2026-10-25, round 3):* CA-89 Lemma 89.L remains conditional: 95.1–95.3 isolate G1–G3; 95.5 refutes the arbitrary-control collar converse, and 95.10 supplies no geometric contacts. CA-78 Lemma 78.11 gains the explicit two-loss/probe constant (95.1); Target 78.13 points to 95.12. CA-79's calibration gains only 95.8–95.9, not all P0–P3. CA-81's graph, collar and geometric requirements remain distinct. CA-88 changes only the annotations listed above; its round-2 verdict/date remain intact (INTEGRATION:67–108).

## 96.6 What This Does Not Claim

Literal F4(a) and F4(c) remain unproved: density supplies neither quotient descent nor G1–G2 or a uniform accumulated bound. F4(b) is false under VR-1's auxiliary-control convention. No nontrivial common-system VR-1 certificate, all-bracket fermion estimate, standard local KS identification,

Wilson state, local conformal theory, category fidelity or functoriality is constructed (REVIEW:21–42,148–178). The general construction remains the target.

## 96.7 Examples (Instances Only)

| Instance | Result, with grade and tags inherited from the indicated statement |
|---|---|
| Any $\mathcal{C}$, declared periodic realization | 95.1–95.3, 95.11: analytic lifting applies; geometric/state data remain to be constructed. |
| Vec, $O = \mathbf{1} \oplus \mathbb{C}^2$ | 95.5: local prescribed KS, zero limiting stress, divergent subcritical collars. |
| Complex CAR, $K = 1 + Q$ | 95.8: at $M = 8, 16, 32$, full-Fock $q_1 := \lvert R_{1,-1}\rvert_{0,2}$ is 0.256416, 0.200672, 0.126705; no geometric Ward certificate. |
| Same CAR, $K = 1 + E_{\text{chiral}}$ | 95.9: neutral even ultraviolet witness gives limiting lower bound 1. |

Checks use Float64, not intervals: CAR tolerance $10^{-10}$, independent review $10^{-9}$ (INTEGRATION: 124–132). Off-diagonal checks restrict input to $Q \le 4$ but retain all generated outputs; the three sizes prove no decay exponent. The checks bundle records independent CAR/Gram/current invariants and rejected mutations (REVIEW:180–211).

## 96.8 Single Next Target

**Target 95.12 [conjectural; full per family; category-free; → R2]**
**(NOTE:713–724; REVIEW:213–225).** *Quantitative two-ordering endpoint realization* (INTEGRATION: 149–172). For specified metric-dependent local gates, their positive reflected quotient, coherent port transports and actual insertion modes $S$, construct a finite contour sweep whose endpoint forms equal the pairings with $S_m S_n$, $S_n S_m$, up to descending errors $e_r^+, e_r^-$ satisfying, for fixed $m, n, s$, fixed $0 \le \alpha_s < 1$, and $0 < t \le 1$,

$$|e_r^+(K^s u, e^{-tK} v)| + |e_r^-(K^s u, e^{-tK} v)| \le b_{r,m,n,s} t^{-s-1-\alpha_s} \|u\| \|v\|, \qquad b_{r,m,n,s} \longrightarrow 0.$$

Compute the endpoint forms from the finite gate/Gram data, never define them by the desired products. The estimate includes every count and transport and is uniform on physical probe spans. This would discharge G1; G2's cylinder comparison and G3's Ward-error accumulation remain. Use the same fields that must match prescribed KS.

# 97 Blocked Dyadic Lifts: Finite-Power Transport and the Failure of Single-Site Centrality

## 97.1 Status, Sources, and Dependencies

**Status: COUNTEREXAMPLE to the brief's F1 equivalence; accepted replacement results.** 96.1, 96.2, 96.4, 96.5, 96.7, 96.8 are *derived*; 96.3 and 96.6 are *negative*, with derived $\mathsf{R0}^u$ witnesses; 96.9 is *conditional*; 96.10 is *conjectural*. REVIEW:1–9 accepts all 35 numbered proof steps. Periodic, symmetry-preserving and canonically selected variants are *open/conjectural*, not conditional theorems (REVIEW:99–115 corrects NOTE:579).

**Grid movement** (INTEGRATION:49–68): (any; $\mathsf{L0}$ + blocked H; $\mathsf{R0}^u$) gains a sufficient certificate on $O^b$; bare $\mathsf{L0}/\mathsf{R0}^u$ gains the two stated classes. General existence/classification stays open. $\mathsf{L1}/\mathsf{R0}^u$ retains CA-90's sharper $r = 0$; $\mathsf{R0}^c$, periodic-seam lifts, the D1-C occupation-compatible target and R1–R5 do not move. Setting *any* means target-chirality independence; all constructions here use an *open ordered chain*. NOTE, REVIEW and INTEGRATION denote the files in `explorations/2026-10-25-astra-r3/F1-dyadic-lift-characterisation/`; `checks/` is its existing run bundle. Dependencies: CA-68 (literal insertion), CA-75 (Link (b1)), CA-76 (76.1, 76.4, 76.12), CA-77 (pointed carrier), CA-88 (grid), CA-90 (90.H, 90.0–90.2, 90.T); CONVENTIONS (a), (b), (r), (t), (am), (an), (aq), (as). New notation is fixed below and proposed for registration in the accompanying `SHARD_META.md`. Existing exact evidence: four baseline/mutation pairs and the independent multiblock check; commands, cutoffs, raw outputs and hashes are in `checks/README.md`:9–39 and REVIEW:168–209. These calibrate the proofs, not their quantifiers.

## 97.2 Setting and the Refuted Equivalence

$\mathcal{C}$ is unitary fusion, $\eta : \mathbf{1} \to O$ is an isometry, $p_0 = \eta\eta^\dagger$, $p_1 = 1 - p_0$, $W_L = O^{\otimes L}$, $W_0 = \mathbf{1}$, $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(W_L)$, $\mathcal{D} = \langle O \rangle$ full. Use (r)'s left bracketing and fixed unitary associators/unitors. $V_L$ inserts $\eta$ at even fine sites. Support means $\mathcal{A}_{[j,k]} = 1_{W_{j-1}} \otimes \mathrm{End}_{\mathcal{C}}(W_{k-j+1}) \otimes 1_{W_{L-k}}$, not merely commutation outside it; single-site algebras need not generate it. No charge-sector Hilbert factorization is assumed. The brief's labels mean (a) all-$L$ unital injective $*$-lifts of $V_L$ with one finite interval slack; (b) $[O]$ central in $K(\mathcal{D})$; (c) pointed unitary relative transport on $O$. The point in (c), and the isometry $\eta$ itself, repair omissions in the brief (REVIEW:149–166). By CA-90, pointed (c) implies (a) with $r = 0$ and (b); 96.3/96.6 refute the other claimed implications, except general (b)⇒(a), which remains open.

## 97.3 Vacancy Shuffle and the General Block Construction

**Lemma 96.1 [derived; any; $\mathsf{L0}$, pointed object; finite → $\mathsf{R0}^u$] (NOTE:59–89; REVIEW: 48–50).** For any pointed object in a unitary monoidal category, put $T = (\eta \otimes 1_O)p_1(1_O \otimes \eta^\dagger)$, $S = p_0 \otimes p_0 + p_1 \otimes p_1 + T + T^\dagger$. Then $S = S^\dagger = S^{-1}$ exchanges $1_O \otimes \eta$ and $\eta \otimes 1_O$; a fixed adjacent schedule gives $U_s(1_{W_s} \otimes \eta^{\otimes s}) = V_s$, with $U_s$ supported on $2s$ sites.

⟨1⟩1. By $\eta^\dagger\eta = 1$, $p_1\eta = 0$ and tensor interchange, $T^\dagger T = p_1 \otimes p_0$, $TT^\dagger = p_0 \otimes p_1$; the four orthogonal corners give $S^2 = 1$.

⟨1⟩2. Separate $1 = p_0 + p_1$ to check both insertion identities. Move each new vacancy left past later old sites in the stable interleaving schedule; every crossing meets $\eta$, proving the $U_s$ identity. QED. No braid relation for $S$ away from the inserted subobject is asserted.

**Theorem 96.2 [derived; any; $\mathsf{L0}$+declared blocked H; $\mathsf{R0}^u$] (NOTE:91–176; REVIEW: 51–57,84–95). Assume** a fixed $b \geq 1$, $P = W_b$, $\xi = \eta^{\otimes b}$, and unitary $t_Y : Y \otimes P \to P \otimes Y$ for

*every* $Y \in \mathcal{D}$, satisfying blocked H:

$$t_{\mathbf{1}} = 1_P, \quad t_Z(f \otimes 1_P) = (1_P \otimes f)t_Y, \quad t_{Y\otimes Z} = (t_Y \otimes 1_Z)(1_Y \otimes t_Z), \quad t_Y(1_Y \otimes \xi) = \xi \otimes 1_Y.$$

Here $f : Y \to Z$ is arbitrary; fixed associators are understood. **Prove** unital injective $*$-maps $\alpha_L : \mathcal{A}_L \to \mathcal{A}_{2L}$ for all $L \geq 1$, with coherent dyadic composites and

$$\alpha_L(a)V_L = V_L a, \qquad \alpha_L(\mathcal{A}_{[j,k]}) \subset \mathcal{A}_{[2j-1-r,\,2k-1+r]}, \qquad r = 2b-1, \tag{12}$$

endpoints clipped to $[1, 2L]$; the original $O$-site algebras and $V_L$ are unchanged.

⟨1⟩1. In a heterogeneous word $JR$, insert $P$ by $a \mapsto (1_J \otimes t_R)(a \otimes 1_P)(1_J \otimes t_R^\dagger)$. CA-90 Lemma 90.0 applies: compression by the point proves injectivity, and H proves intertwining. For an interval $a = 1_A \otimes a_{JR} \otimes 1_{C'}$ straddling the cut, the exact image is

$$1_A \otimes [(1_J \otimes t_R)(a_{JR} \otimes 1_P)(1_J \otimes t_R^\dagger)] \otimes 1_{C'}.$$

Indeed factor $t_{R\otimes C'}$ by H and cancel the exterior factor. Naturality handles intervals wholly to the right; interchange handles those to the left. Endpoints shift by $b$ precisely when after the cut.

⟨1⟩2. Fix the block origin at old site 1; write $L = mb + q$, $0 \leq q < b$, $Q = W_q$. Starting from $P^{\otimes m}Q$, insert a new $P$ after old block $i$, prefix length $(2i-1)b$, using the *entire* suffix, including $Q$. Append a new $Q$ by tensoring $1_Q$. The resulting map $\gamma_L$ intertwines the insertion $v_{\text{block}}$ in order old $P$, new $P$, …, old $Q$, new $Q$. This also covers $m = 0$ (only the tail) and $q = 0$ (no tail).

⟨1⟩3. Let $U_L$ be the tensor product of the $m$ copies of $U_b$ and the final $U_q$, omitted for $q = 0$. By 96.1, $U_L v_{\text{block}} = V_L$; hence $\alpha_L = \mathrm{Ad}_{U_L}\gamma_L$ has all algebra-map properties and $\alpha_L(a)V_L = V_L a$.

⟨1⟩4. Before this conjugation, an interval has endpoints $f(j), f(k)$, $f(s) = s + b\lfloor (s-1)/b \rfloor$. Appending $Q$ changes neither. Set $i(s) = \lfloor (s-1)/b \rfloor$, $a(s) = s - bi(s)$. Cell conjugation enlarges support only to the intersecting cells, so its endpoints obey

$$(2j-1) - (2bi(j)+1) = 2a(j) - 2 \leq 2b-2, \qquad 2b(i(k)+1) - (2k-1) = 2b - 2a(k) + 1 \leq 2b-1.$$

The shorter final cell only improves the bound. Exterior cell unitaries cancel, proving interval *membership*.

⟨1⟩5. Compose adjacent dyadic maps: associativity gives WT-1 A1. QED. The bound is sufficient, not optimal; at $b = 1$ CA-90 gives $r = 0$. Naturality is 96.8; arbitrary spatial boundary-extension or placement coherence is not part of this conclusion.

## 97.4 Two Obstructions to the Proposed Characterization

**Negative 96.3 [negative; any; bare L0, stated family; R0$^u$ witness] (NOTE:178–225; REVIEW:58–61).** The implications (a)⇒(b),(c) fail. For any finite nonabelian $G$, noncentral $g$ of order $b$, take $R = \bigoplus_{h\in G} h$, $O = R \oplus g$ in untwisted unitary $\mathrm{Vec}_G$, with $\eta$ in $R$'s unit summand. Then $\mathcal{D} = \mathcal{C}$; the following construction supplies (a) with $r = 2b-1$.

⟨1⟩1. Choose $x$ with $gx \neq xg$. The graded tensor rule gives $[O][x] - [x][O] = [gx] - [xg] \neq 0$; hence (b),(c) fail (CA-90, 90.6).

⟨1⟩2. With $N = |G|$, $R^2 = NR$, $Rh = hR = R$, so

$$[O]^b = [\mathbf{1}] + K[R], \qquad K = \sum_{k=1}^{b} \binom{b}{k} N^{k-1} = \frac{(N+1)^b - 1}{N}.$$

⟨1⟩3. In $P = O^{\otimes b}$, choose orthonormal grade bases $e_{h,a}$ with $e_{1,1} = \xi$. Set $\rho(k)e_{h,a} = e_{khk^{-1},a}$ and $t_Y(y_k \otimes p_h) = \rho(k)p_h \otimes y_k$. Multiplicities are $K+1$ at 1 and $K$ otherwise; conjugation preserves

them. The grade identity $(khk^{-1})k = kh$, $\rho(kl) = \rho(k)\rho(l)$, and the fixed unit-grade space prove all of pointed H; naturality is identity action on $Y$'s multiplicities. Apply 96.2. QED. For an involution, $b = 2$, $P \cong \mathbf{1} \oplus (N+2)R$, $r = 3$. Noncommuting simple grades exclude a braiding of $\mathcal{C}$.

**Proposition 96.4 [derived; any; L0; finite → R0$^u$] (NOTE:227–276; REVIEW:62–64).** CA-90's one-gap obstruction retains its unchanged-side scope. For insertion of $P$ between pointed words $J, R'$, its finite restriction equation concerns $P$, not its tensor root $O$:

$$\sum_c N_{ij}^c \Lambda_{cc'} = \dim \mathrm{Hom}(c', i \otimes P \otimes j), \qquad \Lambda_{xc'} = \dim \mathrm{Hom}(c', x \otimes P) = \dim \mathrm{Hom}(c', P \otimes x).$$

The second equality applies to simples $x$ occurring in *both* $J$ and $R'$.

⟨1⟩1. CA-90:197–204 fixes both side algebras pointwise; restriction along $\mathrm{End}(J) \otimes \mathrm{End}(R')$ yields the first equation, and the two unit rows the second.

⟨1⟩2. In 96.3, $N_{ij}^c = \delta_{c,ij}$ and $\Lambda_{cc'} = K + \delta_{cc'}$, giving $\sum_c N_{ij}^c \Lambda_{cc'} = K + \delta_{ij,c'} = \dim \mathrm{Hom}(c', i \otimes P \otimes j)$. QED. General dyadic side images are refined algebras with unknown inclusion matrices; interval containment does not give unchanged sides. This is precisely the failed cut extrapolation, refuted by 96.3, not a defect in 90.2(b).

**Theorem 96.5 [derived; any; L0, abelian graded; finite → R0$^u$] (NOTE:278–325; REVIEW:65–68). Assume** finite abelian $A$, normalized unitary 3-cocycle $\omega$, and $O = \bigoplus_g M_g \otimes g$ in $\mathrm{Vec}_A^\omega$, $m_g = \dim M_g$, $\eta \in M_0$. Here additive grade 0 is the unit simple; the associator $(g \otimes x) \otimes y \to g \otimes (x \otimes y)$ is $\omega(g, x, y)$, and products act right first. **Prove** pointed unitary half-braidings $z_x : O \otimes x \to x \otimes O$, inverse to $t_x$, on the *entire ambient category* are exactly

$$\rho_g(0) = 1, \quad \rho_g(y)\rho_g(x) = \beta_g(x,y)\rho_g(x+y), \quad \beta_g(x,y) = \frac{\omega(g,x,y)\omega(x,y,g)}{\omega(x,g,y)}, \quad \rho_0(x)\eta = \eta,$$

with $\rho_g(x) \in U(M_g)$. If $\kappa = \beta_g(x,y)/\beta_g(y,x)$ has finite order $d$, then $d \mid m_g$ for $m_g > 0$.

⟨1⟩1. Grade preservation forces each simple crossing to preserve $M_g$.

⟨1⟩2. From $(g \otimes x) \otimes y$ to $x \otimes (y \otimes g)$, the path via the $(g, x, y)$ associator, $z_{x+y}$, then the $(x, y, g)$ associator has matrix $\omega(g, x, y)\omega(x, y, g)\rho_g(x + y)$. Crossing $z_x$, reassociating by $\omega(x, g, y)$, then crossing $z_y$ gives $\omega(x, g, y)\rho_g(y)\rho_g(x)$. Equality yields the displayed orientation.

⟨1⟩3. Orthogonal sums, acting identically on external multiplicities, extend these blocks naturally to all objects; the same equation gives tensoriality, and the last equation gives the point. This proves sufficiency.

⟨1⟩4. Swapping $x, y$ gives $\rho_g(y)\rho_g(x) = \kappa\rho_g(x)\rho_g(y)$; determinants give $\kappa^{m_g} = 1$. QED. Divisibility is necessary only, and obstructs even invertible nonunitary transport.

**Negative 96.6 [negative; any; bare L0, stated family; R0$^u$ witness via 96.7] (NOTE: 327–365,447–450; REVIEW:69–71,136–147).** For every prime $p$, take $A = (\mathbb{F}_p)^3$, $\zeta = e^{2\pi i/p}$, $\omega(a, b, c) = \zeta^{a_1 b_2 c_3}$, $O = 0 \oplus e_1 \oplus e_2 \oplus e_3$, with the canonical grade-0 point. Then (b) holds, (c) fails, and (a) holds by 96.7 with $b = p^3$, $r = 2p^3 - 1$.

⟨1⟩1. For $F(a, b, c) = a_1 b_2 c_3$, expand $F(b, c, d) - F(a + b, c, d) + F(a, b + c, d) - F(a, b, c + d) + F(a, b, c) = 0$. Normalization is immediate; the sourced twisted construction is unitary fusion. The basis grades generate $A$, so $\mathcal{D} = \mathcal{C}$, and its fusion ring is commutative.

⟨1⟩2. In 96.5, $\beta_g(x, y) = \zeta^{g_1 x_2 y_3 + g_3 x_1 y_2 - g_2 x_1 y_3}$. At $(g, x, y) = (e_1, e_2, e_3)$, $\kappa = \zeta$ has order $p$, while $m_{e_1} = 1$; the required $p \mid 1$ fails. QED. Failure of single-site transport therefore does not obstruct a finite-slack algebra lift.

## 97.5 Bare Fusion-Data Existence and Naturality

**Theorem 96.7 [derived; any; bare L0, abelian $p$-group graded; R0$^u$] (NOTE:367–445; REVIEW:72–77). Assume** $|A| = p^d > 1$, exponent $p^e$, any normalized unitary $\omega$, and any

pointed $O \in \mathrm{Vec}_A^\omega$. **Prove** $b = p^{e+d-1}$ supplies pointed transport on $P = O^{\otimes b}$ over the entire ambient category, hence 96.2 gives a lift. No braiding is supplied or required.

⟨1⟩1. Put $Q(a,b,c,d) = \omega(b,c,d)\omega(a+b,c,d)^{-1}\omega(a,b+c,d)\omega(a,b,c+d)^{-1}\omega(a,b,c) = 1$. Direct cancellation, using abelianness, gives the normalized 2-cocycle identity

$$\frac{\beta_g(y,z)\beta_g(x,y+z)}{\beta_g(x+y,z)\beta_g(x,y)} = Q(g,x,y,z)^{-1}Q(x,g,y,z)Q(x,y,g,z)^{-1}Q(x,y,z,g) = 1.$$

⟨1⟩2. $c_g(x,y) = \beta_g(y,x)$ is also a 2-cocycle. On the $|A|$-dimensional basis $v_t$, set $R_g(x)v_t = c_g(x,t)v_{x+t}$. These phased permutations are unitary; the cocycle equation gives $R_g(x)R_g(y) = c_g(x,y)R_g(x+y)$, hence exactly the reversed product law of 96.5.

⟨1⟩3. In $\mathbb{Z}[A]$, set $s = \sum_g m_g[g]$, $n = \sum_g m_g$. Frobenius modulo $p$ and $p^e g = 0$ give $s^{p^e} \equiv n[0] \pmod p$. Inductively, $s^{p^{e+k-1}} \equiv n^{p^{k-1}}[0] \pmod{p^k}$: raise $C[0] + p^{k-1}T$ to $p$; for $k \geq 2$, nonconstant binomial terms have divisibility at least $1 + j(k-1) \geq k$ for $j < p$, and $p(k-1) \geq k$ for $j = p$. At $k = d$, every nonzero-grade multiplicity of $P$ is divisible by $|A|$.

⟨1⟩4. Fill those multiplicity spaces with copies of $R_g$. Since $\beta_0 = 1$, use the identity on the whole grade-zero space, fixing the actual $\xi = \eta^{\otimes b}$. Apply 96.5 and 96.2. QED. For trivial $A$, the flip gives $b = 1, r = 0$. Multiplicity-space identifications are choices, not canonical selections.

**Proposition 96.8 [derived; any; L0+transported blocked H; R0$^u$] (NOTE:452–517; REVIEW:78–82,104–106).** Let $F : \mathcal{C} \to \mathcal{C}'$ be a linear dagger-preserving strong monoidal equivalence with unitary tensorators $T_{Y,Z} : F(Y) \otimes F(Z) \to F(Y \otimes Z)$, $T_0 : \mathbf{1}' \to F(\mathbf{1})$, and unitary $u : F(O) \to O'$ with $\eta' = uF(\eta)T_0$. Transport $b, t$ and the fixed swap schedule. Then $F_{2L}\alpha_L = \alpha'_L F_L$, compatibly with composition and pointed unitary monoidal natural isomorphisms.

⟨1⟩1. Put $J_0 = T_0^\dagger$, $J_1 = u$, $J_{L+1} = (J_L \otimes u)T^\dagger_{W_L,O}$, $F_L(a) = J_L F(a) J_L^\dagger$. Tensorator associativity inductively gives compatibility with contiguous splittings and interval algebras.

⟨1⟩2. On image objects the transported datum is

$$t'_{F(Y)} = (J_b \otimes 1)T^\dagger_{P,Y}F(t_Y)T_{Y,P}(1 \otimes J_b^\dagger).$$

Every object of $\mathcal{D}' = \langle O' \rangle$ is unitarily isomorphic to an image from $\mathcal{D}$; full faithfulness and naturality make extension independent of this presentation. H and $\xi$ transport by the tensorator axioms.

⟨1⟩3. The formulas for $S, \gamma_L, U_L$ transport termwise; substitution gives the stated square and $J_{2L}F(V_L)J_L^\dagger = V'_L$. For a second equivalence $G$, recursion gives $J_L^{GF} = J_L^G G(J_L^F)$, hence $(GF)_L = G_L F_L$; identity point identification gives identity transport.

⟨1⟩4. For unitary monoidal $\lambda : F \Rightarrow G$ with $u_G \lambda_O = u_F$, induction gives $J_L^G \lambda_{W_L} = J_L^F$; naturality on $a, t$ identifies both transported lifts. QED. This is not R5.

**Proposition 96.9 [conditional; any; L0+blocked H+declared states; → R1] (NOTE: 600–605; REVIEW:111–115).** Along $L_n = 2^n L_0$, supply states $\omega_n$ and put $P_n = V_{L_n}V_{L_n}^\dagger$, $p_n = \omega_{n+1}(P_n)$, $e_n = \|\omega_{n+1}\mathrm{Ad}_{V_{L_n}}/p_n - \omega_n\|$ when $p_n > 0$. For the dyadic composites, $d_{k,n} := \|\omega_{n+1}\alpha_{n+1}^k - \omega_n\alpha_n^k\| \leq p_n e_n + 2(1-p_n)$, with $p_n e_n := 0$ at $p_n = 0$.

⟨1⟩1. Theorem 96.2 supplies exactly Proposition 76.1(iii)–(iv)'s intertwining hypothesis.

⟨1⟩2. *If* these bounds are summable, they supply the WT-1 drift hypothesis. QED by CA-76 and (an),(x). No nontrivial state estimate is supplied; the compatible vacancy state is only CA-90's inherited observation.

## 97.6 Effect on other shards

*Amendment (2026-10-25, round 3):* CA-90 Target 90.T's centrality deduction is refuted by 96.3; Proposition 90.2(b) keeps its unchanged-side one-gap scope (96.4), and (90.5) still characterizes one-site transport. CA-88's R0$^u$ certificate broadens as stated above; its round-2 next-target pointer

becomes 96.10. CA-75 Link (b1), CA-76 Proposal 76.5 and CA-68's unital-completion pointer gain blocked H and the bare-L0 families, with the literal $V_L$ retained. Proposition 76.1 applies unchanged; Link (b3) and Question 76.12's state/witness conditions remain open. CA-77's field-algebra/state-compatible problem is not solved. These are pointers (INTEGRATION:70–111); CA-91–CA-94 receive no mathematical amendment.

### 97.7 What This Does Not Claim

General (b)⇒(a), necessity of finite-power transport and classification of all local lifts remain *conjectural/open* (any; L0; → R0$^u$). The $p$-group theorem is not a theorem for all finite abelian groups or fusion categories. Periodic seams, occupation-degree preservation, translation/reflection, spatial boundary-extension consistency and canonical selection are *open/conjectural*, not conditional results (REVIEW:99–115). No higher-rung state, mode, conformal or fidelity certificate follows. The original three-summand $\mathrm{Vec}_{S_3}$ carrier remains undecided; the different carriers below do not decide it.

### 97.8 Examples (Instances Only)

First three rows: derived instances (any; bare L0; R0$^u$). Last: a finite check of 96.2 (derived; any; L0+declared H; → R0$^u$); it does not prove an all-$L$ statement.

| Instance | Exact datum or finite check | Evidence |
|---|---|---|
| $S_3$, $O = R \oplus (12)$ | $O^2 \cong \mathbf{1} \oplus 8R$, $b = 2, r = 3$; noncentral site. | NOTE:652; 96.3 |
| $(\mathbb{F}_p)^3$, trilinear $\omega$ | $O = 0 \oplus e_1 \oplus e_2 \oplus e_3$; (a),(b), not (c); $b = p^3$. | NOTE:653; 96.6–96.7 |
| $\mathbb{Z}/4$, $\mathbb{Z}/4 \times \mathbb{Z}/2$ | Any normalized unitary $\omega$, pointed $O$; $b = 8, 16$, respectively. | NOTE:655; 96.7 |
| $\mathrm{Vec}_{\mathbb{Z}/2}$, signed H | $b = 3, L \leq 7$: 832 interval units, 803076 entries; all intervals for $L \leq 4$, lengths $\leq 3$ otherwise. Tail mutation fails locality despite exact $V$. | REVIEW:196–205 |

### 97.9 Next Target: Finite-Power Transport Extraction

**Target 96.10 [conjectural; any; L0; → R0$^u$] (NOTE:626–628; REVIEW:211–222).** Assume exactly (a) for arbitrary unitary fusion $\mathcal{C}$ and pointed $O$: all $L$, exact $V_L$, and one finite slack in (12). Prove or refute the existence of a finite $b$ and pointed unitary transport on $(O^{\otimes b}, \eta^{\otimes b})$ over full $\langle O \rangle$. No unchanged-side, occupation or translation hypothesis may be added. Together with 96.2 a positive answer would give a replacement characterization; the unknown side inclusions and associator coherence must be reconstructed (INTEGRATION:145–155).

# 98 What the Tower Remembers: Rooted AF Invariants and Marked Refinements

## 98.1 Status, Sources, and Dependencies

**Status: WEAKER-RESULT-PROVED** (REVIEW:3–12). Statements 97.1–97.2, 97.4–97.8 and 97.11–97.13 are *derived*; 97.3, 97.9–97.10 and 97.14 are *negative*; 97.15 is *conditional*; 97.16 is *conjectural*, an open alternative. No reviewed GAP is promoted. NOTE, REVIEW and INTEGRATION denote files in `explorations/2026-10-25-astra-r3/F6-what-the-tower-remembers/`; all statement locators refer there. **CA-88 cells (INTEGRATION:43–62):** (any; $\mathsf{L0}$; $\mathsf{R0}^u$) gains classification in *trailing geometry*; (any; $\mathsf{L0}+H$; $\mathsf{R0}^u$) gains fixed-$H$ dyadic compatibility and marked extraction; (any; $\mathsf{L0}+H$; $\mathsf{R1}$) gains only the canonical tracial zero-drift instance on that schedule. The $\mathsf{R4}$ insufficiency annotation strengthens; no $\mathsf{R2}$–$\mathsf{R5}$ attainment, $\mathsf{R0}^c$, $\mathsf{L3}$ fidelity or full/periodic cell moves. Bare-$\mathsf{L0}$ dyadic existence is not proved. Dependencies: CA-77 (77.1–77.3), CA-80 (80.8–80.10), CA-88 (grid), CA-90 (90.H, 90.1, 90.3), CA-93 (93.2, 93.8); CA-71–CA-73 and CA-94 supply boundary comparisons only. CONVENTIONS (a), (b), (r), (am), (aq), (as), (av) apply; the additional scoped notation below is proposed in SHARD_META.md. Convention (ar) is unrelated Ward notation.

The checks bundle records six exact author groups, two restored review runs and five intended assertion failures (REVIEW:165–194). These test finite identities, not classification or infinite reconstruction. No local primary AF classification source was found (NOTE:291–300; REVIEW:241–245); 97.8 uses the reviewed finite-stage proof.

## 98.2 Setting and Rooted Invariant

$\mathcal{C}$ is unitary fusion, $O \neq 0$, $\mathcal{D} = \langle O \rangle$, $I = \operatorname{Irr}\mathcal{D}$, $W_0 = \mathbf{1}$, $W_L = O^{\otimes L}$ left-associated, $A_L = \operatorname{End}_{\mathcal{C}}(W_L)$, $i_L(a) = a \otimes 1_O$. Use column incidence $N[y,x] = \dim \operatorname{Hom}(y, x \otimes O)$, $m_0 = e_{\mathbf{1}}$, $m_L = N^L e_{\mathbf{1}}$, $S_L = \{x : m_L[x] > 0\}$; absent blocks are omitted. Let $\kappa_L : A_L \to A_\infty$ be the trailing limit maps. An AF algebra is the norm completion of an increasing finite-dimensional union. $K_0$ is the group completion of stable Murray–von Neumann projection classes; its cone comes from projections and its scale is [1]. Tr below is ordinary unnormalised matrix trace, rank-one value one; $d_x$ is the canonical positive dimension. No point is needed until stated; then $\eta : \mathbf{1} \to O$, $\eta^\dagger \eta = 1$, as in (am).

**Theorem 97.1 [derived; any; $\mathsf{L0}$; $\mathsf{R0}^u$, trailing] (NOTE:67–153; REVIEW:29–37). Assume** the setting above. **Prove** the maps $i_L$ form a local unital injective tower, with

$$A_L \cong \bigoplus_{x \in S_L} M_{m_L[x]}(\mathbb{C}), \qquad (K_0(A_\infty), K_0(A_\infty)^+, [1]) = \varinjlim_L (\mathbb{Z}^{S_L}, \mathbb{Z}_+^{S_L}, m_L; N|_{S_{L+1} \times S_L}).$$

Its Bratteli diagram is the *rooted reachable unfolding*: vertices $S_L$, edges $N[y,x]$.

⟨1⟩1. Orthogonal splitting gives $\operatorname{Hom}(y, W_L \otimes O) = \bigoplus_x \operatorname{Hom}(x, W_L) \otimes \operatorname{Hom}(y, x \otimes O)$; $i_L(a)$ acts as $a_x \otimes 1$. Thus multiplicities are $N[y,x]$, and $\sum_y N[y,x] d_y = d_x d_O > 0$ prevents any source block disappearing. Tensor dagger/interchange gives unitality and locality; the maximum block norm proves isometry and the AF completion.

⟨1⟩2. Stable projection classes in a finite sum of matrices are nonnegative block ranks; the unit has rank vector $m_L$, and inclusions act by $N$.

⟨1⟩3. Continuity uses only finite-dimensional stages. Approximate a limit projection $p$ by self-adjoint finite-stage $b$, $\|b - p\| < \epsilon < 1/4$; $q = 1_{[1/2,\infty)}(b)$ has $\|q - p\| < 2\epsilon < 1$. For $z = qp + (1-q)(1-p)$, $z - 1 = (q-p)(2p-1)$, so $z(z^*z)^{-1/2}$ conjugates $p$ to $q$. Conversely, approximate

a limit partial isometry $v$ between finite projections $p, q$ by finite $w$. For $x = qwp$, both $x^*x$ and $xx^*$ are invertible in their corners; $x(x^*x)^{-1/2}$ witnesses equivalence there.

⟨1⟩4. Every class and relation is therefore witnessed at a finite stage; group completion gives the displayed cone and unit. Common-stage coordinate order also proves directedness, unperforation and interpolation. QED.

**Proposition 97.2 [derived; any; L0; R0$^u$, trailing] (NOTE:155–210; REVIEW:38–41).** Put $G_N = \varinjlim(\mathbb{Z}^I, \mathbb{Z}^I_+; N)$, $u = [e_{\mathbf{1}}, 0]$. Then $K_0(A_\infty) \cong I_u := \{g \in G_N : -nu \le g \le nu \text{ for some } n \in \mathbb{N}\}$, with inherited cone and unit $u$. If $N$ is primitive, $I_u = G_N$; a pointed $O$ makes $N$ primitive on $I$.

⟨1⟩1. Extend supported vectors by zero. Eventual equality and positivity show an order embedding. Every supported $v$ is bounded by $\pm n m_L$; conversely such bounds, witnessed at one later level, force support in $S_L$.

⟨1⟩2. All simples of $\mathcal{D}$ occur in nonnegative powers. Indeed their union $S$ is tensor/summand closed. For $x \in S$, $P[z, y] = N^z_{xy} d_z/(d_x d_y)$ is column-stochastic with strictly positive stationary weights proportional to $d^2_y$, by reciprocity. No edge exits $S$; stationarity forbids entry, so the edge $x^* \to \mathbf{1}$ forces $x^* \in S$. Thus $S = I$.

⟨1⟩3. A point makes $S_L$ increasing, hence eventually full. If $O^K$ contains every simple, for each $x, y$ choose $z \subset x^* \otimes y$; reciprocity gives $N^y_{xz} > 0$, so $(N^K)[y, x] > 0$. Conversely $N^p > 0$ gives a full supported tail (no zero rows); the limits then share that tail. QED.

**Negative 97.3 [negative; any; L0; R0] (NOTE:212–222; REVIEW:42–43).** For an invertible generator $O = g$ of a cyclic group of order $q > 1$, $A_L = \mathbb{C}$, $i_L = \mathrm{id}$, so rooted $K_0 = \mathbb{Z}$. But $N$ is a cyclic permutation and $G_N = \mathbb{Z}^q$ with coordinate cone; under $N^{-L}$, $u$ lies on one axis and is not an order unit of $G_N$. Tensor generation alone is insufficient. Also $O = 0$ kills $A_0 = \mathbb{C}$ at the first map and has $d_O = 0$; the nonzero hypothesis is essential (NOTE:38–39).

## 98.3 Trace and Classification

**Theorem 97.4 [derived; any; L0; R0 with trace] (NOTE:226–278; REVIEW:44–48).** With $\lambda = d_O$, $\tau_L(a) = \lambda^{-L} \sum_{x \in S_L} d_x \operatorname{Tr}(a_x)$ is a compatible tracial state, unique if $N$ is primitive. Rank-one weights are $d_x/\lambda^L$; block-identity weights are $m_L[x] d_x/\lambda^L$.

⟨1⟩1. $d^T N = \lambda d^T$, $d^T m_L = \lambda^L$ prove normalization, positivity and compatibility.

⟨1⟩2. Any trace has nonnegative rank-one weights $w_L = N^T w_{L+1}$, $m_L^T w_L = 1$. If $N^p > 0$, then $m_{L+p}[y]/\lambda^{L+p} \ge c_y := \min_x (N^p)[y, x]/(\lambda^p d_x) > 0$. Consequently $r_L[y] = \lambda^L w_L[y]/d_y \le 1/(c_y d_y)$ eventually.

⟨1⟩3. $P[x, y] = N[y, x] d_y/(\lambda d_x)$ is row-stochastic, $r_L = P r_{L+1}$. For $s = |I|$, $\delta_0 = \min(P^p) > 0$, subtracting $\delta_0$ from each entry gives $\operatorname{osc}(P^p v) \le (1 - s\delta_0) \operatorname{osc}(v)$. Iteration and the bound force $r_L$ constant; normalization makes it one. Compatibility fixes earlier levels; norm continuity extends the state. QED.

**Proposition 97.5 [derived; any, open; L0 + $H$; R1, tracial only] (NOTE:280–287; REVIEW:111–114).** On the fixed pointed-$H$ schedule of 97.11, $\tau_{2L} \alpha_L = \tau_L$: spherical trace is invariant under the shuffle conjugation and trailing insertion $L$ times preserves $\tau$. Thus every Wilson drift is zero, giving the compatible limit state on this declared schedule. This selects neither a critical state nor a conformal vacuum.

**Lemma 97.6 [derived; any; category-free; → R0] (NOTE:302–319; REVIEW:49–50).** For $B = \bigoplus_i M_{b_i}$, $C = \bigoplus_j M_{c_j}$, all sizes positive, unital $*$-maps $B \to C$ modulo target-unitary conjugacy correspond exactly to nonnegative integer matrices $T$ with $Tb = c$; the induced $K_0$ map is $T$.

⟨1⟩1. In each target block, matrix units split the source's central summands into equivalent orthogonal ranges of dimensions $T[j,i]$, giving $a_i \mapsto a_i \otimes 1_{T[j,i]}$.

⟨1⟩2. Dimension counting gives $Tb = c$; these blocks construct every such map. Matching bases give target-unitary conjugacy, and minimal projection ranks give $T$. QED.

**Proposition 97.7 [derived; any; L0; R0$^u$, trailing] (NOTE:321–342; REVIEW:51–53).** A unit-preserving based-ring isomorphism carrying $[O]$ to $[O']$ gives a stage-filtered trailing-tower isomorphism.

⟨1⟩1. It matches $N, m_L, S_L$. Inductively choose a block-label-preserving isomorphism at the next stage; its two maps from the preceding stage have equal multiplicities. Conjugate the new target using 97.6.

⟨1⟩2. The resulting commuting isometries extend to the completions. No interval, concatenation, placement or dyadic mark is promised. Canonical positive dimensions already depend only on the based ring. QED.

**Theorem 97.8 [derived; any; category-free; → R0] (NOTE:344–413; REVIEW:54–61).** For sequential unital AF limits $A = \varinjlim A_n$, $B = \varinjlim B_m$ of *finite-dimensional* algebras with injective connecting maps, $A \cong_* B$ iff $(K_0(A), K_0(A)^+, [1_A]) \cong (K_0(B), K_0(B)^+, [1_B])$.

⟨1⟩1. A $*$-isomorphism preserves all projection data. Conversely, lift the finitely many positive columns of a unit-preserving stage-to-limit group map to one finite stage, by 97.1's continuity proof. Its unit discrepancy vanishes at a later stage; there $Tb = c$ exactly. Lemma 97.6 realises this lift as a unital $*$-map.

⟨1⟩2. Equality of two stage group maps into a limit is equality of finitely many columns at some later stage; 97.6 upgrades this to target-unitary conjugacy of their algebra maps.

⟨1⟩3. Alternately lift the group isomorphism and its inverse along increasing cofinal stages. Using step 2, conjugate each new target to impose exact triangles $\psi_k\phi_k = i_{n_k,n_{k+1}}$, $\phi_{k+1}\psi_k = i'_{m_k,m_{k+1}}$, leaving earlier triangles unchanged. No stage $K_0$ map is assumed injective. Each algebra map is injective because its completed triangle is.

⟨1⟩4. The triangles give compatible isometric maps and inverse maps on dense unions, hence inverse maps on the completions. QED. This classifies the *unmarked limit*, not its filtration or endomorphisms.

## 98.4 Information Loss

**Negative 97.9 [negative; any; L0; R0] (NOTE:422–475; REVIEW:62–66).** For any finite group $G$, $|G| = q$, normalized unitary cocycle $\omega$, and regular site $O = \bigoplus_{g\in G} g$ in $\mathrm{Hilb}_{\mathrm{fd}}(G,\omega)$, put $J_q = (1)_{x,y}$. Then

$$N = J_q, \quad m_L = q^{L-1}\mathbf{1}\ (L \geq 1), \quad (K_0, K_0^+, [1]) = (\mathbb{Z}[1/q], \mathbb{Z}[1/q]_{\geq 0}, 1),$$
$$A_\infty \cong \varinjlim (M_{q^L}, b \mapsto b \otimes 1_q).$$

⟨1⟩1. Exactly one $g = x^{-1}y$ gives each edge; $[v, L] \mapsto q^{-L}\sum_x v_x$ gives the ordered group: zero-sum vectors die under $J_q$, and positive sums become positive vectors; the unit maps to one.

⟨1⟩2. For $B_L = \bigoplus_g M_{q^{L-1}}$, block-diagonal inclusion into $M_{q^L}$ and replication into $B_{L+1}$ give exact interleaving triangles. This is also the limit for Vec, $O' = q\mathbf{1}$, with fusion-ring rank one versus $q > 1$. Both inputs are pointed, generating and primitive. For fixed Vec, $q\mathbf{1}$ and $q^k\mathbf{1}$, $q, k \geq 2$, have identical limits by cofinal blocking, but different site classes. QED.

**Negative 97.10 [negative; any; L0; R0, obstruction to decoding] (NOTE:477–508,668–686; REVIEW:67–69,79–81).** If two normalized cocycles on $G$ are unrelated by automorphism and coboundary, their regular-site categories are monoidally inequivalent but have isomorphic stage-filtered trailing towers with matching canonical trace.

⟨1⟩1. Any tensor equivalence permutes invertible simples by a group isomorphism; normalized tensorator scalars $\beta$ force the cocycle ratio, after relabelling, to be

$$(\partial\beta)(a,b,c) = \frac{\beta(b,c)\beta(a,bc)}{\beta(ab,c)\beta(a,b)}.$$

The assumed obstruction forbids it.

⟨1⟩2. Apply 97.7 and the identical dimension weights of 97.4. The explicit E-F instance below realises the obstruction. A decoder depending only on this output record would assign one record two inequivalent inputs, impossible. QED. No Morita-inequivalence claim is made. Fully marked CA-77 reconstruction can already retain $F$ at R0; a compatible trace need not restore it. Thus "$F, R$ first enter at R1" is false.

## 98.5 What the Refinement Record Adds

Here $O$ is pointed and $H = (t_Y)$ is CA-90's unitary relative half-braiding on the full $\mathcal{D}$: $t_Y : Y \otimes O \to O \otimes Y$, natural, $t_{\mathbf{1}} = 1$, $t_{Y\otimes Z} = (t_Y \otimes 1_Z)(1_Y \otimes t_Z)$, $t_Y(1_Y \otimes \eta) = \eta \otimes 1_Y$. Coherent associators are understood. L1 supplies such $H$; $H$ need not supply a braiding on $\mathcal{D}$. Set $\alpha_L = \mathrm{Ad}_{B_L}(\cdot\otimes 1_{W_L})$, where $B_L$ interleaves old/new strands, crossing old $j$ with new $i$ exactly when $i < j$ (CA-90). **Proposition 97.11 [derived; any, open; L0 + $H$; R0$^u$] (NOTE:520–558; REVIEW:70–72,259–265).** For *one fixed* $H$ at every length,

$$\alpha_{L+1}(a \otimes 1_O) = \alpha_L(a) \otimes 1_O^{\otimes 2}, \qquad \delta\kappa_L = \kappa_{2L}\alpha_L, \qquad \delta_* = \mathrm{id}, \quad \tau\delta = \tau. \tag{13}$$

Here $\delta$ is an injective unital AF endomorphism; its powers form a discrete semigroup.

⟨1⟩1. If $\Sigma_L : O \otimes W_L \to W_L \otimes O$ moves the last old strand across the first $L$ new ones, $B_{L+1} = (B_L \otimes 1_O^{\otimes 2})(1_{W_L} \otimes \Sigma_L \otimes 1_O)$. Only disjoint crossings are reordered.

⟨1⟩2. The right factor commutes with $a$ on the first $L$ factors, proving the square; the isometric maps therefore define $\delta$ on the union and its completion.

⟨1⟩3. Inner conjugation fixes projection classes, so $\delta_*[v, L] = [N^L v, 2L] = [v, L]$. The trace assertion is 97.5's conjugation calculation. QED; no continuous flow is constructed. Reviewer correction: $H$ already implies $t_{12}t_{23}t_{12} = t_{23}t_{12}t_{23}$, by tensoriality and naturality for $t_O$; step 1 needs no extra braid axiom (REVIEW:259–265).

**Proposition 97.12 [derived; any, open; L0 + $H$; R0$^u$] (NOTE:560–588; REVIEW:73–75).** The *abstract* dyadic diagram at $L_n = 2^n L_0$, $L_0 \geq 1$, is isomorphic to the trailing cofinal diagram.

⟨1⟩1. Put $j_n(a) = a \otimes 1_{W_{L_n}}$, $U_0 = 1$, $U_{n+1} = j_n(U_n)B_{L_n}^\dagger$. Direct multiplication gives $\mathrm{Ad}_{U_{n+1}}\alpha_{L_n} = j_n \mathrm{Ad}_{U_n}$. QED. Its negative consequence is loss of $H$ and $F$ discrimination (E-H, E-F); comparison requires $H$ on both inputs. These conjugacies need not also preserve trailing maps, interval inclusions or named junctions.

**Theorem 97.13 [derived; any, open; L0 + $H$ and named marks; R0$^u$] (NOTE:590–634; REVIEW:76–78).** Retain the pointed tensor record and the following typed placements, adjoints and compression identifications:

$$r = \eta \otimes 1_O,\ l = 1_O \otimes \eta : O \to W_2, \quad h = rl^\dagger \in A_2,$$
$$i = 1_O \otimes 1_O \otimes \eta \otimes \eta,\ j = \eta \otimes 1_O \otimes 1_O \otimes \eta : W_2 \to W_4.$$

For the supplied-$H$ formula, $t_O = j^\dagger \alpha_2(h) i$; the tensor marks determine all of $H$.

⟨1⟩1. Naturality for $\eta$ gives $t_O(\eta \otimes 1) = 1 \otimes \eta$. With $B = 1 \otimes t_O \otimes 1$, $B^\dagger i = 1 \otimes \eta \otimes 1 \otimes \eta$, $(h \otimes 1 \otimes 1)B^\dagger i = j$, and $Bj = jt_O$. Thus $j^\dagger B(h \otimes 1 \otimes 1)B^\dagger i = t_O$, since $j^\dagger j = 1$.

⟨1⟩2. Tensoriality fixes $t_{W_n}$; for each isometric summand $v : Y \to W_n$, $t_Y = (1_O \otimes v^\dagger) t_{W_n} (v \otimes 1_O)$. Proposition 97.2 exhausts simples; naturality makes this independent of splitting, and direct sums finish. QED. This recovers supplied $H$, not $H$ from arbitrary local lifts.

**Negative 97.14 [negative; any, open; L0 and lengthwise data; R0] (NOTE:646–651,891–914; REVIEW:88–90).** Lengthwise finite CA-90 axioms and dyadic composability do not imply (13). In E-H below, $\alpha_2^+(h) \otimes 1^{\otimes 2}$ has coefficient $+1$ where $\alpha_3^-(h \otimes 1)$ has $-1$. Both maps separately obey the finite axioms. This refutes automatic descent, not the fixed-$H$ construction.

**Theorem 97.15 [conditional; chiral/named chiral subnet; L0+$L4$ / L1+$L4$; R4 given R3] (NOTE:688–707; REVIEW:82–83,131–137). Assume** a declared completely rational chiral net $\mathcal{A}$ on a separable Hilbert space and a fully faithful unitary tensor functor $\Phi : \mathcal{C} \to \mathrm{DHR}_{\mathrm{fin}}(\mathcal{A})$. On its actual four-interval inclusion $\mathcal{N} = \mathcal{A}(I_1) \vee \mathcal{A}(I_3) \subset \mathcal{M} = (\mathcal{A}(I_2) \vee \mathcal{A}(I_4))'$, assume increasing represented $\mathcal{N}_k \subset \mathcal{M}_k$ boundedly exhaust both algebras, normal expectations $E_k : \mathcal{M}_k \to \mathcal{N}_k$, and one normal state $\psi$ on $\mathcal{M}$, faithful on $\mathcal{N}$, preserved by every $E_k$. Require $b := \liminf K(E_k) \leq \operatorname{Dim} \mathcal{C} = \sum_x d_x^2$, with cost $K$ as in (av). **Prove** $\mu_{\mathcal{A}} = \operatorname{Dim} \mathcal{C}$ and $\Phi$ is onto all finite sectors up to equivalence; if $\Phi$ is braided at L1, the equivalence is braided and $\mathcal{C}$ modular. BY CA-93 Lemma 93.2 and Theorem 93.8. This is a conditional restatement, not an application: an AF trace supplies none of these represented certificates. The full setting needs an identified chiral subnet or CA-93's specified doubled net, not the diagonal extension's DHR category.

## 98.6 Effect on other shards

*Amendment pointers (2026-10-25, round 3; INTEGRATION:64–114).* CA-77 keeps its corner and fully marked reconstructions distinct from 97.8; CA-71–CA-73 keep their corner qualifications. CA-90 gains 97.11–97.14, without resolving its converse; 97.11 records the reviewer's correction to 90.3's self-Yang–Baxter parenthetical. CA-80's insufficiency warning extends by 97.10; CA-93 gains only the boundary in 97.15; CA-88 gains exactly the cells listed above. CA-94 retains its reaction/tensor marks and its distinction from R5. CA-74 is unaffected.

## 98.7 What This Does Not Claim

No classification of interval nets or AF endomorphisms by $K_0$, no effective dimension-group decision algorithm, no universal loss of $F$ at R0, and no recovery from arbitrary lifts. No periodic seam, parity covariance, critical-state selection, modes, conformal limit, central-charge lift, characters or coherent CFT synthesis is constructed.

## 98.8 Examples (Instances Only)

All entries have (any; L0 plus the displayed declarations; R0); their negative conclusions are 97.3, 97.9, 97.10, 97.14. Exact data: NOTE:839–914; REVIEW:84–90. These are tests of the general statements.

| Instance | Exact calibration / counterexample |
|---|---|
| E-P: $\mathrm{Hilb}_{\mathrm{fd}}(C_3), O = g$ | $A_L = \mathbb{C}$; rooted $K_0 = \mathbb{Z}$, full stationary $G_N = \mathbb{Z}^3$, $u$ on one axis. |
| E-R / E-V | Regular sites of $C_4$ and $C_2 \times C_2$: $N = J_4$, $m_L = 4^{L-1}\mathbf{1}$; only the former has a basis element of order four. Both limits equal that for $\mathrm{Vec}, O' = 4\mathbf{1}$, of ring rank one. |
| E-O / E-0 | $\mathrm{Vec}, O = 2\mathbf{1}, 4\mathbf{1}$: same limit by even-stage blocking, different $[O]$. Any $\mathcal{C}, O = \mathbf{1}$: $A_\infty = \mathbb{C}$, $\mathcal{D} = \mathrm{Vec}$; the ambient complement is invisible. |
| E-F: twisted $C_2$, $O = \mathbf{1} \oplus g$ | Labels $a, b, c, d \in \{0, 1\}$, $\omega_+ = 1$, $\omega_-(a, b, c) = (-1)^{abc}$. Pentagon exponents both reduce to $bcd + abd + acd + abc$ mod 2; normalized $\partial\beta(1, 1, 1) = 1$, whereas $\omega_-(1, 1, 1) = -1$, with no nontrivial automorphism. Both give the same traced $J_2$ tower. Both admit $H$: crossing scalar $r_+ = 1$; $r_- = 1$ except $r_-(1, 1) = i$, whose only nontrivial tensor identity is $1 = -i^2$. |
| E-H: untwisted $C_2$, $O = \mathbf{1} \oplus g$ | $t^{\pm}(a, b) = (\pm 1)^{ab}$ swap. Bilinearity gives $H$; a pointed block-scalar gauge cannot change the $g, g$ sign. The $(g, g, \mathbf{1}, \mathbf{1}) \to (\mathbf{1}, g, g, \mathbf{1})$ entry of $\alpha_2^{\pm}(h)$ is $\pm 1$. Appending two vacancies gives the same $+1/-1$ mismatch for $\alpha_2^+ \otimes 1^2$ versus $\alpha_3^-$ in 97.14. |

## 98.9 Single Next Target: Intermediate-Record Faithfulness

**Target 97.16 [conjectural; any, open; L0+$H$; R0 information only] (NOTE:636–644,811–827; REVIEW:196–214).** For arbitrary pointed tensor-generating inputs $(\mathcal{D}, O, \eta, H)$, retain exactly $\mathcal{R} = (A_\infty, \{A_{[j,k]}\}_{1 \le j \le k}, \delta)$, where $A_{[j,k]} = \kappa_k(1_{W_{j-1}} \otimes \mathrm{End}_{\mathcal{C}}(W_{k-j+1}))$, extending (as). Record isomorphisms must carry each named interval algebra onto its counterpart and intertwine $\delta$; no point, corner, $h, i, j$ or concatenation marks remain. Prove that record isomorphism forces a unitary tensor equivalence transporting $\eta, H$, by recovering 97.13's missing marks up to that gauge, or give an explicit counterexample preserving *all* these constraints simultaneously. Neither 97.12 nor 97.13 answers this question.

# 99 Canonical Seams, Reflection Defects, and Conditional Spatial Positivity

## 99.1 Status, Sources, and Dependencies

**Status: WEAKER-RESULT-PROVED; grades carried per statement.** Statements 98.1–98.5 and 98.8 are *derived*; 98.6 and 98.9 are *negative*; 98.7 and the categorical application of 98.8 are *conditional*; Target 98.10 is *conjectural*. The explicit OS forms in 98.9 carry the reviewer's *derived* repair; the unrestricted post-OS claim has a GAP and is not lifted. NOTE, REVIEW and INTEGRATION below mean the files in `explorations/2026-10-25-astra-r3/F3-reflection-coherence/`. **CA-88 cell change (INTEGRATION:47–65):** split the CA-92 data/comment cell: the individual algebraic seam and bent-kernel/Schur construction are now *(any;* L1*; finite →* R3*)*, with $s_{\mathrm{can}}$ determined; 98.4 also supplies an L0+$\psi$ branch. Whole-history RP remains *(full;* L0*+coherent reflected pairing; finite →* R3*) [conditional]*, including at L1. Finite OS normalization is category-free. No cell moves up a target rung; R0–R2, R4–R5, CA-90 and CA-80's temporal conclusions are unchanged. Dependencies: CA-80 (80.1–80.5), CA-86 (linear history), CA-88 (tags/grid), CA-91 (linear parity distinction), CA-92 (92.1–92.9); CONVENTIONS (b), (r), (ak), (al), (am), (aq), (at), (au). Checks in the exploration's `checks/`: `check_core.py`, `check_pointed.py`, `check_nonpointed.py`, `review_duality.py`, `review_history.py`, `review_hexagons.py`; six normal cases reproduce retained outputs, five targeted mutations fail. These verify finite identities, not automatic coherence or a continuum limit (REVIEW:210–238).

## 99.2 Algebraic Seams and Oriented Cut Spaces

Work with any unitary fusion $\mathcal{C}$, nonzero $O$, specified unitary spherical duals and all charge multiplicities. Write $D(X) = X^\vee$, $D(f) = (f^\vee)^\dagger$, $j_X : X \to D^2X$; its unitary tensorator reverses word order. The supplied braiding is $c_{X,Y} : X \otimes Y \to Y \otimes X$. These are shard-local declarations, proposed in `SHARD_META.md`; (au)'s seam convention is distinct from (at)'s linear parity. Put $u = \mathrm{coev}_O$, $v = \mathrm{ev}_O^\dagger$, $W_+ = O \otimes DO$, $W_- = DO \otimes O$. An *algebraic seam* is a unitary $s : W_+ \to W_-$ with $su = v$, fixed before $h$; it does not include (RH). Keep coherent associators in every displayed composition; [P] denotes Penneys at the anchors above.

**Theorem 98.1 [derived; any; L1; finite → R3] (NOTE:50–152; REVIEW:56–65).** For isotypic projections $p_a$ of $O$, the following determines a unitary algebraic seam, natural under unitary braided equivalences with their unitary spherical-dual comparisons:

$$\lambda_a = d_a^{-1}\mathrm{ev}_a c_{a,Da}\mathrm{coev}_a, \qquad t_O = \sum_a \lambda_a^{-1} p_a, \qquad s_{\mathrm{can}} = c_{O,DO}(t_O \otimes 1_{DO}). \tag{14}$$

⟨1⟩1. Rigidity gives $\dim \mathrm{Hom}(\mathbf{1}, Da \otimes a) = 1$; both $c\,\mathrm{coev}_a$ and $\mathrm{ev}_a^\dagger$ have norm $\sqrt{d_a}$. Hence $c\,\mathrm{coev}_a = \lambda_a \mathrm{ev}_a^\dagger$, $|\lambda_a| = 1$; applying $\mathrm{ev}_a$ gives the formula. BY [P]:425–430,1012–1030.

⟨1⟩2. $t_O$ is central unitary and natural in categorical morphisms. For orthogonal simple-copy inclusions $i_{a\mu}$, put $k_{a\mu} = (i_{a\mu}^\vee)^\dagger$. Direct-sum duality gives $u = \sum (i_{a\mu} \otimes k_{a\mu})\mathrm{coev}_a$, $v = \sum (k_{a\mu} \otimes i_{a\mu})\mathrm{ev}_a^\dagger$. Braiding naturality and step 1 give $s_{\mathrm{can}} u = v$. BY [P]:219–237,1029–1030; products of unitaries give unitarity.

⟨1⟩3. Currying $f \mapsto (f \otimes 1)u$ is an isomorphism by the snakes, so the cup equation uniquely determines $t_O$ *within this construction*, not all seams. Transporting $d_a, \lambda_a, p_a$ under a unitary braided equivalence proves naturality. BY steps 1–2 and [P]:1017–1030.

⟨1⟩4. Inverting currying gives $t_O = (1_O \otimes \mathrm{ev}_O)(c_{O,DO}^{-1} \otimes 1_O)(v \otimes 1_O)$. The pivotal cup identity is $\mathrm{coev}_{DO} = (1_{DO} \otimes j_O)v$ ([P]:1004). Thus the typed Drinfeld map $u_O = (\mathrm{ev}_O \otimes 1_{D^2O})(c_{O,DO} \otimes$

$1_{D^2O})(1_O \otimes \mathrm{coev}_{DO})$ equals $j_O t_O^\dagger$, giving $t_O = u_O^{-1} j_O$. QED. For simple self-dual $a$, keep the oriented cup sign: $\lambda_a = \nu_a R_1^{aa}$, $t_a = \nu_a (R_1^{aa})^{-1} = \theta_a$ in RSW:2311–2314's ribbon convention, *not* $\theta_a^{-1}$.

**Theorem 98.2 [derived; any; L0; finite → R3] (NOTE:215–259; REVIEW:71–74).** $D$ gives antiunitaries $\mathrm{Hom}(a, W) \to \mathrm{Hom}(Da, DW)$, respects inverse-Gram leg sewing, and $D^2(f) = j_W f j_a^\dagger$. Both labels change; this is a mirror between oriented cuts.

⟨1⟩1. Dual and dagger each reverse composition, so $D(gf) = D(g)D(f)$, $D(f^\dagger) = D(f)^\dagger$; only dagger conjugates scalars. BY [P]:682–708,992–1008.

⟨1⟩2. For simple $a$, $D(f)^\dagger D(g) = D(f^\dagger g) = \overline{\langle f, g\rangle} 1_{Da}$; dual equivalence gives surjectivity.

⟨1⟩3. The cut Gram $G$ and its inverse become $\overline{G}, \overline{G^{-1}}$. Apply $D$ to CA-80's identity resolution and compositions: leg sewing is preserved. Pivotal naturality gives the displayed double-dual formula; compare both labels back by $j$. QED. Braiding supplies no $O \cong DO$.

**Proposition 98.3 [derived; any; L0+identifications/algebra; finite] (NOTE:261–308; REVIEW:75–78).** Declare unitary $r_X : X \to DX$, $\rho_X(A) = r_X^\dagger D(A) r_X$, $z_X = r_X^\dagger D(r_X^\dagger) j_X$. Then $\rho_X^2 = \mathrm{Ad}_{z_X}$, identity on $\mathrm{End}_{\mathcal{C}}(X)$ iff $z_X$ is central; for a smaller named algebra use its commutant. On identified self-dual simple-charge cuts, $J_{a,W}(f) = r_W^\dagger D(f) r_a$ has $J_{a,W}^2(f) = z_W f z_a^\dagger$.

⟨1⟩1. Substitute 98.2's $D^2$ into $\rho_X^2$ and $J^2$; all factors of $z_X$ are unitary. Commutation with every observable is precisely the stated iff.

⟨1⟩2. For simple self-dual $X$, $r_X^\vee j_X = \nu_X r_X$, $\nu_X = \pm 1$ ([P]:917–933), so $z_X = \nu_X 1$. Where both defects are scalar signs, the cut square is the *ratio* $\nu_W/\nu_a$; even a pseudoreal $a$ gives $J^2 = +1$ on $\mathrm{Hom}(a, a)$. QED. Every admitted word $W$ needs its own defect check. Non-self-dual external charges require paired comparisons, not an invented $r_a$; centrality alone supplies no seam-Hermiticity identity.

**Proposition 98.4 [derived; any; L0 + ψ; finite → R3] (NOTE:310–329; REVIEW: 79–81).** Any declared unitary $\psi : O \to DO$ gives the algebraic seam $s_\psi = \psi \otimes [j_O^\dagger D(\psi)]$.

⟨1⟩1. Each factor is unitary.

⟨1⟩2. The dual-mate identity gives $(\psi \otimes (\psi^\vee)^{-1})u = \mathrm{coev}_{DO}$; unitarity gives $(\psi^\vee)^{-1} = D(\psi)$. Apply $1 \otimes j_O^\dagger$ and 98.1's pivotal cup identity to obtain $s_\psi u = v$. QED. On the unit-multiplicity corner $s_\psi = U \otimes \overline{U}$; $\psi = 1$ gives the factorwise identity, while $s_{\mathrm{can}}$ is the flip. Neither seam identity asserts a central square defect.

**Lemma 98.5 [derived; any; L0+$s$, L1 for $s_{\mathrm{can}}$; finite → R3] (NOTE:331–363; REVIEW: 82–83).** With $K_s(T) = s^\dagger \,\mathrm{rot}(T)$, the map $\sharp_s = K_s^{-1} \circ \dagger \circ K_s$ is a conjugate-linear involution and $K_s(h) = K_s(h)^\dagger$ iff $\sharp_s(h) = h$. The inverse is of the *linear map*, not a kernel matrix.

⟨1⟩1. The typed bend and inverse are

$$\mathrm{rot}(T) = (1_{DO} \otimes 1_O \otimes u^\dagger)(1_{DO} \otimes T \otimes 1_{DO})(v \otimes 1_O \otimes 1_{DO}),$$
$$\mathrm{rot}^{-1}(r) = (u^\dagger \otimes 1_O \otimes 1_O)(1_O \otimes r \otimes 1_O)(1_O \otimes 1_O \otimes v).$$

Both inverse composites cancel by the dagger snakes ([P]:398–430); hence $K_s$ is a linear bijection.

⟨1⟩2. Transport dagger across that bijection. QED. No multiplicative variance or geometric reflection follows: even in the same multiplicity corner the identity and flip seams give different operations.

## 99.3 Reflection Obstructions

**Negative 98.6 [negative; any; L0, L1 for the braided witness; finite] (NOTE:154–212,365–384,648–649; REVIEW:66–70,84–86).** Arbitrary unitary self-duality need not give an observable involution; conjugate-dual invariance is not canonical-seam Hermiticity. A same-tile-algebra

reflection with the Hermiticity iff for *all complex tiles* need not exist with either ordinary composition variance.

⟨1⟩1. In every $\mathcal{C}$, take $O = \mathbf{1} \oplus \mathbf{1}$, standard duality, coordinate conjugation $K_0$,

$$\psi = \frac{1}{5}\begin{pmatrix} 3 & -4 \\ 4 & 3 \end{pmatrix}, \qquad J = \psi^\dagger K_0, \qquad J^2 = \frac{1}{25}\begin{pmatrix} -7 & 24 \\ -24 & -7 \end{pmatrix}.$$

The square does not commute with $Z = \mathrm{diag}(1, -1)$. The pointed variant $O = \mathbf{1}^{\oplus 3}$, $\psi' = 1 \oplus \psi$, fixes the first vacuum inclusion and retains this defect.

⟨1⟩2. With $\psi = 1$, factor swap $P$ and $Y = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix}\right)$, $D_2(h) = P\overline{h}P$, whereas $\sharp_{\mathrm{flip}}(h) = Ph^\dagger P$ (92.7). For $h = Y \otimes Z - Z \otimes Y = h^\dagger$, both conjugation and swap negate $h$, so $D_2(h) = h$, but $K_{\mathrm{flip}}(h) = |\mathrm{vec}Y\rangle\langle\mathrm{vec}Z| - |\mathrm{vec}Z\rangle\langle\mathrm{vec}Y|$ is nonzero anti-Hermitian. Conversely $g = Y \otimes 1 + 1 \otimes Y = g^\dagger$ has Hermitian kernel but $D_2(g) = -g$.

⟨1⟩3. If a conjugate-linear involution $R$ has exactly the fixed tiles of $\sharp_s$, write $h = x + iy$, $x = (h + \sharp_s h)/2$, $y = (h - \sharp_s h)/(2i)$. Then $R(h) = x - iy = \sharp_s h$. The braided instance in the table has $\sharp(AB) \neq \sharp(A)\sharp(B), \sharp(B)\sharp(A)$; neither variance works. QED. This does not exclude graph-dependent port/composition changes or a restricted tile grammar.

## 99.4 Conditional Histories and Finite Transfer Operators

**Proposition 98.7 [conditional; full (chiral only with supplied twist); L0+G1–G3; finite → R3] (NOTE:386–422; REVIEW:87–88). Assume** a reflection-symmetric CA-86 linear brick patch with disjoint crossing tiles and declared boundary/flux data. G1: reflected half-amplitudes, read on the cut, are the original adjoints (RH). G2: crossing identifications tensor/reassociate to that *same* cut identification; charge restriction uses CA-80's inverse-Gram resolution, and the reverse identification is adjoint. G3: crossing tiles $1 + th$ use fixed $h$, independent of $N, M$, or a family with a common step bound; extra seam insertions lie inside a positive reflected factorization; bulk and outer boundary obey G1. The local certificate, with $e = u/\sqrt{d_O}$, $Q = 1 - |e\rangle\langle e|$, $A = K_s(h)$, $a = \langle e, Ae\rangle$, $b = QAe$, $C = QAQ|_{Q\mathcal{H}(W_+)}$, is

$$A = A^\dagger, \qquad C \geq 0, \qquad b \in \mathrm{Ran}\, C, \qquad d_O + t(a - b^\dagger C^+ b) \geq 0,$$

where $C^+$ is the inverse on its range and $t > 0$. **Prove** all-insertion spatial RP for every admissible patch, with this same step interval independent of its brick count.

⟨1⟩1. Theorem 92.5 gives positive crossing kernels from the certificate.

⟨1⟩2. Proposition 92.4 gives the reflected compression $V^\dagger(\bigotimes_i K_{s,i})V \geq 0$. QED. This is its conditional application. Neither 98.1 nor 98.2 proves G1/G2 for the canonical seam; the automatic-coherence GAP remains at L1.

**Lemma 98.8 [derived; any; category-free; finite → R3] (NOTE:424–454; REVIEW: 89–91). Assume** finite $V$, nonzero PSD Hermitian $q_0$, a shift $\tau : V \to V$, and PSD $q_1(f, g) = q_0(f, \tau g)$. **Prove** $T[f] = [\tau f]$ is positive on $V/\operatorname{rad} q_0$ and $T/\varrho$ is contractive for $\varrho \geq \max(1, \|T\|)$, without assuming $q_1 \leq q_0$.

⟨1⟩1. If $z \in \operatorname{rad} q_0$, then $q_1(z, f) = 0$. Hermiticity gives $q_1(f, z) = q_0(f, \tau z) = 0$, so $\tau z$ is null.

⟨1⟩2. On the quotient $\langle [f], T[g]\rangle = q_1(f, g)$, so $T = T^\dagger \geq 0$. Divide by $\varrho$: $0 \leq q_1/\varrho \leq q_0$, and Lemma 92.8 applies. QED. A logarithm on the whole quotient requires injectivity; otherwise restrict to positive spectral support, which does not preserve arbitrary zero-time probes. No uniform normalization or zero-energy vacuum follows. The categorical application remains *conditional*: shift closure and a positive compatible translated form must still be constructed from the same history.

**Negative 98.9 [negative; full finite torus; L0, already symmetric L1; finite] (NOTE: 456–502; REVIEW:92–140).** For every $\mathcal{C}$, $O = \mathbf{1}^{\oplus d}$, $d \geq 2$, and $h = 0$, universal unitary conjugacy of the complete temporal/spatial transfers fails, including the explicit even-torus OS forms below.

⟨1⟩1. On $N, M \geq 2$ periodic sites/slices, identity strands give $T_t = 1$ on $(\mathbb{C}^d)^{\otimes N}$. For a column $\sigma = (\sigma_1, \ldots, \sigma_M)$, let $w(\sigma) = \prod_r \delta_{\sigma_r, \sigma_{r+1}}$, $\sigma_{M+1} = \sigma_1$, and $v_\sigma = w(\sigma)$. The symmetric column transfer is $T_x = vv^\dagger$, $v^\dagger v = d$; $\operatorname{Tr} T_t^M = \operatorname{Tr} T_x^N = d^N$. Their ranks are $d^N$ and 1, even when $N = M$ makes dimensions equal.

⟨1⟩2. *Reviewer repair [derived, same tags; REVIEW:116–140].* Take even $N, M$, unnormalised torus weight $w(\sigma) = \prod_{x,r} \delta_{\sigma_{x,r}, \sigma_{x,r+1}}$, all indicator functions of half histories, and reflections $x \mapsto N - 1 - x$, $r \mapsto M - 1 - r$. For spatial half histories $a, b$, let $c(a) = 1$ iff each column is constant in time, else 0. Direct contraction gives $q_{0,x}(a, b) = c(a)c(b)$, rank 1. Reading the second temporal half in reflected time order gives $q_{0,t}(a, b) = \delta_{ab} c_t(a)$, where $c_t$ enforces constant columns in that half; exactly $d^N$ survive. Both forms are PSD, so their OS quotient dimensions are 1 and $d^N$. QED. This uses the displayed forms, not an inference from transfer rank to OS radical; other forms are not covered. At $d = N = M = 2$, $T_t = I_4$, $T_x = vv^\dagger$, $v = (1, 0, 0, 1)^T$, spectra $\{1, 1, 1, 1\}$, $\{2, 0, 0, 0\}$, contraction 4. All local Schur data vanish. The obstruction concerns cut-carrier intertwiners, including nonzero scalar normalizations; Hilbert–Schmidt unitary reshuffling of matrix entries is not refuted (REVIEW:100–108).

### 99.5 Effect on other shards

*Amendment pointers (2026-10-25, round 3; INTEGRATION:67–101).* CA-92's 92.2 gap closes only for the individual seam (98.1); 92.7 and its coherence proposal require the distinctions in 98.2–98.6 and Target 98.10. Lemma 92.8 gains finite normalization (98.8); 92.9 gains the carrier-level whole-history obstruction with explicit OS forms (98.9). CA-80's coverage paragraph and CA-75 Link (a2) gain an algebraic seam and dual-oriented leg comparison, not a general spatial-sewing or anomaly-line limit. CA-91's linear parity is not replaced by this mirror. CA-88 receives the split stated above, preserving E3's historical grade; CA-86's theorems need no alteration.

### 99.6 What This Does Not Claim

No automatic G1/G2, positive translated strip form, uniform OS normalization or exact-exponential RP is proved. The general post-OS claim without specified forms remains conditional; 98.9 proves only the reviewer repair. No new tower, Wilson state, KS modes, conformal limit, fidelity or functorial pipeline reaches R0–R5; seam naturality alone is not R5 (NOTE:576–619; REVIEW: 142–170). The chiral variant requires its own twisted cuts, boundary comparisons and translated forms.

### 99.7 Examples (Instances Only)

All rows are *derived finite calibrations*; tags and exact inputs are local to the row (NOTE:641–650; REVIEW:109–114,172–190). Pointed labels are $0, 1, 2$ modulo 3, $\zeta = e^{2\pi i/3}$, $O = \mathbf{1} \oplus g$; the associator arrow is $a(bc) \to (ab)c = \omega(a, b, c)$, not its inverse.

| Instance / tags | Exact calibration or falsifier | Check output |
|---|---|---|
| $\mathrm{Vec}_{\mathbb{Z}_3}^{\omega_p}$; any; L0; finite | $\omega_p(a,b,c) = \zeta^{pa\lfloor (b+c)/3 \rfloor}$, $p = 1, 2$. With $x = r(1,1)$, the hexagons give $1 = \zeta^p x^3 = x^3/\zeta^p$, impossible. These nonzero classes are not braided tests; no FS sign for $g$. | pointed:2–17; hexagons:2–7 |
| Braided $\mathbb{Z}_3$; any; L1; finite | $\omega = \zeta^{abc}$, $\beta(a,b) = \zeta^{ab^2}$, $r(a,b) = \zeta^{ba^2-ab^2+ab}$ give the nonconstant coboundary test. Also for $\omega = 1, r = \zeta^{ab}$, in basis $(00, 01, 10, 11)$: $A = E_{01,10}$, $B = E_{10,01}$, $AB = P_{01}$; $\sharp(A) = A$, $\sharp(B) = \zeta^2 B$, $\sharp(AB) = P_{10}$, violating both variances. | pointed:18–40; duality:13–20 |
| $\mathrm{Rep}^\dagger(Q_8)$; any; symmetric L1; finite | $Q_8 = \langle i\,\mathrm{diag}(1,-1), U\rangle$, $U = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$; its irreducible $V = \mathbb{C}^2$ has $\psi = U, \nu_V = -1$, yet $J^2 = 1$ on $\mathrm{Hom}(V,V)$. On connected $O = \mathbf{1} \oplus V$, $b = \lvert 00\rangle$, $v = \lvert 12\rangle - \lvert 21\rangle$, $h = i(bv^\dagger - vb^\dagger)$ is categorical, self-adjoint, $D_2$-fixed, and $K_{s_{\mathrm{can}}}(h)$ nonzero anti-Hermitian. | nonpointed: 2–18 |

Output abbreviations mean the corresponding `checks/*_output.txt` files anchored above; *hexagons* and *duality* mean `review_hexagons` and `review_duality`.

## 99.8 Next Target: Real-Self-Duality Two-Tile Reflection

**Target 98.10 [conjectural; any algebraically/full for RP; L0+declared real self-duality; finite → R3] (NOTE:623–632; REVIEW:192–208).** Assume unitary $\psi$ with $\psi^\vee j_O = \psi$, use $s_\psi$ from 98.4, and construct word identifications using $\psi$ and the unitary anti-monoidal tensorator, displaying associators and external-charge comparisons. First prove $K_{s_\psi}(\rho_2(h)) = K_{s_\psi}(h)^\dagger$ for every categorical two-site $h$. Then prove equality of the induced cut identification for the two sewing orders of adjacent bricks, using the actual inverse Gram and reflected boundary maps. State the remaining generating moves needed for induction. The real-structure assumption and changed seam avoid the known obstructions; neither proves this target. Positive translated forms and continuum reconstruction remain later tasks (INTEGRATION:145–160).

# 100 Signed Contact Classification on the Pointed Carrier: Robin Limits, Dirichlet Limits and Full-Fock Boundaries

## 100.1 Status, Sources, and Dependencies

**Status: COUNTEREXAMPLE to F5 as posed; PROOF-COMPLETE exact-family replacement** (REVIEW:1–9). Statements 99.1–99.6, 99.8–99.9 are *derived*; 99.7, 99.10–99.11 are *negative*; 99.12 and the open Target 99.13 are *conditional*. Global $H^2$ regularity is *conditional*; independent renormalized triple-cluster dynamics is *conjectural*. **CA-88 cells (INTEGRATION:72–91):** the CA-94 row's comment and D1-C collision-sewing discussion gain the restricted classification. Their setting *any (open)*, data $\mathsf{L0}$ plus declarations, and rung $\mathsf{R0}^c$ plus off-ladder $z = 2$ stay unchanged. Neither the restated $\mathsf{R0}^u + \mathsf{R1}$ target, its dressed-refinement lemma, nor any $\mathsf{R1}$–$\mathsf{R5}$ certificate advances. No chiral/full target is selected.

NOTE, REVIEW and INTEGRATION below mean the files in `explorations/2026-10-25-astra-r3/F5-contact-trichotomy/`; their line anchors are part of each statement. Dependencies: CA-77 (77.8 carrier), CA-88 (grid), CA-94 (94.2–94.6, 94.10–94.14); CONVENTIONS (am), (aq), (aw). New local notation is proposed for (ax) in the accompanying `SHARD_META.md`; existing collision conventions are retained. No applicable contact-classification theorem was identified in the recorded local search (REVIEW:268–270); the analytic claims below use the reviewed local derivations, without a novelty claim.

## 100.2 Exact Family and Classification

Fix unitary fusion $\mathcal{C}$, pointed $O = \mathbf{1} \oplus X$, $X \neq 0$, and $t, \mu > 0$. Write $E_{c,n} = \mathrm{Hom}(c, X^{\otimes n})$, $\Delta_n = \{0 < x_1 < \cdots < x_n < 1\}$, $\mathcal{F} = \widehat{\bigoplus}_{c,n} L^2(\Delta_n; E_{c,n})$, with isometry inner products, no factorial and $\Delta_0$ a point. Use CA-77's isometric box maps $K_\varepsilon$, $\varepsilon = 1/L$; smooth samples have amplitudes $\varepsilon^{n/2} u(x_{j_1}, \dots, x_{j_n})$, $x_j = (j - \frac{1}{2})\varepsilon$. Declare finite fixed lists $b_\rho : X \to X^{\otimes 2}$, $v_\sigma : \mathbf{1} \to X^{\otimes 2}$, $\lambda_\rho, \kappa_\sigma \geq 0$, and moments $a = \sum \lambda_\rho + \sum \kappa_\sigma$, $\beta = \sum \lambda_\rho b_\rho$, $\nu = \sum \kappa_\sigma v_\sigma$, $Q_b = \sum \lambda_\rho b_\rho^\dagger b_\rho$, $\gamma = \sum \kappa_\sigma v_\sigma^\dagger v_\sigma$. With $\pi_{00}, \pi_{X0}, \pi_{0X}, \pi_{XX}$ the summand coisometries of $O^{\otimes 2}$ (0 denotes $\mathbf{1}$), set $D_h = \pi_{X0} - \pi_{0X}$, $D_b = \pi_{XX} - \sqrt{\varepsilon} b \pi_{X0}$, $D_v = \pi_{XX} - \varepsilon v \pi_{00}$, $P_{XX} = \pi_{XX}^\dagger \pi_{XX}$. The *exact family* is

$$H_\varepsilon = \sum_{j=1}^{L-1} \left[ t\varepsilon^{-2} D_h^\dagger D_h + \sum_\rho \lambda_\rho \varepsilon^{-1} D_{b_\rho}^\dagger D_{b_\rho} + \sum_\sigma \kappa_\sigma \varepsilon^{-1} D_{v_\sigma}^\dagger D_{v_\sigma} + (\alpha_\varepsilon - a)\varepsilon^{-1} P_{XX} \right]_j + \mu \hat{N}. \quad (15)$$

Here $\alpha_\varepsilon = (c_\varepsilon + 2t)/\varepsilon$ is the *total* contact; the added excess is $\alpha_\varepsilon - a$. Kinetic/contact action is fibre-scalar, with the displayed open endpoint completion and no outer penalty. Use $H_\varepsilon^{[2]} = P_{\leq 2} H_\varepsilon P_{\leq 2}|_{\mathrm{Ran}\, P_{\leq 2}}$ on $\mathcal{F}_{\leq 2} = P_{\leq 2}\mathcal{F}$.

**Theorem 99.1 [derived; any (open); $\mathsf{L0}$+declared family; $\mathsf{R0}^c$+off-ladder $z = 2$] (NOTE:746–800; REVIEW:126–129,147–163).** For (15), $K_\varepsilon (H_\varepsilon^{[2]} - i)^{-1} K_\varepsilon^*$ converges strongly to a self-adjoint resolvent on the entire $\mathcal{F}_{\leq 2}$ iff either $\alpha_\varepsilon \to \alpha \in \mathbb{R}$, giving $H_\alpha$ below, or $|\alpha_\varepsilon| \to \infty$, giving $H_D^W$ below. Arbitrary sign oscillation is allowed.

$\langle 1 \rangle 1$. Sufficiency: 99.4 and 99.6. Necessity: rigidity gives $X^2 \neq 0$, since the unit summand of $X^* \otimes X$ would otherwise give $X = 0$. Thus finite-contact forms differ by $(\alpha - \alpha')\|Tu_2\|^2 \neq 0$ on a suitable smooth vector, and the Dirichlet form has a proper smaller domain.

$\langle 1 \rangle 2$. A sufficiently shifted resolvent $R = (H + z)^{-1}$ recovers its form, including its domain, by $q(u) + z\|u\|^2 = \sup_f \{2\,\mathrm{Re}\langle u, f\rangle - \langle f, Rf\rangle\}$, with value $+\infty$ off-domain (complete the square on the form-dense range of $R$; NOTE:774–788). The limiting operators are distinct.

⟨1⟩3. Two finite clusterpoints, or a finite and an absolute-divergent subsequence, contradict uniqueness of the strong limit. No finite clusterpoint means $|\alpha_\varepsilon| \to \infty$; otherwise the unique finite clusterpoint is the limit. QED. The contact-invisible case $X = 0$ is excluded only from necessity.

**Proposition 99.2 [derived; any (open); L0+declared family; finite, → $z = 2$] (NOTE:66–128; REVIEW:42–45,177–180).** Compression retains lower squares whose reaction outputs have $n > 2$. Put $\mathcal{V} = \bigoplus_{c,n\le 2} H^1(\Delta_n; E_{c,n})$, with $H^1(\Delta_0) = E_{c,0}$, and $S_c(u)(y) = B_{\beta,1,1}u_{c,1}(y) + V_{\nu,0,0}u_{c,0}$, with the insertions of (aw), abbreviated $B_\beta, V_\nu$. Then the smooth-sample limit for finite $\alpha$ is

$$q_\alpha(u) = t\sum_{c,n\le 2}\|\nabla u_{c,n}\|^2 + \sum_{c,n\le 2}\langle u_{c,n}, W_n u_{c,n}\rangle + \alpha\sum_c \|Tu_{c,2}\|^2 - 2\,\mathrm{Re}\sum_c \langle Tu_{c,2}, S_c(u)\rangle, \tag{16}$$
$$W_0 = \gamma, \qquad W_1 = (\mu+\gamma)I + Q_b, \qquad W_2 = (2\mu+\gamma)I + (Q_b)_1 + (Q_b)_2.$$

All trace pairings use $dy = dS/\sqrt{2}$; $D = \partial_1 - \partial_2$. Endomorphisms act by coherent postcomposition on Hom fibres, without a tensor-product factorization of the physical fibre.

⟨1⟩1. Expand the squares (94.2, 94.10): upper coefficient $a$ becomes $\alpha_\varepsilon$, cross maps are $\beta, \nu$, lower diagonals $Q_b, \varepsilon\gamma$. At fixed $n$, empty bonds number $L + O(n)$; missing $Q_b$ insertions occupy only adjacent/outer strips. Smooth recovery gives (16).

⟨1⟩2. Compression of the resolvent is different: for $H = \left(\begin{smallmatrix} 0 & g \\ g & 0 \end{smallmatrix}\right)$, $g \in \mathbb{R}\setminus\{0\}$, and first-coordinate $P$, $P(H - z)^{-1}P = -z/(z^2 - g^2)$, whereas $(PHP - z)^{-1} = -1/z$. Direct inversion proves the distinction. QED.

## 100.3 Finite Contacts and the Weak Generator

**Lemma 99.3 [derived; any; category-free, fixed finite fibre; → off-ladder $z = 2$] (NOTE:302–452; REVIEW:66–81).** For rescaled arrays $f_{jk} \in E$, $j < k$, let $N_\varepsilon = \varepsilon^2\sum\|f_{jk}\|^2$, $E_\varepsilon = \sum_{p\sim q}\|f_p - f_q\|^2$ over unordered legal one-step edges, and $C_\varepsilon = \varepsilon\sum_{j<L}\|f_{j,j+1}\|^2$. For every $\eta > 0$, $C_\varepsilon \le \eta E_\varepsilon + C_0(1+\eta^{-1})N_\varepsilon$, with $C_0$ mesh/fibre independent. Bounded $N_\varepsilon + E_\varepsilon$ gives a subsequence with strongly convergent bulk box steps and full collision steps $A_\varepsilon \to Tu$ in $L^2$, for $u \in H^1(\Delta_2; E)$. The fibre $E$ is fixed.

⟨1⟩1. Reflect to a square array, filling $g_{jj} = f_{j,j+1}$, $g_{LL} = f_{L-1,L}$. Two-edge routes give square mass $\le 4N_\varepsilon$, energy $\le 6E_\varepsilon$. Clamped affine interpolation $P_\varepsilon$ has mass $\le 8N_\varepsilon$, gradient energy $\le 6E_\varepsilon$, and square-step error $\le 72\varepsilon^2 E_\varepsilon$.

⟨1⟩2. Its diagonal trace differs from the filled diagonal step by squared norm $(\varepsilon/12)\sum\|g_{j+1,j+1} - g_{jj}\|^2 \le \varepsilon E_\varepsilon/6$. Apply 94.3's adjustable trace bound; for coarse meshes use $C_\varepsilon \le \varepsilon^{-1}N_\varepsilon$. This proves the displayed inequality.

⟨1⟩3. Cell-average approximations give bulk compactness. Apply the trace bound to differences, first taking the bulk limit, then shrinking the collar: strong full traces follow. Absolute continuity removes the last cell; omitted diagonal half-box mass is $\le \varepsilon C_\varepsilon$. QED, including both collision corners.

**Theorem 99.4 [derived; any (open); L0+declared family; R0$^c$+off-ladder $z = 2$] (NOTE:455–517; REVIEW:87–92).** If $\alpha_\varepsilon \to \alpha \in \mathbb{R}$, (16) on $\mathcal{V}$ is closed and semibounded, with self-adjoint operator $H_\alpha$, and the projected nonreal resolvents of $H_\varepsilon^{[2]}$ converge to its resolvent.

⟨1⟩1. Bounded sources and 99.3 absorb signed traces: $q_\varepsilon \ge (t/2)E_\varepsilon - C\|u\|^2$. The continuous counterpart gives a complete shifted $H^1$ form norm and the variational operator (94.4).

⟨1⟩2. Bounded shifted energies give strong bulk and full traces by 99.3, so signed and mixed terms converge. Missing $Q_b$ strips vanish by its outer-row corollary (NOTE:441–452); the $\gamma$ error is $O_n(\varepsilon)$. The interior difference-quotient proof of 94.5 gives kinetic liminf with coefficient $t$, not the interpolation factor six.

⟨1⟩3. Smooth recovery and density give recovery on $\mathcal{V}$. The shifted minimizer argument of 94.6 gives strong resolvent convergence; its bounded rational transform gives the nonreal version (NOTE:511–517). QED.

**Proposition 99.5 [derived; any (open); L0+declared moments; off-ladder $z = 2$] (NOTE:519–569; REVIEW:93–101).** Exactly $D(H_\alpha) = \{u \in \mathcal{V} : \exists f \in \mathcal{F}_{\le 2},\ q_\alpha(v,u) = \langle v, f\rangle\ \forall v \in \mathcal{V}\}$. For sufficiently regular vectors (charge sums implicit),

$$(H_\alpha u)_2 = -t\Delta u_2 + W_2 u_2, \quad (H_\alpha u)_1 = -tu_1'' + W_1 u_1 - B_\beta^* T u_2, \quad (H_\alpha u)_0 = \gamma u_0 - \int_0^1 V_\nu^* T u_2(y)\, dy.$$

The collision equation is $tDu_2 + \alpha T u_2 - B_\beta u_1 - V_\nu u_0 = 0$; outer fluxes are zero.

⟨1⟩1. Polarize (16) (second-slot-linear convention), then integrate by parts: $\partial_\nu = D/\sqrt{2}$ and $dS = \sqrt{2}dy$ cancel. Without regularity these equations mean the variational boundary functional.

⟨1⟩2. The homogeneous coefficient is $\alpha I_{E_{c,2}}$; reactions and $W_n$ need not be scalar. Robin coefficients are $\alpha/t$ for $D$, $\alpha/(\sqrt{2}t)$ for unit normal. Total $c_\varepsilon = -2t + \kappa\varepsilon + o(\varepsilon)$ gives $\alpha = \kappa$; an external coefficient $-2t + \delta\varepsilon + o(\varepsilon)$ before adding squares gives $\alpha = \delta + a$. Occupation is conserved iff $\beta = \nu = 0$, by the mixed-form argument of 94.7; individual reactions can cancel. QED.

## 100.4 Both Divergent Signs and the Refuted Branch

**Theorem 99.6 [derived; any (open); L0+declared family; R0$^c$+off-ladder $z = 2$] (NOTE:571–738; REVIEW:107–120).** For $|\alpha_\varepsilon| \to \infty$, the limit is $H_D^W$, associated with $q_D = t\sum \|\nabla u_n\|^2 + \sum\langle u_n, W_n u_n\rangle$ on $\mathcal{V}_D = \{u \in \mathcal{V} : Tu_2 = 0\}$. Sectors decouple, with all $W_n$ retained. No common lower bound or restriction on sign/rate is assumed.

⟨1⟩1. An operator core uses arbitrary $u_0$, finite Neumann cosine sums $u_1$, and smooth spatially odd $u_2$, outer-Neumann and zero near the diagonal corners. Cosine determinants are complete; they vanish as $O(\rho^2)$ at these corners. Radial cutoffs cost $O(d)$ in the Laplacian norm, proving graph-core density.

⟨1⟩2. For a fixed core vector set $s = (x+y)/2$, $r = y - x$, $g(s) = \partial_r u_2(s,0)$, $G(x,y) = g(s)\chi(r)$, with $\chi$ smooth/even, one near zero and supported so $G$ avoids the outer walls. Put $\overline{S}(x,y) = B_\beta[u_1(x) + u_1(y)]/2 + V_\nu u_0$. Sample $F_\varepsilon = u_2 + (2t/\alpha_\varepsilon - \varepsilon)G + \overline{S}/\alpha_\varepsilon$, but overwrite each adjacent rescaled sample by $a_j = [2tg(j\varepsilon) + S(u)(j\varepsilon)]/\alpha_\varepsilon$. The overwrite error is $O(\varepsilon^3 + \varepsilon^2/|\alpha_\varepsilon|)$.

⟨1⟩3. On interior adjacent rows, $2a_j - F_{\text{left}} - F_{\text{right}} = -2\varepsilon g + O(\varepsilon^3 + \varepsilon^2/|\alpha_\varepsilon|)$. Thus kinetic/contact/source action is $[S(u)(s) - S_\varepsilon(s)]/\varepsilon + O(\varepsilon + |\alpha_\varepsilon|^{-1}) = O(1)$, where $S_\varepsilon(s) = B_\beta u_1(s - \varepsilon/2) + V_\nu u_0$. Exact overwriting cancels every growing contact term.

⟨1⟩4. Degree-one endpoint rows use $u_2 = g = 0$ and $u_1'(0) = u_1'(1) = 0$. Other rows have Neumann difference consistency. Feedback $-B_\beta^* a_j$, $-\varepsilon\sum V_\nu^* a_j$ is $O(|\alpha_\varepsilon|^{-1})$. Counting strip area gives $\|K_\varepsilon v_\varepsilon - u\| + \|K_\varepsilon H_\varepsilon^{[2]} v_\varepsilon - H_D^W u\| \le C_u(\sqrt{\varepsilon} + |\alpha_\varepsilon|^{-1}) \to 0$.

⟨1⟩5. For $f = (H_D^W - i)u$, the resolvent norm $\le 1$ bounds the convergence error by this graph error. Such $f$ are dense by the operator core; the same bound extends convergence to every input. QED.

**Negative 99.7 [negative; any (open); L0+declared contact; off-ladder $z = 2$] (NOTE:740–744; REVIEW:171–174).** Take $c_\varepsilon = -6t$, so $\alpha_\varepsilon = -4t/\varepsilon$. Theorem 99.6 gives Dirichlet even with reactions. Without reactions, the endpoint adjacent-pair basis vector has energy $-3t\varepsilon^{-2} + 2\mu \to -\infty$. This refutes F5(a)/Target 94.16: spectral diving and failure of a self-adjoint resolvent limit are different assertions.

## 100.5 Full-Fock Results and General-Density Boundaries

**Theorem 99.8 [derived; any (open); L0+nonreacting data; R0$^c$+off-ladder $z=2$] (NOTE:139–261,852–859; REVIEW:46–57,165–167).** With no reactions, $H_\varepsilon = t\varepsilon^{-2}\sum D_h^\dagger D_h + s_L\varepsilon^{-6}\sum n_j n_{j+1}$, $s_L \in \{-1,1\}$ arbitrary, converges on all $\mathcal{F}$ to collision-Dirichlet, outer-Neumann chamber Laplacians. Here $c_\varepsilon = -2t + s_L\varepsilon^{-4}$, $\alpha_\varepsilon = s_L\varepsilon^{-5}$; $s_L = -1$ gives Rayleigh quotient $t\varepsilon^{-2} - \varepsilon^{-6}$ at $(1,2)$, while $s_L = (-1)^L$ gives no signed contact limit.

⟨1⟩1. At fixed $n$, the exclusion kinetic matrix has norm $\le 4nt\varepsilon^{-2}$. Split off adjacent configurations $B$, with adjacency count $A \ge 1$. Its block resolvent has norm $\le 2\varepsilon^6$; Schur elimination changes the complementary resolvent by $\le 32n^2t^2\varepsilon^2$, with cross blocks $O_n(\varepsilon^4)$.

⟨1⟩2. On the complement, $l_i = j_i - (i-1)$, $m = L - n + 1$, identify the principal kinetic matrix with the exterior-power Neumann path sum. Cosine determinants have eigenvalues $\sum_a 4t\varepsilon^{-2}\sin^2(\pi p_a/(2m)) \to t\pi^2\sum_a p_a^2$, $0 \le p_1 < \cdots < p_n < m$. Their original-box limits form a complete chamber basis. This spatial computation imposes no categorical statistics.

⟨1⟩3. Fixed-sector convergence and resolvent norm $\le 1$ control Fock tails by density. QED. More generally $|\alpha_\varepsilon|\varepsilon^3 \to \infty$ suffices, with Schur error $32n^2t^2/(|\alpha_\varepsilon|\varepsilon^3)$; adding $\mu\hat{N}$ adds $\mu n$ sectorwise. No $n$-uniform estimate is used.

**Theorem 99.9 [derived; any (open); L0+positive family; R0$^c$+off-ladder $z=2$] (NOTE:804–850; REVIEW:130–132).** For uncompressed (15), eventual $\alpha_\varepsilon \ge a$ and either $\alpha_\varepsilon \to \alpha \in [a,\infty)$ or $\alpha_\varepsilon \to +\infty$ give full-Fock projected-resolvent convergence.

⟨1⟩1. The finite limit is 94.6's form, kinetic coefficient $t$, plus $(\alpha - a)\sum \|T_{n,i}u_n\|^2$. Positive finite exhaustion, closedness and the smooth core give liminf/recovery.

⟨1⟩2. Infinite contact forces every collision trace to zero, retaining $\mu n I + \sum_i (Q_b)_i + \gamma I$. At fixed $n$, zero all adjacent samples: the graph-energy cost is $O_n(\varepsilon)$, independently of contact growth. Finite-sector thresholds and diagonal recovery prove the claim (94.6). QED.

**Negative 99.10 [negative; any (open); L0+negative total contact; off-ladder stability obstruction] (NOTE:861–896; REVIEW:133–135).** For fixed total $\alpha < 0$, the formal full-Fock form has no common lower bound, even with fixed reactions.

⟨1⟩1. Choose real normalized $h \in C_c^\infty(-1/2,1/2)$, $A = \int |h'|^2$, $B = \int |h|^4$, $h_\ell(x) = \sqrt{\ell}h(\ell(x - 1/2))$, and a unit fibre vector $e_n$. The normalized pure-sector trial $u_n = \sqrt{n!}\prod_i h_\ell(x_i)e_n$ has kinetic energy $tn\ell^2 A$ and total trace norm squared $n(n-1)\ell B$.

⟨1⟩2. Mixed sources vanish; $\ell = |\alpha|(n-1)B/(2tA)$ gives $q_\alpha(u_n) \le -\alpha^2 B^2 n(n-1)^2/(4tA) + n(\mu + \|Q_b\|) + \gamma \to -\infty$. QED. This concerns negative *total* contact, not merely negative excess; it is not a no-resolvent theorem.

**Negative 99.11 [negative; any (open); L0+internal diagonals; off-ladder classification obstruction] (NOTE:263–300,907–959; REVIEW:58–60,136–138).** Arbitrary diagonals and $(G_r)$ alone give neither the scalar classification nor aligned internal bands.

⟨1⟩1. For any $Y \ne 0$, let $X = Y \oplus Y$, $p$ project onto its second copy, and $a_O = 0 \oplus p$ on $O$. Add $\varepsilon^{-2}(a_O \otimes I_O + I_O \otimes a_O)/2$ to scalar hopping with critical contact, no reactions. A scalar band-bottom subtraction leaves internal gap $\varepsilon^{-2}$. A higher-band packet $u_\varepsilon(j) = \sqrt{\varepsilon}f(x_j)e$, $0 \ne f \in C_c^\infty(0,1)$, satisfies $H_\varepsilon u_\varepsilon = \varepsilon^{-2}u_\varepsilon + w_\varepsilon$, $\sup\|w_\varepsilon\| < \infty$. Its resolvent response tends to zero on the nonzero limit $fe$, contradicting resolvent injectivity.

⟨1⟩2. Instead project $p$ onto one of the four $Y^2$ summands of $X^2$. Add $w\varepsilon^{-2}\pi_{XX}^\dagger p \pi_{XX}$, $w > 0$, to a nonreacting finite-contact family. The $p$ channel has contact $\alpha_\varepsilon + w/\varepsilon$, hence Dirichlet; its complement stays Robin inside each relevant fibre.

⟨1⟩3. If $Y$ satisfies $(G_r)$, so does $Y \oplus Y$ by matrix amplification of Hom spaces. The preceding diagonals remain allowed. $(G_r)$ forces occupation conservation (94.11), not these missing restrictions. QED.

**Proposition 99.12 [conditional; any (open); L0+range data; → off-ladder $z = 2$] (NOTE:938–948; REVIEW:138–142).** The positive range-$r$ argument extends only with nearest-neighbour graph control, bounded-length contact paths with bounded edge multiplicity, codimension-one traces, bounded sources, a closed core and smooth recovery (94.13). Then $\|A_\varepsilon - A'_\varepsilon\|^2 \le C_{n,r}\varepsilon E_{\varepsilon,n} \to 0$. An either-sign range classification additionally needs a boundary-row consistency argument like 99.6; none is proved here.

### 100.6 Effect on other shards

*Amendment (2026-10-25, round 3):* CA-94 Target 94.16 is refuted and replaced by 99.1; its full-Fock boundary is 99.8–99.10. CA-77's caveat and CA-88's D1-C discussion must distinguish a finite Robin window from existence of any limit. CA-88 gains only the restricted comment-column classification stated above; its round-2 E6 next-target pointer is superseded. CA-89–CA-93 receive no mathematical amendment (INTEGRATION:93–131); exact append-only text is in `SHARD_META.md`.

### 100.7 What This Does Not Claim

No arbitrary-density or signed reacting full-Fock classification, no unital refinement, WT-1 state, KS/Ward modes, conformality or fidelity follows. Global $H^2$ boundary regularity remains *conditional*; independent renormalized triple-cluster dynamics remains *conjectural* (both: any/open; L0+decl.; off-ladder $z = 2$; REVIEW:140–142, INTEGRATION:23,32–33). Reaching $n = 3$ forces no three-body counterterm: 94.14 concerns a separately added tile. Finite checks calibrate identities, not infinite convergence: the four reviewer outputs and `checks/review_manifest.json` record the independent trace and complex-source checks, and caught mutations (REVIEW:188–210). CA-94.5 uses no $n$-uniform constants.

### 100.8 Examples (Instances Only)

| Instance | Calibration of the general statements (grades/tags inherited) |
|---|---|
| Any $\mathcal{C}$, $X = 0$ | Vacuum only; contact invisible, excluded from 99.1's necessity (NOTE:799–800). |
| Any $\mathcal{C}$, no reactions | $c = -6t$: 99.7 [negative]; $\alpha = (-1)^L\varepsilon^{-5}$: 99.8 [derived]. |
| Any $Y \neq 0$, $X = Y \oplus Y$ | Unaligned internal bands or mixed Dirichlet/Robin channels: 99.11 [negative]. |
| $\mathrm{Hilb}_{\mathrm{fd}}$, $X = \mathbb{C}^2$ | At $L = 4$, the $33\times33$ compression agrees with an independent $81\times81$ chain oracle ([derived; any; L0+decl.; finite], NOTE:970–977; REVIEW:196). |

### 100.9 Single Next Target

**Target 99.13 [conditional; any (open); L0+signed reacting data; → off-ladder $z = 2$] (NOTE:1082–1083; REVIEW:214–223).** *Signed reacting full-Fock realization and graph-core lemma (open).* For (15), $X \neq 0$, finite total $\alpha < 0$, and $\beta \neq 0$ or $\nu \neq 0$, construct a self-adjoint full-Fock operator satisfying CA-94's local source equations, with a dense operator core $\mathcal{D}$ and approximants for every $u \in \mathcal{D}$ obeying $K_\varepsilon u_\varepsilon \to u$, $K_\varepsilon H_\varepsilon u_\varepsilon \to H_\alpha u$; alternatively exhibit a fixed nonzero source obstructing such a resolvent limit. Control occupation tails and feedback: the truncated corrections feed sectors three and four before compression. A finite-sector form core is not automatically an operator core; 99.10 rules out a common lower-bound proof.

# 101 Zero Modes and Lattice Chirality: Parity-Free KS Ledger Balance

## 101.1 Status, Sources, and Dependencies

**Status: WEAKER-RESULT-PROVED (REVIEW:1–11).** Grades are *derived* for 100.1, 100.7, 100.11; *conditional* for 100.2–100.4, 100.8, 100.10; *negative* (derived counterexamples) for 100.5, 100.6, 100.9; and *conjectural* for Target 100.12. Automatic ZC, the unrestricted physical no-go and sharp F2(c) remain open. NOTE, REVIEW and INTEGRATION below mean the files in `explorations/2026-10-25-astra-r3/F2-zero-mode-and-lattice-chirality/`. **CA-88 placement (INTEGRATION:62–94):** the (full; category-free; conditional R2) cell gains parity-free cubic balance and the single-ZC reduction; the (any; category-free; → R2) interface gains the transport tail. No cell moves up a rung: no new local R0, Wilson state R1, or R3–R5 certificate. Definition 88.1's general chirality source gap is narrowed, not closed. Dependencies: CA-79 (79.1, compression/leakage), CA-81 (VR-1), CA-88 (grid), CA-89 (graph calculus), CA-91 (91.5–91.7, 91.10), LC-4; CONVENTIONS (m), (s), (v), (w), (ap), (aq), (at). New notation is shard-scoped here, with proposed registration in the accompanying `SHARD_META.md`.

## 101.2 Common-System Hypotheses and Transport

At scale $r$, $N = N_r \to \infty$; declare a periodic representation of translated self-adjoint two-site densities $k_j = -h_j/v_r$, $v_r > 0$, with nonadjacent densities commuting. Indices are cyclic, $a_N = N/(4\pi)$, $z = e^{2\pi i/N}$, $\mathcal{H}_r = \sum_j k_j$, $\mathcal{J}_j = i[k_j, k_{j+1}]$, $\mathcal{Q}_r = \sum_j \mathcal{J}_j$. Reuse (ap), (at):

$$E_{r,n} = a_N \sum_j z^{nj}(k_j - \varepsilon_N) + b_N \delta_{n0}, \quad B_{r,n} = a_N \sum_j z^{nj} \mathcal{J}_j, \quad L_{r,n} = E_{r,n} - B_{r,n}, \quad S_{r,n} = \bar{T}_{r,-n} = E_{r,n} + B_{r,n}.$$

$\varepsilon_N, b_N$ are real; $D_r = L_{r,0} - S_{r,0} = -2B_{r,0}$. No categorical identity enters; for any unitary fusion $\mathcal{C}$, a categorical realization needs L0 and the declared periodic action. **Common A1–A4** means finite Hilbert carriers $H_r$, isometries $J_r$, $K_r \geq 1$, $K_{r+1}J_r = J_r K_r$; for $A = L, S$, adjoints $A^*_{r,n} = A_{r,-n}$, uniform bounds $|A_{r,n}|_{s,1} \leq C^A_{n,s}$, summable defects $|A_{r+1,n}J_r - J_r A_{r,n}|_{s,1} \leq e^A_{r,n,s}$, and vanishing loss-two internal residuals. Here $|Y|^{t\leftarrow r}_{s,d} = \|K_t^s Y K_r^{-s-d}\|$, $s \in \mathbb{Z}_{\geq 0}$; within a scale omit $t, r$. VR-1 supplies the common smooth invariant domain $\mathcal{D}$, embeddings $i_r$, and all mixed-word limits. The limiting brackets are $[L_m, L_n] = (m-n)L_{m+n} + z^L_{mn} I$, $[S_m, S_n] = (n-m)S_{m+n} - z^R_{mn} I$, with $z^A_{mn} = \delta_{m+n,0}(m^3 c^*_A - m c_A)/12$; the two coefficients remain independent.

**Lemma 100.1 [derived; any; category-free; → R2] (NOTE:63–100; REVIEW:89–100). Assume** only common A1–A3 for two families. For each fixed $m, n$, $[L_m, S_n] = 0$ on $\mathcal{D}$ iff $|[L_{r,m}, S_{r,n}]|_{s,2} \to 0$ for every $s$. When the limit is zero, the explicit bound is

$$|[L_{r,m}, S_{r,n}]|_{s,2} \leq t_{r,m,n,s} := \sum_{k \geq r} \big( e^L_{k,m,s} C^S_{n,s+1} + C^L_{m,s} e^S_{k,n,s+1} + e^S_{k,n,s} C^L_{m,s+1} + C^S_{n,s} e^L_{k,m,s+1} \big) \longrightarrow 0. \tag{17}$$

⟨1⟩1. Telescope the words $LS$, $SL$, replacing one letter at a time; A1 transports each graph norm exactly. Their combined error on $i_r v$ is at most $t_{r,m,n,s} \|K_r^{s+2} v\|$.

⟨1⟩2. If the limit bracket is zero, the remaining vector has norm $\|K_r^s [L_{r,m}, S_{r,n}] v\|$; take the supremum over $\|K_r^{s+2} v\| = 1$. A3 makes the tail vanish.

⟨1⟩3. Conversely use mixed-word convergence and graph density of the coarse union to pass vanishing finite bounds to $\mathcal{D}$. QED. A4 is unused; fixed-vector convergence alone cannot replace A3.

### 101.3 Parity-Free Balance on the Common Domain

**Lemma 100.2 [conditional; full; category-free; R2 restriction] (NOTE:149–219; REVIEW:117–125). Assume** common A1–A4 for the standard KS families. Put $C_{mn} = [L_m, S_n]$, $X_q = C_{q0}$, $\gamma_m = [m^3(c_R^* - c_L^*) - m(c_R - c_L)]/12$. Then on $\mathcal{D}$,

$$C_{mn} - C_{nm} = \delta_{m+n,0}\gamma_m I, \qquad C_{mn} = X_{m+n} + \tfrac{1}{2}\delta_{m+n,0}\gamma_m I. \tag{18}$$

⟨1⟩1. At finite scale $[E_m, E_n] = \kappa_{mn}B_{m+n}$, $\kappa_{mn} = -ia_N(z^n - z^m)$ (79.1). Subtract the internal brackets from $[L_m + S_m, L_n + S_n]$: $C^{(r)}_{mn} - C^{(r)}_{nm} = (4\kappa_{mn} - 2(n-m))B_{r,m+n} - F^L_{r,mn} - F^R_{r,-m,-n} + \delta_{m+n,0}\gamma_m I$, where $F^L, F^R$ are the unbarred/barred A4 residuals. The coefficient has modulus $\le 2\pi(m^2+n^2)/N$; A2 for $B = (S-L)/2$, A4 and word convergence give the first identity.

⟨1⟩2. L-Jacobi and cyclic summation give $(m-n)C_{m+n,k} + (n-k)C_{n+k,m} + (k-m)C_{k+m,n} = 0$: the leftover commutators have scalar arguments. Set $k = 0$; for $m \neq n$, this is $(m-n)(X_{m+n} - C_{mn}) - n\delta_{m+n,0}\gamma_m I = 0$, giving the second identity off the diagonal.

⟨1⟩3. For $n \neq 0$, the triple $(p, n-p, n)$, $p \notin \{0, n, n/2\}$, gives $(2p-n)(C_{nn} - X_{2n}) = 0$. At $n = 0$ use the definition of $X_0$. QED; no ZC or parity was used.

**Theorem 100.3 [conditional; full; category-free; R2 restriction] (NOTE:279–354; REVIEW:136–148). Assume** two opposite internal brackets as above and the first identity of (18) on a common invariant domain with $I \neq 0$. Then $c_L^* = c_R^*$. Thus a *nonzero* common standard-KS VR-1 system has cubic balance without parity or ZC (100.2).

⟨1⟩1. The algebraic steps of 100.2 give its second identity. Set $A_q = [L_0, X_q]$, $B'_q = [S_0, X_q]$. Unreduced Jacobi gives

$$[L_p, X_q] = pX_{p+q} + A_{p+q} + \tfrac{p}{2}\delta_{p+q,0}\gamma_p I, \qquad [S_p, X_q] = -pX_{p+q} + B'_{p+q} + \tfrac{p}{2}\delta_{p+q,0}\gamma_p I.$$

⟨1⟩2. Write $u = m+n+q$. Jacobi gives $[L_m, B'_{n+q}] = [X_m, X_{n+q}] + mB'_u + [S_0, A_u]$, $[S_n, A_{m+q}] = -[X_n, X_{m+q}] - nA_u + [L_0, B'_u]$, and $[S_0, A_u] - [L_0, B'_u] = -[X_0, X_u]$. Substitute these in $[[L_m, S_n], X_q]$; the $A, B'$ terms cancel:

$$[X_{m+n}, X_q] = [X_m, X_{n+q}] + [X_n, X_{m+q}] - [X_0, X_u] - \tfrac{mn}{2}\delta_{u,0}(\gamma_m + \gamma_n)I.$$

⟨1⟩3. Set $q = -m - n$, $F_n = [X_n, X_{-n}]$: $F_{m+n} = F_m + F_n - \frac{mn}{2}(\gamma_m + \gamma_n)I$. Only $[X_0, X_0] = 0$ was used, not $X_0 = 0$. The paths $(2,2)$, $(1,3)$ yield respectively $F_4 = 4F_1 - (2\gamma_1 + 4\gamma_2)I$ and $F_4 = 4F_1 - (\frac{7}{2}\gamma_1 + \gamma_2 + \frac{3}{2}\gamma_3)I$.

⟨1⟩4. Their difference forces $\frac{3}{2}(\gamma_1 - 2\gamma_2 + \gamma_3)I = \frac{3}{2}(c_R^* - c_L^*)I = 0$; use $I \neq 0$. QED. If each $c_A = c_A^*$ is separately declared, also $c_L = c_R$; that normalization is not a VR-1 conclusion.

**Theorem 100.4 [conditional; full; category-free; R2 restriction] (NOTE:221–277,356–363; REVIEW:126–134,204–210).** Under 100.3's algebraic assumptions, $X_0 = 0$ implies $C_{mn} = 0$ for every $m, n$ and both ledgers agree. For a nonzero common standard-KS VR-1 tower this is equivalent to finite ZC, by 100.1; every finite mixed commutator then obeys (17). Equal ledgers are an output, not an extra input.

⟨1⟩1. Set $q = 0$ in 100.3's unreduced actions: $A_p = -pX_p$, $B'_p = pX_p$. Hence $[L_p, X_q] = -qX_{p+q} + \frac{p}{2}\delta_{p+q,0}\gamma_p I$, $[S_p, X_q] = qX_{p+q} + \frac{p}{2}\delta_{p+q,0}\gamma_p I$.

⟨1⟩2. For $q \neq 0$, double Jacobi gives $[[L_1, S_{-1}], X_q] = 2qX_q$; but $[L_1, S_{-1}] = \gamma_1 I/2$, so every $X_q = 0$.

⟨1⟩3. Put $q = -p \neq 0$ in the L action: $0 = p\gamma_p I/2$. Since $I \neq 0$, $\gamma_1 = \gamma_2 = 0$ kills both ledger differences; (18) kills all mixed brackets. The converse to $X_0 = 0$ is immediate. QED.

⟨1⟩4. At finite scale, $[L_{r,0}, S_{r,0}] = N_r^2[\mathcal{H}_r, \mathcal{Q}_r]/(8\pi^2)$; exact total-current conservation therefore supplies ZC. If instead the limiting right stress is $S_n = t\delta_{n0}I$, its internal $(1,-1), (2,-2)$ equations give $c_R^* = 0$, $c_R = 24t$, and $X_0 = 0$. Thus $c_L^* = 0$, $c_L = 24t$; scalarity of the left stress is not proved.

## 101.4 Two Exact Obstructions to Automatic Zero-Mode Arguments

**Negative 100.5 [negative; full; L0; trivial R2] (NOTE:102–147; REVIEW:102–115).** For every unitary fusion $\mathcal{C}$, take $O = \mathbf{1} \oplus \mathbf{1}$, $H_r = (\mathbb{C}^2)^{\otimes N_r}$, increasing even $N_r$, append-vacuum isometries, $K_r = I$, $k_j = \varepsilon_N = b_N = 0$, genuine cyclic shift $U_r$, and complex-linear bond-centred reflection $P_r$. All F2(a) hypotheses hold and ZC is identically zero, but every self-adjoint logarithm $e^{2\pi i p_r/N_r} = U_r$ satisfies $|D_r - p_r|_{s,1} \geq N_r/2$.

⟨1⟩1. Unit multiplicities give the carrier for any $\mathcal{C}$ (Penneys:219–237,442); all modes, ledgers and defects are zero. A1 is exact; $U, P$ commute with $K = I$.

⟨1⟩2. The one-excitation basis obeys $Ue_j = e_{j+1}$; its Fourier vectors include phase $-1$. Every logarithm has an eigenvalue in $N(1/2 + \mathbb{Z})$ on that block. Since $D = 0, K = I$, the bound follows. QED. This refutes automatic current–logarithm identification, not ZC. Only a *spectral function* $p(U)$ automatically commutes with every operator commuting with $U$; arbitrary logarithms on degenerate blocks need not.

**Negative 100.6 [negative; full Fourier grading; category-free; formal, no Hilbert R2] (NOTE:365–453; REVIEW:150–161).** There is a formal $*$-Lie algebra satisfying Jacobi, dagger, linear parity and spatial grading with $X_0 \neq 0$. Its vector space is $\mathrm{span}\{L_n, S_n : n \in \mathbb{Z}\} \oplus V$, $V = \bigoplus_q \mathbb{C}[t]X_q$, finite sums only; $V$ is abelian. Define zero-central internal brackets as above and

$$[L_m, S_n] = X_{m+n}, \quad [L_p, f(t)X_q] = (t+p)f(t+p)X_{q+p}, \quad [S_p, f(t)X_q] = (t+q)f(t)X_{q+p}.$$

$L_n^\dagger = L_{-n}$, $S_n^\dagger = S_{-n}$, $(f(t)X_q)^\dagger = -\overline{f}(-t)X_{-q}$, where the bar conjugates coefficients; $RL_n = S_{-n}$, $RS_n = L_{-n}$, $R(f(t)X_q) = -f(t-q)X_{-q}$, with $R$ linear.

⟨1⟩1. On $fX_q$, the L-action commutator has coefficient $(m-n)(t+m+n)f(t+m+n)$; the S-action gives $(n-m)(t+q)f(t)$; both mixed compositions give $(t+m)(t+m+q)f(t+m)$. Thus these are commuting opposite-Witt module actions.

⟨1⟩2. The LLS and SSL cocycle equations reduce to $(t+m)-(t+n) = m-n$, $(t+k+n)-(t+k+m) = n-m$. Pure triples are Witt; two V entries give zero.

⟨1⟩3. Substitution verifies dagger reverses brackets, $R$ preserves them, and both commute and square to one. Degrees $-n, -q$ give a derivation $P$: $[P, L_n] = -nL_n$, $[P, S_n] = -nS_n$, $[P, fX_q] = -qfX_q$, $P^\dagger = P$, $RP = -P$. The basis vector $[L_0, S_0] = X_0$ is nonzero. QED. No positive Hilbert representation, implemented translation, microscopic KS realization or VR-1 certificate is asserted.

## 101.5 Commutation Range, Cut Current and Spatial Support

**Proposition 100.7 [derived; periodic; category-free; finite] (NOTE:455–504; REVIEW:163–175).** Assume translated self-adjoint densities commute beyond cyclic density-label distance $\rho$, fixed, $N > 2\rho$. Set $C_{j,d} = i[k_j, k_{j+d}]$, $h_d(w) = \sum_{\ell=0}^{d-1} w^\ell$, $B_n^{(d)} = a_N \sum_j z^{nj} C_{j,d}$. Then

$$\mathcal{J}_j^{\mathrm{cut}} = \sum_{d=1}^{\rho} \sum_{\ell=0}^{d-1} C_{j-\ell,d}, \quad B_n^{\mathrm{cut}} = \sum_d h_d(z^n) B_n^{(d)}, \quad \mathcal{Q}^{\mathrm{cut}} = \sum_j \mathcal{J}_j^{\mathrm{cut}} = \sum_{j,d} dC_{j,d}.$$

$i[\mathcal{H}, k_j] = \mathcal{J}_{j-1}^{\mathrm{cut}} - \mathcal{J}_j^{\mathrm{cut}}$ and $[E_m, E_n] = \sum_d \kappa_{mn}^d B_{m+n}^{(d)}$, $\kappa_{mn}^d = -ia_N(z^{nd} - z^{md})$.

⟨1⟩1. The cut difference telescopes to $\sum_d (C_{j-d,d} - C_{j,d})$; summing cuts counts each term $d$ times.

⟨1⟩2. Pair $(j, j+d)$ in the energy bracket contributes $a_N^2 z^{(m+n)j}(z^{nd} - z^{md})[k_j, k_{j+d}]$; Fourier-shifting $j - \ell$ gives $h_d$. QED.

**Proposition 100.8 [conditional; full; category-free; → R2] (NOTE:506–536; REVIEW:25–31,169–175).** Use corrected modes $L = E - B^{\mathrm{cut}}$, $S = E + B^{\mathrm{cut}}$. Assume their common VR-1

certificates, a nonzero system, and component bounds $|B_n^{(d)}|_{s,1} \le M_{d,n,s}$ independent of scale. Then 100.3 holds, and ZC implies 100.4 and (17).

⟨1⟩1. The Fourier remainder is $\sum_d r^d_{N,m,n} B^{(d)}_{m+n}$, where

$$r^d_{N,m,n} = 4\kappa^d_{mn} - 2(n-m)h_d(z^{m+n}), \qquad |r^d_{N,m,n}| \le \frac{2\pi}{N}\big((m^2+n^2)d^2 + |n-m||m+n|d(d-1)\big).$$

⟨1⟩2. Taylor's exponential remainder and $\sum_{\ell<d} \ell = d(d-1)/2$ give this bound; fixed $\rho$, the component bounds and A4 give (18). Apply 100.3–100.4. QED. For $\rho = 1$, A2 supplies the component bound; for $\rho > 1$, a bound on their sum is insufficient. *Spatial claim [conditional, same setting/data; finite]:* if the densities also have uniformly bounded spatial supports at their translated locations, the finite union of paired supports bounds each cut current's support (REVIEW D1).

**Negative 100.9 [negative; periodic; category-free; finite support diagnostic] (NOTE:463–475; REVIEW:25–52).** Commutation range alone does not imply bounded spatial support. At each of $N \ge 5$ sites put two qubits; $G = \prod_\ell Z'_\ell$, $h_j = X_jX_{j+1} + Z_j$, $k_j = h_j(I+G)$. These translated self-adjoint densities commute at nonadjacent labels, yet $i[k_j, k_{j+1}] = 4X_jY_{j+1}(I+G)$ fails to commute with every $X'_\ell$. BY $[G, h_j] = 0$, $(I+G)^2 = 2(I+G)$, and the Pauli product $XY = iZ$. Thus the current has global support. This finite witness has an L0 realization with $O = \mathbf{1}^{\oplus 4}$ for any $\mathcal{C}$; no VR-1 conclusion.

## 101.6 A Supplied-Target Construction Outside the KS Class

**Proposition 100.10 [conditional; declared chiral target, no local spatial tower; L4; abstract mode R2] (NOTE:540–611; REVIEW:177–189).** Supply a nonzero positive-energy target $(H, E = L_0, L_n)$ with discrete nonnegative energy spectrum, finite-dimensional eigenspaces, adjoint pairing and a common smooth domain containing finite-energy vectors. Assume $[E, L_n] = -nL_n$ and $\|L_nv\| \le a_n\|(1+E)v\|$; Carpi–Weiner:701–747 supplies such bounds in its positive-energy Diff-representation setting, not existence from formal brackets. Let finite-rank orthogonal $P_r \uparrow I$ commute with $E$ and contain every energy $\le \Lambda_r \ge a2^r$, $a > 0$. Take inclusion refinements, $H_r = P_rH$, $K_r = (1+E|_{H_r})^p$, integer $p > 1$, $T_{r,n} = P_rL_nP_r|_{H_r}$, $\bar{T}_{r,n} = 0$. These satisfy VR-1 with the target left ledger and zero right ledger.

⟨1⟩1. A1 is exact. Each mode shifts energy $e$ to $e-n$; orthogonal energy blocks and the assumed bound give A2 with a spare factor $(1+e)^{1-p}$.

⟨1⟩2. Leakage $(P_{r+1} - P_r)L_nP_r$ vanishes below $\Lambda_r - |n|$, so $e_{r,n,s} = O(\Lambda_r^{1-p})$, summable. The bracket residual is exactly $-P_rL_m(1-P_r)L_nP_r + P_rL_n(1-P_r)L_mP_r$; it vanishes below $\Lambda_r - |m| - |n|$ and has $q_{r,m,n,s} = O(\Lambda_r^{2-2p})$. Fixed finite-energy vectors eventually see the target modes exactly. QED. No local observable tower is supplied; nonzero cubic charge rules out the vanishing standard-KS Fourier remainder by 100.3.

**Lemma 100.11 [derived; periodic Fourier regulator; category-free; finite/asymptotic] (NOTE:613–641; REVIEW:191–200).** For even $N$, AP momenta $r = -N/2 + 1/2, \dots, N/2 - 1/2$, integer sites $x, y$, and $\epsilon = 2\pi/N$,

$$D_{xy} = \frac{1}{N}\sum_r re^{2\pi ir(x-y)/N}, \quad D_{xx} = 0, \quad D_{x,x-d} = \frac{-i(-1)^d}{2\sin(\pi d/N)}, \quad \epsilon D_{x,x-d} \longrightarrow \frac{-i(-1)^d}{d} \quad (d \ne 0 \text{ fixed}).$$

For the finite formula require $0 < |d| < N$. Differentiate $\sum_r e^{ir\theta} = \sin(N\theta/2)/\sin(\theta/2)$; then use $\sin u/u \to 1$. Each dyadic block of the absolute limiting $1/|d|$ tail has sum at least $1/2$; it is not summable. This is a fixed-separation limit, not an infinite-kernel operator-norm theorem.

### 101.7 Effect on other shards

*Amendment (2026-10-25, round 3):* CA-91's equal-ledger input to 91.5 is removed by 100.4; 100.1 makes ZC necessary as well as sufficient and supplies loss-two mixed tails. The loss-four MW certificate 91.6 remains valid. Proposition 91.7's logarithm estimate is additional data (100.5), and 91.10's scalar-left conclusion still uses parity although its cubic exclusion does not (100.3–100.4). Range extensions use 100.7–100.9. *Amendment (2026-10-25, round 3):* CA-79, CA-89 and LC-4 may use one ZC certificate once both standard-KS families satisfy VR-1 on the same system; CA-81 gains 100.1's tail. CA-88 gains only the within-cell strengthenings stated above (INTEGRATION:96–153). Exact insertion anchors are in `SHARD_META.md`.

### 101.8 What This Does Not Claim

Automatic ZC (F2(a)), a general finite-range finite-site physical no-go, and sharp F2(c) remain *conjectural* (full; category-free plus declared microscopic data; $\to$ R2, or broader R3 target). The spectral construction changes the stress prescription as well as locality; no infinite-site-only example is supplied. No positive energy, integrated projective-unitary symmetry, local net or category fidelity follows from the KS algebra. *Source audit [derived; any; category-free; no new rung]* (NOTE:656–690; REVIEW:221–224): the local Li–Lin–McGreevy–Shi–Kim passage at :738–746 states the anomaly objection but does not prove the required microscopic theorem there; :1046 also leaves the stress OPE unproved. The primary theorem/bridge remains a gap in this evidence chain.

### 101.9 Examples (Instances Only)

| Instance | Grade, tags and exact scope |
|---|---|
| Any $\mathcal{C}$, $O = \mathbf{1} \oplus \mathbf{1}$ | Negative; full; L0; trivial R2: 100.5 has every logarithm distance $\geq N/2$, but ZC=0. |
| Any $\mathcal{C}$, $O = \mathbf{1}^{\oplus 4}$ | Negative; periodic; L0; finite: 100.9 separates commutation range from support. |
| Declared complex fermion sea | Conditional; periodic nonlocal regulator, chiral target; L4; abstract R2: $c_L = c_L^* = 1$, right ledger zero; $N$ two-state sites, kernel 100.11 and target-compressed stresses (NOTE:818–849; REVIEW:214–220). Not CA-88's open-chain $\mathsf{S}_{\mathrm{ch}}$ construction. |

### 101.10 Single Next Target

**Target 100.12 [conjectural; full; category-free; $\to$ R2] (NOTE:779–784; REVIEW:318–336).** Under original F2(a): a nonzero common standard nearest-neighbour KS A1–A4 tower, complex-linear unitary $P_r k_j P_r^* = k_{-j}$, $[P_r, K_r] = 0$, genuine translation $U_r^N = I$, $U_r k_j U_r^* = k_{j+1}$, $U_r T_{r,n} U_r^* = z^{-n} T_{r,n}$, $U_r \bar{T}_{r,n} U_r^* = z^n \bar{T}_{r,n}$, $[U_r, K_r] = 0$, prove for fixed $a, v \in H_a$

$$\left\| \frac{N_r^2}{8\pi^2} [\mathcal{H}_r, \mathcal{Q}_r] J_{r \leftarrow a} v \right\| \longrightarrow 0.$$

Here $a$ is a fixed stage, $v \in H_a$, and $J_{r \leftarrow a}$ is ordered refinement. Mixed-word convergence, graph density and A2 would give $X_0 = 0$, then 100.1 all uniform ZC bounds and 100.4 all mixed brackets. Alternatively give a genuine tower and coarse vector with nonzero limiting bracket. Do not assume current conservation or logarithm identification: the missing argument is positive-Hilbert/microscopic (INTEGRATION:197–222).

# 102 Ancilla Transport Lifts and the Failure of Finite-Power Extraction

## 102.1 Status, Sources, and Dependencies

**Status: COUNTEREXAMPLE to G1 as posed and to Target 96.10 exactly as stated.** 101.1, 101.2, 101.4 and 101.6 are *derived*; 101.3, 101.7 and 101.8 are *negative*, with derived witnesses/arguments; the remark-level gap 101.5 is *conditional*; 101.9 is *conjectural*. **Pillar P1:** local unital refinement of the indefinite-particle, vacancy-carrying tower; this feeds **P4** through WT-1 A1, without a state or drift estimate (REVIEW:172–182). **Grid (INTEGRATION:44–59):** (any; $\mathsf{L0}$ + declared ancilla datum; $\mathsf{R0}^u$) gains a derived certificate, and bare $\mathsf{L0}/\mathsf{R0}^u$ gains witnesses without any transportable site power. CA-88's minimal-data wording must become sufficiency: transport on $O$ or $O^b$ is not necessary. General existence remains open; the rung of CA-90, $\mathsf{L1}/\mathsf{R0}^u$, $\mathsf{R0}^c$, $\mathsf{R1}$–$\mathsf{R5}$, the full/periodic entries and D1-C's occupation-compatible target do not move. Setting *any* means no chirality target is selected; the microscopic presentation is an open ordered chain. NOTE, REVIEW, INTEGRATION and `checks/` denote the files in `explorations/2026-10-25-astra-r3/G1-transport-extraction/`. Dependencies: CA-75 (Link (b1)), CA-76 (76.1–76.4, 76.12), CA-77 (pointed carrier), CA-88 (grid), CA-90 (90.H, 90.0–90.2), CA-96 (96.2, 96.4, 96.8, 96.10), CA-97 (97.11–97.13); CONVENTIONS (r), (t), (am), (aq), (as), (ax). New notation is shard-local, proposed in `SHARD_META.md`.

## 102.2 Setting and Ancilla Transport

Throughout, $\mathcal{C}$ is arbitrary unitary fusion, $\eta : \mathbf{1} \to O$ is an isometry, $O \cong \mathbf{1} \oplus X$, $\mathcal{D} = \langle O \rangle$ full, $W_L = O^{\otimes L}$, $W_0 = \mathbf{1}$, and $\mathcal{A}_L = \mathrm{End}_{\mathcal{C}}(W_L)$. Use left bracketing and fixed unitary associators/unitors; $V_L$ inserts $\eta$ at even sites. The occupation operator is $N_L = \sum_j (1 - \eta\eta^\dagger)^{(j)}$, as in (am),(as). $\mathcal{A}_{[j,k]} = 1_{W_{j-1}} \otimes \mathrm{End}_{\mathcal{C}}(W_{k-j+1}) \otimes 1_{W_{L-k}}$: support is membership. A lift means unital injective $*$-homomorphisms $\alpha_L : \mathcal{A}_L \to \mathcal{A}_{2L}$ with $\alpha_L(a)V_L = V_L a$, all $L$, and $\alpha_L(\mathcal{A}_{[j,k]}) \subset \mathcal{A}_{[2j-1-r,2k-1+r]}$ for one finite slack $r$, endpoints clipped. Write $m_c(L) = \dim \mathrm{Hom}(c, W_L)$. $\Lambda_{c,d}(L)$ counts input-block $c$ copies in output charge block $d$, not projection ranks; only occurring blocks are used. A point-fixing gauge is $\mathrm{Ad}_U$, $U \in \mathcal{A}_{2L}$ unitary, $UV_L = V_L$; without a support bound it is global (NOTE:21–35).

**Theorem 101.1 [derived; any; $\mathsf{L0}$ + $(K, \kappa, t, u)$; $\mathsf{R0}^u$] (NOTE:37–89; REVIEW:51–54,128–132). Assume** a pointed ancilla $\kappa : \mathbf{1} \to K$, $\kappa^\dagger\kappa = 1$, unitaries $t_Y : Y \otimes K \to K \otimes Y$ for all $Y \in \mathcal{D}$, and a unitary $u : O \otimes K \to O \otimes O$, with

$$t_{\mathbf{1}} = 1_K, \quad t_Z(f \otimes 1_K) = (1_K \otimes f)t_Y, \quad t_{Y\otimes Z} = (t_Y \otimes 1_Z)(1_Y \otimes t_Z),$$
$$t_Y(1_Y \otimes \kappa) = \kappa \otimes 1_Y, \qquad u(1_O \otimes \kappa) = 1_O \otimes \eta.$$

Here $f : Y \to Z$ is arbitrary; $K \in \mathcal{D}$ follows since $K \subset O \otimes K \cong O^2$. **Prove** the following all-$L$ lifts, with exact $V_L$, slack one and coherent dyadic composites:

$$\alpha_L(a) = T_L(a \otimes 1_{K^{\otimes L}})T_L^\dagger, \quad T_L = u^{\otimes L}B_L^K, \quad \Lambda_{c,d}^K(L) = \dim \mathrm{Hom}(d, c \otimes K^{\otimes L}). \tag{19}$$

⟨1⟩1. Insert $K_i$ after old site $i$, crossing only the old suffix: CA-90 Lemma 90.0's heterogeneous argument gives $B_L^K : W_L \otimes K^{\otimes L} \to (O \otimes K)^{\otimes L}$. Naturality and the tensor axiom give interval membership from old $2j-1$ to old $2k-1$; no $K$-$K$ crossing occurs.

⟨1⟩2. Pointedness gives $T_L(1_{W_L} \otimes \kappa^{\otimes L}) = V_L$. Conjugation gives a unital $*$-map; compression gives intertwining and injectivity.

⟨1⟩3. Exterior $u$-cells cancel, so $\alpha_L(\mathcal{A}_{[j,k]}) \subset \mathcal{A}_{[2j-1,2k]}$. Composition gives the dyadic cocycle, not coherence across arbitrary placements.

⟨1⟩4. Orthogonal semisimplicity decomposes $\mathrm{Hom}(d, W_L \otimes K^{\otimes L})$ as $\bigoplus_c \mathrm{Hom}(c, W_L) \otimes \mathrm{Hom}(d, c \otimes K^{\otimes L})$; restriction multiplicities give the formula. QED. CA-90 is $K = O, u = 1$, with its sharper slack zero. Existence of the datum is an assumption.

### 102.3 Gauge Invariants and Extraction Boundaries

**Proposition 101.2 [derived; any; L0; finite → R0$^u$] (NOTE:130–161,295–327; REVIEW: 57–60,134–139).** For any two lifts of the same $V_L$, equality of $\Lambda$ is equivalent to global point-fixing unitary conjugacy. It is invariant under any output unitary, and under input/output $*$-automorphisms up to block relabelling. Every CA-96 transport lift, regardless of its block size, has $\Lambda^{\mathrm{tr}}_{c,d}(L) = \dim \mathrm{Hom}(d, c \otimes O^{\otimes L})$. Thus a mismatch with (19) excludes every such transport lift up to output unitary gauge.

⟨1⟩1. Conjugation preserves representation multiplicities; automorphisms only relabel blocks. CA-96's inserted blocks and tail total $O^{\otimes L}$, so its map is $Q_L(a \otimes 1_{W_L})Q_L^\dagger$.

⟨1⟩2. $P = V_L V_L^\dagger$ reduces both lifts (76.1); on $P$ both equal $V_L a V_L^\dagger$. Equal $\Lambda$ gives equal complement multiplicities $\Lambda_{c,d} - \delta_{c,d}$. Choose blockwise unitary intertwiners $W$ on $1-P$; $U = P + W$ conjugates the lifts and fixes $V_L$. QED. No locality bound on $U$ follows.

**Negative 101.3 [negative; any; bare L0; R0$^u$ witnesses] (NOTE:171–287; REVIEW: 14–42,61–66,141–143).** For general $\mathcal{C}$, a local lift family need not come from transport on any $O^{\otimes b}$, even as an existence criterion. G1(a)–(c)'s extraction/dichotomy and the positive answer to Target 96.10 are false exactly as posed.

⟨1⟩1. Any relative half-braiding on $O^{\otimes n}$ gives $O^{\otimes n} \otimes Y \cong Y \otimes O^{\otimes n}$; thus $[O]^n$ must be central in $K(\mathcal{D})$, even without pointedness or unitarity (90.6).

⟨1⟩2. The table supplies exact data satisfying 101.1, but with $[O]^n$ noncentral for every $n \geq 1$. Hence all-$L$ slack-one lifts and no transportable power coexist. QED. The general mechanism is $[O]([K] - [O]) = 0$ with $[K]$ central: cancellation is unavailable. The invariant 101.2 separately obstructs recovery of a given lift even when site transport exists (first table row). Occupation-degree covariance fails in the table's finite checks; the occupation-compatible variant of 96.10 remains open.

**Lemma 101.4 [derived; any; L0; finite → R0$^u$] (NOTE:295–338; REVIEW:67–69,145).** An arbitrary fixed-volume lift extracts nonnegative integers $\Lambda_{c,d}$, a distinguished vacuum copy, and

$$m_d(2L) = \sum_c m_c(L)\Lambda_{c,d}, \qquad \Lambda_{c,c} \geq 1, \qquad \sum_c N_{ij}^c \Lambda_{c,d} = T^\alpha_{(ij),d} \quad (L = s + t).$$

Here $T^\alpha$ is the multiplicity of block $(i, j)$ under the *actual* restriction to $\mathrm{End}(W_s) \otimes \mathrm{End}(W_t)$, with $i, j$ occurring in the two sides.

⟨1⟩1. Unitality gives the dimension sum; the reducing vacuum corner supplies the diagonal copy.

⟨1⟩2. Restriction through $\mathcal{A}_L$ multiplies inclusion matrices; its first multiplicities are $N_{ij}^c$. This proves the identity, with no locality or hexagon input. QED. The global classification is 101.2.

**Proposition 101.5 [conditional; any; L0; finite → R0$^u$] (NOTE:340–347; REVIEW: 70).** For a gap object $P_{\mathrm{gap}} \in \mathcal{D}$, the proposed deduction requires an independently justified identification $T^\alpha_{(ij),d} = \dim \mathrm{Hom}(d, i \otimes P_{\mathrm{gap}} \otimes j)$; only then do the two unit rows identify $\dim \mathrm{Hom}(d, P_{\mathrm{gap}} \otimes x) = \dim \mathrm{Hom}(d, x \otimes P_{\mathrm{gap}})$, for $x$ in both sides (90.2(b)). Dyadic interval containment supplies no such identification: the side maps are refined. *Review GAP, NOTE ⟨1⟩4:* the claim that an actual lift directly witnesses failure of this particular formula remains conditional on computing its $T^\alpha$ with the comparison correctly typed. The note supplies no such calculation; its one-gap/dyadic comparison is not used in 101.3 or 101.4.

## 102.4 Finite Equations and Further Obstructions

**Proposition 101.6 [derived; any; L0+finite associator/dagger data; finite → R0$^u$] (NOTE: 361–408; REVIEW:71–73,147–149).** Fix $b \geq 1$, $P = O^{\otimes b}$, $\xi = \eta^{\otimes b}$. For simples $s_i$ of full $\mathcal{D}$, choose complete orthonormal embeddings $f_{ij;k,\mu} : s_k \to s_i \otimes s_j$. Pointed transport is equivalent to the finite system

$$z_i : P \otimes s_i \to s_i \otimes P, \quad z_i^\dagger z_i = 1, \quad z_i z_i^\dagger = 1, \quad z_{\mathbf{1}} = 1_P, \quad z_i(\xi \otimes 1_i) = 1_i \otimes \xi,$$

$$(f_{ij;k,\mu} \otimes 1_P) z_k = (1_i \otimes z_j)(z_i \otimes 1_j)(1_P \otimes f_{ij;k,\mu}),$$

with fixed associators inserted; $z_Y = t_Y^{-1}$, consistently with (90.5).

⟨1⟩1. Necessity and sufficiency follow from 90.2(a): extend over orthogonal simple summands; Schur gives naturality, and summing against $f^\dagger$ gives the tensor axiom. Real/imaginary coordinates are polynomial.

⟨1⟩2. A pointed unitary tensor equivalence and its quasi-inverse transport solutions at the same $b$ (96.8). Changing $\eta$ conjugates by a unitary on the unit multiplicity space of $O$, tensored $b$ times. Thus the admissible set and its minimum ($\infty$ if empty) are invariant. QED. This minimum can be $\infty$ with slack-one lifts (101.3); it is no lift criterion or bound from $r$. Finite equations at each fixed $b$ imply no decision procedure for some $b$: no terminating bound or general exact encoding of complex input is supplied. No decidability or undecidability claim is made.

**Negative 101.7 [negative; any; L0; → R0$^u$] (NOTE:10–19; REVIEW:119–124,151–152).** G1's hypotheses impose no relation between different lengths. Choosing two available lift constructions at different lengths preserves exact $V_L$ and a common slack bound; composition still defines dyadic maps. The first table row alternates inequivalent inclusion matrices at even/odd lengths, refuting recovery by one site-power transport datum. This is the reviewer's derived R1 argument; extraction needs a separate uniformity hypothesis.

**Negative 101.8 [negative; any; L0; → R0$^u$] (NOTE:87–89; REVIEW:154–165).** Existence of a blocked ancilla datum is not universal either. For arbitrary $\mathcal{C}$, such a datum at block size $b$ necessarily supplies a positive integral class $k = [K]$ central in $K(\mathcal{D})$, with $[O]^b k = [O]^{2b}$.

⟨1⟩1. $K \subset O^b \otimes K \cong O^{2b}$ lies in $\mathcal{D}$; transport gives centrality and the unitary gives the class equation. $K = O^b$ includes CA-96's construction.

⟨1⟩2. The final table row rules out every integral central solution for every $b \geq 1$. QED for the negative to these constructions (reviewer R2). This is not a no-go for arbitrary local lifts.

## 102.5 Effect on other shards

*Amendment (2026-10-27, round 4):* CA-96's status, necessity clause and Target 96.10 must point to 101.3 (refuted) and 101.9 (replacement); its unresolved carrier is outside all blocked/ancilla constructions by 101.8. CA-90's Boundary and 90.T gain 101.1 and the failure of site-power necessity; 90.2(b) retains its one-gap scope. CA-97's "supplied H" qualification is sharp: the counterexample has no site-power H; its compression mark fails unitarity in the finite check below. CA-88's entries change only as stated in Status. CA-75 Link (b1) and CA-76's local-unital pointer gain 101.1; Link (b3) and 76.12's state hypotheses remain open (INTEGRATION:61–95).

## 102.6 What This Does Not Claim

No universal existence or classification of local lifts, no extraction of site-power transport, no local-gauge bound from equal inclusion matrices, and no decision procedure for some block size. No occupation-preserving counterexample to 96.10: preservation means $[N_{2L}, \alpha_L(a)] = \alpha_L([N_L, a])$ as

in (am),(as). Trailing descent is only finitely checked for the negative witnesses; no all-$L$ proof is lifted (REVIEW:167–170). No state estimate, Wilson limit, modes, seam, conformal limit or fidelity is supplied; all higher rungs stay open here.

## 102.7 Examples (Instances Only)

These instantiate 101.1–101.3, 101.7–101.8; grades and tags are those statements' tags, with bare L0. All group categories below are untwisted unitary $\mathrm{Vec}_G$, with orthonormal homogeneous bases.

| Instance | Exact data and check of the general statement |
|---|---|
| Finite $G \neq \{1\}$, $N = \lvert G\rvert$ | $O = \bigoplus_{g\in G} g$, $\eta = e_1$, $K = N\mathbf{1}$, $\kappa = k_1$; $t_Y(y \otimes k_h) = k_h \otimes y$, $u(e_g \otimes k_h) = e_{gh^{-1}} \otimes e_h$. This grade-preserving bijection fixes the point and gives the cellwise $T_L = u^{\otimes L} B_L^K$. $\Lambda^K_{c,d} = N^L \delta_{c,d}$, whereas $\Lambda^{\mathrm{tr}}_{c,d} = N^{L-1}$, by choosing the first $L-1$ letters freely. At $L = 1$ images of minimal projections are central for the former, noncentral for the latter. $O$ itself transports by conjugation of grades: classification fails, not transport existence. Taking the former lifts at even $L$, the latter at odd $L$, gives R1 (NOTE:91–169; REVIEW: 119–124). |
| Dihedral family, every $m \geq 3$ | $G = \langle a, b \mid a^2 = b^2 = 1, (ab)^m = 1\rangle$, $\rho_0 = ab$, $O = \mathbf{1} \oplus a \oplus b \oplus \rho_0$, $K = \mathbf{1} \oplus \mathbf{1} \oplus \rho_0 \oplus \rho_0^{-1}$, $\eta = e_1, \kappa = E_0$. The unit-grade vectors are $E_0, E_1$; $R_\pm$ have grades $\rho_0^{\pm 1}$. Let $\rho(x)$ fix $E_0, E_1$ and permute $R_\pm$ by grade conjugation; $t_Y(y_x \otimes v) = \rho(x)v \otimes y_x$. This satisfies H since $(xhx^{-1})x = xh$ and $\rho(xy) = \rho(x)\rho(y)$. Set $u(g, E_0) = (g, 1)$; the other twelve values (outputs mean $e_x \otimes e_y$) are<br>$g$ \| $E_1$ \| $R_+$ \| $R_-$<br>1 \| $(a,a)$ \| $(1,\rho_0)$ \| $(b,a)$<br>$a$ \| $(1,a)$ \| $(1,b)$ \| $(\rho_0,a)$<br>$b$ \| $(a,\rho_0)$ \| $(b,\rho_0)$ \| $(\rho_0,b)$<br>$\rho_0$ \| $(a,b)$ \| $(\rho_0,\rho_0)$ \| $(b,b)$<br>. Every pair occurs once and each product grade is preserved, using only $a^2 = b^2 = 1$. For $x_i = g_{i+1} \cdots g_L$, $x_L = 1$, the exact lift is implemented by $T_L(g_1, \ldots, g_L; h_1, \ldots, h_L) = \bigotimes_i u(g_i, \rho(x_i)h_i)$, $h_i \in \{E_0, E_1, R_+, R_-\}$. Thus 101.1 gives literal $V_L$ and support $[2j-1, 2k]$, every $L$.<br>For the no-power proof put $\theta = 2\pi/m$, $A = \mathrm{diag}(1, -1)$, $R = \left(\begin{smallmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{smallmatrix}\right)$, $B = AR$. These represent $a, \rho_0, b$. With $s = [O] = (1+a)(1+b)$, $\lambda = 2(1+\cos\theta) > 0$, $M = (I+A)(I+B) = \left(\begin{smallmatrix} \lambda & -2\sin\theta \\ 0 & 0 \end{smallmatrix}\right)$, $M^n = \lambda^{n-1} M$ and $[M^n, A]_{12} = 4\lambda^{n-1}\sin\theta \neq 0$. Hence $s^n$ is never central, and no relative half-braiding exists on any $O^n$. Also $s(2+\rho_0+\rho_0^{-1}) = s^2$; singular $M$ explains noncancellation (NOTE:171–287).<br>Finite evidence only: $m = 3, 4$, $L = 1, 2$ trailing descent passes on 92 matrix units; occupation degree fails on 276/244 entries of $\alpha_2$. At $m = 3$, the CA-97 mark $j^\dagger \alpha_2(h) i$ has rank 8/16 and unitarity defect 1 (`checks/review_g1_independent_output.txt`:7–14). |
| Open carrier in $\mathrm{Vec}_{S_3}$ | $O = \mathbf{1} \oplus g \oplus h$, $g = (12), h = (23)$, $\eta = e_1$, $\mathcal{D} = \mathcal{C}$. Any central integral $k$ is $\alpha + \beta C_3 + \gamma C_2$, where $C_3$ sums the two 3-cycles and $C_2$ the three transpositions. Applying trivial/sign representations to $[O]^b k = [O]^{2b}$, and cancelling the nonzero scalars $3^b, (-1)^b$, gives $\alpha + 2\beta + 3\gamma = 3^b$, $\alpha + 2\beta - 3\gamma = (-1)^b$. Thus $6\gamma = 3^b - (-1)^b \not\equiv 0 \pmod 3$, impossible for integral $\gamma$, every $b \geq 1$. No blocked or ancilla datum exists; arbitrary local-lift existence remains open (REVIEW:154–165). Exact enumeration through $b = 5$ agrees (`checks/review_g1_independent_output.txt`:5–6). |

Reproduce from the package: `python3 -B checks/review_g1_independent.py`; output and headline are in `checks/review_g1_independent_output.txt`:1–15. Six mutations `K`, `u`, `transport`, `point`, `dress`, `central0` are RED; see `checks/review_g1_red_*.txt` and REVIEW:96–103. The `authorpoint` control stays GREEN: it changes to another fixed point. Exact combinatorics and symbolic identities, with tolerance $10^{-9}$ only for dense random tests, calibrate the proofs; finite checks do not establish their all-$L$, all-$m$ quantifiers.

### 102.8 Single Next Target

**Target 101.9 [conjectural; any; L0; $\to$ R0$^u$] (NOTE:349–359; REVIEW:196–208; INTEGRATION:121–137).** *Blocked-ancilla extraction under uniformity.* For arbitrary $\mathcal{C}, (O, \eta)$, assume the lift hypotheses above and one fixed $b \geq 1$ with $\alpha_{L+b}(a \otimes 1_{W_b}) = \alpha_L(a) \otimes 1_{W_{2b}}$ for every $L, a$. Prove or refute existence of $b' \geq 1$, pointed $(K, \kappa)$, pointed unitary transport on full $\mathcal{D}$, and unitary $u : O^{\otimes b'} \otimes K \to O^{\otimes 2b'}$, $u(1 \otimes \kappa) = V_{b'}$, whose blocked construction reproduces $\alpha_L$ up to $V_L$-fixing unitaries of uniformly bounded support. No occupation, translation or unchanged-side assumption may be added. First establish sufficiency of the blocked form by 101.1 and 96.2's block/tail argument; its necessary class equation is $[O]^{b'}([K] - [O]^{b'}) = 0$ with $[K]$ central (101.8). The last table row calibrates this target: a positive answer excludes lifts satisfying its uniformity hypothesis there; an arbitrary-lift no-go would additionally require that uniformity, which is not part of 96.10.

# 103 The Interval/Dilation Record Forgets the Point

## 103.1 Status, Sources, and Dependencies

**Status: COUNTEREXAMPLE** to the positive form of CA-97 Target 97.16, exactly as stated. Statements 102.1, 102.3, 102.5–102.7 are *derived*; 102.2 and 102.4 are *negative*; 102.8 is *conditional*, an unproved extension with a Law-1 gap; 102.9 is *conjectural*. NOTE, REVIEW and INTEGRATION mean the files in `explorations/2026-10-25-astra-r3/G6-record-faithfulness/`. **Pillar P1:** the vacancy in $O \cong \mathbf{1} \oplus X$, and hence the indefinite-number interpretation, must be carried along the tower as declared structure. **P4 interface:** this sharpens the R4 insufficiency annotation; it supplies no continuum certificate. **CA-88 cells (INTEGRATION:36–46):** (any/open; $\mathsf{L}0+H$; $\mathsf{R}0^u$), dyadic geometry, gains a negative information annotation, not an existence or attainment change. The R4 insufficiency annotation extends to the full interval/dilation record: the first missing mark proved here is the point. Trailing (any; $\mathsf{L}0$; $\mathsf{R}0^u$), $\mathsf{R}0^c$, R1–R5 attainment, full/periodic and L3 cells do not move; CA-93's fidelity hypotheses remain required. All settings below are *any, ordered open geometry*; no chiral/full continuum target is selected. Dependencies: CA-77 (77.2), CA-88 (grid), CA-90 (90.H, 90.1), CA-97 (97.11, 97.13, 97.16); CONVENTIONS (a), (b), (r), (am), (aq), (as), (ax). Additional notation is scoped here; its proposed registration is in the package's `SHARD_META.md`.

Checks: the exact matrix-unit script and independent NumPy reviewer script reproduce their stored outputs; the latter checks crossing phases and point-block sizes. Their finite checks do not prove the all-length statements.

## 103.2 The General Record and Cut Information

Fix arbitrary unitary fusion $\mathcal{C}$, pointed $(O,\eta)$, and $\mathcal{D} = \langle O\rangle \subset \mathcal{C}$. Let $H = (t_Y)$ be one fixed pointed unitary relative half-braiding on $\mathcal{D}$ (90.H): $t_Y : Y \otimes O \to O \otimes Y$ is natural, tensorial and unital, with $t_Y(1_Y \otimes \eta) = \eta \otimes 1_Y$. Put $W_L = O^{\otimes L}$, $W_0 = \mathbf{1}$, $A_L = \mathrm{End}_{\mathcal{C}}(W_L)$, and let $\kappa_L : A_L \to A_\infty$ be the norm-limit maps for trailing inclusion $a \mapsto a \otimes 1_O$. Bracketing is left-associated; fixed coherent associators are retained. The shuffle $B_L$ crosses old $j$ with new $i$ exactly when $i < j$, and $\alpha_L(a) = B_L(a \otimes 1_{W_L})B_L^\dagger$, $\delta\kappa_L = \kappa_{2L}\alpha_L$ (97.11). The record and its proposed pointed augmentation are

$$\mathcal{R} = (A_\infty, \{A_{[j,k]}\}, \delta), \quad A_{[j,k]} = \kappa_k(1_{W_{j-1}} \otimes A_{k-j+1}), \qquad \mathcal{R}^p = (\mathcal{R}, \{p_j\}), \quad p_j = \kappa_j(1_{W_{j-1}} \otimes \eta\eta^\dagger).$$

An isomorphism preserves exactly the listed marks, every interval label, and $\delta$. The $p_j$ are placed copies of (am)'s $p_0$, not its occupation projection $p_1$. "Natural recovery" means transport by *every* record isomorphism; recovery up to a coherent choice is weaker.

**Lemma 102.1 [derived; any/open; $\mathsf{L}0+H$; R0 finite information] (NOTE:43–99; REVIEW:70–73).** For any cut $m|n$, put $V_m^x = \mathrm{Hom}(x, W_m)$, $F_{xy}^z = \mathrm{Hom}(z, x \otimes y)$; omit zero summands. Under orthogonal simple splittings,

$$\begin{aligned} A_{m+n} &\cong \bigoplus_z \mathrm{End}\Big(\bigoplus_{x,y} V_m^x \otimes V_n^y \otimes F_{xy}^z\Big), \\ (A_{[1,m]} \vee A_{[m+1,m+n]})' \cap A_{m+n} &\cong \bigoplus_{z,x,y} 1_{V_m^x \otimes V_n^y} \otimes \mathrm{End}(F_{xy}^z). \end{aligned} \tag{20}$$

The record preserves these commutants, their inclusion diagrams, and intersections involving $\delta^r(A_I)$.

⟨1⟩1. Compose orthonormal isometric splittings of $W_m, W_n, x \otimes y$ (Penneys:219–237,319,470–478); the two interval algebras act on the two $V$ factors. Their product embedding is faithful: $d_x d_y > 0$, so $x \otimes y \neq 0$ (ENO:1459–1487,1700–1706, including the reviewer's missing unitary bridge).

⟨1⟩2. Within each $z$-block, the two central projections force preservation of $(x, y)$. Commuting with matrix units kills off-diagonal entries on $V_m^x \otimes V_n^y$ and equalises diagonal entries; the remaining operator is $1 \otimes c$. Conversely every such operator commutes.

⟨1⟩3. Isomorphisms preserve multiplication, commutation and intersections; intertwining $\delta$ gives the moved-cut assertion. QED. The formula alone uses L0; the last assertion needs $H$. The displayed simple labels are choices, not intrinsically named charges.

### 103.3 An Exact Counterexample Family

**Negative 102.2 [negative; any/open; L0 + $H$; R0$^u$ information]**
**(NOTE:101–184; REVIEW:74–79,170).** The universal claim that $\mathcal{R}$ determines the pointed input for arbitrary $\mathcal{C}$ is false. The mechanism is proved for the subclass $\mathcal{D} = \mathrm{Hilb}_{\mathrm{fd}}(G, 1)$, finite abelian $G$, with trivial associator. Write $O = \bigoplus_{a \in G} \mathbb{C}^{m(a)} \otimes a$. Choose $s \in G$ with $m(e), m(s^{-1}) > 0$; both $\operatorname{supp} m$ and its translate must generate $G$. Let $O' = sO$ retain multiplicity spaces but replace grade $a$ by $sa$; choose unit points in $O_e, O'_e$, and initially take flip $H$ on both. Then $\mathcal{R}(O) \cong \mathcal{R}(O')$, even preserving translations and endomorphism concatenation. If no group automorphism matches the two multiplicity functions, no tensor equivalence carries $O$ to $O'$.

⟨1⟩1. The flip is grade-preserving since $G$ is abelian, natural on graded maps, unitary and pointed; its tensor identity sends $y \otimes z \otimes v$ to $v \otimes y \otimes z$ in either order. This verifies 90.H in the sourced graded construction (Penneys:277–288,319–321).

⟨1⟩2. The underlying Hilbert unitary $U : O \to O'$ preserves multiplicity bases and has degree $s$; it is *not* a categorical morphism for $s \neq e$. Put $U_L = U^{\otimes L}$, $\phi_L = \operatorname{Ad} U_L$, $\phi_0 = \mathrm{id}$. A word of total charge $x$ acquires charge $s^L x$. Thus $\phi_L$ maps all categorical matrix units bijectively and is an isometric unital $*$-isomorphism.

⟨1⟩3. Factorwise conjugation gives simultaneously, for $b \in A_{k-j+1}$,

$$\phi_{L+1}(a \otimes 1) = \phi_L(a) \otimes 1, \qquad \phi_L(1^{\otimes(j-1)} \otimes b \otimes 1^{\otimes(L-k)}) = 1^{\otimes(j-1)} \otimes \phi_{k-j+1}(b) \otimes 1^{\otimes(L-k)}.$$

These are onto on every interval; also $\phi_{L+M}(a \otimes b) = \phi_L(a) \otimes \phi_M(b)$.

⟨1⟩4. The flip shuffle is a tensor permutation, hence

$$U_{2L} B_L = B'_L (U_L \otimes U_L), \qquad \phi_{2L} \alpha_L = \alpha'_L \phi_L, \qquad \Phi\delta = \delta'\Phi. \tag{21}$$

For the last identity extend the compatible $\phi_L$ and their inverses to the norm completions and use $\Phi\delta\kappa_L = \kappa'_{2L}\phi_{2L}\alpha_L = \delta'\Phi\kappa_L$, then density (97.11).

⟨1⟩5. A tensor equivalence permutes invertible simples by a group automorphism and preserves multiplicities (Penneys:325–348); the stated mismatch forbids it. QED. In (20), $F^z_{xy}$ has dimension one exactly when $z = xy$; the relabelling $(x, y, z) \mapsto (s^m x, s^n y, s^{m+n} z)$ preserves every cut diagram, including $\delta$-moved cuts. This demonstrates insufficiency for this family, not a reconstruction obstruction for every $\mathcal{C}$.

**Proposition 102.3 [derived; any/open; L0 + $H$; R0$^u$ information]**
**(NOTE:154–177; REVIEW:106–121).** Reviewer R-1 extends 102.2 to $t_Y(y_a \otimes v_b) = \chi(a, b) v_b \otimes y_a$, with unitary bicharacter $\chi$, provided $\chi(a, s) = \chi(s, a) = 1$ for every $a \in G$.

⟨1⟩1. Bicharacter identities give 90.H, including $\chi(a, e) = 1$ at the point.

⟨1⟩2. Transparency gives $\chi(sa, sb) = \chi(a, b)$ at each crossing, proving (21) with the same phases. The other steps are unchanged. QED. Failure of this particular $\Phi$ for a nontransparent $s$ excludes no other possible record isomorphism.

**Negative 102.4 [negative; any/open; L0 + $H$; R0 information]**
**(NOTE:186–200; REVIEW:47–63,80–81).** With flip $H$, let $R_G = \bigoplus_{a\in G} a$. *B1:* for every nontrivial finite abelian $G$ and $s \neq e$, take $O = R_G \oplus \mathbf{1}$, $O' = R_G \oplus s$. Both supports generate; translation gives 102.2, but $\dim \mathrm{Hom}(\mathbf{1}, O) = 2 \neq 1 = \dim \mathrm{Hom}(\mathbf{1}, O')$, invariant under every tensor equivalence. *B2:* take nonunit $a, b$ in distinct automorphism orbits, $s = ba^{-1}$, $O = R_G \oplus a$, $O' = R_G \oplus b$. Both are connected; an equivalence would carry the uniquely doubled simple $a$ to $b$. Different orders suffice; this order-based witness requires such elements in $G$. Thus the positive form of 97.16 is false, including for connected sites. For $s \neq e$, $\phi_1(\eta\eta^\dagger)$ lies in grade $s$, whereas the target point lies in grade $e$. The record forgets the point and the marks built from it; $\mathcal{D}$ and the supplied transport rule are unchanged.

## 103.4 Named Points and the Remaining Marks

**Proposition 102.5 [derived; any/open; L0 + $H$ and (G6.5); R0 marks]**
**(NOTE:202–269; REVIEW:82–85).** For arbitrary $\mathcal{C}$ in the general setting, assume a noncentral unitary $u \in \mathrm{End}(O)$ with

$$u\eta = \eta, \qquad t_Y(1_Y \otimes u) = (u \otimes 1_Y)t_Y \quad (Y \in \mathcal{D}). \tag{G6.5}$$

Then $\mathcal{R}^p$ has an automorphism moving the standard translation, concatenation, left-insertion corner and hop $h$, although their relevant range projections stay fixed.

⟨1⟩1. Put $C_0 = C_1 = 1$, $C_L = 1_O \otimes u^{\otimes(L-1)}$, $\gamma_L = \mathrm{Ad}\, C_L$. Since $C_{L+1} = C_L \otimes u$ for $L \geq 1$, these maps respect trailing inclusion and map every interval algebra onto itself; extend to $\gamma$ by norm density.

⟨1⟩2. Naturality transports $u$ on old strands, (G6.5) on ancillary strands; old strand 1 never crosses. Thus $B_L(C_L \otimes u^{\otimes L}) = C_{2L}B_L$. Conjugate $a \otimes 1$: ancillary factors cancel, so $\gamma_{2L}\alpha_L = \alpha_L\gamma_L$ and $\gamma\delta = \delta\gamma$. Also $\gamma(p_j) = p_j$ since $u\eta = \eta$.

⟨1⟩3. Choose $b$ with $ubu^\dagger \neq b$. For $T(b) = 1 \otimes b$, $\gamma_2 T(b) = 1 \otimes ubu^\dagger \neq T\gamma_1(b)$; the same inequality moves concatenation $(a, b) \mapsto a \otimes b$, and $\theta(b) = p_0 \otimes b$, $p_0 = \eta\eta^\dagger$. The latter range $p_0 \otimes 1$ is fixed, and compression by $\eta$ proves the inequality persists.

⟨1⟩4. For $r = \eta \otimes 1, l = 1 \otimes \eta : O \to W_2$, $h = rl^\dagger$, $C_2 r = ru$, $C_2 l = l$, hence $\gamma_2(h) = rul^\dagger \neq h$, with both support projections fixed.

⟨1⟩5. Every input admits such an instance after replacing $O$ by $O \oplus \mathbf{1}$ and extending $H$ by the unit transport. Set $u = +1$ on the chosen point, $-1$ on the new unit copy, and identity elsewhere. Pointedness and unitarity split off the old unit transport; (G6.5) holds blockwise, while a matrix unit between the two unit copies witnesses noncentrality. Generation is unchanged. QED. This proves *non-naturality*, not failure of reconstruction up to a coherent choice.

**Proposition 102.6 [derived; any/open; L0 + $H$; R0 information]**
**(NOTE:281; REVIEW:146–158).**

For arbitrary inputs, an $\mathcal{R}^p$-isomorphism preserves the matrix-block size containing $p_1 \cdots p_L$ for each $L$. Consequently it separates the three tested B1/B2 pairs in the table below (reviewer R-3, derived for these instances only).

⟨1⟩1. The product is the rank-one projection onto $\eta^{\otimes L}$ in the unit block of $A_L$, of size $m_L(\mathbf{1}) = \dim \mathrm{Hom}(\mathbf{1}, W_L)$.

⟨1⟩2. A $*$-isomorphism carries its minimal central support to the central support of the image and preserves that full matrix algebra's size. The displayed unequal sizes prohibit every point-preserving isomorphism. QED. No general sufficiency theorem for $\mathcal{R}^p$ follows.

**Proposition 102.7 [derived; any/open; L0 + $H$; R0 inventory]**
**(NOTE:271–296; REVIEW:85–88,173–174).** The record intrinsically retains named stages,

inclusions, centres, block sizes, commutants and products of *already embedded* observables (definition and 102.1). The corrected refinement entry is only $\delta : A_{[j,k]} \hookrightarrow A_{[2j-1,2k-1]}$ (90.1(ii)); the reviewed GAP was its upgrade to an identification. Given $p_j$, insertion-range projections are products over omitted sites; their domain identifications are extra natural marks (102.5). The point itself is extra (102.4); concatenation alone cannot repair its loss (102.2). CA-77 Proposition 77.2's coherent placements and partial isometries are a sufficient richer record; 97.13 additionally recovers supplied $H$ from typed compressions. No minimal sufficient list is proved.

**Proposition 102.8 [conditional; any/open; $\mathsf{L}0 + H$; $\mathsf{R}0$ information] (NOTE:40–41; REVIEW:123–136).** Reviewer R-2 is an *unproved conjectural sketch*, kept conditional here: for arbitrary unitary fusion $\mathcal{E}$ with pointed generating $(O_\mathcal{E}, \eta_\mathcal{E}, H_\mathcal{E})$, a proposed $\mathcal{E} \boxtimes \mathrm{Hilb}_{\mathrm{fd}}(G)$ realisation would use $O = O_\mathcal{E} \boxtimes O_G$ and $\mathrm{id} \otimes \mathrm{Ad}\, U_L$. It requires a sourced construction and compatible factorisation of endomorphisms, intervals and shuffles; no local source for that product/graded-object construction was supplied. This is not a proved extension to every $\mathcal{C}$ possessing an invertible object, and is not used in 102.2–102.7.

## 103.5 Effect on other shards

*Amendment pointers (2026-10-27, round 4; INTEGRATION:48–88).* CA-97's status, summary and 97.16 must record the negative 102.4 and the replacement 102.9 (97.16′); after 97.13, name the point as a necessary carried mark. CA-77's placements remain part of its sufficient record (102.2, 102.5); the converse to CA-90 90.1(iv) fails for $\mathcal{R}$ (102.4). CA-88 gains only the information annotations above. Worklog 023 §4.4 must limit G6-B's proved mechanism to the abelian graded family with trivial associator and compatible $H$: the same category occurs on both sides, and pointed-record recovery of $F, H$ remains open. The pipeline must carry named vacancy projections, or richer coherent placement data determining them. Exact append-after anchors and dated amendment text are supplied in `SHARD_META.md`; these files are not edited here.

## 103.6 What This Does Not Claim

There is no proof that $\mathcal{R}$ forgets $F$ or $H$ once points are named, and no general charge-translation classification. G6-D's different-associator and different-transport tests remain open (REVIEW:87–89); the nonexistent G6-D citation is discarded. Non-natural marks may still admit coherent choices (102.5). No state, Wilson limit, symmetry generators, conformal limit or $\mathsf{R}4$ certificate is constructed.

## 103.7 Examples (Instances Only)

All rows instantiate the statements above with (any/open; $\mathsf{L}0 + H$; $\mathsf{R}0$ information). Grades are additive, unit 0; displayed site lists include multiplicities. Points are unit basis vectors (the first unit copy when repeated); $\chi = 1$ unless specified. Block sizes and failures are the checked data in `checks/review_g6_output.txt`:1–13, not claims for arbitrary $\mathcal{C}$.

| Instance | Exact data and verdict |
|---|---|
| B1, $\mathbb{Z}_2$ | $O = (0,0,1)$, $O' = (0,1,1)$, $s = 1$: 102.4 [negative]; point-block sizes $(2,1)$ at $L = 1$, 102.6 [derived]. |
| B2, $\mathbb{Z}_4$ | $O = (0,1,1,2,3)$, $O' = (0,1,2,2,3)$, $s = 1$; doubled orders $4, 2$: 102.4 [negative]; point-block sizes $(6,7)$ at $L = 2$ [derived]. |
| B2, $\mathbb{Z}_6$ | $O = (0,1,1,2,3,4,5)$, $O' = (0,1,2,2,3,4,5)$, $s = 1$; doubled orders $6, 3$: 102.4 [negative]; point-block sizes $(57, 58)$ at $L = 3$ [derived]. |
| R-1, $\mathbb{Z}_4$ | $O = (0,0,1,2,3)$, $O' = (0,1,2,2,3)$, $s = 2$, $\chi(a,b) = (-1)^{ab}$: transparent, 102.3 [derived]. Plain $\Phi$ fails dilation at $L = 2$ for this site and $\chi = i^{ab}$; also for the B2 site, $s = 1$, $\chi = (-1)^{ab}$ or $i^{ab}$. |
| C1, $\mathbb{Z}_2$ | $O = (0,1,0)$, $\eta = (1,0,0)$, $u = \mathrm{diag}(1,1,-1)$: 102.5 [derived], $\gamma$ fixes points and moves $T, h$. |

## 103.8 Single Next Target: Pointed-Record Faithfulness

**Target 102.9 [conjectural; any/open; $\mathsf{L}0 + H$; $\mathsf{R}0$ information only] (NOTE:319–323; REVIEW:208–224).** This is the reviewer's 97.16′ (INTEGRATION:117–128). For arbitrary pointed tensor-generating $(\mathcal{D}, O, \eta, H)$ in a unitary fusion $\mathcal{C}$, does $\mathcal{R}^p \cong \mathcal{R}'^p$ force a unitary tensor equivalence $\mathcal{D} \simeq \mathcal{D}'$ transporting $O, \eta, H$? A positive route must correct the isomorphism by a point-preserving automorphism so that coherent placements, $h$ and typed compressions $i, j$ are transported up to unitary gauge; then $t_O = j^\dagger \alpha_2(h) i$ (97.13) and 77.2 apply. A negative route requires one simultaneous point-, interval- and dilation-preserving counterexample. Neither the unpointed intertwiner 102.2 nor the non-naturality result 102.5 settles this question.