Overview

5 Unequal Probability Sampling

5.1 The Horvitz–Thompson Estimation Principle

5.2 Variance Estimation

5.3 Probability Proportional to Size (PPS)

5.4 Selecting a PPS Sample

5 Unequal Probability Sampling

Sampling with **unequal** inclusion probabilities (unequal probability sampling):

- is of **interest in its own right**,
- provides the **basis** for formulating a **general theory of sampling**.

For orientation

In Chapters 1–4, we followed a **top-down approach**, i.e., we

- specified a **probability distribution** $\mathbb{P}$ over the set of all possible samples $\mathscr{S}$ and
- thereby defined the **inclusion probabilities** $\pi_i := \mathbb{P}(i \in s)$, $i \in U$.

In what follows, we take a **bottom-up approach**,

- in which the inclusion probabilities π_i are assumed to be **known** and
- serve as the **starting point** for constructing a sampling design with these inclusion probabilities.

Motivation

- We are interested in estimating the population total $t_y = \sum_{i \in U} y_i$.
- Recall from Chapter 4 that estimators under **stratified** simple random sampling (STSI) can, under suitable conditions, be **more efficient** than under SRS.

- Under STSI, the inclusion probabilities n_h/N_h **differ across strata** $h = 1, \ldots, H$ (except under proportional allocation).

- The estimator $\widehat{t}^{\,\mathrm{strat}}$ under STSI **compensates** for the unequal inclusion probabilities across strata by using the **weights** N_h/n_h, i.e., the **inverse inclusion probabilities**:

$$\widehat{t}_y^{\,\mathrm{strat}} = \sum_{h=1}^{H} \frac{N_h}{n_h} \sum_{i \in s_h} y_i.$$

- We now **generalize** this idea as follows:
  - **all** elements $i \in U$ may have **different** inclusion probabilities π_i,
  - each observation is **weighted** by the **inverse** $1/\pi_i$.

5.1 The Horvitz–Thompson Estimation Principle

Definition 11 (Horvitz and Thompson, 1952).

Let p be a sampling design without replacement with **possibly unequal** inclusion probabilities $\pi_i > 0$ $(i \in U)$. The **Horvitz–Thompson** (HT) estimator of the population total t_y is

$$\widehat{t}_y^{\mathrm{HT}} = \sum_{i \in s} \frac{y_i}{\pi_i}. \tag{5.1}$$

- The estimator should more properly be called the **Narain–Horvitz–Thompson** estimator; Narain (1951) proposed it "solely on intuitive grounds" (Ajgaonkar, 1967, p. 55).
- **Alternative** representation: $\widehat{t}_y^{\mathrm{HT}} = \sum_{i \in s} w_i y_i$ with $w_i = \frac{1}{\pi_i}$

Aside (Unbiasedness of the HT estimator).

Let $\mathbb{1}_i$ denote the sample inclusion indicator $\mathbb{1}\{i \in s\}$. It is straightforward to show that the HT estimator is **unbiased** for the population total t_y for **arbitrary** inclusion probabilities $\pi_i > 0$ $(i \in U)$:

$$\mathbb{E}(\widehat{t}_y^{\mathrm{HT}}) = \mathbb{E}\left(\sum_{i \in U} \mathbb{1}_i \frac{y_i}{\pi_i}\right) = \sum_{i \in U} \mathbb{E}(\mathbb{1}_i) \frac{y_i}{\pi_i} = \sum_{i \in U} \pi_i \frac{y_i}{\pi_i} = \sum_{i \in U} y_i = t_y,$$

where we used $\mathbb{E}(\mathbb{1}_i) = \pi_i$.

- The choice of the π_i **does**, however, affect the variance of the HT estimator.
- All estimators discussed so far are **special cases** of the HT estimator (e.g., $\widehat{t}_y$ under SRS or $\widehat{t}_y^{\mathrm{strat}}$ under STSI).

Estimating the Population Mean

- The population mean $\bar{y}_U$ can be estimated using the HT estimator only if N is **known**:

$$\bar{y}^{\,\mathrm{HT}} = \frac{\widehat{t}_y^{\,\mathrm{HT}}}{N}.$$

- If N is **unknown**, the **Hájek estimator** (HJ) can be used (Hájek, 1971, p. 236):

$$\bar{y}^{\,\mathrm{HJ}} = \frac{\sum_{i\in s} y_i/\pi_i}{\sum_{i\in s} 1/\pi_i} = \frac{\sum_{i\in s} w_i y_i}{\sum_{i\in s} w_i} \qquad \text{with} \quad w_i = \frac{1}{\pi_i},$$

$$= \frac{1}{\widehat{N}} \sum_{i\in s} w_i y_i = \frac{\widehat{t}_y^{\,\mathrm{HT}}}{\widehat{N}} \qquad \text{with} \quad \widehat{N} = \sum_{i\in s} w_i.$$

- The estimator $\widehat{N}$ can be viewed as the HT estimator of N with $y_i \equiv 1$, i.e., $\widehat{N} = \sum_{i \in s} 1 \, w_i$.

- The **Hájek estimator**:
  - is a **ratio estimator**

$$\bar{y}^{\,\mathrm{HJ}} = \frac{\widehat{t}_y^{\,\mathrm{HT}}}{\widehat{N}}$$

    and is generally **biased** in finite samples, although,
  - the bias is typically small for sufficiently **large samples**.
  - is frequently used in practice even when N is **known**.

5.2 Variance Estimation

Definition 12 (Second-order inclusion probabilities).

Let p be a sampling design without replacement with $\pi_i > 0$ for all $i \in U$. The probability that elements i and j are both included in the sample s is called the **second-order inclusion probability** and is defined as

$$\pi_{ij} := \mathbb{P}(i, j \in s) = \mathbb{P}(\mathbb{1}_i \mathbb{1}_j = 1) \qquad (i, j \in U).$$

- For the **special case** $i = j$, $\pi_{ii} = \pi_i$ for all $i \in U$.
- For a population of N elements, the matrix of second-order inclusion probabilities is of size $(N \times N)$. Because it is **symmetric**, it has

$$\underbrace{N}_{\pi_{ii}=\pi_i} + \underbrace{\frac{N(N-1)}{2}}_{\pi_{ij}, i<j}$$ nonredundant entries.

Definition 13 (Covariance).

Let p be a sampling design without replacement with $\pi_i > 0$ for all $i \in U$. The **covariance** Δ_{ij} between the inclusion indicators $\mathbb{1}_i = \mathbb{1}\{i \in s\}$ and $\mathbb{1}_j = \mathbb{1}\{j \in s\}$, $i, j \in U$, is defined as

$$\Delta_{ij} := \text{cov}(\mathbb{1}_i, \mathbb{1}_j) = \pi_{ij} - \pi_i \pi_j.$$

- Depending on the design, the covariances may be **zero**, **positive**, or **negative**.
- For the variance, $\text{var}(\mathbb{1}_i) = \pi_i(1 - \pi_i)$ for all $i \in U$.

Example 18 (SRS).

Under SRS with sample size n $(0 < n < N)$, the inclusion probabilities are $\pi_i = n/N$ for all $i \in U$ and

$$\pi_{ij} = \frac{n(n-1)}{N(N-1)} = \pi_i \pi_j \frac{N(n-1)}{n(N-1)} \qquad (\text{for } i \neq j,\ i,j \in U).$$

The **covariance** between the inclusion indicators $\mathbb{1}_i$ and $\mathbb{1}_j$ is

$$\Delta_{ij} = \pi_{ij} - \pi_i \pi_j = \frac{n(n-1)}{N(N-1)} - \left(\frac{n}{N}\right)^2 = -\frac{f(1-f)}{N-1} \qquad (\text{for } i \neq j, i,j \in U)$$

with $f = n/N$. Since $0 < n < N$, we have $0 < f < 1$. Hence the covariance is strictly **negative** for all $i \neq j$.

- Let p be a sampling design without replacement with $\pi_i > 0$ for all $i \in U$. The theoretical **variance of the HT estimator** is

$$\mathsf{var}\big(\widehat{t}^{\,\mathrm{HT}}\big) = \sum_{i \in U} \sum_{j \in U} \Delta_{ij} \frac{y_i y_j}{\pi_i \pi_j}$$

- If $\pi_{ij} > 0$ for all $i, j \in U$, then $\mathsf{var}\big(\widehat{t}^{\,\mathrm{HT}}\big)$ can be estimated by the unbiased **Horvitz–Thompson variance estimator** (Horvitz and Thompson, 1952):

$$\widehat{\mathsf{var}}_{HT}\big(\widehat{t}^{\,\mathrm{HT}}\big) = \sum_{i \in s} \sum_{j \in s} \frac{\Delta_{ij}}{\pi_{ij}} \frac{y_i y_j}{\pi_i \pi_j}.$$

Special Case: Fixed Sample Size

- For a sampling design with **fixed sample size** n, $\sum_{j \in U} \Delta_{ij} = 0$ for all $i \in U$.
- Hence, the variance of the HT estimator can also be written as

$$\mathrm{var}(\widehat{t}^{\,\mathrm{HT}}) = -\frac{1}{2} \sum_{i \in U} \sum_{j \in U} \Delta_{ij} \left(\frac{y_i}{\pi_i} - \frac{y_j}{\pi_j} \right)^2 .$$

- If $\pi_{ij} > 0$, this leads to the **Sen–Yates–Grundy** (SYG; Sen, 1953; Yates and Grundy, 1953) variance estimator

$$\widehat{\mathrm{var}}_{SYG}(\widehat{t}^{\,\mathrm{HT}}) = -\frac{1}{2} \sum_{i \in s} \sum_{j \in s} \frac{\Delta_{ij}}{\pi_{ij}} \left(\frac{y_i}{\pi_i} - \frac{y_j}{\pi_j} \right)^2 .$$

- If $\Delta_{ij} \leq 0$ for all $i \neq j$, $i, j \in U$, the SYG estimator is **nonnegative**.
- The SYG estimator is **numerically more stable** than $\widehat{\text{var}}_{HT}$, particularly when the coefficient of variation of the y_i/π_i is small.

Aside (Important identities).

For sampling designs p with **fixed sample size** n,

$$\sum_{i \in U} \pi_i = n \qquad \text{and} \qquad \sum_{j \in U} \pi_{ij} = n\pi_i \quad (i \in U).$$

Hence,

$$\sum_{j \in U} \Delta_{ij} = \sum_{j \in U} (\pi_{ij} - \pi_i \pi_j) = 0 \qquad (i \in U).$$

Aside (cont.).

- In other words,

$$\text{fixed } n \implies \sum_j \pi_{ij} = n\pi_i \implies \sum_j \Delta_{ij} = 0 \implies \text{SYG representation}$$

- The above identities are useful for checking inclusion probabilities and lead to the Sen–Yates–Grundy representation of the HT variance.

Approximation of Second-Order Inclusion Probabilities

- Brewer (2000) proposed the following **approximation** of second-order inclusion probabilities for *high-entropy* designs:

$$\pi_{ij} \approx \pi_i \pi_j \left(\frac{c_i + c_j}{2} \right) \qquad (i, j \in U,\, i \neq j), \tag{5.2}$$

  where the choice of the constants c_i, $i \in U$, is discussed below.

- Systematic sampling, for example, has low entropy.

- Based on the approximation in (5.2), Brewer (2000) gave the following approximation to the theoretical **variance of the HT estimator**:

$$\text{var}_{\text{A}}\left(\widehat{t}^{\text{HT}}\right) = \sum_{i \in U} \pi_i (1 - c_i \pi_i) \left(\frac{y_i}{\pi_i} - \frac{t_y}{n} \right)^2.$$

- He also recommended the following **variance estimator**:

$$\widehat{\mathrm{var}}_{\mathrm{A}}\big(\widehat{t}^{\,\mathrm{HT}}\big) = \sum_{i \in s} \left(\frac{1}{c_i} - \pi_i\right) \left(\frac{y_i}{\pi_i} - \frac{\widehat{t}_y^{\,\mathrm{HT}}}{n}\right)^2.$$

  - The variance estimator is asymptotically unbiased for $\mathrm{var}_{\mathrm{A}}\big(\widehat{t}^{\,\mathrm{HT}}\big)$.
  - $\widehat{\mathrm{var}}_{\mathrm{A}}\big(\widehat{t}^{\,\mathrm{HT}}\big)$ is easy to compute.

- Brewer (2000) proposed **three variants** for choosing the **constants** c_i. We use Variant 1:

$$c_i = \frac{n-1}{n-\pi_i} \qquad (i \in U).$$

5.3 Probability Proportional to Size (PPS)

Example 19 (Oracle, Population U14).

i:	1	2	3	4	5	6	7	8	9	10	11	12	13	14
y_i:	175	143	116	85	150	92	108	67	121	99	118	259	75	142
π_i:	$\frac{2}{5}$	$\frac{286}{875}$	$\frac{232}{875}$	$\frac{34}{175}$	$\frac{12}{35}$	$\frac{184}{875}$	$\frac{216}{875}$	$\frac{134}{875}$	$\frac{242}{875}$	$\frac{198}{875}$	$\frac{236}{875}$	$\frac{74}{125}$	$\frac{6}{35}$	$\frac{284}{875}$

A probability sample of size $n = 4$ is selected without replacement from U14. The **realized** sample s is

$$y_1 = 175, \quad y_2 = 143, \quad y_8 = 67, \quad y_{14} = 142$$
$$\pi_1 = \frac{2}{5}, \quad \pi_2 = \frac{286}{875}, \quad \pi_8 = \frac{134}{875}, \quad \pi_{14} = \frac{284}{875}.$$

Example (Cont.).

- For the realized sample, the HT estimator gives

$$\widehat{t}_y^{\mathrm{HT}} = \sum_{i \in s} \frac{y_i}{\pi_i} = 1'750.$$

- The estimate $\widehat{t}_y^{\mathrm{HT}}$ equals the true population total t_y.
- The variance of the estimator is zero, $\mathrm{var}(\widehat{t}_y^{\mathrm{HT}}) = 0$; hence there is **no sampling uncertainty**.
- In this application, the HT estimator has zero variance because the inclusion probabilities π_i were constructed by an **oracle**, assuming that all y_i $(i \in U)$ were already **known before sampling**.
- Clearly, this makes the idea of estimation **absurd**.

Example (Cont.).

- We can nevertheless learn something from this **absurd** situation.
- The first-order inclusion probabilities π_i were chosen **proportional** to the y_i:

$$\pi_i = \frac{ny_i}{\sum_{i \in U} y_i} = \frac{ny_i}{t_y} \qquad (\text{for all } i \in U),$$

so that the HT estimator becomes

$$\widehat{t}_y^{\mathrm{HT}} = \sum_{i \in s} \frac{y_i}{\pi_i} = \sum_{i \in s} \frac{y_i}{ny_i} t_y = t_y.$$


- Let x_i be an **auxiliary variable** known for **all** $i \in U$.
- For example, $x_i =$ number of employees in business i (auxiliary variable known from a register), and $y_i =$ revenue (variable of interest, unknown).

- Assume that y_i and x_i are strongly **positively correlated**.
- We consider a fixed-size sampling design. If $y_i \propto x_i$, then $\text{var}(\widehat{t}_y^{\text{HT}}) = 0$; see the artificial situation in Example 19.
- The first-order **inclusion probabilities** are chosen proportional to size (PPS), i.e.,

$$\pi_i = \frac{n x_i}{\sum_{i \in U} x_i} \qquad (i \in U),$$

  provided that $0 < \pi_i \leq 1$ for all $i \in U$.
- Variable x_i is called the **measure of size** (e.g., number of employees).
- If proportionality is not exact, PPS sampling can still yield **efficiency gains** over SRS when x_i is strongly related to y_i.

Example (Data from Example 19)



5.4 Selecting a PPS Sample

- Hanif and Brewer (1980) classified **47** PPS sampling procedures without replacement; for more recent literature, see Tillé (2006).
- Following their classification, we impose the following **requirements** on PPS designs (for all $i, j \in U$):
  - **fixed** sample size n;
  - the π_i are strictly **proportional** to x_i;
  - **unbiased** variance estimation: $\pi_{ij} > 0$ for all $i \neq j$;
  - the SYG variance estimator is **nonnegative**: $\Delta_{ij} \leq 0$ for all $i \neq j$.
- The sampling procedure of Brewer (1963, 1975) satisfies these requirements (first procedure).

Sampling Procedure of Brewer (1963); Sample Size $n = 2$

Assumption: $x_i < t_x/2$ for all $i \in U$. Then

$$\pi_i = \frac{2x_i}{t_x} < 1 \qquad (i \in U).$$

1) Select the **first element** with probability

$$p_i = \frac{c_i}{\sum_{i \in U} c_i}, \qquad c_i = \frac{x_i(t_x - x_i)}{t_x(t_x - 2x_i)} \qquad (i \in U).$$

Denote the selected element by k; it is not replaced.

2) Select the **second element** with conditional probability

$$p_{j|k} = \frac{x_j}{t_x - x_k}, \qquad j \in U \setminus \{k\}.$$

- Brewer (1963) derived a formula for the **second-order inclusion probabilities**.
- In the **general case**, i.e., when $n > 2$,
  - the prescribed PPS inclusion probabilities

$$\pi_i = \frac{nx_i}{t_x}$$

    require $x_i \leq t_x/n$ for all $i \in U$;
  - units with $\pi_i = 1$ are **certainty units** and are included in the sample with probability one;
  - after removing certainty units, the inclusion probabilities for the remaining units must be recalculated;
  - sample selection is substantially more **involved**; see Brewer (1975).

References I

Ajgaonkar, S. G. P. (1967). Unbiased Estimators of the Variance of the Narain, Horvitz and Thompson Estimator, *Sankhyā. Series A*, **29**, 55–60.

Brewer, K. R. W. (1963). A model of systematic sampling with unequal probabilities, *Australian Journal of Statistics*, **5**, 5–13, DOI:10.1111/j.1467-842X.1963.tb00132.x.

Brewer, K. R. W. (1975). A Simple Procedure for Sampling πpswor, *Australian Journal of Statistics*, **17**, 166–172, DOI:10.1111/j.1467-842X.1975.tb00954.x.

Brewer, K. R. W. (2000). Deriving and estimating an approximate variance for the Horvitz–Thompson estimator using only first order inclusion probabilities, in: *Proceedings of the Second International Conference on Establishment Surveys*, 1417–1422, Buffalo, New York: Session 30.

Hájek, J. (1971). Comment on D. Basu: An Essay on the Logical Foundations of Survey Sampling, Part 1, in: *Foundations of Statistical Inference. Proceedings of the Symposium on the Foundations of Statistical Inference Prepared Under the Auspices of the René Descartes Foundation and Held at the Department of Statistics, University of Waterloo, Ontario, Canada, from March 31 to April 9, 1970* ed. by Godambe, V. P. and Sprott, D. A. Toronto/ Montreal: Holt, Rinehart and Winston of Canada, 236, https://archive.org/details/foundationsofsta0000symp.

References II

Hanif, M. and Brewer, K. R. W. (1980). Sampling with Unequal Probabilities without Replacement: A Review, *International Statistical Review*, **48**, 317–335, DOI:10.2307/1402944.

Horvitz, D. G. and Thompson, D. J. (1952). A Generalization of Sampling Without Replacement From a Finite Universe, *Journal of the American Statistical Association*, **47**, 663–685, DOI:10.1080/01621459.1952.10483446.

Narain, R. D. (1951). On sampling without replacement with varying probabilities, *Journal of the Indian Society of Agricultural Statistics*, **3**, 169–174.

Sen, A. R. (1953). On the estimate of the variance in sampling with varying probabilities, *Journal of the Indian Society of Agricultural Statistics*, **5**, 119–127.

Tillé, Y. (2006). *Sampling Algorithms*, New York: Springer.

Yates, F. and Grundy, P. M. (1953). Selection without replacement from within strata with probability proportional to size, *Journal of the Royal Statistical Society. Series B*, **15**, 253–261, DOI:10.1111/j.2517-6161.1953.tb00140.x.

ETH zürich

My contact details at ETH
Lecturer at the Department of Mathematics
E-mail: tobias.schoch@math.ethz.ch

Primary affiliation
University of Applied Sciences Northwestern Switzerland FHNW
School of Business – Institute ICC
Riggenbachstrasse 16
CH-4600 Olten

Prof. Dr. Tobias Schoch
E-mail: tobias.schoch@fhnw.ch
Phone (direct): +41 (0)62 957 21 02

Version of October 1, 2026