

Optimal discretization in Banach spaces:
Residual minimization, nonlinear Petrov–Galerkin, and monotone mixed methods¹

I. Muga^a and K.G. van der Zee^b

^aPontificia Universidad Católica de Valparaíso, Instituto de Matemáticas, Chile
ignacio.muga@uvc.cl

^b(presenter) The University of Nottingham, School of Mathematical Sciences, UK
kg.vanderzee@nottingham.ac.uk

Is it possible to obtain numerically the best approximation to solutions of linear operator equations in general Banach spaces? In this contribution, we address this question by considering nonstandard Petrov–Galerkin discretisations. We build on ideas of residual minimization and the recent theory of optimal Petrov–Galerkin methods in Hilbert space settings [1–4]. We propose an optimal discretization method in the setting of certain Banach spaces, proof a priori error estimates, and present some numerical experiments [5].

References

- [1] J. L. Guermond, *A finite element technique for solving first-order PDEs in L^p* , SIAM J. Numer. Anal., 42 (2004), pp. 714–737.
- [2] L. Demkowicz, and J. Gopalakrishnan, *A class of discontinuous Petrov–Galerkin methods. Part I: The transport equation*, Comput. Methods Appl. Mech. Engrg., 199 (2010), pp. 1558–1572.
- [3] J. Zitelli, I. Muga, L. Demkowicz, J. Gopalakrishnan, D. Pardo and V. M. Calo, *A class of discontinuous Petrov–Galerkin methods. Part IV: The optimal test norm and time-harmonic wave propagation in 1D*, J. Comput. Phys., 230 (2011), pp. 2406–2432.
- [4] W. Dahmen, C. Huang, C. Schwab, and G. Welper, *Adaptive Petrov–Galerkin methods for first order transport equations*, SIAM J. Numer. Anal., 50 (2012), pp. 2420–2445.
- [5] I. Muga, K. G. van der Zee, *Discretization of linear problems in Banach spaces: Residual minimization, nonlinear Petrov–Galerkin, and monotone mixed methods*, in preparation.

¹Abstract for “New directions in numerical computation: In celebration of Nick Trefethen’s 60th birthday”, 25–28 August 2015, Oxford, UK