

EXTREMAL PROBLEMS FOR POLYNOMIALS INITIATED BY NUMERICAL COMPUTATIONS

Heinz-Joachim Rack

Steubenstr. 26a, 58097 Hagen, Germany, heinz-joachim.rack@drrack.com

Abstract.

Polynomial-related topics of numerical analysis give rise to challenging theoretical optimization problems. We will discuss three examples, and will provide explicit solutions:

Problem 1 (*Condition Number of Polynomials in Power Form*): Determine, in the sense of [1], the condition number γ of real polynomials P_n of degree $\leq n$ represented on $\mathbf{I} = [-1, 1]$ in the power form, i.e. $P_n(x) = \sum_{k=0}^n a_k x^k$. Let T_n with $T_n(x) = \sum_{k=0}^n t_{n,k} x^k$ denote the n -th Chebyshev polynomial of the first kind relative to \mathbf{I} , see [10].

Solution: Gautschi [1, Theorem 3.1] provides a formula for γ which requires to determine three maxima: $\gamma = (n+1) \max\{\max_k |t_{n,k}|, \max_k |t_{n-1,k}|\}$. We economize it by avoiding to compute *all* $n+1$ nonzero absolute coefficients of $T_n + T_{n-1}$, and to search among them for the largest one. For we have identified in [6] the index $k = k^* = k^*(n)$ with $|t_{n,k^*}| = \max_k |t_{n,k}|$, so that only k^* and *one* coefficient of T_n need to be computed, since $\max_k |t_{n,k}| \geq \max_k |t_{n-1,k}|$. In this way we are able to express $\gamma = \gamma(n) = (n+1)|t_{n,k^*(n)}|$ as an explicit function of n alone. It will be specified in the talk, along with its generalization to zero-symmetric intervals $[-\omega, \omega]$ with $\omega > 0$ (replacing \mathbf{I}).

Problem 2 (*Optimal Nodes for Quadratic and Cubic Lagrange Interpolation*): Determine analytically, for the quadratic ($n = 2$) and cubic ($n = 3$) case of Lagrange interpolation with nodes $x_1 < x_2 < x_3$ resp. $x_1 < x_2 < x_3 < x_4$ in \mathbf{I} , all infinitely many (zero-symmetric and zero-asymmetric) optimal node configurations which lead to the minimal Lebesgue constant $\Lambda_3^* = 1.25$ resp. Λ_4^* .

Solution: The case $n = 2$ has been settled in [11], and alternatively in [7]. The value of Λ_4^* and the optimal canonical nodes $-1 = x_1 < x_2 < x_3 < x_4 = 1$ we have explicitly determined in [3], [4]. In the talk we will reveal, based on [8], *all* optimal node configurations $-1 \leq x_1 < x_2 < x_3 < x_4 \leq 1$.

Finally, we will turn to extremal problems initiated by the evaluation of P_n : Worst case scenarios of the naïve and Horner's algorithm lead to V. A. Markov-type [2, p. 246] extremal coefficient functional problems ([5], [9]), and the search for the most stable evaluation algorithm leads to the barycentric interpolation formula [12, p. 38]. In the talk we will amplify the result given in [9] which is referenced to in [10].

References

- [1] W. Gautschi, *The condition of polynomials in power form*, Math. Comp. 33 (1979), 343-352
- [2] W. Markoff (V. A. Markov), *Über Polynome ...*, Math. Ann. 77 (1916), 213-258 (Russian 1892)
- [3] H.-J. Rack, *An example of optimal ...*, Int. J. Math. Educ. Sci. Technol. 15 (1984), 355-357
- [4] H.-J. Rack, *An example of optimal ... revisited*, Springer Proc. Math. Stat. 41 (2013), 117-120
- [5] H.-J. Rack, *Abschätzung der Teilsummen reeller Polynome*, Math. Z. 182 (1983), 549-552
- [6] H.-J. Rack, *On the length and the height ...*, East J. Approx. 16 (2010), 35-91
- [7] H.-J. Rack & R. Vajda, *On optimal quadratic Lagrange ...*, Serdica J. Comput. 8 (2014), 71-96

- [8] H.-J. Rack & R. Vajda, *On optimal cubic . . .*, Stud. Univ. Babeş-Bolyai, Math. 60 (2015), no. 2
- [9] M. Reimer, *Bounds for the Horner sums*, SIAM J. Numer. Anal. 5 (1968), 461-469
- [10] Th. J. Rivlin, *Chebyshev Polynomials*, 2nd Edition, J. Wiley & Sons, New York, 1990
- [11] F. Schurer, *A remark on extremal sets . . .*, Stud. Sci. Math. Hung. 9 (1974), 77-79
- [12] L. N. Trefethen, *Approximation Theory and Approximation Practice*, SIAM, Phila., 2013