

## **Toward integration of high-frequency oscillators at logarithmic cost: QTT tensor approximation of discretized functions**

**Boris N. Khoromskij**

Max-Planck-Institute for Mathematics in the Sciences,  
Inselstr. 22-26, D-04103 Leipzig, Germany  
*bokh@mis.mpg.de*; <http://personal-homepages.mis.mpg.de/bokh>

*This talk is dedicated to Prof. Nick Trefethen on occasion of his 60th birthday.*

The well established approach to the efficient approximation and integration of “complicated” functions, based on the Chebyshev interpolation (Chebfun1, Chebfun2), was developed in the last decade by Nick Trefethen and his group [1].

The recent alternative approach to low complexity representation of multivariate functions is based on the principle of separation of variables [2]. The novel method of quantized tensor approximation (QTT) is proven to provide the compressed representation of logarithmic complexity to the grid-based discretizations of functions and operators [3]. This enables reformulation of the standard discretization schemes for steady-state and dynamical problems by using the low-rank quantized tensor approximation and then solving the resultant multi-dimensional systems with the logarithmic complexity in the initial problem size. Several examples of successful applications of the QTT-based tensor numerical methods will be discussed.

We demonstrate how the grid-based QTT tensor approximation applies to the integration of multidimensional or/and highly-oscillating functions, say, many-electron integrals, integrals with the large lattice sum of interaction potentials [4] or with the complicated high-frequency oscillators [5] including some examples from Chebfun demonstrations. Other topic addresses the rank-structured QTT tensor representation of the electron density and the Fock matrix in electronic structure calculations for large 3D  $L \times L \times L$  lattice systems [6].

## **References**

- [1] Rodrigo Platte and Nick Trefethen. *Chebfun: A New Kind of Numerical Computing*. No. NA-08/13. Oxford University Computing Laboratory. October, 2008.
- [2] Boris N. Khoromskij. *Tensor Numerical Methods for High-dimensional PDEs: Basic Theory and Initial Applications*. E-preprint arXiv:1408.4053, 2014. ESAIM: Proceedings 2014 (to appear).
- [3] B.N. Khoromskij.  *$O(d \log N)$ -Quantics Approximation of  $N$ -d Tensors in High-Dimensional Numerical Modeling*. J. Constr. Approx. v. 34(2), 257-289 (2011); (Preprint 55/2009 MPI MiS, Leipzig 2009).
- [4] V. Khoromskaia and B. N. Khoromskij. *Grid-based lattice summation of electrostatic potentials by assembled rank-structured tensor approximation*. Comp. Phys. Communications, 185 (2014), pp. 3162-3174.
- [5] Boris N. Khoromskij, and A. Veit. *Efficient computation of highly oscillatory integrals by using QTT tensor approximation*. E-preprint arXiv:1408.5224, 2014 (submitted).
- [6] V. Khoromskaia, and B.N. Khoromskij. *Tensor Approach to Linearized Hartree-Fock Equation for Lattice-type and Periodic Systems*. E-preprint arXiv:1408.3839, 2014 (submitted).