

# Stable polefinding and rational least-squares fitting via eigenvalues

*Shinji Ito and Yuji Nakatsukasa*

## Abstract

A common way of finding the poles of a meromorphic function  $f$ , whose explicit expression is unknown but  $f(z)$  can be evaluated at any given  $z$ , is to interpolate  $f$  by a rational function  $\frac{p}{q}$  such that  $\frac{p(\gamma_i)}{q(\gamma_i)} = f(\gamma_i)$  at sample points  $\{\gamma_i\}$ , and then find the roots of  $q$  [1, 3]. This is a two-step process, and the type of the rational interpolant is usually required as inputs. Many other algorithms for polefinding and rational interpolation have been proposed [2, 4, 5, 6], but their numerical stability has remained largely uninvestigated; for example, many methods become unstable when a pole lies very close to a sample point.

This talk describes an algorithm with the following three features: (i) it automatically detects the appropriate type of the rational approximant, thereby allowing the user to input just the function  $f$ , (ii) it finds the poles in a one-step fashion via a generalized eigenvalue problem involving matrices constructed directly from the sampled values  $f(\gamma_i)$ , and (iii) it computes rational approximants  $\hat{p}, \hat{q}$  in a numerically stable manner, in that  $(\hat{p} + \Delta p)/(\hat{q} + \Delta q) = f$  at the sample points with small  $\Delta p, \Delta q$ , making it the first rational interpolation algorithm with guaranteed numerical stability. Moreover, our algorithm executes an implicit polynomial change-of-basis by the QR factorization for improved accuracy, and allows for oversampling combined with a least-squares fitting as in [3]. Through experiments we illustrate the resulting accuracy and stability, which can significantly outperform existing algorithms.

## References

- [1] A. P. Austin, P. Kravanja, and L. N. Trefethen. Numerical algorithms based on analytic function values at roots of unity. *SIAM J. Numer. Anal.*, 52(4):1795–1821, 2014.
- [2] A. P. Austin and L. N. Trefethen. Computing eigenvalues of real symmetric matrices with rational filters in real arithmetic. preprint.
- [3] P. Gonnet, R. Pachón, and L. N. Trefethen. Robust rational interpolation and least-squares. *Electronic Transactions on Numerical Analysis*, 38:146–167, 2011.
- [4] P. Kravanja and M. Van Barel. A derivative-free algorithm for computing zeros of analytic functions. *Computing*, 63(1):69–91, 1999.
- [5] P. Kravanja and M. Van Barel. *Computing the zeros of analytic functions*. Number Lecture Notes in Math. 1727. Springer, 2000.
- [6] T. Sakurai and H. Sugiura. A projection method for generalized eigenvalue problems using numerical integration. *J. Comput. Appl. Math.*, 159(1):119–128, 2003.