

Coherence time in biological oscillator assemblies bounds the rate of state registration

Ian Todd

Sydney Medical School, University of Sydney

Sydney, NSW, Australia

`itod2305@uni.sydney.edu.au`

January 26, 2026

Abstract

Cognitive and biological processing depends on coherent oscillations across distributed assemblies. Yet the physical principles determining how fast these assemblies can synchronize—and thus how fast neural processes associated with cognition can proceed—remain incompletely formalized. We derive a quantitative framework showing that coherence time in coupled oscillator networks scales exponentially with coordination depth. For M semi-independent modules requiring phase alignment within tolerance ε at coherence r , the minimum interval between irreversible state registrations (“commits”) is dominated by rare-event first-passage time into low-measure alignment sets on the configuration manifold. This produces a fundamental speed-flexibility trade-off: increasing coordination depth expands combinatorial flexibility but slows commits exponentially; tighter coherence accelerates synchronization but restricts dynamics to low-dimensional attractors. We situate this coordination limit within a hierarchy of physical bounds—quantum speed limits, Landauer-Margolus-Levitin thermodynamic

constraints, and energy-time uncertainty relations—showing that in biological neural systems, these quantum and thermodynamic bounds are typically non-binding; the dominant constraint is the time required for distributed oscillatory assemblies to achieve sufficient phase coherence. While neural oscillators provide the most empirically accessible measurements, the framework applies to any biological system whose function requires coordinated phase relations and sparse commitment events, from chromosomal segregation to circadian networks. We provide parameter estimation protocols and testable predictions including alpha entrainment effects on temporal acuity and dual-loop dissociations under arousal.

Keywords: biological oscillations, coherence time, coordination depth, temporal resolution, quantum speed limits, Landauer bound, coupled oscillators

1 Introduction

1.1 Processing speed in biological oscillator assemblies

A fundamental insight from systems neuroscience is that cognition emerges from coherent oscillations across neural assemblies, not merely from individual spike rates. Working memory maintenance requires sustained theta-gamma phase-amplitude coupling [1, 4]. Attention selectively enhances inter-areal synchronization in gamma band [2, 5]. Perceptual binding depends on transient phase alignment across sensory cortices [3, 6]. Long-range communication occurs preferentially during coherent states [7]. The “communication through coherence” framework [8] has become central to understanding neural computation.

Yet despite extensive empirical characterization, the physical principles governing *how fast* distributed assemblies can achieve coherence—and thus how quickly the neural processes underlying cognitive operations can proceed—remain incompletely formalized. Why does perceptual binding require 30–50 ms rather than 3 ms or 300 ms? Why do larger assemblies integrating more information process more slowly? What determines the relationship

between oscillation frequency and temporal acuity?

1.2 Coherence time as a physical bound on processing rate

Previous work established that living intelligence is an emergent property of high-dimensional, self-sustaining dynamical systems that maintain coherence against decoherence [9]. The present manuscript addresses the complementary question: given such a coherent high-dimensional system, what sets the fastest timescale on which it can register change and update its state?

We formalize this as the minimum interval between irreversible “commit” events—transitions in which distributed dynamics become readable to downstream mechanisms. This commit interval is bounded by a coordination term (coherence time as a first-passage problem on a product manifold) together with quantum, noise, and power limits.

Operational definition of commits in neural systems. A commit is operationally detected when distributed oscillatory dynamics cross a threshold that enables downstream readout. Empirical signatures include: (i) phase-locking index exceeding threshold across specified modules for a minimum duration, (ii) onset of stable cross-frequency coupling (e.g., theta-gamma PAC), (iii) classifier-detectable state transition in downstream neural populations. We do not claim all neural computation reduces to discrete commits; rather, we claim that the *fastest update of downstream state* is bounded by the time required for multi-module phase alignment. The framework addresses temporal resolution limits, not the full richness of neural dynamics.

Definition (Ontological dimensionality). Throughout this work, “dimensionality” refers to a physical property of the system, not a mathematical measure. Ontological dimensionality is a quantitative measure of effective connectivity: how many independent directions of coordinated variation are accessible to a system given its interaction structure. High-dimensional states correspond to weakly constrained dynamics in which many components vary semi-independently; low-dimensional states correspond to strongly coupled dynamics

restricted to a small number of collective modes. Dimensionality in this sense is determined not by component count, but by the rank of the system’s interaction structure. Different measurement schemes (participation ratios, manifold dimension, information-geometric metrics) provide projections of this underlying physical property, but do not constitute it [10].

This ontological dimensionality is state-dependent: the same physical system may occupy high- or low-dimensional regimes depending on coupling strength and task demands. Coherence time depends on ontological dimensionality—specifically, how many semi-independent constraints must be simultaneously satisfied for a commit to occur—not on any particular estimator.

1.3 Structure of this work

We first derive the coherence time formula from principles of phase synchronization in coupled oscillator networks (§2.1), then embed it in a unified temporal resolution bound combining quantum, noise, and power constraints (§2.2). Section 2.3 situates our framework explicitly within the quantum and thermodynamic limits of computation, addressing the Landauer bound, Margolus-Levitin quantum speed limit, and energy-time uncertainty relation. Section 3 applies the framework to neural systems, demonstrating quantitative agreement with perceptual binding windows, tachypsychia dissociations, and metabolic scaling. Section 4 discusses the speed-flexibility trade-off, delay embedding as a preservation mechanism, generalization beyond neural systems, and testable predictions.

2 Theory and Methods

2.1 The unified temporal resolution bound

We model biological temporal processing as a sequence of *commits*—thermodynamically irreversible events that register high-dimensional internal state as low-dimensional output. A commit requires: (1) dimensional collapse from distributed state (D_{eff} dimensions) to

registered outcome (few dimensions), (2) dissipation of at least $k_B T \ln 2$ per bit (Landauer bound), and (3) creation of a persistent measurement influencing future dynamics.

The minimum time between commits is bounded below by four physical constraints:

$$\tau_{\text{eff}} \gtrsim \max[\tau_{\text{QSL}}, \tau_{\text{SNR}}, \tau_{\text{coh}}, \tau_{\text{power}}] \quad (1)$$

This is a **lower bound**: the effective commit interval must be at least as long as the slowest constraint. When these constraints act on largely independent subsystems and can be satisfied in parallel, the bound is approximately tight and the max operation gives the dominant timescale. The four terms are:

Quantum speed limits (τ_{QSL}). The Mandelstam-Tamm and Margolus-Levitin bounds [13, 14] establish

$$\tau_{\text{QSL}} = \max\left[\frac{\hbar}{2\Delta E}, \frac{\pi\hbar}{2\langle E \rangle}\right]$$

For biological temperatures ($\Delta E \sim k_B T$), $\tau_{\text{QSL}} \sim 10^{-13}$ s.

Signal-to-noise limit (τ_{SNR}). Matched-filter detection in additive white Gaussian noise (two-sided power spectral density N_0 , units: power/Hz) over bandwidth B achieves $\text{SNR}(T) = 2P_s T / N_0$ where P_s is signal power and T is integration time [19]. Reaching detection threshold $\rho_{\text{min}} \sim 5$ –10 requires

$$\tau_{\text{SNR}} = \frac{\rho_{\text{min}} N_0}{2P_s}$$

Power limit (τ_{power}). Dimensional collapse by $\Delta\mathcal{D}$ at dimensional chemical potential $\lambda_{\mathcal{D}} \sim k_B T_{\text{eff}}$ costs energy $\sim \lambda_{\mathcal{D}} \Delta\mathcal{D}$. At sustained power P , commit rate is bounded by

$$\tau_{\text{power}} = \frac{\lambda_{\mathcal{D}} \Delta\mathcal{D}}{P}$$

Coherence time (τ_{coh}). We model commits as threshold crossings requiring M semi-independent modules to achieve simultaneous phase alignment within tolerance ε . Let r be

the mean resultant length (Kuramoto order parameter), and $\kappa(r)$ the von Mises concentration with $r = I_1(\kappa)/I_0(\kappa)$ [21].

Definition of alignment event. We define alignment as **anchored to a reference module**: all modules must fall within $\varepsilon/2$ of a designated reference phase. Formally:

$$A_\varepsilon = \{|\theta_i - \theta_1| < \varepsilon/2 \text{ for } i = 2, \dots, M\}$$

The alignment probability depends on whether the reference is fixed or random:

Case A (anchored/pacemaker reference): If θ_1 is fixed or tightly controlled (e.g., by a pacemaker oscillator, gating signal, or first-crossing module), then

$$P(A_\varepsilon) = \prod_{i=2}^M P(|\theta_i - \theta_1| < \varepsilon/2) = p_1(\varepsilon, \kappa)^{M-1}$$

where $p_1(\varepsilon, \kappa) = \int_{-\varepsilon/2}^{\varepsilon/2} \frac{e^{\kappa \cos \theta}}{2\pi I_0(\kappa)} d\theta$ is the single-variable window probability.

Case B (random reference): If θ_1 is itself drawn from the same von Mises distribution, then $P(A_\varepsilon) = \mathbb{E}_{\theta_1}[q(\theta_1)^{M-1}]$ where $q(\theta_1) = P(|\theta - \theta_1| < \varepsilon/2)$. In the small-window limit, rotational symmetry ensures the scaling remains $\propto \varepsilon^{M-1}$, though the constant differs.

In both cases:

$$P(A_\varepsilon) \propto p_1(\varepsilon, \kappa)^{M-1} \tag{2}$$

Small- ε approximation. For small windows, the von Mises density near the mode gives $p_1(\varepsilon, \kappa) \approx \varepsilon \cdot e^\kappa / (2\pi I_0(\kappa))$, yielding:

$$\tau_{\text{coh}} \approx \frac{1}{\Delta\omega} \left(\frac{2\pi I_0(\kappa)}{\varepsilon e^\kappa} \right)^{\alpha(M-1)}$$

This makes the $\varepsilon^{-(M-1)}$ scaling and the “exponential in M ” dependence explicit.

The exponent is $(M - 1)$, not M . This arises from quotient geometry: the reference module θ_1 serves as anchor (whether fixed or conditioned upon), leaving $M - 1$ independent

constraints. This is analogous to the classic result that the probability of M uniform phases falling in *some* semicircle is $M/2^{M-1}$, where the factor M accounts for the free choice of arc location; our anchored definition removes this degree of freedom.

First-passage time structure. Assuming independence across modules on the commit timescale, the alignment event A_ε is a rare target set on the M -torus T^M . Under fast mixing relative to the target set size, first-passage times are approximately exponential with mean $\tau \approx 1/(\text{attempt rate} \times P(A_\varepsilon))$ —a form of Kac’s recurrence theorem [20]. The attempt rate is set by the phase-exploration rate $\Delta\omega$. Thus:

$$\tau_{\text{coh}} = \frac{1}{\Delta\omega} p_1(\varepsilon, \kappa(r))^{-(M-1)} \quad (3)$$

For coupled modules (weak inter-module coupling), the effective exponent is reduced:

$$\tau_{\text{coh}} \approx \frac{1}{\Delta\omega} p_1(\varepsilon, \kappa(r))^{-\alpha(M-1)}, \quad \alpha \in (0, 1], \quad (4)$$

where α captures *effective independence*: $\alpha = 1$ for fully independent modules; $\alpha < 1$ when inter-module coupling induces correlations that reduce the effective number of constraints. One can define an effective constraint dimension $d_{\text{eff}} = \alpha(M-1)$ and estimate it empirically from the participation ratio of the covariance matrix of phase deviations $\delta\theta_i = \theta_i - \bar{\theta}$.

Key assumptions. Equation 4 relies on:

1. **Stationarity:** The phase distribution is approximately stationary over the commit timescale.
2. **Fast mixing:** The joint phase process mixes rapidly relative to the target set size, so first-passage times are approximately exponential (memoryless after conditioning on slowly varying parameters like r).
3. **Weak inter-module coupling:** Modules are semi-independent on the commit timescale; strong coupling would synchronize phases and reduce effective M .

4. **Attempt rate interpretation:** $\Delta\omega$ represents the inverse correlation time of phase deviations, not the carrier frequency.

Diagnostic: If these assumptions hold, inter-commit intervals (after conditioning on r) should be approximately exponentially distributed. Deviations indicate strong coupling, metastability, or non-stationarity.

Physical interpretation: M is the **coordination depth**—the number of semi-independent modules (functional clusters) whose order parameters must jointly exceed a threshold for a commit to register. *Modules are not individual oscillators:* within-module coherence is absorbed into the module’s aggregate phase estimate. $\Delta\omega$ is the **phase-exploration rate:** the characteristic frequency at which inter-module phases mix (not the carrier frequency). The tolerance ε is the **full phase window width** (in radians); $\varepsilon = \pi$ corresponds to $\pm 90^\circ$ alignment tolerance.

2.2 The speed-flexibility trade-off

The coherence time reveals opposing constraints on commit rate:

Proposition 1 (Speed-Flexibility Frontier). *At fixed power P , systems face a Pareto frontier in (M, r) space:*

- **Increasing M :** *Expands combinatorial flexibility (number of modules whose phases can vary independently) but slows commits via $\tau_{\text{coh}} \propto (\text{const})^M$*
- **Increasing r :** *Speeds commits via reduced circular variance $(1 - r)$ but restricts exploration to low-dimensional synchronized manifold*

Increasing coordination depth M expands the dimensionality of commit-registered outcomes but requires exponentially longer alignment time. Increasing coherence r speeds commits but forces trajectories onto low-dimensional attractors, reducing the set of accessible commit outcomes.

2.3 Quantum and thermodynamic limits of computation

A natural question is how this coordination-based framework relates to fundamental physical limits on computation. We now explicitly address the quantum and thermodynamic bounds that form the basis of information transmission in living systems.

2.3.1 The Landauer bound

Landauer’s principle [11] establishes that any logically irreversible operation—specifically, the erasure or overwriting of one bit of information—dissipates at least $k_{\text{B}}T \ln 2$ of energy as heat. This bound arises from the connection between logical irreversibility and thermodynamic entropy: erasing a bit reduces the system’s information-theoretic entropy, which must be compensated by entropy increase in the environment [12].

The minimum energy per bit at biological temperature ($T \approx 310$ K) is:

$$E_{\text{Landauer}} = k_{\text{B}}T \ln 2 \approx 3 \times 10^{-21} \text{ J/bit}$$

This sets an absolute floor on energy dissipation per irreversible operation. However, the Landauer bound constrains energy per operation, not time per operation. A system could in principle perform operations arbitrarily fast if sufficient power were available—until quantum speed limits intervene.

2.3.2 The Margolus-Levitin bound and quantum speed limits

The Margolus-Levitin theorem [14] establishes that the minimum time to transition between orthogonal quantum states is:

$$\tau_{\text{ML}} = \frac{\pi \hbar}{2 \langle E \rangle}$$

where $\langle E \rangle$ is the average energy above the ground state. This bound, together with the Mandelstam-Tamm relation [13] ($\tau_{\text{MT}} = \hbar/(2\Delta E)$ where ΔE is energy uncertainty), constitutes the quantum speed limit on state evolution.

These bounds originate from the energy-time uncertainty relation $\Delta E \cdot \Delta t \gtrsim \hbar/2$, which is fundamental to understanding information transmission in physical systems [15, 16, 17]. As Bormashenko [12] notes, synthesis of the Landauer bound with the Margolus-Levitin limit yields a minimal computation time scaling as $\tau_{\min} \sim \hbar/k_B T$ —decreasing temperature reduces energy consumption per operation but simultaneously slows computation.

At biological temperatures with $\langle E \rangle \sim k_B T \approx 4 \times 10^{-21}$ J:

$$\tau_{\text{QSL}} \sim \frac{\pi \times 10^{-34}}{2 \times 4 \times 10^{-21}} \sim 10^{-13} \text{ s}$$

This is fourteen orders of magnitude faster than the millisecond timescales of neural processing.

2.3.3 Why coherence time dominates in biological systems

The central claim of this paper is that in biological neural systems, the quantum speed limit and Landauer bound are typically **non-binding constraints**. The dominant bottleneck is coherence time—the time required for distributed oscillatory assemblies to achieve sufficient phase coordination for a commit to occur.

Consider the hierarchy of timescales for cortical visual processing:

$$\tau_{\text{QSL}} \sim 10^{-13} \text{ s} \quad (\text{quantum speed limit}) \tag{5}$$

$$\tau_{\text{Landauer}} \sim 10^{-12} \text{ s} \quad (\text{thermal relaxation}) \tag{6}$$

$$\tau_{\text{SNR}} \sim 10^{-3} \text{ s} \quad (\text{signal detection}) \tag{7}$$

$$\tau_{\text{coh}} \sim 10^{-2} \text{ s} \quad (\text{coordination time}) \tag{8}$$

The coherence time exceeds quantum limits by eleven orders of magnitude. This enormous gap arises because neural computation is not limited by how fast individual elements can switch states, but by how long it takes for *distributed assemblies of semi-independent*

modules to achieve coordinated phase alignment.

Mathematically, this reflects the exponential scaling of coherence time with coordination depth:

$$\tau_{\text{coh}} \propto p_1^{-\alpha(M-1)}$$

For $M = 8$ modules with $p_1 \approx 0.7$ and $\alpha = 0.7$, the exponential factor is $(0.7)^{-4.9} \approx 10$. Even modest coordination requirements ($M \sim 5\text{--}15$) produce millisecond-scale coherence times from microsecond-scale phase dynamics.

This analysis clarifies the relationship between our framework and fundamental physics: quantum and thermodynamic limits set absolute bounds on computation, but biological systems operate in a regime where multi-module coordination—not quantum switching speed or Landauer dissipation—determines processing rate.

2.4 Parameter estimation protocols

Coordination depth (M). M is the count of semi-independent modules whose order parameters must jointly exceed a threshold to register a commit. **Operational estimation:**

1. Cluster channels/units by band-limited coherence or Granger causality into modules.
2. Identify the minimal set of modules whose joint activity predicts behavioral commits.
3. Typical values: occipital visual tasks yield $M \sim 6\text{--}10$; cross-modal binding $M \sim 10\text{--}15$.

Kuramoto coherence (r). Extract instantaneous phase $\theta_j(t)$ from bandpass-filtered LFP/EEG via Hilbert transform. Compute the global order parameter:

$$r(t) = \left| \frac{1}{N} \sum_{j=1}^N e^{i\theta_j(t)} \right|$$

where N is the number of recording channels. For the coherence time model, we need module-level coherence: first cluster channels into M modules, compute each module’s aggregate

phase, then estimate r and κ from the inter-module phase distribution. Typical values: spontaneous cortex ~ 0.2 – 0.4 ; attention ~ 0.5 – 0.7 ; circadian networks ~ 0.7 – 0.9 [2].

Phase alignment window (ε). From spike train or LFP cross-correlograms, measure half-width at half-maximum (HWHM, units: seconds). The **full** phase window is $\varepsilon = 2\omega_0$ HWHM (radians).

Phase-exploration rate ($\Delta\omega$). Estimate from instantaneous frequency spread or phase diffusion. Typical values: $\Delta\omega \sim 2\pi \times (5\text{--}20)$ rad/s for cortical gamma/alpha bands.

3 Results

3.1 Visual perceptual binding windows

Human visual perception exhibits temporal integration windows of 30–50 ms (flicker fusion at 20–30 Hz). We apply Eq. 1 with neural parameters.

Take $M = 8$ occipital modules, $r = 0.7$ (attention [2]), full tolerance $\varepsilon = \pi$ rad, and phase-exploration rate $\Delta\omega = 2\pi \times 20$ rad/s.

First compute p_1 : at $r = 0.7$, $\kappa \approx 2.0$, giving $p_1(\pi, 2.0) \approx 0.62$ (numerical integration). Using Eq. 4 with $\alpha = 0.7$ (modular cortical topology):

$$\tau_{\text{coh}} \approx \frac{1}{125.66} (0.62)^{-0.7 \times 7} = \frac{1}{125.66} (0.62)^{-4.9} \approx 83 \text{ ms.}$$

With higher coherence $r = 0.8$ ($\kappa \approx 3.0$, $p_1(\pi, 3.0) \approx 0.76$):

$$\tau_{\text{coh}} \approx \frac{1}{125.66} (0.76)^{-4.9} \approx 30 \text{ ms.}$$

Computing all terms in Eq. 1:

$$\tau_{\text{QSL}} \sim 10^{-13} \text{ s} \quad (\text{negligible—quantum limit non-binding}) \quad (9)$$

$$\tau_{\text{SNR}} \sim 5\text{--}10 \text{ ms} \quad (\text{task-dependent}) \quad (10)$$

$$\tau_{\text{power}} \ll 1 \text{ ms} \quad (\text{per commit}) \quad (11)$$

$$\tau_{\text{coh}} \approx 30\text{--}83 \text{ ms} \quad (r = 0.7\text{--}0.8, M = 8) \quad (12)$$

The effective commit time is $\tau_{\text{eff}} \gtrsim \max[\text{QSL}, \text{SNR}, \text{power}, \text{coh}] \approx 30\text{--}83 \text{ ms}$, matching human binding windows (30–50 ms). **Coherence time dominates**; quantum and Landauer bounds are not the limiting factors.

Sensitivity analysis. Table 1 shows how τ_{coh} varies across plausible parameter ranges. The 30–100 ms binding window is robust to parameter uncertainty.

Table 1: Sensitivity of coherence time to parameter variations. Base case: $M = 8$, $r = 0.75$, $\varepsilon = \pi$, $\Delta\omega = 2\pi \times 20 \text{ rad/s}$, $\alpha = 0.7$.

Parameter varied	M	r	ε	$\Delta\omega$ (rad/s)	τ_{coh} (ms)
Base case	8	0.75	π	$2\pi \times 20$	52
Lower M	6	0.75	π	$2\pi \times 20$	28
Higher M	10	0.75	π	$2\pi \times 20$	95
Lower r	8	0.65	π	$2\pi \times 20$	98
Higher r	8	0.85	π	$2\pi \times 20$	25
Narrower ε	8	0.75	$2\pi/3$	$2\pi \times 20$	120
Faster $\Delta\omega$	8	0.75	π	$2\pi \times 30$	35

The exponential dependence on M and strong dependence on r mean that modest changes in attention state (affecting r) or task complexity (affecting M) can shift binding windows by factors of 2–4, consistent with empirical observations of attention-dependent temporal

resolution [23].

3.2 Tachypsychia: a dual-loop hypothesis

During acute stress or falls, subjects report subjective time slowing while objective reaction times remain unchanged [22]. We propose a **dual-commit architecture** as one explanation for this dissociation (alternative accounts exist, including attentional sampling rate changes and memory density effects):

Perceptual loop (cortical): High- M (~ 10 – 15 modules) coherent field dynamics. Commits sparse (5–20 Hz). Arousal increases coherence r and thus information rate per commit interval.

Motor loop (cerebellar/basal ganglia): Low- M (~ 3 – 5 modules) primitives. Commits faster (50–150 ms). Arousal modulates decision threshold, preserving reaction time.

Measurable proxies: Perceptual loop commit rate could be indexed by temporal-order judgment thresholds or EEG phase coherence in theta/alpha bands. Motor loop could be indexed by simple RT distributions or EMG onset latencies. The hypothesis predicts these measures should be *statistically dissociable* under arousal manipulation, with temporal-order thresholds changing while simple RT remains stable.

3.3 Alpha oscillations and temporal acuity

Alpha oscillations (8–12 Hz) in visual cortex correlate with temporal acuity: individuals with faster alpha perceive time faster [23, 24]. One interpretation is that alpha acts as a gating rhythm that modulates commit opportunity windows, effectively setting the attempt rate $\Delta\omega$ or the sampling schedule for phase alignment.

Prediction: If alpha frequency modulates commit timing, then entraining alpha via transcranial alternating current stimulation (tACS) should shift perceptual binding windows approximately linearly with frequency change. This prediction distinguishes the gating interpretation from alternatives where alpha reflects post-commit memory encoding.

3.4 Metabolic scaling across species

When power is limiting ($\tau_{\text{power}} \gg \tau_{\text{coh}}$), we obtain $\tau_{\text{eff}} \sim 1/P$. Critical flicker fusion frequency should scale with mass-specific metabolic rate.

Healy et al. [25] measured critical flicker fusion frequency across diverse taxa: flies ~ 240 Hz, humans ~ 60 Hz, leatherback turtles ~ 15 Hz. Their analysis found that mass-specific metabolic rate explains substantial variance in CFF across species spanning three orders of magnitude in body mass, consistent with the predicted power-limited regime for small, high-metabolism animals.

4 Discussion

4.1 Biological examples of the speed-flexibility trade-off

Mind-wandering and creative insight (high M , low r). Default-mode network activity exhibits low inter-regional coherence ($r \sim 0.3$) and high coordination depth ($M \sim 12\text{--}20$). Commits are rare ($\sim 0.1\text{--}1$ Hz), producing subjective “slow thinking” but enabling novel associations.

Focused attention (moderate M , moderate r). Attention increases coherence ($r : 0.3 \rightarrow 0.6$) and narrows coordination to task-relevant modules ($M : 15 \rightarrow 8$) [2]. This produces faster commits (10–30 Hz) in a restricted subspace.

Automaticity and skill learning (low M , high r). Overlearned motor sequences compress to low-dimensional primitives ($M \sim 3\text{--}5$) with tight coherence ($r \sim 0.8$). Commits are rapid (50–150 ms) but inflexible.

4.2 Temporal embedding as a pre-commit representation mechanism (hypothesis)

Coherence time tells us *how long* pre-commit dynamics must be maintained before alignment occurs. A separate question is *how* high-dimensional structure can remain accessible during that interval. We propose—as a hypothesis for future investigation—that biological systems may implement **Takens delay embedding** [26].

Takens’ theorem. For a deterministic system with state dimension d , time-delayed copies of a scalar observable— $x(t), x(t - \tau), x(t - 2\tau), \dots$ —can reconstruct the complete state-space topology. Crucially, $2d + 1$ delay coordinates generically suffice.

A candidate biological implementation: cerebellar parallel fibers. Parallel fibers—unmyelinated granule cell axons—conduct at 0.2–0.5 m/s with systematically varying path lengths, creating conduction delays of 5–15 ms [27, 28, 29]. These delays match the optimal embedding parameter $\tau^* = 1/(4f)$ for motor dynamics at 15–50 Hz.

We hypothesize that the cerebellum may provide one biological implementation of delay-embedded state preservation during pre-commit intervals. This would enable rapid low-dimensional readout (motor commands) from high-dimensional state without requiring simultaneous global coordination. **This remains conjecture**; direct evidence would require demonstrating that cerebellar computations exploit delay structure for state reconstruction.

Robustness to noise. We use Takens embedding as a *design principle* rather than a literal guarantee: delay-line architectures improve state inference under partial observability, even when exact reconstruction is impossible.

Connection to coherence time. Coherence time determines the maximum window over which temporal embeddings remain usable. If coherence decays faster than the embedding

window, reconstruction fails and high-dimensional structure is lost before a commit can occur.

4.3 Generalization beyond neural systems

While we emphasize neural applications given their empirical accessibility, the coherence time framework applies to any biological system whose function requires coordinated phase relations and whose outputs are sparse commitment events.

Scope statement. Neural systems are the best-instrumented instance of a broader class: living, self-maintaining oscillator assemblies that perform irreversible state registration. We do not treat collectives (e.g., societies) as intelligent systems in this sense; they are coordination networks of intelligent organisms and introduce additional channels not modeled here.

Chromosomal coordination. Chromosomal systems provide a non-neural example of the same architecture. High-dimensional continuous dynamics (polymer conformations, electrostatic interactions, topological constraints, phase separation) produce function via sparse low-dimensional commits (origin firing, checkpoint decisions, segregation outcomes).

Anaphase onset is naturally modeled as a commit event: a global irreversible transition triggered only after multiple semi-independent units satisfy a coherence-like constraint. The waiting time should scale steeply with effective coordination depth, consistent with first-passage into a small target set. The relevant “order parameter” is any coordination metric determining whether a global event can proceed—spatial colocalization, topological state, mechanical alignment—not necessarily Kuramoto r .

Circadian networks. Circadian gene networks exhibit $M \sim 3\text{--}5$ core feedback loops, $r \sim 0.8$ (tight coupling), and $\varepsilon \sim 0.5$ rad (loose phase tolerance). These parameters yield coordination times much shorter than the 24-hour period, implying coordination is *not* the

bottleneck. The 24-hour period arises from molecular lags. The framework correctly identifies the dominant regime: circadian clocks are biochemical-delay-limited, not coordination-limited.

4.4 Testable predictions

Alpha entrainment. tACS at alpha frequency should shift perceptual binding windows linearly with frequency change. 20% frequency increase predicts 20% threshold reduction.

Arousal effects on dual-loop dissociation. During simultaneous simple reaction time + temporal-order judgment under acute stress:

- Temporal-order discrimination threshold changes
- Simple reaction time unchanged or slightly faster
- Statistical independence between measures

Coherence measures and consciousness. If phenomenal experience reflects pre-commit coherent dynamics, conscious states should correlate more strongly with phase-locking index and theta-gamma coupling than with spike count or mean firing rate.

4.5 Relationship to existing frameworks

Quantum-like cognition. Recent work has explored connections between oscillatory neural networks and quantum-like cognitive phenomena [18]. Our framework provides a classical high-dimensional explanation for why such quantum-like signatures emerge: high-dimensional dynamics with limited measurement bandwidth produce phenomenology parallel to quantum systems—measurement-like collapse at commits, superposition-like coexistence of states pre-commit, uncertainty-like timing inaccessibility.

Integrated Information Theory [30]: We provide thermodynamic/temporal foundations. Integration costs energy and takes time (exponentially with coordination depth).

Free Energy Principle [31]: We add temporal structure. Prediction updates occur at commit events with rate limited by Eq. 1.

5 Conclusion

We have established that coherence time—the time required for distributed oscillatory assemblies to achieve sufficient phase alignment for irreversible state registration—bounds the rate of neural processes associated with cognition. The bound

$$\tau_{\text{eff}} = \max[\tau_{\text{QSL}}, \tau_{\text{SNR}}, \tau_{\text{coh}}, \tau_{\text{power}}]$$

with coherence time

$$\tau_{\text{coh}} = \frac{1}{\Delta\omega} p_1(\varepsilon, \kappa(r))^{-\alpha(M-1)}$$

unifies quantum, noise, and coordination limits to explain temporal resolution in biological processing systems.

We have situated this framework explicitly within the hierarchy of physical limits on computation: the Landauer bound constrains energy per irreversible operation; the Margolus-Levitin and Mandelstam-Tamm bounds constrain the speed of quantum state evolution; the energy-time uncertainty relation underlies both. In biological neural systems, these quantum and thermodynamic bounds are typically non-binding. The dominant constraint is the time required for multi-module coordination—a classical phenomenon arising from rare-event statistics on product manifolds.

Key findings:

1. **Speed-flexibility trade-off:** Increasing M exponentially slows commits but expands combinatorial flexibility. Increasing r speeds commits but restricts dynamics to low-

dimensional manifolds.

2. **Quantum limits non-binding:** Neural coherence times exceed quantum speed limits by eleven orders of magnitude. Coordination, not quantum switching, limits processing rate.
3. **Generality:** While neural systems provide the empirical testbed, the framework applies to any biological oscillator assembly with coordinated phase relations and sparse commits.
4. **Quantitative predictions:** Visual binding windows (30–50 ms), metabolic scaling of temporal acuity, alpha entrainment linearity, dual-task dissociations.

The central thesis: biological systems trade speed for flexibility by moving on the coherence-dimension frontier, operating in a regime where quantum and thermodynamic limits permit far faster processing than coordination constraints allow.

Data accessibility

This is a theoretical framework paper. Kuramoto simulation code and parameter estimation workflows are available at <https://github.com/todd866/coherence-time>.

Competing interests

The author declares no competing interests.

Funding

No external funding supported this work.

Acknowledgments

The author thanks Abir Igamberdiev and anonymous reviewers at BioSystems for constructive feedback.

References

- [1] Miller, E.K., Lundqvist, M., Bastos, A.M. (2018). Working memory 2.0. *Neuron*, 100(2), 463–475.
- [2] Fries, P. (2015). Rhythms for cognition: communication through coherence. *Neuron*, 88(1), 220–235.
- [3] Singer, W. (1999). Neuronal synchrony: a versatile code for the definition of relations? *Neuron*, 24(1), 49–65.
- [4] Lisman, J.E., Jensen, O. (2013). The theta-gamma neural code. *Neuron*, 77(6), 1002–1016.
- [5] Womelsdorf, T., et al. (2007). Modulation of neuronal interactions through neuronal synchronization. *Science*, 316(5831), 1609–1612.
- [6] Engel, A.K., Fries, P., Singer, W. (2001). Dynamic predictions: oscillations and synchrony in top-down processing. *Nat. Rev. Neurosci.*, 2(10), 704–716.
- [7] Varela, F., et al. (2001). The brainweb: phase synchronization and large-scale integration. *Nat. Rev. Neurosci.*, 2(4), 229–239.
- [8] Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends Cogn. Sci.*, 9(10), 474–480.
- [9] Todd, I. (2026). Intelligence as high-dimensional coherence: The observable dimensionality bound and computational tractability. *BioSystems*, 251, 105704.

- [10] Todd, I. (2025). Alignment probabilities on product statistical manifolds: Fisher information and coordination depth. *Information Geometry* (in preparation).
- [11] Landauer, R. (1961). Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.*, 5(3), 183–191.
- [12] Bormashenko, E. (2024). Landauer bound in the context of minimal physical principles: Meaning, experimental verification, controversies and perspectives. *Entropy*, 26(5), 423.
- [13] Mandelstam, L., Tamm, I. (1945). The uncertainty relation between energy and time in non-relativistic quantum mechanics. *J. Phys. (USSR)*, 9, 249–254.
- [14] Margolus, N., Levitin, L.B. (1998). The maximum speed of dynamical evolution. *Physica D*, 120(1–2), 188–195.
- [15] Igamberdiev, A.U. (2021). Quantum-like model for unconscious–conscious interaction and emotional coloring of perceptions and other conscious experiences. *BioSystems*, 208, 104471.
- [16] Igamberdiev, A.U. (2025). Physical limits of natural computation as the biological constraints of morphogenesis, evolution, and consciousness: On the 100th anniversary of Efim Liberman (1925–2011). *BioSystems*, 249, 105451.
- [17] Igamberdiev, A.U. (2026). Internal quantum constraints of natural computation in autopoietic systems. *BioSystems*, 251, 105707.
- [18] Khrennikov, A. (2025). Constructing a bridge between functioning of oscillatory neuronal networks and quantum-like cognition along with quantum-inspired computation and AI. *BioSystems*, 250, 105573.
- [19] Poor, H.V. (1994). *An Introduction to Signal Detection and Estimation* (2nd ed.). Springer-Verlag.

- [20] Kac, M. (1947). On the notion of recurrence in discrete stochastic processes. *Bull. Amer. Math. Soc.*, 53(10), 1002–1010.
- [21] Mardia, K.V., Jupp, P.E. (2000). *Directional Statistics*. John Wiley & Sons.
- [22] Stetson, C., Fiesta, M.P., Eagleman, D.M. (2007). Does time really slow down during a frightening event? *PLoS ONE*, 2(12), e1295.
- [23] Samaha, J., Postle, B.R. (2015). The speed of alpha-band oscillations predicts the temporal resolution of visual perception. *Current Biology*, 25(22), 2985–2990.
- [24] Cecere, R., Rees, G., Romei, V. (2015). Individual differences in alpha frequency drive crossmodal illusory perception. *Current Biology*, 25(2), 231–235.
- [25] Healy, K., McNally, L., Ruxton, G.D., Cooper, N., Jackson, A.L. (2013). Metabolic rate and body size are linked with perception of temporal information. *Anim. Behav.*, 86(4), 685–696.
- [26] Takens, F. (1981). Detecting strange attractors in turbulence. In: Rand, D., Young, L.S. (eds) *Dynamical Systems and Turbulence*, Lecture Notes in Mathematics, vol 898. Springer, pp. 366–381.
- [27] Braitenberg, V. (1958). Morphology of nerve net in cerebellar cortex. *Journal of Comparative Neurology*, 109, 1–33.
- [28] Llinás, R. (1982). General discussion: Radial connectivity in the cerebellar cortex. In: *The Cerebellum—New Vistas* (pp. 189–194). Springer.
- [29] Wyatt, K.D., et al. (2005). Speed limits in the cerebellum: constraints from myelinated and unmyelinated parallel fibers. *European Journal of Neuroscience*, 21(8), 2285–2290.
- [30] Tononi, G., Boly, M., Massimini, M., Koch, C. (2016). Integrated information theory: from consciousness to its physical substrate. *Nat. Rev. Neurosci.*, 17(7), 450–461.

- [31] Friston, K. (2010). The free-energy principle: a unified brain theory? *Nat. Rev. Neurosci.*, 11(2), 127–138.