On 'Ultrafilters with small generating sets'

Todor Dukov

October 13, 2026

Introduction

The aim of this document is to give a better understanding of [1] by filling in details while also giving a high-level overview of the construction. We will talk about the ultrafilter number $\mathfrak{u}$ and the dominating number $\mathfrak{d}$. The main result of [1] is the consistency of

$$\mathfrak{u} = \kappa < \mathfrak{d} = \mathfrak{c} = \lambda$$

for any regular κ, λ, such that $\kappa < \lambda$.

The way we would do that is by first adding (with finite support iteration) λ-many Cohen reals and then very carefully adding κ-many Mathias reals. We shall treat the construction of the ultrafilter for Mathias with utmost care, so that


1. It does not ruin any work of the Cohen forcing for the dominating number.
2. The same Mathias real would be added no matter what level we are on the Cohen forcing iteration.

It is obvious why we have the first requirement. After Cohen forcing we would have $\mathfrak{d} = \mathfrak{c} = \lambda$. We would like to keep it that way during every step of the Mathias forcing. It is not so obvious why we need to satisfy the second requirement. If we satisfy the second requirement we would get what is called a matrix iteration. This would allow for more precise control of the model because we can attack the problem from two sides instead of one which would also help with the first requirement.

Let me try to explain it better. Suppose the added Cohen reals are $\{r_\alpha\}_{\alpha<\lambda}$ and the added Mathias reals are $\{s_\beta\}_{\beta<\kappa}$. For $\alpha < \lambda$ and $\beta < \kappa$ denote

$$V_{\alpha,\beta} = \mathbb{L}[\{r_\gamma\}_{\gamma<\alpha}][\{s_\delta\}_{\delta<\beta}].$$

What we would like is for $V_{\alpha+1,\beta+1}$ to be related to both $V_{\alpha,\beta+1}$ and $V_{\alpha+1,\beta}$. Being related to both would guarantee the niceties of both our Mathias forcing and Cohen forcing. This is how it looks visually:



The method was introduced in [1]. It was further developed in [2], where the terminology *matrix iteration* appeared for the first time. Since first introduced, it has proven very useful in tackling combinatorial problems concerning cardinal characteristics of the reals.

So we talked about what we want for $\mathfrak{d}$, but what about $\mathfrak{u}$? Why would we have $\mathfrak{u} = \kappa$ in the end? Turns just having each ultrafilter for stage contain the currently added Mathias reals is enough, as we shall see next.

Why $\mathfrak{u} = \kappa$?

We will show this in two directions. First we will show an ultrafilter base of size κ. Then we will prove that no family of size smaller than κ forms an ultrafilter base.

Let $\mathcal{U}_\beta$ be the ultrafilter in step $\beta < \kappa$ of Mathias forcing. If for each $\beta < \kappa$, the ultrafilter $\mathcal{U}_\beta$ contains $\{s_\delta\}_{\delta<\beta}$, the family $\{s_\beta \setminus x \mid \beta < \kappa \ \& \ x \text{ is finite}\}$ will form a base for an ultrafilter. Let $A \in [\omega]^\omega \cap V_{\lambda,\kappa}$. Then since κ is regular, $A \in [\omega]^\omega \cap V_{\lambda,\beta}$ for some $\beta < \kappa$. This is true because in finite support iterations, limit stages of regular cofinality add no new reals. By properties of Mathias forcing, either A or $\omega \setminus A$ almost includes s_β. Therefore, $\mathfrak{u} \leq \kappa$.

We will show that $\kappa \leq \mathfrak{u}$. Let $\mathcal{A}$ be a family of subsets of ω in $V_{\lambda,\kappa}$ of size less than κ. By similar arguments, $\mathcal{A} \in V_{\lambda,\beta}$ for some $\beta < \kappa$. We will show a set $X \in V_{\lambda,\kappa}$ such that neither X nor $\omega \setminus X$ includes a member of $\mathcal{A}$. Define

$$X = \{n \in \omega \mid |n \cap s_\beta| \text{ is even}\}.$$

Assume that either X or $\omega \setminus X$ includes some $A \in \mathcal{A}$. Let (z, Z) be a condition forcing that. Let $m = \min Z$ and let a be an element of A that is larger than m. Then

- $(z, Z \setminus a)$ is an extension of (z, Z) forcing $a \cap s_\beta = z$.
- $(z \cup \{m\}, Z \setminus a)$ is an extension of (z, Z) forcing $a \cap s_\beta = z \cup \{m\}$.

One of the conditions force $a \in X$ and the other $a \notin X$, which is impossible, since

- If (z, Z) forces $A \subseteq X$, then $a \notin X$ can't be forced by any extension of (z, Z).
- If (z, Z) forces $A \subseteq \omega \setminus X$, then $a \in X$ can't be forced by any extension of (z, Z).

We have proved that neither X nor $\omega \setminus X$ includes a member of $\mathcal{A}$, so $\mathcal{A}$ can't form an ultrafilter base.

On normalized names

Before proceeding forward. We shall introduce (as it was done in [1]) a new type of name for functions $f : \omega \longrightarrow \omega$. Let $(\mathbb{P}, \leq)$ be a forcing poset in a model $\mathcal{M}$ of **ZFC**. A normalized $\mathbb{P}$-name for a function in $\mathcal{M}$ will be every $f = (A_n, f_n)_{n\in\omega}$, such that

- A_n is a maximal antichain.
- $f_n : A_n \longrightarrow \omega$.

Let G be a $\mathbb{P}$-generic filter in $\mathcal{M}$. Then the denotation of f with respect to G (written as f_G) is the function

$$\{(n, f_n(p)) \mid n \in \omega \ \& \ p \in G \cap W_n\}.$$

Since for each $n \in \omega$ the set $G \cap W_n$ is a singleton, f_G is a well-defined function.

We will prove the this notion of naming for functions is equivalent to the usual one.

Claim 1. For every normalized $\mathbb{P}$-name $f \in \mathcal{M}$ there exists a standard $\mathbb{P}$-name $\tau^f \in \mathcal{M}$, such that for each $\mathbb{P}$-generic filter G in $\mathcal{M}$:

$$f_G = (\tau_f)_G.$$

Proof. We can define the name explicitly as follows:

$$\tau_f = \{(\mathrm{op}(\check{n}, \check{f_n}(p)), p) \mid n \in \omega \ \& \ p \in W_n\}.$$

The rest is easy to check. □

Claim 2. For every standard $\mathbb{P}$-name $\tau \in \mathcal{M}$ for a function there exists a normalized $\mathbb{P}$-name $f \in \mathcal{M}$, such that for each $\mathbb{P}$-generic filter G in $\mathcal{M}$:

$$\tau_G = f_G.$$

Proof. Since τ is a name for a function, that means

$$\Vdash_{\mathbb{P}} \tau : \omega \longrightarrow \omega.$$

Then for each $n \in \omega$, the following set is dense

$$D_n = \{p \in \mathbb{P} \mid (\exists k \in \omega)(p \Vdash_{\mathbb{P}} \tau(k) = \check{n})\}.$$

For each $n \in \omega$, choose (by Zorn's lemma) a maximal antichain $A_n \subseteq D_n$. Define the function f_n as follows

$$f_n(p) = k_p \text{ for all } p \in A_n, \text{ where } k_p \in \omega \text{ is the unique witness to } p \in D_n.$$

Set $f = (A_n, f_n)_{n\in\omega}$. The rest is easy to check. □

Working below a certain condition, one can obtain an even stronger result than the previous one:

Claim 3. Let $\tau \in \mathcal{M}$ be a standard $\mathbb{P}$-name and $p \in \mathbb{P}$ be a condition such that

$$p \Vdash_{\mathbb{P}} \tau : \omega \longrightarrow \omega.$$

Then there exists a normalized name f such that for each $\mathbb{P}$-generic filter G in $\mathcal{M}$ containing p, we have

$$\tau_G = f_G.$$

Proof. Like previously, for each $n \in \omega$, the following sets are dense below p

$$D_n^p = \{q \in \mathbb{P} \mid q \leq p \ \& \ (\exists k \in \omega)(p \Vdash_{\mathbb{P}} \tau(k) = \check{n})\}.$$

Now choose for each $n \in \omega$, a maximal antichain A_n^p that is a subset of D_n^p. For each $q \in A_n^p$ choose a witness $k_q \in \omega$ to $q \in D_n^p$. Let A be a maximal antichain that is incompatible with p. Then for each $n \in \omega$, the set $A_n = A_n^p \cup A$ is a maximal antichain in $\mathbb{P}$, for an element that is incompatible with A_n must be compatible with one in A due to maximality and vice versa. Define the function f_n as follows

$$f_n(p) = \begin{cases} k_p, & \text{if } p \in A_n^p \\ 0, & \text{if } p \in A. \end{cases}$$

Set $f = (A_n, f_n)_{n \in \omega}$. The rest is easy to check. □

Constructing the ultrafilters (the central idea)

Going up with Cohen forcing adds unbounded reals. What we'd like is to keep it that way when going right with Mathias forcing. Also we would like to have our diagram commute (as shown in the picture before). That what this means formally is the following:

Claim 4. Let $\mathcal{M}_1$ and $\mathcal{M}_2$ be models of **ZFC** such that $\mathcal{M}_1$ is an inner model of $\mathcal{M}_2$. Let $g \in \omega_\omega \cap \mathcal{M}_2$ be unbounded over $\mathcal{M}_1$ and let $\mathcal{U}_1$ be an ultrafilter in $\mathcal{M}_1$. Then there exists and ultrafilter $\mathcal{U}_2$ in $\mathcal{M}_2$ such that:

(1) $\mathcal{U}_1 \subseteq \mathcal{U}_2$ and every maximal antichain in $\mathbb{M}(\mathcal{U}_1)$ is a maximal antichain in $\mathbb{M}(\mathcal{U}_2)$.

(2) If $f \in \mathcal{M}_1$ is a normalized $\mathbb{M}(\mathcal{U}_1)$-name for a real, then

$$\Vdash_{\mathbb{M}(\mathcal{U}_2)} \text{“}\tau_f \text{ does not dominate } \mathring{g}\text{“}.$$

We shall proceed as in [1], albeit more carefully. Before we dive into the proof, let's make some clarifying remarks:

Having (1) means that if G_2 is a $\mathbb{M}(\mathcal{U}_2)$-generic in $\mathcal{M}_2$, then the set $G_1 := G_2 \cap \mathbb{M}(\mathcal{U}_1)$ is $\mathbb{M}(\mathcal{U}_1)$-generic in $\mathcal{M}_1$, since it intersects every maximal antichain in $\mathcal{M}(\mathcal{U}_1)$. Also, the Mathias real $s(G_2)$ from adjoining G_2 to $\mathcal{M}_2$ is the same as the Mathias real $s(G_1)$ from adjoining G_1 to $\mathcal{M}_1$. This is true, because if $(a, A) \in G_2$ forces $a \subseteq s(G_2)$, then $(a, \omega \setminus \min A)$ is in G_2 (because it is closed upwards). Also, $\omega \setminus \min A$ is cofinite, so it trivially belongs to $\mathcal{U}_1$, thus $(a, \omega \setminus \min A) \in G_1$. This condition would force $a \subseteq s(G_1)$. Since $G_1 \subseteq G_2$, the converse direction is trivial.

Another fact that would follow from (1) is that any normalized $\mathcal{M}(\mathcal{U}_1)$-name f for a real is also a normalized $\mathcal{M}(\mathcal{U}_2)$-name. In fact, f is a name for the same real, in the sense that $f_{G_1} = f_{G_2}$. So the third statement is well-defined.

Last thing that we shall comment on before the actual proof, is that statement (2) is equivalent to statement (2′) which is obtained by replacing “dominate“ with “totally dominate“ in statement (2). Clearly, (2′) follows from (2). Now suppose f is a counterexample for (2). Then, there exists a forcing condition p and a $k \in \omega$, such that

$$p \Vdash_{\mathbb{M}(\mathcal{U}_2)} (\forall n \geq \check{k})(\tau_f(n) \geq \mathring{g}(n)).$$

We can easily change f to a normalized name f' (by bumping up the possible values of f for $0, \ldots, k$), so that

$$p \Vdash_{\mathbb{M}(\mathcal{U}_2)} (\forall n \in \omega)(\tau_{f'}(n) \geq \mathring{g}(n)).$$

Thus, a counterexample for (2) will produce a counterexample for (2′).

Constructing the ultrafilters (formally)

T.B.D.

Why $\mathfrak{d} = \mathfrak{c} = \lambda$?

T.B.D.

References

[1] Blass, A., and Shelah, S. Ultrafilters with small generating sets. *Israel Journal of Mathematics 65*, 3 (1989), 259–271.

[2] Brendle, J., and Fischer, V. Mad families, splitting families and large continuum. *The Journal of Symbolic Logic 76*, 1 (2011), 198–208.