


## Two- and Three-dimensional problems: Heat equation.

Derivation of the model problem



$u$ : temperature

$\rho$ : density

$c$ : specific heat capacity

$f$ : heat source / radiation

$\mathcal{L}$ : internal energy

$D$  is arbitrarily chosen  
and  $D \subset \Omega \subset \mathbb{R}^{n_{sd}}$ .

$$\int_D \mathcal{L}|_{t=t_2} - \mathcal{L}|_{t=t_1} dx = \int_{t_1}^{t_2} \int_{\partial D} -\vec{q} \cdot \vec{n} d\mathbf{x} dt + \int_{t_1}^{t_2} \int_D f d\mathbf{x} dt$$

Divergence theorem:  $f \in C^1$

$$\int_{\partial \Omega} f n_i da = \int_{\Omega} f_{,i} dx.$$

$$f_{,i} = \frac{\partial f}{\partial x_i}$$

$$n_{sd} = 2 \\ \{x_i\} = \begin{Bmatrix} x \\ y \end{Bmatrix}$$

$$n_{sd} = 3 \\ \{x_i\} = \begin{Bmatrix} x \\ y \\ z \end{Bmatrix}$$

$$\int_{t_1}^{t_2} \int_D \frac{\partial \mathcal{L}}{\partial t} dx dt = \int_{t_1}^{t_2} \int_D -\nabla \cdot \vec{q} + f dx dt$$

$$\Rightarrow \frac{\partial \mathcal{L}}{\partial t} = -\nabla \cdot \vec{q} + f$$

Remark: We write  $\nabla \cdot \vec{q}$  as  $q_{i,i}$  in index notation.

Constitutive relation:

$$C = \rho c u$$


$$\vec{q} = -\bar{\kappa} \nabla u \quad \text{or} \quad q_i = -\kappa_{ij} u_{,j} \quad (\text{Fourier's law})$$

$\kappa$  is the thermal conductivity, and it is symmetric, positive definite.

Remark: If  $\kappa_{ij}$ 's are constants, we say the material is homogeneous.

If  $\kappa_{ij}(x) = \kappa(x) \delta_{ij}$ , we say the material is isotropic.

Strong-form problem:



$$\left\{ \begin{array}{l} \rho c u_{,t} + q_{i,i} = f \quad \text{in } \Omega \times (0, T) \\ u = g \quad \text{on } \Gamma_g \times (0, T) \\ -q_i n_i = h \quad \text{on } \Gamma_h \times (0, T) \\ u(x, 0) = u_0(x) \quad \text{in } \Omega. \end{array} \right.$$

$g$ : prescribed boundary temperature

$h$ : .. .. heat flux

Integration by parts :

Let  $f, g : \bar{\Omega} \rightarrow \mathbb{R}$  be  $C^1$ , then

$$\int_{\Omega} f_{,i} g \, d\Omega = - \int_{\Omega} f g_{,i} \, d\Omega + \int_{\Gamma} f g \, n_i \, d\Gamma.$$

Proof:  $\int_{\Omega} (fg)_{,i} \, d\Omega = \int_{\Gamma} fg \, n_i \, d\Gamma$

||

$$\int_{\Omega} f_{,i} g + f g_{,i} \, d\Omega.$$

Trial solution space  $\mathcal{S} := \{ u : u \in H^1(\Omega), u = g \text{ on } \Gamma_g \}$

Test function space  $\mathcal{V} := \{ w : w \in H^1(\Omega), w = 0 \text{ on } \Gamma_g \}$

$$(w) \left\{ \begin{array}{l} \text{Given } f, g, h, \text{ find } u \in \mathcal{S} \text{ such that} \\ - \int_{\Omega} w_{,i} g_{,i} \, d\Omega = \int_{\Omega} w f \, d\Omega + \int_{\Gamma_h} w h \, d\Gamma \\ \text{for } \forall w \in \mathcal{V}. \end{array} \right.$$

Theorem: Assuming all functions involved are smooth enough to justify manipulations, then a solution of (S) is a solution of (w) and vice versa.

Proof: (S)  $\Rightarrow$  (w)

$$\begin{aligned}
0 &= \int_{\Omega_1} (g_{i,i} - f) w \, d\Omega_1 \\
&= - \int_{\Omega_1} g_{i,i} w_{,i} \, d\Omega_1 + \int_{\Gamma} w_{,i} g_{i,i} n_i \, d\Gamma - \int_{\Omega_1} w f \, d\Omega_1 \\
&= - \int_{\Omega_1} g_{i,i} w_{,i} - \int_{\Gamma} w h_i \, d\Gamma - \int_{\Omega_1} w f \, d\Omega_1.
\end{aligned}$$

(W)  $\Rightarrow$  (S)

$$\begin{aligned}
0 &= \int_{\Omega_1} w_{,i} g_{i,i} + w f \, d\Omega_1 + \int_{\Gamma_h} w h \, d\Gamma \\
&= \int_{\Omega_1} w \underbrace{(-g_{i,i} + f)}_{\alpha} \, d\Omega_1 + \int_{\Gamma_h} w \underbrace{(n_i g_{i,i} + h)}_{\beta} \, d\Gamma
\end{aligned}$$

Step 1: pick  $w = \alpha \phi$  with  $\phi$  satisfying  
 $\phi > 0$  in  $\Omega_1$ ;  $\phi = 0$  on  $\Gamma$ ;  $\phi$  is smooth.

then  $0 = \int_{\Omega_1} \phi \alpha^2 \, d\Omega_1,$

which implies  $\alpha = 0$ .

Step 2: pick  $w = \beta \psi,$

$\psi > 0$  on  $\Gamma_h$ ;  $\psi = 0$  on  $\Gamma_g$ ;  $\psi$  is smooth.

then  $0 = \int_{\Omega_1} \beta^2 \psi^2 \, d\Gamma$

$\Rightarrow \beta = 0.$





Notation:  $a(w, u) = \int_{\Omega} w_{,i} x_{ij} u_{,j} d\Omega$

$$(w, f) = \int_{\Omega} w f d\Omega$$

$$(w, h)_{\Gamma} = \int_{\Gamma} w h d\Gamma$$

$$\Rightarrow a(w, u) = (w, f) + (w, h)_{\Gamma}.$$

Index-free notation: Consider the case of  $n_{sd} = 2$ .

$\nabla$ : gradient operator

$$\nabla u = \{u_{,i}\} = \begin{Bmatrix} u_{,1} \\ u_{,2} \end{Bmatrix} \quad \text{isotropic case}$$

$$\kappa = [\kappa_{ij}] = \begin{bmatrix} \kappa_{11} & \kappa_{12} \\ \kappa_{21} & \kappa_{22} \end{bmatrix} = \underset{\text{scalar}}{\kappa} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

then  $w_{,i} x_{ij} u_{,j} = (\nabla w)^T \kappa \nabla u$

and 
$$a(w, u) = \int_{\Omega} (\nabla w)^T \kappa \nabla u d\Omega$$

Galerkin approximation:  $\mathcal{Z}^h \subset \mathcal{Z}$ ,  $\mathcal{V}^h \subset \mathcal{V}$ .

We assume  $u^h \in \mathcal{Z}^h$  adopts the decomposition,

$$u^h = v^h + \underset{\cap \mathcal{V}^h}{g^h}$$


and  $g^h$  satisfy (or approximately satisfy)

$$u = g \text{ on } \Gamma_g$$

$$(G) \left\{ \begin{array}{l} \text{Given } f, g, h, \text{ find } u^h = v^h + g^h \text{ such that} \\ \text{for } \forall w^h \in \mathcal{V}^h, \\ a(w^h, v^h) = (w^h, f) + (w^h, g^h)_{\Gamma} \\ \quad - a(w^h, g^h) \end{array} \right.$$

□:  $g$ -node

• node



$\bigcup_{e=1}^{n_{el}} \Omega_4^e$  approximates  $\Omega$ ;  $\eta = \{1, 2, \dots, n_{np}\}$

↪ set of global node numbers

$g$ -node: node at which  $u^h = g$  is prescribed.

$\eta_g$  : set of g-node.

complement of  $\eta_g$  in  $\eta$  :  $\eta - \eta_g$  is the set of nodes at which  $w^h$  is to be determined.

$$\boxed{\dim(\eta - \eta_g) = n_{eg}}$$

$$w^h(x) = \sum_{A \in \eta - \eta_g} N_A(x) C_A$$

↑  
Shape function associated with node A.

$$v^h(x) = \sum_{A \in \eta - \eta_g} N_A(x) d_A$$

↑  
unknown temperature at node A.

$$g^h(x) = \sum_{A \in \eta_g} N_A(x) g_A. \quad g_A = g(x_A)$$

then we have

$$\sum_{B \in \eta - \eta_g} a(N_A, N_B) d_B = (N_A, f) + (N_A, h)_{\Gamma}$$

$$- \sum_{B \in \eta_g} a(N_A, N_B) g_B$$

for  $A \in \eta - \eta_g$ .

Now, we invoke ID array:

$$ID(A) = \begin{cases} P & \text{if } A \in \eta - \eta_g \\ 0 & \text{if } A \in \eta_g \end{cases}$$


where  $1 \leq P \leq n_{eq}$ .

$$\Rightarrow K_{PQ} = a(N_A, N_B) \quad P = ID(A), \quad Q = ID(B)$$

$$F_P = (N_A, f) + (N_A, \bar{u})_P - a(N_A, g^h)$$

$$d_Q = d_B$$

↑      ↑  
Q = ID(B)



and

$$u_{(x)}^h = \sum_{B \in \eta - \eta_g} N_B^{(x)} d_B + g^h(x)$$

Remark: In addition to the function space approximation (i.e.,  $\mathcal{V}^h \subset \mathcal{V}$ ,  $\mathcal{Z}^h \subset \mathcal{Z}$ ),

there are geometrical approximation  $(\bigcup_{e=1}^{n_{el}} \Omega_e^e \approx \Omega)$



and BC data approximation (i.e.,  $g^h(x) \approx g(x)$ ).

Local / element perspective

$$K = \sum_{e=1}^{nel} K^e = \sum_{e=1}^{nel} \int_{\Omega_e} \nabla N_A^T \chi \nabla N_B d\Omega$$

$$F = \sum_{e=1}^{nel} F^e = \sum_{e=1}^{nel} \left\{ \int_{\Omega_e} N_A f d\Omega + \int_{\Gamma_h^e} N_A \chi d\Gamma - a(N_A, N_B)^e g_B \right\}$$

$a(N_A, N_B)^e$


$\Gamma_h^e$

element stiffness & element force vector

$$K^e = [k_{ab}^e] \quad k_{ab}^e = \int_{\Omega_e} \nabla N_a^T \chi \nabla N_b d\Omega$$

$$f^e = [f_a^e] \quad f_a^e = \int_{\Omega_e} N_a f d\Omega + \int_{\Gamma_h^e} N_a \chi d\Gamma - \sum_{b=1}^{nel} k_{ab}^e g_b^e$$

$$1 \leq a, b \leq n_{en}$$



$$A = IEN(a, e)$$

local node number  
↓  
global node number      element number

Again,  $LM(a, e) = ID(IEN(a, e))$ .

Assembly:  $K = \sum_{e=1}^{nel} A k^e \quad F = \sum_{e=1}^{nel} A f^e$