

Single-Crystalline PtTe₂ for Efficient Ethylene Glycol Oxidation Reaction with Bias-free Solar-to-Hydrogen

Yung-Hung Huang^{1,2}, Chun-Wei Chen^{1,3,4}

¹International Graduate Program of Molecular Science and Technology (NTU-MST), National Taiwan University, Taipei, Taiwan

²Molecular Science and Technology Program Taiwan International Graduate Program (TIGP), Academia Sinica, Taipei, Taiwan

³Department of Materials Science and Engineering, National Taiwan University, Taipei, Taiwan

⁴Center for Condensed Matter Sciences and Center of Atomic Initiative for New Materials (AI-MAT), National Taiwan University (NTU), Taipei, Taiwan

e-mail: hyh00.05.16@gmail.com

ABSTRACT

My background in materials science and electrochemistry motivated me to pursue sustainable energy technologies that address both global energy demand and environmental challenges. During my studies, I recognized the limitations of conventional water electrolysis, particularly its dependence on freshwater and the sluggish oxygen evolution reaction (OER). Meanwhile, the growing impact of plastic waste, especially PET, highlighted the potential of integrating clean energy generation with carbon-based waste conversion. These interests aligned with the research focus of Prof. Chun-Wei Chen's laboratory, which specializes in photoelectrochemical (PEC) systems, two-dimensional catalysts, and sustainable electrochemical processes. My previous collaboration with Prof. Chen during my master's studies, which resulted in a co-first-author publication in *Journal of Materials Chemistry A*,¹ further strengthened my motivation to continue in this field.

Ethylene glycol (EG), obtainable from PET hydrolysis, serves as an ideal molecule for value-added oxidation.² Replacing OER with the ethylene glycol oxidation reaction (EGOR) enables simultaneous hydrogen production and chemical upcycling.³ My current work focuses on developing catalysts for EGOR in 1 M NaOH + 1 M EG, where the reaction proceeds at much lower potential and produces high-value glycolic acid (GA), confirmed by ¹H NMR spectroscopy. Single-crystalline PtTe₂ demonstrates excellent EGOR performance with a low onset potential of 0.41 V vs. RHE, along with strong HER activity in acidic media. To improve overall system efficiency, I integrated a PtTe₂ anode with a PtTe₂/graphene/silicon photocathode to construct a bias-free PEC-EC system. According to Nernst equation, the alkaline-acidic pH gradient provides a built-in 0.83 V potential, enabling photocurrent densities up to 37.8 mA/cm² without external bias. Moving forward, I aim to deepen mechanistic understanding and advance PEC-driven plastic upcycling toward scalable solar-to-chemical conversion.

REFERENCES

1. Wu, P.-H. et al. Efficient solar-assisted seawater splitting in alkaline solution using perovskite-graphene-Si tandem integration. *J. Mater. Chem. A* **13**, 28965-28973 (2025).
2. Shi, K. et al. Pd/NiMoO₄/NF electrocatalysts for the efficient and ultra-stable synthesis and electrolyte-assisted extraction of glycolate. *Nat. Commun.* **15**, 2899 (2024).
3. Liu, F. et al. Concerted and selective electrooxidation of polyethylene-terephthalate-derived alcohol to glycolic acid at an industry-level current density over a Pd-Ni(OH)₂ catalyst. *Angew. Chem. Int. Ed.* **62**, e202300094 (2023).