

BROWNIAN MOTION

Time Allowed: 2 Hours

Full marks may be gained by correctly answering 3 complete questions. Candidates may attempt all questions. Marks will be awarded for the best 3 answers only.

Read carefully the instructions on the answer book and make sure that the particulars required are entered on each answer book

1. (a) Give a definition of one-dimensional, standard Brownian motion. [4 marks]
- (b) Construct a process which has the same finite dimensional distributions as one-dimensional Brownian motion but is *almost surely* not continuous. You need to include full justifications of the properties of the process that you construct. [8 marks]
- (c) For a one-dimensional, standard Brownian motion $\beta(\cdot)$, starting from zero, denote by P_0 its distribution. Consider the process $x(t) := \int_0^t \beta(s) ds$ and calculate $E_0[x(t)]$, $E_0[(x(t))^2]$ and $E_0[(x(t))^3]$. [8 marks]

2. Let $\beta(\cdot)$ be one-dimensional Brownian motion starting at 0 and denote by P_0 its distribution. For $n, k_n \geq 1$ let

$$0 = t_0^n \leq t_1^n \leq \dots \leq t_{k_n}^n = t,$$

define partitions of the interval $(0, t)$, such that $\max_{i=1, \dots, k_n} |t_j^n - t_{j-1}^n| \leq e^{-n}$. Define

$$D_n := \sum_{j=1}^{k_n} (\beta(t_j^n) - \beta(t_{j-1}^n))^2.$$

- (a) Compute the mean of D_n . [3 marks]
- (b) Compute the variance of D_n . [6 marks]
- (c) Show that D_n converges *almost surely* as $n \rightarrow \infty$ and compute its limit. [6 marks]
- (d) Show that

$$\sup_{n \geq 1} \sum_{i=1}^{k_n} |\beta(t_i^n) - \beta(t_{i-1}^n)| = +\infty, \quad P_0 - a.s.$$

[5 marks]

Continued ...

3. (a) Consider a cylinder $\mathcal{C} := B(0, 1) \times \mathbb{R}_+$, where $B(0, 1)$ is the disk on \mathbb{R}^2 , with unit radius and centred at zero. Assume that there is a “smooth” solution to the boundary value problem

$$\begin{aligned} u_t &= \frac{1}{2} \Delta u, & \text{in } \mathcal{C}, \\ u(0, x) &= f(x), & \text{for } x \in B(0, 1), \\ u(t, x) &= g(t, x), & \text{for } t > 0, \quad x \in \partial B(0, 1). \end{aligned}$$

Write a representation of the solution as a functional of Brownian motion. Justify all your steps and state the “smoothness” assumptions that you need.

[6 marks]

- (b) Let $P_x, x \in \mathbb{R}$, denote the Wiener measure corresponding to standard, one-dimensional Brownian motion starting from x . Denote $u(t, x) := P_x(\max_{0 \leq s \leq t} \beta(s) < b)$. Set up the boundary value problem that $u(t, x)$ satisfies. Justify your derivation. [6 marks]
- (c) Let $\beta_1(\cdot)$ and $\beta_2(\cdot)$ be two independent, standard, one-dimensional Brownian motions starting from x_1, x_2 , respectively. Compute the probability that they do not meet in the time interval $[0, t], t > 0$. [8 marks]

4. (a) Consider the parabolic problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{1}{2} \Delta u + V(x)u, & t > 0, x \in \mathbb{R}^d, \\ u(0, x) &= f(x), & x \in \mathbb{R}^d. \end{aligned}$$

Show that

$$u(t, x) := E_x \left[f(\beta(t)) \exp \left(\int_0^t V(\beta(s)) ds \right) \right]$$

solves the above boundary value problem. State any assumptions you might need on V, f . [10 marks]

- (b) Let $\beta(\cdot)$ be a one-dimensional Brownian motion. Let $\gamma, \alpha > 0$ and $\tau := \inf\{t: |\beta(t)| \geq \alpha\}$. Compute $E_0[e^{\gamma\tau}]$. Does this exist for all $\gamma > 0$? Justify your answer. [10 marks]

THE END

BROWNIAN MOTION

Time Allowed: 2 Hours

Full marks may be gained by correctly answering 3 complete questions. Candidates may attempt all questions. Marks will be awarded for the best 3 answers only.

Read carefully the instructions on the answer book and make sure that the particulars required are entered on each answer book

1. (a) Give the definition of one dimensional, standard Brownian motion.

[4 marks]

Solution [Bookwork]. Brownian motion is a one dimensional, continuous process with independent increments, which are distributed as normals with mean zero and variance equal to the time increment.

- (b) Construct a process which has the same finite dimensional distributions like one dimensional Brownian motion but it is *almost surely* not continuous. You need to include full justifications of the properties of the process that you will construct.

Solution [Similar example seen]: Let U be a uniform random variable on $[0, 1]$ and $\beta(\cdot)$ a standard one dimensional Brownian motion. Define the process $(x(t) : t \in [0, 1])$ which equals $\beta(t)$ for any $t \in [0, 1]$ different than U and it is equal to zero for $t = U$. For any collection of times $0 \leq t_1 < t_2 < \dots < t_k \leq 1$, we have that *a.s.* $U \notin \{t_1, \dots, t_k\}$ and thus $(x(t_1), \dots, x(t_k)) = (\beta(t_1), \dots, \beta(t_k))$, which implies that the finite dimensional distributions of $x(\cdot)$ agree with those of $\beta(\cdot)$. However, given U the probability that $\beta(U) = 0$ is zero and the continuity of Brownian motion implies that $x(\cdot)$ is *a.s.* not continuous at U [4 marks for the process + 4 for the properties]. [8 marks]

- (c) Given a one dimensional, standard Brownian motion $\beta(\cdot)$, starting from zero, consider the process $x(t) := \int_0^t \beta(s) ds$. Calculate $E_0[x(t)]$, $E_0[(x(t))^2]$ and $E_0[(x(t))^3]$.

[8 marks]

Solution [Unseen]:

(i)

$$E_0 x(t) = E_0 \int_0^t \beta(s) ds = \int_0^t E_0[\beta(s)] ds = 0.$$

(ii)

$$\begin{aligned} E_0(x(t))^2 &= E_0 \int_0^t \int_0^t \beta(r) \beta(s) dr ds = \int_0^t \int_0^t E_0[\beta(r)\beta(s)] dr ds \\ &= \int_0^t \int_0^t \min\{r, s\} dr ds = \int_{0 < r < s < t} r dr ds + \int_{0 < s < r < t} r dr ds \\ &= 2 \int_{0 < r < s < t} r dr ds = \frac{1}{3} t^3. \end{aligned}$$

(d) Show that

$$V_n := \sup_{n \geq 1} \sum_{i=1}^{k_n} |\beta(t_i^n) - \beta(t_{i-1}^n)| = +\infty, \quad P_0 - a.s.$$

[5 marks]

Solution [Unseen example]: Assume that $\sup_n V_n < \infty$. Then

$$D_n \leq \max_{i=1, \dots, k_n} |\beta(t_i^n) - \beta(t_{i-1}^n)| V_n \leq C \max_{i=1, \dots, k_n} |\beta(t_i^n) - \beta(t_{i-1}^n)|.$$

Continuity of the brownian motion and the fact that the mesh of the partition converges to zero implies that the right hand side would converge to zero if $< \infty$, which is a contradiction to the fact that D_n converges to t .

Continued ...

- (c) Let $\beta_1(\cdot)$ and $\beta_2(\cdot)$ be two independent, standard, one dimensional Brownian motions starting from x_1, x_2 , respectively. Compute the probability that they do not meet in the time interval $[0, t]$, $t > 0$. [8 marks]

Solution [Unseen example]: The difference $\beta(s) := (\beta_1(s) - \beta_2(s))/\sqrt{2}$ is a Brownian motion [2 marks]. Therefore, they will not meet by time t , if $\beta(s)$ does not hit zero by time $2t$. This now reduces to the reflection principle: Start from $P(\beta(t) \geq b) = P(\beta(t) \geq b, \tau_b \leq t)$ where τ_b is the hitting time of b [2 marks]. Use the strong Markov property at $\mathcal{F}^+(\tau_b)$. Finally, use the fact that for any t , $P_0(\beta(t) \geq 0) = 1/2$.

$$\begin{aligned} P(\beta(t) \geq b) &= P(\beta(t) \geq b, \tau_b \leq t) = E[P(\beta(t) \geq b, \tau_b \leq t) | \mathcal{F}_{\tau_b}] \\ &= E[P_b(\beta(t - \tau_b) \geq b) 1_{\tau_b \leq t}] = \frac{1}{2} P(\tau_b \leq t) \end{aligned}$$

Apply this formula for $b = x_1 - x_2$. [4 marks]

4. (a) Consider the parabolic problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{1}{2} \Delta u + V(x)u, & t > 0, x \in \mathbb{R}^d, \\ u(0, x) &= f(x) & x \in \mathbb{R}^d. \end{aligned}$$

Show that

$$u(t, x) := E_x \left[f(\beta(t)) \exp \left(\int_0^t V(\beta(s)) ds \right) \right]$$

solves the above boundary value problem. State any assumptions you might need on u, V, f . [10 marks]

Solution [Seen example]: Expand the exponential:

$$\begin{aligned} &1 + \sum_{n \geq 1} \frac{1}{n!} E_x \left[f(\beta(t)) \left(\int_0^t V(\beta(s)) ds \right)^n \right] \\ &= 1 + \sum_{n \geq 1} \frac{1}{n!} E_x \left[f(\beta(t)) \int_0^t \cdots \int_0^t V(\beta(s_1)) \cdots V(\beta(s_n)) ds \right], \end{aligned}$$

which by symmetry writes as

$$1 + \sum_{n \geq 1} E_x \left[f(\beta(t)) \int_{0 < s_1 < \cdots < s_n < t} V(\beta(s_1)) \cdots V(\beta(s_n)) ds_1 \cdots ds_n \right].$$

[2 marks]

Denote the n^{th} term by I_n and change variables $s_i =: t - \eta_i$ to write the above as

$$I_n(t, x) := \int_{0 < \eta_n < \cdots < \eta_1 < t} E_x \left[f(\beta(t)) \int_0^t \cdots \int_0^t V(\beta(t - \eta_1)) \cdots V(\beta(t - \eta_n)) d\eta_1 \cdots d\eta_n \right].$$

The general solution to this equation is $u(x) = A \cos(\sqrt{2\gamma}x) + B \sin(\sqrt{2\gamma}x)$. Satisfying the boundary conditions gives

$$1 = A \cos(\sqrt{2\gamma}a) + B \sin(\sqrt{2\gamma}a)$$

$$1 = A \cos(\sqrt{2\gamma}a) - B \sin(\sqrt{2\gamma}a)$$

(3 marks)

Adding and subtracting the equations gives $A \cos(\sqrt{2\gamma}a) = 1$ and $B \sin(\sqrt{2\gamma}a) = 0$. From this we can choose $B = 0$. However, if $\cos(\sqrt{2\gamma}a) = 0$ then there is no choice of A that can satisfy the boundary condition. So, if $\sqrt{2\gamma}a < \pi/2$ then we can choose

$$A = \frac{\cos(\sqrt{2\gamma}x)}{\cos(\sqrt{2\gamma}a)}.$$

(2 marks)

If γ is such that $\sqrt{2\gamma}a \geq \pi/2$ then there is no solution, either because the boundary condition will force $A = B = 0$ or the above solution takes negative values. So for $\sqrt{2\gamma}a \geq \pi/2$ the exponential moment of τ is infinite **(2 marks)**.

THE END