

学 位 論 文 要 旨				
論 文 提出者	入学年度	平成 29 年度	専 攻	基盤科学系
	氏 名	徳永 優	指導教員	西井 淳
論 文 題 目	<p><b>A Computational Model of Cell Migration of Fish Keratocytes</b></p> <p>Fish keratocytes usually show a circular shape; however, they change their shape to a half-moon shape when they begin migration. This phenomenon suggests that the deformation of the cell shape is a key feature to realize cell migration of keratocytes. The purpose of this research is to clarify the intracellular mechanism that forms a half-moon shape by physical simulation experiments considering intracellular mechanism.</p> <p>Previous studies have reported that during cell migration actin molecules extend their head toward the cell membrane by actin polymerization (AP), which has been suggested as a source of the deformation of the cell membrane and the propulsive force of the cell. The actin retrograde flow (ARF) that pulls the actin molecules back toward the stress fiber (SF), a bundle of actin fibers spreading from side to side of the rear part of the cell, has also been reported.</p> <p>In the computer simulation of this study, the cell membrane was modeled by a network of simple particles interacting with each other and placed on a cylindrical surface as an initial condition. Each particle of the membrane was assumed to receive elastic force from neighboring particles and repulsive force from actin molecules. Actin molecules were modelled by a simple rod, and polymerization and depolymerization were expressed by stochastic elongation of one end of the rod and contraction of the other end, respectively. As an initial condition, actin molecules have no length and were randomly distributed in a U-shaped region, and the direction of the AP of each molecule was randomly determined. As an effect of the ARF, actin molecules were assumed to move toward the SF stochastically and oriented in the direction in which actin molecule is pulled.</p> <p>As a result, the actin molecules aggregated into a half-moon shape under the above simulation conditions. The results also suggested that the effect of aligning the orientation of the AP in the direction of the ARF is important for the formation of a half-moon shape and that the effect of retraction of actin molecule prevents excessive expansion of the cell membrane.</p>			