

CENG 551 Probability and Stochastic Processes for Engineers

Week # 3
Counting & Random Variables

Dr. Deniz Ozdemir
Deniz.ozdemir@yasar.edu.tr
[Web: dozdemir.yasar.edu.tr](http://Web:dozdemir.yasar.edu.tr)



Plan for the Session

- More on Independence
- Counting
- Random variables
- Probability mass function (pmf)
 - Binomial Random Variable
- Expectation
 - Example

Independence

- Simply put $P(A/B) = P(A)$
- This implies that


$$P(A \cap B) = P(A/B) P(B) = P(A) P(B)$$



Conditioning may affect independence

- Assume A and B are independent:
- If we are told that C occurred, are A and B **MAY NOT BE** independent.
- Two independent fair ($p=1/2$) coin tosses.

- Event A: First toss is H
- Event B: Second toss is H
- Event C: The two outcomes are different.



4

10/8/2012

Independence of a Collection of Events



- Intuitive definition:
 - Information about some of the events tells us nothing about probabilities related to remaining events.
 - Example: $P(A_1 \cup (A_2 \cap A_3) | A_4 \cap A_5) = P(A_1 \cup (A_2 \cap A_3))$
- Mathematical definition:
 - For any distinct i, j, \dots, q

$$P(A_i \cap A_j \cap \dots \cap A_q) = P(A_i) P(A_j) \dots P(A_i)$$


10/8/2012

5

Independence vs. Pairwise Independence



- Example 1 revisited
 - Two independent fair ($p=1/2$) coin tosses.
 - Event A: First toss is H
 - Event B: Second toss is H
 - Event C: The two outcomes are different.
 - $P(A) = P(B) = P(C) = 1/2$
- $P(A \cap C) =$
- $P(C | A \cap B) =$



- Pairwise independence **does not** imply independence.

10/8/2012

6



True or False?

- If $P(A|B) = P(A)$, then $P(B|A^c) = P(B)$.
- If 10 out of 10 independent fair coin tosses resulted in tails, the events “first toss was tails” and “10th toss was tails” are independent.
- If 5 out of 10 independent fair coin tosses resulted in tails, the events “first toss was tails” and “10th toss was tails” are independent.
- If the events A_1, \dots, A_n form a partition of the sample space, and if B, C are some other events, then

$$P(B|C) = \sum_i P(A_i|C)P(B|A_i)$$

10/8/2012

7



Discrete Uniform Law

- Let all sample points be equally likely
- Then,

$$P(A) = \frac{\text{number of elements of } A}{\text{total number of sample points}}$$

- Just count ...

10/8/2012


Modelos Probabilistas Aplicados

8

Basic Counting Principle



- r steps
- n_i choices at step i
- Number of choices is:
 $n_1 n_2 \dots n_r$
- Number of license plates with 3 letters a 4 digits =
- ...if repetition is prohibited =
- **Permutations:** Number of ways of ordering n elements is =



10/8/2012

9

Example



- Probability that six rolls of a six-sided die all give different numbers?
 - Number of outcomes that make the event happen =
 - Number of elements in the sample space =
 - Answer =

10/8/2012

10



Combinations

- $\binom{n}{k}$: number of k -element subsets of a given n element set.
 - Two ways of constructing an ordered sequence of k **distinct items**:
 - Choose the k items **one** at a time:
 $n(n-1)\dots(n-k+1) = \frac{n!}{(n-k)!}$ choices
 - Choose k items, then order them ($k!$ possible orders) $\binom{n}{k} k! = \frac{n!}{(n-k)!}$
- $$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

10/8/2012

11



Combinations: Some Properties

- $\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r} \quad 1 \leq r \leq n$
- $(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$
- Identity: $\sum_{k=0}^n \binom{n}{k} =$

10/8/2012

12

Basic Counting Principle



Number of Possible Arrangements of Size r from n Objects:

	Without Replacement	With Replacement
Ordered:	$\frac{n!}{(n-r)!}$	n^r
Unordered:	$\binom{n}{r}$	$\binom{n+r-1}{r}$

10/8/2012

13

Binomial Probabilities



- n independent coin tosses
 - $P(H) = p$
 - $P(HTTHH) =$
 - $P(\text{sequence}) = p^{\# \text{ of heads}} (1-p)^{\# \text{ of tails}}$
 - $P(k \text{ heads}) =$

$$= \sum_i k\text{-head seq}_i. P(\text{seq}_i)$$

$$= (\# k\text{-head seq}) p^k (1-p)^{n-k}$$

$$= \binom{n}{k} p^k (1-p)^{n-k}$$

10/8/2012

14

Birthday Problem

- At a gathering of s randomly chosen students what is the probability that at least 2 will have the same birthday?

$P(\text{at least 2 have same birthday}) = 1 - P(\text{all } s \text{ students have different birthdays})$.

Assume 365 days in a year. Think of students' birthdays as a sample of these 365 days.

The total number of possible outcomes is:

$$N = 365^s \text{ (ordered, with replacement)}$$

The number of ways that s students can have different birthdays is

$$M = 364! / (365 - s)! \text{ (ordered, without replacement)}$$

$$P(\text{all } s \text{ students have different birthdays}) =$$


$$M / N.$$

10/8/2012

15

Birthday Problem

Probability that all students have different birthdays



10/8/2012

16

Coin Tossing Problem



- Event B: 3 out of 10 tosses were “heads”.
 - What is the (conditional) probability that the first 2 tosses were heads, given that B occurred?
- All outcomes in conditioning set B are equally likely:
 - Probability: $p^3(1-p)^7$
 - Conditional probability law is uniform.
- Number of outcomes in B:
- Out of the outcomes in B, how many start with HH?

10/8/2012

17

Partitions



- 52-card deck, dealt to 4 players.
- Find $P(\text{each gets an ace})$
- Count size of the sample space (possible combination of “hands”)
- Count number of ways of distributing the four aces: $4 \cdot 3 \cdot 2$
- Count number of ways of dealing the remaining 48 cards
- Answer:

10/8/2012

18

Another Counting Rule



- The number of ways of classifying n items into k groups with r^i in group i , $r^1 + r^2 + \dots + r^k = n$, is:
- $n! / (r^1! r^2! r^3! \dots r^k!)$

10/8/2012

19

Hogwart's Problem



- How many ways are there to assign 100 incoming students to the 4 houses at Hogwarts?
 $\sim 1.6(10^{57})$

10/8/2012

20

Random Variables



- Before this we talked about “Probabilities” of events and sets of events where in many cases we hand selected the set of fine grain events that made up an event whose probability we were seeking. Now we move onto another more interesting way to group this point: using a function $p_x(x)$ to ascribe values to every point in a sample space (discrete or continuous)
- One example might be the number of heads in 3 tosses of a coin.

10/8/2012

Modelos Probabilistas Aplicados

21

Random Variables



- An assignment of a value (number) to every possible outcome.
- Mathematically: A function from the sample space to the real numbers:
 - –Discrete or Continuous
- Can have several random variables defined on the same sample space

10/8/2012

22

Random Variables



- A random variable (r.v.) associates a **unique numerical value** with each outcome in the sample space.
- Notation:
 - Random Variable X
 - Experimental Value x
- Example: 1 coin toss. Define X :
 - $X = \begin{cases} 1 & \text{if } x = H \\ 0 & \text{if } x = T \end{cases}$ i.e. $X(H) = 1, X(T) = 0$
- Example: $Y = g(X)$

10/8/2012

23

Random Variables



- Temperature in DC on Feb 1.
- Length of queue at Migros
- Amount of water in a “tall Americano”
- The number of points Efes score in a game they win
- The number of words in your emails

10/8/2012

24

Discrete Random Variables



- **Discrete random variables:** number of possible values is finite or countably infinite:

$x_1, x_2, x_3, x_4, x_5, x_6, \dots$

- *Probability mass function (p.m.f.)*

- $f(x) = P(X = x)$ (Sum over all possible values = 1 always)

- *Cumulative distribution function (c.d.f)*

- $F(x) = P(X \leq x) = \sum_{k \leq x} f(k)$

10/8/2012

25

Probability Mass Function



- (“probability law”,
- “probability distribution”)
- Notation:

$$p_X(x) = P(X = x)$$

- **Example:** X = number of coin tosses until first head

- Assume independent tosses, $P(H) = p > 0$

- $p_X(k) = P(X = k)$

$$= P(TT \dots TH)$$

$$= (1-p)^{k-1}p, \quad k=1,2,\dots$$

*Geometric probability distribution
With parameter p*

10/8/2012

26

How to compute a pmf $p_X(x)$

- Collect all possible outcomes for which X is equal to x :
 $\{\omega \in \text{Sample Space} / X(\omega)=x\}$
- Add their probabilities.
- Repeat for all x .
- **Example:** Two independent throws of a fair tetrahedral die:
 - F : outcome of first throw
 - S : outcome of second throw
 - $L = \min(F, S)$

$$p_L(2) =$$

S = Second roll

4				
3				
2				
1				
	1	2	3	4

F = First roll

10/8/2012

27

Probability Mass Function

- $p_x(x_0)$ probability that the experimental value of a random variable X obtained on a performance of the experiment is equal to x_0
- The same story value of Probability Mass Function (pmf). It can extend up to more dimensions which then allows for conditional pmfs

$$\sum_i p_x(x_0) \quad 0 \leq p_x(x_0) \leq 1$$

10/8/2012

28

Binomial pmf with p and n



- X : number of heads in n independent coin tosses
 - $P(H) = p > 0$
- Let $n = 4$
- $p_X(2) =$

$$= 6p^2(1-p)^4$$

$$= \binom{4}{2} p^2(1-p)^4$$

- In general:

$$p_X(k) = \binom{n}{k} p^k(1-p)^{n-k}, \quad k = 1, 2, \dots, n$$

10/8/2012


29

Selected Discrete Distributions:



- Single Coin flip $P(\text{heads/success}) = p$
 - *Probability of success*

$$f(x) = P(X = x) = \begin{cases} p & \text{if } x = 1 \text{ (success)} \\ 1 - p & \text{if } x = 0 \text{ (failure)} \end{cases}$$



- *"Bernoulli Distribution"*
- Multiple coin flips
 - x successes out of n trials
 - $f(x) = P(X = x) = \binom{n}{x} p^x(1-p)^{n-x}$ for $x = 1, \dots, n$
 - *"Binomial distribution"*

10/8/2012


30

Selected Discrete Distributions:

- Number of failures before the first success

- $f(x) = P(X=x) = (1-p)^{x-1}p, \quad x=1,2,\dots$

- “Geometric Distribution”



- N equally likely events

- $f(x) = P(X=x) = 1/N, \quad x=1,2,\dots$

- “Uniform distribution”



U

10/8/2012

31

Selected Discrete Distributions:

- *Hypergeometric distribution*

- Drawing balls from the box without replacing the balls

- *Poisson Distribution*

- number of occurrences of a rare event

- *Multinomial Distribution*

- more than two outcomes

- *Negative Binomial Distribution*

- number of trials to get r successes

10/8/2012

32

Expectation

- The expected value or mean of a discrete r.v. X , denoted by $E(X)$, is defined as:

$$E[X] = \sum_x x \cdot p_X(x)$$

- Interpretations:
 - Center of gravity of pmf.
 - Average in large number of repetitions of the experiment. (to be substantiated later in this course)
- This is essentially a weighted average of the possible values the r.v. can assume, weights = $f(x)$

10/8/2012

33


Expectation

- Example: Uniform on $0, 1, \dots, n$

$$E[X] = \sum_x x \cdot p_X(x)$$

- $E[X] =$

$$= 0 \times \frac{1}{n+1} + 1 \times \frac{1}{n+1} + \dots + n \times \frac{1}{n+1}$$



10/8/2012

34

Properties of Expectations -1



- Let X be a r.v. and let $Y = g(X)$
 - $E(Y) = \sum_y y \cdot p_Y(y)$
 - $E(Y) = \sum_x g(x) \cdot p_X(x)$
- “Second Moment”: $E(X^2) = \sum_x x^2 p_X(x)$
- Caution: In general,

$$E[g(X)] \neq g(E[X])$$



10/8/2012

35

Expected Values



- $E(x)$ given a p.m.f. provides some sense of the center of mass of the pmf.
- Variance is another measure that provides some measure of the distribution of a pmf/pdf around its expected value.
- Variance: $\text{var}(X) = E[(X - E[X])^2]$
$$= \sum_x (x - E[X])^2 p_X(x)$$

10/8/2012

36

Properties of Expectations -2



- If α, β are constants, then:

- $E[\alpha] =$
- $E[\alpha X] =$
- $E[\alpha X + \beta] =$
- $E[X - E(X)] =$




10/8/2012

37

Average Speed vs. Average Time



- Traverse a 200 mile distance at constant but random speed V :



- $d = 200, T = t(V) = 200/V$
 - $E[V] = 1 \cdot \frac{1}{2} + 200 \cdot \frac{1}{2} = 100.5$
 - $E[T] = \sum_v t(v) p_V(v)$
 $= \frac{200}{1} \cdot \frac{1}{2} + \frac{200}{200} \cdot \frac{1}{2} = 100.5$

10/8/2012

38

Average Speed vs. Average Time vs. Expected Distance



- $E[d] =$
- $E[T] \neq 200/E[V]$
- $\text{Var}(V) = \sum_v (v - E[V])^2 p_V(v)$
 $= (1 - 100.5)^2 \cdot \frac{1}{2} + (200 - 100.5)^2 \cdot \frac{1}{2}$
 $\approx 10\,000$
- Standard Deviation $\sigma_V = \sqrt{\text{var}(V)} \approx 100$



10/8/2012

39

Expectations:



- 4 buses carrying 148 job-seeking students arrive at a job fair. The buses carry, respectively, 40, 33, 25, and 50 students. One of the students is randomly selected. Let X denote the number of students that were on the bus carrying this randomly selected student. Also, one of the 4 bus drivers is also randomly selected. Let Y denote the number of students on his bus.

Do you think $E[X]$ and $E[Y]$ are equal?

$$E[X] = \frac{40}{148} \cdot 40 + \frac{33}{148} \cdot 33 + \frac{25}{148} \cdot 25 + \frac{50}{148} \cdot 50 \approx 39.3$$

$$E[Y] = \frac{1}{4} \cdot 40 + \frac{1}{4} \cdot 33 + \frac{1}{4} \cdot 25 + \frac{1}{4} \cdot 50 = 37$$

10/8/2012

40

Variance and Standard Deviation



- The variance of an r.v. X , denoted by $\text{Var}(X)$, or simply σ^2 , is defined as:
 - $\text{Var}(X) = \sigma^2 = E[(X - E[X])^2]$
$$\begin{aligned}\text{Var}(X) &= E(X^2 - 2E[X]X + (E[X])^2) \\ &= E(X^2) - 2E[X].E(X) + E((E[X])^2) \\ &= E(X^2) - 2(E[X])^2 + (E[X])^2 \\ &= E(X^2) - (E[X])^2\end{aligned}$$
- The standard deviation (SD) is the square root of the variance.
- Note that the variance is in the square of the original units, while the SD is in the original units.

10/8/2012

41

Next Time...



- Conditional Expectations
- Bernoulli & Binomial Random Variables

42