



# Minimum Swaps 2 ☆

Problem

Submissions

Leaderboard

Discussions

Editorial

You are given an unordered array consisting of consecutive integers  $\mathbb{E}[1, 2, 3, \dots, n]$  without any duplicates. You are allowed to swap any two elements. You need to find the minimum number of swaps required to sort the array in ascending order.

For example, given the array **arr** = [7, 1, 3, 2, 4, 5, 6] we perform the following steps:

i	arr	swap (indices)
0	[7, 1, 3, 2, 4, 5, 6]	swap (0, 3)
1	[2, 1, 3, 7, 4, 5, 6]	swap (0, 1)
2	[1, 2, 3, 7, 4, 5, 6]	swap (3, 4)
3	[1, 2, 3, 4, 7, 5, 6]	swap (4, 5)
4	[1, 2, 3, 4, 5, 7, 6]	swap (5, 6)
5	[1, 2, 3, 4, 5, 6, 7]	

It took **5** swaps to sort the array.

Function Description

Complete the function minimumSwaps in the editor below. It must return an integer representing the minimum number of swaps to sort the array.

minimumSwaps has the following parameter(s):

- arr: an unordered array of integers

Input Format

The first line contains an integer, **n**, the size of **arr**.

The second line contains **n** space-separated integers **arr[i]**.

Constraints

- $1 \leq n \leq 10^5$
- $1 \leq arr[i] \leq n$

Output Format

Return the minimum number of swaps to sort the given array.

Sample Input 0

```
4
4 3 1 2
```

Sample Output 0

```
3
```

Explanation 0

Given array **arr** : [4, 3, 1, 2]

After swapping (0, 2) we get **arr** : [1, 3, 4, 2]

After swapping (1, 2) we get **arr** : [1, 4, 3, 2]

After swapping (1, 3) we get **arr** : [1, 2, 3, 4]

So, we need a minimum of **3** swaps to sort the array in ascending order.

Sample Input 1

```
5
2 3 4 1 5
```

Sample Output 1

```
3
```

Explanation 1

Given array **arr** : [2, 3, 4, 1, 5]

After swapping (2, 3) we get **arr** : [2, 3, 1, 4, 5]

After swapping (0, 1) we get **arr** : [3, 2, 1, 4, 5]

After swapping (0, 2) we get **arr** : [1, 2, 3, 4, 5]

So, we need a minimum of **3** swaps to sort the array in ascending order.

Sample Input 2

```
7
1 3 5 2 4 6 7
```

Sample Output 2

```
3
```

Explanation 2

Given array **arr** : [1, 3, 5, 2, 4, 6, 7]

After swapping (1, 3) we get **arr** : [1, 2, 5, 3, 4, 6, 7]

After swapping (2, 3) we get **arr** : [1, 2, 3, 5, 4, 6, 7]

After swapping (3, 4) we get **arr** : [1, 2, 3, 4, 5, 6, 7]

So, we need a minimum of **3** swaps to sort the array in ascending order.

Author

rishi\_07

Difficulty

Medium

Max Score

40

Submitted By

162952

NEED HELP?

View discussions

View editorial

View top submissions

RATE THIS CHALLENGE

☆☆☆☆

MORE DETAILS

Download problem statement

Download sample test cases

Suggest Edits