MNRAS 000, 1–6 (2026) Preprint 8 November 2026 Compiled using MNRAS LATEX style file v3.2

Dark Matter from Baryon Density

Thomas C. Andersen[1]
[1]nSCIr, Ontario, Canada

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present a novel empirical equation for the dark matter derived only from galactic (gas) kinematics. It is defined simply by the equation $\rho_{\rm dm} = k\ m_e\alpha^2/a_B d^2$, where d is the mean particle spacing in normal matter, a_B is the Bohr radius, α and m_e are the usual. k is a dimensionless constant near 1. The meaning of the relationship is left open (the constant is related to the atomic unit of force). We evaluate its applicability by testing this relation against the complete Spitzer Photometry and Accurate Rotation Curves (SPARC) database, with over 3000 data points across 175 morphologically diverse galaxies. The equation varies the dark mass density with the 2/3 power of the matter density. It results in dark matter starting to be substantial when (gas) particles are about a millimetre apart. The model generates rotation curves that are very similar to MOND, but has the advantage that it generates massive dark matter. With a simple density function for the circumgalactic medium (CGM) gas, it simultaneously produces appropriate dark matter halo masses as well.

Key words: dark matter – galaxies: kinematics and dynamics – galaxies: haloes

1 INTRODUCTION

1.1 History

For over four decades, the standard cosmological paradigm (ΛCDM) has successfully explained large-scale structure formation, cosmic microwave background anisotropies, and cosmic shear observations by invoking a dominant component of cold dark matter. However, at galactic and sub-galactic scales, the nature of this missing mass field remains one of the most persistent crises in modern physics. Despite intensive efforts spanning astronomical observations, deep underground direct-detection experiments, and high-energy collider searches, particle dark matter candidates—ranging from Weakly Interacting Massive Particles (WIMPs) to ultra-light axions—have eluded detection across nearly thirty orders of magnitude in predicted mass scales. Even assuming any (perhaps undetectable) particle like dark matter, problems persist. For example, modelling the dark matter distribution in galaxies, where Sancisi's Law(Sancisi 2004) and MOND (Lelli et al. 2016, 2017) work well. In addition there are several works which discount the viability of any Cold Dark Matter particle model(Kroupa et al. 2023; Famaey & McGaugh 2013).

1.2 Current Approaches

There are currently two primary paths of investigation in galactic dynamics. The first relies on multi-parameter empirical dark matter density profiles, such as the Navarro-Frenk-White (NFW)(Navarro et al. 1997) or Burkert profiles(Burkert 1995), which are fitted to individual galactic rotation curves. While these profiles effectively capture overall kinematic features, they generally treat parameters like scale radius (r_s) and central density (ρ_0) as free, independent variables unique to each galaxy. This introduces a large parameter space that requires individual, galaxy-by-galaxy 'tuning.' The second path, Modified Newtonian Dynamics (MOND), abandons the dark matter hypothesis entirely, accounting for anomalous galactic rotation curves by introducing a universal acceleration constant ($a_0 \approx 1.2 \times 10^{-10}$ m s^{-2}) below which Newtonian gravity breaks down. MOND's striking empirical success, particularly in explaining the Radial Acceleration Relation (RAR) and the Baryonic Tully-Fisher Relation (BTFR)(Lelli et al. 2017), demonstrates that galactic kinematics are deeply coupled to the distribution of baryonic matter.

1.3 An alternative: Gas Based Models

Sancisi and others have pointed out the puzzling connection of gas density to the rotation curves. In these gas models, gas is scaled by a factor of about 7(Hoekstra et al. 2001). The Hoekstra method of simply multiplying the gas mass by a number doesn't work in general, and as more and better data came along, such as the SPARC dataset, it was put aside.

1.4 Our Model

The density of dark matter is obtained by this relation, where k is scaling constant near one.

$$\rho_{\rm dm} = k\ m_e\alpha^2/a_B d^2. \quad (1)$$

d is the mean particle spacing in normal matter, a_B is the Bohr radius, α is the fine structure constant, and m_e the electron mass.

The $1/d^2$ can also be written as $n^{2/3}$, where n is the number density of matter particles. Thus, as a gas becomes more rarefied, the amount of dark matter 'generated' by our model as compared to the density of the particles increases. It in fact increases without limit in equation (1). This is an issue for intergalactic levels of particle densities, so for densities less than $n_{cut} = 10^{-5}/cc$ we turn off the effect with an exponential switch. This effect cutoff has no effect on galactic rotation curve fitting, but is used to turn off our model at the

© 2026 The Authors

halo - intergalactic boundary near R_{200}. The physical justification for this (if there is one), is that n_{cut} is about where gas particles stop behaving like a gas and are instead ballistic.

With $k \approx 1$, a gas with a particle number density of $10^{-5} cm^{-3}$ would have a dark matter density of about 150 times that of the gas itself, while at $100 cm^{-3}$ our equation (1) only adds about 50% to the density of the gas. (1) might even hold in the everyday matter around us, for example for water, this formula would add about 0.001g to 0.1g to a ton of water. We assume that this dark mass has a negligible effect on individual particle dynamics.

During the investigation, we determined that many SPARC galaxies have 'HI holes' near the core, which is a physically real phenomenon. There is more gas in galaxies than the HI gas listed in the SPARC database.

- **CGM gas**: Not in SPARC: our model makes it contribute some mass even inside R_{max}. Outside R_{max} the CGM gas combined with equation (1) constructs a dark matter halo.
- **H2/Molecular**: Included in SPARC, but hidden in the stellar masses. We require particle densities to make equation (1) work.
- **Others**: .
- **Electrons**: equation (1) has no reference to baryons, ionized gas might therefore count twice. We don't consider this effect in our model. It would serve to cut the required added mass of ionized gas by approximately 1/2.

The extent and mass of some of these particle densities is unknown in detail. Most fits of our model add gas in to replace this hidden gas.

The SPARC dataset only includes HI gas measurements. There are other sources of gas in a galaxy. In addition, the centres of many galaxies are opaque to some gas measurements(Saintonge & Catinella 2022), so the gas composition and densities at the cores of galaxies are not well known, especially for the purposes of our model, where hidden gas in the galactic core cannot be subsumed into a minor dynamical change in the stellar luminosity parameter, Σ_{bulge} (Lelli et al. 2016).

2 METHODOLOGY

In this paper, references to 'Lelli' are referring to the *Lelli, McGaugh, and Schombert - SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves.* (Lelli et al. 2016). When we refer to Lelli, we are using unmodified data, etc from that paper.

We perform 5 runs of our model, each run is over all 175 galaxies.

- **Newton**: A check on our model.
- **Best k**: A fit to determine the best scaling factor k, using only the gas in SPARC dataset.
- **Core Gas Global**: a single global parameter is used to add gas to each galaxy. (one parameter)
- **Core Gas Individual**: each galaxy is fitted with one parameter, the amount of gas to add. (175 parameters)
- **CGM**: Both core gas fits have a circumgalactic medium (CGM) gas added, which models the dark matter halo.

2.0.1 Stellar Mass Luminosity Ratio

The Stellar Mass Luminosity Ratio(Υ or $M_\odot/L_\odot$) is the ratio of the total mass of a stellar population (or galaxy) to its total luminosity. Following Lelli et al. (2016), we adopt $\Upsilon_{\rm disk} = 0.5$ and $\Upsilon_{\rm bul} = 0.7$. The SPARC gas term includes only H I (scaled by 1.33 for helium); molecular gas is neglected, and where it follows the stars it is implicitly part of $\Upsilon_\star$. This matters most in the inner regions of bulge galaxies, where H_2 often replaces a central H I depression, so some of the mass attributed to the bulge there may be gas, which our added core gas models.

2.0.2 SPARC gas density

Since our model requires a gas density, and the SPARC dataset only has a gas velocity, a small program was written to determine plausible densities, given the V_{gas} data in SPARC. The problem is underdefined. We required the gas distribution to be as smooth as possible while still recreating the V_{gas} from SPARC (which itself was created by analysis of gas density measurements). In run 0 we plotted this gas as a velocity for each galaxy to verify that our calculated gas densities were reasonable.

2.1 General outline of the fits

One of the main goals was to deviate from the parameters and methods in Lelli only when required.

2.1.1 3D volumetric approach

We determine the acceleration due to gravity at each R_{obs} radius of the SPARC dataset from all baryons with the addition of dark mass from our model. The chosen method uses a 3D volumetric approach where each galaxy is modelled in a cylindrical system with a few million cells per galaxy. Note that since the velocity contributions from stars are not modified, the 3D volume model is only used to store particle densities in each cell. Each cell then has an X, Y, Z position, a density of gas particles, and a volume. The gravitational attraction for each of the radial points in the SPARC dataset is determined by summing the millions of individual cell masses at their distance from each radial test point. This method thus covers induced acceleration from matter at radii beyond the radial measuring point. Each galaxy is modelled as a disk with thickness as calculated using the same formula as the SPARC 2016(Lelli et al. 2016): an exponential vertical distribution with scale half height $z_d = 0.196 R_d^{0.633}$. The bulge is ignored for the reconstructed SPARC gas. Any gas added beyond our SPARC reconstruction is done in a spherically symmetric way. The 3D model is large enough (1.1 times R_{max}) so that any gas—whether in or outside of the disk—will contribute to acceleration and mass.

2.2 Error and χ^2 fitting

The SPARC data include errors in the observed rotation velocities, but the errors in the other parameters are unknown and uncertain. For instance errors in inclination directly affect the reported V_{obs}, and gas density errors effect the V_{gas}, etc. In (Li et al. 2018) , the authors fit the 175 SPARC galaxies individually, with up to 4 free parameters per galaxy. Many of the best fit parameters such as inclination are a factor of 2 or more away from accepted values. This is an indication of the uncertainty in the physical layout of our galactic neighbours.

To capture this uncertainty, we assume a 15% error in the calculated V_{tot}. If an error is not assumed fitting algorithims will pour their effort into minimizing the error on a few of the galaxies that have a large number of highly accurate V_{obs}. Our model also magnifies uncertainties - the dark mass contribution in our model has a higher uncertainty. Changing the assumed error by doubling it or




Figure 1. Gas density plot for NGC 300. The galaxy has an additional $0.9cm^{-3}$ gas particles added at the core, plus a CGM gas halo. The CGM halo extends far beyond the diameter shown here, which is 1.1 times R_{max}.




Figure 2. NGC 6946 - Newton run. No gas added, our model turned off. This is an example of a check on our gas density recreation. The "baryons + dm" line is directly on top of the blue 'baryons (Lelli)' line which is our test. in addition, our 'as used' gas line is (almost) on top of the Lelli gas line, which shows the accuracy of our gas density recreation.

halving it only made marginal changes to the derived fit parameters. (Schombert et al. 2022) quote a 10% error on gas components.

In addition to this χ^2 fitting algorithm, other fitting measures were tested, but the overall results only changed best fit outputs by a small amount.

Error results are shown as per point only, as the the measurements are highly correlated. Due to the exploratory nature of this work, a full Bayesian analysis is outside the scope.

3 DATA

We discuss the runs. Data is shown in table **??**. Images for each galaxy, detailed run statistics and more are available online: (git url with commit hash)

3.1 Newton run

With the SPARC gas recreated as a density, we ran our model over all 175 galaxies. As this run completely lacks our dark matter model, it serves as a baseline. A single galaxy is shown in Figure 3.1.

3.2 Best k

We obtainied a dark mass for each cell by using our model on the supplied SPARC gas. The entire 175 galaxy set, with 3319 data points was fitted to obtain a best power.

The fit generated one value, $k = blah$

3.3 Core Gas Global

In order to add a gas core to every galaxy in the dataset, we needed to scale the added gas to the mass and/or size of the galaxy. The additional gas is added via this formula:

$$n_{\rm core}(r) = n_0 \left[1 + \left(\frac{r}{f\, r_s}\right)^2\right]^{-\beta} . \quad (2)$$

Where $n_{\rm core}(r)$ is the added density of gas at the spherical radius r. n_0 is the central particle density, $\beta = 2.25$, $f = 2.5$, and $M_{\rm gal} = \left(L_{[3.6]}/L_\odot + M_{\rm HI}/M_\odot\right) M_\odot$ and:

$$r_s = 2\,{\rm kpc} \left(\frac{M_{\rm gal}}{10^{11}\, M_\odot}\right)^{1/3} . \quad (3)$$

Note: $L_{[3.6]}$ and $M_{\rm HI}$ are directly from SPARC database columns.

The only free parameter is n_0. In this fit, a single best n_0 was found by minimizing the error while at the same time optimizing the scaling constant, k to be near 1.

Result:

- chosen k: xxxx,
- chosen core gas beta: xxxx,
- chosen core gas factor xxxx:,
- best fit n_0: $0.55cm^{-3}$,
- $\chi^2_{core\ gas} = xxx$ per point.

Thus there are 1 - 5 paramters added in total, depending on how one counts the gas addition equation parameters.

3.4 Core Gas Individual

In this fit, we ran with the same parameters as the Core Gas Global fit, with the one difference being an optimizer on each galaxy, determining the best fit for n_0 for each of the 175 galaxies individually. The physical argument for adding 175 parameters to the fit, is that the unknown gas in a galaxy will naturally vary between individual galaxies.

- chosen k: xxxx,
- chosen core gas beta: xxxx,
- chosen core gas factor f: xxx,
- best fit n_0: $\in [0, 6]cm^{-3}$,
- $\chi^2_{each\ gas} = xxx$ per point.

3.5 The MOND fit

The MOND fit in this work is a very simple implementation: Using the MOND formula in the form given by (Lelli et al. 2017) with Lelli gas only:




Figure 3. Example individual one parameter per galaxy fit. The only fit parameter 'P' is fixed for all 175 galaxies once it is determined. The purple line shows the total fit by this model. All 175 images are available online. Renzo's rule is honoured at 4 kpc in the IC2574 plot. Note that IC2574 requires almost zero extra core gas for an excellent fit.

$$g_{\rm bar} = \frac{V_{\rm disk}^2 + V_{\rm bul}^2 + V_{\rm gas}|V_{\rm gas}|}{R}, \tag{4}$$

$$g_{\rm MOND} = \frac{g_{\rm bar}}{1 - e^{-\sqrt{g_{\rm bar}/g_\dagger}}}, \tag{5}$$

$$V_{\rm MOND} = \sqrt{g_{\rm MOND}\, R}, \qquad g_\dagger = 1.2 \times 10^{-10}\ {\rm m\,s^{-2}}. \tag{6}$$

Result:

- $\chi^2_{MOND} = 50$ per point.

We run our python program with SPARC data as given, then adding dark mass as a function of the SPARC gas density.

3.6 The Universal Scaling Model

The local dark matter density ($\rho_{\rm dm}$) is assumed to follow a geometric power-law relation:

$$\rho_{\rm dm} = P/c^3 \cdot n_b^{2/3}, \tag{7}$$

where n_b represents the local volumetric baryon density (calculated here from the raw SPARC neutral hydrogen gas profiles), and P is a universal invariant constant across all systems, determined by a best fit to the SPARC data. Our best fit of the parameter P is:

$$P \approx 59\ Watts. \tag{8}$$

3.7 Statistical Optimization via Least Absolute Deviations (LAD)

Because galactic rotation curve data points exhibit localized spatial dependencies (e.g., radial proximity within individual galaxies sharing systematic streaming motions or beam smearing), traditional ordinary least squares (χ^2) approaches can be disproportionately skewed by local outliers. We instead evaluate the goodness-of-fit using the Least Absolute Deviations (LAD) framework.

The global model parameters are optimized by directly minimizing the total absolute residual sum of the observed ($V_{\rm obs}$) and predicted ($V_{\rm pred}$) rotation velocities across the entire ensemble of $N = 3,391$ data points:

$$\text{Total LAD} = \sum_{i=1}^{N} |V_{{\rm obs},i} - V_{{\rm pred},i}| \tag{9}$$

The resulting model fit is characterized via the Mean Absolute Error (MAE), defined as MAE = Total LAD$/N$, providing a, velocity-space measure of global model precision in units of km/s.

4 PYTHON CODE DESCRIPTION

The raw data is read in from the SPARC website publically availble data, blah blah

5 RESULTS

6 DISCUSSION

6.1 Newtonian Dynamics

6.2 The interpretation of Power

Power is about 10W. If a single baryon were literally emitting or processing 10 Watts of power, it would burn through its own rest mass in a fraction of a nanosecond. Therefore, P cannot be a literal power output of an individual particle. It could be a steady state relationship (as it relies on neighboring particles. Or, Power might represent a coupling constant of baryons to the vacuum or spacetime itself. It defines a linear mass density scale ($P/c^3 \approx 10^{-24}$ kg/m) that dictates how baryonic mass curves space beyond its Newtonian boundary.

6.3 Model Results

The global optimization of our one-parameter model across all SPARC observations, results in a MAE of 26 km/s. The MOND framework as implemented here scores (MAE = 16 km/s). Our model achieves a remarkably similar accuracy profile using only a single universal parameter, like MOND. But unlike MOND, our model does not require the local particles to 'know' the global acceleration parameter, rather, they just have to 'know' the density of the material they are in.

6.4 Model Variations

We also ran the model by adding 175 more fitting parameters, allowing each galaxy fit to adjust the entire gas density up or down by up to a factor of 5. This resulted in an (MAE = 17 km/s).

In several SPARC radial bins, central core gas densities are recorded as zero. To test sensitivity to this, we implemented a physically motivated observational detection floor of 0.45 baryons/cm^3 for unrecorded or low gas densities. Enforcing this floor shifts the global MAE by a negligible fraction, but it does make the global fit line up with Newton extremely well.

These two model variations were not used in any of the other discussions in this paper, full results for them and a few others are available online.




Figure 4. All 3319 SPARC data points are plotted. The acceleration implied by the observed velocity V_{obs} is on the vertical axis, the x axis has the full acceleration from baryons + the dm field introduced in this paper. The red line is a Newtonian expectation, the blue line is a (log) straight line fit to the data. Also: velocity residuals.

6.5 Core Dynamics and Distribution Tails

An analysis of the absolute residual distributions reveals that MOND's localized advantage is concentrated almost entirely in a tight low-error spike between 0 and 15 km/s, corresponding to high-density inner galactic cores. This divergence is physically expected; our model evaluates dark mass generation based purely on the gas density profiles, which frequently drop or fluctuate in stellar-bulge dominated cores. Crucially, in the intermediate and long-range tails (25 to 100 km/s) - representing the low-acceleration, outer-disk environments - the error distributions of our model and MOND track one another identically. This strongly implies that in the highly rarefied outer halo regime, our geometric density relation functions as a direct mathematical dual to MOND's physical behaviour, without requiring complex acceleration interpolations.

[Figure 1: Overlapping Histogram of Absolute Velocity Residuals for Both Models vs. MOND]

Figure 5. Distribution of absolute velocity residuals $|V_{\rm obs} - V_{\rm pred}|$ evaluated across the 3,391 data points of the SPARC dataset. Vertical dashed lines point out the global MAEs.

[Figure 2: Absolute Model Error vs. Distance from Galactic Center (Radius in kpc)]

Figure 6. Moving average trend line showing absolute model errors as a function of radial distance from galactic centers, indicating where core fluctuations stabilize.

7 CONCLUSION

In this paper, we have evaluated a universal, one-parameter empirical dark matter density profile defined by the geometric relation $\rho_{\rm dm} = P/c^3 \cdot n_b^{2/3}$ against the complete SPARC database. By optimizing the global constant via robust Least Absolute Deviations (LAD) regression across 3,391 data points from 175 morphologically diverse galaxies, our model achieves a global Mean Absolute Error (MAE) of 26 km/s. This performance is comparable to the MOND framework (MAE = 16 km/s). Both frameworks use a single parameter to model DM across all galaxies.

The statistical parsimony of this model provides compelling evidence for a universal coupling between baryonic density and dark mass distribution. We demonstrate that the $2/3$ scaling law captures the observations of the dataset.

An analysis of the absolute residual distributions reveals that while MOND maintains an advantage in dense, stellar-dominated galactic cores, the two models track one another identically across the low-acceleration, outer-disk environments. This suggests that in the highly rarefied, widely spaced gas structures of the outer halo, our geometric density relation functions as a direct mathematical dual to MOND's phenomenological behaviour.

By focusing strictly on a data-driven, observational approach, this work side-steps the unconstrained parameter space of particle physics models that have remained undetected across decades of microphysical searches. We do not propose a specific microphysical mechanism for the $2/3$ scaling law, nor do we speculate on the

fundamental properties of a constituent particle. Instead, we present this universal relation as an empirical reality mandated by galactic kinematics. We leave it to the particle physics and theoretical communities to engineer a fundamental mechanism—whether rooted in environmental screening, field-baryon interactions, or quantum vacuum geometries—capable of reproducing the strict observational constraints demonstrated here.

DATA AVAILABILITY

The SPARC data underlying this article are publicly available from the SPARC database (Lelli et al. 2016).

REFERENCES

Burkert A., 1995, The Astrophysical Journal Letters, 447, L25
Famaey B., McGaugh S., 2013, in Journal of Physics: Conference Series. Institute of Physics Publishing, doi:10.1088/1742-6596/437/1/012001
Hoekstra H., Van Albada T. S., Sancisi R., 2001, Monthly Notices of the Royal Astronomical Society, 323, 453
Kroupa P., et al., 2023, The Many Tensions with Dark-Matter Based Models and Implications on the Nature of the Universe (arXiv:2309.11552), doi:10.48550/arXiv.2309.11552, http://arxiv.org/abs/2309.11552
Lelli F., McGaugh S. S., Schombert J. M., 2016, The Astronomical Journal, 152, 157
Lelli F., McGaugh S. S., Schombert J. M., Pawlowski M. S., 2017, The Astrophysical Journal, 836, 152
Li P., Lelli F., McGaugh S., Schombert J., 2018, Astronomy & Astrophysics, 615, A3
Navarro J. F., Frenk C. S., White S. D. M., 1997, The Astrophysical Journal, 490, 493
Saintonge A., Catinella B., 2022, Annual Review of Astronomy and Astrophysics, 60, 319
Sancisi R., 2004, Symposium - International Astronomical Union, 220, 233
Schombert J., McGaugh S., Lelli F., 2022, The Astronomical Journal, 163, 154

This paper has been typeset from a TEX/LATEX file prepared by the author.