

The sample size required for each State for various investigations among different target groups of individuals are given below:

Investigations	No. of HHs	Age/Sex/ Physiological Group	Prevalence	C.I*	Relative Precision	Sample size per State
Anthropometry	2400	All the available individuals				
Clinical Examination						
History of morbidity						
Diet survey	1200	All the individuals partaking meals in the HH				
Blood Pressure #	2400	Men ( $\geq 18$ yrs)	10%	95%	20%	1728
		Women ( $\geq 18$ yrs)				1728
Fasting blood glucose	2400	Men ( $\geq 18$ yrs)	5%	95%	20%	1825
		Women ( $\geq 18$ yrs)				1825
Knowledge & Practices on HTN & DM	2400	Men ( $\geq 18$ yrs)	-	-	-	1728
	2400	Women ( $\geq 18$ yrs)	-	-	-	1728

\*CI: Confidence Interval      # Design effect 2

### 3.4.2. Measurement of blood pressure

Earlier studies have revealed that the prevalence of hypertension among the rural adults of  $\geq 18$  years in the State of Gujarat was 10%<sup>3</sup> which was lowest compared to other NNMB States. Assuming 10% prevalence of hypertension among rural adults, with 95% confidence interval, 20% relative precision and design effect of 2, the sample size computed for each State was 1,728 adults in each gender.

### 3.4.3. Estimation of fasting blood glucose

Earlier studies have revealed that the overall prevalence of diabetes among the rural adults of  $\geq 18$  years was about 5%<sup>3</sup>. Assuming 5% prevalence of diabetes among rural adults, with 95% confidence interval and 20% relative precision, the sample size computed was 1,825 adults in each gender.

## 3.5. Investigations

The following investigations were carried out in the selected HHs:

### 3.5.1. Socio-economic and demographic particulars of households

Socioeconomic and demographic particulars, such as age, sex, occupation, annual family income, family size, type of family, literacy level of individuals, information about possession of agricultural land, types of crops raised, their yield during previous year, live stock, type of dwelling, environmental sanitation etc. from all the selected 20 HHs were collected using pre-tested questionnaires.