

MATH1131 Mathematics 1A
MATH1231-41 1B **CALCULUS NOTES**
MATH1141 Higher Mathematics 1A

These Notes form the basis for the calculus strand of MATH1131 and MATH1141. However, not all of the material in these Notes is included in the MATH1131 or MATH1141 calculus syllabuses. A detailed syllabus will be uploaded to Moodle.

In using these Notes, you should remember the following points:

1. Most courses at university present new material at a faster pace than you will have been accustomed to in high school, so it is essential that you start working right from the beginning of the session and continue to work steadily throughout the session. Make every effort to keep up with the lectures and to do problems relevant to the current lectures.
2. These Notes are **not** intended to be a substitute for attending lectures or tutorials. The lectures will expand on the material in the notes and help you to understand it.
3. These Notes may seem to contain a lot of material but not all of this material is equally important. One aim of the lectures will be to give you a clearer idea of the relative importance of the topics covered in the Notes.
4. Use the tutorials for the purpose for which they are intended, that is, to ask questions about both the theory and the problems being covered in the current lectures.
5. Some of the material in these Notes is more difficult than the rest. This extra material is marked with the symbol [H]. Material marked with an [X] is intended for students in MATH1141.
6. Some of the problems are marked [V]. These have a video solution available from Moodle.
7. It is **essential** for you to do **problems** which are given at the end of each chapter. If you find that you do not have time to attempt all of the problems, you should at least attempt a representative selection of them. You will find advice about this on Moodle. You should also work through the Online Tutorials that you will find on Moodle.
8. You will be expected to use the computer algebra package Maple in tests and understand Maple syntax and output for the end of term examination.

Remember that working through a wide range of problems is the key to success in mathematics. However, *solving problems* and *writing mathematics clearly* are two separate skills that need to be developed through practice. We recommend that you keep a workbook to practice *writing* solutions to mathematical problems.

CALCULUS PROBLEM SCHEDULE

Solving problems and writing mathematics clearly are two separate skills that need to be developed through practice. We recommend that you keep a workbook to practice *writing* solutions to mathematical problems. The range of questions suitable for each week will be provided on Moodle along with a suggestion of specific recommended problems to do before your classroom tutorials.

The Online Tutorials will develop your problem solving skills, and give you examples of mathematical writing. Online Tutorials help build your understanding from lectures towards solving problems on your own.

	v
1 Sets, inequalities and functions	1
1.1 Sets of numbers	1
1.2 Solving inequalities	4
1.3 Absolute values	7
1.4 Functions	9
1.5 Polynomials and rational functions	13
1.6 The trigonometric functions	14
1.7 The elementary functions	17
1.8 Implicitly defined functions	18
1.9 Continuous functions	21
1.10 Maple notes	22
Problems for Chapter 1	24
2 Limits	27
2.1 Limits of functions at infinity	27
2.1.1 Basic rules for limits	27
2.1.2 The pinching theorem	29
2.1.3 Limits of the form $f(x)/g(x)$	30
2.1.4 Limits of the form $\sqrt{f(x)} - \sqrt{g(x)}$	30
2.1.5 Indeterminate forms	31
2.2 The definition of $\lim_{x \rightarrow \infty} f(x)$	31
2.3 Proving that $\lim_{x \rightarrow \infty} f(x) = L$ using the limit definition	34
2.4 Proofs of basic limit results (MATH1141 only)	37
2.5 Limits of functions at a point	38
2.5.1 Left-hand, right-hand and two-sided limits	38
2.5.2 Limits and continuous functions	40
2.5.3 Rules for limits at a point	42
2.6 Maple notes	46
Problems for Chapter 2	47
3 Properties of continuous functions	51
3.1 Combining continuous functions	51
3.2 Continuity on intervals	53
3.3 The intermediate value theorem	54
3.4 The maximum-minimum theorem	56
Problems for Chapter 3	60
4 Differentiable functions	63
4.1 Gradients of tangents and derivatives	65
4.2 Rules for differentiation	69
4.3 Proofs of results in Section 4.2	71
4.4 Implicit differentiation	72
4.5 Differentiation, continuity and split functions	74
4.6 Derivatives and function approximation	77
4.7 Derivatives and rates of change	78
4.8 Local maximum, local minimum and stationary points	79
4.9 Maple notes	81

Problems for Chapter 4	82
5 The mean value theorem and applications	85
5.1 The mean value theorem	85
5.2 Proof of the mean value theorem	87
5.3 Proving inequalities using the mean value theorem	88
5.4 Error bounds	90
5.5 The sign of a derivative	91
5.6 The second derivative and applications	93
5.7 Critical points, maxima and minima	96
5.8 Counting zeros	98
5.9 Antiderivatives	101
5.10 L'Hôpital's rule	103
Problems for Chapter 5	107
6 Inverse functions	111
6.1 Some preliminary examples	111
6.2 One-to-one functions	113
6.3 Inverse functions	115
6.4 The inverse function theorem	119
6.5 Applications to the trigonometric functions	120
6.6 Maple notes	125
Problems for Chapter 6	127
7 Curve sketching	131
7.1 Curves defined by a Cartesian equation	131
7.1.1 A checklist for sketching curves	132
7.1.2 Oblique asymptotes	132
7.1.3 Examples	135
7.2 Parametrically defined curves	136
7.2.1 Parametrisation of conic sections	139
7.2.2 Calculus and parametric curves	141
7.2.3 The cycloid and curve of fastest descent	142
7.3 Curves defined by polar coordinates	144
7.3.1 Polar coordinates	144
7.3.2 Basic sketches of polar curves	146
7.3.3 Sketching polar curves using calculus	148
7.4 Maple notes	151
Problems for Chapter 7	153
8 Integration	157
8.1 Area and the Riemann Integral	157
8.1.1 Area of regions with curved boundaries	157
8.1.2 Approximations of area using Riemann sums	160
8.1.3 The definition of area under the graph of a function and the Riemann integral	163
8.2 Integration using Riemann sums	165
8.3 The Riemann integral and signed area	168
8.4 Basic properties of the Riemann integral	169
8.5 The first fundamental theorem of calculus	171
8.6 The second fundamental theorem of calculus	175
8.7 Indefinite integrals	178
8.8 Integration by substitution	180

8.9	Integration by parts	185
8.10	Improper integrals	186
8.11	Comparison tests for improper integrals	189
8.12	Functions defined by an integral	196
8.13	Maple notes	199
	Problems for Chapter 8	200
9	The logarithmic and exponential functions	207
9.1	Powers and logarithms	208
9.2	The natural logarithm function	209
9.3	The exponential function	212
9.4	Exponentials and logarithms with other bases	214
9.5	Integration and the $\ln$ function	215
9.6	Logarithmic differentiation	216
9.7	Indeterminate forms with powers	217
	Problems for Chapter 9	219
10	The hyperbolic functions	223
10.1	Hyperbolic sine and cosine functions	224
10.2	Other hyperbolic functions	227
10.3	Hyperbolic identities	228
10.4	Hyperbolic derivatives and integrals	230
10.5	The inverse hyperbolic functions	231
10.6	Integration leading to the inverse hyperbolic functions	234
10.7	A summary of important hyperbolic formulae	236
10.8	(Appendix):The catenary	237
10.9	Maple notes	240
	Problems for Chapter 10	241
	Answers to selected problems	243
	Chapter 1	243
	Index	252
1	Functions of several variables	1
1.1	Sketching simple surfaces in $\mathbb{R}^3$	1
1.2	Partial differentiation	4
1.3	Tangent planes to surfaces	10
1.4	The total differential approximation	13
1.5	Chain rules	16
1.6	Functions of more than two variables	19
1.7	Maple notes	20
	Problems for Chapter 1	21
2	Integration techniques	25
2.1	Trigonometric integrals	25
2.1.1	Integrating powers of sine and cosine	25
2.1.2	Integrating multiple angles of sine and cosine	28
2.1.3	Integrating powers of tan and sec	29
2.2	Reduction formulae	29
2.2.1	[X] Application: the irrationality of π	32
2.3	Trigonometric and hyperbolic substitutions	35
2.4	Integrating rational functions	37
2.4.1	The overall strategy	38
2.4.2	Partial fractions decompositions	40
2.4.3	Integrating rational functions: two examples	42
2.5	Other substitutions	44
2.6	Maple notes	47

Problems for Chapter 2	48
3 Ordinary differential equations	55
3.1 An introduction	56
3.2 Initial value problems	58
3.3 Separable ODEs	60
3.4 First order linear ODEs	62
3.5 Exact ODEs	65
3.6 Solving ODEs by using a change of variable $[\mathbf{X}]$	69
3.7 Modelling with first order ODEs	71
3.7.1 Mixing problems	71
3.7.2 Population models	73
3.8 Second order linear ODEs with constant coefficients	76
3.8.1 The homogeneous case	77
3.8.2 The non-homogeneous case	80
3.8.3 An application: vibrations and resonance	84
3.8.4 A connection with linear algebra	87
3.9 Maple notes	88
Problems for Chapter 3	90
4 Taylor series	101
4.1 Taylor polynomials	101
4.2 Taylor's theorem	106
4.2.1 Classifying stationary points	112
4.2.2 Some questions arising from Taylor's theorem	113
4.3 Sequences	114
4.3.1 Describing the limiting behaviour of sequences	115
4.3.2 Techniques for calculating limits of sequences	117
4.3.3 Suprema and infima $[\mathbf{X}]$	120
4.4 Infinite series	122
4.5 Tests for series convergence	126
4.5.1 Some preliminary results on series summation	126
4.5.2 The k th term divergence test	126
4.5.3 The integral test	127
4.5.4 The comparison test	128
4.5.5 $[\mathbf{X}]$ The limit form of the comparison test	130
4.5.6 The ratio test	131
4.5.7 Leibniz' test for alternating series	132
4.5.8 Absolute and conditional convergence	136
4.6 Taylor series	138
4.7 Power series	142
4.7.1 Radius of Convergence	144
4.7.2 Convergence of power series at endpoints $[\mathbf{X}]$	146
4.8 Manipulation of power series	146
4.8.1 Proof of theorems in Section 4.8 $[\mathbf{X}]$	151
4.9 Maple notes	154
Problems for Chapter 4	155
5 Averages, arc length, speed and surface area	167
5.1 The average value of a function	167
5.2 The arc length of a curve	171
5.2.1 An intuitive derivation of the arc length formula	172
5.2.2 Arc length for a parametrised curve	174
5.2.3 Arc length for the graph of a function	175
5.2.4 Arc length for a polar curve	177
5.3 The speed of a moving particle	178
5.4 Surface area	179
5.4.1 An heuristic derivation for the surface area of a surface of revolution	179
5.4.2 Surface area formulae and examples	182
Problems for Chapter 5	187

Chapter 1 Sets, inequalities and functions

In the days of the Roman empire, the word ‘calculus’ denoted a pebble that was used for counting and gambling. As time progressed, the word ‘calcularē’ came to mean ‘to compute.’ In the second half of the seventeenth century, two mathematicians, the Englishman Isaac Newton and the German Gottfried Leibniz, independently invented methods for

- calculating gradients of tangents to curves,
- calculating instantaneous acceleration and velocity,
- calculating the area of regions with a curved boundary,
- calculating the volume of solids with curved boundary,
- calculating the length of a curve,
- calculating the work done by a force, and
- calculating the centre of mass of a general solid.

These methods were developed by combining algebra, geometry and trigonometry with the *limiting process* and became known as *the calculus*.

Calculus (as it is known today) has many applications to engineering, physics, chemistry, biology, geology, surveying, sociology, economics and statistics. It includes two major branches: the *differential calculus* (introduced in Chapter 4) and the *integral calculus* (introduced in Chapter 8). These two branches are related by *the fundamental theorem of calculus* (see Theorem 8.5.1). The underlying tool used to develop these branches is the concept of the *limit* (introduced in Chapter 2), which is applied to *functions* (introduced in Chapter 1).

The main goal of this chapter is to introduce functions. Although functions will be familiar to students from high school, important points that deserve students’ attention are emphasised here. Since the functions we study in this course take real numbers as inputs and give real numbers of outputs, we begin the chapter with a review of different sets that consist of real numbers. We also devote a little time to revising inequalities and absolute values, in preparation for our discussion of limits in Chapter 2.

1.1 Sets of numbers

(Ref: SH10 §1.2)

A *set* is a collection of distinct objects. The objects in a set are called the *elements* or *members* of the set. Some commonly used sets of numbers are listed below.


- The set $\mathbb{N}$ of *natural numbers* is given by

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}.$$

- The set $\mathbb{Z}$ of *integers* is given by

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}.$$

- The set $\mathbb{Q}$ of *rational numbers* is the collection of all numbers of the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$.
- The set $\mathbb{R}$ of *real numbers* may be represented as the collection of points lying on the number line.



If A is a set of numbers and the number x is a member of the set A , then we write

$$x \in A.$$

If x is not a member of A then we write

$$x \notin A.$$

The next example illustrates this notation.

Example 1.1.1. The following statements are true:

$$2 \in \mathbb{N}, \quad -2 \notin \mathbb{N}, \quad \frac{3}{4} \notin \mathbb{Z}, \quad \frac{3}{4} \in \mathbb{Q}, \quad \sqrt{2} \in \mathbb{R}, \quad \sqrt{2} \notin \mathbb{Q}.$$

Many other sets of numbers can be described using $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$ and inequalities.

Example 1.1.2. (a) The set


$$\{n \in \mathbb{Z} : n > 0\}$$

is the set of all numbers n belonging to the integers such that n is greater than 0. In other words, it is the set of *positive integers*. (Note that the colon ‘:’ above is read as ‘such that’.)

(b) The set

$$\{x \in \mathbb{R} : -2 \leq x < 3\}$$

is represented by the following interval on the real line.




(c) The set

$$\{x \in \mathbb{R} : x \notin \mathbb{Q}\}$$


is the set of *irrational numbers*.

Sets that are represented by intervals of the real line occur so frequently that we introduce a special notation for them. Suppose that a and b are real numbers and that $a < b$. Then the intervals (a, b) , $[a, b]$, $(a, b]$ and $[a, b)$ are given by


- $(a, b) = \{x \in \mathbb{R} : a < x < b\}$




- $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$



- $[a, b) = \{x \in \mathbb{R} : a \leq x < b\}$




- $(a, b] = \{x \in \mathbb{R} : a < x \leq b\}$




In each case, the numbers a and b are called the *endpoints* of the interval. An interval $[a, b]$ that includes its endpoints is called a *closed interval*, while an interval (a, b) that excludes its endpoints is called an *open interval*. The intervals $[a, b)$ and $(a, b]$ are neither open nor closed.

Interval notation can be extended to describe rays of the real line by using the symbol ∞ for infinity. Thus we have

- $[a, \infty) = \{x \in \mathbb{R} : a \leq x\}$



- $(-\infty, b) = \{x \in \mathbb{R} : x < b\}$



- $(-\infty, \infty) = \mathbb{R}$.

We end by introducing an important set relation.

Definition 1.1.3. Suppose that A and B are two sets. We say that A is a *subset* of B if $x \in A$ implies that $x \in B$. If A is a subset of B then we also say that B *contains* the set A .

The above definition is illustrated by the following examples:

- $\{0, 1, 2, 3\}$ is a subset of $\{0, 1, 2, 3, 4\}$,
- $\mathbb{N}$ is a subset of $\mathbb{Z}$,
- $[-1, 2]$ is a subset of $(-3, \infty)$,
- $\mathbb{R}$ is a subset of $\mathbb{R}$, and
- $[1, 4)$ is not a subset of $(1, 4]$.

1.2 Solving inequalities

(Ref: SH10 §1.3)

When solving an inequality, one must remember that if the inequality is multiplied by a negative number, the sign of the inequality reverses.

Two types of inequalities deserve special attention: quadratic inequalities and inequalities with an unknown quantity in the denominator.

Example 1.2.1. Solve the inequality

$$x^2 > 6 - x.$$

Solution. Rearranging the inequality gives

$$x^2 + x - 6 > 0.$$

The values of x for which $x^2 + x - 6$ is positive are the same as those for which the graph of

$$y = x^2 + x - 6$$

lies above the x -axis.


To graph

$$y = x^2 + x - 6,$$

we factorise the right-hand side to obtain

$$y = (x - 2)(x + 3).$$

Hence $y = 0$ when $x = 2$ and when $x = -3$. We can now easily sketch its graph.



Since the graph lies *above* the x -axis when $x < -3$ and when $x > 2$, the solution to the inequality is

$$x < -3 \quad \text{or} \quad x > 2.$$

□

Example 1.2.2. Solve the inequality

$$\frac{1}{3} < \frac{1}{x-1}.$$

Solution. We begin by noting that $x \neq 1$.

To solve the inequality, we cannot simply multiply the inequality through by $x - 1$, since we do not know whether $x - 1$ is positive or negative. (Taking the reciprocal of both sides is similarly problematic.) Instead, we multiply through by the *positive* number $(x - 1)^2$ to obtain

$$\frac{1}{3}(x-1)^2 < x-1.$$

Multiplication by 3 and rearrangement gives

$$(x-1)^2 - 3(x-1) < 0$$


and by factorisation we obtain

$$(x-1)[(x-1)-3] = (x-1)(x-4) < 0.$$

So proceeding in a manner similar to the previous example, we seek to identify the values of x for which the graph of

$$y = (x-1)(x-4)$$

lies *below* the x -axis.



Hence the solution to the inequality is given by

$$1 < x < 4.$$

□

The next example is slightly more difficult, but uses techniques illustrated in the previous two examples.

Example 1.2.3. Solve the inequality

$$x + 2 \geq \frac{4}{x - 1}$$

Solution. First observe that $x \neq 1$.

Multiplying the inequality through by $(x - 1)^2$ gives

$$(x - 1)^2(x + 2) \geq 4(x - 1).$$

Hence

$$(x - 1)^2(x + 2) - 4(x - 1) \geq 0.$$

Now

$$\begin{aligned} (x - 1)^2(x + 2) - 4(x - 1) &= (x - 1)[(x - 1)(x + 2) - 4] \\ &= (x - 1)[x^2 + x - 6] \\ &= (x - 1)(x - 2)(x + 3). \end{aligned}$$


So the original inequality is equivalent to

$$(x - 1)(x - 2)(x + 3) \geq 0, \quad x \neq 1.$$

We now sketch the graph of

$$y = (x - 1)(x - 2)(x + 3)$$

and see for which values of x the graph lies on or above the x -axis.



Hence the solution to the inequality is

$$-3 \leq x < 1 \quad \text{or} \quad x \geq 2.$$

□

Remark 1.2.4. A variety of methods (some good and others bad) exist for solving inequalities where an unknown lies in the denominator. It is strongly recommended that students use the method shown in this section, or that shown by their first year calculus lecturer.

1.3 Absolute values

(Ref: SH10 §§1.2, 1.3)

The magnitude, or *absolute value*, of a real number x is denoted by $|x|$ and defined by

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases} \quad (1.1)$$

Absolute values of real numbers interact in the following way.

Proposition 1.3.1. *Suppose that x and y are real numbers. Then*

- (i) $|-x| = |x|$,
- (ii) $|xy| = |x||y|$,
- (iii) $\left|\frac{x}{y}\right| = \frac{|x|}{|y|}$, provided that $y \neq 0$, and
- (iv) $|x + y| \leq |x| + |y|$.

Property (iv) is known as *the triangle inequality*.

Three other useful facts about absolute values are noted below.

(F1) For every real number x ,

$$|x| = \sqrt{x^2} \quad \text{and} \quad |x|^2 = x^2. \quad (1.2)$$

(F2) If a and x are real numbers then $|x - a|$ is equal to the distance between x and a on the real number line.

(F3) For any positive real number a ,

$$|y| < a \quad \text{if and only if} \quad -a < y < a$$

and

$$|y| > a \quad \text{if and only if} \quad y < -a \quad \text{or} \quad y > a.$$

These facts give different methods for solving inequalities involving absolute values.

Example 1.3.2. Solve the following inequalities:

- (a) $|x - 3| \leq 5$
- (b) $|2x + 5| > 8$
- (c) $\left|\frac{x - 3}{x - 1}\right| < 1.$


Solution. (a) *An algebraic solution.* Using (F3) we have

$$\begin{aligned} |x - 3| &\leq 5 \\ -5 &\leq x - 3 \leq 5 \\ -2 &\leq x \leq 8. \end{aligned}$$

(a) *A geometric solution.* By (F2) the inequality $|x - 3| \leq 5$ is equivalent to saying that

‘the distance between x and 3 is less than or equal to 5.’

The points x on the number line that have this property are shown below.



Hence $-2 \leq x \leq 8$.

(b) *An algebraic solution.* Using (F3) the inequality is equivalent to

$$2x + 5 < -8 \quad \text{or} \quad 2x + 5 > 8.$$

Hence

$$2x < -13 \quad \text{or} \quad 2x > 3,$$

which gives the solution

$$x < -\frac{13}{2} \quad \text{or} \quad x > \frac{3}{2}.$$

(c) *An algebraic solution.* By Prop.1.3.1 (iii), we can write, for $x \neq 1$,

$$\left| \frac{x-3}{x-1} \right| = \frac{|x-3|}{|x-1|}$$

and since the denominator is positive, the inequality can be written as

$$|x-3| < |x-1|.$$

Squaring both (positive) sides), and using (F1), we have

$$(x-3)^2 < (x-1)^2.$$

Expanding and solving, we have $x > 2$.


(c) *A geometric solution.* As above, the inequality is equivalent to

$$|x-3| < |x-1|, \quad x \neq 1.$$

By (F2) the inequality is interpreted geometrically as

‘the distance from x to 3 is less than the distance from x to 1.’

In other words, x is closer to 3 than to 1. Points that have this property are shown on the numberline below.



Hence $x > 2$. □

1.4 Functions

(Ref: SH10 §1.5)

A function $f : A \rightarrow B$ is a rule which assigns to every element x belonging to a set A exactly one element $f(x)$ belonging to a set B . The set A is called the *domain* of the function f and the set B is called the *codomain* of f . In this course, A and B are always sets of real numbers.

Example 1.4.1. A function $f : [0, \infty) \rightarrow \mathbb{R}$ is given by

$$f(x) = \sqrt{x}$$

for all x in $[0, \infty)$. Informally, this means that f takes a number x from $[0, \infty)$ as input and gives the number $\sqrt{x}$ as output. More formally, we say that f *maps* an element x of $[0, \infty)$ to $\sqrt{x}$. We offer a few comments on terminology employed to describe f .

- The expression $f : [0, \infty) \rightarrow \mathbb{R}$ says that $[0, \infty)$ is the domain of f and that $\mathbb{R}$ is the codomain of f . We write

$$\text{Dom}(f) = [0, \infty) \quad \text{and} \quad \text{Codom}(f) = \mathbb{R}.$$

- The domain of f is the set of all inputs for the function.
- The codomain of f is a set that contains all the output values of the function. Since the output values are all real numbers, we say that f is a *real-valued function*. While all the outputs must lie in $\text{Codom}(f)$, not every number in $\text{Codom}(f)$ need be an output value.
- The expression $f(x)$ (read as ‘ f of x ’) is the *value of f* at the point x . That is, $f(x)$ is the unique number in $\mathbb{R}$ that corresponds to the input x . To emphasise the point: $f(x)$ is a real number, *not* a function.
- The statement

$$‘f(x) = \sqrt{x} \text{ for all } x \text{ in } [0, \infty)’$$

can be abbreviated as

$$f(x) = \sqrt{x} \quad \forall x \in [0, \infty).$$

The symbol $\forall$ means ‘for all.’

In high school, functions are often described by specifying a rule without specifying a domain. For example, the following sentence was taken from an HSC study guide:


‘Consider the function $f(x) = x^2$.’

There are two problems with this sentence. First, neither $f(x)$ nor $f(x) = x^2$ is a function. As mentioned above, $f(x)$ is a real number, and $f(x) = x^2$ is an equation. (This may seem picky, but being able to distinguish between a function f and its value $f(x)$ at the point x is crucial to understanding key concepts in both algebra and calculus.) Second, the quoted sentence does not specify the domain of the function. The next example gives two different functions that obey the same squaring rule.

Example 1.4.2. Consider the functions f and g , defined by

$$\begin{array}{ll} f : \mathbb{R} \rightarrow \mathbb{R} & g : [0, \infty) \rightarrow \mathbb{R} \\ f(x) = x^2 \quad \forall x \in \text{Dom}(f) & \text{and} \quad g(x) = x^2 \quad \forall x \in \text{Dom}(g). \end{array}$$

Even though they are both squaring functions, f and g are *not* the same function. For example, f can square negative numbers while g does not. On the other hand, g is *invertible* (this important and desirable property is discussed in Chapter 6) while f is not. Their graphs, shown below, are also noticeably different.



If, for whatever reason, the domain of a function is not specified, but the function rule is, then the default domain, known as the *maximal* or *natural domain*, is the largest possible domain for which the rule makes sense. We give two examples.

- If

$$f(x) = \sqrt{x-1}$$

then the rule makes sense only if $x-1 \geq 0$, that is, if $x \geq 1$. Therefore the maximal domain of the function f is $[1, \infty)$.

- If

$$f(x) = \frac{2}{(x-2)(x-8)}$$

then the rule only makes sense if $x \neq 2$ or $x \neq 8$. So the maximal domain is the set

$$\{x \in \mathbb{R} : x \neq 2, x \neq 8\}.$$

Another set associated to a function is its *range*.

Definition 1.4.3. Suppose that $f : A \rightarrow B$ is a function. The *range* of f , denoted by $\text{Range}(f)$, is defined by


$$\text{Range}(f) = \{f(x) \in B : x \in A\}.$$

In other words, the range of f is the set of all output values. Equivalently, if the rule of a function is described by the equation $y = f(x)$, then the range of f is the set of all corresponding y -values. The range is always a subset of the codomain, but the range need not equal the codomain. The difference between range and codomain is illustrated by the following examples.

Example 1.4.4. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by the rule

$$f(x) = \sin x \quad \forall x \in \mathbb{R}.$$

Then $\text{Dom}(f) = \mathbb{R}$ and $\text{Codom}(f) = \mathbb{R}$ (this is obtained from the expression $f : \mathbb{R} \rightarrow \mathbb{R}$). To determine the range, consider the graph of f is sketched below.




The set of all output values (equivalently, the set of all corresponding y -values) is $[-1, 1]$. Hence $\text{Range}(f) = [-1, 1]$.

Example 1.4.5. Suppose that $f : [-1, 1] \rightarrow \mathbb{R}$ is given by the rule

$$f(x) = \sqrt{1 - x^2} \quad \forall x \in [-1, 1].$$

Then $\text{Dom}(f) = [-1, 1]$ and $\text{Codom}(f) = \mathbb{R}$ (again, we are simply reading this from $f : [-1, 1] \rightarrow \mathbb{R}$). To determine the range, we observe that the graph of f is a semicircle.



The set of all outputs (or y -values) is $[0, 1]$. Hence $\text{Range}(f) = [0, 1]$.

If f and g are two functions with the same domain, then one can combine f and g to form new functions.

Definition 1.4.6. Suppose that $f : A \rightarrow B$ and $g : A \rightarrow B$ are real-valued functions. Then the functions $f + g$, $f - g$, $f \cdot g$ and f/g are defined by the rules

$$(f + g)(x) = f(x) + g(x) \quad \forall x \in A$$

$$(f - g)(x) = f(x) - g(x) \quad \forall x \in A$$

$$(f \cdot g)(x) = f(x)g(x) \quad \forall x \in A$$

$$(f/g)(x) = \frac{f(x)}{g(x)} \quad \forall x \in A, \text{ provided that } g(x) \neq 0.$$

Another way of constructing new functions is given below.

Definition 1.4.7. Suppose that $f : C \rightarrow D$ and $g : A \rightarrow B$ are functions such that $\text{Range}(g)$ is a subset of $\text{Dom}(f)$. Then the *composition* $f \circ g : A \rightarrow D$ is defined by the rule

$$(f \circ g)(x) = f(g(x)) \quad \forall x \in A.$$

Example 1.4.8. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are given by the rules

$$f(x) = x^2 + 1 \quad \forall x \in \mathbb{R} \quad \text{and} \quad g(x) = x - 2 \quad \forall x \in \mathbb{R}.$$

Find $(f + g)(5)$ and $(f \circ g)(x)$. Is 2 in the domain of the function f/g ? Give reasons.

Solution. First,

$$\begin{aligned} (f + g)(5) &= f(5) + g(5) \\ &= (5^2 + 1) + (5 - 2) \\ &= 29. \end{aligned}$$

Second, if x in $\mathbb{R}$ then

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) \\ &= (x - 2)^2 + 1 \\ &= x^2 - 4x + 5. \end{aligned}$$

Finally, $g(2) = 0$ so 2 is not in the domain of f/g . □

The final example illustrates the importance of bearing in mind domain and range when composing two functions.

Example 1.4.9. Suppose that $f : [0, \infty) \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are given by

$$f(x) = \sqrt{x} \quad \forall x \in [0, \infty) \quad \text{and} \quad g(x) = \sin x - 2 \quad \forall x \in \mathbb{R}.$$

Find, if they exist, $(f \circ g)(x)$ and $(g \circ f)(x)$.

Solution. Consider first $f \circ g$. For this function to exist, we require that $\text{Range}(g)$ is a subset of $\text{Dom}(f)$. Now

$$-1 \leq \sin x \leq 1 \quad \forall x \in \mathbb{R}$$

so

$$-3 \leq \sin x - 2 \leq -1 \quad \forall x \in \mathbb{R};$$

that is, $\text{Range}(g) = [-3, -1]$. Since $\text{Dom}(f) = [0, \infty)$, $\text{Range}(g)$ is not a subset of $\text{Dom}(f)$ and hence $f \circ g$ does not exist. (The point here is that, if

$$(f \circ g)(x) = f(g(x)) = \sqrt{\sin x - 2},$$

then the expression under the square root sign is always negative.)

Now consider $g \circ f$. Since f is a real-valued function and $\text{Dom}(g) = \mathbb{R}$, it is clear that $\text{Range}(f)$ is a subset of $\text{Dom}(g)$. (The fact that $\text{Range}(f) = [0, \infty)$ is not needed because $\text{Dom}(g)$ is so large.) Hence

$$(g \circ f)(x) = g(f(x)) = \sin(\sqrt{x}) - 2 \quad \forall x \in \text{Dom}(f) = [0, \infty).$$

□

1.5 Polynomials and rational functions

(Ref: SH10 §1.6)

In the next three sections we discuss some important classes of functions. Since these were introduced in high school, we give a brief survey only.

Definition 1.5.1. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called a *polynomial* if





$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0 \quad \forall x \in \mathbb{R},$$

where n is a natural number, the constants $a_0, a_1, \dots, a_n$ are real numbers and $a_n \neq 0$. We call n the *degree* of f , and the constants $a_0, a_1, \dots, a_n$ are called the *coefficients* of f . The constant a_n is called the *leading coefficient* of f .

Polynomials of small degree have special names and properties.

- If $n = 0$ then $f(x) = a_0$. Hence f is a constant function and the graph of f is a horizontal straight line.
- If $n = 1$ then $f(x) = a_1 x + a_0$. The graph of f is a straight line with gradient a_1 and y -axis intercept a_0 .
- If $n = 2$ then f is called a *quadratic function* and its graph is a parabola.
- If n is 3, 4 or 5 then f is called a *cubic*, *quartic* or *quintic*, polynomial respectively.

The general shape of the graph of a polynomial function of degree greater than 1 is indicated by the table below.

	Positive leading coefficient	Negative leading coefficient
Odd degree		
Even degree		

The number of turning points in the graph is always strictly less than the degree of the polynomial. Exactly how many turning points the graph has depends on the coefficients of the polynomial, and can be determined using calculus (see, for example, Chapters 4 and 5).

By dividing two polynomials, one obtains a new type of function.

Definition 1.5.2. Suppose that p and q are polynomials. A function f is called a *rational function* if

$$\text{Dom}(f) = \{x \in \mathbb{R} : q(x) \neq 0\}$$

and

$$f(x) = \frac{p(x)}{q(x)}, \quad \forall x \in \text{Dom}(f).$$

Hence the expressions


$$f(x) = \frac{x}{x^2 + 2} \quad \text{and} \quad f(x) = \frac{x^3 - 4x^2 + 1}{x - 2}$$

both give rise to rational functions, as does

$$f(x) = x - 1 + \frac{3}{x^2 + 3}$$

(to see why, rewrite right-hand side with a common denominator of $x^2 + 3$).

The graphs of two simple rational functions are shown below.



The graphs of more sophisticated rational functions are studied in Chapter 7.

1.6 The trigonometric functions

(Ref: SH10 §1.6)

The trigonometric functions will already be familiar to you. This section gives a brief summary of their definitions and some of their properties.


Remark 1.6.1. In this course, angles are always measured in *radians* rather than degrees. (Radian measure is a more natural way of measuring angles; moreover, important formulae used in calculus, such as

$$\frac{d}{d\theta}(\sin \theta) = \cos \theta,$$

only make sense geometrically when θ is interpreted as an angle measured in radians rather than degrees.) We remind readers that

$$2\pi \text{ radians} = 360 \text{ degrees}.$$

Consider an angle θ and the corresponding point $P(x, y)$ that lies on the unit circle centred at the origin, as shown below.




The functions $\cos : \mathbb{R} \rightarrow \mathbb{R}$ and $\sin : \mathbb{R} \rightarrow \mathbb{R}$ are defined by


$$\cos \theta = x \quad \text{and} \quad \sin \theta = y.$$

Other trig functions are defined in terms of the sine and cosine functions:

$$\begin{aligned} \tan \theta &= \frac{\sin \theta}{\cos \theta}, & \text{provided that } \cos \theta \neq 0 \\ \sec \theta &= \frac{1}{\cos \theta}, & \text{provided that } \cos \theta \neq 0 \\ \operatorname{cosec} \theta &= \frac{1}{\sin \theta}, & \text{provided that } \sin \theta \neq 0 \\ \cot \theta &= \frac{\cos \theta}{\sin \theta}, & \text{provided that } \sin \theta \neq 0. \end{aligned}$$

The graphs of $\sin$, $\cos$ and $\tan$ are shown below.





The sine and cosine functions are 2π -periodic, while the tangent function is π -periodic, which means that


$$\sin(\theta + 2\pi) = \sin \theta, \quad \cos(\theta + 2\pi) = \cos \theta \quad \text{and} \quad \tan(\theta + \pi) = \tan \theta$$

whenever θ is in the domain of the respective function. Sine and cosine both have an amplitude of 1.

It is not hard to obtain functions whose graphs have the same general shape as the sine (or cosine) curve with a period other than 2π and amplitude other than 1. For example, if A and ω are positive real numbers then the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = A \sin(\omega x) \quad \forall x \in \mathbb{R},$$

has an amplitude of A and a period of $\frac{2\pi}{\omega}$. Its graph is shown below.



The six trigonometric functions are related to one another by various identities and formulae:

- complementary identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x$$

$$\cos\left(\frac{\pi}{2} - x\right) = \sin x$$

- Pythagorean identities

$$\cos^2 x + \sin^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$\cot^2 x + 1 = \operatorname{cosec}^2 x$$

- the sum and difference formulae

$$\sin(x \pm y) = \sin x \cos y \pm \cos x \sin y$$

$$\cos(x \pm y) = \cos x \cos y \mp \sin x \sin y$$

$$\tan(x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y}$$

- double-angle formulae

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

$$\tan(2x) = \frac{2 \tan x}{1 - \tan^2 x}.$$


The above formulae, especially the double-angle formulae for $\sin(2x)$ and $\cos(2x)$, should be learnt carefully.

The inverse trigonometric functions will be discussed in Chapter 6.

1.7 The elementary functions


(Ref: SH10 §1.6)

Other important functions that have not yet been mentioned include the exponential and logarithm functions. Their graphs are shown below.



These functions will be studied in greater depth in Chapter 9.

Functions arising from the study of roots of polynomials are also important to mathematics, especially those functions f of the form $f(x) = \sqrt[n]{x}$, where n is a positive integer. It is important that students are familiar with the graphs of the square root and cube root functions, shown below.



The *elementary functions* are all those functions that can be constructed by combining a finite number of polynomials, exponentials, logarithms, roots and trigonometric functions (including the inverse trigonometric functions) via function composition, addition, subtraction, multiplication and division. Hence the following expressions give rise to elementary functions:

$$\begin{aligned}f(x) &= e^{\sin x} + x^2 \\g(x) &= \frac{\ln x - \tan x}{\sqrt{x}} \\h(x) &= \sqrt[3]{x^4 - 2x^2 + 5}.\end{aligned}$$

It also follows that every rational function is an elementary function.


The absolute values function, given by

$$\begin{aligned}\text{abs} : \mathbb{R} &\rightarrow \mathbb{R} \\ \text{abs}(x) &= |x| \quad \forall x \in \mathbb{R},\end{aligned}$$

is an elementary function by equation (1.2). It follows that if f is an elementary function then $\text{abs} \circ f$ is also an elementary function. It is standard to denote function $\text{abs} \circ f$ by $|f|$. Thus

$$|f|(x) = (\text{abs} \circ f)(x) = |f(x)| \quad \forall x \in \text{Dom}(f).$$

The graph of $|f|$ is obtained by reflecting in the x -axis any part of the graph of f that lies below the x -axis. This is illustrated below.




Most, if not all functions studied in high school were elementary functions. In this course we shall meet some useful functions that are not elementary functions. (These are defined by integrals; see Section 8.12.)

1.8 Implicitly defined functions

Many curves on the plane can be described as all those points (x, y) on the plane that satisfy some equation involving x and y . For example, consider the equation

$$(x^2 + y^2 - 1)^3 - x^2 y^3 = 0. \quad (1.3)$$

The set of points (x, y) satisfying this equation are shown on the graph below.




Note that the curve cannot be described by one function since some x -values have two corresponding y -values. However, the curve can be described by combining the graphs of two functions $f : [-a, a] \rightarrow \mathbb{R}$ and $g : (-a, a) \rightarrow \mathbb{R}$, which are defined by the rules

$$\begin{cases} y = f(x) \\ (x^2 + y^2 - 1)^3 - x^2 y^3 = 0 \\ y \geq b \end{cases}$$

and

$$\begin{cases} y = g(x) \\ (x^2 + y^2 - 1)^3 - x^2 y^3 = 0 \\ y < b \end{cases}$$

(where $a \approx 1.139$ and $b \approx 0.545$), as illustrated below.



We say that f and g are *implicitly* defined by equation (1.3).

Example 1.8.1. The set of points (x, y) that satisfy the equation


$$x^2 + y^2 = 1$$

describes a circle of radius 1 centred at the origin. The above equation could be used to implicitly define many different functions. We give two examples.

- The function $f : [0, 1] \rightarrow \mathbb{R}$ is defined by the rule

$$\begin{cases} y = f(x) \\ x^2 + y^2 = 1 \\ y \geq 0. \end{cases}$$


Its graph is shown below.




- The function $g : (-1, 1) \rightarrow \mathbb{R}$ is defined by the rule

$$\begin{cases} y = f(x) \\ x^2 + y^2 = 1 \\ y \leq 0. \end{cases}$$


Its graph is shown below.




Many heavenly bodies (such as planets and comets) trace out paths that are roughly elliptic, parabolic or hyperbolic. (The fact that planets trace out elliptic paths around the sun was discovered by Johannes Kepler at the beginning of the sixteenth century, and was later used by Newton, in conjunction with calculus, to establish his universal law of gravitation.) Circles, ellipses, parabolas and hyperbolas are known as *conic sections* because they can be obtained by intersecting a cone with a plane. Equations that describe conic sections, accompanied by corresponding diagrams, are shown below.



Circle: $x^2 + y^2 = r^2$



Ellipse: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$




Students should be able to recognise such curves.

1.9 Continuous functions


The sine, cosine, exponential, logarithmic and polynomial functions are examples of *continuous functions*. Intuitively, a function f is continuous on its domain if small changes in the input variable x produce small changes in $f(x)$ (assuming, of course, that x always remains within $\text{Dom}(f)$).

The intuitive notion that a function is continuous if ‘its graph can be drawn without lifting the pencil off the page’ works well for some functions, but not for others. For example, the tangent function is a continuous function, even though its graph (part of which is shown below) cannot be drawn without lifting the pencil off the page.



The break in the graph when, say, $x = -\pi/2$ or $x = \pi/2$, is a consequence of a break in the domain of the function rather than a discontinuity of the function. Similarly, the trigonometric functions sec, cosec and cot are continuous, as are the rational functions. The notion of continuity shall be discussed in more precise terms in Chapters 2 and 3.

Intuitively, a function f has a *discontinuity* at a point a if $a \in \text{Dom}(f)$ and there is a break in the graph of f at a . It is important to realise that there are different types of discontinuities. We compare three examples.



The functions f , g and h each have a discontinuity at a .

- For f , the discontinuity can be easily removed by redefining the value of $f(a)$ to ‘plug up the hole’ at a . This is an example of a *removable discontinuity*.
- For g , the discontinuity is a result of a finite jump in the graph of g at the point a . The discontinuity for g , which is an example of a *jump discontinuity*, is worse than that for f but not nearly as bad as that for h .
- For h , the discontinuity is a result of rapid oscillation on the right-hand side of a and an infinite jump on the left-hand side of a . This is an example of an *essential discontinuity*.

While students are not expected to learn or use the terms removable, jump and essential discontinuity, it is expected that they will appreciate that some discontinuities are worse than others.


1.10 Maple notes

Many chapters in these notes conclude with a summary of relevant Maple commands. Other sources of information are the First Year Computing Notes and the online Maple **help** command. While Maple can relieve you of the tedium of calculations, it is not a substitute for understanding the mathematics behind the calculations, which is what you need in order to interpret Maple output intelligently.


The following Maple command is relevant to the material of this chapter:

`plot(f(x),x=a..b);` draws a (two-dimensional) plot of the graph of $y = f(x)$ for $a \leq x \leq b$. To find out about the many options available, use `?plot`. Implicitly defined functions can be plotted using the `smartplot` command. For example,


```
> # draw a rational function
> f:=x^5-2*x^4+3*x^3-6*x^2-4*x+8:
> g:=4*x^4-8*x^3-12*x^2+16*x+16:
> plot(f/g, x=-10..10,y=-10..10,numpoints=1000,discont=true);
```



```
> # draw a hyperbola, via implicit definition
> smartplot(x^2/9-y^2/4=1);
```



```
> # use the Maple plot command implicitplot --- need more plotting tools
> with(plots):
> implicitplot(y^5+x^2*y^3+16,x=-10..10,y=-2..0);
```



Problems for Chapter 1

Questions marked with [R] are routine, with [H] are harder and with [X] are for MATH1141 only. Questions marked with [V] have a video solution available from Moodle. You should make sure that you can do the easier questions before you tackle the more difficult questions.

Problems 1.1 : Sets of numbers

- [R] Express the following sets in words. Graph the sets on the number line (if possible).
 - $\{x \in \mathbb{Z} : -\pi < x < \pi\}$
 - $\{x \in \mathbb{R} : x^2 - x - 1 < 0\}$
 - $\{x \in \mathbb{Q} : x^2 = 2\}$
- [R] Graph on the number line the following sets.
 - $[3, \infty), (-\infty, 3), (-\infty, \infty), (-3, 3]$
 - $\{x : |x - 2| < 5\}$
 - $\{x : x^2 + 4x - 5 > 0\}$
- [R] Sketch the set of points (x, y) which satisfy the following relations.
 - $0 \leq y \leq 2x$ and $0 \leq x \leq 2$
 - $y/2 \leq x \leq 2$ and $0 \leq y \leq 4$

Problems 1.2 : Solving inequalities and 1.3 : Absolute values

- [R] Solve the following inequalities.
 - $x(x - 1) > 0$
 - $(x - 1)(x - 2) < 0$
 - $\frac{1}{x} > -\frac{1}{2}$
 - $\frac{1}{1 - x} > \frac{1}{2}$
 - $x \geq \frac{6}{x - 1}$
- [R] Solve the following inequalities.
 - $|x + 1| < 3$
 - $|x + 2| > 3$
 - $|3x + 2| < 1$
 - $\left| \frac{x - 1}{x + 1} \right| < 1$
- [R] [V]
 - By expanding $(x - y)^2$, prove that $x^2 + y^2 \geq 2xy$ for all real numbers x and y .
 - Deduce that $\frac{a + b}{2} \geq \sqrt{ab}$ for all non-negative real numbers a and b . When does equality hold?
 - Use the result above to find the minimum value of $y = x^2 + \frac{1}{x^2}$.

7. [R] True or false:
- a) If $a > b$ then $\frac{1}{a} < \frac{1}{b}$.
 - b) If $a < b$ then $a^2 < b^2$.
 - c) If $0 < a < b$ then $a^2 < b^2$.
 - d) If $a^2 + b^2 = 0$ then $a = b = 0$.
 - e) If $-1 < a < b$ then $a^2 < b^2$.
8. [H] Prove that $(x + y)^2 \geq 4xy$ and hence deduce that $\frac{1}{x^2} + \frac{1}{y^2} \geq \frac{4}{x^2 + y^2}$.
9. [H] [V]
- a) Prove that $f(x) = 1 + x + x^2$ is positive for all real numbers x .
 - b) By considering cases (or otherwise) prove that $1 + x + x^2 + x^3 + x^4$ is always positive.
 - c) Generalise the above results.

Problems 1.4 : Functions

10. [R] Determine the (maximal) domain and corresponding range for each function f described below.
- a) $f(x) = \sqrt{5 - x^2}$
 - b) $f(x) = \sqrt{x^2 - 5}$
 - c) $f(x) = (x - 8)^{-1/3}$
 - d) $f(x) = \sqrt{x - 1}$
 - e) $f(x) = \frac{1}{\sqrt{x - 1}}$
 - f) $f(x) = \sqrt{\sin x}$
 - g) $f(x) = \sqrt{1 - 2 \sin x}$
 - h) $f(x) = 1 + \tan^2 x$
 - i) $f(x) = \begin{cases} \cos x & \text{if } x < 0 \\ \sqrt{1 - x} & \text{if } 0 \leq x \leq 1 \\ |x| & \text{if } x > 1 \end{cases}$
11. [R] If $f(x) = x + 5$ and $g(x) = x^2 - 3$ then find
- a) $(g \circ f)(0)$
 - b) $(g \circ f)(x)$
 - c) $(f \circ g)(2)$
 - d) $(f \circ g)(x)$.
12. [R] If $f(x) = x - 1$ and $g(x) = \frac{1}{\sqrt{x - 1}}$, then give the explicit forms of
- a) $(f + g)(x)$
 - b) $(fg)(x)$
 - c) $\left(\frac{f}{g}\right)(x)$
 - d) $(f \circ g)(x)$.

Problems 1.5 : Polynomials and rational functions, 1.6 : The trigonometric functions and 1.7 : The elementary functions

13. [R] Draw neat sketches (preferably without using calculus) of the graphs given by the following equations.

- a) $y = x^2 - 5x + 6$ b) $y = 2x^3 - 16$ c) $y = \frac{4}{x-3}$
 d) $y = 2e^{x-1}$ e) $y = \frac{1}{x^2+4}$ f) $y = 3 \sin 2x$
 g) $y = \sqrt{x} - 1$
14. [R]
 a) Sketch the graph of $y = \sqrt{x+1}$ and use your graph to sketch (on the same diagram)
 $y = \frac{1}{\sqrt{x+1}}$.
 b) Repeat for $y = x^2 - 4x + 3$.
15. [R] Sketch the graph of $y = x^2 - 7x - 8$ and hence sketch the graph of $y = |x^2 - 7x - 8|$.
16. [R] What range of values will $x^2 + 4$ take if $-2 \leq x \leq 3$?
17. [R] Use a graphical approach to solve $|2x - 5| = x + 2$.
18. [R]
 a) Show that if p and q are polynomials then $p \circ q$ is again a polynomial.
 b) [H] Is the same true for rational functions?

Problems 1.8 : Implicitly defined functions

19. [R] Sketch the graphs given by the following equations.
- a) $\frac{x^2}{9} + \frac{y^2}{4} = 1$ b) $\frac{x^2}{9} - \frac{y^2}{4} = 1$
 c) $4x^2 + 9y^2 = 36$ d) $\frac{y^2}{9} - \frac{x^2}{4} = 1$

Chapter 2

Limits

In many situations, it is desirable to know what the long term state of a system will be. For example, suppose that an initially unpolluted lake containing 10^9 litres of water has a river flowing through it at a rate of 10^6 litres per day. A factory is built next to the lake and discharges 10^4 litres of pollution per day. By making some simple assumptions, it can be shown that the amount $P(t)$ of pollution in the lake after t days of the factory's operation is given by

$$P(t) = \frac{10^9}{101}(1 - e^{-101t/10^5}). \quad (2.1)$$

Environmental authorities want to know what the long term level of the pollution will be. To see whether the pollution in the lake eventually stabilises, and what the level of pollution will be, one studies the behaviour of $P(t)$ as $t \rightarrow \infty$.

Understanding the limiting behaviour of functions at infinity is important for practical and mathematical reasons. That is what we study in the next four sections.

2.1 Limits of functions at infinity

(Ref: SH10 §4.7)

In this section, we examine some techniques for calculating limits of the form $\lim_{x \rightarrow \infty} f(x)$.

2.1.1 Basic rules for limits

There are a few basic things to bear in mind when working with limits.

- If $f(x)$ gets arbitrarily close to some real number L as x tends to infinity, then we write

$$\lim_{x \rightarrow \infty} f(x) = L.$$

We can also write $f(x) \rightarrow L$ as $x \rightarrow \infty$.

- If $f(x)$ gets arbitrarily large (that is, approaches ∞) as x tends to ∞ then we write

$$f(x) \rightarrow \infty \text{ as } x \rightarrow \infty.$$

We do *not* write $\lim_{x \rightarrow \infty} f(x) = \infty$ since ∞ is not a real number. (The $\lim$ notation is only used when the limit exists and is a real number.)

- If $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ then

$$\lim_{x \rightarrow \infty} \frac{1}{f(x)} = 0.$$

This is intuitively obvious and gives limits such as

$$\lim_{x \rightarrow \infty} \frac{1}{x} = 0, \quad \lim_{x \rightarrow \infty} \frac{5}{4x^6} = 0, \quad \lim_{x \rightarrow \infty} e^{-x} = 0$$

(for the last example, recall that $e^{-x} = \frac{1}{e^x}$).

The following proposition shows that the arithmetic of limits behaves nicely in many situations.

Proposition 2.1.1. *Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then*

$$(i) \quad \lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$$

$$(ii) \quad \lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$$

$$(iii) \quad \lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$$

$$(iv) \quad \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}, \text{ provided that } \lim_{x \rightarrow \infty} g(x) \neq 0.$$

Part of this proposition will be proved in Section 2.4. For now, we demonstrate how the proposition is used to calculate limits.

Example 2.1.2. Calculate

$$\lim_{x \rightarrow \infty} \frac{3 - 1/x^2}{2 + e^{-x}}$$

if the limit exists. Justify each step of your working with reference to Proposition 2.1.1.

Solution.


$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{3 - 1/x^2}{2 + e^{-x}} &= \frac{\lim_{x \rightarrow \infty} (3 - 1/x^2)}{\lim_{x \rightarrow \infty} (2 + e^{-x})} && \text{(rule (iv))} \\ &= \frac{\lim_{x \rightarrow \infty} 3 - \lim_{x \rightarrow \infty} 1/x^2}{\lim_{x \rightarrow \infty} 2 + \lim_{x \rightarrow \infty} e^{-x}} && \text{(rules (i) and (ii))} \\ &= \frac{3 - 0}{2 + 0} \\ &= \frac{3}{2}. \end{aligned}$$

□

In practice, you may use the rules of Proposition 2.1.1 without referring to them, and leave out some of the steps shown above.

2.1.2 The pinching theorem

The idea behind the following theorem is intuitively obvious. Suppose, as illustrated in the diagram below, that the graph of a function h always lies above the graph of a function f . Suppose also that f and h have the same limit as $x \rightarrow \infty$. If the graph of a function g always lies between the graphs of h and f (so that g is ‘pinched’ between f and h) then g has the same limit also.



This idea is stated precisely below.

Theorem 2.1.3 (The pinching theorem). *Suppose that f , g and h are all defined on the interval (b, ∞) , where $b \in \mathbb{R}$. If*

$$f(x) \leq g(x) \leq h(x) \quad \forall x \in (b, \infty)$$

and

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} h(x) = L$$

then

$$\lim_{x \rightarrow \infty} g(x) = L.$$

One can modify the theorem to cover the case when $x \rightarrow -\infty$.

Example 2.1.4. Use the pinching theorem to find $\lim_{x \rightarrow \infty} \frac{\sin x}{x}$.

Solution. We begin with the basic inequality

$$-1 \leq \sin x \leq 1,$$

which is valid for every real number x . If we restrict x to the interval $(0, \infty)$ then we have

$$\frac{-1}{x} \leq \frac{\sin x}{x} \leq \frac{1}{x}.$$

Now

$$\lim_{x \rightarrow \infty} -\frac{1}{x} = \lim_{x \rightarrow \infty} \frac{1}{x} = 0,$$

and so

$$\lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0$$

by the pinching theorem. □

2.1.3 Limits of the form $f(x)/g(x)$

Suppose that we want to calculate a limit of the form

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where both $f(x)$ and $g(x)$ tend to infinity as $x \rightarrow \infty$. In this situation, we cannot apply the rules of Proposition 2.1.1 directly because the numerator and denominator don't have finite limits. Instead, we find an equivalent form of $\frac{f(x)}{g(x)}$ for which Proposition 2.1.1 can be applied. The key idea is to divide both f and g by the *fastest growing term* appearing in the denominator g .

Example 2.1.5. Evaluate $\lim_{x \rightarrow \infty} \frac{4x^2 - 5}{2x^2 + 3x}$, if it exists.

Solution. There are two terms appearing in the denominator: $2x^2$ and $3x$. As $x \rightarrow \infty$, the term which grows fastest is the one involving x^2 . So we divide the numerator and denominator by x^2 to evaluate the limit:

$$\begin{aligned} \frac{4x^2 - 5}{2x^2 + 3x} &= \frac{4 - 5/x^2}{2 + 3/x} \\ &\rightarrow \frac{4 - 0}{2 + 0} \end{aligned}$$

as $x \rightarrow \infty$. Therefore

$$\lim_{x \rightarrow \infty} \frac{4x^2 - 5}{2x^2 + 3x} = 2.$$

□

In general, we divide the numerator and denominator by the highest power of x in the *denominator*.

Example 2.1.6. Find $\lim_{x \rightarrow \infty} \frac{5x^3 + 6x^2 - 4 \sin x}{\cos 3x + 5x - 2x^3}$.

Example 2.1.7. Find $\lim_{x \rightarrow \infty} \frac{x^2 + 3x}{\sqrt{2x^4 + 3} - 4x}$.

2.1.4 Limits of the form $\sqrt{f(x)} - \sqrt{g(x)}$

Consider the limit of $\sqrt{x+5} - \sqrt{x+2}$ as $x \rightarrow \infty$. Since both $\sqrt{x+5}$ and $\sqrt{x+2}$ tend to infinity (rather than a finite number) as $x \rightarrow \infty$, one cannot apply Proposition 2.1.1. As the next example shows, a simple algebraic trick is used to overcome this difficulty.

Example 2.1.8. Evaluate $\lim_{x \rightarrow \infty} (\sqrt{x+5} - \sqrt{x+2})$, if it exists.

Solution. The algebraic trick is to multiply both the numerator and denominator by $\sqrt{x+5} + \sqrt{x+2}$ and expand the numerator as a difference of squares:

$$\begin{aligned} \sqrt{x+5} - \sqrt{x+2} &= \frac{(\sqrt{x+5} - \sqrt{x+2})(\sqrt{x+5} + \sqrt{x+2})}{\sqrt{x+5} + \sqrt{x+2}} \\ &= \frac{(x+5) - (x+2)}{\sqrt{x+5} + \sqrt{x+2}} \\ &= \frac{3}{\sqrt{x+5} + \sqrt{x+2}} \\ &\rightarrow 0 \end{aligned}$$

as $x \rightarrow \infty$. □

One should not be fooled into thinking, on the basis of the above example, that all limits of this type are zero.

Example 2.1.9. Show that $\lim_{x \rightarrow \infty} (\sqrt{x^2 + x} - x) = \frac{1}{2}$.

2.1.5 Indeterminate forms

We have already mentioned that the rules of Proposition 2.1.1 only apply when the limits involved are finite. In Section 2.1.3, we studied limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)},$$

when $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$. We say that such a limit is of the form $\frac{\infty}{\infty}$. We cannot say in advance whether or not a limit of the form $\frac{\infty}{\infty}$ exists, and if it does exist, what its value is.

For example, while the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behaviour as $x \rightarrow \infty$:

- $\frac{x^2}{x} \rightarrow \infty$ as $x \rightarrow \infty$
- $\frac{x}{x^2} \rightarrow 0$ as $x \rightarrow \infty$
- $\frac{2x^2}{x^2} \rightarrow 2$ as $x \rightarrow \infty$
- $\frac{x^2}{2x^2} \rightarrow \frac{1}{2}$ as $x \rightarrow \infty$.

Because we cannot determine in advance what kind of limiting behaviour something of the form $\frac{\infty}{\infty}$ has, we say that $\frac{\infty}{\infty}$ is an *indeterminate form*.

Other types of indeterminate forms (for example, limits of the form $\frac{0}{0}$ and $\infty - \infty$) arise naturally in many applications. Techniques for evaluating such limits will be developed at various stages of the course.


2.2 The definition of $\lim_{x \rightarrow \infty} f(x)$

Up to this point, we have treated limits in a rather intuitive way. It is now time to establish a rigorous basis for our treatment of limits. That is, we seek a definition for $\lim_{x \rightarrow \infty} f(x) = L$ that is mathematically precise and agrees with our intuitive notion of limits. How do we do this? A good strategy is to think deeply about a few simple examples before considering the general case.

Consider the real-valued functions f , g and h defined by

$$f(x) = \frac{1}{x}, \quad g(x) = \sin x \quad \text{and} \quad h(x) = 0.05$$

whenever $x > 0$. Their graphs are sketched below.



Intuitively, we see that $f(x)$ tends to 0 as x tends to infinity, while neither g nor h have this property. What is it about f that distinguishes its limiting behaviour from that of g or h ?


The above graph shows that the distance between $f(x)$ and its limit 0 is small whenever x is large. However, to get to the core difference between f and the other two functions, we must be more precise.

- The distance between $h(x)$ and 0 is also small, but its limit is not 0. The reason why $f(x)$ tends to 0 as $x \rightarrow \infty$ is that the distance between $f(x)$ and 0 can be made *as small as we like* whenever x is large enough.
- The distance between $g(x)$ and 0 can also be made as small as we like, by taking x sufficiently close to an integer multiple of π . However the limit of $g(x)$ is not 0. The difference between f and g is that the distance between $f(x)$ and 0 *remains* small for *all* sufficiently large values of x . The same is not true for g .


In summary, $f(x)$ tends to 0 as $x \rightarrow \infty$ because the distance between $f(x)$ and 0 can be made *as small as we like* for *all* values of x that are sufficiently large.

The next step in crafting a good definition is to express phrases like ‘can be made as small as we like’ and ‘sufficiently large’ in concrete mathematical language. The following list is one attempt to do this. (As a reminder, $f(x) = 1/x$.)

- The distance between $f(x)$ and 0 is less than 1 whenever $x > 1$.




- The distance between $f(x)$ and 0 is less than 0.2 whenever $x > 5$.



- The distance between $f(x)$ and 0 is less than 0.1 whenever $x > 10$.
- The distance between $f(x)$ and 0 is less than 0.01 whenever $x > 100$.
- The distance between $f(x)$ and 0 is less than 0.0001 whenever $x > 10000$.
- *etc.*

Of course, to express the fact that ‘the distance between $f(x)$ and 0 can be made as small as we like’ in this way would require an infinite list. Instead, we say that, for *every* small positive number ϵ , there is real number M such that

- the distance between $f(x)$ and 0 is less than ϵ whenever $x > M$.



By looking at the graph we can see that $M = \frac{1}{\epsilon}$. (The symbol ϵ is the Greek letter ‘epsilon’ and is traditionally used by mathematicians in this context.)

So far we have agreed that $f(x) \rightarrow 0$ as $x \rightarrow \infty$ precisely because

for every positive real number ϵ , there is a real number M such that the distance between $f(x)$ and 0 is less than ϵ whenever $x > M$.

This statement forms a basis for defining what is meant by $\lim_{x \rightarrow \infty} f(x) = 0$. A general definition for what is meant by $\lim_{x \rightarrow \infty} f(x) = L$ is now obtained by


- replacing the limit 0 with the limit L , and then
- replacing ‘the distance between $f(x)$ and L ’ with $|f(x) - L|$ (see Section 1.3).

We are now ready to give a general definition for the limit of a function at infinity.

Definition 2.2.1. Suppose that L is a real number and f is a real-valued function defined on some interval (b, ∞) . We say that $\lim_{x \rightarrow \infty} f(x) = L$ if


for every positive real number ϵ , there is a real number M such that if $x > M$ then $|f(x) - L| < \epsilon$.

Remark 2.2.2. Suppose that $\lim_{x \rightarrow \infty} f(x) = L$. Definition 2.2.1 can be interpreted geometrically in the following way. For every small number ϵ , you can find a marker M on the x -axis such that the distance between $f(x)$ and the limit L is less than ϵ whenever x lies to the right of M .



Remark 2.2.3. In Definition 2.2.1, the number M depends on ϵ . In general, the smaller the value of ϵ , the larger the value of M . We note, however, that for any particular value of ϵ the choice of M is not unique. For example, the M chosen for the above diagram is different from the M chosen for the diagram below, but both choices guarantee that

$$|f(x) - L| < \epsilon \quad \text{whenever} \quad x > M.$$



Remark 2.2.4. Students who find Definition 2.2.1 difficult to understand can take comfort in the fact that a rigorous formulation of limits evaded mathematicians for over 2000 years. The problems and paradoxes involving limiting processes tabled around 450 BC by the ancient Greek philosopher Zeno of Elea were only satisfactorily resolved in the nineteenth century, when a rigorous definition of the limit was given by Bolzano (in 1817) and Weierstrass (in the 1850s).

2.3 Proving that $\lim_{x \rightarrow \infty} f(x) = L$ using the limit definition

In the previous section we gave a definition for $\lim_{x \rightarrow \infty} f(x) = L$. In this section this definition is used to prove limits that were previously taken for granted.

Example 2.3.1. Prove from Definition 2.2.1 that $\lim_{x \rightarrow \infty} \frac{4}{x^2} = 0$.

Proof. Suppose that $f(x) = 4/x^2$, $L = 0$ and $\epsilon > 0$. We begin by calculating the distance between the function and its limit:

$$\begin{aligned} |f(x) - L| &= \left| \frac{4}{x^2} - 0 \right| \\ &= \frac{4}{x^2}. \end{aligned} \tag{2.2}$$

We need to find a condition on x such that $|f(x) - L| < \epsilon$. By (2.2), we require that

$$\frac{4}{x^2} < \epsilon.$$

By rearrangement, we see that this is satisfied whenever

$$x > \frac{2}{\sqrt{\epsilon}}.$$

So far we have shown that $|f(x) - L| < \epsilon$ whenever $x > \frac{2}{\sqrt{\epsilon}}$. Hence if

$$M = \frac{2}{\sqrt{\epsilon}}. \quad (2.3)$$

then $|f(x) - L| < \epsilon$ whenever $x > M$. This completes the proof. $\square$

The above proof can be streamlined by omitting some explanatory commentary. In the next example, explanatory comments (which are not part of the proof) are inserted inside square brackets.

Example 2.3.2. Prove that

$$\lim_{x \rightarrow \infty} \frac{5x}{x+3} = 5$$

by using Definition 2.2.1.

Proof. Suppose that $\epsilon > 0$. For convenience, suppose also that $x > 0$. [This is allowed since we are considering the behaviour of $f(x)$ for large positive values of x .] Now

$$\begin{aligned} |f(x) - L| &= \left| \frac{5x}{x+3} - 5 \right| \\ &= \left| \frac{5x - 5(x+3)}{x+3} \right| \\ &= \left| \frac{-15}{x+3} \right| \\ &= \frac{15}{x+3} && [\text{since } x > 0] \\ &< \frac{15}{x} && [\text{to make algebra simpler later on}]. \end{aligned}$$

In summary,

$$|f(x) - L| < \frac{15}{x}.$$

[This inequality gives an *upper bound* for the distance between $f(x)$ and L .] Hence $|f(x) - L| < \epsilon$ whenever

$$\frac{15}{x} < \epsilon.$$

This condition is equivalent to

$$x > \frac{15}{\epsilon}.$$

Hence if $M = \frac{15}{\epsilon}$ then

$$|f(x) - L| < \epsilon \quad \text{whenever} \quad x > M,$$

as required. $\square$

Before attempting one more proof, we reflect on the general strategy employed. Given ϵ , we need to find a number M such that

$$|f(x) - L| < \epsilon \quad \text{whenever} \quad x > M.$$

The number M can be found by following the procedure below.

1. Find a good upper bound for $|f(x) - L|$.
2. Find a simple condition on x such that this upper bound is less than ϵ .
3. Use this condition to state an appropriate value for M (in terms of ϵ).

Example 2.3.3. Prove that

$$\lim_{x \rightarrow \infty} \frac{\sin x}{9x^2 + e^{3x}} = 0$$

by using Definition 2.2.1.

Proof. Suppose that $\epsilon > 0$. Now

$$\begin{aligned} |f(x) - L| &= \left| \frac{\sin x}{9x^2 + e^{3x}} - 0 \right| \\ &= \frac{|\sin x|}{|9x^2 + e^{3x}|} \\ &\leq \frac{1}{9x^2 + e^{3x}} && \text{[using the fact that } -1 \leq \sin x \leq 1\text{]} \\ &< \frac{1}{9x^2} && \text{[decreasing the denominator increases the RHS].} \end{aligned}$$

In summary,

$$|f(x) - L| < \frac{1}{9x^2}.$$

[This is our upper bound for $|f(x) - L|$.] Now

$$\frac{1}{9x^2} < \epsilon$$

whenever

$$x > \frac{1}{3\sqrt{\epsilon}}.$$

Set

$$M = \frac{1}{3\sqrt{\epsilon}}.$$

Then $|f(x) - L| < \epsilon$ whenever $x > M$, as required. $\square$

Remark 2.3.4. As pointed out in Remark 2.2.3, the value of M is not unique. For instance, the upper bound for $|f(x) - L|$ in our solution to Example 2.3.3 was $1/(9x^2)$ and the corresponding value for M was $1/(3\sqrt{\epsilon})$. If instead we use the upper bound $1/e^{3x}$ then the corresponding value for M is $\frac{1}{3} \ln \frac{1}{\epsilon}$.

2.4 Proofs of basic limit results (MATH1141 only)

By using Definition 2.2.1, one can now prove the limit rules stated in Section 2.1. The selection of proofs given in this section is designed to give MATH1141 students a taste of how such results are proved.

Proof of Proposition 2.1.1 (i). Suppose that

$$\lim_{x \rightarrow \infty} f(x) = L_1 \quad \text{and} \quad \lim_{x \rightarrow \infty} g(x) = L_2.$$

We need to show, using Definition 2.2.1, that

$$\lim_{x \rightarrow \infty} (f(x) + g(x)) = L_1 + L_2.$$

We begin by fixing a positive real number ϵ .

Since $\lim_{x \rightarrow \infty} f(x) = L$, there is a positive M_1 such that

$$|f(x) - L_1| < \frac{\epsilon}{2} \quad \text{whenever} \quad x > M_1.$$

Similarly, there is a positive M_2 such that

$$|g(x) - L_2| < \frac{\epsilon}{2} \quad \text{whenever} \quad x > M_2.$$

Hence, by the triangle inequality,

$$\begin{aligned} |(f(x) + g(x)) - (L_1 + L_2)| &\leq |f(x) - L_1| + |g(x) - L_2| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon \end{aligned}$$

whenever both $x > M_1$ and $x > M_2$. Now set $M = \max\{M_1, M_2\}$ so that

$$|(f(x) + g(x)) - (L_1 + L_2)| < \epsilon \quad \text{whenever} \quad x > M.$$

This proves that $\lim_{x \rightarrow \infty} (f(x) + g(x)) = L_1 + L_2$. $\square$

The other rules in Proposition 2.1.1 can be proved similarly, except that sometimes the desired inequalities are harder to obtain. We move on to prove the pinching theorem.

Proof of Theorem 2.1.3. Suppose that

$$f(x) \leq g(x) \leq h(x) \quad \forall x \in (b, \infty)$$

and that

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} h(x) = L.$$

Fix a positive real number ϵ .

Since $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} h(x) = L$, there are real numbers M_1 and M_2 , both greater than b , such that

$$|f(x) - L| < \epsilon \quad \text{whenever} \quad x > M_1$$

and

$$|h(x) - L| < \epsilon \quad \text{whenever} \quad x > M_2.$$

In other words,

$$L - \epsilon < f(x) < L + \epsilon \quad \text{whenever} \quad x > M_1$$

and

$$L - \epsilon < h(x) < L + \epsilon \quad \text{whenever} \quad x > M_2.$$

Hence

$$L - \epsilon < f(x) \leq g(x) \leq h(x) < L + \epsilon$$

whenever $x > \max\{M_1, M_2\}$. This shows that

$$|g(x) - L| < \epsilon \quad \text{whenever} \quad x > \max\{M_1, M_2\},$$

completing the proof. □


2.5 Limits of functions at a point

(Ref: SH10 §§2.1–2.3, 2.5)

So far we have only discussed limits of the form $\lim_{x \rightarrow \infty} f(x)$. In this section we study limits of the form $\lim_{x \rightarrow a} f(x)$, where a is a finite real number.

2.5.1 Left-hand, right-hand and two-sided limits

Consider the function f whose graph is shown below.



With reference to this graph, we will discuss the behaviour of $f(x)$ when x is near the points 2, 4, 6 and 8.

- As x approaches 2 from the left-hand side, $f(x)$ approaches 3. In this situation, we write

$$\lim_{x \rightarrow 2^-} f(x) = 3$$

and say that ‘the limit of $f(x)$ as x approaches 2 from the left is 3.’ (Note that $f(2) = 1 \neq 3$. One should not confuse a limit with a function value.) As x approaches 8 from the left-hand side, $f(x)$ approaches 1 and hence we write

$$\lim_{x \rightarrow 8^-} f(x) = 1.$$

Both of these limits are called *left-hand limits*.

- As x approaches 2 from the right-hand side, $f(x)$ approaches 1. In this situation, we write

$$\lim_{x \rightarrow 2^+} f(x) = 1$$

and say that ‘the limit of $f(x)$ as x approaches 2 from the right is 1.’ As x approaches 6 from the right-hand side, $f(x)$ approaches 2 and hence we write

$$\lim_{x \rightarrow 6^+} f(x) = 2.$$

Both of these limits are called *right-hand limits*.

- As x approaches 4 from the left-hand side, $f(x)$ is positive and grows large without bound. In this case, $\lim_{x \rightarrow 4^-} f(x)$ does not exist because $f(x)$ does not approach a real number. However, we can write

$$f(x) \rightarrow \infty \quad \text{as} \quad x \rightarrow 4^-.$$

- As x approaches 4 from the right-hand side, $f(x)$ is negative and grows large without bound. In this case, $\lim_{x \rightarrow 4^+} f(x)$ does not exist but we can write

$$f(x) \rightarrow -\infty \quad \text{as} \quad x \rightarrow 4^+.$$

Once an understanding of left- and right-hand limits is grasped, we can talk about *two-sided limits*.

Definition 2.5.1. If the left-hand limit $\lim_{x \rightarrow a^-} f(x)$ and the right-hand limit $\lim_{x \rightarrow a^+} f(x)$ both exist and equal the same real number L , then we say that *the limit of $f(x)$ as $x \rightarrow a$ exists and is equal to L* , and we write

$$\lim_{x \rightarrow a} f(x) = L.$$

If any one of these conditions fails then we say that $\lim_{x \rightarrow a} f(x)$ does *not* exist.

Recall the graph of the function f introduced at the beginning of this section.

- The two-sided limit $\lim_{x \rightarrow 2} f(x)$ does not exist because

$$\lim_{x \rightarrow 2^-} f(x) \neq \lim_{x \rightarrow 2^+} f(x).$$

- The two-sided limit $\lim_{x \rightarrow 4} f(x)$ does not exist because $\lim_{x \rightarrow 4^-} f(x)$ does not exist.

- Since the left-hand and right-hand limits both exist at 6 and

$$\lim_{x \rightarrow 6^-} f(x) = \lim_{x \rightarrow 6^+} f(x) = 2,$$

the two-sided limit $\lim_{x \rightarrow 6} f(x)$ exists and

$$\lim_{x \rightarrow 6} f(x) = 2.$$

Note that $f(6) = 4$ and so

$$\lim_{x \rightarrow 6} f(x) \neq f(6).$$

That is, the two-sided limit of a function at a point need not equal the value of the function at that point.

- Since the left-hand and right-hand limits at 8 both exist and equal 1, the two-sided limit $\lim_{x \rightarrow 8} f(x)$ exists and

$$\lim_{x \rightarrow 8} f(x) = 1.$$

In this case, the two-sided limit of f at 8 equals the value of the function at 8. That is,

$$\lim_{x \rightarrow 8} f(x) = f(8).$$

The last two points emphasise an important feature of limits and functions. If $\lim_{x \rightarrow a} f(x) = L$ then $f(x)$ approaches L as x tends to a . The function f may not be defined at a , or if it is defined at a , we can have the situation where $f(a) = L$ or where $f(a) \neq L$. In other words, the value of $f(a)$ does not determine the value of $\lim_{x \rightarrow a} f(x)$. As far as the limit is concerned, all that matters is how f behaves when it is *very close* to the point a .


Remark 2.5.2. In this section we have not given a proper definition for limits of the form $\lim_{x \rightarrow a^-} f(x)$ or $\lim_{x \rightarrow a^+} f(x)$. At this stage it is enough to know that such a rigorous definition exists and is similar to the definition for limits of the form $\lim_{x \rightarrow \infty} f(x)$ (see Definition 2.2.1). This definition can be used to prove the basic limit results presented in the remainder of this chapter.

2.5.2 Limits and continuous functions

Using limits we can now give a definition for continuity.

Definition 2.5.3. Suppose that f is defined on some open interval containing the point a . If $\lim_{x \rightarrow a} f(x) = f(a)$, then we say that f is *continuous* at a ; otherwise, we say that f is *discontinuous* at a .

To illustrate, consider the function f whose graph is sketch below.



The function f is continuous at 2, discontinuous at 4 and undefined at 6. (We do not say that f is discontinuous at 6 since $6 \notin \text{Dom}(f)$.)

We now extend our formal notion of continuity at a point to continuity on the real line.

Definition 2.5.4. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at every point a in $\mathbb{R}$ then we say that f is *continuous everywhere*.

If a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous everywhere then it has the property that its graph can be drawn on the Cartesian plane as an unbroken curve. This coincides with intuitive notions about continuity gained in high school.

As mentioned in Section 1.9, polynomials, rational functions, the trigonometric functions, exponentials and logarithms are continuous at every point in the respective domains. Moreover, if f is continuous at a point a then $|f|$ is also continuous at a (see Section 1.7 for the definition of $|f|$). The proof of these facts requires a rigorous definition of the limit at a point, and, as already mentioned, such a definition is not presented in this course. However, we will discuss aspects of the proof in Chapter 3.

A consequence of the above definition is that both one-sided and two-sided limits of f at a point a are easy to evaluate provided that (a) $a \in \text{Dom}(f)$ and (b) we know that f is a continuous at a . Hence, in view the facts mentioned above,

- $\lim_{x \rightarrow 2} \sin(x) = \sin(2)$
- $\lim_{x \rightarrow 3^+} (x^2 - x + 1) = \lim_{x \rightarrow 3} (x^2 - x + 1) = 3^2 - 3 + 1 = 7$
- $\lim_{x \rightarrow 4} |x - 10| = |4 - 10| = 6$.

Remark 2.5.5. Continuity is a deep property for a function to have. Contrary to the impression that many students form from their study of functions at school, most functions are *not* continuous. Three interesting functions with discontinuities are mentioned below.

- The function $f : \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q}, \end{cases}$$

is discontinuous at every point in its domain.

- The function $f : \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$f(x) = \begin{cases} x & \text{if } x \in \mathbb{Q} \\ -x & \text{if } x \notin \mathbb{Q}, \end{cases}$$

is continuous at 0 but discontinuous everywhere else.

- It is not too difficult to construct a function $f : \mathbb{R} \rightarrow \mathbb{R}$ which is continuous at every irrational point but discontinuous at every rational point.

Hence, one should not assume that the graph of a general function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at even a majority of points in its domain. Continuous functions are quite special and deserve much attention. We will study them in greater depth in Chapter 3.

2.5.3 Rules for limits at a point

The following proposition, which is analogous to Proposition 2.1.1, shows that limits at a point have nice arithmetic properties.

Proposition 2.5.6. *Suppose that $a \in \mathbb{R}$ and that $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist and are finite real numbers. Then*

$$(i) \quad \lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

$$(ii) \quad \lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

$$(iii) \quad \lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$$

$$(iv) \quad \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \text{ provided that } \lim_{x \rightarrow a} g(x) \neq 0.$$

A similar set of rules hold for left- and right-hand limits.

Proposition 2.5.6 says that limits interact nicely with function addition, subtraction, multiplication and division. The next proposition shows that limits interact nicely with function composition.

Proposition 2.5.7. *If $\lim_{x \rightarrow a} f(x) = L$ and g is continuous at L then*

$$\lim_{x \rightarrow a} g(f(x)) = g\left(\lim_{x \rightarrow a} f(x)\right).$$

If the functions f and g are continuous everywhere, then Proposition 2.5.7 implies that

$$\lim_{x \rightarrow a} g(f(x)) = g\left(\lim_{x \rightarrow a} f(x)\right)$$

for any point a in $\mathbb{R}$.

Example 2.5.8. Evaluate the limit

$$\lim_{x \rightarrow \sqrt{\pi/3}} \cos\left(x^2 + \frac{\pi}{6}\right),$$


justifying each step of your calculation.

Solution. Since the cosine function and polynomials are continuous everywhere,

$$\begin{aligned}
 \lim_{x \rightarrow \sqrt{\pi/3}} \cos \left(x^2 + \frac{\pi}{6} \right) &= \cos \left(\lim_{x \rightarrow \sqrt{\pi/3}} x^2 + \frac{\pi}{6} \right) && \text{(by Proposition 2.5.7)} \\
 &= \cos \left(\frac{\pi}{3} + \frac{\pi}{6} \right) && \text{(since polynomials are continuous)} \\
 &= \cos \left(\frac{\pi}{2} \right) \\
 &= 0.
 \end{aligned}$$

□

A version of the pinching theorem for two-sided limits is given below. Similar versions exist for left- and right-hand limits.



Theorem 2.5.9 (The pinching theorem). *Let I be an open interval containing the point a . Suppose that f , g and h are all defined on I except possibly at the point a . If*

$$f(x) \leq g(x) \leq h(x) \quad \forall x \in I$$

and

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$$

then

$$\lim_{x \rightarrow a} g(x) = L.$$

Remark 2.5.10. The pinching theorem can be used to prove the well-known limit

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1.$$

See the problem set of Chapter 2 for details.

We close this chapter with four examples involving limits of the form $\lim_{x \rightarrow a} f(x)$, where f is not necessarily continuous at the point a . Determining whether or not the limit exists, and if so what its value is, requires special analysis of the function f .

Example 2.5.11. Determine whether or not the limit

$$\lim_{x \rightarrow 0} x \sin(1/x)$$

exists. If it exists, find its value.

Solution. We begin with the inequality

$$-1 \leq \sin \theta \leq 1$$

which is valid for all θ in $\mathbb{R}$. In particular, if $\theta = 1/x$ then

$$-1 \leq \sin(1/x) \leq 1 \quad \text{whenever } x \neq 0.$$

Case 1: Suppose that $x > 0$. Then we have

$$-x \leq x \sin(1/x) \leq x.$$

Since $\lim_{x \rightarrow 0^+} -x = \lim_{x \rightarrow 0^+} x = 0$, it follows from the pinching theorem that $\lim_{x \rightarrow 0^+} x \sin(1/x) = 0$.

Case 2: Suppose that $x < 0$. Then


$$-x \geq x \sin(1/x) \geq x.$$

Since $\lim_{x \rightarrow 0^-} -x = \lim_{x \rightarrow 0^-} x = 0$, it follows from the pinching theorem that $\lim_{x \rightarrow 0^-} x \sin(1/x) = 0$.

Conclusion: Since the left- and right-hand limits exist and are both equal to 0, the two-sided limit exists and

$$\lim_{x \rightarrow 0} x \sin(1/x) = 0.$$

This limit is illustrated graphically below.



□

Example 2.5.12. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula


$$f(x) = \begin{cases} \frac{x^2 - 4}{x - 2} & \text{if } x \neq 2 \\ 0 & \text{if } x = 2. \end{cases}$$

Discuss the limiting behaviour of $f(x)$ as x approaches 2.

Solution. Since we are discussing limiting behaviour, the value of f at 2 is irrelevant. All that matters is how $f(x)$ behaves *near* 2. So suppose that $x \neq 2$. Then

$$\begin{aligned} f(x) &= \frac{x^2 - 4}{x - 2} \\ &= \frac{(x - 2)(x + 2)}{x - 2} \\ &= x + 2. \end{aligned}$$

The graph of f is now easy to sketch.



Since

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = 4,$$

the two-sided limit exists and we have $\lim_{x \rightarrow 2} f(x) = 4$. (Note that f is not continuous at 2 since $\lim_{x \rightarrow 2} f(x) \neq f(2)$.) $\square$

Example 2.5.13. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula

$$f(x) = \begin{cases} \frac{|x^2 - 9|}{x - 3} & \text{if } x \neq 3 \\ 2 & \text{if } x = 3. \end{cases}$$

Discuss the limiting behaviour of $f(x)$ as x approaches 3.

Example 2.5.14. Suppose that

$$f(x) = \sin\left(\frac{\pi}{x}\right), \quad x \neq 0.$$

Discuss the limiting behaviour of $f(x)$ as x approaches 0.

Solution. Since f is an odd function, we initially restrict our attention to positive values of x . Now $\sin \theta = 0$ whenever $\theta \in \{\pi, 2\pi, 3\pi, \dots\}$. If $\theta = \pi/x$ then we have $\sin(\pi/x) = 0$ whenever


$$x = \frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \dots$$

Similarly, $\sin \theta = 1$ whenever $\theta \in \{\pi/2, 5\pi/2, 9\pi/2, 13\pi/2, \dots\}$ so $\sin(\pi/x) = 1$ whenever

$$x = \frac{2}{1}, \frac{2}{5}, \frac{2}{9}, \frac{2}{13}, \dots$$

Consequently, no matter how close x is to 0 on the right, we can *always* find two closer points x_1 and x_2 such that $f(x_1) = 0$ and $f(x_2) = 1$. We conclude that f has no right-hand limit at 0. Since f is odd, it follows that f has no left-hand limit at 0. Hence $\lim_{x \rightarrow 0^+} f(x)$ does not exist.

The graph of f near the origin is shown below.



The $f(x)$ oscillates wildly between 1 and -1 as x tends to 0, and $\lim_{x \rightarrow 0^+} f(x)$ does not exist. $\square$

2.6 Maple notes

The following Maple command is relevant to the material of this chapter:

`limit(f(x),x=a);` finds the (two-sided) limit of $f(x)$ as $x \rightarrow a$. One-sided limits can be specified using `limit(f(x),x=a,right);` or `limit(f(x),x=a,left);`. For example,

```
> limit(sin(x)/x, x=0);
1

> limit(sin(x)/x, x=infinity);
0

> # the next limit needs methods from chapter 5 for hand calculations
> limit((1-cos(x))/x^2, x=0, left);
1/2

> # Maple produces results that are not in strict agreement with our definitions
as x goes to plus or minus infinity
> limit(x^2, x=-infinity);
infinity

> limit(sin(x), x=-infinity);
-1...1

> limit(x^2*sin(x), x=-infinity);
undefined
```

Problems for Chapter 2

Problems 2.1 : Limits of functions at infinity

1. [R] [V] Evaluate the following limits if they exist.
 - a) $\lim_{x \rightarrow \infty} \frac{x^2 - 1}{x^2 + 1}$
 - b) $\lim_{x \rightarrow \infty} \frac{2x^2 + x - 1}{x^2 + 4x - 3}$
 - c) $\lim_{x \rightarrow \infty} \frac{2x^2 + 5x - 1}{x^3 + x}$
 - d) $\lim_{x \rightarrow \infty} \frac{x^5 + 5x + 1}{x^4 + 3}$
 - e) $\lim_{x \rightarrow \infty} \frac{5x^2 - 3x + \cos 7x}{4 + \sin 2x + x^2}$
 - f) $\lim_{x \rightarrow \infty} \sin x$
2. [R] Use the pinching theorem to find the following limits.
 - a) $\lim_{x \rightarrow \infty} \frac{\sin x}{x}$
 - b) $\lim_{x \rightarrow \infty} \frac{\cos x}{x^2}$
3. [R] [V]
 - a) Prove that $\lim_{x \rightarrow \infty} (\sqrt{x+1} - \sqrt{x}) = 0$.
 - b) Show that $\lim_{x \rightarrow \infty} (\sqrt{x^2 + x} - x) = \frac{1}{2}$.

Problems 2.2 : The definition of $\lim_{x \rightarrow \infty} f(x)$ and2.3 : Proving that $\lim_{x \rightarrow \infty} f(x) = L$ using the limit definition

4. [R]
 - a) Write down the formal definition for the statement

$$\lim_{x \rightarrow \infty} f(x) = L.$$
 - b) Evaluate $\lim_{x \rightarrow \infty} \frac{1}{2x^2}$.
 - c) Verify from the formal definition that your answer in (b) is correct.
5. [R]
 - a) Evaluate $\lim_{x \rightarrow \infty} \frac{x^2 + 1}{x^2}$.
 - b) Find a real number M such that the distance between $\frac{x^2 + 1}{x^2}$ and its limit is less than 0.01 whenever $x > M$.
 - c) Suppose that $\epsilon > 0$. Find a real number M (expressed in terms of ϵ) such that the distance between $\frac{x^2 + 1}{x^2}$ and its limit is less than ϵ whenever $x > M$.

6. [R] [V] For each of the following, find the limit of $f(x)$ as x tends to infinity and prove from the definition that your answer is correct.

a) $f(x) = \frac{4x}{x+7}$ b) $f(x) = \frac{x-3}{x^2+3}$ c) $f(x) = e^{-2x}$
 d) $f(x) = \frac{\sin x}{x}$ e) $f(x) = \frac{\sin 3x}{x^2+4}$

7. [X]

- a) With ϵ in $(0, 1)$, Sarah solves the inequality $|f(x) - 4| < \epsilon$ and finds that the required x values satisfy

$$x \in \left(\frac{1}{\epsilon}, \infty\right).$$

Does $\lim_{x \rightarrow \infty} f(x)$ exist? Give reasons for your answer.

- b) With ϵ in $(0, 1)$, Lyndal solves the inequality $|g(x) - 5| < \epsilon$ and finds that the inequality holds for all x satisfying

$$x \in \left(\frac{1}{\epsilon}, \infty\right).$$

Does $\lim_{x \rightarrow \infty} g(x)$ exist? Give reasons for your answer.

8. [R] [V] A parcel is dropped from an aeroplane. A simple model, taking into account gravity and air resistance, suggests that the parcel's velocity $v(t)$ (in metres per second) is given by $v(t) = 50(1 - e^{-t/5})$, where t is the number of seconds since leaving the plane.
- a) Calculate the terminal velocity of the parcel (that is, find $\lim_{t \rightarrow \infty} v(t)$).
- b) The parcel never attains its terminal velocity. How long does it take to come within 1 metre per second of its terminal velocity?

Problems 2.4 : Proofs of basic limit results (MATH1141 only)

9. [X] For each question below, give reasons for your answer. [In some cases a single example will be sufficient while in other cases a general proof will be required. As a reminder, if $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ then $\lim_{x \rightarrow \infty} f(x)$ does not exist.]
- a) If $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ do not exist, can $\lim_{x \rightarrow \infty} [f(x) + g(x)]$ or $\lim_{x \rightarrow \infty} f(x)g(x)$ exist?
- b) If $\lim_{x \rightarrow \infty} f(x)$ exists and $\lim_{x \rightarrow \infty} [f(x) + g(x)]$ exists, must $\lim_{x \rightarrow \infty} g(x)$ exist?
- c) If $\lim_{x \rightarrow \infty} f(x)$ exists and $\lim_{x \rightarrow \infty} g(x)$ does not exist, can $\lim_{x \rightarrow \infty} [f(x) + g(x)]$ exist?
- d) If $\lim_{x \rightarrow \infty} f(x)$ exists and $\lim_{x \rightarrow \infty} f(x)g(x)$ exists, does it follow that $\lim_{x \rightarrow \infty} g(x)$ exists?

Problems 2.5 : Limits of functions at a point

10. [R] Evaluate the following limits.

a) $\lim_{x \rightarrow 3} 2x + 4$ b) $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$ c) $\lim_{x \rightarrow 1} \frac{x^3 - 1}{x - 1}$ d) $\lim_{x \rightarrow 3} \frac{\frac{1}{x} - \frac{1}{3}}{x - 3}$

11. [R] [V]


- a) Find the left-hand limit $\lim_{x \rightarrow 2^-} \frac{|x-2|}{x-2}$.
- b) Find the right-hand limit $\lim_{x \rightarrow 2^+} \frac{|x-2|}{x-2}$.
- c) Does $\lim_{x \rightarrow 2} \frac{|x-2|}{x-2}$ exist?

12. [R] By finding the left- and right-hand limits first, decide whether or not each of the following limits exist and if so find their values.

- a) $\lim_{x \rightarrow 0} \frac{x}{|x|}$ b) $\lim_{x \rightarrow 2} \frac{|x^2-4|}{x-2}$ c) $\lim_{x \rightarrow 4} \frac{x-4}{|x-4|}$ d) $\lim_{x \rightarrow 0} \frac{4}{x}$

13. [R] [V]

- a) Use the pinching theorem to find $\lim_{x \rightarrow 0} x \sin \frac{1}{x}$.
- b) Repeat for $\lim_{x \rightarrow 0} x^2 \sin \frac{1}{2x}$.

14. [R] [V] Suppose that θ is a (positive) angle measured in radians and consider the diagram below.

The curve segment CB is the arc of a circle of radius 1 centre O .

- a) Write down, in terms of θ , the length of arc CB and the lengths of the line segments CA and DB .
- b) By considering areas, deduce that $\sin \theta \cos \theta \leq \theta \leq \tan \theta$ whenever $0 < \theta < \frac{\pi}{2}$.
- c) Use the pinching theorem to show that $\lim_{\theta \rightarrow 0^+} \frac{\theta}{\sin \theta} = 1$.
- d) Deduce that $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$.
15. [H] Discuss the limiting behaviour of $\cos \frac{1}{x}$ as $x \rightarrow 0$.

Chapter 3

Properties of continuous functions

In the world of functions, there is a small civilised country inhabited by the race of continuous functions. Citizens of this country, such as the polynomials and exponentials, are studied (and perhaps even loved) by students in high schools throughout the human world. However, outside this small civilised country is a vast untamed universe of all sorts of discontinuous functions. From this perspective, the continuous functions are a very rare breed.

However, continuous functions are far from endangered. They are among the most useful functions for modelling real-life phenomena, particularly when the quantity that is being modelled occurs in nature and changes smoothly with time. For example,

- the distance from the earth to the sun at time t ,
- the height of Mount Everest at time t ,
- the volume of water in a dam at time t ,
- the mass of an earthworm at time t , and
- the temperature at the back of a lecture theatre at time t

can all be modelled by continuous functions of the variable t .

In addition to their usefulness for modelling naturally occurring quantities, continuous functions have important mathematical characteristics. The combination of the usefulness of continuous functions and their rarity among the world of functions means that continuity is a deep property. In this chapter we begin to explore the ramifications of continuity.

3.1 Combining continuous functions

(Ref: SH10 §2.4)

If two continuous functions are combined via function addition, subtraction, multiplication, division or composition, when is the resulting function also continuous? The two next propositions answer this question.

Proposition 3.1.1. *Suppose that the functions f and g are continuous at a point a . Then $f + g$, $f - g$ and fg are continuous at a . If $g(a) \neq 0$ then f/g is also continuous at a .*

Proof. Suppose that f and g are continuous at a . Then, by the definition of continuity at a point (see Definition 2.5.3), $\lim_{x \rightarrow a} f(x) = f(a)$ and $\lim_{x \rightarrow a} g(x) = g(a)$. Therefore

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} (f(x) + g(x)) && \text{(by the definition of } f + g) \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) && \text{(by Proposition 2.5.6)} \\ &= f(a) + g(a) && \text{(since } f \text{ and } g \text{ are continuous)} \\ &= (f + g)(a) && \text{(by the definition of } f + g). \end{aligned}$$

Hence $f + g$ is continuous at a .

The proofs that the functions $f - g$, fg and f/g are continuous at a are similar. $\square$

Proposition 3.1.2. *Suppose that f is continuous at a and that g is continuous at $f(a)$. Then $g \circ f$ is continuous at a .*

Proof. Combine Definition 2.5.3 with Proposition 2.5.7. $\square$

In Chapters 1 and 2 it was stated (without proof) that the polynomials, rational functions and trigonometric functions are continuous at every point in their respective domains. The next example shows that the continuity of these functions can be deduced from the continuity of

- the constant functions
- the sine function, and
- the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x$.

Example 3.1.3. Suppose that a is a point in $\mathbb{R}$. By assuming that the constant functions, the sine function and the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = x$, are continuous everywhere, show that

- (a) every polynomial is continuous at a ,
- (b) every rational function r is continuous at a (provided that $a \in \text{Dom}(r)$), and
- (c) the cosine and tangent functions are continuous at a (provided that $a \in \text{Dom}(\tan)$).

Solution. (a) Every polynomial can be constructed from constant functions and the function f by a finite number of function multiplications and additions. (For example,

$$x^3 - 4x^2 + 5 = (x \times x \times x) + ((-4) \times x \times x) + 5$$

for all x in $\mathbb{R}$.) Since the constant functions and f are continuous at a , it follows from Proposition 3.1.1 that every polynomial is continuous at a .

(b) Every rational function r can be written in the form p/q , where p and q are polynomials. By the result of part (a), p and q are continuous at a . It follows from Proposition 3.1.1 that p/q is also continuous at a , provided that $q(a) \neq 0$.

(c) Recall that

$$\cos x = \sin(\pi/2 - x) \quad \forall x \in \mathbb{R}.$$

That is, $\cos x = h(g(x))$ where $h(x) = \sin x$ and $g(x) = \pi/2 - x$. Since the polynomial g is continuous at a and the sine function h is continuous everywhere, the cosine function must also be continuous at a by Proposition 3.1.2.

Finally, since

$$\tan x = \frac{\sin x}{\cos x},$$

the tangent function is also continuous at a (provided that $a \in \text{Dom}(\tan)$) by Proposition 3.1.1. $\square$

The next example justifies an assertion made in Section 2.5.2. We continue working under the assumption that the constant functions and the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = x$, are continuous everywhere.

Example 3.1.4. Suppose that $g : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at the point a and let $|g|$ denote the function given by

$$|g|(x) = |g(x)| \quad \forall x \in \mathbb{R}.$$

Show that $|g|$ is continuous at the point a .

Remark 3.1.5. Throughout this section we have taken for granted that the constant functions, the sine function and the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = x$, are continuous everywhere. To prove this, one needs a rigorous definition of the limit of a function at a point. Such a definition is similar in flavour to the definition for the limit of a function at infinity, but will not be presented in these notes.

Example 3.1.6. Suppose that a and b are real numbers and consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = \begin{cases} e^{ax+b} & \text{if } x \geq 0 \\ \cos x & \text{if } x < 0. \end{cases}$$

For what values of a and b will the function f be continuous at 0?

Solution. We require that

$$\lim_{x \rightarrow 0^-} f(x) = f(0) = \lim_{x \rightarrow 0^+} f(x).$$

Now

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \cos x = 1, \quad f(0) = e^b \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^{ax+b} = e^b.$$

So we require that $e^b = 1$, which implies that $b = 0$. There is no restriction on a . $\square$

3.2 Continuity on intervals

(Ref: SH10 §2.4)


In section 2.5.2, a definition for continuity at a point was given. We now extend this definition to define what is meant by continuity on an interval.

Definition 3.2.1. Suppose that f is a real-valued function defined on an open interval (a, b) . We say that f is a *continuous on* (a, b) if f is continuous at every point in the interval (a, b) .

Definition 3.2.2. Suppose that f is a real-valued function defined on a closed interval $[a, b]$. We say that

- (a) f is continuous at the endpoint a if $\lim_{x \rightarrow a^+} f(x) = f(a)$,
- (b) f is continuous at the endpoint b if $\lim_{x \rightarrow b^-} f(x) = f(b)$, and
- (c) f is continuous on the closed interval $[a, b]$ if f is continuous on the open interval (a, b) and at each of the endpoints a and b .

To illustrate these definitions, consider the functions f, g and h , whose graphs are shown below.




All three functions are defined on the interval $[a, b]$. We see that

- f is continuous on the open interval (a, b) and at the endpoint b ;
- g is continuous at the endpoints a and b but not continuous on the open interval (a, b) ; and
- h is continuous on the closed interval $[a, b]$ (and, by implication, on the open interval (a, b) and at both endpoints a and b).

3.3 The intermediate value theorem

(Ref: SH10 §§2.6, B.1)

The idea of this section is very simple. Suppose that a function f is continuous on some closed interval $[a, b]$ and that z is any real number that lies between $f(a)$ and $f(b)$, as illustrated below.




It is clear from the graph that one can find a real number c in $[a, b]$ such that $f(c) = z$. The *intermediate value theorem*, which is stated below, says that this can always be done for any function that is continuous on $[a, b]$.


Theorem 3.3.1 (The intermediate value theorem). *Suppose that f is continuous on the closed interval $[a, b]$. If z lies between $f(a)$ and $f(b)$ then there is at least one real number c in $[a, b]$ such that $f(c) = z$.*

The intermediate value theorem is proven using *the least upper bound property* of the real numbers. (It is this property that distinguishes the real numbers from the rational numbers.) We shall not examine the proof here.

We note three important points about the theorem. First, there may be more than one number c in $[a, b]$ that satisfies the conclusion of the intermediate value theorem. The diagram below illustrates this possibility.



Second, the condition in Theorem 3.3.1 that f is continuous cannot be relaxed. For example, consider the function $f : [a, b] \rightarrow \mathbb{R}$ whose graph is shown below.



Because of the discontinuity, z is not in the range of f , even though z lies between $f(a)$ and $f(b)$. That is, there is no real number c in $[a, b]$ such that $f(c) = z$.

Third, the intermediate value theorem is a theorem about continuous functions defined on the *real numbers*. There is no analogous theorem for continuous functions defined on the rational numbers. For example, consider the functions $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{Q} \rightarrow \mathbb{Q}$, given by

$$f(x) = x^2 - 2 \quad \text{and} \quad g(x) = x^2 - 2.$$

Both functions are continuous on their domains. Also,

$$f(0) = g(0) = -2 \quad \text{and} \quad f(2) = g(2) = 2.$$

However, while there is a number c in $[0, 2]$ such that $f(c) = 0$, there is no number c in $[0, 2]$ such that $g(c) = 0$ (since $\sqrt{2} \notin \text{Dom}(g)$).

One of the chief applications of the intermediate value theorem is determining whether an equation has a solution. Moreover, if a solution exists, then the theorem helps us determine an approximate location of this solution. This is particularly helpful when the equation cannot be solved by simple algebra.

Example 3.3.2. Show that the equation

$$e^{2x} = \sin x + 4 \quad (3.1)$$

has at least one positive solution.

Solution. Consider the function f given by

$$f(x) = e^{2x} - \sin x - 4.$$

Now

$$f(0) = e^0 - \sin 0 - 4 = -3$$

while

$$f(\pi/2) = e^\pi + 1 - 4 \approx 20.14.$$

Since f is continuous on the closed interval $[0, \pi/2]$ and 0 lies between $f(0)$ and $f(\pi/2)$, the intermediate value theorem implies that there is a real number c in $[0, \pi/2]$ such that $f(c) = 0$. That is,

$$e^{2c} = \sin c + 4$$

for some c in $[0, \pi/2]$. Moreover, one can check that $c \neq 0$. Hence the equation has at least one positive solution c . $\square$

Note that in equation (3.1), one cannot isolate x to find an explicit solution. However, now that it is known a solution exists in the interval $[0, \pi/2]$, one could use Newton's method to find a good approximation to this solution.

3.4 The maximum-minimum theorem

(Ref: SH10 §§2.6, B.2)


In many situations, it is desirable to know the maximum (or minimum) value of a quantity that varies over an interval of time. Examples include the maximum temperature in a 24 hour period or the minimum stock price over a seven day period. Depending on the quantity and the time interval, a maximum or minimum value may not be attained. The *maximum-minimum theorem* gives conditions under which a maximum and minimum is attained. To clarify discussion, we begin with a definition.

Definition 3.4.1. Suppose that f is defined on a closed interval $[a, b]$.

- (a) We say that a point c in $[a, b]$ is an *absolute minimum point* for f on $[a, b]$ if $f(c) \leq f(x)$ for all x in $[a, b]$. The corresponding value $f(c)$ is called the *absolute minimum value* of f on $[a, b]$. If f has an absolute minimum point on $[a, b]$ then we say that f *attains its minimum* on $[a, b]$.
- (b) We say that a point d in $[a, b]$ is an *absolute maximum point* for f on $[a, b]$ if $f(x) \leq f(d)$ for all x in $[a, b]$. The corresponding value $f(d)$ is called the *absolute maximum value* of f on $[a, b]$. If f has an absolute maximum point on $[a, b]$ then we say that f *attains its maximum* on $[a, b]$.

An absolute maximum point and an absolute minimum point are sometimes referred to as a *global maximum point* and a *global minimum point*.

Example 3.4.2. Consider the functions g and h , whose graphs are illustrated below.



The absolute minimum and maximum points of g and h on $[1, 5]$ are recorded in the following table.

	g	h
Absolute minimum points	none	3
Absolute minimum value	n.a.	1
Absolute maximum points	2, 4	none
Absolute maximum value	4	n.a.

The above example shows that a function $f : [a, b] \rightarrow \mathbb{R}$ need not have an absolute maximum point (or an absolute minimum point) on a closed interval $[a, b]$. However, if f is continuous on $[a, b]$ then such points always exist.

Theorem 3.4.3 (The maximum-minimum theorem). *If f is continuous on a closed interval $[a, b]$ then f attains its minimum and maximum on $[a, b]$. That is, there exist points c and d in $[a, b]$ such that*

$$f(c) \leq f(x) \leq f(d)$$

for all x in $[a, b]$.


As with the intermediate value theorem, the least upper bound property of the real numbers is used to prove the maximum-minimum theorem. We omit the proof.

Example 3.4.4. Suppose that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by the rule

$$f(x) = x^2 - 10x + 16.$$

Find the absolute maximum and absolute minimum values of f on the interval $[1, 10]$.

Solution. Since $f(x) = (x - 2)(x - 8)$, the function f has zeros at 2 and 8.




By symmetry, we know that the axis of the parabola is the line $x = 5$. Hence 5 is an absolute minimum point for f on $[1, 10]$. Since $f(5) = -9$, the absolute minimum value of f on $[1, 10]$ is -9 .

It is clear from the sketch of f that at least one of the endpoints of the interval $[1, 10]$ must be an absolute maximum point for f on $[1, 10]$. Now $f(1) = 7$ while $f(10) = 16$. So 10 is an absolute maximum point for f on $[1, 10]$ and the corresponding maximum value of f is 16. $\square$

Remark 3.4.5. While continuity on a closed interval guarantees the existence of at least one absolute maximum point and absolute minimum point, locating such points may not be easy. A systematic approach to finding maximum and minimum points will be developed in Chapters 4 and 5.

Remark 3.4.6. All the hypotheses stated in the maximum-minimum theorem must be satisfied for the conclusion of the theorem to be valid. If the interval is closed but the function is not continuous, then an absolute maximum (or minimum) point may not exist (see Example 3.4.2). The next example shows that if the function is continuous on the interval but the interval is not closed then an absolute maximum (or minimum) point may not exist. It is expected that students will know the precise hypotheses of the maximum-minimum theorem (and other important theorems, such as the intermediate value theorem) and know how to apply the theorem correctly.

Example 3.4.7. Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x + 1$. Since f is a polynomial, it is continuous on the open interval $(0, 1)$. However, f does not have an absolute maximum point on $(0, 1)$. To see why, consider the graph of f on $(0, 1)$.



As $x \rightarrow 1^-$, $f(x) \rightarrow 2$ but there is no real number d in $(0, 1)$ such that $f(d) = 2$. By a similar observation, f has no absolute minimum point on $(0, 1)$.

The final result of this chapter is a corollary of maximum-minimum theorem. We begin with a definition.

Definition 3.4.8. A function f is said to be *bounded* on an interval I if there is some positive number M such that $|f(x)| \leq M$ for all x in I .

In other words, f is bounded if the y -values of its graph lie between $-M$ and M for some positive number M . The *sine* and *cosine* functions are obvious examples of functions that are bounded on the whole of the real line.

Corollary 3.4.9. *If $f : [a, b] \rightarrow \mathbb{R}$ is continuous on $[a, b]$ then f is bounded on $[a, b]$.*

Proof. Suppose that f is continuous on $[a, b]$. Then the function $|f|$ is also continuous on $[a, b]$ (see Example 3.1.4). By the maximum-minimum theorem, $|f|$ attains its maximum on $[a, b]$. If M denotes the absolute maximum value of $|f|$ on $[a, b]$ then

$$|f(x)| \leq M \quad \forall x \in [a, b].$$

Hence f is bounded. □

Problems for Chapter 3

Problems 3.1 : Combining continuous functions and 3.2 : Continuity on intervals

- [R] Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = |x|$.
 - Show that f is continuous at 0.
 - Is f continuous everywhere? Give brief reasons for your answer.
- [R] Determine at which points each function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Give reasons.

$$\text{a) } f(x) = \begin{cases} e^{2x} & \text{if } x < 0 \\ \cos x & \text{if } x \geq 0 \end{cases} \quad \text{b) } f(x) = \begin{cases} e^{-2x} & \text{if } x < 0 \\ \sin x + 1 & \text{if } 0 \leq x \leq \pi/2 \\ 2x - \pi & \text{if } x > \pi/2 \end{cases}$$

- [R] Suppose that

$$f(x) = \begin{cases} \frac{x^2-16}{x-4} & \text{if } x \neq 4 \\ k & \text{if } x = 4, \end{cases}$$

where k is a real number. For which values of k (if any) will f be continuous everywhere?

- [H] Use the pinching theorem for limits to show that if f , g and h are three functions defined on an open interval I , such that
 - $f(x) \leq g(x) \leq h(x)$ for all $x \in I$,
 - $f(a) = g(a) = h(a)$ for some $a \in I$, and
 - f and h are continuous at a ,

then g is also continuous at a .

Problems 3.3 : The intermediate value theorem

- [R] Show that the function f , given by $f(x) = x^3 - 5x + 3$, has a zero in each of the intervals $[-3, -2]$, $[0, 1]$ and $[1, 2]$.
- [R] [V] Use the intermediate value theorem to show that the equation $e^x = 2 \cos x$ has at least one positive real solution.
- [H] Suppose that f is continuous on $[0, 1]$ and that $\text{Range}(f)$ is a subset of $[0, 1]$. By using $g(x) = f(x) - x$, prove that there is a real number c in $[0, 1]$ such that $f(c) = c$.
- [X] Suppose that f is a continuous function such that $f(0) = 1$ and $\lim_{x \rightarrow \infty} f(x) = -1$. Show that f has a zero somewhere in $(0, \infty)$.

Problems 3.4 : The maximum-minimum theorem

9. [R] In each case, determine whether or not f attains a maximum on the given interval. Give reasons for your answer.
- a) $f(x) = x^2 - 4$ on $[-3, 5]$ b) $f(x) = \left| \sin(e^x) + \frac{\ln x}{x^2 - 1} \right|$ on $[2, 4]$
- c) $f(x) = x^2 - 4$ on $(-3, 5)$ d) $f(x) = -(x^2 - 4)$ on $(-3, 5)$
10. [H] [V] Suppose that f is a continuous function on $\mathbb{R}$ and that $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow -\infty} f(x) = 0$.
- a) Give an example of such a function which has both a maximum value and a minimum value.
- b) Give an example of such a function which has a minimum value but no maximum value.
- c) [X] Show that if there is a real number ξ such that $f(\xi) > 0$ then f attains a maximum value on $\mathbb{R}$. [Note that the maximum-minimum theorem only applies to *finite* closed intervals $[a, b]$.]


Chapter 4

Differentiable functions

A cyclist accelerates from rest in such a way that her displacement s (in metres) from her starting position after t seconds is given by

$$s = t^2 \quad \forall t \in [0, 3].$$

The corresponding displacement–time graph for the first three seconds is shown below.



What is the cyclist's instantaneous velocity when $t = 1$?

At present, we do not have a formula for calculating instantaneous velocity. However, we can calculate the average velocity $\bar{v}$ of the cyclist over any time interval by using the formula

$$\bar{v} = \frac{\Delta s}{\Delta t},$$

where Δs denotes the change in displacement corresponding to a change Δt in time. For example, the average velocity over the time interval $[1, 1.5]$ is 2.5 metres per second, as calculated below:

$$\bar{v}_{[1,1.5]} = \frac{\Delta s}{\Delta t} = \frac{(1.5)^2 - (1)^2}{1.5 - 1} = 2.5. \quad (4.1)$$


Average velocities over small time intervals are recorded in the following table.

Time interval	Δt	Average velocity
$[1, 1.5]$	0.5	2.5
$[1, 1.4]$	0.4	2.4
$[1, 1.3]$	0.3	2.3
$[1, 1.2]$	0.2	2.2
$[1, 1.1]$	0.1	2.1
$[1, 1.01]$	0.01	2.01
$[1, 1.001]$	0.001	2.001

The table suggests that the average velocity approaches 2 meters per second as $\Delta t \rightarrow 0$. Hence it appears that instantaneous velocity of the cyclist when $t = 1$ is 2 meters per second.

One of the goals of this chapter is put these kinds of ideas on a *rigorous* footing in a *general* context (where they can be applied to understanding rates of change of any quantity, not just displacement). Hence we reframe these ideas in terms of functions and analytic geometry. More precisely, we shall now show that the problem of determining instantaneous velocity is equivalent to the problem of finding the gradient of a tangent line to the graph of a function.

To begin, the problem of calculating average velocity is equivalent to the problem of calculating the gradient of a secant to the graph of a function. (A secant is a straight line that intersects a curve at least twice.) For example, compare the calculation of average velocity in (4.1) with the calculation of the gradient of the secant shown in the following diagram.



In the same way, the above table of average velocities can be reinterpreted geometrically as a table of gradients corresponding to secants of the curve $s = t^2$.

Intersection points	Δt	Gradient of secant
$t = 1, t = 1.5$	0.5	2.5
$t = 1, t = 1.4$	0.4	2.4
$t = 1, t = 1.3$	0.3	2.3
$t = 1, t = 1.2$	0.2	2.2
$t = 1, t = 1.1$	0.1	2.1

The secants used in the table are illustrated over the page in Figure 4.1. The figure suggests that as $\Delta t \rightarrow 0$, the gradient of the corresponding secant approaches the gradient of the tangent line to the curve. Thus the problem of determining instantaneous velocity is equivalent to the problem of determining the gradient of a tangent to the graph of a function. It is to this general problem that we now turn.


4.1 Gradients of tangents and derivatives

(Ref: SH10 §3.1)

Suppose that a function f has a ‘smooth’ graph. In this section, we give a general approach to calculating the gradient of tangents to f . This leads the concept of a gradient function, also known as a *derivative* (since it is derived from f), whose values give the gradient of the tangent to f at any point. We begin with an example.

Example 4.1.1. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = x^3$. Find the gradient of the tangent to the graph of f when $x = 2$.

Solution. The tangent to the graph of f passes through the point $P(2, 2^3)$. We will approximate the gradient of the tangent with the gradient of a secant passing through the points $P(2, 2^3)$ and $Q(2 + h, (2 + h)^3)$, where h is a very small real number. (See the illustration below for the case when $h > 0$.)



Using the ‘rise over run’ formula, the gradient of the secant is

$$\frac{(2 + h)^3 - 2^3}{h}.$$

Now, as h approaches 0, the point Q moves along the curve towards P , and hence the gradient of




Figure 4.1: Using secants to determine the gradient of a tangent.

the secant approaches the gradient of the tangent. In other words,

$$\begin{aligned}
 (\text{gradient of tangent}) &= \lim_{h \rightarrow 0} \frac{(2+h)^3 - 2^3}{h} \\
 &= \lim_{h \rightarrow 0} \frac{8 + 12h + 6h^2 + h^3 - 8}{h} \\
 &= \lim_{h \rightarrow 0} \frac{12h + 6h^2 + h^3}{h} \\
 &= \lim_{h \rightarrow 0} 12 + 6h + h^2 \\
 &= 12.
 \end{aligned}$$

Thus the gradient of the tangent to f at the point 2 is 12. This gives a quantitative description of the rate of change of f at the point 2. $\square$

There is nothing special about the point 2 in the previous example; we could calculate the gradient of the tangent to f at any point x . The calculation would go something like this:

$$\begin{aligned}
 (\text{gradient of tangent at } x) &= \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h} \\
 &= \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2}{h} \\
 &= \lim_{h \rightarrow 0} 3x^2 + 3xh \\
 &= 3x^2.
 \end{aligned}$$

So the gradient of the tangent to f when $x = 5$ is 75 (since $3 \times 5^2 = 75$), while the gradient of the tangent when $x = -4$ is 48. The gradient function of f , denoted by f' and given by

$$f'(x) = 3x^2 \quad \forall x \in \mathbb{R},$$

is called the *derivative* of f .

We can repeat the same procedure for functions other than $f(x) = x^3$. The gradient function (or derivative) may be calculated using the limit given in the next definition.

Definition 4.1.2. Suppose that f is defined on some open interval containing the point x . We say that f is *differentiable* at x if

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. If the limit exists, we denote it by $f'(x)$. We call $f'(x)$ the derivative of f at x .

Other notation for $f'(x)$ includes $\frac{df}{dx}(x)$ and $\frac{d}{dx}f(x)$. If $y = f(x)$ then the derivative is often denoted by y' or $\frac{dy}{dx}$. The ratio

$$\frac{f(x+h) - f(x)}{h}$$

is called the *difference quotient* for f at the point x .

Remark 4.1.3. The notation $\frac{dy}{dx}$ originates from the gradient formula $\frac{\Delta y}{\Delta x}$, where Δy represents a change in y corresponding to a small change Δx in x . (The symbol Δ represents the Greek letter ‘delta’, which is equivalent to our capital ‘D’.) Some calculus texts prefer the use of delta notation and write the difference quotient for a function f as

$$\frac{f(x + \Delta x) - f(x)}{\Delta x}.$$

Example 4.1.4. Suppose that $f : [0, \infty) \rightarrow \mathbb{R}$ is given by $f(x) = \sqrt{x}$. Find the equation of the tangent to f when $x = 16$.

Solution. First we shall calculate the derivative of f (if it exists) at 16. Note that, in the working below, the limit as h approaches 0 is difficult to evaluate immediately, so we use a standard trick (see, for example, Subsection 2.1.4):

$$\begin{aligned} \frac{f(16+h) - f(16)}{h} &= \frac{\sqrt{16+h} - \sqrt{16}}{h} \\ &= \frac{\sqrt{16+h} - \sqrt{16}}{h} \times \frac{\sqrt{16+h} + \sqrt{16}}{\sqrt{16+h} + \sqrt{16}} \\ &= \frac{16+h-16}{h(\sqrt{16+h} + \sqrt{16})} \\ &= \frac{h}{h(\sqrt{16+h} + \sqrt{16})} \\ &= \frac{1}{\sqrt{16+h} + \sqrt{16}} \\ &\rightarrow \frac{1}{8} \end{aligned}$$

as $h \rightarrow 0$. So the gradient of the tangent is $1/8$. Using the point-gradient formula, the equation of the tangent is

$$\begin{aligned} y - y_1 &= m(x - x_1) \\ y - 4 &= \frac{1}{8}(x - 16) \\ 8y - x - 16 &= 0. \end{aligned}$$

□

Example 4.1.5. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = |x|$. Determine whether or not f is differentiable at the point 0.

Solution. We calculate the difference quotient of f at 0:

$$\begin{aligned} \frac{f(0+h) - f(0)}{h} &= \frac{|h|}{h} \\ &= \begin{cases} \frac{h}{h} & \text{if } h > 0 \\ \frac{-h}{h} & \text{if } h < 0. \end{cases} \\ &= \begin{cases} 1 & \text{if } h > 0 \\ -1 & \text{if } h < 0. \end{cases} \end{aligned}$$

Because of the ‘split formula,’ we must consider the left- and right-hand limits as h approaches 0 separately. Now


$$\lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = -1$$

while

$$\lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = 1.$$

Since the left- and right-hand limits are different, the two-sided limit doesn’t exist and hence f is *not* differentiable at 0.

Note that this conclusion makes sense geometrically, since the graph of f has a ‘vertex’ at 0 and there is no unique tangent that touches the graph at this point.



□

Example 4.1.6. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0. \end{cases}$$

Determine whether or not f is differentiable at 0.

4.2 Rules for differentiation

(Ref: SH10 §§3.2, 3.5, 3.6)

So far we have calculated derivatives working directly from the difference quotient. In this section we see that there are fast ways of calculating derivatives without referring to the difference quotient and limits at all. This fast method is part of what is known as ‘the calculus.’ We begin with the derivatives of some standard functions, and then show how derivatives of other functions are obtained by applying some simple rules.

The following table gives the derivatives for some standard functions.

$f(x)$	$f'(x)$
C , where C is a constant	0
x^n , where n is a positive integer	nx^{n-1}
$\sin x$	$\cos x$
e^x	e^x

With these basic functions, it is possible to construct many other functions via function addition, subtraction, multiplication and division. The next theorem describes what happens to the resulting derivatives.

Theorem 4.2.1 (Rules for differentiation). *Suppose that f and g are differentiable at x . Then $f + g$, $f - g$ and fg are differentiable at x . If $g(x) \neq 0$ then $\frac{f}{g}$ is also differentiable at x . Moreover,*

$$(i) \quad (f + g)'(x) = f'(x) + g'(x),$$

$$(ii) \quad (C.f)'(x) = C.f'(x), \text{ where } C \text{ is a constant,}$$

$$(iii) \quad (fg)'(x) = f'(x)g(x) + f(x)g'(x), \text{ and}$$

$$(iv) \quad \left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}, \text{ provided that } g(x) \neq 0.$$

Rules (iii) and (iv) are usually called the product and quotient rules. A familiar way of expressing these rules is

$$\frac{d(uv)}{dx} = v \frac{du}{dx} + u \frac{dv}{dx}$$

and

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2},$$

where u and v are both functions of x .

Given two functions f and g , a new function $f \circ g$ may be constructed by function composition. The next theorem describes the derivative of this new function.

Theorem 4.2.2 (The chain rule). *Suppose that g is differentiable at the point x and f is differentiable at the point $g(x)$. Then $f \circ g$ is differentiable at x and*

$$(f \circ g)'(x) = f'(g(x))g'(x). \quad (4.2)$$

You are probably more familiar with the chain rule expressed in the following way: if $y = f(u)$ and $u = g(x)$ then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}. \quad (4.3)$$

While formula (4.3) looks simpler than (4.2), it does not specify the relationship between the variables or the points at which each derivative is to be evaluated. It should therefore be used with care.

The next example illustrates how the chain rule formula (4.2) is applied.

Example 4.2.3. Suppose that $y = (\sin x + 1)^7$. Find the derivative of y at the point x .

Solution. Suppose that $f(x) = x^7$ and $g(x) = \sin x + 1$. Then $f'(x) = 7x^6$ and $g'(x) = \cos x$. Then $y = (f \circ g)(x)$. By the chain rule, the derivative of y at x is given by

$$\begin{aligned} (f \circ g)'(x) &= f'(g(x))g'(x) \\ &= 7(g(x))^6 \times \cos x \\ &= 7(\sin x + 1)^6 \cos x. \end{aligned}$$

□

As mentioned at the beginning of this section, derivatives of many functions can be obtained from those of a few basic functions by the application of simple rules. The next example illustrates this point.

Example 4.2.4. Show that

$$(a) \quad \frac{d}{dx}(x^m) = mx^{m-1}, \text{ where } m \text{ is an integer;}$$

$$(b) \quad \frac{d}{dx}(\cos x) = -\sin x; \text{ and}$$

$$(a) \quad \frac{d}{dx}(\tan x) = \sec^2 x.$$

4.3 Proofs of results in Section 4.2

In Section 4.2 we stated two theorems and gave a table of derivatives. In this section we prove some of these results. We begin with the derivative of the sine function, using the standard result that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1. \quad (4.4)$$

For a geometric argument for this limit, see Question 14 from the problems for Chapter 2.

Proposition 4.3.1. *If $f(x) = \sin x$ then $f'(x) = \cos x$ for all real x .*

Proof. The difference quotient for the sine function and the angle sum formula for sine gives

$$\begin{aligned} \frac{\sin(x+h) - \sin x}{h} &= \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\ &= \sin x \frac{\cos h - 1}{h} + \cos x \frac{\sin h}{h}. \end{aligned} \quad (4.5)$$

Now $\cos^2 h + \sin^2 h = 1$ so

$$\begin{aligned} \frac{\cos h - 1}{h} &= \frac{\cos^2 h - 1}{h(\cos h + 1)} \\ &= \frac{-\sin^2 h}{h(\cos h + 1)} \\ &= -\frac{\sin h}{h} \frac{\sin h}{\cos h + 1}, \end{aligned}$$

and by (4.4) we have

$$-\frac{\sin h}{h} \frac{\sin h}{\cos h + 1} \rightarrow -1 \times 0 \quad \text{as } h \rightarrow 0.$$

If we combine this with (4.5) then

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} &= \sin x \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \times 0 + \cos x \times 1 \\ &= \cos x. \end{aligned}$$

Thus $f'(x) = \cos x$. □

The derivative of e^x will be dealt with in Chapter 9. The derivative of constant functions can be easily computed from Definition 4.1.2.

Suppose that f is given by $f(x) = x^n$, where n is a positive integer. The derivative of f can be computed using the same method as Example 4.1.1, with the assistance of the binomial theorem. Alternatively, one can use mathematical induction and the product rule for differentiation.

Proof of the product rule for differentiation. Suppose that f and g are differentiable at the point x . The difference quotient of fg at x gives

$$\begin{aligned} & \frac{(fg)(x+h) - (fg)(x)}{h} \\ &= \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \frac{f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x)}{h} \\ &= g(x+h) \frac{f(x+h) - f(x)}{h} + f(x) \frac{g(x+h) - g(x)}{h}. \end{aligned} \quad (4.6)$$

Now g is differentiable at x , so g is also continuous at x (see Theorem 4.5.1). Hence $g(x+h) \rightarrow g(x)$ as $h \rightarrow 0$. Therefore (4.6) implies that

$$\frac{(fg)(x+h) - (fg)(x)}{h} \rightarrow g(x)f'(x) + f(x)g'(x)$$

as $h \rightarrow 0$. □

Proofs of other differentiation rules are found in most undergraduate calculus textbooks.

4.4 Implicit differentiation

(Ref: SH10 §3.7)

In Section 1.8 we introduced functions that are defined by an equation relating the variables x and y . Using the chain rule, one can calculate the derivative of such functions without first expressing y explicitly as a function of x . We begin with a simple example to illustrate the principle. The point to keep in mind is that y is a function of x .

Example 4.4.1. Suppose that y is a function of x , and that y and x are related by the formula

$$x^2 + y^2 = 1.$$

Calculate $\frac{dy}{dx}$.

Solution. We begin with the equation

$$x^2 + y^2 = 1$$

and differentiate both sides with respect to x to obtain

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = \frac{d}{dx}(1). \quad (4.7)$$

Now clearly, $\frac{d}{dx}(x^2) = 2x$ and $\frac{d}{dx}(1) = 0$. To simplify the central term of (4.7), we regard y as a function of x and suppose that $g(y) = y^2$. The chain rule gives

$$\begin{aligned}\frac{d}{dx}(y^2) &= \frac{dg}{dx} \\ &= \frac{dg}{dy} \frac{dy}{dx} \\ &= 2y \frac{dy}{dx}.\end{aligned}$$

So continuing from (4.7) we have

$$2x + 2y \frac{dy}{dx} = 0.$$

By isolating $\frac{dy}{dx}$ we obtain

$$\begin{aligned}\frac{dy}{dx} &= \frac{-2x}{2y} \\ &= -\frac{x}{y}\end{aligned}$$

as required.

(We note in passing that this answer coincides with the derivative obtained by first expressing y as a function of x and then differentiating. In particular, if

$$y = (1 - x^2)^{1/2}$$

then

$$\frac{dy}{dx} = \frac{1}{2}(1 - x^2)^{-1/2}(-2x) = -\frac{x}{(1 - x^2)^{1/2}} = -\frac{x}{y}$$

as above.) □

In general, if g is a function of y and y is a function of x then

$$\frac{d}{dx}g(y) = g'(y) \frac{dy}{dx} \tag{4.8}$$

by the chain rule. The use of this formula is illustrated below.

Example 4.4.2. Suppose that y is a function of x , implicitly related by the equation

$$y^4 + x^3 - x^2 e^{3y} = 4. \tag{4.9}$$

Find the equation of the tangent to the corresponding curve at the point where $(x, y) = (2, 0)$.

In Section 4.2 we saw that $\frac{d}{dx}x^n = nx^{n-1}$ whenever n is an integer. A similar formula holds when n is any rational number. This can be easily proved using implicit differentiation.

Proposition 4.4.3. Suppose that q is a rational number. Then

$$\frac{d}{dx}x^q = qx^{q-1}.$$

Proof. Since q is a rational number, there are integers m and n such that $q = \frac{m}{n}$ and $n \neq 0$. Now if $y = x^q$ then $y = x^{m/n}$ and taking the n th power of both sides gives

$$y^n = x^m.$$

Differentiating both sides with respect to x yields

$$ny^{n-1} \frac{dy}{dx} = mx^{m-1}.$$

Hence

$$\begin{aligned} \frac{dy}{dx} &= \frac{mx^{m-1}}{ny^{n-1}} \\ &= \frac{m}{n} \frac{x^{m-1}}{x^{q(n-1)}} \\ &= qx^{(m-1)-qn+q} \\ &= qx^{q-1} \end{aligned}$$

as required. □

4.5 Differentiation, continuity and split functions

(Ref: SH10 §3.1)

Not every function that is continuous at a point a is differentiable at a . (Consider, for example, the function f , given by $f(x) = |x|$, at the point 0.) However, we do have the following result.

Theorem 4.5.1. *If f is differentiable at a then f is continuous at a .*

Proof. Suppose that f is differentiable at the point a . Then

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists. To show that f is continuous at a , we need to show that

$$\lim_{x \rightarrow a} f(x) = f(a),$$

or equivalently, that

$$\lim_{h \rightarrow 0} f(a+h) = f(a).$$

Now

$$\begin{aligned} \lim_{h \rightarrow 0} f(a+h) - f(a) &= \lim_{h \rightarrow 0} \left(\frac{f(a+h) - f(a)}{h} \times h \right) \\ &= \left(\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \right) \times \lim_{h \rightarrow 0} h \\ &= f'(a) \times 0 \\ &= 0. \end{aligned} \tag{4.10}$$

(Note that we can separate the limits in (4.10) because we know that the limit of the difference quotient exists.) Hence


$$\lim_{h \rightarrow 0} f(a+h) = f(a),$$

proving the theorem. □

The *contrapositive* form of the theorem is stated below.

Corollary 4.5.2. *If f is not continuous at a then it is not differentiable at a .*

Remark 4.5.3. Differentiability is a much stronger property than continuity. While every differentiable function is continuous, there exist functions that are continuous everywhere but differentiable nowhere. In fact, functions that are differentiable everywhere are a very rare breed, even among the continuous functions. An example of a function that is continuous everywhere but differentiable nowhere is the *Weierstrass function* whose graph is shown below.



The next function studied is an example of a ‘split function’. Whether or not the function is differentiable at the ‘split point’ can be determined by calculating left- and right-hand limits of the difference quotient.

Example 4.5.4. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} \sin x & \text{if } x \geq 0 \\ x^2 + bx + c & \text{if } x < 0, \end{cases}$$

where b and c are real numbers. Find all possible values of b and c such that f is (a) continuous at 0 and (b) differentiable at 0.

Solution. (a) For f to be continuous at 0, we require that

$$\lim_{x \rightarrow 0^-} f(x) = f(0) = \lim_{x \rightarrow 0^+} f(x).$$

Now $f(0) = \sin 0 = 0$ while

$$\lim_{x \rightarrow 0^-} f(x) = c \quad \text{and} \quad \lim_{x \rightarrow 0^+} f(x) = 0.$$

Hence f is continuous if and only if $c = 0$ and $b \in \mathbb{R}$.

(b) If f is differentiable at 0 then it is continuous at 0 and hence $c = 0$ by part (a). Also, for differentiability we require that

$$\lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h}.$$


Now,

$$\lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{\sin h - \sin 0}{h} = \lim_{h \rightarrow 0^+} \frac{\sin h}{h} = 1$$

while

$$\lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{h^2 + bh - 0}{h} = \lim_{h \rightarrow 0^-} h + b = b.$$

Hence we require that $b = 1$. So f is differentiable at 0 if and only if $b = 1$ and $c = 0$. The graph of the corresponding function is sketched below.



□

An important application of split functions is the theory of splines. A *spline* is a function defined piecewise by polynomials. They are frequently used in computer graphics and elsewhere because of their simplicity and their capacity to approximate complex curves. The following is an example of a spline.

Example 4.5.5.

$$f(x) = \begin{cases} (x-1)^2 - 1 & \text{if } -2 \leq x < 0 \\ 1 - (x-1)^2 & \text{if } 0 \leq x \leq 2. \end{cases}$$

In this example, each of the functions $p(x) = (x-1)^2 - 1$ and $q(x) = 1 - (x-1)^2$ have had their domains restricted and then ‘glued’ together at the point 0.

If the constituent functions are continuous and differentiable in some interval containing the point a , then the following theorem tells us when the corresponding split function is differentiable.

Theorem 4.5.6. *Suppose a is a fixed real number and the function f is defined by*

$$f(x) = \begin{cases} p(x) & \text{if } x \geq a \\ q(x) & \text{if } x < a, \end{cases}$$

where $p(x)$ and $q(x)$ are continuous and differentiable in some open interval containing a . Then if f is continuous at a and $p'(a) = q'(a)$, then f is differentiable at $x = a$.

This theorem in particular can be applied to splines since the constituent functions are polynomials.

Example 4.5.7.

$$f(x) = \begin{cases} \sin x & \text{if } x \geq 0 \\ x^2 + x & \text{if } x < 0. \end{cases}$$


The functions $\sin x$ and $x^2 + x$ are continuous and differentiable everywhere. We saw in Example 4.5.4 that f is continuous at $x = 0$. Also, with $p(x) = \sin x$ and $q(x) = x^2 + x$, we have $p'(0) = \cos 0 = 1$ and $q'(0) = 1$ so f is differentiable at $x = 0$.

4.6 Derivatives and function approximation

(Ref: SH10 §4.11)

The last three sections of this chapter are devoted to some applications of the derivative.

Suppose that a function f is differentiable at a . Consider its graph and the tangent to f at a , shown below.



The equation of the tangent to f at a may be calculated using the point-gradient formula:

$$\begin{aligned} y - f(a) &= f'(a)(x - a) \\ y &= f'(a)(x - a) + f(a). \end{aligned}$$

Observe in the above diagram that the tangent line is close to the graph of f for all points that are close to a . In other words,

$$f(x) \approx f'(a)(x - a) + f(a) \quad (4.11)$$

when x is close to a .

Formula (4.11) reflects the fact that every differentiable function can be *locally* approximated by a linear function. We use this principle in the following example.

Example 4.6.1. Estimate $\sqrt{9.001}$ without using a calculator.

Solution. Suppose that $f(x) = \sqrt{x}$. Since 9.001 is close to 9 and f is a continuous function, we could use the approximation

$$\sqrt{9.001} = f(9.001) \approx f(9) = 3. \quad (4.12)$$

However, we can do better than this. We shall approximate f with a linear function at the point 9 using (4.11). Since $f'(9) = 1/6$ we have

$$f(x) \approx (x - 9)/6 + 3$$

whenever x is close to 9. When $x = 9.001$ this gives

$$\sqrt{9.001} = f(9.001) \approx (9.001 - 9)/6 + 3 = 3 + 1/6000. \quad (4.13)$$

The table below gives the error involved in each approximation.

Approximation	Error
Approximation (4.12)	1.667^{-4}
Approximation (4.13)	4.637^{-9}

Clearly (4.13) is the superior approximation. □

4.7 Derivatives and rates of change

(Ref: SH10 §3.4)

Many physical processes involve quantities (such as temperature, volume, concentration, velocity) that change with time. If Q is a quantity that varies with time, then the derivative $\frac{dQ}{dt}$ gives the rate of change of that quantity with respect to time. The chain rule is a major tool for answering questions about rates of quantities which are related.

Example 4.7.1. A spherical balloon is being inflated and its radius is increasing at a constant rate of 6 mm/sec. At what rate is its volume increasing when the radius of the balloon is 20 mm?

Solution. Suppose that $V(t)$ is the volume of the balloon and $r(t)$ is its radius at time t . We are told that $\frac{dr}{dt} = 6$ and we need to find $\frac{dV}{dt}$ when $r = 20$. By the chain rule,

$$\frac{dV}{dt} = \frac{dV}{dr} \frac{dr}{dt}.$$

Now $V = \frac{4}{3}\pi r^3$ so $\frac{dV}{dr} = 4\pi r^2$. Hence

$$\frac{dV}{dt} = 4\pi r^2 \times 6.$$

When $r = 20$,

$$\frac{dV}{dt} = 24\pi(20)^2 = 9600\pi.$$

So the volume is increasing at a rate of 9600π mm³/sec when the radius is 20 mm. □

The above example illustrates an approach to solving such problems.

1. Define variables for the quantities involved.
2. Write down what is known in terms of these variables and their derivatives.
3. Write down what you need to find in terms of the these variables and their derivatives.
4. Write down anything else you know that relates the variables (for example, a volume or area formula).
5. Use the chain rule (or implicit differentiation) to find the relevant derivative.

Sometimes drawing a diagram helps in the first few steps.

The next example follows this procedure.


Example 4.7.2. A point P moves to the right along the positive x -axis at a constant rate of 5 cm per second, and a point Q moves up the positive y -axis at a constant rate of 10 cm per second. How fast is the distance between them changing when $OP = 30$ cm and $OQ = 40$ cm?

4.8 Local maximum, local minimum and stationary points

(Ref: SH10 §4.3)

In Example 3.4.4, we determined the absolute maximum and minimum points of a continuous function over a closed interval. While the existence of these extreme points is given by the max-min theorem, the theorem provides no systematic way of locating these points. In this section we begin to develop a systematic approach to locating maxima and minima. A complete approach will be presented in the next chapter.

We begin by defining what is meant by a *local minimum point* c . This is a point such that for all x sufficiently close to c , $f(c) \leq f(x)$. A more precise statement is below.



Local minimum point

Definition 4.8.1. Suppose that f is defined on some interval I . We say that a point c in I is a *local minimum point* if there is a positive number h such that $f(c) \leq f(x)$ whenever $x \in (c-h, c+h)$ and $x \in I$. We say that a point d in (a, b) is a *local maximum point* if there is a positive number h such that $f(x) \leq f(d)$ whenever $x \in (d-h, d+h)$ and $x \in I$.

The following theorem will be familiar from high school.

Theorem 4.8.2. Suppose that f is defined on (a, b) and has a local maximum or minimum point at c for some c in (a, b) . If f is differentiable at c then $f'(c) = 0$.

Proof. Suppose that f is a local maximum at c . To show that $f'(c) = 0$ we consider the difference quotient

$$\frac{f(c+h) - f(c)}{h}.$$

Since f is a maximum at c , $f(c+h) \leq f(c)$ for h sufficiently close to 0. So if h is sufficiently small and positive,

$$\frac{f(c+h) - f(c)}{h} \leq 0,$$

while if h is sufficiently small and negative,

$$\frac{f(c+h) - f(c)}{h} \geq 0.$$

That is,

$$\lim_{h \rightarrow 0^+} \frac{f(c+h) - f(c)}{h} \leq 0$$

while

$$\lim_{h \rightarrow 0^-} \frac{f(c+h) - f(c)}{h} \geq 0.$$

Hence

$$\lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h} = 0$$

which is equivalent to saying that $f'(c) = 0$.

The proof when c is a local minimum is similar and will be omitted. $\square$

The set of points where the derivative of a function is zero is important. In particular, tangents to the graph of the function at these points are horizontal lines. This motivates the following definition.

Definition 4.8.3. If a function f is differentiable at a point c and $f'(c) = 0$ then c is called a *stationary point* of f .


Example 4.8.4. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = 4x^5 - 5x^4 - 40x^3 - 2$. Find all the stationary points of f .

Solution. We begin by differentiating f and factorising the derivative:

$$\begin{aligned} f'(x) &= 20x^4 - 20x^3 - 120x^2 \\ &= 20x^2(x^2 - x - 6) \\ &= 20x^2(x+2)(x-3). \end{aligned}$$

Hence the solutions to $f'(x) = 0$ are $x = -2, 0, 3$. So the points $-2, 0$ and 3 are the stationary points of f . $\square$

The graph of the function f appearing in Example 4.8.4 is shown below.



It is clear that -2 is a local maximum point of f and that 3 is a local minimum point of f . However, 0 is neither a local minimum point nor local maximum point, and none of these points are global maximum points or global minimum points. So while the local maxima and minima of a function occur where f' is zero, it does not follow that any point c satisfying $f'(c) = 0$ is a local maximum point or local minimum point of f .

Tools for identifying whether a stationary point is a local maximum, local minimum or neither are developed in the next chapter using an important result called the *mean value theorem*.

4.9 Maple notes

The Maple `diff` command is used to compute derivatives: `diff(f(x),x);` calculates $\frac{df}{dx}$ and `diff(f(x),x$n);` calculates $\frac{d^n f}{dx^n}$. For example,

```
> # Note Maple use of csc to denote function we usually call cosec
```

```
> diff(sqrt(csc(x^12-53*x^5-1)),x);
```

$$-\frac{1}{2} \sqrt{\csc(x^{12} - 53x^5 - 1)} \cot(x^{12} - 53x^5 - 1) (12x^{11} - 265x^4)$$

```
> diff(x^3*exp(5*x),x);
```

$$3x^2e^{5x} + 5x^3e^{5x}$$

```
> diff(%,x);
```

$$6xe^{5x} + 30x^2e^{5x} + 25x^3e^{5x}$$

```
> f2ndDeriv:=diff(x^3*exp(5*x),x$2);
```

$$f2ndDeriv := 6xe^{5x} + 30x^2e^{5x} + 25x^3e^{5x}$$

Problems for Chapter 4

Problems 4.1 : Gradients of tangents and derivatives

1. [R] Using the definition of the derivative, show that:

- a) if $f(x) = x^2$ then $f'(x) = 2x$;
- b) if $f(x) = x^3$ then $f'(x) = 3x^2$;
- c) if $f(x) = \frac{1}{x}$ then $f'(x) = \frac{-1}{x^2}$;
- d) [H] if $f(x) = \sqrt{x}$ then $f'(x) = \frac{1}{2\sqrt{x}}$.

Problems 4.2 : Rules for differentiation

2. [R] Find the derivative in each case.

- a) $f(x) = 5(x^4 + 3x^7)$
- b) $g(x) = (x^4 - 2x)(4x^2 + 2x + 5)$
- c) $h(y) = \frac{y^2}{y^3 + 8}$
- d) $f(x) = x(x^2 - 4)^{1/2}$
- e) $f(t) = t/\sqrt{t^2 - 4}$
- f) $g(y) = \sin 3y - 3 \cos^2 2y$
- g) $g(x) = x^4 e^{-x}$
- h) $f(x) = (x^2 + 1) \ln \sqrt{x^3 + 1}$
- i) $f(x) = \ln(e^{\tan x})$
- j) $f(x) = \ln(\cos x)$

3. [H] Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x|x|$ for all x in $\mathbb{R}$.

- a) If it exists, evaluate $\lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h}$.
- b) If it exists, evaluate $\lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h}$.
- c) State the value of $f'(0)$ or explain why f is not differentiable at 0.

4. [R] [V] Determine at which points each function f is (i) differentiable; (ii) continuous.

- a) $f(x) = |x|$
- b) $f(x) = \begin{cases} \sin x & \text{if } x \leq 0 \\ x & \text{if } x > 0 \end{cases}$
- c) $f(x) = \frac{x^3 - 6x + 4}{x^2 + 4x + 4}$

5. [R] Sketch the graph of f , where $f(x) = x^{1/3}$. Is f differentiable at 0? Give reasons.

6. [X] Prove that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0, \end{cases}$$

is continuous and differentiable everywhere, but that f' is not continuous at 0.

7. [X] The function f is differentiable at a . Find

$$\lim_{h \rightarrow 0} \frac{f(a + ph) - f(a - ph)}{h}.$$

8. [R] (*An exercise on notation.*) Suppose that $f(x) = x + \cos 2x$. Write down
- a) $f(x + 17\pi)$ b) $f'(x + n\pi)$ c) $f(2 - x^2)$
- d) $f'(2 - x^2)$ e) $\frac{d}{dx} f(2 - x^2)$.

Problems 4.4 : Implicit differentiation

9. [R] Find $\frac{dy}{dx}$ in terms of x and y if
- a) $x^3 + y^3 = xy$ b) $x^2 - \sqrt{xy} + y^2 = 6$.
10. [R] Find $\frac{dy}{dx}$ for the curve $x^4 + y^4 = 16$. Sketch the graph of the curve.
11. [R] [V] Find the equation of the line tangent to the curve $x^3 + y^3 = 3(x + y)$ at the point $(1, 2)$.

Problems 4.5 : Differentiation, continuity and split functions

12. [R] Suppose that a and b are real numbers. Find all values of a and b (if any) such that the functions f and g , given by

$$\text{a) } f(x) = \begin{cases} ax + b & \text{if } x < 0 \\ \sin x & \text{if } x \geq 0 \end{cases} \quad \text{and} \quad \text{b) } g(x) = \begin{cases} ax + b & \text{if } x < 0 \\ e^{2x} & \text{if } x \geq 0 \end{cases},$$

are (i) continuous at 0 and (ii) differentiable at 0.

13. [H] The function $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} \sqrt{x} \sin \sqrt{x} & \text{if } x \geq 0 \\ ax + b & \text{if } x < 0, \end{cases}$$


where a and b are real numbers. Find all values of a and b (if any) such that f is differentiable at 0.

Problems 4.6 : Derivatives and function approximation

14. [R] [V] Suppose that $f(x) = \sqrt[3]{x}$.
- a) Without using a calculator, give a rough estimate of $f(8.01)$.
- b) i) Find the equation of the tangent to f at the point $(8, 2)$.
 ii) Use your answer to part (i) to find a different approximation for $f(8.01)$.
- c) Using a calculator, determine the error for the approximation in (a) and in (b). Which approximation is better?

Problems 4.7 : Derivatives and rates of change

15. **[R]** At a certain instant the side length of an equilateral triangle is a cm and this length is increasing at r cm/s. How fast, in cm^2/s , is the area increasing?
16. **[R]** **[V]** A 5 m ladder is leaning against a vertical wall. Suppose that the bottom of the ladder is being pulled away from the wall at a rate of 1 m/s. How fast is the area of the triangle underneath the ladder changing at the instant that the top of the ladder is 4 m from the floor?
17. **[R]** A spherical balloon is to be filled with water so that there is a constant increase in the rate of its surface area of $3 \text{ cm}^2/\text{s}$.
(The surface area A and volume V of a sphere of radius r is given by $A = 4\pi r^2$ and $V = \frac{4}{3}\pi r^3$.)
- Find the rate of increase in the radius when the radius is 3 cm.
 - Find the volume when the volume is increasing at a rate of $10 \text{ cm}^3/\text{s}$.
18. **[R]**
- A container in the shape of a right circular cone, of semi-vertical angle $\tan^{-1}(\frac{1}{2})$, is placed vertex downwards with its axis vertical.



Water is poured in at the rate of $10 \text{ mm}^3/\text{s}$. Find the rate at which the depth, h mm, is increasing when the depth of water in the cone is 50 mm.

- [H]** The cone is filled to a depth of 100 mm and pouring is then stopped. A hole is then opened at the vertex of the cone and water flows out of the hole at the rate of $50\pi\sqrt{h} \text{ mm}^3$ per second, where h is the depth at time t . Show that it takes 200 seconds to empty the cone.

Chapter 5

The mean value theorem and applications

The mean value theorem is one of the most important results for establishing the theoretical structure of calculus. Several results that you are familiar with from high school calculus are based on the mean value theorem. Applications of the mean value theorem include

- identifying where a function is increasing or decreasing,
- identifying different types of stationary points,
- determining how many zeros a polynomial has,
- evaluating limits which are indeterminate forms of type $\frac{\infty}{\infty}$ and $\frac{0}{0}$,
- proving useful inequalities and
- estimating errors in approximations.

We begin the chapter by introducing the theorem.

5.1 The mean value theorem

(Ref: SH10 §4.1)

The basic idea behind the mean value theorem is straightforward. Consider a differentiable function f and choose an interval $[a, b]$. We construct a chord AB joining the points $(a, f(a))$ and $(b, f(b))$ as shown in Figure 5.1. Because the function is continuous and differentiable, there is a point c in (a, b) such that tangent to the graph of f at c lies parallel to the chord AB .

Since the tangent and the chord are parallel, their gradients are equal. The gradient of the tangent at c is $f'(c)$ while the gradient of the chord is given by

$$\frac{f(b) - f(a)}{b - a}$$

(using the ‘rise over run’ formula). Hence

$$\frac{f(b) - f(a)}{b - a} = f'(c).$$

A precise statement of the theorem is given below.




Figure 5.1: The mean value theorem – AB is parallel to the tangent at C .

Theorem 5.1.1 (The mean value theorem). *Suppose that f is continuous on $[a, b]$ and differentiable on (a, b) . Then there is at least one real number c in (a, b) such that*

$$\frac{f(b) - f(a)}{b - a} = f'(c).$$

The proof of the theorem will be presented in the next section. For now we illustrate the theorem with a simple example.

Example 5.1.2. Suppose that $f : [1, 8] \rightarrow \mathbb{R}$ is given by $f(x) = \sqrt[3]{x}$. Find a number c in $(1, 8)$ that satisfies the conclusions of the mean value theorem for f on $[1, 8]$.

Solution. Note that f is continuous on $[1, 8]$ and differentiable on $(1, 8)$. By the mean value theorem there is a real number c in $(1, 8)$ such that

$$\frac{f(8) - f(1)}{8 - 1} = f'(c).$$


Now $f'(x) = \frac{1}{3}x^{-2/3}$ so the above equation becomes

$$\frac{2 - 1}{7} = \frac{1}{3\sqrt[3]{c^2}}.$$

Rearranging to find c gives

$$\begin{aligned}\sqrt[3]{c^2} &= \frac{7}{3} \\ c &= \sqrt{\frac{343}{27}}.\end{aligned}$$

Since $c \approx 3.56$ it is clear that $c \in (1, 8)$. This is illustrated graphically below.



□

5.2 Proof of the mean value theorem

(Ref: SH10 §4.1)

In this section we prove the mean value theorem. The proof is part of MATH1141 only, so MATH1131 students may want to jump straight to Section 5.3.

To prove the mean value theorem, we begin by considering the special case when $f(a) = f(b)$. This case is known as *Rolle's theorem* and is illustrated in Figure 5.2.


Theorem 5.2.1. *Suppose that g is continuous on $[a, b]$ and differentiable on (a, b) . Suppose also that $g(a) = g(b) = 0$. Then there is a real number c in (a, b) such that $g'(c) = 0$.*

Proof. The proof considers three cases.

Case 1: Suppose that g is the constant function given by $g(x) = 0$. Then $g'(c) = 0$ for every c in (a, b) (see Section 4.2).

Case 2: Suppose that there is a point d in (a, b) such that $g(d) > 0$. By the max-min theorem (Theorem 3.4.3), g attains a maximum value at some point c in $[a, b]$. Moreover, c cannot be a or b since $g(d) > g(a) = g(b) = 0$. Hence c lies in (a, b) , and since g is differentiable on (a, b) , we have $g'(c) = 0$ (see Theorem 4.8.2).

Case 3: Suppose that $g(x) \leq 0$ for all x in $[a, b]$ and that g is not constant on $[a, b]$. Then g attains a minimum at a point c in (a, b) , and similarly to Case 2 one can show that $g'(c) = 0$. □

Figure 5.2: Rolle's theorem – the tangent at C is horizontal.

From here it is not hard to see how to prove the mean value theorem. Consider the graph of the function f in Figure 5.1 and ‘subtract’ the chord AB . The result will look something like the graph of g in Figure 5.2, to which we can apply Rolle's theorem and obtain a horizontal tangent at C . We then ‘add’ the chord back again to obtain the tangent to f of Figure 5.1. This rough geometric argument is made rigorous below.

Proof of the mean value theorem. Suppose that f is continuous on $[a, b]$ and differentiable on (a, b) . We consider the function g given by

$$g(x) = f(x) - \left[\frac{f(b) - f(a)}{b - a}(x - a) + f(a) \right].$$

(The part in square brackets is taken from the equation of the chord AB in Figure 5.1.) One can check that g is continuous on $[a, b]$, differentiable on (a, b) and that $g(a) = g(b) = 0$. By Rolle's theorem there is a c in (a, b) such that $g'(c) = 0$; that is, such that

$$f'(c) - \left[\frac{f(b) - f(a)}{b - a} \right] = 0.$$

If we rearrange this equation then we obtain

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

as required. □

5.3 Proving inequalities using the mean value theorem

Our first application of the mean value theorem will be to prove some useful inequalities. We begin with an example.

Example 5.3.1. By using the mean value theorem, show that $\ln x < x - 1$ whenever $x > 1$.

Solution. Suppose that $x > 1$ and consider the closed interval $[1, x]$. We define a function $f : [1, x] \rightarrow \mathbb{R}$ by $f(t) = \ln t$. Now f is continuous on $[1, x]$ and differentiable on $(1, x)$. So we may apply the mean value theorem to f on the interval $[1, x]$. Now $f'(t) = 1/t$ so

$$\frac{f(x) - f(1)}{x - 1} = \frac{1}{c}$$

for some c in $(1, x)$ by the mean value theorem. That is,

$$\frac{\ln x}{x-1} = \frac{1}{c}$$

for some c between 1 and x . Since $c > 1$ we have $1/c < 1$ and hence

$$\frac{\ln x}{x-1} < 1.$$

Rearranging the inequality gives $\ln x < x - 1$ as desired. $\square$

The technique used in the above example is as follows. Consider a function f that is continuous on the interval $[a, x]$ and differentiable on the interval (a, x) . The mean value theorem gives

$$\frac{f(x) - f(a)}{x - a} = f'(c) \quad (5.1)$$

for some c between a and x . If $f'(c) < M$ then

$$\frac{f(x) - f(a)}{x - a} < M$$

and hence

$$f(x) < M(x - a) + f(a). \quad (5.2)$$

whenever $x > a$.

Sometimes it is desirable for the inequality to go the other way. In this case we find m such that $f'(c) > m$. Then, instead of (5.2), we obtain

$$f(x) > m(x - a) + f(a).$$

Often, the tricky part of the process is determining which function f will give the desired inequality.

Example 5.3.2. Use the mean value theorem to show that

$$\sqrt{x+4} - 2 < \frac{x}{4}$$

whenever $x > 0$.

Solution. Fix positive x . The first step is to identify a good choice of f . If we rewrite the inequality as

$$\sqrt{x+4} < \frac{x}{4} + 2$$

then we have a similar form to inequality (5.2). This suggests that we should define $f : [0, x] \rightarrow \mathbb{R}$ by the formula $f(t) = \sqrt{t+4}$. Then f is continuous on $[0, x]$ and differentiable on $(0, x)$. An application of the mean value theorem gives

$$\frac{f(x) - f(0)}{x - 0} = \frac{1}{2\sqrt{c+4}}$$

for some c in $(0, x)$. Now the right-hand side is bounded above by $\frac{1}{2\sqrt{4}}$, so we have

$$\frac{\sqrt{x+4} - \sqrt{0+4}}{x - 0} < \frac{1}{2\sqrt{4}}.$$

Simplifying and rearranging gives

$$\sqrt{x+4} - 2 < \frac{x}{4}$$

as required. $\square$

5.4 Error bounds

A second application of the mean value theorem is in calculating error bounds. Suppose that a calculation involves $\sqrt{27}$. Clearly $\sqrt{27}$ is close to $\sqrt{25}$, but the latter is easier to work with because it is equal to 5. If we use 5 instead of $\sqrt{27}$ then calculations are made easier with the tradeoff that we introduce an error. How bad is this error?

Example 5.4.1. Use the mean value theorem to find an upper bound for the error involved if we approximate $\sqrt{27}$ by $\sqrt{25}$.

Solution. The precise error in the approximation is given by

$$\sqrt{27} - \sqrt{25}.$$

We apply the mean value theorem to the function f , given by $f(x) = \sqrt{x}$, on the interval $[25, 27]$. This gives

$$\frac{\sqrt{27} - \sqrt{25}}{27 - 25} = \frac{1}{2\sqrt{c}}$$

for some c in $(25, 27)$. Rearranging this gives us

$$\sqrt{27} - \sqrt{25} = \frac{1}{\sqrt{c}}$$

where $25 < c < 27$. Hence

$$\begin{aligned} \text{error} &= \sqrt{27} - \sqrt{25} \\ &= \frac{1}{\sqrt{c}} \\ &< \frac{1}{\sqrt{25}} \\ &= \frac{1}{5}, \end{aligned}$$

since $c > 25$. Hence an upper bound for the error is $1/5$. □

We illustrate the process with one more example.

Example 5.4.2. Use the mean value theorem to find an upper bound for the error involved if we approximate $\sin \frac{5\pi}{7}$ by $\sin \frac{2\pi}{3}$.

Solution. Note that since

$$\frac{2\pi}{3} = \frac{14\pi}{21} \approx \frac{15\pi}{21} = \frac{5\pi}{7}$$

and the sine function is continuous,

$$\sin \frac{2\pi}{3} \approx \sin \frac{5\pi}{7}.$$

The (absolute) error of the approximation is exactly

$$\left| \sin \frac{5\pi}{7} - \sin \frac{2\pi}{3} \right|.$$

If we apply the mean value theorem to the sine function on the interval $[2\pi/3, 5\pi/7]$ then

$$\frac{\sin \frac{5\pi}{7} - \sin \frac{2\pi}{3}}{\frac{5\pi}{7} - \frac{2\pi}{3}} = \cos c$$

for some c in $(2\pi/3, 5\pi/7)$. Taking absolute values of both sides and rearranging gives

$$\left| \sin \frac{5\pi}{7} - \sin \frac{2\pi}{3} \right| = \left| \frac{5\pi}{7} - \frac{2\pi}{3} \right| |\cos c|$$

where $2\pi/3 < c < 5\pi/7$. Now $|\cos c| < |\cos \frac{5\pi}{7}|$ (because the absolute value of the cosine is increasing over the interval $(\pi/2, \pi)$) but we don't have an exact value for $\cos \frac{5\pi}{7}$. A simple way out of this situation is to just use the fact that $|\cos c| < 1$ for all c in $(2\pi/3, 5\pi/7)$. Hence we have shown that

$$\text{absolute error} = \left| \sin \frac{15\pi}{21} - \sin \frac{2\pi}{3} \right| < \left| \frac{5\pi}{7} - \frac{2\pi}{3} \right| \times 1.$$

This proves that

$$\text{absolute error} < \frac{\pi}{21} < 0.15.$$

If we aren't so lazy with our estimates we can quickly spot that

$$\left| \cos \frac{5\pi}{7} \right| = \left| \cos \frac{15\pi}{21} \right| < \left| \cos \frac{15\pi}{20} \right| = \frac{1}{\sqrt{2}}.$$

This allows us to prove the more accurate statement that

$$\text{absolute error} < \frac{\pi}{21\sqrt{2}} < 0.11.$$

□

5.5 The sign of a derivative


(Ref: SH10 §§4.2, 4.3)

Suppose that a function f is differentiable at 4 and that $f'(4) = -2$. In high school, it was taught that, since the derivative of f at 4 is negative, the function f is decreasing at the point 4. This makes intuitive sense since the gradient of the tangent to f at the point 4 is -2 and so the tangent slopes downward. In this section, these ideas are given a rigorous foundation by using the mean value theorem.

Definition 5.5.1. Suppose that a function f is defined on an interval I . We say that

- a) f is *increasing* on I if for every two points x_1 and x_2 in I ,
 $x_1 < x_2$ implies that $f(x_1) < f(x_2)$;
- b) f is *decreasing* on I if for every two points x_1 and x_2 in I ,
 $x_1 < x_2$ implies that $f(x_1) > f(x_2)$.

Example 5.5.2. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^2$ is increasing on $[0, \infty)$ and decreasing on $(-\infty, 0]$.



The following theorem should be familiar to you. It is proved using the mean value theorem.

Theorem 5.5.3. Suppose that f is continuous on $[a, b]$ and differentiable on (a, b) .

- (i) If $f'(x) > 0$ for all x in (a, b) then f is increasing on $[a, b]$.
- (ii) If $f'(x) < 0$ for all x in (a, b) then f is decreasing on $[a, b]$.
- (iii) If $f'(x) = 0$ for all x in (a, b) then f is constant on $[a, b]$.

Proof. We only prove statement (i) since the proofs of statements (ii) and (iii) are similar. Suppose that $f'(x) > 0$ for all x in (a, b) and choose two points x_1 and x_2 in $[a, b]$ such that $x_1 < x_2$. By Definition 5.5.1, it suffices to show that $f(x_1) < f(x_2)$.

Since f is differentiable on I , it is continuous on $[x_1, x_2]$ (see Theorem 4.5.1) and differentiable on (x_1, x_2) . Hence, by the mean value theorem, there is a c in (x_1, x_2) such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(c).$$

But $f'(c) > 0$ and $x_2 - x_1 > 0$. This means that $f(x_2) - f(x_1) > 0$ and hence $f(x_1) < f(x_2)$ as required. $\square$

Theorem 5.5.3 has several important applications. The first of these is classifying different types of stationary points. The basic idea is as follows. Suppose that f is differentiable on the open interval $(1, 3)$ and that for some small positive number h we have the following:

- $f'(x) > 0$ for all x in $(2 - h, 2)$,
- $f'(2) = 0$, and
- $f'(x) < 0$ for all x in $(2, 2 + h)$.

Now f is differentiable on $(2 - h, 2 + h)$ so it is also continuous on this interval. Moreover, f increases on the interval $(2 - h, 2)$, is stationary at 2 and then decreases on the interval $(2, 2 + h)$. From these facts we conclude that the stationary point 2 is a local maximum point for f .

Example 5.5.4. Find and classify all stationary points of the function $f : \mathbb{R} \rightarrow \mathbb{R}$ whose derivative is given by

$$f'(x) = (x - 4)(x - 1)(x + 5)^2.$$

Solution. The stationary points, which are found by solving the equation $f'(x) = 0$, are 4, 1 and -5 . To classify each stationary point, we examine the sign of f' on small intervals either side of each stationary point. This process is concisely documented in the following table.

	-5^-	-5	-5^+	1^-	1	1^+	4^-	4	4^+
$x - 4$	$-$	$-$	$-$	$-$	$-$	$-$	$-$	0	$+$
$x - 1$	$-$	$-$	$-$	$-$	0	$+$	$+$	$+$	$+$
$(x + 5)^2$	$+$	0	$+$	$+$	$+$	$+$	$+$	$+$	$+$
$f'(x)$	$+$	0	$+$	$+$	0	$-$	$-$	0	$+$
Gradient	$/$	$-$	$/$	$/$	$-$	$\backslash$	$\backslash$	$-$	$/$

For example, the notation 4^- is shorthand for those points x lying in a small interval to the immediate left of 4. For such a point x , the factor $x - 4$ is negative, while the factors $x - 1$ and $(x + 5)^2$ are positive. Hence $f'(x)$ is negative when x is slightly to the left of 4. Other columns of the table are filled by a similar process.

In conclusion, 4 is a local minimum point, 1 is a local maximum point and -5 is neither a maximum nor minimum point. (In fact, -5 is called a *horizontal point of inflexion* since the tangent to f at -5 is horizontal and the concavity of the function changes about this point. See any standard undergraduate calculus text or any advanced high school calculus text for further details.) $\square$

5.6 The second derivative and applications

(Ref: SH10 §4.3)

In this section we give another method for classifying the stationary points of a function f . To do so, we use the *second derivative of f* , which is denoted by f'' .

Example 5.6.1. Find the second derivative of the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by the rule

$$f(x) = 3x^4 + 2x - \sin x.$$

Solution. The *first derivative* of f is given by

$$f'(x) = 12x^3 + 2 - \cos x.$$

To find the second derivative f'' of f , we simply differentiate f' . Hence

$$f''(x) = 36x^2 + \sin x.$$

$\square$

Remark 5.6.2. If $y = f(x)$ then the first derivative of f is often written as $\frac{dy}{dx}$ or y' while the second derivative of f is often written as $\frac{d^2y}{dx^2}$ or y'' .

Remark 5.6.3. Even if a function is differentiable at a point a , the function may not have a second derivative at a . (Consider, for example, the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = x^{4/3},$$

at the point 0.) A function which has a second derivative at a is called *twice differentiable* at a .

If f is twice differentiable, the sign of f'' can often be used to determine whether a stationary point is a local maximum point or a local minimum point.

Theorem 5.6.4 (The second derivative test). *Suppose that a function f is twice differentiable on (a, b) and that $c \in (a, b)$.*

(i) *If $f'(c) = 0$ and $f''(c) > 0$ then c is a local minimum point of f ;*

(ii) *If $f'(c) = 0$ and $f''(c) < 0$ then c is a local maximum point of f .*

The proof of the theorem uses the following result.

Lemma 5.6.5. *Suppose that g is differentiable at a point c .*

(i) *If $g'(c) > 0$ then $g(c - h) < g(c) < g(c + h)$ for all positive h sufficiently small.*

(ii) *If $g'(c) < 0$ then $g(c + h) < g(c) < g(c - h)$ for all positive h sufficiently small.*

Sketch proof. We give an intuitive argument for statement (i) and leave (ii) as an exercise. Suppose that $g'(c) > 0$. By the definition of the derivative,

$$\lim_{h \rightarrow 0} \frac{g(c + h) - g(c)}{h} = g'(c). \quad (5.3)$$

That is,

$$\frac{g(c + h) - g(c)}{h} \approx g'(c)$$

when h is close to 0. Now, if $h > 0$ then

$$g(c + h) - g(c) \approx hg'(c) > 0$$

Hence

$$g(c + h) - g(c) > 0 \quad (5.4)$$

provided that $h > 0$ and that h is sufficiently small.

On the other hand, the definition of the derivative also implies that

$$\lim_{h \rightarrow 0} \frac{g(c - h) - g(c)}{-h} = g'(c)$$

(to see this, simply replace h with $-h$ in (5.3)). Now, if $h > 0$ and h is close to 0 then

$$g(c) - g(c - h) \approx hg'(c) > 0.$$

Hence

$$g(c) - g(c - h) > 0 \quad (5.5)$$

provided that $h > 0$ and that h is sufficiently small.

If we combine (5.4) and (5.5) then we have

$$g(c - h) < g(c) < g(c + h)$$

for all positive h sufficiently small. □

Proof of Theorem 5.6.4. We prove statement (i) and leave the proof of (ii) as an exercise. Suppose that $f'(c) = 0$, $f''(c) > 0$ and that $g = f'$. Then $g'(c) > 0$ and

$$g(c - h) < g(c) < g(c + h)$$

for all positive h sufficiently small, by Lemma 5.6.5. Since $f'(c) = g(c) = 0$ we conclude that

$$f'(c - h) < 0 < f'(c + h)$$

for all positive h sufficiently small. So f' is positive on a small interval immediately to the right of c , zero at x and negative on a small interval immediately to the left of c . By Theorem 5.5.3 we now conclude that f has a local minimum at the point c . $\square$


Example 5.6.6. Find and classify the stationary points of the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = x^3 - 6x^2 + 9x - 5.$$

Solution. First we calculate the first and second derivatives of f :


$$\begin{aligned} f'(x) &= 3x^2 - 12x + 9 = 3(x - 1)(x - 3) \\ f''(x) &= 6x - 12. \end{aligned}$$

Hence f has stationary points at 1 and 3. Now $f''(1) = -6 < 0$, so f has a local maximum point at 1. For second stationary point, $f''(3) = 6 > 0$, so f has a local minimum stationary point at 3. (To illustrate these conclusions, we have included the graph of f , shown below.)



$\square$

Remark 5.6.7. Suppose that $f'(c) = 0$ and that $f''(c) = 0$. Then we cannot apply the second derivative test. In fact, the point c may be a local maximum, a local minimum or neither of these. The examples below illustrate various possibilities.



- If $f(x) = x^4$ then $f'(0) = f''(0) = 0$ and there is a local *minimum* at 0.
- If $f(x) = -x^4$ then $f'(0) = f''(0) = 0$ and there is a local *maximum* at 0.
- If $f(x) = x^3$ then $f'(0) = f''(0) = 0$ and there is neither a local maximum and minimum at 0 (in fact, we have a horizontal point of inflexion at 0).

Hence if $f'(c) = f''(c) = 0$ then it is best to classify the stationary point c by examining the sign of the derivative on either side of c , as illustrated in Section 5.5.

5.7 Critical points, maxima and minima

(Ref: SH10 §§4.3–4.5)

By combining results from Chapters 3, 4 and 5, we now give a comprehensive presentation of how to locate global maxima and minima for any real-valued continuous function defined on a closed interval.

Suppose that a function $f : [a, b] \rightarrow \mathbb{R}$ is continuous on $[a, b]$. By the max-min theorem (Theorem 3.4.3), f attains a global maximum and a global minimum on $[a, b]$. Suppose that the global maximum occurs at the point c . Now either c is one of the endpoints a or b , or c lies in the open interval (a, b) . In the latter case, f is either differentiable at c or it is not. If it is differentiable at c , then $f'(c) = 0$ by Theorem 4.8.2.

Thus we have identified that a global maximum point c must satisfy one of the three following properties:

- c is one of the endpoints a or b of the closed interval $[a, b]$,
- f is differentiable at c and $f'(c) = 0$, or
- f is not differentiable at c .

The set of points satisfying these properties is useful so we give it a special name.

Definition 5.7.1. Suppose that f is defined on $[a, b]$. We say that a point c in $[a, b]$ is a *critical point* for f on $[a, b]$ if c satisfies one of the following properties:

- (a) c is an endpoint a or b of the interval $[a, b]$,
- (b) f is not differentiable at c , or
- (c) f is differentiable at c and $f'(c) = 0$.

Theorem 5.7.2. Suppose that f is continuous on $[a, b]$. Then f has a global maximum and global minimum on $[a, b]$. Moreover, the global maximum point and the global minimum point are both critical points for f on $[a, b]$.

The proof of the theorem follows the same reasoning given at the beginning of this section. The theorem provides a systematic way of finding global minima and maxima of functions.




Figure 5.3: Graphs of functions and their critical points.

Example 5.7.3. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$f(x) = 3(x - 1)^{2/3}.$$

Find the absolute maximum and minimum values of f on $[0, 2]$.

Solution. The derivative of f is given by

$$f'(x) = 2(x - 1)^{-1/3} = \frac{2}{\sqrt[3]{x - 1}}.$$

Clearly f is not differentiable at 1. Moreover, its derivative is never zero. So the only critical points of f on $[0, 2]$ are 1 and the endpoints 0 and 2. By Theorem 5.7.2, the global maximum and minimum points must be among the points 0, 1 and 2. But

$$f(0) = 3, \quad f(1) = 0 \quad \text{and} \quad f(2) = 3.$$

Hence the global minimum value of f on $[0, 2]$ is 0 while the global maximum value of f on $[0, 2]$ is 3. (See Figure 5.3 (a) for a sketch of f that shows its critical points.) $\square$

Example 5.7.4. Suppose that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by the rule

$$f(x) = |x^2 - 3x - 4|.$$

Find the absolute maximum and absolute minimum values of f on the interval $[0, 5]$.

Example 5.7.5. Consider the parabolic arc

$$\{(x, y) \in \mathbb{R}^2 : y = x^2 - x - 1, -2 \leq x \leq 3\}$$

(see Figure 5.4 (a)). Find where the arc is (i) closest to and (ii) furthest from the origin.

Solution. If (x, y) is a point on the arc then its distance from the origin is given by

$$\sqrt{x^2 + y^2}.$$




Figure 5.4: Graphs for Example 5.7.5.

To find where the arc is closest to the origin, we want to minimise $\sqrt{x^2 + y^2}$ subject to the constraints $y = x^2 - x - 1$ and $-2 \leq x \leq 3$. In other words, we want to find a global minimum for the function

$$f(x) = \sqrt{x^2 + (x^2 - x - 1)^2}$$

on the interval $[-2, 3]$. Because calculations involving square roots can get messy, we instead consider the function g , given by

$$g(x) = x^2 + (x^2 - x - 1)^2$$

on $[-2, 3]$. (Note that $g(x)$ may be interpreted as the square of the distance between the origin and a point on the arc with coordinate $(x, x^2 - x - 1)$. The global minimum of both f and g will occur at the same point x .)

We seek the critical points of g , so we compute its derivative:

$$\begin{aligned} g'(x) &= 2x + 2(x^2 - x - 1)(2x - 1) \\ &= 2(x + 2x^3 - 2x^2 - 2x - x^2 + x + 1) \\ &= 2(2x^3 - 3x^2 - x^2 + 1) \\ &= 2(x - 1)(2x^2 - x - 1) \\ &= 2(x - 1)(x - 1)(2x + 1). \end{aligned}$$

Hence the critical points of g on $[-2, 3]$ are -2 , $-1/2$, 1 and 3 . Now

$$g(-2) = 29, \quad g(-1/2) = 5/16, \quad g(1) = 2, \quad g(3) = 34$$

(see Figure 5.4 (b)). Therefore the arc is closest to the origin when $x = -1/2$ and furthest from the origin when $x = 3$. $\square$

5.8 Counting zeros

In this section we combine the intermediate value theorem with results about the sign of derivatives to determine how many solutions there are to various equations. The main ideas are illustrated in the next example.

Example 5.8.1. Determine how many real numbers satisfy the equation

$$x^3 - 6x^2 - 15x + 8 = 0. \quad (5.6)$$

Solution. Suppose that $f(x) = x^3 - 6x^2 - 15x + 8$ for all real x . Ideally, we would look for one root of f , factor this out and solve a quadratic equation to determine the remaining roots (if any). However, finding one root by inspection is not straightforward, so we need another approach.

Note that f is differentiable (and hence continuous) everywhere. We begin by identifying intervals where f is monotonically decreasing and intervals where f is monotonically increasing. The derivative of f is given by

$$f'(x) = 3x^2 - 12x - 15 = 3(x - 5)(x + 1).$$


So the stationary points of f are 5 and -1 . Moreover, the table below shows that f is increasing on the intervals $(-\infty, -1]$ and $[5, \infty)$, while it is decreasing on $[-1, 5]$.

Interval	$(-\infty, -1)$	$(-1, 5)$	$(5, \infty)$
$(x - 5)$	—	—	+
$(x + 1)$	—	+	+
$f'(x)$	+	—	+

We evaluate f at the (finite) endpoints of these intervals and see that

$$f(-1) = 16 \quad \text{and} \quad f(5) = -92.$$

Since f is a cubic, we can now give a rough sketch of the graph of f .



We now argue that f has *exactly* one root on each of the intervals mentioned above.

- On the interval $[-1, 5]$, the function f is decreasing. Hence f cannot have more than one root on $[-1, 5]$. On the other hand, since $f(-1) > 0$ and $f(5) < 0$ there is at least one c in $[-1, 5]$ such that $f(c) = 0$ by the intermediate value theorem. Hence f has exactly one root on $[-1, 5]$.
- On the interval $(-\infty, -1]$, the function f is increasing. Hence f cannot have more than one root on $(-\infty, -1]$. On the other hand, since $f(-1) > 0$ and $f(-5) < 0$ there is at least one c in $[-5, -1]$ such that $f(c) = 0$ by the intermediate value theorem. Hence f has exactly one root on $(-\infty, -1]$.

- On the interval $[5, \infty)$, the function f is increasing. Hence f cannot have more than one root on $[5, \infty)$. On the other hand, since $f(5) < 0$ and $f(10) > 0$ there is at least one c in $[5, 10]$ such that $f(c) = 0$ by the intermediate value theorem. Hence f has exactly one root on $[5, \infty)$.

In conclusion, equation (5.6) has exactly three real solutions, one in each of the intervals $[-5, -1]$, $[-1, 5]$ and $[5, 10]$. $\square$

Guided by our solution to Example 5.8.1, we now outline a general approach for solving problems of this type.

Suppose that f is continuous and differentiable everywhere and that its derivative is also continuous. To determine how many real solutions the equation $f(x) = 0$ has, one may follow the procedure below.

1. Calculate f' and solve $f'(x) = 0$.
2. Determine the intervals where the derivative is positive and the intervals where it is negative.
3. By step 2 we know the intervals where the function f is monotonically increasing, and the intervals where f is monotonically decreasing.
4. Evaluate the function f at the endpoints of each interval. If f changes sign on the interval, there is exactly one root on that interval. If it does not change sign, there are no roots on that interval. (This step must be modified slightly for intervals of the form $(-\infty, a]$ or $[b, \infty)$.)

Example 5.8.2. Determine how many real numbers satisfy the equation

$$2x^3 - 9x^2 + 12x - 1 = 0. \quad (5.7)$$

Solution. We follow the steps outlined above. Suppose that

$$f(x) = 2x^3 - 9x^2 + 12x - 1.$$


By differentiating f we obtain

$$f'(x) = 6x^2 - 18x + 12 = 6(x-1)(x-2).$$

The stationary points of f are 1 and 2. By studying the sign of f' , it is clear that f is increasing on the intervals $(-\infty, 1)$ and $(2, \infty)$ while it is decreasing on $(1, 2)$. Now

$$f(1) = 4 \quad \text{and} \quad f(2) = 3,$$

as illustrated in the sketch below.



So it is clear that f has no zeros on $[1, 2]$ or on $[2, \infty)$. On $(-\infty, 1)$, the function f has no more than one zero (since it is increasing on this interval). Moreover, f changes sign on the interval $[0, 1]$ (since $f(0) = -1 < 0$ and $f(1) > 0$). Hence f has exactly one root on $[0, 1]$ by the intermediate value theorem.

In conclusion, equation (5.7) has exactly one real solution and this solution lies somewhere in the interval $[0, 1]$. $\square$

5.9 Antiderivatives

We open with an example.

Example 5.9.1. While filming a movie, a stunt man jumps out of a stationary helicopter 900 meters above the ground. The scriptwriter wants the man to make a 20 second mobile phone call before opening the parachute. Physical considerations (gravity and air resistance) suggest that the man's velocity $f(t)$ (in metres per second) without an open parachute is given by

$$f(t) = 50(1 - e^{-t/5}),$$

where t is the number of seconds after jumping from the helicopter. How far will the man fall 20 seconds?

Solution. Write $F(t)$ for the distance fallen (in metres) after t seconds. We note that $F(0) = 0$. Since velocity is the rate of change of displacement, F satisfies the equation

$$F'(t) = f(t).$$

(Any equation like this involving differentiation is called a *differential equation*). The differential equation may be rewritten as

$$\frac{dF}{dt} = 50 - 50e^{-t/5}. \quad (5.8)$$

One can check by differentiation that if

$$F(t) = 50t + 250e^{-t/5} + C, \quad (5.9)$$

where C is a real constant, then F satisfies (5.8). Let's *assume* (for the moment) that all possible solutions to the differential equation (5.8) are of the form (5.9). Imposing the condition that $F(0) = 0$ allows us to evaluate the unknown constant C :

$$0 = F(0) = 50 \times 0 + 250e^0 + C,$$

whence $C = -250$. Therefore $F(t) = 50t + 250e^{-t/5} - 250$. To complete the solution,

$$F(20) = 50 \times 20 + 250e^{-4} - 250 \approx 754.6.$$

Hence the stunt man will fall about 755 metres in the first 20 seconds. $\square$

To be certain of this solution, we need to show that *every* function F satisfying the differential equation (5.8) is of the form (5.9). This can be done using the mean value theorem. We begin by introducing the notion of an *antiderivative*.

Definition 5.9.2. Suppose that f is continuous on an open interval I . A function F is said to be an *antiderivative* (or a *primitive*) of f on I if $F'(x) = f(x)$ for all x in I . The process of finding an antiderivative of a function is called *antidifferentiation*.

Example 5.9.3. (a) Suppose that

$$f(t) = 50 - 50e^{-t/5} \quad \forall t \in \mathbb{R}$$

Then the function F , given by

$$F(t) = 50t + 250e^{-2t} - 250$$

is an antiderivative of f on $(-\infty, \infty)$.

(b) Suppose that n is a positive integer and $f(x) = x^n$. If

$$F(x) = \frac{x^{n+1}}{n+1} \quad \text{and} \quad G(x) = \frac{x^{n+1}}{n+1} + 5$$

then F and G are both antiderivatives of f on the interval $(-\infty, \infty)$. So f has more than one antiderivative on $(-\infty, \infty)$. In fact, if C is any real number and

$$H(x) = x^{n+1}/(n+1) + C$$

then H is an antiderivative of f on $(-\infty, \infty)$. Thus the function f has infinitely many antiderivatives on $(-\infty, \infty)$.

Example 5.9.3 (b) illustrates the general principle that if F is an antiderivative of f on I then

$$F + C$$

is also an antiderivative of f on I , for any real constant C . The next theorem says that all antiderivatives are of this form.

Theorem 5.9.4. Suppose that f is a continuous function on an open interval I and that F and G are two antiderivatives of f on I . Then there is a real constant C such that

$$G(x) = F(x) + C$$

for all x in I .

Proof. Suppose that F and G are two antiderivatives of f on I and let H denote the function given by

$$H(x) = G(x) - F(x)$$

for all x in I . Then H is differentiable on I and

$$\begin{aligned} H'(x) &= G'(x) - F'(x) \\ &= f(x) - f(x) \\ &= 0 \end{aligned}$$

for all x in I . Hence there is a constant C such that $H(x) = C$ for all x in I (by Theorem 5.5.3 (c)). Therefore

$$G(x) = F(x) + H(x) = F(x) + C$$

for all x in I . □

Theorem 5.9.4 justifies the assumption made in the solution to Example 5.9.1. It also enables us to write down all possible antiderivatives of some well-known functions. In the table below, C is any real constant.

Function	Antiderivative
x^r , where r is rational and $r \neq -1$	$\frac{1}{r+1}x^{r+1} + C$
$\sin x$	$-\cos x + C$
$\cos x$	$\sin x + C$
e^{ax}	$\frac{1}{a}e^{ax} + C$
$\frac{f'(x)}{f(x)}$	$\ln f(x) + C$

5.10 L'Hôpital's rule

(Ref: SH10 §§11.5, 11.6)

Consider the limit

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{5x + x^2}.$$

Since both the numerator and denominator approach 0 as $x \rightarrow 0^+$, the limit is an example of an indeterminate form of type $\frac{0}{0}$. None of the limit rules of Chapter 2 can be easily applied to evaluate the limit. However, the mean value theorem gives another rule which helps in this situation.

Suppose that $f(x) = e^x - 1$ and $g(x) = 5x + x^2$. Instead of considering the quotient $f(x)/g(x)$ (whose limit is difficult to calculate), we consider the quotient $f'(x)/g'(x)$ of derivatives (whose limit is easier to calculate):

$$\lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow 0} \frac{e^x}{5 + 2x} = \frac{1}{5}.$$

Since this second limit exists, *l'Hôpital's rule* (which is stated below) implies that

$$\lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)}.$$

Hence

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{5x + x^2} = \frac{1}{5}.$$

Theorem 5.10.1 (l'Hôpital's rule). *Suppose that f and g are both differentiable functions and a is a real number. Suppose also that either one of the two following conditions hold:*

- $f(x) \rightarrow 0$ and $g(x) \rightarrow 0$ as $x \rightarrow a$;
- $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow a$.

If

$$\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

exists then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}.$$

Remark 5.10.2. The theorem also holds for

- limits as $x \rightarrow \infty$ or $x \rightarrow -\infty$;
- one-sided limits (that is, as $x \rightarrow a^+$ or as $x \rightarrow a^-$).

L'Hôpital's rule is proved using the mean value theorem (see the sketch proof at the end of this section).

Example 5.10.3. Calculate $\lim_{x \rightarrow 1} \frac{\ln x}{1 - x}$.

Solution. If $f(x) = \ln x$ and $g(x) = 1 - x$ then both $f(x)$ and $g(x)$ approach 0 as $x \rightarrow 1$. Moreover, f and g are both differentiable. So we look at the quotient of the derivatives:

$$\frac{f'(x)}{g'(x)} = \frac{1/x}{-1} = -\frac{1}{x} \rightarrow -1$$

as $x \rightarrow 1$. We conclude by l'Hôpital's rule that

$$\lim_{x \rightarrow 1} \frac{f(x)}{g(x)} = -1.$$

(*Note:* for short we may write

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{\ln x}{1 - x} &= \lim_{x \rightarrow 1} \frac{1/x}{-1} \\ &= \lim_{x \rightarrow 1} -\frac{1}{x} \\ &= -1 \end{aligned} \tag{5.10}$$

provided that we check that our application of l'Hôpital's rule in (5.10) is valid.) □

Sometimes l'Hôpital's rule must be applied more than once to calculate a limit. The next example illustrates this.

Example 5.10.4. Determine the limiting behaviour of $\frac{x^2}{e^{2x}}$ as $x \rightarrow \infty$.

Solution. Observe that both the numerator and denominator approach ∞ as $x \rightarrow \infty$. Differentiating the numerator and denominator gives

$$\frac{2x}{2e^{2x}}.$$

Again, as $x \rightarrow \infty$ we have an indeterminate form of type $\frac{\infty}{\infty}$. If we differentiate the numerator and denominator once more we have

$$\frac{2}{4e^{2x}}$$

and this limit can be evaluated. Piecing this together, we have

$$\lim_{x \rightarrow \infty} \frac{x^2}{e^{2x}} = \lim_{x \rightarrow \infty} \frac{2x}{2e^{2x}} = \lim_{x \rightarrow \infty} \frac{2}{4e^{2x}} = 0$$

by l'Hôpital's rule. □

As the next example shows, l'Hôpital's rule can be used to solve limits that are indefinite forms of type $0 \times \infty$.

Example 5.10.5. Determine the limiting behaviour of $x \ln x$ as $x \rightarrow 0^+$.

The next example warns against the improper application of l'Hôpital's rule.

Example 5.10.6. Determine the limiting behaviour of $\frac{2x - \cos x}{3x + \cos x}$ as $x \rightarrow \infty$.

We shall now sketch the proof of l'Hôpital's rule for the case when $x \rightarrow a^+$. Other cases are done similarly. (A complete proof uses a generalisation of the mean value theorem known as *Cauchy's mean value theorem*.)

Sketch proof of Theorem 5.10.1. Suppose that f and g are differentiable (and hence continuous) everywhere, that $f(x) \rightarrow 0$ and $g(x) \rightarrow 0$ as $x \rightarrow a^+$, and that

$$\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)}$$

exists. We need to show that

$$\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)}. \quad (5.11)$$

We begin by applying the mean value theorem to both functions f and g on the interval $[a, x]$. So there are real numbers c and d in (a, x) such that

$$\frac{f(x) - f(a)}{x - a} = f'(c) \quad \text{and} \quad \frac{g(x) - g(a)}{x - a} = g'(d). \quad (5.12)$$

Since f is continuous and $f(x) \rightarrow 0$ as $x \rightarrow a^+$, we have $f(a) = 0$. Similarly $g(a) = 0$. So by (5.12) we have

$$f(x) = (x - a)f'(c) \quad \text{and} \quad g(x) = (x - a)g'(d)$$

for some c and d in (a, x) . Hence

$$\frac{f(x)}{g(x)} = \frac{(x-a)f'(c)}{(x-a)g(d)} = \frac{f'(c)}{g'(d)}. \quad (5.13)$$

Now $c \rightarrow a^+$ and $d \rightarrow a^+$ as $x \rightarrow a^+$. Hence (5.13) becomes (5.11) as required. $\square$

Problems for Chapter 5

Problems 5.1 : The mean value theorem

1. [R] Find a real number c which satisfies the conclusions of the mean value theorem for each function f on the given interval.

a) $f(x) = x^3$ on $[1, 2]$ b) $f(x) = \sqrt{x}$ on $[0, 2]$.

2. [R] Suppose that $f(x) = 1/x$. Show that there is no real number c in $[-1, 2]$ such that

$$f'(c) = \frac{f(2) - f(-1)}{2 - (-1)}.$$

Why does this not contradict the mean value theorem?

3. [R] Consider the function f given by $f(x) = (x-2)^4 \cos(x^2 - 4x + 4)$. Use the mean value theorem to show that f' has a zero on the interval $[1, 3]$.

Problems 5.3 : Proving inequalities using the mean value theorem

4. [R] [V] By using the mean value theorem, show that

- a) $\ln(1+x) < x$ whenever $x > 0$;
 b) $-\ln(1-x) < x/(1-x)$ whenever $0 < x < 1$;
 c) $1+x < e^x$ whenever $x > 0$.

5. [R]

- a) Use the mean value theorem to show that $\sin t \leq t$ whenever $t > 0$.
 b) Hence show that $\sin t < t$ whenever $t > 0$.
 c) Using the pinching theorem and part (a), evaluate the limit $\lim_{x \rightarrow \infty} \sin \frac{1}{x}$.

6. [H] Prove that

$$1 + \frac{x}{2\sqrt{1+x}} < \sqrt{1+x} < 1 + \frac{x}{2} \quad \text{whenever } x > 0.$$

Problems 5.4 : Error bounds

7. [R] [V] Use the mean value theorem to find an upper bound for the error involved if we approximate

- a) $\sqrt{17}$ by $\sqrt{16} = 4$;
 b) $\left(\frac{1998}{1000}\right)^2$ by $2^2 = 4$;
 c) $\frac{1}{1002}$ by $\frac{1}{1000}$.

Problems 5.5 : The sign of a derivative,**5.6 : The second derivative and applications and****5.7 : Critical points, maxima and minima**

8. [R] The derivative of a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f'(x) = 3(x+1)(x-1)^2(x-4)^3$. Locate all stationary points of f and identify any local maximum or minimum points of f .

9. [X] The function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0, \end{cases}$$

is continuous but not differentiable at 0. Does f have a local maximum or a local minimum at 0? Prove your answer.

10. [R] Find the maximum and minimum values for each function f over the given interval.

- a) $f(x) = 3 - x^3$ over $[-2, 4]$ b) $f(x) = 3 - x^4$ over $[-2, 4]$
 c) $f(x) = x^3 - x^4$ over $[-5, 5]$ d) $f(x) = 2x(x+4)^3$ over $[-2, 1]$
 e) $f(x) = |x^2 - 3x + 2|$ over $[0, 3]$

11. [R] Find the point on the straight line $2x + 3y = 6$ which is closest to the origin.

12. a) i) [R] Show that the polynomial p_3 , where $p_3(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}$, has *at least* one real root.

- ii) [H] Show that the polynomial p_2 , where $p_2(x) = 1 + x + \frac{x^2}{2!}$, has no real roots and deduce that p_3 has *exactly* one real root.

- iii) [X] Deduce that $p_4(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} > 0$ for all real numbers x .

- b) [X] Suppose that $p_n(x) = \sum_{k=0}^n \frac{x^k}{k!}$ whenever $n = 1, 2, 3, \dots$. Use induction to prove that

- i) if n is even then $p_n(x) > 0$ for all real numbers x , and
 ii) if n is odd then $p_n(x)$ has exactly one real root and this root is negative.

13. [R] A wire of length 100 cm is cut into two pieces of length x cm and y cm. The piece of length x cm is bent into the shape of a square and the piece of length y cm into the shape of a circle. Find x and y so that the sum of the areas enclosed by the shapes will be

- a) a minimum b) a maximum.

14. [X] Suppose that $a \geq 0$. Find the greatest and least distances from the point $(a, 0)$ to the ellipse

$$\frac{x^2}{4} + \frac{y^2}{1} = 1.$$

(Have a precise answer before comparing with the given answer.)

15. [X] Find all the values of a and x , both in $[0, 2\pi]$, where

$$f(x) = \cos a + 2\cos(2x) + \cos(4x - a)$$

has a horizontal point of inflexion.

Problems 5.8 : Counting zeros

16. [R] Show that $x^3 + x - 9 = 0$ has only one real solution.
17. [R] [V] Suppose that $p(x) = x^3 - 12x^2 + 45x - 51$ whenever $x \in \mathbb{R}$. How many real zeros does p have?

Problems 5.9 : Antiderivatives

18. [R]

a) Find a function f that has the following properties:

$$\begin{aligned} f'(t) &= \sin t + t && \text{whenever } t \in \mathbb{R}, \\ f(0) &= 2. \end{aligned}$$

b) Are there any other functions with these properties? Explain your answer.

19. [R] [V] A particle moving along the x -axis has velocity $2t - t^2$ units per second after t seconds. Find
- a) the distance from the starting point after three seconds;
- b) the total distance travelled after three seconds.

Problems 5.10 : L'Hôpital's rule

20. [R] Calculate the following limits.

$$\begin{aligned} \text{a) } \lim_{x \rightarrow 0} \frac{e^x - 1}{x(3 + x)} & \quad \text{b) } \lim_{x \rightarrow 1} \frac{x^m - 1}{x^n - 1}, \quad n \neq 0 & \quad \text{c) } \lim_{x \rightarrow \pi/2} \frac{x - \pi/2}{\cos x} \\ \text{d) } \lim_{x \rightarrow 0} \frac{\ln(1 + x) - x}{x^2} & \quad \text{e) } \lim_{x \rightarrow \pi/2} \left(\frac{1 - \sin x}{1 + \cos 2x} \right) & \quad \text{f) } \lim_{x \rightarrow 0} \frac{\tan x - x}{x^3} \end{aligned}$$

21. [R] Determine the limiting behaviour in the following cases.

$$\begin{aligned} \text{a) } \frac{x^3 + 1}{x^4 + 1} \text{ as } x \rightarrow \infty & \quad \text{b) } \frac{e^{5x}}{x^3} \text{ as } x \rightarrow \infty \\ \text{c) } \frac{e^{5x}}{x^3} \text{ as } x \rightarrow -\infty & \quad \text{d) } x \sin(1/x) \text{ as } x \rightarrow \infty \\ \text{e) } \frac{\sqrt{x^4 + 1}}{\sqrt[3]{x^6 + 1}} \text{ as } x \rightarrow \infty & \quad \text{f) } \frac{\ln(x^3 + 1)}{\ln(x^2 + 1)} \text{ as } x \rightarrow \infty \end{aligned}$$

22. [H] Find the value of $\lim_{t \rightarrow 0} \left(\frac{1}{\ln(1+t)} + \frac{1}{\ln(1-t)} \right)$.
23. [H] Find (a, b) such that $\lim_{x \rightarrow 0} \frac{ax - 1 + e^{bx}}{x^2} = 1$.
24. [R] Explain why l'Hôpital's Rule cannot be used to find $\lim_{x \rightarrow \infty} \frac{4x + \sin x}{2x - \sin x}$. Use another method to find this limit.
25. [R] [V] Show that the function f , given by

$$f(x) = \begin{cases} e^{2x} & \text{if } x \geq 0 \\ 2x + 1 & \text{if } x < 0, \end{cases}$$

is differentiable at 0.

26. [R]

- a) Evaluate $\lim_{h \rightarrow 0^+} \frac{\cos \sqrt{h} - 1}{h}$.
- b) A function f is defined by

$$f(x) = \begin{cases} \cos \sqrt{x} & \text{if } x \geq 0 \\ ax + b & \text{if } x < 0, \end{cases}$$

where a and b are real numbers. By using the limit calculated in (a), find all possible values of a and b such that f is differentiable at 0.

27. [H] [V]

- a) Use l'Hôpital's rule to show that $\lim_{x \rightarrow 0^+} x \ln x = 0$.
- b) By using part (a), or otherwise, show that $\lim_{x \rightarrow 0^+} x^2 \ln x = 0$.
- c) A function f is defined by

$$f(x) = \begin{cases} x^2 \ln x & \text{if } x > 0 \\ ax + b & \text{if } x \leq 0, \end{cases}$$

where a and b are real numbers. Find all possible values of a and b such that f is differentiable at 0.

Chapter 6

Inverse functions

We return again to the pollution example (see the introductions to Chapter 2). According to our model, the volume of pollution $P(t)$ in the lake after t days of the factory's operation is given by

$$P(t) = \frac{10^9}{101}(1 - e^{-101t/10^5}).$$

Environmental authorities, after hearing that the eventual amount of pollution in the lake will be $10^9/101$ litres, determine that this level would be environmentally devastating. Recommendations are made that the amount of pollution should not exceed 8×10^6 litres. If factory operations do not change, in how many days will pollution levels in the lake exceed this amount?

Note that the function P expresses pollution as a function of time. One approach to solving this problem is to find a function T that expresses time as a function of pollution. Such a function is called an *inverse function* of P . This chapter is devoted to a study of inverse functions.

6.1 Some preliminary examples

The examples of this section give an introduction to some of the key concepts appearing in Chapter 6. (A rigorous treatment of these concepts will be given in later sections.)

Example 6.1.1. Jack and Jill are playing a number game. Jack says, 'A positive number, when squared, is equal to 16. What was the original number?' Jill correctly answers that the number was 4.

Jack goes again: 'A positive number, when squared, is equal to 49. What was the original number?' This time Jill answers '7.'

This game may be modelled using two functions. Jack's function f takes a positive number and squares it. That is, f is given by

$$f : (0, \infty) \rightarrow (0, \infty), \quad f(x) = x^2.$$

The function g that Jill uses to answer Jack's question is given by

$$g : (0, \infty) \rightarrow (0, \infty), \quad g(x) = \sqrt{x}.$$

So f takes an initial number and squares it, while g takes the square root and recovers the original number. A way of writing down the fact that g undoes (or reverses) what f does to any positive number x is:

$$g(f(x)) = x \quad \forall x \in \text{Dom}(f). \quad (6.1)$$




Figure 6.1: Correspondence between inputs and outputs

In fact, f also undoes what g does to any positive number. This is easily verified by checking that

$$f(g(x)) = x \quad \forall x \in \text{Dom}(g). \quad (6.2)$$

Because f and g have this special relationship, they are called *inverses* of each other.

The relationship between any function f and its inverse function g can also be viewed in the following way. If $y = f(x)$ and we want to express x as a function of y , then we find that $x = g(y)$. For the functions of Example 6.1.1, this is easily verified by the following calculation:

$$y = f(x) \iff y = x^2 \iff \sqrt{y} = x \iff x = g(y).$$

The next example shows that changing the domain of f gives a different inverse function g .

Example 6.1.2. Jack changes the game slightly. He says, ‘A negative number, when squared, is equal to 36. What was the original number?’ Jill correctly answers that the number was -6 .

Again, we model the game with two functions. Jack’s function f is given by

$$f : (-\infty, 0) \rightarrow (0, \infty), \quad f(x) = x^2,$$

while Jill’s function g is given by

$$g : (0, \infty) \rightarrow (-\infty, 0), \quad g(x) = -\sqrt{x}.$$

Note that g undoes what f does to any negative number, and vice-versa. That is, f and g satisfy equations (6.1) and (6.2) and are therefore inverse functions of each other.

Each function f in Example 6.1.1 and Example 6.1.2 had the property that for every output there is a unique corresponding input (See Figure 6.1 (a) and (b)). However, the function f graphed in Figure 6.1 (c) does not have a one-to-one correspondence between its outputs and inputs. As we see in the next example, this function f has no inverse function.

Example 6.1.3. Jack changes the game one more time. This time he says, ‘A real number, when squared, is equal to 64. What was the original number?’ This time Jill cannot give a definite answer to his question. The answer is either 8 or -8 , but she has no way of telling which it is.

The model for this situation is the following. Jack’s function f is now given by

$$f : \mathbb{R} \rightarrow [0, \infty), \quad f(x) = x^2;$$

but there is no function g that undoes what f does to a real number x . That is, there is no function g that satisfies the equation (6.1). Therefore f has no inverse function. The reason why this is the case is clear: the domain of f now includes the positive *and* negative real numbers, and therefore some (in fact most) outputs of f have two possible corresponding inputs. That is, there is no *one-to-one correspondence* between inputs and outputs of f .

The ideas introduced through these examples are summarised below.

- If f has an inverse function g , then g undoes (or reverses) what f does to elements of $\text{Dom}(f)$.
- Suppose f and g are inverse functions. Then $y = f(x)$ if and only if $x = g(y)$.
- A function f has an inverse function only if there is a one-to-one correspondence between inputs and outputs of f .
- The domain of a function f plays an important role in determining whether f has an inverse function g , and if so, exactly what the inverse function is.
- If a function f does not have an inverse function, then we can sometimes modify the domain of f so that it does have an inverse function. (For example, if $f(x) = x^2$ then $f : \mathbb{R} \rightarrow [0, \infty)$ does not have an inverse function while the function $f : (0, \infty) \rightarrow (0, \infty)$, with domain *restricted* to the positive real numbers, does have an inverse function.)

In the following sections we treat these ideas more rigorously.

6.2 One-to-one functions

(Ref: SH10 §7.1)

As was demonstrated in Section 6.1, the property that a function has a one-to-one correspondence between inputs and outputs is critical to our discussion.

Definition 6.2.1. A function f is said to be *one-to-one* if

$$f(x_1) = f(x_2) \quad \text{implies that} \quad x_1 = x_2$$

whenever $x_1, x_2 \in \text{Dom}(f)$.

In other words, a function is one-to-one if every output has a unique input. Equivalently, f is one-to-one if

$$f(x_1) \neq f(x_2) \quad \text{whenever} \quad x_1 \neq x_2,$$

provided that both x_1 and x_2 belong to $\text{Dom}(f)$. One-to-one functions are sometimes called *injective* functions.

Example 6.2.2. Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = 2x - 3$, is one-to-one.

Solution. Suppose that $f(x_1) = f(x_2)$. Then

$$2x_1 - 3 = 2x_2 - 3.$$

If we add 3 to both sides and afterwards divide by 2 then we obtain

$$x_1 = x_2.$$

Hence f is one-to-one. □

Example 6.2.3. Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = x^2$, is not one-to-one.


Proof. Clearly $f(-2) = f(2)$. However, $-2 \neq 2$ and so f is not one-to-one. □

There are other ways of identifying one-to-one functions. The following method is a simple geometric test.

Proposition 6.2.4 (The horizontal line test). *Suppose that f is a real-valued function defined on some subset of $\mathbb{R}$. Then the following statements are equivalent:*

- (a) f is one-to-one;
- (b) every horizontal line in the Cartesian plane intersects the graph of f at most once.

Example 6.2.5. Consider the functions graphed below.



- f is not one-to-one because the horizontal line passing through the point A cuts the graph of f more than once;
- g is one-to-one (in fact, since g is increasing, every horizontal line can cut the graph of g no more than once);
- h is also one-to-one (even though it is not always increasing).

Although not every one-to-one function is increasing (or decreasing), it is true that every increasing function is one-to-one.

Proposition 6.2.6. *If a function f is either increasing or decreasing, then f is one-to-one.*

Sketch proof. If f is increasing and $x_1, x_2 \in \text{Dom}(f)$, then

$$x_1 < x_2 \iff f(x_1) < f(x_2). \quad (6.3)$$

Now compare (6.3) with Definition 6.2.1. □

Example 6.2.7. Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = 2x^5 + x^3 + x - 10,$$

is one-to-one.

Solution. The derivative of f is given by $f'(x) = 10x^4 + 3x^2 + 1$. Since $f'(x) \geq 1$ for all real x , the function f is increasing and hence one-to-one. $\square$

Remark 6.2.8. Not every function whose derivative is only positive (or only negative) on its domain is one-to-one. For example, on $\text{Dom}(\tan)$

$$\frac{d}{dx} \tan x = \sec^2 x \geq 1,$$

but $\tan$ is *not* one-to-one, nor is it increasing on its domain. The problem here is that the domain of $\tan$ has gaps.

6.3 Inverse functions

(Ref: SH10 §7.1)

We have already mentioned the term *inverse function* in Section 6.1. In this section we define precisely what is meant by an inverse function and explore some of the relationships between a function and its inverse. Our starting point is the following theorem. It says that if f is one-to-one, then there is a function g which undoes what f does to any input x (c.f. equation (6.1)).

Theorem 6.3.1. Suppose that f is a one-to-one function. Then there is a unique function g with $\text{Dom}(g) = \text{Range}(f)$ satisfying

$$g(f(x)) = x \quad \forall x \in \text{Dom}(f) \tag{6.4}$$

and

$$f(g(x)) = x \quad \forall x \in \text{Range}(f). \tag{6.5}$$

Moreover,

$$\text{Range}(g) = \text{Dom}(f)$$

and g is one-to-one.

Proof. Suppose that f is one-to-one and let D and R denote the domain and range of f respectively. Since f is one-to-one, for every y in R there is a unique x in D such that $y = f(x)$.

Now define a function $g : R \rightarrow D$ by the following rule: if $y \in R$ then $g(y)$ is the unique x in D such that $f(x) = y$.

It is left as an exercise to show that g is one-to-one and that equations (6.4) and (6.5) hold. $\square$

The theorem allows us to define the term *inverse function*.

Definition 6.3.2. Suppose that f is a one-to-one function. Then the *inverse function* of f is the unique function g given by Theorem 6.3.1. The inverse function for f is often written as f^{-1} .

Remark 6.3.3. If f^{-1} denotes the inverse function of a one-to-one function f , then equations (6.4) and (6.5) can be expressed as

$$f^{-1}(f(x)) = x \quad \forall x \in \text{Dom}(f)$$

and

$$f(f^{-1}(x)) = x \quad \forall x \in \text{Range}(f).$$

(*Warning:* This notation is potentially confusing if not understood correctly. The function f^{-1} is *not* the reciprocal function of f (that is, $1/f(x)$). To denote the reciprocal of $f(x)$ using index notation we write $[f(x)]^{-1}$ rather than $f^{-1}(x)$.)

Remark 6.3.4. If f is a bijection and g is the inverse of a function f , then f is the inverse of g . This is due to the symmetry between f and g in the statements of Theorem 6.3.1.

If f is a one-to-one function then equation (6.5) can often be used to find an explicit formula for its inverse g .

Example 6.3.5. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = 4 - \frac{1}{3}x^3$.

- Explain why f has an inverse function.
- Find the inverse function g of f .
- Sketch f and g on the Cartesian plane.
- Use the inverse function g to solve the equation $f(x) = 13$.

Solution. (a) Since $f'(x) = -x^2 \leq 0$ for all x in $\mathbb{R}$ (with $f'(x) = 0$ only when $x = 0$), the function f is decreasing and is therefore one-to-one. Hence f has an inverse function.

(b) Let g denote the inverse function of f . Then $\text{Dom}(g) = \text{Range}(f) = \mathbb{R}$ and $\text{Range}(g) = \text{Dom}(f) = \mathbb{R}$. To find an explicit formula for $g(x)$, suppose that

$$y = g(x).$$

If we apply f to both sides of the equation, and use (6.5), then

$$f(y) = f(g(x)) = x.$$

Using the definition of f , this simplifies to

$$4 - \frac{1}{3}y^3 = x,$$


whereupon rearrangement gives

$$y = \sqrt[3]{12 - 3x}.$$

Therefore $g : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$g(x) = \sqrt[3]{12 - 3x}. \quad (6.6)$$

(c) The graphs of f and g are shown below. Note that each graph is the reflection the other in the line $y = x$.



(d) To solve the equation $f(x) = 13$, apply g to both sides. This gives

$$g(f(x)) = g(13).$$

By (6.4) and (6.6) this simplifies to

$$\begin{aligned} x &= \sqrt[3]{12 - 3 \times 13} \\ &= -3. \end{aligned}$$

□

Remark 6.3.6. As was illustrated in Example 6.3.5 (c), the graph of a one-to-one function f and its inverse g are reflections of each other in the line $y = x$. We give a proof of this fact below:

$$\begin{aligned} (x, y) \text{ lies on the graph of } f &\iff y = f(x) \\ &\iff g(y) = g(f(x)) \\ &\iff g(y) = x \\ &\iff (y, x) \text{ lies on the graph of } g. \end{aligned}$$

If a function f is not one-to-one, then we can sometimes *restrict* the domain of f so that it becomes one-to-one. The function f with restricted domain then has an inverse function.

Example 6.3.7. Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = 4 - x^2.$$

Find a restriction of f such that f becomes one-to-one. Find the inverse function g of this restriction.

Solution. A quick sketch of the graph of f (see Figure 6.2 (a)) shows that f is not one-to-one. If we restrict the domain of f to $[0, \infty)$, then the restricted function $f : [0, \infty) \rightarrow \mathbb{R}$ passes the horizontal line test and is therefore one-to-one. Let g denote the inverse of $f : [0, \infty) \rightarrow \mathbb{R}$. Then

$$g : (-\infty, 4] \rightarrow [0, \infty)$$




Figure 6.2: Two different restrictions of the function f , and their corresponding inverses (see Example 6.3.7).

since $\text{Dom}(g) = \text{Range}(f) = (-\infty, 4]$ and $\text{Range}(g) = \text{Dom}(f) = [0, \infty)$. We find an explicit formula for g :

$$\begin{aligned}
 y &= g(x) \\
 f(y) &= x && \text{(by equation 6.5)} \\
 4 - y^2 &= x \\
 y &= \sqrt{4 - x} && \text{(taking the positive root since } \text{Range}(g) = [0, \infty)).
 \end{aligned}$$

Hence $g(x) = \sqrt{4 - x}$. The graphs of f and g are sketched in Figure 6.2 (b). $\square$

Remark 6.3.8. In the solution to Example 6.3.7, one could have instead chosen the restriction $f : (-\infty, 0] \rightarrow \mathbb{R}$. The corresponding inverse g is given by

$$g : (-\infty, 4] \rightarrow (-\infty, 0], \quad g(x) = -\sqrt{4 - x^2}.$$

The graphs of f and g are given in Figure 6.2 (c).

Example 6.3.9. Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = -x^3 + 3x^2 + 24x - 13.$$

Find all intervals I , each as large as possible, such that the function $f : I \rightarrow \mathbb{R}$ (with domain restricted to I) has an inverse function.

Remark 6.3.10. If you study MATH1081 you will see that an abstract function $f : A \rightarrow B$ is said to be invertible if it is a bijection from the set A onto the set B . That is, it is both one-to-one and its range is all of its codomain B and so the problem of solving $f(x) = b$ has a unique solution for all $b \in B$. In the algebra strand of this course you will see that an $m \times n$ matrix A is invertible if the matrix equation $A\mathbf{x} = \mathbf{b}$ has a unique solution for all $\mathbf{b} \in \mathbb{R}^m$, and we'll see the idea again with linear transformations in Mathematics 1B.

In calculus we often initially define a function without precisely knowing its range, and so we usually say,

“Define $f : (a, b) \rightarrow \mathbb{R}$ by ...”.

For example, a very important function is $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \int_0^x e^{-t^2} dt$. It is easy to check that this function f is one-to-one, but it is hard to see that the range of f is $(-\sqrt{\pi}/2, \sqrt{\pi}/2)$. In calculus, when we say that this function f is invertible, we are *not* saying that f is a bijection from its domain onto its codomain $\mathbb{R}$, but rather that f is a bijection from its domain onto its *range*.

6.4 The inverse function theorem

(Ref: SH10 §§7.1, B.3)

If a function f is one-to-one then its inverse function g is also one-to-one (see Theorem 6.3.1). What other properties of f does its inverse function g inherit?

First, if f is continuous, then so is g . (A rough argument for this fact would go like this: if f is continuous then its graph can be drawn as an unbroken line. Now reflect this graph in the line $y = x$. The reflected graph is also an unbroken line. Moreover it corresponds to the graph of g . Hence g is continuous.)

What about differentiability? If a one-to-one function f is differentiable then its inverse g is not always differentiable. For example, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = x^3$, then f is differentiable everywhere. Its inverse function g is given by


$$g : \mathbb{R} \rightarrow \mathbb{R}, \quad g(x) = \sqrt[3]{x}.$$

Since

$$g'(x) = \frac{1}{\sqrt[3]{x^2}},$$

g is not differentiable at 0. So while f is differentiable everywhere, its inverse g is not.

The problem with the above example occurs where the derivative of f is zero. When a horizontal tangent to f is reflected in the line $y = x$, it becomes a vertical tangent to g and hence g is not differentiable at this point.



However, if f is a differentiable function whose derivative is never zero, its inverse g will also be differentiable.

Theorem 6.4.1 (The inverse function theorem). *Suppose that I is an open interval, $f : I \rightarrow \mathbb{R}$ is differentiable and $f'(x) \neq 0$ for all x in I . Then*

- (i) f is one-to-one and has an inverse function $g : \text{Range}(f) \rightarrow \text{Dom}(f)$,
- (ii) g is differentiable at all points in $\text{Range}(f)$, and
- (iii) the derivative of g is given by the formula

$$g'(x) = \frac{1}{f'(g(x))}$$

for all x in $\text{Range}(f)$.

Some parts of the inverse function theorem are easy to prove while others are hard. Statement (i) is true since $f'(x) \neq 0$ for x in I implies that f is either increasing on I or decreasing on I (by

the mean value theorem). Hence f is one-to-one. Statement (ii) is difficult to prove and involves a delicate limiting argument with the difference quotients of f and g .

It is important that students understand the proof of statement (iii). Suppose (by statement (ii)) that g is differentiable. Beginning with the equation

$$f(g(x)) = x$$

(see 6.5), differentiation with respect to x gives

$$f'(g(x)) \cdot g'(x) = 1$$

by the chain rule. Since f' is never zero on I , we can divide by $f'(g(x))$ to obtain

$$g'(x) = \frac{1}{f'(g(x))}$$


as required.

In the next section the inverse function theorem will be applied to the trigonometric functions.

6.5 Applications to the trigonometric functions

(Ref: SH10 §7.7)

In elementary trigonometry, the sine function gives the ratio $\frac{b}{c}$ of two side lengths of a right-angle triangle corresponding to the angle θ .



However, if we know the ratio and instead want to find the corresponding angle, we use the inverse sine function. Thus relationship between the sine and inverse sine functions is given by

$$\sin \theta = \frac{b}{c}, \quad \sin^{-1} \frac{b}{c} = \theta. \quad (6.7)$$

This relationship works in elementary trigonometry where the angle θ is acute. However, if θ is allowed to be any real number then the relationship between $\sin$ and $\sin^{-1}$ is not so straightforward. This is because the function $\sin : \mathbb{R} \rightarrow \mathbb{R}$ is not one-to-one and therefore has no inverse (see Figure 6.3).


In the next example, we define exactly what we mean by $\sin^{-1}$ and compute its derivative.

Example 6.5.1 (The inverse sine function). Since the usual sine function is not a one-to-one function, we consider instead a *restricted* sine function

$$\sin : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]$$

which is one-to-one (see Figure 6.3 (a)). This restricted function has an inverse

$$\sin^{-1} : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right],$$

Figure 6.3: Graphs of $\sin$ and $\sin^{-1}$

which is graphed in Figure 6.3 (b).

We note that $\sin^{-1}$ is an odd function, that is,

$$\sin^{-1}(-x) = -\sin^{-1} x \quad \forall x \in [-\frac{\pi}{2}, \frac{\pi}{2}].$$

We now examine the differentiability of $\sin^{-1}$. If $I = (-\frac{\pi}{2}, \frac{\pi}{2})$ then the sine function is differentiable on I and

$$\frac{d}{dx}(\sin x) = \cos x \neq 0$$

for all x in I . So by the inverse function theorem (Theorem 6.4.1), $\sin^{-1}$ is differentiable on $(-1, 1)$. While the derivative of $\sin^{-1}$ could be computed using the formula of Theorem 6.4.1 (iii), we prefer to compute it directly by implicit differentiation.

If $y = \sin^{-1} x$ then

$$\begin{aligned} \sin y &= x && \text{(by applying } \sin \text{ to both sides)} \\ \cos y \frac{dy}{dx} &= 1 && \text{(by implicit differentiation)} \\ \frac{dy}{dx} &= \frac{1}{\cos y} && \text{(since } \cos y > 0 \text{ when } -\frac{\pi}{2} < y < \frac{\pi}{2}). \end{aligned} \quad (6.8)$$

We seek a simple expression for $\cos y$ in terms of x . By the trigonometric identity $\sin^2 y + \cos^2 y = 1$, and the fact that $y = \sin^{-1} x$, we have

$$\cos y = \sqrt{1 - \sin^2 y} = \sqrt{1 - (\sin(\sin^{-1} x))^2} = \sqrt{1 - x^2}, \quad (6.9)$$

where the positive square root was taken since $\cos y > 0$. Substituting (6.9) into (6.8) gives

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}$$


and hence

$$\frac{d}{dx}(\sin^{-1}) = \frac{1}{\sqrt{1 - x^2}}$$

whenever $-1 < x < 1$.

As a corollary, note that the derivative of $\sin^{-1}$ is positive, so $\sin^{-1}$ is an increasing function.

The following table summarises the domain, range and derivatives of the restricted (and hence one-to-one) trigonometric functions and their inverses. The derivatives exist on the largest open

Figure 6.4: Graphs of $\cos^{-1}$ and $\tan^{-1}$

intervals contained in the domain of each function. Proofs of the derivatives for the inverse trigonometric functions are done in much the same way as illustrated in Example 6.5.1. It is expected that students will be able to derive these derivatives, using the method of Example 6.5.1, if required. The graphs of $\cos^{-1}$ and $\tan^{-1}$ are shown in Figure 6.4.

Function	Domain	Range	Derivative
$\sin$	$[-\frac{\pi}{2}, \frac{\pi}{2}]$	$[-1, 1]$	$\frac{d}{dx}(\sin x) = \cos x$
$\sin^{-1}$	$[-1, 1]$	$[-\frac{\pi}{2}, \frac{\pi}{2}]$	$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$
$\cos$	$[0, \pi]$	$[-1, 1]$	$\frac{d}{dx}(\cos x) = -\sin x$
$\cos^{-1}$	$[-1, 1]$	$[0, \pi]$	$\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$
$\tan$	$(-\frac{\pi}{2}, \frac{\pi}{2})$	$(-\infty, \infty)$	$\frac{d}{dx}(\tan x) = \sec^2 x$
$\tan^{-1}$	$(-\infty, \infty)$	$(-\frac{\pi}{2}, \frac{\pi}{2})$	$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$

We note also that $\sin^{-1}$ and $\tan^{-1}$ are both odd functions, but that

$$\cos^{-1}(-x) = \pi - \cos^{-1} x \quad \forall x \in [-1, 1].$$

The next example illustrates that one must be careful with the notation $\sin$ and $\sin^{-1}$. It is not always true that $\sin^{-1}(\sin x) = x$. This is because $\sin^{-1}$ is *not* the inverse of $\sin : \mathbb{R} \rightarrow \mathbb{R}$. It is only the inverse of $\sin : [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow \mathbb{R}$. In particular,

$$\sin^{-1}(\sin x) = x$$

only when $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Similar comments apply to the other trigonometric functions.

Example 6.5.2. Evaluate the following numbers without using a calculator:

- (a) $\sin^{-1}(\sin \frac{5\pi}{4})$,
- (b) $\sin(\cos^{-1} \frac{3}{5})$.

Solution. (a) We cannot write $\sin^{-1}(\sin \frac{5\pi}{4}) = \frac{5\pi}{4}$, since $\frac{5\pi}{4}$ is not in the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$. However, since $\sin^{-1}$ is an odd function,

$$\sin^{-1}\left(\sin \frac{5\pi}{4}\right) = \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right) = -\sin^{-1}\left(\frac{1}{\sqrt{2}}\right) = -\frac{\pi}{4}.$$

(b) Let θ denote $\cos^{-1} \frac{3}{5}$. Then we seek $\sin \theta$. By the trigonometric identity

$$\sin^2 \theta + \cos^2 \theta = 1,$$

we have

$$\sin \theta = \pm \sqrt{1 - \cos^2 \theta},$$

which simplifies to


$$\sin \theta = \pm \sqrt{1 - \left(\frac{3}{5}\right)^2} = \pm \frac{4}{5}.$$

Since $\theta = \cos^{-1} \frac{3}{5} > 0$ we have $\sin \theta > 0$ and hence

$$\sin \theta = +\frac{4}{5},$$

completing the question.

An alternate solution to (b) is the following. Once again, let θ denote the angle $\cos^{-1} \frac{3}{5}$. This means that $\cos \theta = \frac{3}{5}$, whereupon we construct the right-angled triangle shown below.




By Pythagoras' theorem, the length of side AB is 4. Hence $\sin \theta = \frac{4}{5}$. □

The final example uses theory from all chapters covered in the course so far.

Example 6.5.3. A man and a woman sit on opposite edges of a canyon which is 50 metres wide and 100 metres deep. The man drops a small stone from rest into the canyon. The woman watches the stone fall to the bottom of the canyon through a telescope. At what time during the stone's fall will the angle that the telescope makes with the horizontal change the fastest? (That is, when will the angular velocity of the telescope be greatest?) Assume that the air resistance experienced by the stone is negligible.

Solution. Let $x(t)$ denote the distance (in metres) that the stone has travelled exactly t seconds after the stone was dropped. Let $\theta(t)$ denote the angle that the telescope makes with the horizontal at t seconds. Let g denote acceleration due to gravity ($g \approx 9.8 \text{ ms}^{-2}$).



If $v(t)$ denotes the velocity of the stone at time t , then we have

$$\frac{dx}{dt} = v, \quad x(0) = 0, \quad (6.10)$$

$$\frac{dv}{dt} = g, \quad v(0) = 0. \quad (6.11)$$

From (6.11) we obtain $v(t) = gt + C_1$. By imposing the initial condition that $v(0) = 0$ we see that $C_1 = 0$. Hence $v(t) = gt$ and substituting this into (6.10) gives

$$\frac{dx}{dt} = gt, \quad x(0) = 0.$$

It follows that $x(t) = \frac{1}{2}gt^2 + C_2$ and one easily shows that $C_2 = 0$.

To find out when the stone hits the canyon floor, we solve the equation

$$100 = \frac{1}{2}gt^2$$

for t . Hence $t = \sqrt{200/g} \approx 4.5$. So it takes about 4.5 seconds for the stone to hit the canyon floor.

By examining the triangle in the above diagram, we see that

$$\tan \theta = \frac{x(t)}{50} = \frac{gt^2}{100}.$$

To solve the problem, we need to determine the value of t when $\frac{d\theta}{dt}$ is a maximum. Now

$$\theta = \tan^{-1} \left(\frac{gt^2}{100} \right)$$

and so

$$\frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{gt^2}{100} \right)^2} \times \frac{gt}{50}$$

by the derivative of $\tan^{-1}$ and the chain rule. This simplifies to

$$\frac{d\theta}{dt} = \frac{200gt}{10^4 + g^2t^4}.$$

For simplicity, denote $\frac{d\theta}{dt}$ by ω . We need to find when ω attains its maximum on the interval $[0, \sqrt{200/g}]$ (that is, over the time interval when the stone is falling). To locate the critical points of ω , we calculate $\frac{d\omega}{dt}$:

$$\begin{aligned}\frac{d\omega}{dt} &= \frac{d^2\theta}{dt^2} \\ &= \frac{(10^4 + g^2t^4)200g - 200gt(4g^2t^3)}{(10^4 + g^2t^4)^2} \\ &= \frac{(10^4)200g - 600g^3t^4}{(10^4 + g^2t^4)^2}.\end{aligned}$$

The stationary points of ω occur when

$$(10^4)200g - 600g^3t^4 = 0,$$

that is, when

$$t^4 = \frac{10^4}{3g^2}.$$

Since $t \geq 0$, we obtain $t = 10/\sqrt[4]{3g^2} \approx 2.42$. So the critical points of ω on $[0, \sqrt{200/g}]$ are

$$0, \quad 10/\sqrt[4]{3g^2} \quad \text{and} \quad \sqrt{200/g}.$$

The value of ω at each of these points is calculated below:

$$\omega(0) = 0, \quad \omega(10/\sqrt[4]{3g^2}) \approx 1.10, \quad \omega(\sqrt{200/g}) \approx 0.26.$$


Clearly $\omega(t)$ attains its maximum when $t = 10/\sqrt[4]{3g^2}$.

In conclusion, the angular velocity of the telescope is greatest $10/\sqrt[4]{3g^2}$ (or about 2.26) seconds after the stone is dropped. At this time, the angular velocity is about 1.10 radians per second (that is, about 62.9 degrees per second). $\square$

6.6 Maple notes

Maple knows about the inverse trigonometric functions. The functions $\sin^{-1}$, $\cos^{-1}$, $\tan^{-1}$, $\operatorname{cosec}^{-1}$, $\sec^{-1}$, $\cot^{-1}$ are written respectively as: `arcsin`, `arccos`, `arctan`, `arccsc`, `arcsec` and `arccot`. For example,

```
> plot([arctan(x), arccsc(x)], x=-10..10, y=-2..2, linestyle=[dash,solid]);
```

> # Note the gap around zero in the graph of $\arcscsc(x)$ --- please explain!

Problems for Chapter 6

Problems 6.1 : Some preliminary examples

1. [R] [V] Suppose that the functions $f : [0, \infty) \rightarrow [1, \infty)$ and $g : [1, \infty) \rightarrow [0, \infty)$ are given by $f(x) = \sqrt{1+x^2}$ and $g(x) = \sqrt{x^2-1}$.
 - a) By calculating $(f \circ g)(x)$ and $(g \circ f)(x)$, verify that g is the inverse function to f .
 - b) What are the domains of $f \circ g$ and $g \circ f$?
2. [R]
 - a) Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = 3x + 1$. Find $f^{-1}(x)$. Sketch the graph of f and the graph of its inverse function, f^{-1} , on the same diagram.
 - b) The function $g : (-\infty, 0] \rightarrow \mathbb{R}$ is defined by $g(x) = x^2 + 1$. Write down the domain and range of the inverse function g^{-1} and find a formula for $g^{-1}(x)$. Find the derivative of g^{-1} .

Problems 6.2 : One-to-one functions and 6.3 : Inverse functions

3. [R] Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = x^3 + 3x + 1$, has an inverse function whose domain is $\mathbb{R}$.
4. [R] [V] Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = 4x + \cos x$.
 - a) Show that f has an inverse function g .
 - b) By using the inverse function theorem, find $g'(2\pi)$.
5. [R] Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x^3 - 3x + 1$.
 - a) Show that $f : \mathbb{R} \rightarrow \mathbb{R}$ is not a one-to-one function.
 - b) Find all possible intervals I of $\mathbb{R}$, each as large as possible, such that the restricted function $f : I \rightarrow \mathbb{R}$ has an inverse. What is the domain of each of corresponding inverse function?
6. [H]
 - a) Can you find a quadratic function from $\mathbb{R}$ to $\mathbb{R}$ which is one-to-one?
 - b) Can you find a cubic function from $\mathbb{R}$ to $\mathbb{R}$ which is not one-to-one?

Problems 6.4 : The inverse function theorem

7. [H] For each function $f : \mathbb{R} \rightarrow \mathbb{R}$ given below, find all possible intervals I of $\mathbb{R}$, each as large as possible, such that the restricted function $f : I \rightarrow \mathbb{R}$ is one-to-one. State the range of each restricted function $f : I \rightarrow \mathbb{R}$. What can you say about existence, domain of definition, continuity and differentiability of the corresponding inverse functions?

- a) $f(x) = x(x^2 - 1)(x + 2)$
 b) $f(x) = (x + 1)^{17}$
 c) $f(x) = |x| |x + 1|$

Problems 6.5 : Applications to the trigonometric functions

8. [R] Simplify each expression without using a calculator.
 a) $\sin^{-1}(\sqrt{3}/2)$ b) $\cos(\cos^{-1}(2/5))$ c) $\sin^{-1}(\sin(5\pi/3))$
 d) $\cos^{-1}(\cos(-\pi/3))$ e) $\cos(\sin^{-1}(3/5))$ f) $\sin(\tan^{-1}(3/5))$
 g) $\sec^{-1}(2)$ h) $\sin^{-1}(\sin x)$ when $\frac{\pi}{2} \leq x \leq \frac{3\pi}{2}$
9. [R] Sketch the graph of $f : [1, 3] \rightarrow \mathbb{R}$, where $f(x) = \cos^{-1}(x - 2)$.
10. [R] Show that
 a) $\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$ b) $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$.
11. [R] Differentiate
 a) $\cos^{-1}(2x)$ b) $\sin^{-1} \sqrt{x}$ c) $\tan^{-1}(2x - 3)$.
12. [R] Prove that $\sin^{-1} x + \cos^{-1} x$ is constant. For what values of x is this valid and what is the constant?
13. [H] [V] Suppose that $f(x) = \tan^{-1} x + \tan^{-1}(1/x)$ whenever $x \neq 0$.
 a) Show that $f'(x) = 0$ whenever $x \neq 0$.
 b) Hence evaluate f on the intervals $(0, \infty)$ and $(-\infty, 0)$.
 c) How do you account for this result geometrically?
14. [H]
 a) Draw the graph of $\operatorname{cosec} x$.
 b) Show that cosec restricted to the interval $(0, \frac{\pi}{2}]$ has an inverse function. Sketch the graph of the inverse and calculate its derivative.
15. [X]
 a) Show that $2 \tan^{-1} 2 = \pi - \cos^{-1}(3/5)$.
 b) Show that $\cos^{-1}(1 - 2x^2) = 2 \sin^{-1} x$ whenever $0 \leq x \leq 1$.
 c) Suppose that $q(x) = \cos^{-1}(1 - 2x^2)$. Is q differentiable at 0?
16. [H] A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} x \tan^{-1} \left(\frac{1}{\sqrt{x}} \right) & \text{if } x > 0 \\ ax + b & \text{if } x \leq 0, \end{cases}$$

where a and b are real numbers. Find all values of a and b such that f is differentiable at 0.

17. [H] A lighthouse containing a revolving beacon is located 3 km from P , the nearest point on a straight shoreline. The beacon revolves with a constant rotation rate of 4 revolutions per minute and throws a spot of light onto the shoreline. How fast is the spot of light moving when it is (a) at P and (b) at a point on the shoreline 2 km from P ?
18. [H] [V] A picture 2 metres high is hung on a wall with its bottom edge 6 metres above the eye of the viewer. How far from the wall should the viewer stand for the picture to subtend the largest possible vertical angle with her eye?


Chapter 7

Curve sketching

There are a variety of ways to describe curves that lie in the plane. In this chapter we study curves which are described by using


- a Cartesian equation (for example, $y = x^2$ or $y = \sqrt{1 - x^2}$),
- parametric equations, and
- polar coordinates.

Each section of this chapter is devoted to one of these methods. The use of Cartesian equations will already be familiar. However, many curves cannot be easily described by Cartesian equations. The following diagrams illustrate two curves that are better described by a parameter or with polar coordinates.



An example of a parametric curve:

$$\begin{cases} x(t) = \sin t \cos t \ln |t| \\ y(t) = \sqrt{|t|} \cos t \\ t \in [-1, 1], \quad t \neq 0 \end{cases}$$



An example of a polar curve:

$$r = |\cos(4\theta)|$$

7.1 Curves defined by a Cartesian equation

(Ref: SH10 §§4.7, 4.8)

In this section we survey techniques for sketching curves that are described by a Cartesian equation of the form $y = f(x)$.

7.1.1 A checklist for sketching curves

Students will already be familiar with techniques for sketching the graph of a function f . For convenience we summarise some of the most useful here.


- Identify the domain of f . (In particular, $\text{Dom}(f)$ should not include any points that lead to division by 0 or the square root of a negative number.)
- Identify any symmetries:

$$\begin{array}{lll} f \text{ is odd if} & f(-x) = -f(x) & \forall x \in \text{Dom}(f) \\ f \text{ is even if} & f(-x) = f(x) & \forall x \in \text{Dom}(f). \end{array}$$

- Find x - and y -axis intercepts.
- Identify vertical asymptotes.
- Examine the behaviour of $f(x)$ as $x \rightarrow \pm\infty$. (This will identify any other asymptotes that exist).
- Use calculus to identify stationary points and other features, if necessary.

7.1.2 Oblique asymptotes

Consider the graph of the function f that is sketched below.



It appears that the graph of f approaches the line $y = ax + b$ as $x \rightarrow \infty$. If this is indeed the case, we say that f is *asymptotic* to the line. Since this line is neither vertical, nor horizontal, we call it an *oblique asymptote* to the function f .

Definition 7.1.1. Suppose that a and b are real numbers and that $a \neq 0$. We say that a straight line, given by the equation

$$y = ax + b,$$

is an *oblique asymptote* for a function f if

$$\lim_{x \rightarrow \infty} (f(x) - (ax + b)) = 0.$$

Example 7.1.2. Suppose that f is defined by the equation

$$f(x) = x + 1 + \frac{2}{x-2}$$

whenever $x \neq 2$. Identify all asymptotes to f and hence sketch its graph.

Solution. Clearly $f(x) \rightarrow \infty$ as $x \rightarrow 2^+$, therefore there is a vertical asymptote when $x = 2$. To identify the other asymptotes (if any), we examine the behaviour of $f(x)$ as $x \rightarrow \infty$ and as $x \rightarrow -\infty$.

Now as $x \rightarrow \infty$, $\frac{2}{x-2} \rightarrow 0$ from above. Hence we conclude that $f(x)$ approaches $x + 1$ from above. Therefore the line $y = x + 1$ is an oblique asymptote to f . (The calculation

$$\lim_{x \rightarrow \infty} (f(x) - (x + 1)) = \lim_{x \rightarrow \infty} \frac{2}{x-2} = 0$$

demonstrates from Definition 7.1.1 that this is the case.)

On the other hand, as $x \rightarrow -\infty$, $\frac{2}{x-2} \rightarrow 0$ from below. Therefore $f(x)$ approaches $x + 1$ from below.


The asymptotes for f help us construct the sketch (which is pictured below). To complete the sketch, we identify the axes intercepts. For the y -axis, we have

$$f(0) = 0.$$

For the x -axis,

$$f(x) = 0 \Rightarrow (x + 1)(x - 2) + 2 = 0 \Rightarrow x^2 - x = 0 \Rightarrow x = 0, 1.$$

Thus we have the following sketch:



(Note that we have obtained sufficient information to make a useful sketch of the graph without resorting to calculus. However, calculus would help to refine this by locating the turning points and confirming where the graph is increasing or decreasing.) $\square$

If f is a rational function of the form p/q , where p and q are polynomials and $\deg(p) > \deg(q)$, then the asymptotic behaviour of f may be determined by polynomial division.

Example 7.1.3. Suppose that f is defined by the equation

$$f(x) = \frac{x^2 - 4}{x + 1}$$

whenever $x \neq -1$. Identify all asymptotes to f and hence sketch its graph.

Solution. As in the previous example, clearly there is a vertical asymptote when $x = -1$. It is also clear that $f(x) \rightarrow \infty$ as $x \rightarrow \infty$, but the precise asymptotic behaviour of $f(x)$ as $x \rightarrow \infty$ is still not clear. However, polynomial division gives

$$\begin{array}{r} x-1 \\ x+1 \overline{) x^2 - 4} \\ \underline{x^2 + x} \\ -x - 4 \\ \underline{-x - 1} \\ -3. \end{array}$$

Hence

$$f(x) = x - 1 - \frac{3}{x+1}. \quad (7.1)$$

(See Remark 7.1.4 for an alternative method of establishing (7.1).)

Since $\frac{3}{x+1} \rightarrow 0$ from above as $x \rightarrow \infty$, we conclude that $f(x)$ approaches $x - 1$ from below as $x \rightarrow \infty$. A similar observation shows that $f(x)$ approaches $x - 1$ from above as $x \rightarrow -\infty$. So the line $y = x - 1$ is an oblique asymptote to the graph of f at both ∞ and $-\infty$.


Finally, we calculate the axes intercepts:

$$f(0) = -4$$

and

$$f(x) = 0 \quad \Rightarrow \quad x^2 - 4 = 0 \quad \Rightarrow \quad x = \pm 2.$$

Hence we obtain the following sketch.



(Again, we have obtained sufficient information for a useful sketch of the graph without the use of calculus.) □

Remark 7.1.4. We give a method for establishing equation (7.1) that uses an algebraic ‘trick’ rather than polynomial ‘long division’:

$$\begin{aligned} f(x) &= \frac{x^2 - 4}{x + 1} \\ &= \frac{x^2 - 1 - 3}{x + 1} \\ &= \frac{(x - 1)(x + 1) - 3}{x + 1} \\ &= (x - 1) - \frac{3}{x + 1}. \end{aligned}$$

7.1.3 Examples

We complete this section with three examples.

Example 7.1.5. Sketch the graph of f , where f is defined by

$$f(x) = xe^{-x^2}.$$

Solution. It is clear that $\text{Dom}(f) = \mathbb{R}$. Moreover,

$$f(-x) = -xe^{-x^2} = -f(x) \quad \forall x \in \mathbb{R},$$

so f is an odd function. Therefore, it suffices to consider $x \geq 0$; the rest of the graph can be obtained by symmetry.

Since $e^{-x^2} > 0$, it is easy to see that the only axis intercept occurs at $(0, 0)$. We look now at the behaviour of $f(x)$ as $x \rightarrow \infty$. By l'Hôpital's rule,

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{x}{e^{x^2}} = \lim_{x \rightarrow \infty} \frac{1}{2xe^{x^2}} = 0.$$

So $y = 0$ is an horizontal asymptote.

Finally, we use calculus to determine the shape of the curve. By the product rule,

$$\begin{aligned} f'(x) &= x(-2xe^{-x^2}) + e^{-x^2}(1) \\ &= (1 - 2x^2)e^{-x^2}. \end{aligned}$$

Since $e^{-x^2} > 0$ for all x in $\mathbb{R}$, we conclude that

$$f'(x) = 0 \text{ when } x = \frac{1}{\sqrt{2}},$$

while


$$f'(x) > 0 \text{ when } 0 \leq x < \frac{1}{\sqrt{2}} \quad \text{and} \quad f'(x) < 0 \text{ when } x > \frac{1}{\sqrt{2}}.$$

So f is increasing on the interval $(0, \frac{1}{\sqrt{2}})$, stationary at $\frac{1}{\sqrt{2}}$ and decreasing on $(\frac{1}{\sqrt{2}}, \infty)$. Now

$$f(1/\sqrt{2}) = \frac{1}{\sqrt{2}}e^{-1/2} = \frac{1}{\sqrt{2}e} \approx 0.429.$$

Moreover, $f'(0) = 1$, implying that the tangent line to the graph at the origin has a gradient of 1.

This information can be used to sketch the graph when $x \geq 0$, and the fact that f is odd allows us to deduce the shape of the graph when $x < 0$.



□

Example 7.1.6. Sketch the graph of f , where f is defined by

$$f(x) = x^2 - 4 + \frac{3}{x^2}.$$

The next function is important since it is later used to define the Si function (see Example 8.12.1). The Si function is used by electrical engineers for digital signal processing and by surveyors using GPS (Global Positioning System).

Example 7.1.7. Sketch the graph of f , where f is defined by


$$f(x) = \begin{cases} \frac{\sin x}{x} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0. \end{cases}$$

7.2 Parametrically defined curves

(Ref: SH10 §§10.5, 10.6)

Many moving bodies trace out a path that lies in a plane. An important problem is to describe the curve of the trajectory. We give an example.

Example 7.2.1. Suppose that a stone is thrown horizontally from the deck of Sydney Harbour Bridge at 20 metres per second, and that the air resistance experienced by the stone is negligible. Let $x(t)$ denote the horizontal distance (in metres) travelled by the stone t seconds after being thrown and let $y(t)$ denote its height (in metres) above the water. The deck is 50 metres above sea level.






Figure 7.1: Sketch for Example 7.1.6.

If we take the acceleration due to gravity to be 10 m/s^2 , then Newton's laws of motion show that

$$x(t) = 20t \quad \text{and} \quad y(t) = 50 - 5t^2 \quad (7.2)$$

when $0 < t < \sqrt{10}$ (that is, after the stone is thrown and just before it hits the water). Sketch the trajectory of the stone when $0 < t < \sqrt{10}$.

Solution. We want to express y as a function of x . By (7.2) we have

$$t = \frac{x(t)}{20} \quad \text{and} \quad y(t) = 50 - 5t^2.$$


If we substitute the equation on the left into the equation on the right then

$$y(t) = 50 - 5 \left(\frac{x(t)}{20} \right)^2$$

which becomes

$$y = 50 - \frac{x^2}{80}.$$

So the trajectory is a parabola. It is easy to see that $y = 50$ when $x = 0$ and that $x = 20\sqrt{10}$ when $y = 0$. We therefore obtain the following sketch.



□

Remark 7.2.2. The equation

$$(x(t), y(t)) = (20t, 50 - 5t^2), \quad t \in \mathbb{R}$$

is called a *parametrisation* of the parabola

$$y = 50 - \frac{x^2}{80}.$$

The variable t is called a *parameter*.

Parametrisations of curves are important to study for several reasons.

- They arise naturally with moving bodies, as illustrated in Example 7.2.1. In this context they also contain more information than a corresponding Cartesian equation (since a Cartesian equation only records the trajectory of the body, not the position of the body at any given time).

- Many curves that cannot be described by an equation of the form $y = f(x)$ can be described parametrically.
- Even if a curve is the graph of a function, it may be much easier to describe the curve parametrically than to write down an equation describing the function. (This is the case with the cycloid, which will be studied in Example 7.2.7.)
- Curves in three (or higher) dimensional space are most easily described using parameters.

7.2.1 Parametrisation of conic sections

The trajectories of many heavenly bodies (such as planets and comets) can be modelled as conic sections. In this section we will give some standard parametrisations of conic sections.

Example 7.2.3. Show that the curve given by the functions

$$x(t) = a \cos t, \quad y(t) = b \sin t, \quad 0 \leq t \leq 2\pi \quad (7.3)$$

is an ellipse. Sketch the curve, showing how the point $(x(t), y(t))$ moves as t varies from 0 to 2π .

Solution. Suppose that $t \in [0, 2\pi]$. From (7.3) we see that

$$\frac{x(t)^2}{a^2} = \cos^2 t, \quad \frac{y(t)^2}{b^2} = \sin^2 t.$$

Since $\cos^2 t + \sin^2 t = 1$,


$$\frac{x(t)^2}{a^2} + \frac{y(t)^2}{b^2} = \cos^2 t + \sin^2 t = 1.$$

Hence the point $(x(t), y(t))$ lies on the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

(Conversely, any point (x, y) that satisfies this equation can be expressed as $x = a \cos t$, $y = b \sin t$ for some t in $\mathbb{R}$.)

In fact, it is easy to see that, as t moves from 0 to 2π , $(x(t), y(t))$ traces out the entire ellipse. This is illustrated in the diagram below.



□

The table below lists some commonly used parametrisations of conic sections.

Conic section	Cartesian equation	Parametric equation
Parabola	$4ay = x^2$	$x(t) = 2at$ $y(t) = at^2$
Circle	$x^2 + y^2 = a^2$	$x(t) = a \cos t$ $y(t) = a \sin t$
Ellipse	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	$x(t) = a \cos t$ $y(t) = b \sin t$
Hyperbola	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	$x(t) = a \sec t$ $y(t) = b \tan t$

There are also other parametrisations for each of these curves. (We give a second parametrisation of the hyperbola in Chapter 10.)

In Examples 7.2.1 and 7.2.3, we sketched a parametrically defined curve by first eliminating the parameter t . It is not always possible to do this, as the next example illustrates. Such curves can be sketched with the aid of a table of values.

Example 7.2.4. Sketch the curve defined parametrically by

$$(x(t), y(t)) = (t^2 \cos t, t^2 \sin t), \quad 0 \leq t \leq 3\pi. \quad (7.4)$$

Solution. We begin by noting that the above parametric equation is similar to the parametrisation


$$(x(t), y(t)) = (a \cos t, a \sin t) \quad (7.5)$$

of the circle. The only difference is that the points in (7.5) have a fixed distance a from the origin, whereas the points in (7.4) have a variable distance t^2 from the origin. So intuitively, as t increases, the point $(x(t), y(t))$ in (7.4) rotates about origin, while at the same time getting further away from the origin. Therefore it is likely that the curve is a spiral.

The following table of values helps verify this reasoning.

t	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π	$\frac{5\pi}{2}$	3π
x	0	0	$-\pi^2$	0	$4\pi^2$	0	$-9\pi^2$
y	0	$\frac{\pi^2}{4}$	0	$\frac{9\pi^2}{4}$	0	$\frac{25\pi^2}{4}$	0

We plot these points on the Cartesian plane and interpolate between them to produce the following sketch.



□

7.2.2 Calculus and parametric curves

Given a smooth curve defined using a parameter t , one would like to calculate its gradient at any point along the curve. The following proposition enables us to do this without having to eliminate t .

Proposition 7.2.5. *Suppose that x and y are both differentiable functions of t and that y is a function of x . If $x'(t) \neq 0$ then*

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{y'(t)}{x'(t)}. \quad (7.6)$$

Proof. Since y is a function of x , we write $y = f(x)$. Hence

$$y(t) = f(x(t)).$$

Differentiating both sides with respect to t gives

$$y'(t) = \frac{df}{dx}(x(t)) x'(t)$$

by the chain rule. Since $x'(t) \neq 0$,

$$\frac{df}{dx}(x(t)) = \frac{y'(t)}{x'(t)}$$

which is equivalent to (7.6). □

Example 7.2.6. A curve is described parametrically by

$$(x(t), y(t)) = (t^2 - 1, t^3 + 2t), \quad t \in \mathbb{R}.$$

Find the equation of the tangent to the curve at the point $(0, 3)$.

Solution. By Proposition 7.2.5,

$$\begin{aligned}\frac{dy}{dx} &= \frac{y'(t)}{x'(t)} \\ &= \frac{3t^2 + 2}{2t}\end{aligned}\tag{7.7}$$

provided that $t \neq 0$.

It is easy to see that

$$t^2 - 1 = 0 \quad \text{and} \quad t^3 + 2t = 3$$


only when $t = 1$. By substituting this into (7.7), we find that the gradient of the curve at $(0, 3)$ is $5/2$. Hence the equation of the tangent at $(0, 3)$ is given by

$$y = \frac{5x}{2} + 3.$$

□

7.2.3 The cycloid and curve of fastest descent

Suppose that a point A is higher than a point B . What curve, starting at A and ending at B , would enable a particle moving under the influence of gravity (and ignoring friction) to move from A to B in the shortest time?




Curve of fastest descent.

Such a curve is known as a *curve of fastest descent* or a *brachistochrone* (which, in Greek, means ‘shortest time’).


In 1638 Galileo stated that the curve of fastest descent is the arc of a circle. This was shown to be false by Johann Bernoulli who found the correct description of the curve. In 1696 he issued a public challenge for others to find this description. Four mathematicians responded with a correct answer: Jakob Bernoulli (Johann’s brother), Gottfried Leibniz, Isaac Newton and Guillaume de l’Hôpital. The curve is the arc of a *cycloid*.

Example 7.2.7 (The cycloid). A circle of radius r rolls along the x -axis, starting from the origin as shown below. Describe the locus $(x(t), y(t))$ of the point P on the edge of the circle that satisfies $(x(0), y(0)) = (0, 0)$



The cycloid.

Solution. The solution will be built up over three steps, of which the first two correspond to the diagrams below.



Step 1. First we consider a circle of radius r centred at the origin. If the circle rotates about its centre by t radians in a clockwise direction (as shown) then

$$\begin{aligned} x(t) &= r \cos \theta = r \cos\left(\frac{3\pi}{2} - t\right) = r\left(\cos \frac{3\pi}{2} \cos t + \sin \frac{3\pi}{2} \sin t\right) = -r \sin t, \\ y(t) &= r \sin \theta = r \sin\left(\frac{3\pi}{2} - t\right) = r\left(\sin \frac{3\pi}{2} \cos t - \cos \frac{3\pi}{2} \sin t\right) = -r \cos t \end{aligned}$$

and the point P sweeps out a distance of rt .

Step 2. We now consider what happens when the circle rolls along the line $y = -r$. If the circle rotates by t radians, then the centre of the circle moves to the right by rt (corresponding to the distance that P sweeps out – see Step 1). Hence

$$x(t) = -r \sin t + rt.$$

By Step 1 we still have

$$y(t) = -r \cos t.$$

Step 3. We retain the same situation as in Step 2, except that we shift the circle up by r (so that the circle rolls along the line $y = 0$ rather than along the line $y = -r$. Thus

$$y(t) = -r \cos t + r.$$

By Step 2 we still have

$$x(t) = -r \sin t + rt.$$

Final solution. The locus of the curve is given by

$$\begin{aligned} x(t) &= r(t - \sin t), \\ y(t) &= r(1 - \cos t) \end{aligned}$$

where $t \geq 0$. □

Remark 7.2.8. The curve of fastest descent from A to B is the unique arc of an (inverted) cycloid whose tangent at A is vertical.

7.3 Curves defined by polar coordinates


(Ref: SH10 §§10.2, 10.3)

In this final section, we introduce a different coordinate system for the plane. Many equations in this system correspond to beautiful curves.

7.3.1 Polar coordinates

Every point P in plane can be specified by

- (x, y) , where x is the horizontal signed distance of P from the origin, and y is the vertical distance, or
- (r, θ) , where r is the distance of P from the origin and θ is the angle (taken in the anticlockwise direction) between OP and the positive horizontal axis.




The pair (x, y) is called the *Cartesian coordinates* of P and the pair (r, θ) are *polar coordinates* of P . (If P is the origin then $r = 0$ and θ is not defined.)

The polar coordinates (r, θ) of P are related to the Cartesian coordinates (x, y) of P by the formulae

$$\begin{aligned} x &= r \cos \theta & r &= \sqrt{x^2 + y^2} \\ y &= r \sin \theta & \tan \theta &= \frac{y}{x}, \text{ provided } x \neq 0. \end{aligned} \quad (7.8)$$


This is easily seen using trigonometry, Pythagoras' theorem and the diagram below.



When converting between the polar coordinates and Cartesian coordinates, it is best to draw a diagram.

Example 7.3.1. Suppose that the polar coordinates of P and Q are given by $(2, \frac{\pi}{4})$ and $(3, \frac{7\pi}{6})$ respectively. Find the Cartesian coordinates for P and Q .

Solution. First we plot these points on the Cartesian plane.



From the diagram, it is clear that the Cartesian coordinate of P is given by

$$\left(2 \cos \frac{\pi}{4}, 2 \sin \frac{\pi}{4}\right) = \left(\frac{2}{\sqrt{2}}, \frac{2}{\sqrt{2}}\right) = (\sqrt{2}, \sqrt{2}),$$


while the Cartesian coordinate of Q is given by

$$\left(-3 \cos \frac{\pi}{6}, -3 \sin \frac{\pi}{6}\right) = \left(-\sqrt{3}, -\frac{3}{2}\right).$$

□

Example 7.3.2. Suppose that the Cartesian coordinate of R is given by $(-2, 2\sqrt{3})$. Find polar coordinates for R .

Solution. We begin with the following diagram.



In the diagram, $\tan \alpha = \frac{2\sqrt{3}}{2} = \sqrt{3}$, so $\alpha = \frac{\pi}{3}$. Hence

$$\theta = \pi - \alpha = \frac{2\pi}{3}.$$

By Pythagoras' theorem,

$$r^2 = 2^2 + (2\sqrt{3})^2 = 16.$$

Hence in polar coordinates, R can be written as $(4, \frac{2\pi}{3})$.

□

Remark 7.3.3. Note that the point R does not have unique polar coordinates. For example

$$\left(4, \frac{2\pi}{3} + 2\pi\right) \quad \text{and} \quad \left(4, \frac{2\pi}{3} - 2\pi\right)$$

are also polar coordinates for R . In practice, we often choose the value for θ such that $-\pi < \theta \leq \pi$.

7.3.2 Basic sketches of polar curves

Many curves can be described by an equation relating the polar variables r and θ . We begin with the simplest case: when either r or θ are constant.



Example 7.3.4. Sketch the curves corresponding to the the equations

(a) $r = 3$

(b) $\theta = \frac{3\pi}{4}$.

Solution. (a) Since $r = 3$, we seek all points in the plane whose distance from the origin is 3. This is a circle of radius three centred at $(0, 0)$.

(b) This is the collection of all points P such that the angle between OP and the positive x -axis is $\frac{3\pi}{4}$. Hence the curve is the ray sketched below with the origin excluded.




□

To sketch a curve, described by the polar equation $r = f(\theta)$, in the xy -plane, it may be helpful to first sketch $r = f(\theta)$ in the $r\theta$ -plane.


Example 7.3.5. Sketch the curve whose polar equation is

$$r = 2 - \cos \theta.$$

Solution. First we graph r against θ .




The r -value of each point $a, b, c, \dots, i$ in the $r\theta$ -plane corresponds to the distance from the origin to each point $a', b', \dots, i'$ in the xy -plane.



Our task now is to interpolate between the points $a', b', \dots, i'$ to obtain a sketch of the curve. Consider what happens to r as θ moves from 0 to 2π .

- On the $r\theta$ -plane, as θ moves from 0 to π , r increases from 1 to 3. So on the xy -plane, if we travel along the curve in an anticlockwise direction from the positive to the negative x -axis, then the distance r from the origin to points on the curve increases from 1 to 3.
- On the $r\theta$ -plane, as θ moves from π to 2π , r decreases from 3 to 1. So on the xy -plane, if we travel along the curve in an anticlockwise direction from the negative to the positive x -axis, then the distance r from the origin to points on the curve decreases from 3 to 1.

These considerations lead to the final sketch.



□

We list a few more helpful tips for sketching a curve described by the polar equation $r = f(\theta)$.

- If f is an even function (that is, $f(-\theta) = f(\theta)$) then the polar curve is symmetric about the x -axis.


- If $f(\pi - \theta) = f(\theta)$ then the polar curve is symmetric about the y -axis.
- If f is 2π -periodic then it suffices to consider θ in the range $0 \leq \theta < 2\pi$.
- Sometimes it is possible to rewrite the equation $r = f(\theta)$ in terms of x and y only.

The use of calculus will be discussed in the next subsection.


Example 7.3.6. Sketch the curve that is described by the polar equation

$$r = 2 \sin \theta, \quad 0 \leq \theta \leq \pi.$$

Solution. Graphing r against θ , followed by y against x leads to the following two diagrams.



$$r = 2 \sin \theta$$



$$r = 2 \sin \theta$$

The curve looks as though it is a circle, an observation that can be confirmed by rewriting $r = 2 \sin \theta$ in terms of x and y :

$$\begin{aligned} r &= 2 \sin \theta \\ r^2 &= 2r \sin \theta && \text{(multiply by } r) \\ x^2 + y^2 &= 2y && \text{(by (7.8))} \\ x^2 + y^2 - 2y + 1 &= 1 && \text{(completing the square)} \\ x^2 + (y - 1)^2 &= 1. \end{aligned}$$

Therefore the polar curve is a circle with Cartesian centre $(0, 1)$ and radius 1. $\square$

Example 7.3.7. Sketch the curve that is described by the polar equation

$$r = 4|\sin(3\theta)|.$$

7.3.3 Sketching polar curves using calculus

Suppose that a curve can be expressed in polar form as $r = f(\theta)$. Since $x = r \cos \theta$ and $y = r \sin \theta$, the curve can also be written in parametric form (with parameter θ) as

$$x(\theta) = f(\theta) \cos \theta \quad \text{and} \quad y(\theta) = f(\theta) \sin \theta.$$

Hence,

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{y'(\theta)}{x'(\theta)} \\ &= \frac{f(\theta) \cos \theta + f'(\theta) \sin \theta}{-f(\theta) \sin \theta + f'(\theta) \cos \theta} \end{aligned}$$

by the product rule. Since $r = f(\theta)$, this is usually written as


$$\frac{dy}{dx} = \frac{r \cos \theta + \frac{dr}{d\theta} \sin \theta}{-r \sin \theta + \frac{dr}{d\theta} \cos \theta}.$$

(Rather than memorising this formula, we recommend that students understand how it is obtained.)


Knowing the value of derivative of the curve at various points allows for greater accuracy in sketching. We illustrate with one example.

Example 7.3.8. Sketch the curve described by the polar equation $r = 1 - \sin \theta$.

Solution. First we graph r against θ and then give a preliminary sketch of y against x .



$$r = 1 - \sin \theta$$



$$r = 1 - \sin \theta$$

To obtain a clearer idea of the behaviour of the curve, we calculate $\frac{dy}{dx}$. We have

$$x = r \cos \theta = (1 - \sin \theta) \cos \theta$$

and

$$y = r \sin \theta = (1 - \sin \theta) \sin \theta.$$

Hence

$$\begin{aligned} x'(\theta) &= \cos \theta (-\cos \theta) + (1 - \sin \theta)(-\sin \theta) \\ &= -\cos^2 \theta - \sin \theta + \sin^2 \theta \\ &= 2 \sin^2 \theta - \sin \theta - 1 \\ &= (2 \sin \theta + 1)(\sin \theta - 1) \end{aligned}$$

and

$$\begin{aligned} y'(\theta) &= \cos \theta - 2 \sin \theta \cos \theta \\ &= \cos \theta (1 - 2 \sin \theta). \end{aligned}$$

Thus

$$\frac{dy}{dx} = \frac{y'(\theta)}{x'(\theta)} = \frac{\cos \theta (1 - 2 \sin \theta)}{2 \sin^2 \theta - \sin \theta - 1}.$$

To locate horizontal and vertical tangents, we examine where

$$x'(\theta) = 0 \quad \text{and} \quad y'(\theta) = 0.$$

Solving $y'(\theta) = 0$ gives

$$\cos \theta = 0 \quad \text{or} \quad \sin \theta = \frac{1}{2},$$

that is,

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}.$$

Solving $x'(\theta) = 0$ gives

$$\sin \theta = 1 \quad \text{or} \quad \sin \theta = -\frac{1}{2},$$

that is

$$\theta = \frac{\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}.$$

So horizontal tangents occur when

$$\theta = \frac{3\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}$$

and vertical tangents occur when

$$\theta = \frac{7\pi}{6}, \frac{11\pi}{6}.$$


(Understanding how the derivative behaves when θ is close to $\frac{\pi}{2}$ is harder since

$$x'(\frac{\pi}{2}) = y'(\frac{\pi}{2}) = 0.$$

A full analysis is given in Remark 7.3.9.)

Finally, we observe that $\frac{dy}{dx} = -1$ when $\theta = 0$ and that $\frac{dy}{dx} = 1$ when $\theta = \pi$.

Hence we obtain the following sketch. (The abbreviations HT and VT, which appear in the diagram, stand for ‘horizontal tangent’ and ‘vertical tangent.’)



□

Remark 7.3.9. (MATH1141 only) In the above example, the curve has a ‘vertical cusp’ at the origin. This is seen by analysing $\frac{dy}{dx}$ when θ is close to $\frac{\pi}{2}$. By l’Hôpital’s rule we see that

$$\begin{aligned}\lim_{\theta \rightarrow \frac{\pi}{2}^-} \frac{x'(\theta)}{y'(\theta)} &= \lim_{\theta \rightarrow \frac{\pi}{2}^-} \frac{2 \sin^2 \theta - \sin \theta - 1}{\cos \theta - 2 \sin \theta \cos \theta} \\ &= \lim_{\theta \rightarrow \frac{\pi}{2}^-} \frac{4 \sin \theta \cos \theta - \cos \theta}{-\sin \theta - 2(\cos \theta \cos \theta - \sin \theta \sin \theta)} \\ &= \lim_{\theta \rightarrow \frac{\pi}{2}^-} \frac{0}{1} \\ &= 0.\end{aligned}$$

Hence

$$\frac{dy}{dx} = \frac{y'(\theta)}{x'(\theta)} \rightarrow \infty \quad \text{as} \quad \theta \rightarrow \frac{\pi}{2}^-.$$

By symmetry,

$$\frac{dy}{dx} = \frac{y'(\theta)}{x'(\theta)} \rightarrow -\infty \quad \text{as} \quad \theta \rightarrow \frac{\pi}{2}^+.$$


Therefore the derivative has an infinite discontinuity when $\theta = \frac{\pi}{2}$. Such a point is called a ‘cusp point.’

Remark 7.3.10. In most cases, a reasonable sketch of a polar curve can be made without appealing to calculus. As seen in the example above, while employing calculus gives a more accurate sketch, it also dramatically increases the amount of work and time required.


7.4 Maple notes

It is easy to draw parametric curves using Maple. Maple also knows about polar coordinates, but you should be aware that Maple allows $r < 0$. For example,


```
> # Draw a hyperbola via parametric specification
> plot([3*sec(t), 2*tan(t), t=-Pi/2..3*Pi/2], x=-10..10, y=-8..8, numpoints=100);
```



- > # What is the reason for the straight lines through the origin in graph above?
 > `plot([sin(5*theta), theta, theta=-Pi..Pi], coords=polar);`



- > `plot([2+cos(5*t), t, t=-Pi..Pi], coords=polar);`



Problems for Chapter 7

Problems 7.1 : Curves defined by a Cartesian equation

1. [R] Find the maximal domain and range of the function f , given by $f(x) = \sqrt{5 + 4x - x^2}$, and sketch its graph.
2. [R] Write down the period of each of the following functions f (where possible). Determine which are odd or even. Sketch the graph of each function.
 - a) $f(x) = \sin 3x$
 - b) $f(x) = 1 + \sin(2x/3)$
 - c) $f(x) = x \sin x$
 - d) $f(x) = \tan 3x$
 - e) $f(x) = \cos^2 x$
 - f) $f(x) = \sin x + \cos x$
3. [R] Suppose that f is an odd function (not everywhere zero). Determine whether each function g below is odd, even or neither.
 - a) $g(x) = x^2 f(x)$
 - b) $g(x) = x^3 f(x)$
 - c) $g(x) = x^2 + f(x)$
 - d) $g(x) = x^3 + f(x)$
 - e) $g(x) = \sin(f(x))$
 - f) $g(x) = f(\cos x)$
4. [R] For each function f , identify any vertical and oblique asymptotes and hence sketch the graph. (Do *not* use calculus.)
 - a) $f(x) = x + 2 + \frac{1}{x-3}$
 - b) $f(x) = \frac{x^2 - 2}{x + 1}$
 - c) [H] $f(x) = \frac{x^3 - 7x + 8}{x^2 + x - 6}$
5. [R] Sketch the following curves, showing their main features.
 - a) $y = x^2 + \frac{1}{x^2}$
 - b) [V] $y = \frac{x-1}{x-2}$
 - c) $y = e^{-x^2/2}$
 - d) $y = xe^{-x}$
 - e) $y^2 = x(x-4)^2$
 - f) $y = \frac{x^2}{x-2}$
 - g) $y = \frac{x^2 - 1}{x^2 - 2x}$
 - h) $y = x \cos^{-1} x$
6. [H] (*Longer rather than difficult*) Suppose that $y = \frac{3x^2 - 10x + 3}{3x^2 + 10x + 3}$.
 - a) Find the values of x for which $y \geq 0$.
 - b) Find the asymptotes.
 - c) Find the turning points.
 - d) Find the domain and range.
 - e) Sketch the graph.

Problems 7.2 : Parametrically defined curves

7. [R] Sketch the curves given by the following parametric equations. Also find, where possible, a Cartesian equation for the curve.
 - a) $x = 4 \cos t, \quad y = 5 \sin t$
 - b) $x = 3 \sec t, \quad y = 2 \tan t$
 - c) $x = t^3, \quad y = t^2$
 - d) $x = e^t \cos t, \quad y = e^t \sin t$

8. [R] For each of the curves given in parametric form by

a) $\begin{cases} x = 1 - t \\ y = 1 + t \end{cases}$ b) $\begin{cases} x = 3t + 2 \\ y = t^4 - 1 \end{cases}$ c) $\begin{cases} x = \cos t \\ y = \sin t, \end{cases}$

- i) find the points on the curve corresponding to $t = -1, 0, 1$, and 2 ;
 ii) find any point on the curve where $y = 0$;
 iii) find $\frac{dy}{dx}$ as a function of t .

9. [R] [V]

- a) Find the equation of the normal to the curve $x = \frac{t}{t+1}$, $y = \frac{t}{t-1}$ at the point P when $t = 2$.
 b) Eliminate t from the above equations and find the gradient of the normal at P using the Cartesian form.


10. [X] A curve is given in terms of the parameter t by $x = t^3$, $y = 3t^2$.

- a) What is the equation of the curve? Can you sketch it?
 b) Show that the equation of the chord joining the points with parameters t_1, t_2 is

$$(t_1^2 + t_1t_2 + t_2^2)y = 3(t_1 + t_2)x + 3t_1^2t_2^2.$$

- c) Show that the equation of the tangent at t is $ty = 2x + t^3$.
 d) Suppose that P is a point with coordinates (a, b) and that P does not lie on the curve or on the y -axis.
 i) Show that either one or three tangents may be drawn from P to the given curve. Illustrate on a sketch the region in which P must lie so that there are three tangents from P to the curve.
 ii) Assume that P lies in this region and let Q_1, Q_2, Q_3 denote the points of contact of the tangents from P to the curve. Show that the centroid of the triangle $Q_1Q_2Q_3$ is the point $(-2a, 2b)$.

11. [H] Consider a fixed circle of radius 1 centred at the origin and a smaller circle of radius $\frac{1}{4}$ initially centred at $(\frac{3}{4}, 0)$. The smaller circle rolls (without slipping) around the inside rim of the larger circle such that the centre Q of the smaller circle moves in an anticlockwise direction. A point P , fixed on the rim of the smaller circle and initially with coordinates $(1, 0)$, traces out a curve as the smaller circle moves inside the larger circle.



The goal of this question is to find the Cartesian form of the trajectory of P . Let θ denote the angle (in radians) between OQ and the positive x -axis, as shown in the above diagram.

- Explain why $\vec{OQ} = \frac{3}{4}(\cos \theta, \sin \theta)$.
- [X] Explain why $\vec{QP} = \frac{1}{4}(\cos(-3\theta), \sin(-3\theta))$.
- Show that $\vec{OP} = (\cos^3 \theta, \sin^3 \theta)$.
(You may find techniques from MATH1131 Algebra useful here.)
- Hence the trajectory of P is given by

$$x = \cos^3 \theta, \quad y = \sin^3 \theta, \quad 0 \leq \theta \leq 2\pi.$$

By using an appropriate trigonometric identity, eliminate θ to find the cartesian equation of the trajectory of P .

- Sketch the curve corresponding to this equation. (This curve is called an *astroid* after the Greek word for 'star'.)
12. [R] [V] In the 1960s two French car engineers, Paul de Casteljau and Pierre Bezier, independently discovered a remarkable new approach to parameterising curves. Let's introduce their approach by finding a quadratic curve determined by the control points $A(1, 1)$, $B(2, 2)$ and $C(3, 1)$ with respective coordinate vectors $\mathbf{a}$, $\mathbf{b}$ and $\mathbf{c}$.
- The *linear de Casteljau Bezier* curve determined by A and B is given parametrically as

$$\mathbf{p}(t) = (1 - t)\mathbf{a} + t\mathbf{b}.$$

This is called the *linear interpolation* of the vectors $\mathbf{a}$ and $\mathbf{b}$. Find a parametric vector equation for $\mathbf{p}(t)$ and evaluate $\mathbf{p}(0)$ and $\mathbf{p}(1)$.

- Similarly, the linear de Casteljau Bezier determined by B and C is

$$\mathbf{q}(t) = (1 - t)\mathbf{b} + t\mathbf{c}.$$

Find a Cartesian equation for this line, and evaluate $\mathbf{q}(1/2)$.

- c) The *quadratic de Casteljau Bezier* curve $\mathbf{r}(t)$ determined A, B and C , is the linear interpolation of the vectors $\mathbf{p}(t)$ and $\mathbf{q}(t)$:

$$\mathbf{r}(t) = (1 - t)\mathbf{p}(t) + t\mathbf{q}(t).$$

Find an explicit expression for $\mathbf{r}(t)$ and find the polynomials p_0, p_1 and p_2 such that

$$\mathbf{r}(t) = p_0(t)\mathbf{a} + p_1(t)\mathbf{b} + p_2(t)\mathbf{c}.$$

Problems 7.3 : Curves defined by polar coordinates

13. [R] The following points are given in polar coordinate form. Plot them on a diagram and find their Cartesian coordinates.
 a) $(3, 0)$ b) $(6, 7\pi/6)$ c) $(2, 7\pi/4)$
14. [R] Convert these Cartesian coordinates into polar forms with $r \geq 0$ and $-\pi < \theta \leq \pi$.
 a) $(-3, 0)$ b) $(-1, -1)$ c) $(-2, 2\sqrt{3})$
 d) $(0, 1)$ e) $(-2\sqrt{3}, 2)$ f) $(-2\sqrt{3}, -2)$
15. [R] Sketch the graph corresponding to each polar equation.
 a) $r = 4$ b) $\theta = 2$ c) $r = 3\theta$, for $\theta \geq 0$.
16. [R]
 a) Express $r = 6 \sin \theta$, where $0 \leq \theta \leq \pi$, in Cartesian form and hence draw its graph.
 b) Repeat this for $r = 2 \cos \theta$, where $-\pi/2 \leq \theta \leq \pi/2$.
17. [R] Sketch the graph corresponding to each polar equation.
 a) $r = 2 + \sin \theta$ b) $r = 3 + \cos \theta$ c) [V] $r = 2 - 2 \cos \theta$
 d) $r = 2|\cos \theta|$ e) $r = 3|\sin 6\theta|$ f) $r = |\tan \frac{\theta}{2}|$ ($-\pi < \theta < \pi$)
18. [H] [V] The hyperbolic spiral is described by the equation $r\theta = a$ whenever $\theta > 0$, where a is a positive constant. Using the fact that $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$, show that the line $y = a$ is a horizontal asymptote to the spiral. Sketch the spiral.
19. [H] Show that $r = \frac{5}{3 - 2 \cos \theta}$ is the polar equation of an ellipse by finding the Cartesian equation of the curve (and completing the square).
20. [X]
 a) For what values of θ is $r^2 = 25 \cos 2\theta$ defined?
 b) Sketch the graph of this curve. What difference would it make if you allowed negative values of r ?
21. [H] Find $\frac{dy}{dx}$ for the curves in Q16a,b,c.

Chapter 8

Integration

The problem of understanding and calculating the area of regions in the plane has a long history. Some of the main contributors to this field are Archimedes (the great Greek mathematician of antiquity), Isaac Barrow (Isaac Newton's mentor), Isaac Newton (one of the greatest mathematicians and physicists of all time), Gottfried Leibniz (Newton's contemporary), Bernhard Riemann (a German mathematician of the nineteenth century) and Henri Lebesgue (a French mathematician of the early twentieth century). In this chapter we see a method for calculating the area of regions with curved boundaries known as *Riemann integration*. We then give a startling result, known as *the fundamental theorem of calculus*, which connects the problem of calculating areas with antidifferentiation. This theorem was known to Barrow and its implications were developed by Newton, Leibniz and their disciples. A consequence of this theorem and other related results, is that the problem of calculating area is much the same as that of calculating mass, volume, work and probability. The unifying feature is a body of theory known as the *integral calculus*.

8.1 Area and the Riemann Integral

(Ref: SH10 §§5.1, 5.2, B.5)

In this section we examine the problem of defining the area of regions with curved boundaries. The main technique introduced is called *integration*, which leads to defining 'the area under the graph of a function' via the *Riemann integral*.

8.1.1 Area of regions with curved boundaries

We all know how to calculate the area of a rectangle or a triangle. The area of more exotic polygons can be computed by partitioning the polygon into triangles and then summing the areas of these triangles (see Figure 8.1 (a)). Such calculations are based on an intuitive understanding of area that takes for granted that

- the area of a rectangle is the product of its length and height,
- areas of congruent regions are equal, and
- the area of a whole region is the sum of the areas of its parts.

When precisely formulated, these points could be taken together as a *definition* of area. With this definition, one can derive a formula for the area of a right-angled triangle (see Figure 8.1 (b)), then




Figure 8.1: Calculating the area of polygons

for an arbitrary triangle (see Figure 8.1 (c)) and then calculate the area of an arbitrary polygon (as in Figure 8.1 (a)).

When it comes to calculating the area of a region with curved boundaries (see, for example, Figures 8.2 and 8.3), a more sophisticated definition of area is needed. Before asking what shape this definition might take, it is worthwhile asking the question, What *is* area?

First, we recognise that the area of a region is a non-negative real number. Also, our intuition of area informs us that it should satisfy the following properties (see Figure 8.2 for a diagram illustrating each property). We assume throughout that the regions considered are bounded.

(A1) If Ω is a region of the plane then

$$\text{area}(\Omega) \geq 0.$$

(A2) If one region Ω_1 is contained in another region Ω_2 , then

$$\text{area}(\Omega_1) \leq \text{area}(\Omega_2).$$

(A3) If the area of a region Ω is partitioned into two smaller disjoint regions Ω_1 and Ω_2 , then

$$\text{area}(\Omega) = \text{area}(\Omega_1) + \text{area}(\Omega_2).$$

(A4) If Ω_1 and Ω_2 are congruent regions then

$$\text{area}(\Omega_1) = \text{area}(\Omega_2).$$

(A5) If Ω is a rectangle of length a and height b then

$$\text{area}(\Omega) = ab.$$

These properties are *axioms* that any definition of area should satisfy.

Rather than give a definition for the area of any bounded region of the plane, we shall only give a definition for ‘the area under the graph of a function.’ The following example gives some justification for this decision.

Example 8.1.1. Consider the region Ω of Figure 8.3. By partitioning Ω with straight lines as shown, one can see by axiom (A3) that

$$\text{area}(\Omega) = \text{area}(\Omega_1) + \text{area}(\Omega_2) + \text{area}(\Omega_3).$$

By rotating and translating each subregion, we see by axiom (A4) that each of $\text{area}(\Omega_1)$, $\text{area}(\Omega_2)$ and $\text{area}(\Omega_3)$ is equal to the area under the graph of a function.

This procedure can be done for any region in the plane with a ‘reasonable’ boundary. Hence the problem of defining the area of a region with a curved boundary can be reduced to defining what is meant by ‘the area under the graph of a function.’




Figure 8.2: Properties of area





Figure 8.3: Partitioning of a region with curved boundary

Figure 8.4: Area under the graph of $f(x) = x^2$.

8.1.2 Approximations of area using Riemann sums

To guide us towards a definition for ‘the area under the graph of a function,’ we study the following example.

Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by the rule $f(x) = x^2$. Let Ω denote the region bounded by the graph of f , the x -axis and the lines $x = 0$ and $x = 1$. The region Ω is shaded in Figure 8.4 (a). The region Ω may be ‘approximated’ with rectangles as in Figure 8.4 (b) and (c). Since the regions Ω_1 and Ω_2 are composed of disjoint rectangles, their areas are easily calculated using axioms (A3) and (A5) (see Subsection 8.1.1). Moreover, however we eventually choose to define $\text{area}(\Omega)$, it is clear by axiom (A2) that we want

$$\text{area}(\Omega_1) \leq \text{area}(\Omega) \leq \text{area}(\Omega_2).$$

This gives a lower and upper bound for $\text{area}(\Omega)$.

The regions Ω_1 and Ω_2 each consist of ten rectangles. By changing the number of rectangles used, we obtain different upper and lower bounds for $\text{area}(\Omega)$.

Example 8.1.2. Find different upper and lower bounds for the area of the shaded region Ω of Figure 8.4 (a).

The following formula will help our solution:

$$\sum_{k=1}^n k^2 = 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1). \quad (8.1)$$

(This may be proved using mathematical induction. If students need to use such a formula in an exam, they will be given it).

Solution to Example 8.1.2. Let A denote the (as yet undefined) area of the region Ω . Rather than calculating an upper bound for A by using 10 rectangles (as in Figure 8.4 (c)), we use n rectangles, where n is a positive integer (see Figure 8.5). The bases of the rectangles are constructed by subdividing the interval $[0, 1]$ into n subintervals

$$\left[0, \frac{1}{n}\right], \left[\frac{1}{n}, \frac{2}{n}\right], \left[\frac{2}{n}, \frac{3}{n}\right], \dots, \left[\frac{n-1}{n}, 1\right].$$

The set $\mathcal{P}_n$, given by

$$\mathcal{P}_n = \left\{0, \frac{1}{n}, \frac{2}{n}, \frac{3}{n}, \dots, \frac{n-1}{n}, 1\right\},$$

that divides the interval $[0, 1]$ into these subintervals is called a *partition* of $[0, 1]$.




Figure 8.5: An upper Riemann sum for Example 8.1.2.

Let $\overline{R}_k$ denote the area of the k th rectangle in Figure 8.5. Then

$$\begin{aligned}
 \overline{R}_k &= \text{width} \times \text{height} \\
 &= \frac{1}{n} \times f\left(\frac{k}{n}\right) \\
 &= \frac{1}{n} \times \left(\frac{k}{n}\right)^2 \\
 &= \frac{1}{n^3} \times k^2.
 \end{aligned}$$

If $\overline{S}_{\mathcal{P}_n}(f)$ denotes the total area of the shaded region in Figure 8.5, then

$$\begin{aligned}
 \overline{S}_{\mathcal{P}_n}(f) &= \sum_{k=1}^n \overline{R}_k \\
 &= \sum_{k=1}^n \frac{1}{n^3} \times k^2 \\
 &= \frac{1}{n^3} \sum_{k=1}^n k^2 \\
 &= \frac{1}{n^3} (1^2 + 2^2 + 3^2 + \cdots + (n-1)^2 + n^2) \\
 &= \frac{1}{n^3} \times \frac{1}{6} n(n+1)(2n+1) && \text{(by formula (8.1))} \\
 &= \frac{1}{6} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right) \\
 &= \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2}.
 \end{aligned}$$

The quantity $\overline{S}_{\mathcal{P}_n}(f)$ is called the *upper Riemann sum* of f with respect to the partition $\mathcal{P}_n$. Axiom (A2) implies that

$$A \leq \overline{S}_{\mathcal{P}_n}(f) = \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2},$$

and by taking different values of n we obtain different upper bounds for A .

To compute a lower bound corresponding to n rectangles, consider Figure 8.6. The area $\underline{R}_k$ of the k th rectangle is given by

$$\underline{R}_k = \frac{1}{n} \times f\left(\frac{k-1}{n}\right) = \frac{1}{n^3}(k-1)^2.$$

The sum of all the areas of the rectangles is called the *lower Riemann sum* for the function f over the partition $\mathcal{P}_n$ and is denoted by $\underline{S}_{\mathcal{P}_n}(f)$. We see that

$$\begin{aligned} \underline{S}_{\mathcal{P}_n}(f) &= \sum_{k=1}^n \underline{R}_k \\ &= \sum_{k=1}^n \frac{1}{n^3} \times (k-1)^2 \\ &= \frac{1}{n^3} \sum_{k=1}^n (k-1)^2 \\ &= \frac{1}{n^3} (1^2 + 2^2 + 3^2 + \cdots + (n-1)^2) \\ &= \frac{1}{n^3} \times \frac{1}{6} (n-1)n(2n-1) \quad (\text{by formula (8.1)}) \\ &= \frac{1}{6} \left(1 - \frac{1}{n}\right) \left(2 - \frac{1}{n}\right) \\ &= \frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2}. \end{aligned}$$

By axiom (A2),

$$A \geq \underline{S}_{\mathcal{P}_n}(f) = \frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2}.$$

Hence for each positive integer n , the inequality

$$\frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2} \leq A \leq \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2} \quad (8.2)$$

gives an upper and lower bound for A . □

From inequality (8.2) we see that, however the area A of Ω is eventually defined, it must be squeezed between

$$\frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2} \quad \text{and} \quad \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2}$$

for every value of n . But as $n \rightarrow \infty$,

$$\frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2} \rightarrow \frac{1}{3} \quad \text{and} \quad \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2} \rightarrow \frac{1}{3}. \quad (8.3)$$

So there is only *one* number A that satisfies (8.2) for every positive integer n , namely $A = \frac{1}{3}$. Our definition of the area under the graph of a function must agree with this calculation.




Figure 8.6: A lower Riemann sum for Example 8.1.2.

Remark 8.1.3. The process of calculating upper and lower Riemann sums and taking a limit as in (8.3) is called *integration*.

Guided by the ideas introduced in this subsection, we are now ready to give a definition for the area under the graph of a function.

8.1.3 The definition of area under the graph of a function and the Riemann integral

Suppose that f is a bounded function on $[a, b]$ and that $f(x) \geq 0$ for all x in $[a, b]$. In this subsection we define what is meant by ‘the area under the graph of f from a to b ’. This is done by constructing upper and lower Riemann sums with respect to partitions of $[a, b]$.

Definition 8.1.4. A finite set $\mathcal{P}$ of points in $\mathbb{R}$ is said to be a *partition* of $[a, b]$ if

$$\mathcal{P} = \{a_0, a_1, a_2, \dots, a_n\}$$

and

$$a = a_0 < a_1 < a_2 < \dots < a_n = b.$$

Suppose that $\mathcal{P}$ is a partition of $[a, b]$. In a manner similar to Example 8.1.2, $\mathcal{P}$ is used to construct rectangles that approximate the region under the graph of f (see Figure 8.7). As illustrated in the diagram, the points of $\mathcal{P}$ need not be evenly spaced.

The area of the k th rectangle in Figure 8.7 is

$$\text{width} \times \text{height} = (a_k - a_{k-1}) \times \overline{f}_k,$$




Figure 8.7: An example of an upper Riemann sum

where $\bar{f}_k$ is the maximum value of the function f on the subinterval $[a_{k-1}, a_k]$. The *upper Riemann sum* $\bar{S}_{\mathcal{P}}(f)$ for f with respect to the partition $\mathcal{P}$ is defined by the formula

$$\bar{S}_{\mathcal{P}}(f) = \sum_{k=1}^n (a_k - a_{k-1}) \bar{f}_k. \quad (8.4)$$

Hence the upper Riemann sum corresponds to the total area of the rectangles in Figure 8.4. Likewise, the *lower Riemann sum* $\underline{S}_{\mathcal{P}}(f)$ for f with respect to the partition $\mathcal{P}$ is defined by the formula

$$\underline{S}_{\mathcal{P}}(f) = \sum_{k=1}^n (a_k - a_{k-1}) \underline{f}_k, \quad (8.5)$$

where $\underline{f}_k$ is the minimum value of f on the subinterval $[a_{k-1}, a_k]$.

We are now ready to give a definition of the area under the graph of a function f .

Definition 8.1.5 (Definition of the area under the graph of a function). Suppose that a function f is bounded on $[a, b]$ and that $f(x) \geq 0$ for all x in $[a, b]$. If there exists a *unique* real number A such that

$$\underline{S}_{\mathcal{P}}(f) \leq A \leq \bar{S}_{\mathcal{P}}(f) \quad \text{for every partition } \mathcal{P} \text{ of } [a, b], \quad (8.6)$$

then we say that A is the *area under the graph of f from a to b* .

It may not seem obvious now but, as we shall see in Section 8.2, not every bounded function f has an area under its graph in the sense of Definition 8.1.5. The following definition gives a special name to those functions that do have a well-defined area under their graph. At this point, we lift the restriction that $f(x) \geq 0$ for all x in $[a, b]$.

Definition 8.1.6 (Definition of the Riemann integral). Suppose that a function f is bounded on $[a, b]$. If there exists a *unique* real number I such that

$$\underline{S}_{\mathcal{P}}(f) \leq I \leq \overline{S}_{\mathcal{P}}(f) \quad \text{for every partition } \mathcal{P} \text{ of } [a, b], \quad (8.7)$$

then we say that f is *Riemann integrable* on the interval $[a, b]$. If f is Riemann integrable, then the unique real number I satisfying (8.7) is called the *definite integral* of f from a to b and we write

$$I = \int_a^b f(x) dx.$$

The function f is called the *integrand* of the definite integral, while the points a and b are called the *limits* of the definite integral.

Remark 8.1.7. The notation $\int_a^b f(x) dx$ is due to Leibniz. It evolved from a slightly different way of writing down lower and upper Riemann sums. For example, $\overline{S}_{\mathcal{P}}(f)$ may be written as

$$\overline{S}_{\mathcal{P}}(f) = \sum_{k=1}^n f(\overline{x}_k) \Delta x_k,$$

where $\Delta x_k = a_k - a_{k-1}$ and f attains its maximum value on $[a_{k-1}, a_k]$ at the point $\overline{x}_k$. When taking a limit, as in (8.3), Δx_k was replaced with dx and the symbol $\sum$ was replaced with an elongated ‘S’ (‘S’ stands for ‘sum’).

Remark 8.1.8. If f is Riemann integrable on $[a, b]$, then the real number $\int_a^b f(x) dx$ is the area under the graph of f from a to b , provided that $f(x) \geq 0$ for all x in $[a, b]$. A geometric interpretation of the real number $\int_a^b f(x) dx$ when f takes negative values will be discussed in Section 8.3.

Remark 8.1.9. (MATH1141 only.) If f is bounded but not continuous, then f may not attain a maximum or minimum value on the closed interval $[a_k, a_{k+1}]$. This technical difficulty can be overcome by defining the upper and lower Riemann sums in terms of suprema and infima rather than maxima and minima. The concept of a supremum and infimum will be introduced in MATH1241.

Remark 8.1.10. One should think about why Definition 8.1.5 is a good definition for the area under the graph of a function. The following questions need to be answered:

- If the area under the graph of f is well-defined by Definition 8.1.5, would it agree with any other reasonable definition for the area under the graph of f ?
- Is Definition 8.1.5 consistent with the axioms of area listed in Subsection 8.1.1?

The answer to each of these questions is ‘yes,’ but we leave it to the reader to ponder why this might be so. Some of the results given over the next few sections do confirm that our definition of area under a graph is consistent with the axioms of area.

8.2 Integration using Riemann sums

(Ref: SH10 §5.2)

In this section we calculate the area under the graph of functions by computing Riemann sums. Our notation for the upper and lower Riemann sums follows that of equations (8.4) and (8.5).




Figure 8.8: Diagrams for examples in Section 8.2.

Example 8.2.1. Suppose that $f(x) = C$ where C is a positive constant. Show that f is Riemann integrable on $[a, b]$ and calculate the area under the graph of f from a to b (see Figure 8.8 (a)).

Solution. Suppose that $\mathcal{P}$ is a partition of $[a, b]$ and that $\mathcal{P} = \{a_0, a_1, \dots, a_n\}$. Since f is constant, $\bar{f}_k = \underline{f}_k = C$ for every k between 1 and n . Therefore

$$\begin{aligned} \bar{S}_{\mathcal{P}}(f) &= \sum_{k=1}^n \bar{f}_k (a_k - a_{k-1}) \\ &= \sum_{k=1}^n C (a_k - a_{k-1}) \\ &= C(a_1 - a_0 + a_2 - a_1 + a_3 - a_2 + \dots + a_{n-1} - a_{n-2} + a_n - a_{n-1}) \\ &= C(a_n - a_0) \\ &= C(b - a). \end{aligned}$$

The same calculation gives $\underline{S}_{\mathcal{P}}(f) = C(b - a)$. Since $\mathcal{P}$ is an arbitrary partition of $[a, b]$, we have

$$\underline{S}_{\mathcal{P}}(f) = C(b - a) = \bar{S}_{\mathcal{P}} \quad \text{for every partition } \mathcal{P} \text{ of } [a, b].$$

Hence there is a unique number A satisfying (8.7), namely $A = C(b - a)$. Therefore f is Riemann integrable,

$$\int_a^b f(x) dx = C(b - a)$$

and the area under the graph of f is $C(b - a)$. □

Remark 8.2.2. The area under the graph of f is consistent with axiom (A5).

There are many other kinds of functions which are Riemann integrable. One of the largest classes consists of those functions that are bounded and piecewise continuous.

Definition 8.2.3. A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *piecewise continuous* if it is continuous on $[a, b]$ at all except perhaps a finite number of points.

Figure 8.9 illustrates the graphs of two functions f and g that are piecewise continuous on $[a, b]$. The function f is bounded on $[a, b]$ while the function g is not. According to the next theorem, the area of the shaded region under the graph of f is well-defined.




Figure 8.9: Piecewise continuous functions

Theorem 8.2.4. *If f is bounded and piecewise continuous on $[a, b]$ then f is integrable on $[a, b]$.*

Since the proof of this theorem is difficult, we omit it.

In practice, it is often hard to prove from first principles that a function f is Riemann integrable. However, if we know in advance that f is integrable on $[a, b]$ then it is much easier to calculate its definite integral from a to b . All one needs to do is show that

$$\lim_{n \rightarrow \infty} \overline{S}_{\mathcal{P}_n}(f) = \lim_{n \rightarrow \infty} \underline{S}_{\mathcal{P}_n}(f)$$

for some sequence of partitions $\mathcal{P}_n$ of $[a, b]$. If I denotes this common limit, then

$$\int_a^b f(x) dx = I.$$

Example 8.2.5. If $f(x) = x^2$, find the area under the graph of f from 0 to 1 (see Figure 8.8 (b)).

Solution. We first note that f is integrable on $[0, 1]$ since it is bounded and continuous on $[0, 1]$. Suppose that

$$\mathcal{P}_n = \{0, \frac{1}{n}, \frac{2}{n}, \frac{3}{n}, \dots, \frac{n-1}{n}, \frac{n}{n}\}.$$

Then

$$\overline{S}_{\mathcal{P}_n}(f) = \frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2} \quad \text{and} \quad \underline{S}_{\mathcal{P}_n}(f) = \frac{1}{3} - \frac{1}{2n} + \frac{1}{6n^2}$$

(see Example 8.1.2 for these calculations). Since

$$\lim_{n \rightarrow \infty} \overline{S}_{\mathcal{P}_n}(f) = \frac{1}{3} = \lim_{n \rightarrow \infty} \underline{S}_{\mathcal{P}_n}(f),$$

we conclude that

$$\int_0^1 f(x) dx = \frac{1}{3}.$$

So the area underneath the graph of f from 0 to 1 is $\frac{1}{3}$. □

Example 8.2.6. Suppose that $f(x) = 3 - x$. Calculate $\int_1^2 f(x) dx$ (see Figure 8.8 (c)).

Our final example shows that some functions are not Riemann integrable. For such a function f , the area under the graph of f is *not* defined according to Riemann integration.

Example 8.2.7. Suppose that $f : [0, 1] \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} 2 & \text{if } x \in \mathbb{Q} \\ 1 & \text{if } x \notin \mathbb{Q}. \end{cases}$$

(see Figure 8.8 (d)). Show that f is not Riemann integrable.

Solution. Suppose that $\mathcal{P}_n$ is an arbitrary partition of $[0, 1]$. Then $\overline{f}_k = 2$ while $\underline{f}_k = 1$ for every k between 1 and n . Hence

$$\underline{S}_{\mathcal{P}_n}(f) = 1 \quad \text{and} \quad \overline{S}_{\mathcal{P}_n}(f) = 2.$$

(Exercise: show that this is the case by following a method similar to Example 8.2.1). Therefore there is no *unique* real number I satisfying (8.7). We conclude that f is not Riemann integrable. $\square$

Remark 8.2.8. A more sophisticated definition of the integral, known as the *Lebesgue integral*, was developed at the beginning of the twentieth century by the French mathematician Henri Lebesgue. Lebesgue's definition of the integral allows us to integrate some functions which are not Riemann integrable. For example, if f is the function of Example 8.2.7 then

- f is not Riemann integrable and the Riemann integral $\int_0^1 f(x) dx$ is not defined;
- f is Lebesgue integrable and the Lebesgue integral $\int_0^1 f(x) dx$ is equal to 1.

The Lebesgue integral is studied in some third year mathematics courses.

8.3 The Riemann integral and signed area

(Ref: SH10 §5.5)

The definite integral

$$\int_a^b f(x) dx$$

can only be interpreted as the area under the graph of f from a to b if $f(x) \geq 0$ for all x in $[a, b]$. How should we interpret the definite integral if the integrand f has negative values on $[a, b]$?

Consider the function f graphed in Figure 8.10 (a). The upper sum from a to b , illustrated in Figure 8.10 (b), approximates $\text{area}(\Omega_1)$. However, since $f(x) < 0$ when $b < x < c$, each term in the upper sum from b to c will be negative. Hence the upper sum from b to c approximates $-\text{area}(\Omega_2)$. By increasing the number of subdivisions of $[a, c]$ and decreasing the size of each subdivision, these approximations get better and better. It follows that

$$\int_a^c f(x) dx = \text{area}(\Omega_1) - \text{area}(\Omega_2).$$

We call this quantity the *signed area* under the graph of f from a to c .

Thus there are two ways to calculate the (unsigned or absolute) area of the entire shaded region Ω of Figure 8.10 (a):

- $\text{area}(\Omega) = \int_a^c |f(x)| dx$; or
- $\text{area}(\Omega) = \int_a^b f(x) dx - \int_b^c f(x) dx.$

This kind of calculation will be illustrated in Section 8.6.




Figure 8.10: Signed area and the definite integral

8.4 Basic properties of the Riemann integral

(Ref: SH10 §5.8)

The following proposition gives a list of some of the basic properties of the Riemann integral.

Proposition 8.4.1 (Basic properties of the Riemann integral). *Suppose that f and g are integrable functions over $[a, b]$.*

(i) *If α and β are real numbers then $\alpha f + \beta g$ is integrable and*

$$\int_a^b (\alpha f + \beta g)(x) dx = \alpha \int_a^b f(x) dx + \beta \int_a^b g(x) dx.$$

(ii) *If $a < c < b$ then*

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

(iii) *If $f(x) \geq 0$ for all x in $[a, b]$ then $\int_a^b f(x) dx \geq 0$.*

(iv) *If $f(x) \leq g(x)$ for all x in $[a, b]$ then $\int_a^b f(x) dx \leq \int_a^b g(x) dx$.*

(v) *If $m \leq f(x) \leq M$ for all x in $[a, b]$ then*

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a).$$

(vi) *If $|f|$ is integrable on $[a, b]$ then*

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

Each of these properties can be interpreted geometrically (see Figure 8.11). For example, under the assumption that $f(x) \geq 0$, property (iv) says that if the graph of f lies beneath the graph of g then the area under the graph of f will be less than the area under the graph of g . Except for (i), each of these properties is easy to prove.




Figure 8.11: Geometric interpretation of Proposition 8.4.1.

Proof. We omit the proof of (i) and (ii).

(iii) If $f(x) \geq 0$ for all x in $[a, b]$, then each term in lower Riemann sum (8.5) is non-negative for any partition $\mathcal{P}$ of $[a, b]$. Hence

$$0 \leq \underline{S}_{\mathcal{P}}(f) \leq \int_a^b f(x) dx$$

by Definition 8.1.6.

(iv) Suppose that $h = g - f$. Then $h(x) \geq 0$ for all x in $[a, b]$ and by applying (iii) and (i) we have

$$0 \leq \int_a^b h(x) dx = \int_a^b (g(x) - f(x)) dx = \int_a^b g(x) dx - \int_a^b f(x) dx.$$

Rearranging the inequality gives the result.

(v) By (iv) we have

$$\int_a^b m dx \leq \int_a^b f(x) dx \leq \int_a^b M dx.$$

By Example 8.2.1, the integral on the left is $m(b-a)$ and the integral on the right is $M(b-a)$, completing the proof.

(vi) Now

$$-|f(x)| \leq f(x) \leq |f(x)| \quad \forall x \in [a, b],$$

so

$$\int_a^b -|f(x)| dx \leq \int_a^b f(x) dx \leq \int_a^b |f(x)| dx$$

by (iv). By (i), $\int_a^b -|f(x)| dx = -\int_a^b |f(x)| dx$ and so

$$-\int_a^b |f(x)| dx \leq \int_a^b f(x) dx \leq \int_a^b |f(x)| dx$$

which is equivalent to

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

by a property of the absolute value function. □

So far the definite integral

$$\int_a^b f(x) dx$$

has only been defined when $a < b$. We now extend this definition.

Definition 8.4.2. Suppose that $b < a$ and that f is integrable on $[b, a]$. Then we define

$$\int_a^b f(x) dx$$

to be the real number

$$- \int_b^a f dx$$

and we define

$$\int_a^a f(x) dx$$

to be 0.

This definition gives a more general version of Proposition 8.4.1 (ii). We omit the proof.

Proposition 8.4.3. Suppose that a , b and c are real numbers and that f is integrable over some interval containing a , b and c . Then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

8.5 The first fundamental theorem of calculus

In this section we present a surprising theoretical result that indicates a strong connection between integration (calculating areas) and differentiation (calculating gradients of tangents). This result has enormous ramifications for the problem of calculating area.

We begin by studying the area under the graph of a function. Suppose that f is a continuous function on the interval $[a, b]$. Since f is integrable on $[a, b]$ (see Theorem 8.2.4) we can define the function $F : [a, b] \rightarrow \mathbb{R}$ by the formula

$$F(x) = \int_a^x f(t) dt \quad \forall x \in [a, b]. \quad (8.8)$$

For x in $[a, b]$, the number $F(x)$ represents the (signed) area captured underneath the graph of f on the interval $[a, x]$ (see Figure 8.12 (a)). Thus F is an ‘area function’.

(Note that the limit x in (8.8) is a different symbol from the ‘dummy variable’ t . It would be incorrect and confusing to write


$$F(x) = \int_a^x f(x) dx$$

for formula (8.8).)

We investigate some of the properties of F . First, it is easy to see that if $f(t) \geq 0$ for all t in $[a, b]$ then the shaded area under f increases as x increases. That is, if f is a non-negative function then F is an increasing function.

Second, we ask whether F is continuous on $[a, b]$ and differentiable on (a, b) . Since differentiability on (a, b) implies continuity on (a, b) , we address the question of differentiability first. Consider the difference quotient

$$\frac{F(x+h) - F(x)}{h}$$

Figure 8.12: The area function F

when $x \in (a, b)$ and h is small. Now, if h is a small positive number then Figure 8.12 (b) shows that

$$\begin{aligned} F(x+h) - F(x) &= (\text{area under } f \text{ on the interval } [x, x+h]) \\ &\approx f(x) \times h, \end{aligned}$$

since $f(t) \approx f(x)$ for all t in $[x, x+h]$ by the continuity of f . Dividing both sides by h gives

$$\frac{F(x+h) - F(x)}{h} \approx f(x).$$

As h gets smaller, this approximation gets better and so we expect that

$$\lim_{h \rightarrow 0^+} \frac{F(x+h) - F(x)}{h} = f(x).$$

A similar argument may be repeated when h is a small negative number to obtain the corresponding left-hand limit. Therefore it seems true that

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = f(x).$$

This intuitive argument can be made rigorous (see the proof at the end of this section). Hence we conclude that F is continuous and differentiable on (a, b) and that $F'(x) = f(x)$ for all x in (a, b) .

The fact that a continuous function f and its ‘area function’ F are related by the differential equation

$$F'(x) = f(x)$$

has far-reaching consequences. This result is so important that it has come to be known as the *fundamental theorem of calculus*.

Theorem 8.5.1 (The first fundamental theorem of calculus). *If f is continuous function defined on $[a, b]$ then the function $F : [a, b] \rightarrow \mathbb{R}$, defined by*

$$F(x) = \int_a^x f(t) dt, \tag{8.9}$$

is continuous on $[a, b]$, differentiable on (a, b) and has derivative F' given by

$$F'(x) = f(x)$$

for all x in (a, b) .

The first fundamental theorem of calculus is a deep mathematical result which relates differentiation to integration. From the statement of the theorem we note the following points.

- The fact that F satisfies the equation $F' = f$ on (a, b) says that F is an antiderivative of f on (a, b) . Hence every continuous function f has an antiderivative F , given by formula (8.9).
- Since any two antiderivatives of f differ by a constant (see Theorem 5.9.4), every antiderivative of f is of the form

$$F + \text{constant},$$

where F is given by the integral formula (8.9). Hence the process of integration and of antidifferentiation are essentially the same. (This is surprising from a geometric point of view, where integration is used to calculate areas while differentiation is used to calculate gradients of tangents.)

- The above point suggests that integration and differentiation are inverse processes. If one takes a function f , integrates it and differentiates the result, then one obtains f again. That is, differentiation undoes what integration does to f . This is precisely expressed by the formula

$$f(x) = \frac{d}{dx} \left(\int_a^x f(t) dt \right).$$

Whether or not the converse statement is true (that is, that integration undoes what differentiation does to a function f) will be discussed in the next section.

Thus the first fundamental theorem of calculus suggests that there may be a different method for calculating the area under the graph of f ; rather than integrating via limits of Riemann sums, try integrating via antidifferentiation. This approach is investigated in the next section.

We end this section by presenting a proof of the first fundamental theorem of calculus. The proof of the differentiability of F is based on the intuitive argument given immediately prior to the statement of Theorem 8.5.1.

Proof of Theorem 8.5.1. We begin by showing that F is differentiable on (a, b) . Suppose that $x \in (a, b)$ and consider a small positive number h . By Proposition 8.4.1 (ii) one can show that

$$F(x+h) - F(x) = \int_x^{x+h} f(t) dt. \quad (8.10)$$

Since f is continuous on $[a, b]$, it attains a minimum value m_h and a maximum value M_h on $[x, x+h]$; that is,

$$m_h \leq f(t) \leq M_h \quad \forall t \in [x, x+h].$$

By Proposition 8.4.1 (v) (see also Figure 8.13),

$$m_h h \leq \int_x^{x+h} f(t) dt \leq M_h h$$

and hence

$$m_h h \leq F(x+h) - F(x) \leq M_h h.$$

by (8.10). Now $h > 0$ so

$$m_h \leq \frac{F(x+h) - F(x)}{h} \leq M_h.$$




Figure 8.13: Diagram for the proof of Theorem 8.5.1.

Since f is continuous on $[x, x+h]$,

$$\lim_{h \rightarrow 0^+} m_h = f(x) = \lim_{h \rightarrow 0^+} M_h$$

and hence

$$\lim_{h \rightarrow 0^+} \frac{F(x+h) - F(x)}{h} = f(x)$$

by the pinching theorem (Theorem 2.5.9).

In a similar manner, one can show that

$$\lim_{h \rightarrow 0^-} \frac{F(x+h) - F(x)}{h} = f(x).$$

Hence F is differentiable on (a, b) and $F'(x) = f(x)$ for all x in (a, b) .

To complete the proof, we need to show that F is continuous on $[a, b]$. Since F is differentiable on (a, b) , it is also continuous on (a, b) by Theorem 4.5.1. So it suffices to show that F is continuous at the endpoints of $[a, b]$.

We will show that F is continuous at a . Since f is continuous on $[a, b]$ there is a positive constant M such that $|f(x)| \leq M$ for all x in $[a, b]$ (see Corollary 3.4.9). Suppose that $x \in (a, b)$. By using the properties of Proposition 8.4.1,

$$\begin{aligned} |F(x) - F(a)| &= \left| \int_a^x f(t) dt - \int_a^a f(t) dt \right| \\ &= \left| \int_a^x f(t) dt \right| && \text{(Definition 8.4.2)} \\ &\leq \int_a^x |f(t)| dt && \text{(Proposition 8.4.1 (vi))} \\ &\leq \int_a^x M dt && \text{(f is bounded)} \\ &= M|x - a| && \text{(Example 8.2.1).} \end{aligned}$$

Hence

$$|F(x) - F(a)| \leq M|x - a| \rightarrow 0$$

as $x \rightarrow a^+$, or in other words, $\lim_{x \rightarrow a^+} F(x) = F(a)$. So F is continuous at a .

The proof of continuity at the endpoint b is similar. □

8.6 The second fundamental theorem of calculus

(Ref: SH10 §5.4)

The first fundamental theorem of calculus says that the processes of integration and antidifferentiation are essentially the same. The second fundamental theorem of calculus gives a fast way of calculating integrals by exploiting this similarity.

Theorem 8.6.1 (The second fundamental theorem of calculus). *Suppose that f is a continuous function on $[a, b]$. If F is an antiderivative of f on $[a, b]$ then*

$$\int_a^b f(t) dt = F(b) - F(a). \quad (8.11)$$

The second fundamental theorem of calculus is an astonishing result. To calculate the signed area under the graph of a continuous function f , all one needs to know is the value of any antiderivative of f at the endpoints a and b . To illustrate the power of this result, compare the lengthy calculation of Riemann sums in Example 8.1.2 with the example below.

Example 8.6.2. Suppose that $f(x) = x^2$ for all real numbers x . Calculate the area of the region bounded by the x -axis, the graph of f and lines $x = 0$ and $x = 1$.

Solution. Essentially we are being asked to evaluate

$$\int_0^1 f(t) dt$$

where $f(t) = t^2$ (see Figure 8.4 (a)). An antiderivative F of f is given by $F(t) = \frac{t^3}{3}$. Hence, by the second fundamental theorem of calculus,

$$\int_0^1 f(t) dt = F(1) - F(0) = \frac{1^3}{3} - \frac{0^3}{3} = \frac{1}{3}.$$

So the area of the region is $1/3$. □

Remark 8.6.3. The expression $F(b) - F(a)$ appearing in (8.11) is used so frequently that it is often abbreviated to

$$F(x) \Big|_a^b \quad \text{or} \quad \left[F(x) \right]_a^b.$$


Thus the calculation of Example 8.6.2 may be written as

$$\int_0^1 t^2 dt = \left[\frac{t^3}{3} \right]_0^1 = \frac{1}{3} - 0 = \frac{1}{3}.$$

When calculating areas, it is important to bear in mind that the integral in (8.11) represents the *signed* area.

Example 8.6.4. Calculate the area of the region bounded by the x -axis, the lines $x = 0$ and $x = 4$ and the function f given by

$$f(x) = (x - 2)(x - 4).$$



Solution. The above sketch shows that part of the region lies beneath the x -axis. So the area we seek corresponds to the integral

$$\begin{aligned}
 \int_0^4 |f(x)| dx &= \int_0^2 f(x) dx + \int_2^4 (-f(x)) dx \\
 &= \int_0^2 x^2 - 6x + 8 dx - \int_2^4 x^2 - 6x + 8 dx \\
 &= \left[\frac{x^3}{3} - 3x^2 + 8x \right]_0^2 - \left[\frac{x^3}{3} - 3x^2 + 8x \right]_2^4 \\
 &= \left[\left(\frac{8}{3} - 12 + 16 \right) - 0 \right] \\
 &\quad - \left[\left(\frac{64}{3} - 48 + 32 \right) - \left(\frac{8}{3} - 12 + 16 \right) \right] \\
 &= 8.
 \end{aligned}$$

So the area is 8 square units. □

Note that this is a different problem to: Evaluate $\int_0^4 f(x) dx$, whose solution is:

$$\begin{aligned}
 \int_0^4 f(x) dx &= \int_0^4 x^2 - 6x + 8 dx \\
 &= \left[\frac{x^3}{3} - 3x^2 + 8x \right]_0^4 \\
 &= \frac{16}{3}.
 \end{aligned}$$

We now present a proof of the second fundamental theorem of calculus. The tools used in the proof are the first fundamental theorem of calculus and a corollary of the mean value theorem.

Proof of Theorem 8.6.1. Suppose that f is continuous on $[a, b]$ and that F is an antiderivative of f on $[a, b]$. Define a function $G : [a, b] \rightarrow \mathbb{R}$ by the formula

$$G(x) = \int_a^x f(t) dt \quad \forall x \in [a, b].$$

By the first fundamental theorem of calculus (see Theorem 8.5.1), G is continuous on $[a, b]$ and $G'(x) = f(x)$ for all x in (a, b) . Hence G is also an antiderivative of f on $[a, b]$. By Theorem 5.9.4, there is a constant C such that

$$G(x) = F(x) + C \quad (8.12)$$

for all x in $[a, b]$. Note that

$$G(a) = \int_a^a f(t) dt = 0.$$

So if $x = a$ then equation (8.12) becomes

$$0 = F(a) + C.$$

Hence $C = -F(a)$. Therefore

$$\begin{aligned} \int_a^b f(t) dt &= G(b) \\ &= F(b) + C && \text{(by (8.12))} \\ &= F(b) - F(a) \end{aligned}$$

as required. $\square$

We return to the question of whether integration and differentiation are inverse processes. Suppose that f is continuous on $[a, b]$. The first fundamental theorem of calculus states that if one integrates a function f and differentiates the result, then one obtains f again. This is expressed by the formula

$$f(x) = \frac{d}{dx} \left(\int_a^x f(t) dt \right) \quad \forall x \in (a, b).$$

To be able to say that integration and differentiation are inverse operations of each other, we must ask whether the converse is true. That is, if one differentiates a function f and integrates the result, then does one obtain f again? In precise terms, we would like the formula

$$f(x) = \int_a^x f'(t) dt \quad \forall x \in (a, b) \quad (8.13)$$

to hold. Unfortunately, this converse is false. The following corollary to the second fundamental theorem of calculus says that one obtains f again only when $f(a) = 0$. Otherwise formula (8.13) is ‘out by a constant.’

Corollary 8.6.5. *Suppose that f is continuous on $[a, b]$ and has a continuous derivative on (a, b) . Then*

$$\int_a^x f'(t) dt = f(x) - f(a)$$

for all x in $[a, b]$

Proof. Apply the second fundamental theorem of calculus to the function f' on the interval $[a, x]$. $\square$

The second fundamental theorem of calculus allows a fruitful interaction between Riemann sums and antidifferentiation.

Example 8.6.6. Suppose that $f(x) = \frac{1}{1+x^2}$. By considering the lower Riemann sum of f with respect to the partition

$$\{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, \frac{n}{n}\}$$

of $[0, 1]$, show that

$$\frac{\pi}{4} = \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n}{n^2 + k^2} = \lim_{n \rightarrow \infty} \left(\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \frac{n}{n^2 + 3^2} + \dots + \frac{n}{n^2 + n^2} \right).$$

Solution. Let $\mathcal{P}_n$ denote the above partition of $[0, 1]$. Since f is a decreasing function on $[0, 1]$, the lower Riemann sum is given by

$$\begin{aligned} \underline{S}_{\mathcal{P}_n}(f) &= \sum_{k=1}^n \frac{1}{n} \frac{1}{1 + (k/n)^2} \\ &= \sum_{k=1}^n \frac{n}{n^2 + k^2}. \end{aligned}$$

Since f is bounded and continuous, it is integrable on $[0, 1]$ and hence

$$\lim_{n \rightarrow \infty} \underline{S}_{\mathcal{P}_n}(f) = \int_0^1 \frac{1}{1+x^2} dx$$

(see Remark 8.6.7). But by the second fundamental theorem of calculus,

$$\int_0^1 \frac{1}{1+x^2} dx = \left[\tan^{-1}(x) \right]_0^1 = \frac{\pi}{4}.$$

Hence

$$\frac{\pi}{4} = \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n}{n^2 + k^2}$$

as desired. □

Remark 8.6.7. The solution to Example 8.6.6 uses the following result. Suppose that f is Riemann integrable on $[a, b]$ and that $\{\mathcal{P}_n\}_{n=1}^{\infty}$ is a sequence of partitions of $[a, b]$. Let s_n denote the maximum size of the subintervals generated by $\mathcal{P}_n$. If $\lim_{n \rightarrow \infty} s_n = 0$ then

$$\lim_{n \rightarrow \infty} \underline{S}_{\mathcal{P}_n}(f) = \int_a^b f(x) dx = \lim_{n \rightarrow \infty} \overline{S}_{\mathcal{P}_n}(f).$$

8.7 Indefinite integrals

(Ref: SH10 §5.6)

The fundamental theorems of calculus say that integration and antidifferentiation are, in a certain sense, the same process. In particular, if a F is an antiderivative of f then

$$\int_a^x f(t) dt = F(x) + C$$

for some suitable constant C . If we have no particular interest in the interval $[a, x]$ and merely want to indicate that F is an antiderivative of f , then we write

$$\int f(t) dt = F(t) + C. \quad (8.14)$$

An integral expressed in this way, without limits, is called an *indefinite integral*. The constant C in (8.14) is called the *constant of integration*.

The table of antiderivatives given in Section 5.9 is equivalent to the following table of indefinite integrals.

Indefinite integrals	
$\int x^r dx = \frac{1}{r+1}x^{r+1} + C,$	where r is a rational number and $r \neq -1$
$\int \sin x dx = -\cos x + C$	
$\int \cos x dx = \sin x + C$	
$\int e^{ax} dx = \frac{1}{a}e^{ax} + C$	
$\int \frac{f'(x)}{f(x)} dx = \ln f(x) + C$	

Since differentiation and indefinite integration (or antidifferentiation) are inverse processes, they have many analogous properties. For example, differentiation behaves *linearly*, which means that

$$(f + g)'(x) = f'(x) + g'(x)$$

and

$$(\alpha f)'(x) = \alpha \times f'(x)$$

for every real number α and every differentiable function f and g . The next proposition says that indefinite integration is also behaves linearly. (We have already shown in Proposition 8.4.1 that definite integration is linear.)

Proposition 8.7.1. *Suppose that f and g are integrable functions and that α is a real constant. Then*

$$\int (f + g)(x) dx = \int f(x) dx + \int g(x) dx \quad (8.15)$$

and

$$\int (\alpha f)(x) dx = \alpha \int f(x) dx. \quad (8.16)$$

Proof. We prove (8.15) only and leave the proof of 8.16 as an exercise. Suppose that F , G and H are antiderivatives of f , g and $f + g$ respectively. Then

$$H'(x) = (f + g)(x) = f(x) + g(x) = F'(x) + G'(x)$$

by the linearity of differentiation. By expressing the equation $H' = F' + G'$ in terms of indefinite integrals we obtain (8.15) as desired. $\square$

Example 8.7.2. Find

$$\int (10x^4 + 8 \sin x - 9e^x) dx.$$

Proof. By Proposition 8.7.1,

$$\begin{aligned} \int (10x^4 + 8 \sin x - 9e^x) dx &= 10 \int x^4 dx + 8 \int \sin x dx - 9 \int e^x dx \\ &= 10 \left(\frac{1}{5}x^5 + C_1 \right) + 8(-\cos x + C_2) - 9(e^x + C_3) \\ &= 2x^5 - 8 \cos x - 9e^x + C \end{aligned}$$

where C_1 , C_2 and C_3 are constants of integration and $C = 10C_1 + 8C_2 - 9C_3$. $\square$

Remark 8.7.3. It is standard practice to combine all constants of integration and simply write ‘ $+C$ ’. Thus the calculation in Example 8.7.2 may be written as

$$\begin{aligned} \int (10x^4 + 8 \sin x - 9e^x) dx &= 10 \int x^4 dx + 8 \int \sin x dx - 9 \int e^x dx \\ &= 10 \times \frac{1}{5}x^5 + 8(-\cos x) - 9e^x + C \\ &= 2x^5 - 8 \cos x - 9e^x + C. \end{aligned}$$

8.8 Integration by substitution

(Ref: SH10 §5.7)

When differentiating the composition of two functions, we apply the chain rule. By reversing this process with integration, we obtain a technique known as *integration by substitution*.

The basic idea is the following. Suppose that F is an antiderivative for f . Then

$$\int f(g(x)) \cdot g'(x) dx = F(g(x)) + C, \quad (8.17)$$

as can be easily verified by differentiating the right-hand side via the chain rule. On the other hand,

$$\int f(u) du = F(u) + C. \quad (8.18)$$

If we make the substitution $u = g(x)$ then the right-hand sides of (8.17) and (8.18) are the same. This suggests the substitution $du = g'(x) dx$, so that the left-hand side of (8.17) is transformed into the left-hand side of (8.18). These mechanical substitutions can be used to transform the complicated integral into a simpler one.

Example 8.8.1. Calculate the indefinite integral

$$\int (2x - 3) \cos(x^2 - 3x + 4) dx$$

by the method of substitution.

Solution. We begin with the substitution

$$u = x^2 - 3x + 4.$$

Since

$$\frac{du}{dx} = 2x - 3,$$

we also make the substitution

$$du = (2x - 3) dx.$$

Then

$$\begin{aligned} \int (2x - 3) \cos(x^2 - 3x + 4) dx &= \int \cos(u) du \\ &= \sin u + C \\ &= \sin(x^2 - 3x + 4) + C. \end{aligned}$$

We can verify our answer by differentiating $\sin(x^2 - 3x + 4)$ via the chain rule. □

A similar technique can be used to evaluate definite integrals.

Example 8.8.2. Evaluate the definite integral

$$\int_0^2 3x^2 \sqrt{1 + x^3} dx.$$

Solution. If $u = 1 + x^3$ then $\frac{du}{dx} = 3x^2$. Hence we make the following substitutions:

$$\begin{aligned} u &= 1 + x^3 \\ du &= 3x^2 dx \\ u &= 1 \text{ (when } x = 0\text{)} \\ u &= 9 \text{ (when } x = 2\text{)}. \end{aligned}$$

Thus

$$\begin{aligned} \int_0^2 3x^2 \sqrt{1 + x^3} dx &= \int_1^9 \sqrt{u} du \\ &= \left[\frac{2}{3} u^{3/2} \right]_1^9 \\ &= \frac{2}{3} (9^{3/2} - 1^{3/2}) \\ &= \frac{52}{3}. \end{aligned}$$

□

In general, the reversal of the chain rule via integration gives rise to the *change of variables formula*

$$\boxed{\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du,} \quad (8.19)$$

The substitutions

$$u = g(x), \quad du = g'(x) dx,$$

are only to be understood within this context. Precise conditions under which formula (8.19) holds is given by the following theorem.

Theorem 8.8.3 (Change of variables formula). *Suppose that g is a differentiable function such that g' is continuous on $[a, b]$. If f is continuous on any interval I containing $g(a)$ and $g(b)$ then the change of variables formula (8.19) holds.*

Proof. Since f is continuous on I it has an antiderivative $F : I \rightarrow \mathbb{R}$ by the first fundamental theorem of calculus. By the chain rule (Theorem 4.2.2),

$$\begin{aligned} (F \circ g)'(x) &= F'(g(x))g'(x) \\ &= f(g(x))g'(x). \end{aligned}$$

So by two applications of the second fundamental theorem of calculus,

$$\begin{aligned} \int_a^b f(g(x))g'(x) dx &= (F \circ g)(b) - (F \circ g)(a) \\ &= F(g(b)) - F(g(a)) \\ &= \int_{g(a)}^{g(b)} f(u) du, \end{aligned}$$

completing the proof. □

Finding a fruitful substitution is not always easy (and sometimes not even possible). In principle, one looks for a function g in the integrand whose derivative g' is also in the integrand. Once such a function g is identified, try the substitution $u = g(x)$. As illustrated below, it may be necessary to manipulate the integrand to implement this strategy.

Example 8.8.4. Evaluate

$$\int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx.$$

Solution. Note that

$$\frac{d}{dx}(\sqrt{x}) = \frac{1}{2\sqrt{x}}$$

and so if $g(x) = \sqrt{x}$ then $g'(x)$ appears in the integrand (up to the constant factor $1/2$). So the substitution we use is

$$u = \sqrt{x}, \quad du = \frac{1}{2\sqrt{x}} dx.$$

Hence

$$\begin{aligned} \int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx &= 2 \int \frac{e^{\sqrt{x}}}{2\sqrt{x}} dx \\ &= 2 \int e^u du \\ &= 2e^u + C \\ &= 2e^{\sqrt{x}} + C. \end{aligned}$$

□

There are occasions when we make a substitution of the form $x = g(u)$ rather than $u = g(x)$.

Example 8.8.5. Evaluate the integral I , where

$$I = \int_4^9 \frac{dx}{2 + \sqrt{x}}.$$

Solution. If the integral were

$$\int_4^9 \frac{dx}{2 + x},$$

then it would be easy to evaluate. The difficulty with I clearly lies with the square root appearing in the integrand. To remove the square root, we use the substitution $x = u^2$. This gives

$$\begin{aligned} x &= u^2 \\ dx &= 2u \, du \\ u &= 2 \text{ when } x = 4 \\ u &= 3 \text{ when } x = 9. \end{aligned}$$

(Note that we are implicitly assuming that $u > 0$. It would be incorrect to take the limits in the variable u to be -2 and 3 , for example.) Hence

$$\begin{aligned} I &= \int_2^3 \frac{2u \, du}{2 + u} \\ &= 2 \int_2^3 \frac{u}{2 + u} \, du \\ &= 2 \int_2^3 \frac{2 + u - 2}{2 + u} \, du \\ &= 2 \int_2^3 1 - \frac{2}{2 + u} \, du \\ &= 2 \left[u - 2 \ln |2 + u| \right]_2^3 \\ &= 2 + 4 \ln \left(\frac{4}{5} \right). \end{aligned}$$

□

If an integrand possesses symmetry, then the corresponding definite integral may be easy to evaluate. For example, if f is an odd function and we integrate f over a balanced interval $[-a, a]$, then the ‘negative area’ cancels out the ‘positive area’ and consequently the integral is zero (see Figure 8.14). This result is proved using integration by substitution.


Proposition 8.8.6. Suppose that f is a continuous function and a is a real number.

(i) If f is even then

$$\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx.$$

(ii) If f is odd then

$$\int_{-a}^a f(x) \, dx = 0.$$

Figure 8.14: Integration with an odd function f over $[-a, a]$

(iii) If f is periodic with period T then

$$\int_a^{a+T} f(x) dx = \int_0^T f(x) dx.$$

Proof. We prove (ii) only and leave the proofs of (i) and (iii) as an exercise. Suppose that f is an odd function. By breaking up the integral and using a substitution we have

$$\begin{aligned} \int_{-a}^a f(x) dx &= \int_{-a}^0 f(x) dx + \int_0^a f(x) dx \\ &= -\int_a^0 f(-u) du + \int_0^a f(x) dx && \text{(substituting } u = -x) \\ &= \int_0^a f(-u) du + \int_0^a f(x) dx \\ &= -\int_0^a f(u) du + \int_0^a f(x) dx && \text{(since } f(-u) = -f(u)) \\ &= 0, \end{aligned}$$

completing the proof of (ii). □

Example 8.8.7. Find

$$\int_{\frac{\pi}{17}}^{\frac{\pi}{17}+2\pi} \sin^5 x dx.$$

Solution. If $f(x) = \sin^5 x$ then f is both odd and periodic with period 2π . Hence

$$\begin{aligned} \int_{\frac{\pi}{17}}^{\frac{\pi}{17}+2\pi} \sin^5 x dx &= \int_0^{2\pi} \sin^5 x dx && \text{(by Proposition 8.8.6 (iii) when } a = \frac{\pi}{17}) \\ &= \int_{-\pi}^{\pi} \sin^5 x dx && \text{(by Proposition 8.8.6 (iii) when } a = -\pi) \\ &= 0 && \text{(by Proposition 8.8.6 (ii)).} \end{aligned}$$

□

8.9 Integration by parts

(Ref: SH10 §8.2)

Suppose that u and v are two functions of a variable x . The product rule for differentiation gives

$$\frac{d}{dx}(uv) = v \frac{du}{dx} + u \frac{dv}{dx}.$$

By integrating both sides with respect to x we obtain

$$uv = \int v \frac{du}{dx} dx + \int u \frac{dv}{dx} dx.$$

Rearranging this equation gives the *integration by parts* formula

$$\boxed{\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx.} \quad (8.20)$$

For definite integrals the formula becomes

$$\boxed{\int_a^b u \frac{dv}{dx} dx = [uv]_a^b - \int_a^b v \frac{du}{dx} dx.} \quad (8.21)$$

by the second fundamental theorem of calculus.

Integration by parts allows one to evaluate the integral of a product of two functions. Its success depends on choosing u and v such that

$$\int_a^b v \frac{du}{dx} dx \quad \text{is easier to calculate than} \quad \int_a^b u \frac{dv}{dx} dx.$$

Once u and v are chosen, simply apply formula (8.20) or (8.21).

Remark 8.9.1. The shorthand version

$$\boxed{\int uv' = uv - \int vu'}$$

of formulae (8.20) and (8.21) is easier to remember.

Example 8.9.2. Evaluate the integral

$$\int x e^{2x} dx.$$

Solution. We set

$$\begin{aligned} u &= x & v &= \frac{1}{2}e^{2x} \\ u' &= 1 & v' &= e^{2x}. \end{aligned}$$

Then integration by parts (see formula (8.20)) gives

$$\begin{aligned} \int x e^{2x} dx &= \frac{1}{2} x e^{2x} - \int \frac{1}{2} e^{2x} \times 1 dx \\ &= \frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} + C, \end{aligned}$$

completing the solution.

(Note that if we had instead set

$$\begin{aligned} u &= e^{2x} & v &= \frac{1}{2}x^2 \\ u' &= 2e^{2x} & v' &= x. \end{aligned}$$

then integration by parts gives

$$\int x e^{2x} dx = \frac{1}{2} x^2 e^{2x} - \int x^2 e^{2x} dx$$

and the integral on the right-hand side is harder to evaluate than the one we started with.) $\square$

Sometimes integration by parts must be applied several times to evaluate the integral.

Example 8.9.3. Find the integral I , where

$$I = \int_0^\pi x^2 \sin x dx.$$

Example 8.9.4. Evaluate the indefinite integral I , where

$$I = \int e^x \sin x dx$$

Integration by parts can also be used to evaluate some integrals of the form $\int f(x) dx$. The trick is to rewrite the integral as

$$\int f(x) \times 1 dx.$$

Example 8.9.5. Evaluate the integral

$$\int \cos^{-1} x dx.$$

8.10 Improper integrals


(Ref: SH10 §11.7)

So far we have defined definite integrals for bounded functions over finite intervals. In this section we look at integrals over infinite intervals. Such integrals have many applications in probability, statistics, physics and engineering.

To give meaning to the *improper integral*,

$$\int_a^\infty f(x) dx,$$

we examine the behaviour of $\int_a^R f(x) dx$ as $R \rightarrow \infty$, as illustrated below.



Definition 8.10.1. (a) Suppose that there is a real number L such that

$$\int_a^R f(x) dx \rightarrow L$$

as $R \rightarrow \infty$. Then f is said to be *integrable* over $[a, \infty)$ and the integral $\int_a^\infty f(x) dx$ is said to be *convergent*. Moreover, we write

$$\int_a^\infty f(x) dx = L.$$

(b) Suppose that

$$\int_a^R f(x) dx$$

does not have a limit as $R \rightarrow \infty$. Then f we say that f *not integrable* over (a, ∞) and we say that the integral $\int_a^\infty f(x) dx$ is *divergent*.

One can write down a similar definition for improper integrals of the form $\int_{-\infty}^b f(x) dx$.

Example 8.10.2. Evaluate the following improper integrals or show that they diverge:

(a) $\int_0^\infty \frac{1}{x^2 + 1} dx;$

(b) $\int_1^\infty \frac{1}{\sqrt{x}} dx;$

(c) $\int_{-\infty}^0 e^{2x} dx.$

Solution. (a) If R is a real number then

$$\begin{aligned} \int_0^R \frac{1}{x^2 + 1} dx &= \left[\tan^{-1} \right]_0^R \\ &= \tan^{-1} R - 0 \\ &\rightarrow \frac{\pi}{2} \end{aligned}$$

as $R \rightarrow \infty$. Hence the improper integral converges and

$$\int_0^\infty \frac{1}{x^2 + 1} dx = \frac{\pi}{2}.$$

(b) If R is a real number then

$$\begin{aligned} \int_1^R x^{-1/2} dx &= \left[2x^{1/2} \right]_1^R \\ &= 2\sqrt{R} - 2 \\ &\rightarrow \infty \end{aligned}$$

as $R \rightarrow \infty$. Hence the improper integral diverges.

□

We now consider improper integrals whose interval of integration is the entire real line.

Definition 8.10.3. We say f is integrable over $(-\infty, \infty)$ if f is integrable over both $(-\infty, 0)$ and $(0, \infty)$. In this case we write

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^0 f(x) dx + \int_0^{\infty} f(x) dx.$$

If f is not integrable on either of the intervals $(-\infty, 0)$ or $(0, \infty)$, then we say that the improper integral

$$\int_{-\infty}^{\infty} f(x) dx$$

diverges.

The following example shows that the improper integral $\int_{-\infty}^{\infty} f(x) dx$ may diverge even if

$\lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$ exists.

Example 8.10.4. Consider the improper integral

$$\int_{-\infty}^{\infty} x dx.$$

If R is a positive number then

$$\int_{-R}^R x dx = 0$$

since the function f , given by $f(x) = x$, is odd (see Proposition 8.8.6). Hence

$$\lim_{R \rightarrow \infty} \int_{-R}^R x dx = 0.$$

However

$$\int_0^R x dx = \left[\frac{1}{2} x^2 \right]_0^R = \frac{R^2}{2} \rightarrow \infty$$

as $R \rightarrow \infty$. Hence

$$\int_0^{\infty} x dx$$

diverges, and by Definition 8.10.3

$$\int_{-\infty}^{\infty} x dx$$

diverges also.

The following proposition determines the convergence or divergence of improper integrals whose integrand is a power of x . Such integrals will be used frequently in Section 8.11

Proposition 8.10.5 (Convergence and divergence of p -integrals). *The improper integral*

$$\int_1^{\infty} \frac{1}{x^p} dx$$

is convergent if $p > 1$ and divergent if $p \leq 1$.

Proof. If $p \neq 1$ then

$$\begin{aligned}\int_1^R x^{-p} dx &= \left[\frac{x^{1-p}}{1-p} \right]_1^R \\ &= \frac{R^{1-p} - 1}{1-p} \\ &\rightarrow \begin{cases} \frac{1}{p-1} & \text{when } 1-p < 0 \\ \infty & \text{when } 1-p > 0 \end{cases}\end{aligned}$$

as $R \rightarrow \infty$. Hence the integral converges when $1-p < 0$ (that is, when $p > 1$) and diverges when $1-p > 0$ (that is, when $p < 1$).

In the case when $p = 1$, we have

$$\int_1^R \frac{1}{x} dx = \left[\ln x \right]_1^R = \ln R - \ln 1 \rightarrow \infty \quad \text{as } R \rightarrow \infty.$$

Hence the integral diverges when $p = 1$. □

Remark 8.10.6. While you may be familiar with the fact that the $\ln$ function is unbounded, the proof that $\ln R \rightarrow \infty$ as $R \rightarrow \infty$ requires the use of Riemann sums and is given later in §9.2.

8.11 Comparison tests for improper integrals

(Ref: SH10 §11.7)

Some of the most important improper integrals, such as


$$\int_0^\infty e^{-x^2} dx$$

have integrands without any antiderivative in the elementary functions. In these cases, determining convergence via a computation like

$$\int_a^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_a^R f(x) dx = \lim_{R \rightarrow \infty} F(R) - F(a),$$

(where F is an antiderivative of f) is not a viable strategy. Instead, we compare the integral of f with another integral $\int_a^\infty g(x) dx$ whose behaviour is known. The basic idea is as follows.

Suppose that $0 \leq f(x) \leq g(x)$ whenever $x > a$, as illustrated below.



Obviously the area under the graph of g is greater than that under the graph of f . Hence if the area under g is finite then the area under f is also finite. If the area under f is infinite then so too is the area under g . The next theorem expresses this conclusion in terms of improper integrals.

Theorem 8.11.1 (The comparison test). *Suppose that f and g are integrable functions and that $0 \leq f(x) \leq g(x)$ whenever $x > a$.*

(i) *If $\int_a^\infty g(x) dx$ converges then $\int_a^\infty f(x) dx$ converges.*

(ii) *If $\int_a^\infty f(x) dx$ diverges then $\int_a^\infty g(x) dx$ diverges.*

Proof. We begin by proving (i). Suppose that f and g are integrable functions, that $0 \leq f(x) \leq g(x)$ for all x in $[a, \infty)$ and that $\int_a^\infty g(x) dx$ converges. Then

$$0 \leq \int_a^R f(x) dx \leq \int_a^R g(x) dx$$

whenever $R > 0$ by Proposition 8.4.1. Since the integral

$$\int_a^R f(x) dx$$

increases as R increases and is bounded above by $\int_a^\infty g(x) dx$, it has a limit as $R \rightarrow \infty$. Hence $\int_a^\infty f(x) dx$ is also convergent.

The proof of (ii) is left as an exercise. □

When applying the comparison test, we often compare an improper integral I with

$$\int_1^\infty \frac{1}{x^p} dx,$$

since the behaviour of this second integral is known (see Proposition 8.10.5). The value of p is chosen by analysing the dominant terms appearing in the integrand of I . The next example illustrates this procedure.

Example 8.11.2. Determine whether or not the following improper integrals converge.

(a) $\int_1^\infty \frac{1}{x^3 + 1} dx$

(b) $\int_2^\infty \frac{x + 2}{x^{3/2} - 1} dx.$

Solution. (a) When x is large, the dominant term in the denominator of

$$\frac{1}{x^3 + 1}$$

is x^3 . So intuitively,

$$\frac{1}{x^3 + 1} \approx \frac{1}{x^3}$$

for large x . Consequently we aim to compare

$$\int_1^\infty \frac{1}{x^3 + 1} dx \quad \text{with} \quad \int_1^\infty \frac{1}{x^3} dx.$$

To make this comparison rigorous, we seek an appropriate inequality. Note that $x^3 + 1 > x^3$ for all $x > 1$ and so

$$\frac{1}{x^3 + 1} < \frac{1}{x^3}$$

for all $x > 1$. Since

$$\int_1^\infty \frac{1}{x^3} dx$$

converges (see Proposition 8.10.5),

$$\int_1^\infty \frac{1}{x^3 + 1} dx$$

also converges by the comparison test. □

There are many examples where the ‘dominant term analysis’ is straightforward but it is difficult to obtain an appropriate inequality for successful use of the comparison test. This difficulty can sometimes be overcome by multiplying one of the integrands by a ‘fudge factor’. The name ‘fudge factor’ might sound like we’re about to ‘cheat’ and do something that is not mathematically rigorous, but despite the name we really will be finding an inequality that will rigorously prove convergence or divergence of our improper integral.

Example 8.11.3. Determine whether or not the improper integral

$$\int_2^\infty \frac{1}{x^{3/2} - 1} dx.$$

converges.

Solution. Dominant term analysis suggests that

$$\frac{1}{x^{3/2} - 1} \approx \frac{1}{x^{3/2}}$$

when x is large. So we compare

$$\int_2^\infty \frac{1}{x^{3/2} - 1} dx \quad \text{with} \quad \int_2^\infty \frac{1}{x^{3/2}} dx.$$

Now the integral to the right converges (by Proposition 8.10.5) but it is clear that

$$\frac{1}{x^{3/2} - 1} > \frac{1}{x^{3/2}}$$

when $x > 2$, so we cannot immediately apply the comparison test.

However, by multiplying the integrand on the right with the ‘fudge factor’ 2, we see that

$$\frac{1}{x^{3/2} - 1} < \frac{2}{x^{3/2}} \tag{8.22}$$

when $x > 2$. (To see why, observe that when $x > 2$,

$$x^{3/2} - 1 > x^{3/2} - \frac{1}{2}x^{3/2} = \frac{1}{2}x^{3/2}$$

whence (8.22) follows.) Since

$$\int_2^\infty \frac{2}{x^{3/2}} dx$$

converges,

$$\int_2^\infty \frac{1}{x^{3/2} - 1} dx.$$

must also. □

Finding a fudge factor can be a lot of work in itself. Later we give a method for determining the convergence of this integral that does not rely on the use of fudge factors.

Dominant term analysis is not always straightforward, and at times may not even be useful. One must develop an intuition of how functions decay at infinity and of what comparisons to use.

Example 8.11.4. Determine whether or not the following improper integrals converge.

(a) $\int_1^\infty e^{-\sqrt{x}} dx$

(b) $\int_1^\infty \frac{\sin x + 2}{\sqrt{x} + 1} dx.$

Solution. (a) Dominant term analysis does not apply since there is only one term appearing in the integrand. Instead, we recognise that the decay of the exponential $e^{-\sqrt{x}}$ is very rapid; it is certainly much faster than the decay of $\frac{1}{x^2}$. This can be verified by the following limit calculation:

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{e^{-\sqrt{x}}}{\frac{1}{x^2}} &= \lim_{x \rightarrow \infty} \frac{x^2}{e^{x^{1/2}}} \\ &= \lim_{x \rightarrow \infty} \frac{2x}{\frac{1}{2}x^{-1/2}e^{x^{1/2}}} && \text{(by l'Hôpital's rule)} \\ &= \lim_{x \rightarrow \infty} \frac{4x^{3/2}}{e^{x^{1/2}}} \\ &= \lim_{x \rightarrow \infty} \frac{4 \cdot \frac{3}{2}x^{1/2}}{\frac{1}{2}x^{-1/2}e^{x^{1/2}}} && \text{(by l'Hôpital's rule)} \\ &= \lim_{x \rightarrow \infty} \frac{12x}{e^{x^{1/2}}} \\ &\vdots && \text{(two more applications of l'Hôpital's rule)} \\ &= \lim_{x \rightarrow \infty} \frac{24}{e^{x^{1/2}}} \\ &= 0. \end{aligned}$$

Hence we conclude that

$$e^{-\sqrt{x}} < \frac{1}{x^2}$$

whenever x is sufficiently large. Since

$$\int_1^{\infty} \frac{1}{x^2} dx$$

converges,

$$\int_1^{\infty} e^{-\sqrt{x}} dx$$

also converges by the comparison test.

(b) While the numerator of

$$\frac{\sin x + 2}{\sqrt{x} + 1}$$

has no dominant term, intuitively we see that, since the numerator is bounded and the dominant term of the denominator is $\sqrt{x}$, we should compare the integrand with

$$\frac{1}{\sqrt{x}}.$$

Since

$$\int_1^{\infty} \frac{1}{\sqrt{x}} dx$$

diverges, we seek an inequality of the form

$$f(x) \leq \frac{\sin x + 2}{\sqrt{x} + 1}$$

where $f(x)$ is $1/\sqrt{x}$ with an appropriate fudge factor.

Now, $-1 \leq \sin x$, and so

$$\begin{aligned} \frac{\sin x + 2}{\sqrt{x} + 1} &\geq \frac{1}{\sqrt{x} + 1} \\ &\geq \frac{1}{2\sqrt{x}} \end{aligned}$$

for all $x \geq 1$. Since

$$\int_1^{\infty} \frac{1}{2\sqrt{x}} dx$$

diverges (see Proposition 8.10.5),

$$\int_1^{\infty} \frac{\sin x + 2}{\sqrt{x} + 1} dx$$

also diverges by the comparison test. □

The next theorem provides an alternative approach for constructing a comparison of integrals. It is particularly useful if dominant term analysis is straightforward but one wants to avoid the use of fudge factors.

Theorem 8.11.5 (The limit form of the comparison test). *Suppose that f and g are non-negative and bounded on $[a, \infty)$. If*

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$$

and $0 < L < \infty$ then either

$$\int_a^\infty f(x) dx \quad \text{and} \quad \int_a^\infty g(x) dx \quad \text{both converge}$$

or

$$\int_a^\infty f(x) dx \quad \text{and} \quad \int_a^\infty g(x) dx \quad \text{both diverge.}$$

The hypothesis

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$$

indicates that f and g exhibit similar behaviour at infinity. The proof of the theorem will be given at the end of this section.

Example 8.11.6. Determine whether or not the following improper integrals converge.

$$(a) \int_1^\infty \frac{1}{\sqrt{1+x+x^2}} dx$$

$$(b) \int_2^\infty \frac{x^2+3x}{5x^4-2} dx.$$

Solution. (a) Suppose that $f(x) = \frac{1}{\sqrt{1+x+x^2}}$. We need to find a suitable function g to compare f with. As x gets large, the dominant term under the square root is x^2 . Therefore it seems that

$$\frac{1}{\sqrt{1+x+x^2}} \approx \frac{1}{\sqrt{x^2}} = \frac{1}{x}$$

when x is large. So define g by $g(x) = \frac{1}{x}$. The following calculation verifies that our choice of g is suitable:

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{1+x+x^2}} = \lim_{x \rightarrow \infty} \sqrt{\frac{x^2}{1+x+x^2}} = \lim_{x \rightarrow \infty} \sqrt{\frac{1}{\frac{1}{x^2} + \frac{1}{x} + 1}} = 1.$$

Since this limit is a positive real number, we can apply the limit form of the comparison test. By Proposition 8.10.5, $\int_1^\infty g(x) dx$ diverges. Hence $\int_1^\infty f(x) dx$ also diverges. □

Remark 8.11.7. As a general rule, students will find the limit form of the comparison test easier to use than the inequality form of the comparison test when the integrand is a rational function, or is a ratio of roots of polynomials.

Proof of Theorem 8.11.1 (MATH1141 only). Suppose that f and g are non-negative and bounded on $[a, \infty)$. (It follows, for later use, that if $a < M < R$ and R is increasing then the integrals

$$\int_M^R f(x) dx \quad \text{and} \quad \int_M^R g(x) dx,$$

are also increasing.) Suppose also that

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$$

where $0 < L < \infty$. Choose a positive number ϵ such that $0 < \epsilon < L$. By Definition 2.2.1, there is a positive real number M such that

$$\left| \frac{f(x)}{g(x)} - L \right| < \epsilon$$

whenever $x > M$. Hence

$$-\epsilon < \frac{f(x)}{g(x)} - L < \epsilon \quad \forall x > M$$

and rearranging gives

$$0 < (L - \epsilon)g(x) < f(x) < (L + \epsilon)g(x) \quad \forall x > M.$$

By integrating with respect to x we obtain

$$0 < (L - \epsilon) \int_M^R g(x) dx < \int_M^R f(x) dx < (L + \epsilon) \int_M^R g(x) dx$$

whenever $R > M$. By letting R approach infinity it is seen that

- if $\int_M^\infty g(x) dx$ is convergent then $\int_M^R f(x) dx$ is also convergent (since $\int_M^R f(x) dx$ increases as R increases and it is bounded above by $(L + \epsilon) \int_M^\infty g(x) dx$);
- if $\int_M^\infty f(x) dx$ is convergent then $\int_M^R g(x) dx$ is also convergent (since $\int_M^R g(x) dx$ increases as R increases and is bounded above by $\frac{1}{L - \epsilon} \int_M^\infty f(x) dx$);
- if $\int_M^\infty f(x) dx$ is divergent then $\int_M^R g(x) dx \rightarrow \infty$ as $R \rightarrow \infty$; and
- if $\int_M^\infty g(x) dx$ is divergent then $\int_M^R f(x) dx \rightarrow \infty$ as $R \rightarrow \infty$.

Hence either

$$\int_M^\infty f(x) dx \quad \text{and} \quad \int_M^\infty g(x) dx \quad \text{both converge}$$

or

$$\int_M^\infty f(x) dx \quad \text{and} \quad \int_M^\infty g(x) dx \quad \text{both diverge.}$$

The proof of the theorem now easily follows by first observing that

$$\int_a^R f(x) dx = \int_a^M f(x) dx + \int_M^R f(x) dx$$

and

$$\int_a^R g(x) dx = \int_a^M g(x) dx + \int_M^R g(x) dx$$

whenever $a < M < R$, and then letting R approach infinity. □




Figure 8.15: The Si function.

8.12 Functions defined by an integral

Many important functions are defined by an integral. If f is a continuous function then we may define a function F by

$$F(x) = \int_a^x f(t) dt.$$

We note the following points.

- The derivative of F may be calculated by the first fundamental theorem of calculus.
- The value of $F(x)$ may be approximated by computing Riemann sums.
- Symmetries of f can help in understanding the behaviour of F (see, for example, Proposition 8.8.6).
- The behaviour of $F(x)$ as $x \rightarrow \infty$ or as $x \rightarrow -\infty$ may be ascertained by studying an improper integral of f .

We give one example. Others may be found in the problem set for Chapter 8 and at the beginning of Chapter 9.

Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(t) = \begin{cases} \frac{\sin t}{t} & \text{if } t \neq 0 \\ 1 & \text{if } t = 0. \end{cases}$$

Since $\lim_{t \rightarrow 0} \frac{\sin t}{t} = 1$, the function f is continuous everywhere. We can now define the function $\text{Si} : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\text{Si}(x) = \int_0^x f(t) dt \quad \forall x \in \mathbb{R}.$$

The name Si comes from ‘sine integral.’ This function is used in electrical engineering for signal processing and in surveying for the Global Positioning System (GPS). Its graph is plotted in Figure 8.15.

Example 8.12.1. Consider the functions f and Si defined above.

- (a) Find the value of $\text{Si}(0)$.
- (b) Show that Si is an odd function.
- (c) Find and classify the critical points of Si on $(0, 3\pi)$.
- (d) A function $G : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula

$$G(x) = \int_0^{x^2} f(t) dt.$$

Calculate $G'(x)$.

Solution. (a) $\text{Si}(0) = \int_0^0 f(t) dt = 0$.

(b) We need to show that $\text{Si}(-x) = -\text{Si}(x)$. Now

$$\begin{aligned} \text{Si}(-x) &= \int_0^{-x} \frac{\sin t}{t} dt \\ &= \int_0^x \frac{\sin(-u)}{-u} (-du) \quad (\text{using the substitution } t = -u) \\ &= - \int_0^x \frac{-\sin u}{-u} du \\ &= - \int_0^x \frac{\sin u}{u} du \\ &= -\text{Si}(x). \end{aligned}$$

Hence Si is an odd function.

(c) By the first fundamental theorem of calculus,

$$\text{Si}'(x) = \frac{\sin x}{x}$$

for all x in $(0, 3\pi)$. Critical points occur when $\text{Si}'(x) = 0$, that is, when $x = \pi$ or when $x = 2\pi$. To classify the critical points, we calculate the second derivative:

$$\text{Si}''(x) = \frac{x \cos x - \sin x}{x^2}.$$

Now $\text{Si}''(\pi) = -\frac{1}{\pi}$ and $\text{Si}''(2\pi) = \frac{1}{2\pi}$. Therefore Si has a local maximum point when $x = \pi$ and a local minimum point when $x = 2\pi$.

(d) Note that $G(x) = \text{Si}(x^2)$, so to calculate $G'(x)$ we use the chain rule:

$$G'(x) = \frac{d}{dx}(\text{Si}(x^2)) = 2x \text{Si}'(x^2) = 2x \times \frac{\sin(x^2)}{x^2} = \frac{2 \sin(x^2)}{x}.$$

(If this calculation is too dense, then consider

$$y = G(x), \quad u = x^2$$

so that

$$y = \int_0^u f(t) dt \quad \text{and} \quad \frac{dy}{du} = f(u)$$

by the first fundamental theorem of calculus. Then by the chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = f(u) \times 2x = 2xf(x^2) = \frac{2x \sin(x^2)}{x^2} = \frac{2 \sin(x^2)}{x}.$$

Hence $G'(x) = \frac{2 \sin(x^2)}{x}$.)

□

8.13 Maple notes

The command `int(f(x),x);` will cause Maple to attempt to find the indefinite integral $\int f(x) dx$, and the command `int(f(x),x=a..b);` is used to find definite integrals $\int_a^b f(x) dx$. If Maple is unable to find a primitive, then numerical integration can be carried out by combining the `int` and `evalf` commands.

For example,

```
> int(1/(x*log(x)^2),x);
                                      $-(\ln(x))^{-1}$ 
> int(1/(x*log(x)^2),x=2..infinity);
                                      $(\ln(2))^{-1}$ 
> int( exp(-x^2)*ln(x), x=0..1 );
                                      $\int_0^1 e^{-x^2} \ln(x) dx$ 
> evalf(%);
                                     -0.9059404763
```

Problems for Chapter 8

Problems 8.1 : Area and the Riemann Integral and 8.2 : Integration using Riemann sums

1. [R] [V]

a) By taking the partition $\mathcal{P}_n = \left\{0, \frac{1}{n}, \frac{2}{n}, \dots, 1\right\}$ of the interval $[0,1]$, calculate the lower sum $\underline{S}_{\mathcal{P}_n}(f)$ and the upper sum $\overline{S}_{\mathcal{P}_n}(f)$ for each function f .

i) $f(x) = 1$

ii) $f(x) = x$

iii) $f(x) = x^2$

[You may need $\sum_{k=1}^n k^2 = \frac{1}{6}n(n+1)(2n+1)$.]

iv) $f(x) = x^3$

[You may need $\sum_{k=1}^n k^3 = \frac{1}{4}n^2(n+1)^2$.]

v) $f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$

b) By taking the limit as $n \rightarrow \infty$ for each sum $\underline{S}_{\mathcal{P}_n}(f)$ and $\overline{S}_{\mathcal{P}_n}(f)$ calculated in (a), either calculate $\int_0^1 f(x) dx$, or show that f is not Riemann integrable.

2. [R] An electrical signal $S(t)$ has its amplitude $|S(t)|$ tested (sampled) every $\frac{1}{10}$ of a second. It is desired to estimate the energy over a period of half a second, given exactly by

$$\left(\int_0^{\frac{1}{2}} |S(t)|^2 dt \right)^{\frac{1}{2}}.$$

The results of the measurement are shown in the following table:

t	.1	.2	.3	.4	.5
$ S(t) $	60	50	50	45	55
$e(t)$	5	3	7	4	10

- a) Using the above data for $S(t)$, set up an appropriate Riemann sum and compute an approximate value for the energy.
- b) It is known that the signal varies by an amount of at most $\pm e(t)$, as shown above, in each $\frac{1}{10}$ second period. Calculate upper and lower bounds for the energy.

3. [X] Consider the partition $\mathcal{P}_n$ of $[1, 2]$, given by $\mathcal{P}_n = \{q^0, q^1, q^2, \dots, q^n\}$ where $q^n = 2$. (Notice that (i) the divisions are not of equal width and (ii) $1 < q < 2$ and $q \rightarrow 1$ as $n \rightarrow \infty$.) If $f(x) = x^j$ for some positive integer j , then evaluate the integral

$$\int_1^2 f(x) dx$$

by calculating the limit $\lim_{n \rightarrow \infty} \underline{S}_{\mathcal{P}_n}(f)$ of the corresponding lower Riemann sums.

Problems 8.3 : The Riemann integral and signed area and 8.4 : Basic properties of the Riemann integral

4. [R] Find the area of the region bounded by the line $y = x$ and the parabola $y = x^2 - 2$.
5. [R] Find

a) $\int_4^9 \frac{x^3 - x}{x^{3/2}} dx$ b) $\int_{-4}^2 |x| dx$.

6. [H] Find a function f which satisfies the integral equation

$$\int_0^x t f(t) dt = \int_x^0 (t^2 + 1) f(t) dt + x.$$

7. [R] Explain why $\int_{-1}^1 \frac{1}{x^2} dx = \left[-\frac{1}{x} \right]_{-1}^1 = -1 - 1 = -2$ is not valid.

8. [H]

- a) Suppose that f is a continuous increasing (and hence invertible) function on $[a, b]$. If $c = f(a)$, $d = f(b)$ and $a, b, c, d \geq 0$, then explain why

$$\int_c^d f^{-1}(t) dt = bd - ac - \int_a^b f(x) dx.$$

- b) Use this to find $\int_{1/2}^1 \sin^{-1} x dx$.

9. [H] Suppose that $U'(x) = u(x)$.

- a) Find $V'(x)$ if $V(x) = (a - x)U(x) + \int_0^x U(t) dt$ where a is a constant.

- b) Hence show that $\int_0^a U(x) dx = aU(0) + \int_0^a (a - x)u(x) dx$.

Problems 8.5 : The first fundamental theorem of calculus

10. [H] Suppose that $f(t) = \lfloor t \rfloor$ and $F(x) = \int_0^x f(t) dt$, where $\lfloor t \rfloor$ is the greatest integer less than or equal to t . Use a graph of f to sketch F on the interval $[-1, 3]$. Is F continuous? Where is F differentiable?
11. [H] Suppose that $f(t) = \sin(t^2)$. Sketch the graph of f on the interval $[0, 3]$. Use this to sketch the graph of F on the interval $[0, 3]$, where $F(x) = \int_0^x f(t) dt$. Indicate where F has local maxima and minima.
12. [R] [V] Find $F'(x)$ for each function $F : \mathbb{R} \rightarrow \mathbb{R}$ given below.
- a) $F(x) = \int_0^x \sin(t^2) dt$ b) $F(x) = \int_0^{x^3} \sin(t^2) dt$
- c) $F(x) = \int_{x^3}^1 \sin(t^2) dt$ d) $F(x) = \int_x^{x^3} \sin(t^2) dt$
13. [R] Find $\frac{d}{dx} \int_x^4 (5 - 4t)^5 dt$.

Problems 8.6 : The second fundamental theorem of calculus

14. [R]
- a) Suppose that $f(x) = \frac{1}{x}$. By considering the lower Riemann sum for f with respect to the partition

$$\left\{ \frac{n}{n}, \frac{n+1}{n}, \frac{n+2}{n}, \dots, \frac{2n}{n} \right\}$$

of $[1, 2]$, show that

$$\ln 2 = \lim_{n \rightarrow \infty} \left(\frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} \right).$$

- b) Suppose that $f(x) = \frac{1}{\sqrt{1-x^2}}$.
- i) Show that f is increasing on the interval $[0, \frac{1}{2}]$.
- ii) Find the upper Riemann sum for f with respect to the partition

$$\left\{ \frac{0}{2n}, \frac{1}{2n}, \frac{2}{2n}, \frac{3}{2n}, \dots, \frac{n}{2n} \right\}$$

of $[0, \frac{1}{2}]$.

- iii) Hence evaluate

$$\lim_{n \rightarrow \infty} \left(\frac{1}{\sqrt{4n^2 - 1^2}} + \frac{1}{\sqrt{4n^2 - 2^2}} + \frac{1}{\sqrt{4n^2 - 3^2}} + \dots + \frac{1}{\sqrt{4n^2 - n^2}} \right).$$

Problems 8.7 : Indefinite integrals

15. [R] Evaluate the following integrals by *inspection*.

- a) $\int x e^{x^2} dx$ b) $\int \frac{\sin \sqrt{x}}{\sqrt{x}} dx$
 c) $\int_0^1 2x(1+x^2)^3 dx$ d) $\int_{-a}^a x^2 \sqrt{a^3 - x^3} dx \quad (a > 0)$
 e) $\int_0^{\pi/2} \cos^3 x \sin x dx$ f) [H] $\int_{-1}^0 \sqrt{t^2 + t^4} dt$

Problems 8.8 : Integration by substitution

16. [R] Use a substitution to evaluate the following integrals.

- a) $\int \frac{dx}{1 + \sqrt{x}}$ b) $\int x(5x - 1)^{19} dx$
 c) [V] $\int \frac{1 - x}{(1 + x)^3} dx$ d) [V] $\int_0^4 \frac{dx}{5 + \sqrt{x}}$

17. [X] Use the substitution $u = t - t^{-1}$ to find $\int \frac{1 + t^2}{1 + t^4} dt$.

Problems 8.9 : Integration by parts

18. [R] Use integration by parts to evaluate the following integrals.

- a) $\int_0^1 x e^{5x} dx$ b) $\int x^2 \cos x dx$ c) $\int \ln x dx$
 d) $\int_0^{0.5} \sin^{-1} x dx$ e) $\int_1^e x^7 \ln x dx$ f) $\int_0^\pi x^2 \cos 2x dx$
 g) [V] $\int e^x \cos x dx$ h) $\int \tan^{-1} x dx$ i) [H] $\int_0^{\pi/4} \sec^3 \theta d\theta$

Problems 8.10 : Improper integrals

19. [R] [V] Evaluate the following improper integrals or show that they diverge.

- a) $\int_0^\infty e^{-5x} dx$ b) $\int_{-\infty}^1 e^{-0.01x} dx$ c) $\int_0^\infty \frac{dx}{4 + x^2}$
 d) $\int_{-\infty}^\infty x^3 e^{-x^4} dx$ e) $\int_2^\infty \frac{dx}{(x - 1)^{3/2}}$ f) $\int_e^\infty \frac{dx}{x \ln x}$

20. [H] Prove that $\int_0^\infty x^n e^{-x} dx = n!$ whenever $n = 0, 1, 2, \dots$.

21. [H]

- a) Find $\lim_{R \rightarrow \infty} \int_{-R}^R \frac{x}{1+x^2} dx$.
- b) Find $\lim_{R \rightarrow \infty} \int_{-R}^{2R} \frac{x}{1+x^2} dx$.
- c) Does $\int_{-\infty}^{\infty} \frac{x}{1+x^2} dx$ converge? Explain.

Problems 8.11 : Comparison tests for improper integrals

22. [R] Use the inequality form of the comparison test to determine whether or not the following improper integrals converge.

a) [V] $\int_1^{\infty} \frac{1}{\sqrt{1+x^4}} dx$ b) $\int_2^{\infty} \frac{1}{\sqrt[3]{x^2-x}} dx$ c) $\int_2^{\infty} \frac{1}{\ln x} dx$

23. [R] Use the limit form of the comparison test to determine whether or not the following improper integrals converge.

a) $\int_2^{\infty} \frac{x}{2x^3-1} dx$ b) $\int_1^{\infty} \frac{2x-1}{x^2+2} dx$ c) $\int_2^{\infty} \frac{1}{\sqrt{x^6-1}} dx$

24. [R] Use a comparison test to determine whether or not the following improper integrals converge.

a) $\int_1^{\infty} \frac{3x + \sin x + 2}{2x^3 - x + 8} dx$ b) $\int_4^{\infty} \frac{4x^3 - x + 5}{x^4 - x^2 + 1} dx$ c) [H] $\int_2^{\infty} \frac{\ln t}{t^{3/2}} dt$

25. [H] Find all real numbers s such that the improper integral

$$\int_1^{\infty} \frac{x^s}{1+x} dx$$

is convergent.

26. [H] Find all real numbers p such that $\int_2^{\infty} \frac{1}{x(\ln x)^p} dx$ converges.

27. [H] For which pairs of numbers (a, b) does the improper integral $\int_1^{\infty} \frac{x^b}{(1+x^2)^a} dx$ converge?

Problems 8.12 : Functions defined by an integral

28. [R] [V] Given a positive real number x , let $\pi(x)$ denote the number of primes less than or equal to x . The function Li with domain $(1, \infty)$ is given by

$$\text{Li}(x) = \int_2^x \frac{1}{\ln t} dt$$

and is known as the ‘logarithmic integral function’. It has the property that

$$\frac{\text{Li}(x)}{\pi(x)} \approx 1$$

when x is sufficiently large.

- Evaluate $\pi(10)$, $\pi(20)$ and $\pi(3.14159)$.
- Suppose that $x > 0$. What does $\frac{\pi(x)}{x}$ represent?
- Find $\frac{d}{dx} \text{Li}(x)$ and $\text{Li}(2)$.
- By applying the mean value theorem to Li on the interval $[2, 10^6]$, find a lower bound for $\text{Li}(10^6)$.
- If x is large then

$$\frac{\pi(x)}{x} \approx \frac{\pi(x)}{x} \frac{\text{Li}(x)}{\pi(x)} = \frac{\text{Li}(x)}{x}.$$

Using this approximation and your answer to part (d), find an approximate lower bound for $\frac{\pi(10^6)}{10^6}$.

Note: There are 78,498 primes less than one million so the actual value of $\frac{\pi(10^6)}{10^6}$ is 0.078498.

29. **[R]** The function $\text{erf} : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt.$$

The function erf is an error function and can be used to calculate the probability that a measurement has an error in a given range of values.

- Calculate $\text{erf}'(x)$.
- Explain why erf is an increasing function on $\mathbb{R}$.
- [H]** Show that erf is an odd function.
- By calculating Riemann sums with respect to the partition $\{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$, find upper and lower bounds for $\text{erf}(1)$.
 - Explain why $e^{-t^2} < e^{-t}$ whenever $t > 1$.
 - Hence show that $\int_1^\infty e^{-t^2} dt$ converges and find an upper bound for this improper integral.
 - Using your answers to (i) and (iii), find an upper bound for $\lim_{x \rightarrow \infty} \text{erf}(x)$. (In fact, $\lim_{x \rightarrow \infty} \text{erf}(x) = 1$ but this is not so easy to prove.)
- Sketch the graph of erf .
- Explain why erf has an inverse function erf^{-1} and sketch its graph.

Chapter 9

The logarithmic and exponential functions

The volume of a colony of bacteria on an agar plate is 1 cubic millimetre and doubles every day. Let $V(t)$ denote the volume of bacteria after t days. It is clear that

$$V(0) = 1, \quad V(1) = 2, \quad V(2) = 4, \quad V(3) = 8, \quad V(4) = 16,$$

and so on. In general

$$V(t) = 2^t$$

if t is a nonnegative integer. The corresponding points are plotted in Figure 9.1 (a).

If one wanted to determine the volume of the colony part-way through a day, it would be natural to use the formula

$$V(t) = 2^t,$$

where t is a nonnegative real number. This formula makes sense when t is a rational number. If $t = p/q$ where p and q are integers and $q > 0$, then 2^t is defined to be the unique positive q th root of 2^p . However, if t is an irrational number, this definition does not apply. So what do we mean by $2^{\sqrt{3}}$ or 2^π ? We have not yet seen a definition for such numbers.

That this shortcoming in our current definition of 2^t is a serious problem can be illustrated graphically. The graph corresponding to $V(t) = 2^t$, where t is confined to the nonnegative rational numbers, has an infinite number of ‘gaps’ (see Figure 9.1 (b)). Clearly this is not a satisfactory state of affairs. One of the aims of this chapter is to rectify this deficiency and show that, when t is an irrational number, 2^t can be defined in such a way that the graph of $V : [0, \infty) \rightarrow \mathbb{R}$ is continuous (see in Figure 9.1 (c)).

Our plan of attack is the following.

1. In Section 9.2, we define the function $\ln$ as an integral (by the fundamental theorem of calculus).
2. In Section 9.3, we define the function $\exp$ as the inverse of $\ln$ (by the inverse function theorem).
3. In Section 9.4, we define the number b^t , where $b > 0$ and t is an irrational number, by a formula involving both $\ln$ and $\exp$.




Figure 9.1: Growth in volume of bacteria on an agar plate.

We shall see that when the definition of b^t (for rational t) is combined with the definition of b^t (for irrational t), the resulting function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(t) = b^t \quad \forall t \in \mathbb{R},$$

is continuous.

9.1 Powers and logarithms

The content of this section is revision, and is placed here primarily as reference. It summarises a few known definitions and facts about powers and logarithms.

Definition 9.1.1. Suppose that b is a positive real number. If p and q are integers and $q > 0$ then $b^{p/q}$ is defined to be the unique positive q th root of b^p .

The following is an elementary definition of the logarithm to the base b .

Definition 9.1.2. Suppose that b is a positive real number not equal to one, c is a rational number and $a = b^c$. Then $\log_b a$ is defined by the formula

$$c = \log_b a.$$

Note that, according to Definition 9.1.2, one cannot write down $\log_b a$ for *any* positive number a (even if a is rational); $\log_b a$ is only defined if $a = b^c$ for some rational number c . (This shortcoming in the domain of the function $\log_b$ will be rectified later in the chapter when Definition 9.1.2 is superseded by a superior definition.)

The elementary properties of logarithms are summarised below: if $x > 0$, $y > 0$ and $b > 0$ with $b \neq 1$ then

$$\begin{aligned}\log_b 1 &= 0, & \log_b b &= 1, \\ \log_b(xy) &= \log_b x + \log_b y, & \log_b\left(\frac{x}{y}\right) &= \log_b x - \log_b y, \\ \log_b(b^r) &= r, & \log_b(x^r) &= r \log_b x.\end{aligned}\tag{9.1}$$

Remark 9.1.3. In the properties listed above, it is implicitly assumed that r is a rational number, and that x and y are of the form b^c for some rational number c . One of the aims of this chapter is to remove these awkward assumptions.

9.2 The natural logarithm function

(Ref: SH10 §7.2)


The $\ln$ function will already be familiar to you from high school. However, it was probably never defined rigorously. In order to address the problems identified in the introduction to Chapter 9, it will be necessary to start from scratch. Fortunately, by the fundamental theorem of calculus, this is not difficult.

Definition 9.2.1. The function $\ln : (0, \infty) \rightarrow \mathbb{R}$ is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

We read $\ln$ as either ‘ell en’ or as ‘log’.

Since the function $\ln$ is defined by an integral, $\ln x$ is simply the area of the shaded region shown below in the case when $x > 1$.



One can also immediately see, by the first fundamental theorem of calculus, that

$$\frac{d}{dx}(\ln x) = \frac{1}{x}.$$

This and other properties of $\ln$ are stated in the proposition below.

Proposition 9.2.2. *The function $\ln$ has the following properties:*

- (i) $\ln$ is differentiable on $(0, \infty)$ and $\frac{d}{dx}(\ln x) = \frac{1}{x}$;
- (ii) $\ln x > 0$ if $x > 1$, $\ln 1 = 0$ and $\ln x < 0$ if $0 < x < 1$;
- (iii) $\ln x \rightarrow -\infty$ as $x \rightarrow 0^+$ and $\ln x \rightarrow \infty$ as $x \rightarrow \infty$;
- (iv) $\ln(xy) = \ln(x) + \ln(y)$ for all positive real numbers x and y ;
- (v) $\ln\left(\frac{x}{y}\right) = \ln(x) - \ln(y)$ for all positive real numbers x and y ; and
- (vi) $\ln(x^r) = r \ln(x)$ whenever r is a rational number and x is a positive real number.


Proof. (i) Apply the first fundamental theorem of calculus (Theorem 8.5.1) to the definition of $\ln$.

(ii) This follows from Definition 8.4.2 and the fact that $\frac{1}{t} > 0$ when $t > 0$.

(iii) We need to show that the improper integral $\int_1^\infty \frac{1}{t} dt$ diverges to infinity. The diagram below shows that

$$\int_1^2 \frac{dt}{t} \geq 1 \times \frac{1}{2}, \quad \int_2^4 \frac{dt}{t} \geq 2 \times \frac{1}{4}, \quad \int_4^8 \frac{dt}{t} \geq 4 \times \frac{1}{8}$$

and so on.



Hence

$$\begin{aligned} \int_1^{2^n} \frac{dt}{t} &\geq \underbrace{\frac{1}{2} + \frac{1}{2} + \cdots + \frac{1}{2}}_{n \text{ terms}} \\ &= \frac{n}{2} \\ &\rightarrow \infty \end{aligned}$$

as $n \rightarrow \infty$. Hence the integral diverges and therefore $\ln x \rightarrow \infty$ as $x \rightarrow \infty$.

This argument can be adapted to show that

$$\int_{2^{-n}}^1 \frac{dt}{t} \rightarrow -\infty$$

as $n \rightarrow \infty$. Hence $\ln x \rightarrow -\infty$ as $x \rightarrow 0^+$.

(iv) Suppose that y is some fixed positive real number and that $x \in (0, \infty)$. Then, by the chain rule and part (i),

$$\frac{d}{dx}(\ln(xy)) = y \ln'(xy) = y \times \frac{1}{xy} = \frac{1}{x} = \frac{d}{dx}(\ln(x)).$$

Hence

$$\ln(xy) = \ln(x) + C$$

for some constant C . When $x = 1$ we obtain

$$\ln(y) = \ln(1) + C = 0 + C = C$$

by part (ii). Hence

$$\ln(xy) = \ln(x) + C = \ln(x) + \ln(y)$$

as required.

(v) The proof uses the same technique as the proof of (vi) and is left to the reader.

(vi) This proof also uses the same technique as the proof of (iv). □

The fact that $\ln(1) = 0$ and that $\ln$ satisfies properties (iv), (v) and (vi) suggests that $\ln$ coincides with one of the logarithm functions $\log_b$ for some base b (with the advantage that $\text{Dom}(\ln) = (0, \infty)$). We now aim to identify the base.

The function $\ln$ is increasing and continuous with $\text{Range}(\ln) = \mathbb{R}$ (see Proposition 9.2.2 (i) and (ii)). Hence, by the intermediate value theorem, there is exactly one real number x that satisfies the equation

$$\ln(x) = 1.$$

We call this solution e (after Leonard Euler, who was probably the greatest mathematician of the eighteenth century).

Definition 9.2.3. The real number e is defined to be the unique number x that satisfies the equation


$$\int_1^x \frac{1}{t} dt = 1.$$

Thus $\ln(e) = 1$. One can show that e is irrational and that $e \approx 2.71828$. From Proposition 9.2.2, we see now that

$$\ln(e^r) = r$$

for every rational number r . This equation shows that $\ln$ coincides with the logarithm to the base e . Since the number e arises naturally in many contexts, $\ln$ is usually called the *natural logarithm*.

By using the properties of $\ln$ given by Proposition 9.2.2, one can draw the graph of $\ln$.



We note that, although $\ln(x) \rightarrow \infty$ as $x \rightarrow \infty$, we have $\frac{d}{dx} \ln(x) = \frac{1}{x} \rightarrow 0$ as $x \rightarrow \infty$. This means the graph of $\ln$ becomes flatter towards infinity and hence approaches infinity very slowly.


9.3 The exponential function

(Ref: SH10 §7.4)

In Section 9.2 it was shown that $\frac{d}{dx}(\ln x) > 0$ for all $x > 0$. Therefore, by the inverse function theorem (Theorem 6.4.1), $\ln$ has an inverse function.

Definition 9.3.1. The function $\exp : \mathbb{R} \rightarrow (0, \infty)$ is defined to be the inverse function of $\ln : (0, \infty) \rightarrow \mathbb{R}$.

By reflecting the graph of $\ln$ in the line $y = x$, one obtains the graph of $\exp$.



Proposition 9.3.2. The function $\exp : \mathbb{R} \rightarrow (0, \infty)$ has the following properties:

- (i) $\exp(\ln x) = x$ for all x in $(0, \infty)$ and $\ln(\exp x) = x$ for all x in $\mathbb{R}$;

- (ii) $\exp(1) = e$ and $\exp(0) = 1$;
- (iii) $\exp x \rightarrow \infty$ as $x \rightarrow \infty$ and $\exp x \rightarrow 0$ as $x \rightarrow -\infty$.
- (iv) $\exp$ is differentiable on $\mathbb{R}$ and $\frac{d}{dx}(\exp x) = \exp x$ for all x in $\mathbb{R}$;
- (v) $\exp(x + y) = \exp(x) \exp(y)$ for all x and y in $\mathbb{R}$; and
- (vi) $\exp(rx) = (\exp x)^r$ for every real number x and every rational number r .

Proof. The proof of (i) follows immediately from Definition 9.3.1 and the proof of (ii) and (iii) then follow from (i) and the continuity of $\ln$ and $\exp$.

That $\exp$ is differentiable on $\mathbb{R}$ follows from the inverse function theorem. To complete the proof of (iii), we differentiate both sides of the equation

$$\ln(\exp x) = x$$

with respect to x to obtain

$$\frac{1}{\exp x} \times \frac{d}{dx}(\exp x) = 1$$

by the chain rule. Rearranging gives $\frac{d}{dx}(\exp x) = \exp x$ as required.

If x and y are real numbers then

$$\begin{aligned} \exp(x + y) &= \exp\left(\ln(\exp x) + \ln(\exp y)\right) && \text{(by part (i))} \\ &= \exp\left(\ln(\exp x \times \exp y)\right) && \text{(by Proposition 9.2.2 (iv))} \\ &= \exp x \times \exp y && \text{(by part (i)).} \end{aligned}$$

This completes the proof of (iv)

To prove (v), suppose that r is a rational number and x is a real number. Then

$$\begin{aligned} \exp(rx) &= \exp\left(r \ln(\exp x)\right) && \text{(by part (i))} \\ &= \exp\left(\ln((\exp x)^r)\right) && \text{(by Proposition 9.2.2 (v))} \\ &= (\exp x)^r && \text{(by part (i)).} \end{aligned}$$

This completes the proof of the proposition. □

Note that by Proposition 9.3.2 (ii) and (v) we have

$$\exp(r) = (\exp 1)^r = e^r \tag{9.2}$$

for every rational number r . This inspires the following definition.

Definition 9.3.3. Suppose that x is an irrational number. Then we define the real number e^x by the formula

$$e^x = \exp x.$$

Thus Definition 9.3.3 extends Definition 9.1.1, in the case when $b = e$, to include irrational powers. Moreover, since $\exp x = e^x$ and $\exp$ is a continuous function, the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = e^x$, is also continuous. We have thus made significant progress towards the problem outlined at the beginning of the chapter. Finally, since $\exp x = e^x$, we call $\exp$ the *exponential function*.

9.4 Exponentials and logarithms with other bases

(Ref: SH10 §7.5)

At the end of the last section, we gave a definition of e^x , where x is an irrational number. Moreover, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = e^x$, then f is a continuous and differentiable function. In this section, we do the same for b^x , where b is any positive real number.

To begin, consider a rational number r and any positive real number b . Then

$$b^r = \exp(\ln(b^r)) = \exp(r \ln b) = e^{r \ln b} \quad (9.3)$$

by the definitions and properties established in the last two sections. This inspires the following definition.

Definition 9.4.1. Suppose that b is a positive real number and x is an irrational number. We define the number b^x by the formula

$$b^x = e^{x \ln b}.$$

By combining equation (9.3) and Definition 9.4.1 we see that

$$b^x = e^{x \ln b} \quad \forall x \in \mathbb{R}. \quad (9.4)$$

Since $\exp$ and $\ln$ are continuous functions, the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = b^x \quad \forall x \in \mathbb{R},$$

is also continuous (see Proposition 3.1.2). This solves the problem articulated at the beginning of the chapter. Thus

$$2^{\sqrt{3}} = e^{\sqrt{3} \ln 2} \simeq 1.492106 \dots$$


Moreover, we can discard our old definition of $\log_b$ (see Definition 9.1.2), together with its technical difficulties (see Remark 9.1.3), and replace it with a simpler and more powerful definition.

Definition 9.4.2. Suppose that b is a positive real number, $b \neq 1$. Then the function $\log_b : (0, \infty) \rightarrow \mathbb{R}$ is defined to be the inverse of the function $f : \mathbb{R} \rightarrow (0, \infty)$, where

$$f(x) = b^x$$

for all x in $\mathbb{R}$.

We leave it as an exercise to show that the function f of Definition 9.4.2 is one-to-one and hence invertible. The graphs of various exponential functions and their corresponding logarithmic functions are shown below.



The following proposition gives a formula for $\log_b$ in terms of $\ln$. Its proof is left as an exercise.

Proposition 9.4.3. *Suppose that b is a positive real number, $b \neq 1$. Then*

$$\log_b x = \frac{\ln x}{\ln b} \quad \forall x \in (0, \infty). \quad (9.5)$$

Using equations (9.4) and (9.5), one can easily establish familiar properties for the exponentials and logarithms to the base b , with the advantage that we no longer need the technical restrictions of Remark 9.1.3.

9.5 Integration and the $\ln$ function

We have already seen that

$$\frac{d}{dx}(\ln x) = \frac{1}{x}$$

provided that $x > 0$. It follows from the chain rule that

$$\frac{d}{dx}(\ln(f(x))) = \frac{1}{f(x)} \times f'(x) = \frac{f'(x)}{f(x)},$$

provided that f is differentiable and $f(x) > 0$. Hence

$$\int \frac{f'(x)}{f(x)} dx = \ln(f(x)) + C.$$

It is not difficult to remove the restriction that $f(x) > 0$. To take a simple example, suppose that $x \neq 0$. Then

$$\begin{aligned} \frac{d}{dx}(\ln|x|) &= \begin{cases} \frac{d}{dx}(\ln x) & \text{if } x > 0 \\ \frac{d}{dx}(\ln(-x)) & \text{if } x < 0 \end{cases} \\ &= \begin{cases} \frac{1}{x} & \text{if } x > 0 \\ \frac{1}{-x} \times (-1) & \text{if } x < 0 \end{cases} \\ &= \frac{1}{x}. \end{aligned}$$

Thus

$$\int \frac{1}{x} dx = \ln |x| + C$$

provided x is restricted to an interval not containing zero. By generalising this calculation we obtain the formula

$$\boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + C,}$$

provided that f is never zero over the interval of integration.

Example 9.5.1. Find

(a) $\int \tan x dx$

(b) $\int \sec x dx$.

Solution. (a) By using the fact that $\tan x = \frac{\sin x}{\cos x}$ we have

$$\begin{aligned} \int \tan x dx &= - \int \frac{-\sin x}{\cos x} dx \\ &= - \ln |\cos x| + C. \end{aligned}$$

(The indefinite integral of $\cot$ can be found similarly.)

(b) This example involves a special trick:

$$\begin{aligned} \int \sec x dx &= \int \sec x \frac{\tan x + \sec x}{\tan x + \sec x} dx \\ &= \int \frac{\tan x \sec x + \sec^2 x}{\sec x + \tan x} dx \\ &= \ln |\sec x + \tan x| + C. \end{aligned}$$

(A similar trick can be used to find the indefinite integral of cosec .) □

9.6 Logarithmic differentiation

(Ref: SH10 §7.3)

Logarithms are powerful because they transform powers into products, products into sums and quotients into differences. In this section, we illustrate the use of the logarithm for differentiating functions that are defined by (combinations of) powers, products or quotients.

Example 9.6.1. Find $\frac{dy}{dx}$ in each case:

(a) $y = 10^x$

(b) $y = \left(\frac{(3x^2 + 4)(x + 2)}{x^3 + 5x} \right)^{3/5}$

(c) $y = (\sin x)^{\cos x}$.

Solution. (a) If we take $\ln$ of both sides of the equation $y = 10^x$ then

$$\begin{aligned}\ln y &= \ln 10^x \\ &= x \ln 10.\end{aligned}$$

By differentiating with respect to x , we obtain

$$\frac{1}{y} \frac{dy}{dx} = \ln 10$$

and hence

$$\begin{aligned}\frac{dy}{dx} &= (\ln 10) \times y \\ &= (\ln 10)10^x.\end{aligned}$$

Alternatively: $\frac{d}{dx}(10^x) = \frac{d}{dx}(e^{x \ln 10}) = \ln 10 e^{x \ln 10} = (\ln 10)10^x$. $\square$

Remark 9.6.2. Recall that one can only take the logarithm of a positive number. Hence our solution to Example 9.6.1 (b) and (c) is only valid when $y > 0$.

9.7 Indeterminate forms with powers

(Ref: SH10 §11.6)

Consider the limits

$$\lim_{x \rightarrow 0^+} x^x \quad \text{and} \quad \lim_{x \rightarrow \infty} x^{1/x}. \quad (9.6)$$

The first limit is of the form 0^0 while the second is of the form ∞^0 . Both of these forms are examples of indeterminate forms and hence we cannot say what the corresponding limits are in each case without further calculation. Since each limit involves a power, it is natural to first take the logarithm of the limit and then bring l'Hôpital's rule into play.

Example 9.7.1. Evaluate the limit

$$\lim_{x \rightarrow 0^+} x^{2x}.$$

Solution. The limit is an indeterminate form of the type 0^0 . By taking the natural logarithm, we can transform the limit into an indeterminate form of the type $\frac{\infty}{\infty}$:

$$\begin{aligned}\lim_{x \rightarrow 0^+} x^{2x} &= \lim_{x \rightarrow 0^+} \exp(\ln x^{2x}) && \text{(since } \ln \text{ and } \exp \text{ are inverses)} \\ &= \lim_{x \rightarrow 0^+} \exp(2x \ln x) \\ &= \exp\left(\lim_{x \rightarrow 0^+} 2x \ln x\right) && \text{(since } \exp \text{ is continuous)} \\ &= \exp\left(\lim_{x \rightarrow 0^+} \frac{\ln x}{1/(2x)}\right).\end{aligned}$$

We can now apply l'Hôpital's rule to the problem. By differentiating the numerator and denominator and then simplifying we obtain

$$\begin{aligned}\lim_{x \rightarrow 0^+} x^{2x} &= \exp \left(\lim_{x \rightarrow 0^+} \frac{1/x}{-1/(2x^2)} \right) \\ &= \exp \left(\lim_{x \rightarrow 0^+} -2x \right) \\ &= \exp(0) \\ &= 1.\end{aligned}$$

□

Each of the limits in (9.6) can be evaluated by using the ideas in this example.

Remark 9.7.2. The limit of Example 9.7.1 is of the form 0^0 and has the value 1. The limit

$$\lim_{x \rightarrow 0^+} \left(e^{-\frac{1}{x}} \right)^x$$

is also of the form 0^0 . However, its value is not 1:

$$\lim_{x \rightarrow 0^+} \left(e^{-\frac{1}{x}} \right)^x = \lim_{x \rightarrow 0^+} e^{-1} = \frac{1}{e}.$$

The fact that limits of the type 0^0 can have different values explains why 0^0 is an indeterminate form.

Problems for Chapter 9

Problems 9.1 : Powers and logarithms and
9.2 : The natural logarithm function

1. [R]
 - a) Write down the definition of $\ln x$, where $x > 0$.
 - b) Explain why $\frac{d}{dx} \ln x = \frac{1}{x}$ whenever $x > 0$.
 - c) Suppose that r is a rational number and that x and y are positive real numbers.
 - i) By first differentiating $\ln(xy)$ with respect to x , show that $\ln(xy) = \ln x + \ln y$.
 - ii) Use the same technique to show that

$$\ln\left(\frac{x}{y}\right) = \ln x - \ln y \quad \text{and} \quad \ln(x^r) = r \ln x.$$

2. [R]
 - a) Prove, using upper and lower Riemann sums and the definition of $\ln x$, that $\ln 2 < 1 < \ln 4$, and hence that $2 < e < 4$.
 - b) [H] Use Maple and the method of part (a) to prove that $\frac{5}{2} < e < 3$. How many partition points do you need?
3. [R] Find the derivatives of
 - a) $f(x) = \ln \sqrt{x^3 + 1}$
 - b) $g(x) = e^{|x|}$
 - c) $h(x) = \ln(\ln(\ln x))$
 - d) $q(x) = e^{\ln(x^5+6)}$

Problems 9.3 : The exponential function and
9.5 : Integration and the $\ln$ function

4. [R] Find
 - a) $\int \frac{e^{2x}}{1 + e^{2x}} dx$
 - b) $\int \frac{e^{1/x}}{x^2} dx$
 - c) $\int 3^x dx$
 - d) $\int \frac{e^{\sqrt{x}}}{8\sqrt{x}} dx$
 - e) $\int \frac{\ln x}{x} dx$
 - f) $\int \cot x dx$.


(Hint for part (f): express $\cot$ in terms of $\sin$ and $\cos$.)

5. [R] Sketch the curves
 - a) $y = \ln(1 + e^x)$
 - b) $y = \frac{(e^x + x)}{(e^x - x)}$.

6. [R]

- a) Sketch the curve $y = \frac{\ln x}{x}$, noting any turning points and asymptotes.
 b) By using (a) or otherwise, prove that $\pi^e < e^\pi$.

7. [R] [V]



- a) From the graph, explain why $\frac{1}{1+t} \leq \ln\left(1 + \frac{1}{t}\right) \leq \frac{1}{t}$ whenever $t \geq 0$.
 b) Deduce that $\lim_{t \rightarrow \infty} \ln\left(1 + \frac{1}{t}\right)^t = 1$ and hence find the value of $\lim_{t \rightarrow \infty} \left(1 + \frac{1}{t}\right)^t$.

Problems 9.6 : Logarithmic differentiation8. [R] Use logarithmic differentiation to find $\frac{dy}{dx}$ if

- a) $y = 3^x$ b) $y = \left(\frac{x^3 - 3}{1 + x^2}\right)^{1/5}$
 c) $y = (\sin x)^{\sin x}$ d) $y = \sin(x^{\sin x})$.

Problems 9.7 : Indeterminate forms with powers

9. [R] Calculate the following limits:

- a) [V] $\lim_{x \rightarrow \infty} \frac{\ln x}{x^a}$, $a > 0$ b) $\lim_{x \rightarrow 0+} x^a \ln x$, $a > 0$
 c) $\lim_{x \rightarrow 0+} x^x$ d) $\lim_{x \rightarrow 0+} x^{2/\ln x}$
 e) $\lim_{x \rightarrow \infty} x^{1/x}$ f) $\lim_{x \rightarrow \infty} a^{1/x}$, $a > 0$
 g) $\lim_{x \rightarrow \infty} \left(1 + \frac{a}{x}\right)^x$ h) $\lim_{x \rightarrow \infty} x^{100} e^{-x}$
 i) $\lim_{x \rightarrow \infty} p(x) e^{-x}$, where p is any polynomial.

10. [H] Prove that the functions $f : (-1, \infty) \rightarrow \mathbb{R}$ and $g : (-1, \infty) \rightarrow \mathbb{R}$, given by

$$f(x) = \ln(1+x) - \left(x - \frac{x^2}{2}\right) \quad \text{and} \quad g(x) = \left(x - \frac{x^2}{2} + \frac{x^3}{3}\right) - \ln(1+x),$$

are increasing on $(0, \infty)$. Deduce that

$$x - \frac{x^2}{2} < \ln(1+x) < x - \frac{x^2}{2} + \frac{x^3}{3}$$

whenever $x > 0$.

Chapter 10

The hyperbolic functions

A telecommunications company wishes to suspend a cable from a sequence of telegraph poles in a cost effective manner. There are at least two competing considerations. First, it is desirable to minimise the length of cable used. Second, it is desirable to minimise the amount of tension in the cable (since an increase in tension requires an increase in cable strength). The configuration in Figure 10.1 (a) requires a short length of strong cable while that in Figure 10.1 (b) reduces the tension in the cable at the cost of increasing its length. Is there an optimal configuration for this problem?

In order to solve this problem, one needs an equation which describes the curve of a suspended cable. Such a curve is called a *catenary*, after the Latin word for ‘chain.’ Galileo claimed that every catenary was a parabola, but this was later disproved in 1669. Two decades later, the Swiss mathematician Jakob Bernoulli issued a public challenge to find the general equation that describes a catenary. Three mathematicians Gottfried Leibniz, Christiaan Huygens and Johann Bernoulli (Jakob’s brother) independently derived the correct equation:

$$y = \frac{a}{2}(e^{x/a} + e^{-x/a}) \quad (10.1)$$

(see Figure 10.1 (c)). In this chapter we study the function described by equation (10.1) and other functions related to it. Such functions are important to mathematics, engineering, physics (especially the theory of relativity) and architecture.

We note from the outset that a large number of new formulae will be introduced in this chapter. However, students do not need to memorise all of them. Those which should be memorised are listed in Section 10.7.




Figure 10.1: Catenaries.

10.1 Hyperbolic sine and cosine functions

(Ref: SH10 §7.8)

We mentioned in the introduction that the equation describing a freely suspended chain is given by $y = f(x)$, where

$$f(x) = \frac{a}{2}(e^{x/a} + e^{-x/a}).$$

To begin we study the simplest case: when $a = 1$.

Definition 10.1.1. The *hyperbolic cosine function* $\cosh : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula

$$\cosh x = \frac{1}{2}(e^x + e^{-x}) \quad \forall x \in \mathbb{R}.$$

The reasoning behind the name ‘hyperbolic cosine’ will gradually become apparent. Since $\cosh$ is a linear combination of exponential functions, it is differentiable and

$$\frac{d}{dx}(\cosh x) = \frac{1}{2} \frac{d}{dx}(e^x + e^{-x}) = \frac{1}{2}(e^x - e^{-x}).$$

The derivative of $\cosh$ is important in its own right and is given a special name.

Definition 10.1.2. The *hyperbolic sine function* $\sinh : \mathbb{R} \rightarrow \mathbb{R}$ is defined by the formula

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) \quad \forall x \in \mathbb{R}.$$

Some people pronounce $\sinh$ as ‘shine.’ One can easily show that the derivative of $\sinh$ is $\cosh$. Thus we have

$$\boxed{\frac{d}{dx}(\sinh x) = \cosh x \quad \text{and} \quad \frac{d}{dx}(\cosh x) = \sinh x.} \quad (10.2)$$

We now consider the graphs of these functions.

Example 10.1.3. Sketch the graph of $\sinh$.

Solution. We begin by identifying a few key features of the $\sinh$ function.

- Since

$$\sinh(-x) = \frac{1}{2}(e^{-x} - e^x) = -\frac{1}{2}(e^x - e^{-x}) = -\sinh x,$$

we see that $\sinh$ is an odd function. As a corollary, $\sinh(0) = 0$.

- Since

$$\frac{d}{dx}(\sinh x) = \cosh x = \frac{1}{2}(e^x + e^{-x}) > 0,$$

$\sinh$ is an increasing function.

- Since $\lim_{x \rightarrow \infty} e^{-x} = 0$, $\sinh x$ gets arbitrarily close to $\frac{1}{2}e^x$ as $x \rightarrow \infty$.




Figure 10.2: Graphs of sinh and cosh.

From these observations we obtain a sketch of the graph of sinh (see Figure 10.2). □

Example 10.1.4. Sketch the graph of cosh.

Solution. Some properties of the cosh function are listed below:

- cosh is an even function;
- $\cosh(0) = 1$;
- cosh is decreasing on $(-\infty, 0)$, stationary at 0 and increasing on $(0, \infty)$;
- $\cosh x \geq 1$ for all x in $\mathbb{R}$;
- $\cosh x$ gets arbitrarily close to $\frac{1}{2}e^x$ as $x \rightarrow \infty$.

The reader should verify each of these properties by appealing to Definition 10.1.1. From these observations, we obtain a sketch of the graph of cosh (see Figure 10.2). □

The functions cosh and sinh are related by the following important identity.

Proposition 10.1.5. *If x is a real number then*

$$\cosh^2 x - \sinh^2 x = 1.$$

Proof. If $x \in \mathbb{R}$ then

$$\begin{aligned} \cosh^2 x - \sinh^2 x &= \left(\frac{1}{2}(e^x + e^{-x}) \right)^2 - \left(\frac{1}{2}(e^x - e^{-x}) \right)^2 \\ &= \frac{1}{4} \times 2e^x \times 2e^{-x} \quad \text{difference of two squares} \\ &= 1, \end{aligned}$$

completing the proof. □

We are now in a position to appreciate why these functions are called ‘hyperbolic cosine’ and ‘hyperbolic sine.’ First, we examine their similarity to the trigonometric cosine and sine functions. The sine and cosine functions have the following properties:

$$\begin{array}{ll} \frac{d}{dx}(\cos x) = -\sin x & \frac{d}{dx}(\sin x) = \cos x \\ \cos \text{ is an even function} & \sin \text{ is an odd function} \\ \cos^2 x + \sin^2 x = 1. \end{array}$$

The functions $\sinh$ and $\cosh$ have analogous properties (with the occasional adjustment of a negative sign):

$$\begin{array}{ll} \frac{d}{dx}(\cosh x) = \sinh x & \frac{d}{dx}(\sinh x) = \cosh x \\ \cosh \text{ is an even function} & \sinh \text{ is an odd function} \\ \cosh^2 x - \sinh^2 x = 1. \end{array}$$

The next example explains the origin of the name *hyperbolic* in ‘hyperbolic cosine.’

Example 10.1.6. Sketch the curve defined by the parametric equation

$$\begin{cases} x(t) = \cosh t \\ y(t) = \sinh t, \end{cases} \quad (10.3)$$

where $t \in \mathbb{R}$.

Solution. The parameter t in equation (10.3) can be eliminated by Proposition 10.1.5 to obtain

$$\begin{aligned} 1 &= \cosh^2 t - \sinh^2 t \\ &= (x(t))^2 - (y(t))^2, \end{aligned}$$

or simply

$$x^2 - y^2 = 1.$$

This equation describes an hyperbola. Since $\cosh t > 0$ for every real number t , it follows that $x > 0$ and hence the curve is the branch of the hyperbola that lies in the right-half plane (see the black curve in Figure 10.3 (b)). Its asymptotes are the lines $y = x$ and $y = -x$.

(The other branch of the hyperbola is parameterised by

$$\begin{cases} x(t) = -\cosh t \\ y(t) = \sinh t \end{cases}$$

where $t \in \mathbb{R}$, and is shown in Figure 10.3 (b) in gray.) □

Hence we use trigonometric sine and cosine to parameterise a circle or ellipse, and hyperbolic sine and cosine to parameterise an hyperbola.




Figure 10.3: Sketch for Example 10.1.6.

10.2 Other hyperbolic functions

(Ref: SH10 §7.9)

Other hyperbolic functions are defined in analogy to the trigonometric functions. Thus we have

$$\begin{aligned}\tanh x &= \frac{\sinh x}{\cosh x}, & \coth x &= \frac{\cosh x}{\sinh x}, \\ \operatorname{sech} x &= \frac{1}{\cosh x}, & \operatorname{cosech} x &= \frac{1}{\sinh x}.\end{aligned}$$

Some people pronounce $\tanh$ as ‘than.’

The derivative of each of these functions is easy to compute.

Example 10.2.1. Compute the derivative of $\tanh$.


Solution. By the definition of $\tanh$,

$$\begin{aligned}\frac{d}{dx}(\tanh x) &= \frac{d}{dx} \left(\frac{\sinh x}{\cosh x} \right) \\ &= \frac{\cosh x \frac{d}{dx}(\sinh x) - \sinh x \frac{d}{dx}(\cosh x)}{\cosh^2 x} && \text{(by the quotient rule)} \\ &= \frac{\cosh^2 x - \sinh^2 x}{\cosh^2 x} && \text{(by (10.2))} \\ &= \frac{1}{\cosh^2 x} && \text{(by Proposition 10.1.5)} \\ &= \operatorname{sech}^2 x.\end{aligned}$$

□

A complete list of derivatives will be given in Section 10.4.

Example 10.2.2. Sketch the graph of the function $\tanh$.

Figure 10.4: The graph of $\tanh$.

Solution. Since $\tanh x = \frac{\sinh x}{\cosh x}$ and $\cosh x > 0$ for all x in $\mathbb{R}$, we see that $\text{Dom}(\tanh) = \mathbb{R}$. Some properties of $\tanh$ are listed below.

- By using the fact that $\cosh$ is even and $\sinh$ is odd, we see that

$$\tanh(-x) = \frac{\sinh(-x)}{\cosh(-x)} = \frac{-\sinh x}{\cosh x} = -\tanh(x).$$

Hence $\tanh$ is an odd function. It follows that $\tanh(0) = 0$.

- Since $\frac{d}{dx}(\tanh x) = \text{sech}^2 x > 0$ for all x in $\mathbb{R}$, the function $\tanh$ is increasing everywhere.
- By writing $\tanh x$ in terms of exponentials we find that

$$\begin{aligned} \tanh x &= \frac{\sinh x}{\cosh x} \\ &= \frac{e^x - e^{-x}}{e^x + e^{-x}} \\ &= \frac{1 - e^{-2x}}{1 + e^{-x}} \\ &\rightarrow 1 \end{aligned}$$

as $x \rightarrow \infty$.

- Finally, the slope of the graph of $\tanh$ at the origin is

$$\left. \frac{d}{dx}(\tanh x) \right|_{x=0} = \text{sech}^2 0 = 1.$$

These features are shown in the following sketch in Figure 10.4. □

10.3 Hyperbolic identities

(Ref: SH10 §7.8)

It has already been observed that some identities involving $\cosh$ and $\sinh$ are *almost* analogous to

those for $\cos$ and $\sin$. This analogy can also be seen in the following larger list of identities. There are the ‘difference of squares’ identities

$$\begin{aligned}\cosh^2 x - \sinh^2 x &= 1, \\ 1 - \tanh^2 x &= \operatorname{sech}^2 x, \\ \coth^2 x - 1 &= \operatorname{cosech}^2 x,\end{aligned}$$

the ‘sum and difference’ formulae

$$\begin{aligned}\sinh(x \pm y) &= \sinh x \cosh y \pm \cosh x \sinh y, \\ \cosh(x \pm y) &= \cosh x \cosh y \pm \sinh x \sinh y, \\ \tanh(x \pm y) &= \frac{\tanh x \pm \tanh y}{1 \pm \tanh x \tanh y},\end{aligned}$$

and the ‘double-angle’ formulae

$$\begin{aligned}\sinh(2x) &= 2 \sinh x \cosh x, \\ \cosh(2x) &= \cosh^2 x + \sinh^2 x, \\ \tanh(2x) &= \frac{2 \tanh x}{1 + \tanh^2 x}.\end{aligned}$$

The following remark gives an easy method for remembering these identities.

Remark 10.3.1. There is a convenient mnemonic for these hyperbolic identities, provided that you know the corresponding trigonometric identities. Simply replace $\cos$ with $\cosh$ and $\sin$ with $i \sinh$. For example,

$$\begin{aligned}\cos^2 x + \sin^2 x = 1 &\longrightarrow \cosh^2 x + i^2 \sinh^2 x = 1 \\ &\longrightarrow \cosh^2 x - \sinh^2 x = 1,\end{aligned}$$

and

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ &\longrightarrow i \sinh(x + y) = i \sinh x \cosh y + i \cosh x \sinh y \\ &\longrightarrow \sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y.\end{aligned}$$

Since $\tan = \frac{\sin}{\cos}$ we replace $\tan$ with $\frac{i \sinh}{\cosh} = i \tanh$. Similarly $\sec$ is replaced with sech . Thus

$$\begin{aligned}1 + \tan^2 x = \sec^2 x &\longrightarrow 1 + i^2 \tanh^2 x = \operatorname{sech}^2 x \\ &\longrightarrow 1 - \tanh^2 x = \operatorname{sech}^2 x.\end{aligned}$$

We emphasise that Remark 10.3.1 provides a way to *remember* each formula, not to *prove* it. Proofs are constructed using techniques that are illustrated in the following example.

Example 10.3.2. (a) Prove the identity

$$\cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y.$$

(b) Hence show that

$$\cosh(x - y) = \cosh x \cosh y - \sinh x \sinh y.$$

(c) By using the result of (a), prove the double-angle formula for cosh.

(d) By assuming that

$$\sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y, \quad (10.4)$$

use the result of (a) to show that

$$\tanh(x + y) = \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y}.$$

Solution. (a) By the definition of cosh,

$$\text{LHS} = \frac{1}{2}(e^{x+y} + e^{-(x+y)})$$

while

$$\begin{aligned} \text{RHS} &= \frac{1}{2}(e^x + e^{-x})\frac{1}{2}(e^y + e^{-y}) + \frac{1}{2}(e^x - e^{-x})\frac{1}{2}(e^y - e^{-y}) \\ &= \frac{1}{4}(e^{x+y} + e^{x-y} + e^{-x+y} + e^{-x-y}) + \frac{1}{4}(e^{x+y} - e^{x-y} - e^{-x+y} + e^{-x-y}) \\ &= \frac{1}{2}e^{x+y} + \frac{1}{2}e^{-x-y} \\ &= \text{LHS}. \end{aligned}$$

This proves the identity. □

Remark 10.3.3. Note that in Example 10.3.2, each proof started by simplifying *either* the left-hand side of the equation, *or* the right-hand side of the equation. It would be wrong to begin the proof of (a) by

$$\begin{aligned} \cosh(x + y) &= \cosh x \cosh y + \sinh x \sinh y \\ \frac{1}{2}(e^{x+y} + e^{-(x+y)}) &= \frac{1}{2}(e^x + e^{-x})\frac{1}{2}(e^y + e^{-y}) + \frac{1}{2}(e^x - e^{-x})\frac{1}{2}(e^y - e^{-y}) \\ &\dots = \dots \end{aligned}$$

Such a method is flawed and can be used to ‘prove’ erroneous statements such as ‘ $1 = 0$ ’.

10.4 Hyperbolic derivatives and integrals

(Ref: SH10 §§7.8, 7.9)

Below is a list of derivatives for the hyperbolic functions:

$$\begin{aligned} \frac{d}{dx}(\sinh x) &= \cosh x & \frac{d}{dx}(\cosh x) &= \sinh x \\ \frac{d}{dx}(\tanh x) &= \text{sech}^2 x & \frac{d}{dx}(\coth x) &= -\text{cosech}^2 x \\ \frac{d}{dx}(\text{sech } x) &= -\text{sech } x \tanh x & \frac{d}{dx}(\text{cosech } x) &= -\text{cosech } x \coth x. \end{aligned}$$

These derivatives can be established by using the definition of each function, as in Example 10.2.1. The derivatives of sinh, cosh and tanh are the most important to remember.

Corresponding to these derivatives are the indefinite integrals

$$\int \sinh x \, dx = \cosh x + C, \quad \int \text{sech}^2 x \, dx = \tanh x + C$$

and so on.

Example 10.4.1. Calculate the derivative of the function f given by

$$f(x) = \ln(\operatorname{sech}(2x)).$$

Solution. First we have

$$\frac{d}{dx}(\operatorname{sech} 2x) = -2 \operatorname{sech}(2x) \tanh(2x).$$

Hence the chain rules gives

$$f'(x) = \frac{-2 \operatorname{sech}(2x) \tanh(2x)}{\operatorname{sech}(2x)} = -2 \tanh(2x).$$

□

Example 10.4.2. Find the following integrals:

$$(a) \int_0^{\frac{1}{5} \ln 3} \sinh 5x \, dx$$

$$(b) \int x \cosh x \, dx$$

$$(c) \int \frac{\operatorname{sech}^2(\sqrt{x})}{\sqrt{x}} \, dx$$

$$(d) \int e^x \sinh x \, dx.$$

Solution. (a) We have

$$\begin{aligned} \int_0^{\frac{1}{5} \ln 3} \sinh 5x \, dx &= \left[\frac{1}{5} \cosh 5x \right]_0^{\frac{1}{5} \ln 3} \\ &= \frac{1}{10} \left[e^{5x} + e^{-5x} \right]_0^{\frac{1}{5} \ln 3} \\ &= \frac{1}{10} \left(e^{\ln 3} + e^{-\ln 3} - e^0 - e^0 \right) \\ &= \frac{1}{10} \left(3 + \frac{1}{3} - 1 - 1 \right) \\ &= \frac{2}{15}. \end{aligned}$$

□

10.5 The inverse hyperbolic functions

(Ref: SH10 §7.9)

The functions $\sinh : \mathbb{R} \rightarrow \mathbb{R}$ and $\tanh : \mathbb{R} \rightarrow (-1, 1)$ are both increasing on $\mathbb{R}$ and hence have well-defined inverses. However, $\cosh : \mathbb{R} \rightarrow [1, \infty)$ is not one-to-one on $\mathbb{R}$. To obtain an inverse, one must restrict the domain of $\cosh$. The following definition gives the conventional restriction.




Figure 10.5: Graphs of the hyperbolic functions (in gray) and their inverses (in black).

Definition 10.5.1. The function $\cosh^{-1} : [1, \infty) \rightarrow [0, \infty)$ is defined to be the inverse function of the restricted hyperbolic cosine function

$$\cosh : [0, \infty) \rightarrow [1, \infty).$$

Graphs of the functions $\sinh^{-1}$, $\cosh^{-1}$ and $\tanh^{-1}$ are illustrated in Figure 10.5.

Expressions involving the hyperbolic functions and their inverses can be simplified using the identities of Section 10.3. It is important to note that

$$\cosh^{-1}(\cosh x) = x$$

only when $x \geq 0$.

Example 10.5.2. Simplify

- (a) $\sinh\left(\cosh^{-1}\frac{4}{3}\right)$,
- (b) $\cosh^{-1}(\cosh(-7))$, and
- (c) $\cosh(2\sinh^{-1}3)$.

Solution. (a) The identity

$$\cosh^2 t - \sinh^2 t = 1$$

can be rearranged to give

$$\sinh^2 t = \cosh^2 t - 1.$$

By substituting $\cosh^{-1}\frac{4}{3}$ for t , we obtain

$$\sinh^2\left(\cosh^{-1}\frac{4}{3}\right) = \left(\frac{4}{3}\right)^2 - 1 = \frac{7}{9}.$$

We need to decide whether to take the positive or negative square root. Since $\cosh^{-1}$ is a nonnegative function, we have $\cosh^{-1}\frac{4}{3} > 0$ and hence

$$\sinh\left(\cosh^{-1}\frac{4}{3}\right) > 0.$$

Therefore

$$\sinh\left(\cosh^{-1}\frac{4}{3}\right) = +\frac{\sqrt{7}}{3}.$$

(b) We cannot write $\cosh^{-1}(\cosh(-7)) = -7$ because $\cosh^{-1}$ is only the inverse function of $\cosh$ with domain restricted to $[0, \infty)$. However, since $\cosh$ is an even function,

$$\cosh^{-1}(\cosh(-7)) = \cosh^{-1}(\cosh(7)) = 7.$$

(c) We use the double-angle formula

$$\cosh(2t) = 1 + 2 \sinh^2 t$$

for $\cosh$:

$$\begin{aligned} \cosh(2 \sinh^{-1} 3) &= 1 + 2(\sinh(\sinh^{-1} 3))^2 \\ &= 1 + 2 \times 3^2 \\ &= 19. \end{aligned}$$

□

Each of the functions $\sinh$, $\cosh$ and $\tanh$ can be expressed in terms of the exponential function. It is not surprising, then, that their inverses can be expressed in terms of the natural logarithm.

Proposition 10.5.3. *The following identities hold:*

$$\begin{aligned} \sinh^{-1} x &= \ln(x + \sqrt{x^2 + 1}) & \forall x \in \mathbb{R}, \\ \cosh^{-1} x &= \ln(x + \sqrt{x^2 - 1}) & \forall x \in [1, \infty), \\ \tanh^{-1} x &= \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right) & \forall x \in (-1, 1). \end{aligned}$$

Proof. We prove the formula for $\sinh^{-1}$ only; the others can be proved similarly.

Suppose that $y = \sinh^{-1} x$. Then

$$\sinh y = x,$$

which means that

$$\frac{1}{2}(e^y - e^{-y}) = x.$$

If we multiply this equation through by $2e^y$ then we obtain

$$e^{2y} - 1 = 2xe^y,$$

which is a quadratic equation in e^y . Further rearrangement gives

$$(e^y)^2 - 2x(e^y) = 1,$$

to which we complete the square to obtain

$$(e^y - x)^2 = 1 + x^2.$$

Since $e^y > 0$, we take the positive square root. Thus

$$e^y = x + \sqrt{x^2 + 1}.$$

(Note that this may also be obtained by using the quadratic formula rather than completing the square). Taking the natural logarithm of both sides of this equation gives

$$y = \ln(x + \sqrt{x^2 + 1})$$

as required. □

By the inverse function theorem, the functions

$$\sinh^{-1} : \mathbb{R} \rightarrow \mathbb{R}, \quad \cosh^{-1} : (1, \infty) \rightarrow (0, \infty) \quad \text{and} \quad \tanh^{-1} : (-1, 1) \rightarrow \mathbb{R}$$

are differentiable on their respective domains. A formula for each derivative is given in the next proposition.

Proposition 10.5.4. *The derivatives of $\sinh^{-1}$, $\cosh^{-1}$ and $\tanh^{-1}$ are given by*

$$\begin{aligned} \frac{d}{dx}(\sinh^{-1} x) &= \frac{1}{\sqrt{x^2 + 1}}, \\ \frac{d}{dx}(\cosh^{-1} x) &= \frac{1}{\sqrt{x^2 - 1}}, \\ \frac{d}{dx}(\tanh^{-1} x) &= \frac{1}{1 - x^2}. \end{aligned}$$

Proof. We prove the formula for $\frac{d}{dx}(\sinh^{-1} x)$ only; the others can be proved similarly.

Suppose that $y = \sinh^{-1} x$. Then

$$\sinh y = x$$

and differentiating with respect to x gives

$$\cosh y \frac{dy}{dx} = 1.$$

Hence

$$\frac{dy}{dx} = \frac{1}{\cosh y} = \frac{1}{\cosh(\sinh^{-1} x)}. \quad (10.5)$$

The identity $\cosh^2 t - \sinh^2 t = 1$ gives

$$\cosh t = \sqrt{1 + \sinh^2 t}$$

(where we have taken the positive square root since $\cosh t > 0$ for all t). Hence

$$\cosh(\sinh^{-1} x) = \sqrt{1 + x^2}.$$

If we substitute this into (10.5) then we obtain

$$\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1 + x^2}}$$

as required. □

Remark 10.5.5. Students should be familiar with the proofs of Propositions 10.5.3 and 10.5.4.

10.6 Integration leading to the inverse hyperbolic functions

When propositions 10.5.3 and 10.5.4 for $\sinh^{-1}$ are restated in terms of indefinite integration we obtain

$$\int \frac{dx}{\sqrt{x^2 + 1}} = \sinh^{-1} x + C = \ln(x + \sqrt{x^2 + 1}) + C.$$

A more general version may be obtained by observing that, when $a > 0$,

$$\frac{d}{dx} \left(\sinh^{-1} \frac{x}{a} \right) = \frac{1}{a} \frac{1}{\sqrt{\left(\frac{x}{a}\right)^2 + 1}} = \frac{1}{\sqrt{x^2 + a^2}}.$$

In this manner, one is able to prove the formulae

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 + a^2}} &= \sinh^{-1} \frac{x}{a} + C \\ &= \ln(x + \sqrt{x^2 + a^2}) + (C - \ln a), \end{aligned} \quad a > 0,$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \cosh^{-1} \frac{x}{a} + C \\ &= \ln(x + \sqrt{x^2 - a^2}) + (C - \ln a), \end{aligned} \quad x \geq a > 0,$$

$$\begin{aligned} \int \frac{dx}{a^2 - x^2} &= \begin{cases} \frac{1}{a} \tanh^{-1} \frac{x}{a} + C, & |x| < a \\ \frac{1}{a} \coth^{-1} \frac{x}{a} + C, & |x| > a > 0 \end{cases} \\ &= \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + C, \end{aligned} \quad x^2 \neq a^2.$$

These formulae are included in the table of standard integrals that is issued at the final examination.

Example 10.6.1. Find

$$\begin{aligned} \text{(a)} \quad & \int \frac{dx}{\sqrt{1+9x^2}} \\ \text{(b)} \quad & \int \frac{dx}{\sqrt{x^2-2x+10}}. \end{aligned}$$

Solution. (a) A little algebraic manipulation shows that

$$\frac{1}{\sqrt{1+9x^2}} = \frac{1}{3\sqrt{(1/3)^2 + x^2}}.$$

Hence

$$\begin{aligned} \int \frac{dx}{\sqrt{1+9x^2}} &= \frac{1}{3} \int \frac{dx}{\sqrt{(1/3)^2 + x^2}} \\ &= \frac{1}{3} \sinh^{-1} \frac{x}{1/3} + C \\ &= \frac{1}{3} \sinh^{-1}(3x) + C. \end{aligned}$$

Alternatively, one could approach this problem by using the substitution $u = 3x$.

(b) The technique we apply to this example is known as ‘completing the square’:

$$x^2 - 2x + 10 = x^2 - 2x + 1 + 9 = (x-1)^2 + 3^2 = u^2 + 3^2,$$

where $u = x - 1$. Hence

$$\begin{aligned}\int \frac{dx}{\sqrt{x^2 - 2x + 10}} &= \int \frac{du}{\sqrt{u^2 + 3^2}} \\ &= \sinh^{-1} \frac{u}{3} + C \\ &= \sinh^{-1} \frac{x-1}{3} + C.\end{aligned}$$

□

10.7 A summary of important hyperbolic formulae

A large number of formulae was introduced in this chapter. Below is a list of formulae, with the more important appearing earlier in the list, that students are expected to know.

- The definitions of $\sinh$ and $\cosh$:

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) \qquad \cosh x = \frac{1}{2}(e^x + e^{-x})$$

- The derivatives of $\sinh$ and $\cosh$:

$$\frac{d}{dx}(\sinh x) = \cosh x \qquad \frac{d}{dx}(\cosh x) = \sinh x$$

- The definition of $\tanh$:

$$\tanh x = \frac{\sinh x}{\cosh x}$$

- The hyperbolic identity

$$\cosh^2 x - \sinh^2 x = 1$$

- The derivative of $\sinh^{-1}$:

$$\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1+x^2}}$$

- The derivative of $\tanh$:

$$\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$$

Other formulae need not be memorised, though students will do well to note the following points.

- Derivatives for the inverse hyperbolic functions can be read from the table of standard integrals.
- Identities which express the inverse hyperbolic functions in terms of the natural logarithm (see Proposition 10.5.3) can be read from the table of standard integrals.
- The definition of $\coth$, sech and cosech are in exact analogy with the definition of $\cot$, $\sec$ and cosec .
- The hyperbolic identities listed in Section 10.3 are easily remembered from their trigonometric counterparts (see Remark 10.3.1).




Figure 10.6: Forces acting on a suspended cable.

- It is not expected that students will memorise the derivatives of $\coth$, sech and cosech . However, they can be easily derived from first principles provided that the derivatives of $\sinh$ and $\cosh$ are known.

A table of standard integrals will be issued at the final examination.

10.8 (Appendix):The catenary

We end this chapter showing that the $\cosh$ function describes the curve of a suspended cable.

Suppose that a cable of uniform mass is suspended from two fixed points. To describe the equation of the resulting curve in terms of x and y , we fix a coordinate system such that vertex of the curve passes through the point $A(0, y_0)$ as in Figure 10.6. (The ordinate y_0 can be any real number; its choice only fixes the coordinate system and does not alter the shape of the catenary.) Consider a point $P(x, y)$ that lies on the curve and suppose that

- T is the tension in the cable at $P(x, y)$,
- H is the tension in the cable at $A(0, y_0)$,
- $\ell(x)$ is the length of the cable from $A(0, y_0)$ to $P(x, y)$, and
- w is the weight per unit length of the cable.

(see Figure 10.6). Thus the gravitational force acting on the cable along the segment AP is given by $w\ell(x)$. We note that the quantities T , θ and ℓ depend on x but that w and H do not. We also make the natural assumption that the catenary is described by a differentiable function.

Theorem 10.8.1. *There is a choice of coordinate system such that the catenary in Figure 10.6 is described by the equation*

$$y = \frac{H}{w} \cosh\left(\frac{wx}{H}\right).$$

To prove the theorem, it suffices to show that if the catenary is described by $y = f(x)$, where f is a differentiable function, then

$$f(x) = \frac{H}{w} \cosh\left(\frac{wx}{H}\right)$$


for a particular choice of y_0 .

We begin with a general formula that relates a function f to its arc length function ℓ by a differential equation.

Lemma 10.8.2. *Suppose that f is a differentiable function. Then the arc length function ℓ is also differentiable and*

$$\ell'(x) = \sqrt{1 + (f'(x))^2} \quad (10.6)$$

Sketch proof. Suppose that h is a small nonzero real number and consider the diagram below.



By Pythagoras' theorem,

$$\ell(x+h) - \ell(x) \approx \sqrt{h^2 + (f(x+h) - f(x))^2}.$$

If we divide both sides by h then

$$\frac{\ell(x+h) - \ell(x)}{h} \approx \sqrt{1 + \left(\frac{f(x+h) - f(x)}{h}\right)^2}.$$

As $h \rightarrow 0$ the approximation gets better and we obtain

$$\ell'(x) = \sqrt{1 + (f'(x))^2}$$

as required. □

The next lemma uses facts specific to the catenary.

Lemma 10.8.3. *If the equation of the catenary in Figure 10.6 is given by $y = f(x)$, then f satisfies the following conditions:*

$$f(0) = y_0, \quad (10.7)$$

$$f'(0) = 0, \quad (10.8)$$

$$f''(x) = \frac{w}{H} \sqrt{1 + (f'(x))^2}. \quad (10.9)$$

Proof. The first two conditions are a result of our choice of coordinate system and the assumption that f is differentiable. It remains to establish the third condition.

Since the cable is stationary, the forces $w \ell(x)$ and $T \sin \theta$ that act vertically cancel each other out. That is,

$$T \sin \theta = w \ell(x).$$

Similarly, the horizontal forces H and $T \cos \theta$ also cancel each other out and so

$$T \cos \theta = H.$$

Using these equations, and the fact that $f'(x) = \tan \theta$ at the point $P(x, y)$, gives

$$f'(x) = \tan \theta = \frac{T \sin \theta}{T \cos \theta} = \frac{w \ell(x)}{H}.$$

So far we have

$$f'(x) = \frac{w \ell(x)}{H}.$$

Since w and H are constants, differentiating this equation with respect to x gives

$$f''(x) = \frac{w}{H} \ell'(x).$$

Therefore

$$f''(x) = \frac{w}{H} \sqrt{1 + (f'(x))^2}.$$

by equation (10.6). □

Proof of Theorem 10.8.1. It remains to find all functions f that satisfy the conditions of Lemma 10.8.3. If we let u denote $f'(x)$ and b denote w/H , then equation (10.9) becomes

$$\frac{du}{dx} = b \sqrt{1 + u^2}.$$

Hence

$$\frac{dx}{du} = \frac{1}{b \sqrt{1 + u^2}}.$$

Every solution to this differential equation is of the form

$$x = \frac{1}{b} \sinh^{-1} u + C_1.$$

Hence we have

$$f'(x) = u = \sinh(bx - bC_1).$$

Enforcing condition (10.8) shows that either $b = 0$ or $C_1 = 0$. Now if $b = 0$ then $f'(x) = 0$ and f is simply a constant function (that is, the cable is pulled tight and exhibits no sag). On the other hand, if $b \neq 0$ then $C_1 = 0$ and hence

$$f'(x) = \sinh(bx).$$

Therefore

$$f(x) = \frac{\cosh(bx)}{b} + C_2$$

and enforcing the condition (10.7) yields $C_2 = \frac{1}{b} - y_0$. If we choose y_0 to be $1/b$ then $C_2 = 0$ and the function f that describes the catenary is given by

$$f(x) = \frac{\cosh(bx)}{b} = \frac{H}{w} \cosh\left(\frac{wx}{H}\right),$$

proving the theorem. □

10.9 Maple notes

Maple knows about the hyperbolic functions `sinh`, `cosh`, `tanh`, `csch`, `sech`, and `coth`, and inverse hyperbolic functions `arsinh`, `arcosh`, `artanh`, `arccsch`, `arcsech`, and `arcoth`. For example,

```
> int(arctanh(x),x);
```

$$x \operatorname{arctanh}(x) + \frac{1}{2} \ln(1 - x^2)$$

```
> # which is closely analogous to
```

```
> int(arctan(x),x);
```


$$x \operatorname{arctan}(x) - \frac{1}{2} \ln(1 + x^2)$$

```
> # BUT the following result does not follow the analogy between trig and hyperbolic functions
```

```
> int(sech(x),x);
```

$$\operatorname{arctan}(\sinh(x))$$

```
> plot(arccsch(x),x=-6..6, y=-6..6, discontinuity=true, numpoints=100);
```




Problems for Chapter 10

Problems 10.1 : Hyperbolic sine and cosine functions and
10.2 : Other hyperbolic functions

1. [R] [V] Define $\sinh x$ and $\cosh x$. Hence show that
 - a) $\frac{d}{dx}(\cosh 6x) = 6 \sinh 6x$;
 - b) $\ln(\sinh x) < x - \ln 2$ whenever $x > 0$.
2. [R] By expressing the following hyperbolic functions in terms of $\sinh x$ and $\cosh x$, find the derivative of each function f given below.
 - a) $f(x) = \tanh x$
 - b) $f(x) = \operatorname{sech} x$
 - c) $f(x) = \coth x$
3. [R] In each case, find $f'(x)$.
 - a) $f(x) = \sinh(3x^2)$
 - b) $f(x) = \cosh(\frac{1}{x})$
 - c) $f(x) = \sinh(\ln x)$

Problems 10.3 : Hyperbolic identities

4. [R]
 - a) Given the formula $\sinh(A+B) = \sinh A \cosh B + \cosh A \sinh B$, find a formula for $\sinh 2x$. By differentiation or otherwise, find a formula for $\cosh 2x$.
 - b) [H] Using the results of part (a), express $\sinh 3x$ as a cubic polynomial in $\sinh x$. Hence, or otherwise, find $\int \sinh^3 x \, dx$.
5. [R] Show that $\cosh x + \sinh x = e^x$. Deduce that $(\cosh x + \sinh x)^n = \cosh nx + \sinh nx$.
6. [H] [V] Consider the hyperbola $x^2 - y^2 = 1$, where $x \geq 1$.



- a) Using the definitions of $\cosh$ and $\sinh$, prove that, for every real number t , the point $(\cosh t, \sinh t)$ lies on the hyperbola.

- b) When $t > 0$, let $A(t)$ denote the shaded region in the diagram. Explain why

$$A(t) = \frac{1}{2} \cosh t \sinh t - \int_1^{\cosh t} \sqrt{x^2 - 1} \, dx.$$

- c) By first calculating $A'(t)$, prove that $A(t) = \frac{t}{2}$.

Problems 10.4 : Hyperbolic derivatives and integrals

7. [R] Evaluate the following integrals.

$$\begin{array}{ll} \text{a)} \quad \int \cosh(4x) \, dx & \text{b)} \quad \int_0^{\frac{1}{3} \ln 2} \sinh 3x \, dx \\ \text{c)} \quad \int \cosh^2 x \, dx & \text{d)} \quad \int \frac{\sinh(\sqrt{x})}{\sqrt{x}} \, dx \end{array}$$

Problems 10.5 : The inverse hyperbolic functions

8. [R] Simplify $\cosh(\sinh^{-1}(3/4))$, $\cosh^{-1}(\cosh(-3))$ and $\sinh(\tanh^{-1}(5/13))$.

9. [R] Show that

$$\text{a)} \quad \frac{d}{dx}(\cosh^{-1} x) = \frac{1}{\sqrt{x^2 - 1}}, \text{ for } x > 1 \qquad \text{b)} \quad \frac{d}{dx}(\tanh^{-1} x) = \frac{1}{1 - x^2}.$$

10. [R] Show that

$$\begin{array}{ll} \text{a)} \quad \cosh^{-1} x = \ln(x + \sqrt{x^2 - 1}) & \forall x \in [1, \infty) \\ \text{b)} \quad [\mathbf{V}] \quad \tanh^{-1} x = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) & \forall x \in (-1, 1). \end{array}$$

11. [R] Find $\frac{dy}{dx}$ if

$$\begin{array}{ll} \text{a)} \quad y = \sinh^{-1}(2x) \\ \text{b)} \quad y = \tanh^{-1}(1/x) \\ \text{c)} \quad y = \cosh^{-1}(\sec x) & \text{whenever } 0 < x < \pi/2. \end{array}$$

Problems 10.6 : Integration leading to the inverse hyperbolic functions

12. [R] Find

$$\begin{array}{lll} \text{a)} \quad [\mathbf{V}] \quad \int \frac{dx}{\sqrt{1+4x^2}} & \text{b)} \quad \int_0^{1/2} \frac{dx}{1-x^2} & \text{c)} \quad \int \frac{dx}{\sqrt{x^2+4x+13}}. \end{array}$$

13. [X] Sketch the function sech^{-1} . What is its maximal domain? For $y = \operatorname{sech}^{-1} x$, show that

$$\text{a)} \quad \frac{dy}{dx} = \frac{-1}{x\sqrt{1-x^2}} \qquad \text{b)} \quad y = \ln \left(\frac{1 + \sqrt{1-x^2}}{x} \right).$$

Answers to selected problems

Chapter 1

1. a) The set of integers between $-\pi$ and π .
c) The empty set.
3. Answer for both: the interior and boundary of the triangle with vertices at $(0, 0)$, $(2, 0)$ and $(2, 4)$.
4. a) $x < 0$ or $x > 1$ b) $1 < x < 2$ c) $x < -2$ or $x > 0$
d) $-1 < x < 1$ e) $-2 \leq x < 1$ or $x \geq 3$
5. a) $-4 < x < 2$ b) $x < -5$ or $x > 1$
c) $-1 < x < -1/3$ d) $x > 0$
6. c) From (a) we have $x^2 + \frac{1}{x^2} \geq 2$ with equality if and only if $x = \pm 1$.
7. a) F b) F c) T d) T e) F
8. Hint: $(x^2 + y^2)^2 \geq 4x^2y^2$.
10. a) $-\sqrt{5} \leq x \leq \sqrt{5}; \quad 0 \leq y \leq \sqrt{5}$
b) $x \leq -\sqrt{5}$ or $x \geq \sqrt{5}; \quad y \geq 0$
c) $x \neq 8; \quad y \neq 0$
d) $[1, \infty); [0, \infty)$
e) $(1, \infty); (0, \infty)$
f) $\{x \in \mathbb{R} : 2n\pi \leq x \leq (2n+1)\pi; n \in \mathbb{Z}\}; [0, 1]$
g) The union of the intervals $[-\frac{7\pi}{6} + 2k\pi, \frac{\pi}{6} + 2k\pi]$ where $k \in \mathbb{Z}; \quad 0 \leq y \leq \sqrt{3}$
h) $\{x \in \mathbb{R} : x \neq (2n+1)\pi/2, n \text{ an integer}\}; [1, \infty)$
i) $\mathbb{R}; [-1, \infty)$
11. a) 22 b) $x^2 + 10x + 22$ c) 6 d) $x^2 + 2$
12. a) $x - 1 + 1/\sqrt{x-1}$ b) $\sqrt{x-1}$ c) $(x-1)^{3/2}$ d) $(1/\sqrt{x-1}) - 1$
16. $[4, 13]$
17. $x = 1, 7$

18. a) If $p(x) = a_0 + a_1x + \cdots + a_nx^n$ then $p(q(x)) = a_0 + a_1q(x) + a_2(q(x))^2 + \cdots + a_n(q(x))^n$.
Products and sums of polynomials are again polynomials.
b) Yes.

Chapter 2

1. a) 1 b) 2 c) 0
d) Doesn't exist ($\rightarrow \infty$). e) 5 f) Doesn't exist.
2. a) 0 b) 0
4. b) 0
5. a) 1 b) $M = 10$ (best possible) c) $M = 1/\sqrt{\epsilon}$ will do.
6. a) 4 b) 0 c) 0 d) 0 e) 0
7. a) Not necessarily, as the information given indicates only that the inequality holds for a subset of (ϵ^{-1}, ∞) .
b) Yes. In fact one can prove that $\lim_{x \rightarrow \infty} g(x) = 5$ from the definition of the limit by taking M to be $\frac{1}{\epsilon}$.
8. a) 50 metres per second b) $5 \ln 50 \approx 19.56$ seconds after leaving the plane.
9. a) Yes. If limit of $f(x)$ as $x \rightarrow \infty$ does not exist and $f(x) \neq 0$, then
 $\lim_{x \rightarrow \infty} (f(x) - f(x)) = 0$ and $\lim_{x \rightarrow \infty} (f(x)/f(x)) = 1$.
b) Yes, since $g(x) = (f(x) + g(x)) - f(x)$.
c) No, as in (b).
d) No. For example if $f(x) = 0$ for all x and $\lim_{x \rightarrow \infty} g(x)$ does not exist, we have
 $\lim_{x \rightarrow \infty} (f(x)g(x)) = 0$.
10. a) 10 b) 4 c) 3 d) $-1/9$
11. a) -1 b) 1 c) No
12. a) Doesn't exist. b) Doesn't exist. c) Doesn't exist. d) Doesn't exist.
13. a) 0 b) 0
14. a) $|CB| = \theta$, $|CA| = \sin \theta$, $|DB| = \tan \theta$.
15. Neither the left-hand nor right-hand limits exist due to wild oscillatory behaviour.

Chapter 3

1. b) Yes
2. a) Continuous everywhere. b) Continuous everywhere except at $\pi/2$.
3. $k = 8$
5. Use the intermediate value theorem.
9. a) Yes b) Yes c) No d) Yes

Chapter 4

2. a) $5(4x^3 + 21x^6)$ b) $(4x^3 - 2)(4x^2 + 2x + 4) + (x^4 - 2x)(8x + 2)$
 c) $(16y - y^4)/(y^3 + 8)^2$ d) $(2x^2 - 4)/(x^2 - 4)^{1/2}$
 e) $-4/(t^2 - 4)^{3/2}$ f) $3 \cos 3y + 12 \cos 2y \sin 2y$
 g) $(4x^3 - x^4)e^{-x}$ h) $x \ln(x^3 + 1) + 3x^2(x^2 + 1)/2(x^3 + 1)$
 i) $\sec^2 x$ j) $-\tan x$
3. a) 0 b) 0 c) $f'(0) = 0$
4. a) i) $x \neq 0$ ii) all x b) i) all x ii) all x
 c) i) $x \neq -2$ ii) $x \neq -2$
7. $2pf'(a)$
8. a) $x + 17\pi + \cos 2x$ b) $1 - 2 \sin 2x$ c) $2 - x^2 + \cos 2(2 - x^2)$
 d) $1 - 2 \sin 2(2 - x^2)$ e) $-2x(1 - 2 \sin 2(2 - x^2))$
9. a) $\frac{dy}{dx} = \frac{3x^2 - y}{x - 3y^2}$ b) $\frac{dy}{dx} = (y - 4x\sqrt{xy})/(4y\sqrt{xy} - x)$
11. $y = 2$
12. a) (i) $b = 0$ (ii) $a = 1, b = 0$
 b) (i) $b = 1$ (ii) $a = 2, b = 1$.
13. $a = 1, b = 0$
14. a) $f(8.01) \approx f(8) = 2$
 b) i) $y = (x - 8)/12 + 2$
 ii) $f(8.01) \approx (8.01 - 8)/12 + 2 = 2 + \frac{1}{1200}$
 c) The approximation in (b) is much better.
15. $\sqrt{3}ar/2$
16. $7/8 \text{ m}^2/\text{s}$

17. a) $\frac{1}{8\pi}$ cm/s
 b) $\frac{32000\pi}{81}$ cm³
18. a) $\frac{dh}{dt} = \frac{2}{125\pi}$ mm/s when $h = 50$ mm.

Chapter 5

1. $\sqrt{\frac{7}{3}}$ b) $\frac{1}{2}$
5. c) 0
7. a) By the Mean Value Theorem, for some c with $16 < c < 17$, $\sqrt{17} - \sqrt{16} = \frac{1}{2\sqrt{c}} < \frac{1}{2\sqrt{16}} = 0.125$.
 b) 0.008
 c) 2×10^{-6} .
8. -1 , 1 and 4 are stationary points; 4 is a local minimum point; -1 is a local maximum point.
9. No
10. a) $11, -61$ b) $3, -253$ c) $27/256, -750$
 d) $250, -54$ e) $2, 0$
11. $(12/13, 18/13)$
12. $p'_n(x) = p_{n-1}(x)$, and if $p_{n-1}(x) = 0$ then $p_n(x) = x^n/n!$. These hints are all you need!
13. a) $(400)/(4 + \pi)$, $100\pi/(4 + \pi)$ b) $0, 100$
14. The greatest distance is $a + 2$; the least distance is $\begin{cases} \sqrt{1 - a^2/3} & \text{if } 0 \leq a \leq 3/2 \\ |a - 2| & \text{if } a > 3/2. \end{cases}$
15. $a = \frac{\pi}{2}, x = \frac{3\pi}{4}, \frac{7\pi}{4}$; $a = \frac{3\pi}{2}, x = \frac{\pi}{4}, \frac{5\pi}{4}$. The Maple commands
`with(plots):`
`animate(plot, [cos(a) + 2*cos(2*x) + cos(4*x-a)], x=0..2*Pi, a=0..2*Pi);`
 should confirm your answers.
17. Three real zeros
18. a) $f(t) = -\cos t + t^2/2 + 3$ b) No
19. a) 0 b) $8/3$
20. a) $\frac{1}{3}$ b) $\frac{m}{n}$ c) -1 d) $-\frac{1}{2}$ e) $\frac{1}{4}$ f) $\frac{1}{3}$
21. a) $\rightarrow 0$ b) $\rightarrow \infty$ c) $\rightarrow 0$
 d) $\rightarrow 1$ e) $\rightarrow 1$ f) $\rightarrow \frac{3}{2}$

22. Combine the two fractions and apply l'Hôpital twice only. You will need to simplify the quotient obtained after the first application of l'Hôpital. Maple can confirm your answer.
23. $(a, b) = (-\sqrt{2}, \sqrt{2})$ or $(\sqrt{2}, -\sqrt{2})$
26. a) $-1/2$
b) $a = -1/2, b = 1$
27. c) $a = b = 0$


Chapter 6

2. a) $f^{-1}(x) = \frac{1}{3}(x - 1)$
b) $g^{-1}(x) = -\sqrt{x - 1}$, $\text{Dom}(g^{-1}) = [1, \infty)$,
 $\text{Range}(g^{-1}) = (-\infty, 0]$, $(g^{-1})'(x) = \frac{-1}{2\sqrt{x-1}}$
4. b) $1/3$
5. b) The restriction of f to $(-\infty, -1]$ has an inverse with domain $(-\infty, 3]$, the restriction of f to $[-1, 1]$ has an inverse with domain $[-1, 3]$, and the restriction of f to $[1, \infty)$ has an inverse with domain $[-1, \infty)$.
6. a) No b) Yes
7. a) The graph is symmetric about $x = -\frac{1}{2}$, which surely gives a local maximum of $f(x)$. There will be four (maximal) intervals where f will have an inverse. Try this exercise on Maple. The commands `plot`, `diff` and `solve` should suffice.
b) f is one-to-one; $f^{-1}(x) = x^{1/17} - 1$ is not differentiable when $x = 0$.
c) I can be one of four intervals.
8. a) $\pi/3$ b) $2/5$ c) $-\pi/3$ d) $\pi/3$
e) $4/5$ f) $3/\sqrt{34}$ g) $\pi/3$ h) $\pi - x$
11. a) $-2/\sqrt{1 - 4x^2}$ b) $1/(2\sqrt{x - x^2})$ c) $2/(4x^2 - 12x + 10)$
12. Differentiate; $-1 \leq x \leq 1$; $\pi/2$.
13. b) $f(x) = \pi/2$ when $x > 0$ and $f(x) = -\pi/2$ when $x < 0$.
14. b) The derivative of the inverse is $-1/x\sqrt{x^2 - 1}$ when $x > 1$.
16. $a = \pi/2, b = 0$
17. a) 24π km/min b) $104\pi/3$ km/min
18. $\sqrt{48}$ metres

Chapter 7

1. $[-1, 5]$, $[0, 3]$, upper half of circle.
2. a) period $2\pi/3$, odd b) period 3π , neither
 c) not periodic, even d) period $\pi/3$, odd
 e) period π , even f) 2π
3. odd, even, neither, odd, odd, even.
4. The asymptotes are
 a) $x = 3$, $y = x + 2$ b) $x = -1$, $y = x - 1$ c) $x = -3$, $x = 2$, $y = x - 1$.
6. a) $x \geq 3$, $-\frac{1}{3} < x \leq \frac{1}{3}$ b) $x = -\frac{1}{3}$, $x = -3$, $y = 1$ c) $(1, -\frac{1}{4})$, $(-1, -4)$ d) Domain:
 $x \neq 3$, $-\frac{1}{3}$, Range: $(-\infty, -4]$, $[-\frac{1}{4}, \infty)$.
7. a) $\frac{x^2}{16} + \frac{y^2}{25} = 1$, ellipse b) $\frac{x^2}{9} - \frac{y^2}{4} = 1$, hyperbola
 c) $y = x^{2/3}$ d) spiral
8. a) ii) $(2, 0)$ iii) -1
 b) ii) $(5, 0)$, $(-1, 0)$ iii) $4t^3/3$
 c) ii) $(1, 0)$, $(-1, 0)$ iii) $-\cot t$
9. a) $3x - 27y + 52 = 0$ b) $\frac{1}{9}$
10. a) $y = 3x^{\frac{2}{3}}$.
11. b) Hint: the length of one particular arc of the larger circle equals the length of one arc on the smaller circle.
 d) $x^{2/3} + y^{2/3} = 1$
12. a) $\mathbf{p}(t) = \mathbf{a} + t(\mathbf{b} - \mathbf{a})$, $\mathbf{p}(0) = \mathbf{a}$, $\mathbf{p}(1) = \mathbf{b}$ b) $y = 4 - x$, $\mathbf{q}(1/2)$ is the coordinate vector for the midpoint of B and C c) $p_0(t) = (1 - t)^2$, $p_1(t) = 2t(1 - t)$, $p_2(t) = t^2$
13. a) $(3, 0)$ b) $(-3\sqrt{3}, -3)$ c) $(\sqrt{2}, -\sqrt{2})$
14. a) $(3, \pi)$ b) $(\sqrt{2}, -3\pi/4)$ c) $(4, 2\pi/3)$
 d) $(1, \pi/2)$ e) $(4, 5\pi/6)$ f) $(4, -5\pi/6)$
15. a) Circle, centre $(0, 0)$, radius 4
 b) A ray in the second quadrant
 c) A spiral of Archimedes
16. a) Circle, centre $(0, 3)$, radius 3
 b) Circle, centre $(1, 0)$, radius 1

17. The following sketches are a guide to shape only.



19. $\frac{(x-2)^2}{9} + \frac{y^2}{5} = 1$

Chapter 8

1. a) i) $\overline{S}_{\mathcal{P}_n}(f) = \underline{S}_{\mathcal{P}_n}(f) = 1$
 ii) $\underline{S}_{\mathcal{P}_n}(f) = \frac{1}{2} \left(1 - \frac{1}{n}\right)$, $\overline{S}_{\mathcal{P}_n}(f) = \frac{1}{2} \left(1 + \frac{1}{n}\right)$
 iii) $\underline{S}_{\mathcal{P}_n}(f) = \frac{1}{6} \left(1 - \frac{1}{n}\right) \left(2 - \frac{1}{n}\right)$, $\overline{S}_{\mathcal{P}_n}(f) = \frac{1}{6} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right)$
 v) $\overline{S}_{\mathcal{P}_n}(f) = 1$, $\underline{S}_{\mathcal{P}_n}(f) = 0$
 b) i) 1 ii) $\frac{1}{2}$ iii) $\frac{1}{3}$ iv) $\frac{1}{4}$ (v) Not Riemann integrable
2. a) $\sqrt{1365} = 36.95$
 b) $\sqrt{1690.9} = 41.12$ and the lower bound is $\sqrt{1078.9} = 32.85$
4. 4.5
5. a) 82.4 b) 10
6. $f(x) = \frac{1}{x^2 + x + 1}$
7. $\frac{1}{x}$ is not differentiable on all of $[-1, 1]$ so the FTC doesn't apply.
8. a) Draw a picture! b) $5\pi/12 - \sqrt{3}/2$
10. F is continuous everywhere, but not differentiable at the integers.
12. a) $\sin x^2$ b) $3x^2 \sin x^6$ c) $-3x^2 \sin x^6$ d) $3x^2 \sin x^6 - \sin x^2$
13. $-(5 - 4x)^5$
14. biii) $\frac{\pi}{6}$.
15. a) $\frac{1}{2} e^{x^2} + C$ b) $-2 \cos \sqrt{x} + C$ c) $15/4$
 d) $4\sqrt{2} a^{9/2}/9$ e) $1/4$ f) $(2\sqrt{2} - 1)/3$
16. a) $2\sqrt{x} - 2 \ln(1 + \sqrt{x}) + C$ b) $\frac{1}{25} \left(\frac{1}{21} (5x - 1)^{21} + \frac{1}{20} (5x - 1)^{20} \right) + C$
 c) $x/(x+1)^2 + C$ d) $4 - 10 \ln(7/5)$

17. $\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{t^2 - 1}{\sqrt{2}t} \right)$ for $t \neq 0$
18. a) $\frac{4e^5 + 1}{25}$ b) $x^2 \sin x + 2x \cos x - 2 \sin x + C$
 c) $x(\ln(x) - 1) + C$ d) $\frac{\pi}{12} + \frac{\sqrt{3}}{2} - 1$
 e) $\frac{7e^8 + 1}{64}$ f) $\frac{\pi}{2}$
 g) $\frac{e^x}{2}(\cos x + \sin x)$ h) $x \tan^{-1} x - \ln \sqrt{1 + x^2} + C$
 i) $\frac{\sqrt{2}}{2} + \frac{1}{2} \ln(1 + \sqrt{2})$
19. a) $1/5$ b) diverges c) $\pi/4$
 d) 0 e) 2 f) diverges
21. a) 0 b) $\ln 2$ c) No
22. a) convergent b) divergent c) divergent
23. a) convergent b) divergent c) convergent
24. a) convergent b) divergent c) convergent
25. $s < 0$
26. $p > 1$
27. The integral converges whenever $2a - b > 1$.
28. a) $4, 8, 2$
 c) $\text{Li}'(x) = \frac{1}{\ln x} > 0$ so Li is an increasing function; $\text{Li}(2) = 0$.
 d) $\text{Li}(10^6) \geq \frac{10^6 - 2}{6 \ln 10}$.
 e) $\frac{\pi(10^6)}{x} \gtrapprox 0.07238$.
29. a) $\frac{2}{\sqrt{\pi}} e^{-x^2}$
 d) (i) $0.749 < \text{erf}(1) < 0.928$ (iii) $1/e$ (iv) 1.344

Chapter 9

2. a) A partition into 7 equal parts will suffice
3. a) $3x^2/2(x^3 + 1)$ b) e^x for $x > 0$, $-e^{-x}$ for $x < 0$
 c) $\frac{1}{(\ln(\ln x))(\ln x)x}$ d) $5x^4$ (where $x > -6^{1/5}$)
4. a) $\frac{1}{2} \ln(1 + e^{2x})$ b) $-e^{1/x}$
 c) $3^x / \ln 3$ d) $\frac{e^{\sqrt{x}}}{4}$
 e) $\frac{(\ln x)^2}{2}$ f) $\ln |\sin x|$
7. b) e

8. a) $3^x \ln 3$ b) $\left(\frac{x^3 - 3}{x^2 + 1}\right)^{1/5} \left(\frac{3x^2}{5(x^3 - 3)} - \frac{2x}{5(1 + x^2)}\right)$
 c) $(\sin x)^{\sin x} \cos x (1 + \ln(\sin x))$ d) $\cos(x^{\sin x}) x^{\sin x} \left(\cos x \ln x + \frac{\sin x}{x}\right)$
9. a) 0 b) 0 c) 1 d) e^2 e) 1
 f) 1 g) e^a h) 0 i) 0

Chapter 10

2. a) $\operatorname{sech}^2 x$ b) $-\operatorname{sech} x \tanh x$ c) $-\operatorname{cosech}^2 x$
3. a) $6x \cosh(3x^2)$ b) $\frac{-\sinh(1/x)}{x^2}$ c) $\frac{1}{2} + \frac{1}{2x^2}$
4. a) $\sinh 2x = 2 \cosh x \sinh x$; $\cosh 2x = \cosh^2 x + \sinh^2 x$
b) $\frac{1}{4}(\frac{1}{3} \cosh 3x - 3 \cosh x)$ or $\frac{1}{3} \cosh^3 x - \cosh x$
7. a) $\frac{\sinh 4x}{4}$ b) $\frac{1}{12}$ c) $(2x + \sinh 2x)/4$ d) $2 \cosh \sqrt{x}$
8. $5/4, 3, 5/12$
11. a) $2/\sqrt{1+4x^2}$ b) $\frac{1}{1-x^2}$ for $|x| > 1$ c) $\sec x$
12. a) $\frac{1}{2} \sinh^{-1} 2x$ b) $\tanh^{-1} \frac{1}{2} = \frac{1}{2} \ln 3$ c) $\sinh^{-1} \left(\frac{x+2}{3} \right)$

Index

- absolute maximum point, 57
- absolute maximum value, 57
- absolute minimum point, 57
- absolute minimum value, 57
- absolute value, 7
- answers, 243
- antiderivative, 102
- antidifferentiation, 102, 177
- area, 157
 - absolute, 168
 - signed, 168
 - under the graph of f , 164
 - unsigned, 168
- astroid, 155
- asymptote, 132
 - oblique, 132
 - vertical, 132

- Barrow, Isaac, 157
- Bernoulli, Jakob, 142, 223
- Bernoulli, Johann, 142, 223
- Bolzano, 34
- bounded function, 59
- brachistochrone, 142

- Cartesian coordinates, 144
- catenary, 223, 237
- chain rule, 70, 78
- change of variables formula, 181, 182
- circle, 20, 140
- closed interval, 3
- codomain, 9
- composition (of functions), 12
- conic sections, 20
- contains, 3
- continuity, 40, 75
 - and differentiability, 75
 - at an end point, 54
 - everywhere, 41
 - on an interval, 53, 54
- continuous, 74
- continuous functions, 21
 - and limits, 40
 - combination of, 51
 - piecewise, 166
- critical point, 96
- cubic, 13
- curve of fastest descent, 142
- cusp point, 151
- cycloid, 142

- decreasing function, 91
- definite integral, *see* integral(s), definite
- derivative(s), 65, 67
 - definition of, 67
- difference of squares identities, 229
- difference quotient, 67
- differentiability, 75
 - and continuity, 75
 - local extrema, 79
- differentiable, 74
 - at a point, 67
 - local approximation, 77
 - maximum point, 79
 - mean value theorem, 85
 - minimum point, 79
 - twice, 93
- differential calculus, 1
- differential equation, 101
- differentiation, 69
 - chain rule, 70
 - product rule, 72
 - quotient rule, 70
- discontinuity, 21
- distance, 7
- domain, 9
 - maximal, 10
 - natural, 10
- dominant term analysis, 190, 193
- double-angle formulae, 229
- dummy variable, 171

- e , 211
- elementary functions, 18
- elements (of set), 2
- ellipse, 20, 140
- endpoints
 - critical points, 96
- endpoints of intervals, 3
- equations
 - solutions, 98
- erf, 205
- error bounds, 90
- error function, 205
- even function, 132
- exp, 212
- exponential function, 214
- fastest growing term, 30
- function(s), 1, 9
 - composition of, 12, 70
 - decreasing, 91
 - error function, 205
 - even, 132
 - exponential, 213
 - implicitly defined, 19
 - increasing, 91
 - inverse function, 115
 - limit at infinity, 33
 - local approximation, 77
 - natural logarithm, 209
 - odd, 132
 - quadratic, 13
 - range of, 10
 - rational, 14
 - real-valued, 9
 - sine integral, 196
 - value of, 9
- fundamental theorem of calculus, 1, 196
 - first fundamental theorem, 172
 - second fundamental theorem, 175
- Galileo, 142, 223
- horizontal line test, 114
- horizontal point of inflexion, 93
- Huygens, Christiaan, 223
- hyperbola, 20, 140, 226
- hyperbolic functions, 227
 - double-angle formulae, 229
 - hyperbolic cosine, 224
 - hyperbolic sine, 224
 - sum and difference formulae, 229
- improper integral
 - p -integrals, 188
 - convergence of, 187
- indeterminate form(s), 31, 103, 217, 218
- inequalities, 4
 - and absolute value, 7
 - triangle inequality, 7
- inequality, 4
- infinity, symbol for, 3
- injective, 113
- integers, 2
- integral calculus, 1, 157
- integral(s), 165
 - definite
 - limits of, 165
 - improper
 - p -integrals, 186
 - comparison test, 190
 - convergence of, 186
 - limit comparison test, 193
 - indefinite, 179
 - over (a, ∞) , 187
 - Riemann sum, 162, 164
 - table of, 237
- integrand, 165
- integration, 163
 - and differentiation, 177
 - by parts, 185
 - by substitution, 180
 - constant of, 179
 - dummy variable, 171
- intermediate value theorem, 55
 - number of solutions, 100
- intervals, 3
 - closed, 3
 - open, 3
- inverse function(s), 115
 - hyperbolic functions, 231
 - inverses of, 231
 - restriction, 118
 - trigonometric functions, 120, 121
- irrational numbers, 2
- l'Hôpital's rule, 104, 217
- l'Hôpital, Guillaume, 142
- least upper bound property, 55, 57

INDEX

- Lebesgue, Henri, 157, 168
- left-hand limits, 39
- Leibniz, Gottfried, 1, 142, 157, 223
- Leonard Euler, 211
- limit(s), 1, 27
 - at infinity, 33
 - composition of function, 42
 - left-hand, 39
 - of continuous functions, 40
 - of the form $\lim_{x \rightarrow \infty} f(x)$, 27
 - pinching theorem, 37, 43
 - right-hand, 39
 - two-sided, 39
- $\ln$, 209
- logarithmic functions, 214
- logarithms, 209
- maximum
 - global, 96
 - local, 79, 94
 - second derivative test, 94
- maximum-minimum theorem, 56, 57
- mean value theorem, 85, 86
- members (of set), 2
- minimum
 - global, 96
 - local, 94
 - second derivative test, 94
- natural domain, 10
- natural logarithm, 211, 233
- natural numbers, 2
- Newton, Isaac, 1, 142, 157
- oblique asymptote, 132
- odd function, 132
- one-to-one correspondence, 112, 113
- one-to-one function, 113
- open interval, 3
- parabola(s), 20, 140
- parametric curves, 136
 - conic sections, 139
 - parameter, 138
 - parametrisation, 138
 - polar curves, 146
- partition, 160, 163
- piecewise continuous, 166
- pinching theorem, 29, 37, 43
- polar coordinates, 144
- polar curves, 146
- polynomial, 13
 - coefficients of, 13
 - degree of, 13
 - leading coefficient, 13
- polynomial division, 134
- primitive, 102
- product rule, 72
- quadratic function, 13
- quartic, 13
- quintic, 13
- radians, 14
- range of a function, 10
- rates of change, 78
- rational function, 14
- rational numbers, 2, 55
- real numbers, 2, 55
- Riemann integrable, 165
- Riemann integral, 163
 - area under the graph, 163
 - basic properties of, 169
- Riemann sum
 - upper, 162, 164
- Riemann sums, 177, 196
 - lower, 162, 164
- Riemann, Bernhard, 157
- right-hand limits, 39
- Rolle's theorem, 87
- secant, 64
- second derivative, 93, 95
- set(s), 2
- sine integral, 196
- stationary point, 80
 - classification of, 92
- subset, 3
- sum and difference formulae, 229
- tangent to a curve, 65
- triangle inequality, 7
- trigonometric functions, 14
 - inverse functions, 121
- vertical asymptotes, 132
- Weierstrass, 34
- Zeno of Elea, 34

Chapter 1

Functions of several variables

Most functions which arise in real world applications depend on more than one variable. In this chapter, we give a brief introduction to functions of two (or more) variables. Examples include functions defined by the following well known formulae:

- $A(b, h) = \frac{1}{2}bh$ (the area of a triangle);
- $D(x, y) = \sqrt{x^2 + y^2}$ (the distance of a point (x, y) from the origin);
- $\ell(x, y) = 2(x + y)$ (the perimeter of a rectangle of dimensions x and y units); and
- $F(m_1, m_2, \mathbf{x}_1, \mathbf{x}_2) = \frac{Gm_1m_2}{|\mathbf{x}_1 - \mathbf{x}_2|^2}$ (the gravitational force between two bodies of mass m_1 and m_2 positioned at points $P(\mathbf{x}_1)$ and $Q(\mathbf{x}_2)$).

Geometrically, a function of two variables represents a surface in $\mathbb{R}^3$, just as a function of one variable represents a curve in $\mathbb{R}^2$. We will introduce the *partial derivative* of a function of two variables and use this to calculate the equation of the tangent plane to a surface at a point. Other applications of the partial derivative include function and error estimation.

1.1 Sketching simple surfaces in $\mathbb{R}^3$

(Ref: SH10 §15.2, 15.3)

The graph of a function f of one variable, given by $y = f(x)$, gives rise to a curve in $\mathbb{R}^2$. The graph of a function F of two variables, given by $z = F(x, y)$, gives rise to a surface in $\mathbb{R}^3$. In this section we introduce some simple techniques for sketching surfaces given by an equation $z = F(x, y)$.


Sketching a surface in $\mathbb{R}^3$ can be challenging because the sketch must be represented in $\mathbb{R}^2$. Topographic maps solve this problem by using *contour lines* to represent the height (above sea level) of the surface of the earth at various points. We adapt this idea to sketching a surface $z = F(x, y)$ described by a function F . Here, z represents the height of the surface above the xy -plane, and the contours of the surface are defined as follows.

Definition 1.1.1. A *contour* or *level curve* of a function $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a curve in $\mathbb{R}^2$ corresponding to an equation of the form $F(x, y) = C$, where C is a constant.

For each level curve, the corresponding value of C gives the height of the curve above the xy -plane.

Example 1.1.2. Sketch level curves for the function $F : \mathbb{R}^2 \rightarrow \mathbb{R}$, where $F(x, y) = x^2 + y^2$.

Solution. The level curves of F are of the form $x^2 + y^2 = C$, where C is a *nonnegative* constant (since if $C < 0$ then there is no solution to the equation $x^2 + y^2 = C$). The level curves given by $x^2 + y^2 = 0$, $x^2 + y^2 = 1$, $x^2 + y^2 = 2$, $x^2 + y^2 = 3$ and $x^2 + y^2 = 4$ are shown below.




Level curves can be interpreted in the following way. If one walked around the circle $x^2 + y^2 = 4$, then one would remain a constant height of 4 units above the xy -plane. If one started at the origin and walked 2 units ‘east,’ then one would rise from ‘sea level’ to 4 units above sea level. $\square$


Some simple surfaces may be sketched in $\mathbb{R}^3$ by using the level curves of the surface and considering the intersection of the surface with the yz -plane. The final sketch is drawn using perspective.

Example 1.1.3. Sketch the surface in $\mathbb{R}^3$ described by the equation $z = x^2 + y^2$.

Solution. The level curves for the surface are circles (see the sketch in the previous example). When $x = 0$, we have $z = y^2$. This gives the intersection of the surface with the yz -plane. The profile of this intersection is shown below.



The yz -profile and level curves help us produce the sketch of the surface.



□


Remark 1.1.4. The previous diagram illustrates the conventional orientation for the x -, y - and z -axes.

Example 1.1.5. A surface in $\mathbb{R}^3$ is described by the equation $x^2 + y^2 - z^2 = 1$. Sketch some level curves and hence sketch the surface in $\mathbb{R}^3$.


Solution. Each level curve is obtained by setting z equal to C , for some constant C .

z	level curve
0	$x^2 + y^2 = 1$
± 1	$x^2 + y^2 = 2$
± 2	$x^2 + y^2 = 5$
± 3	$x^2 + y^2 = 10$


Each level curve is a circle. Those given in the table are sketched below.



If $x = 0$ then $y^2 - z^2 = 1$, which is a hyperbola.



Putting these two sketches together gives the following surface.



This surface is called a *hyperboloid of one sheet*. □

1.2 Partial differentiation

(Ref: SH10 §15.4, 15.6)

Suppose that f is a differentiable function of one variable. In MATH1131 we

- investigated techniques for calculating the derivative of f ,
- interpreted the derivative in terms of the rate of change of f ,
- used the derivative to calculate the equation of the tangent line to the graph of f at a point a , and
- used the tangent line to give a *linear approximation* to f near a .

Over the next few sections, we generalise some of these ideas and techniques to functions F of two variables. In particular, we introduce the notion of a *partial derivative* and then

- show how to calculate the partial derivatives of F ,
- interpret the partial derivatives in terms of the rate of change of F ,
- use the partial derivatives to calculate the equation of the tangent plane to the graph of F at a point (a, b) , and
- use the tangent plane to give a *linear approximation* for F near (a, b) .

Our discussion will assume that we have an *intuitive* idea of what is meant by a ‘tangent plane to a surface.’ A rigorous discussion of the existence of tangent planes is connected with the formal definition of the *derivative* (as opposed to *partial derivative*) of a function of two variables. This rigour is pursued in some second year courses.

To introduce the notion of a partial derivative, consider the function F given by

$$F(x, y) = x^2 + y^2.$$

The surface $z = x^2 + y^2$, corresponding to the graph of F , was sketched in Example 1.1.3. Our immediate goal is to quantify the rate of change of $F(x, y)$ at the point $(1, 2)$.

First, we find the rate of change of $F(x, y)$ at $(1, 2)$ in the y -direction. This means that we hold x constant and only allow y to vary. Since we are considering the rate of change at the point $(1, 2)$, we have $x = 1$ and

$$z = F(x, y) = F(1, y) = 1 + y^2$$

(since x is constant but y is not). Geometrically, we may interpret $z = 1 + y^2$ as the intersection of the surface $z = x^2 + y^2$ with the plane $x = 1$. This intersection is shown in Figure 1.1. The rate of change of $F(x, y)$ at $(1, 2)$ in the y -direction is equal to the gradient of the dashed tangent line shown in the figure. This gradient may be calculated in the usual way:

$$\text{gradient} = \frac{d}{dy}(1 + y^2)\Big|_{y=2} = 2y\Big|_{y=2} = 4.$$

Therefore the rate of change of $F(x, y)$ at $(1, 2)$ in the y -direction is 4.

A fast way of finding this rate of change is the following. First, differentiate F with respect to y , treating x as a constant, to obtain

$$F_y(x, y) = 2y.$$

Then the rate of change of $F(x, y)$ at $(1, 2)$ in the y -direction is

$$F_y(1, 2) = 2 \times 2 = 4.$$

The function F_y is called the *partial derivative of F with respect to y* .

In a similar manner, one may find the rate of change of $F(x, y)$ at $(1, 2)$ in the x -direction. First, differentiate F with respect to x , treating y as a constant, to obtain

$$F_x(x, y) = 2x.$$

Then the rate of change of $F(x, y)$ at $(1, 2)$ in the x -direction is

$$F_x(1, 2) = 2 \times 1 = 2.$$

The function F_x is called the *partial derivative of F with respect to x* .

The partial derivatives of a function F may be defined formally by using limits.




Figure 1.1: The intersection of the surface $z = x^2 + y^2$ with the plane $x = 1$. The dashed line is the tangent line to the cross-section at the point $(1, 2, 5)$.

Definition 1.2.1. Suppose that F is a function of two variables x and y . The partial derivatives of F with respect to x and y are defined by

$$F_x(x, y) = \lim_{h \rightarrow 0} \frac{F(x + h, y) - F(x, y)}{h}$$

and

$$F_y(x, y) = \lim_{h \rightarrow 0} \frac{F(x, y + h) - F(x, y)}{h},$$

wherever these limits exist.

Remark 1.2.2. If F is a function of x and y , then $F_x(x, y)$ is calculated by treating y as a constant and differentiating F with respect to x . On the other hand, $F_y(x, y)$ is calculated by treating x as a constant and differentiating F with respect to y .

Remark 1.2.3. As illustrated at the beginning of this section, $F_y(a, b)$ gives the rate of change of F at the point (a, b) in the y -direction. Interpreted geometrically, the number $F_y(a, b)$ is the gradient of the tangent to the cross-section at (a, b) when the surface $z = F(x, y)$ is intersected with the plane $x = a$. Similarly, $F_x(a, b)$ gives the rate of change of F at the point (a, b) in the x -direction.

Example 1.2.4. Suppose that $F(x, y) = x^2y + 2y + 4$. Find F_x and F_y .

Solution. To calculate F_x , we treat y as a constant and differentiate F with respect to x :

$$F_x(x, y) = 2xy.$$

To calculate F_y , we treat x as a constant and differentiate F with respect to y :

$$F_y(x, y) = x^2 + 2.$$

□

Partial derivatives may be denoted in a variety of ways.

Remark 1.2.5 (Notation). Suppose that a function F has partial derivatives F_x and F_y . Then F_x may be denoted by

$$\frac{\partial F}{\partial x} \quad \text{or} \quad D_1F,$$

while F_y may be denoted by

$$\frac{\partial F}{\partial y} \quad \text{or} \quad D_2F.$$

The notation involving the ‘curly d’ is used most frequently. However, the notation D_1F and D_2F is less ambiguous. (For example, to calculate $D_1F(y, x)$, we differentiate F with respect to its first variable and then evaluate this partial derivative at the point (y, x) . In ‘curly d’ notation, one would write $\frac{\partial F}{\partial x}(y, x)$. Here, the first x represents the first variable of the function while the x in parentheses represents the second ordinate of the point (y, x) .)

Example 1.2.6. Suppose that

$$F(x, y) = 3e^{xy^3} \sin y.$$

Find $\frac{\partial F}{\partial x}$ and $\frac{\partial F}{\partial y}$.

Solution. To calculate $\frac{\partial F}{\partial x}$, we treat y as a constant and differentiate F with respect to x :

$$\frac{\partial F}{\partial x} = 3y^3 e^{xy^3} \sin y.$$

To calculate $\frac{\partial F}{\partial y}$, we treat x as a constant and differentiate F with respect to y :

$$\begin{aligned} \frac{\partial F}{\partial y} &= 3 \sin(y) \frac{\partial}{\partial y} (e^{xy^3}) + 3e^{xy^3} \frac{\partial}{\partial y} (\sin y) && \text{(by using the product rule)} \\ &= 3 \sin(y) (3xy^2 e^{xy^3}) + 3e^{xy^3} (\cos y) \\ &= 3e^{xy^3} (3xy^2 \sin y + \cos y). \end{aligned}$$

□

So far we have seen examples of first order partial derivatives. Second order partial derivatives can be computed by differentiating first order partial derivatives. Some notation is given below:

$$\begin{aligned} \frac{\partial^2 F}{\partial x^2} &\text{ means } \frac{\partial}{\partial x} \left(\frac{\partial F}{\partial x} \right); \\ \frac{\partial^2 F}{\partial y^2} &\text{ means } \frac{\partial}{\partial y} \left(\frac{\partial F}{\partial y} \right); \\ \frac{\partial^2 F}{\partial x \partial y} &\text{ means } \frac{\partial}{\partial x} \left(\frac{\partial F}{\partial y} \right); \quad \text{and} \\ \frac{\partial^2 F}{\partial y \partial x} &\text{ means } \frac{\partial}{\partial y} \left(\frac{\partial F}{\partial x} \right). \end{aligned}$$

Second order partial derivatives may be denoted using other notation. For example,

$$\frac{\partial^2 F}{\partial x^2} = F_{xx} = D_1^2 F$$

while

$$\frac{\partial^2 F}{\partial y \partial x} = F_{xy} = D_2 D_1 F.$$

Example 1.2.7. Compute all second order partial derivatives of F , where $F(x, y) = 3xy^4 + e^x \sin y$.

Solution. The first order partial derivatives are given by

$$F_x(x, y) = 3y^4 + e^x \sin y \quad \text{and} \quad F_y(x, y) = 12xy^3 + e^x \cos y.$$

The second order partial derivatives are obtained by further differentiation. Thus

$$\begin{aligned} F_{xx}(x, y) &= \frac{\partial}{\partial x} (3y^4 + e^x \sin y) = e^x \sin y \\ F_{yy}(x, y) &= \frac{\partial}{\partial y} (12xy^3 + e^x \cos y) = 36xy^2 - e^x \sin y \\ F_{yx}(x, y) &= \frac{\partial}{\partial x} (12xy^3 + e^x \cos y) = 12y^3 + e^x \cos y \\ F_{xy}(x, y) &= \frac{\partial}{\partial y} (3y^4 + e^x \sin y) = 12y^3 + e^x \cos y. \end{aligned}$$

□

Note in the previous example that $F_{xy} = F_{yx}$. This is no accident.

Theorem 1.2.8 (The mixed derivative theorem). *Suppose that F is a function of two variables. If F and all its first and second order partial derivatives are continuous then*

$$\frac{\partial^2 F}{\partial x \partial y} = \frac{\partial^2 F}{\partial y \partial x}.$$

Two questions arise. First, what does it mean for a function F of two variables to be continuous? Second, why is the theorem true? We give a partial answer to each of these questions in the following remarks.

Remark 1.2.9. Suppose that F is a function of two variables. While a formal definition for continuity is not given in this course, the following rule may often be used to verify that F is continuous.

If F can be constructed by combining (via function addition, multiplication, division and composition) a finite number of continuous functions of a single variable, then F is continuous on its domain.

Thus the function F , given by

$$F(x, y) = 3xy^4 + e^x \sin y,$$

is continuous since

$$F(x, y) = f(x)g(y) + h(x)k(y),$$

where the functions given by $f(t) = 3t$, $g(t) = t^4$, $h(t) = e^t$ and $k(t) = \sin t$ are continuous. Similarly, the function G given by

$$G(x, y) = e^{3x^4y}$$

is continuous since $G(x, y) = h(g(x)f(y))$. Suffice to say, most functions of two variables given in this course are continuous on their domains.

Remark 1.2.10. To begin proving the mixed derivative theorem, one begins with the definition of the partial derivative. Now

$$\begin{aligned} F_{xy}(a, b) &= \lim_{h_2 \rightarrow 0} \frac{F_x(a, b + h_2) - F_x(a, b)}{h_2} \\ &= \lim_{h_2 \rightarrow 0} \lim_{h_1 \rightarrow 0} \frac{\frac{F(a + h_1, b + h_2) - F(a, b + h_2)}{h_1} - \frac{F(a + h_1, b) - F(a, b)}{h_1}}{h_2} \\ &= \lim_{h_2 \rightarrow 0} \lim_{h_1 \rightarrow 0} \frac{F(a + h_1, b + h_2) - F(a, b + h_2) - F(a + h_1, b) + F(a, b)}{h_1 h_2}. \end{aligned}$$

Similarly,

$$F_{yx}(a, b) = \lim_{h_1 \rightarrow 0} \lim_{h_2 \rightarrow 0} \frac{F(a + h_1, b + h_2) - F(a + h_1, b) - F(a, b + h_2) + F(a, b)}{h_1 h_2}.$$

To show that $F_{yx}(a, b) = F_{xy}(a, b)$, one need only swap the order in which the limits are taken. This step can be justified by the continuity hypothesis of the mixed derivative theorem. The details are not easy and will be omitted here.

1.3 Tangent planes to surfaces

In this section, we give an heuristic derivation of the Cartesian equation for the tangent plane to a surface at a given point. Our treatment relies on intuition rather than rigour; a full presentation requires a rigorous account of what is meant by a ‘tangent plane’ and conditions under which a tangent plane exists. Such questions are tackled in some second year courses.

As a side benefit of our derivation for equation of the tangent plane, we also obtain a formula for a *normal vector* to the surface at a given point. A normal vector is defined as follows. Suppose that a surface has a tangent plane at a point P . We say that a vector $\mathbf{n}$ is *normal* to the surface at P if $\mathbf{n}$ is normal to the tangent plane to the surface at P .

The main ideas for deriving a formula for the tangent planes and normal vectors are contained in the following example.

Example 1.3.1. Suppose that $F(x, y) = x^2 + y^2$. By using vector geometry, find the Cartesian equation of the tangent plane to the surface $z = F(x, y)$ at the point where $(x, y, z) = (1, 2, 5)$. Find also a vector $\mathbf{n}$ that is normal to the surface at this point.

Solution. We will complete the solution in four steps.

Step 1. First, intersect the surface $z = x^2 + y^2$ with the plane $x = 1$, as in Figure 1.1. This gives the cross-sectional profile

$$\begin{cases} z = 1 + y^2 \\ x = 1 \end{cases}$$

as illustrated in Figure 1.1. The dashed line passing through the point $(1, 2, 5)$ lies in the plane $x = 1$ and is tangent to the parabola $z = 1 + y^2$. Its gradient is

$$F_y(1, 2) = 4.$$

By using the point-gradient formula for a straight line, the Cartesian equation for this tangent is given by

$$z - 5 = 4(y - 2), \quad x = 1.$$

If $\lambda = y - 2$ then the equation of the tangent line in parametric vector form is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} \quad (1.1)$$

whenever $\mu \in \mathbb{R}$. Note that this line lies in the tangent plane to the surface at $(1, 2, 5)$.

Step 2. Now we intersect the surface with the plane $y = 2$ and repeat the method of Step 1. The cross-sectional profile of this intersection is given by

$$\begin{cases} z = x^2 + 4 \\ y = 2. \end{cases}$$

Consider the line that lies in the plane $y = 2$ and is tangent to $z = x^2 + 4$ at $(1, 2, 5)$. Its gradient is given by

$$F_x(1, 2) = 2$$

and hence its Cartesian equation by

$$z - 5 = 2(x - 1), \quad y = 2,$$

If $\mu = x - 1$ then we find that the parametric vector form of the tangent line is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \quad (1.2)$$

whenever $\lambda \in \mathbb{R}$. Note that this line also lies in the tangent plane to the surface at $(1, 2, 5)$.

Step 3. Since the lines given by (1.1) and (1.2) both lie in the tangent plane, and since the vectors

$$\begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$$

are nonparallel, the tangent plane to the surface at $(1, 2, 5)$ is given by

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$$

where λ and μ are arbitrary real numbers.

Step 4. We now convert the parametric vector form of the plane to point-normal form. A vector $\mathbf{n}$ that is normal to the plane is given by

$$\mathbf{n} = \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} \times \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 4 \\ -1 \end{pmatrix}.$$

Hence the point-normal form for the tangent plane is

$$\begin{pmatrix} 2 \\ 4 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} x - 1 \\ y - 2 \\ z - 5 \end{pmatrix} = 0.$$

By expanding the dot product, one obtains $z = 5 + 2(x - 1) + 4(y - 2)$, which simplifies to

$$2x + 4y - z = 5.$$

This is the cartesian form of the plane, and the vector $(2, 4, -1)^T$ is normal to the surface at the point $(1, 2, 5)$. $\square$

Note in the above example that $F_x(1, 2) = 2$ and $F_y(1, 2) = 4$. Each of these numbers appear as coefficients in the equation of the tangent plane and as components of the normal vector. By generalising the above example, one obtains the formulae in the following proposition.

Proposition 1.3.2. *Suppose that F is a function of two variables and (x_0, y_0, z_0) is a point that lies on the surface $z = F(x, y)$. If the surface has a tangent plane at the point (x_0, y_0, z_0) , then the tangent plane is given by the equation*

$$z = z_0 + F_x(x_0, y_0)(x - x_0) + F_y(x_0, y_0)(y - y_0)$$

and a normal vector to the surface at (x_0, y_0, z_0) is given by

$$\begin{pmatrix} F_x(x_0, y_0) \\ F_y(x_0, y_0) \\ -1 \end{pmatrix}.$$

Remark 1.3.3. The above proposition is similar to what is already known for functions of a single variable. In fact, we have the following. Suppose that f is a function of one variable and (x_0, y_0) is a point that lies on the curve $y = f(x)$. If f is differentiable at x_0 , then the tangent line to the curve at (x_0, y_0) is given by the equation

$$y = y_0 + f'(x_0)(x - x_0)$$

and a normal vector to the curve at (x_0, y_0) is given by

$$\begin{pmatrix} f'(x_0) \\ -1 \end{pmatrix}.$$

Remark 1.3.4. The proposition implicitly assumes that the first order partial derivatives for F exist at (x_0, y_0) . It can be shown that if a tangent plane exists then these first order partial derivatives also exist. Hence this implicit assumption causes no problems. (Be warned however, that the existence of first order partial derivatives at (x_0, y_0) does not guarantee the existence of a tangent plane at (x_0, y_0, z_0) .)

Example 1.3.5. Suppose that $F(x, y) = \sin(\pi xy^2)$. Write down the Cartesian equation for the tangent plane to the surface $z = F(x, y)$ at the point $(2, -1, 0)$ and find a vector that is normal to the surface at this point.

Solution. The partial derivatives of F are given by

$$F_x(x, y) = \pi y^2 \cos(\pi xy^2) \quad \text{and} \quad F_y(x, y) = 2\pi xy \cos(\pi xy^2).$$

Hence

$$F_x(2, -1) = \pi \quad \text{and} \quad F_y(2, -1) = -4\pi.$$

So the equation of the tangent plane is given by

$$z = 0 + \pi(x - 2) - 4\pi(y + 1) \text{ or } \pi x - 4\pi y - z = 6\pi.$$

and a normal vector is

$$\begin{pmatrix} \pi \\ -4\pi \\ -1 \end{pmatrix}.$$

□

Notes:

- One could also use the point normal form directly to find the equation of the tangent plane.

Thus the vector normal to the plane is $\mathbf{n} = \begin{pmatrix} f_x \\ f_y \\ -1 \end{pmatrix}$ (evaluated at the point P), and so the equation of the tangent plane is $(\mathbf{x} - P) \cdot \mathbf{n} = 0$.

Hence, in the previous example, we could simply expand $\left(\mathbf{x} - \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix} \right) \cdot \begin{pmatrix} \pi \\ -4\pi \\ -1 \end{pmatrix} = 0$ to find the equation of the tangent plane.

1.4. THE TOTAL DIFFERENTIAL APPROXIMATION

- More generally, if a surface is given in the form $g(x, y, z) = 0$, then the vector $\mathbf{n} = \begin{pmatrix} g_x \\ g_y \\ g_z \end{pmatrix}$ evaluated at some point P on the surface, is a vector normal to the surface at P .

Example 1.3.6. Find the equation of the tangent plane to the ellipsoid $\frac{x^2}{4} + \frac{y^2}{2} + \frac{z^2}{8} = 1$ at the point $P(0, 1, 2)$

Solution. A vector normal to the ellipsoid at P is given by

$$\mathbf{n} = \begin{pmatrix} \frac{x}{2} \\ y \\ \frac{z}{4} \end{pmatrix}_P = \begin{pmatrix} 0 \\ 1 \\ 1/2 \end{pmatrix}.$$

Hence the equation of the tangent has the form

$$0x + 1y + \frac{1}{2}z = d$$

for some constant d . Substituting in P , we find that $d = 2$ and so the desired equation is $2y + z = 4$. $\square$

1.4 The total differential approximation

Suppose that f is a differentiable function of one variable. The equation of the tangent to the graph of f at a point x_0 is given by

$$y = y_0 + f'(x_0)(x - x_0),$$

where $y_0 = f(x_0)$. When x is close to x_0 , the tangent line is close to the graph of f . In other words,

$$f(x) \approx f(x_0) + f'(x_0)(x - x_0).$$

Now writing $\Delta x = x - x_0$ and $\Delta f = f(x) - f(x_0)$ we have

$$\Delta f \approx f'(x_0)\Delta x.$$

This is called the *differential approximation* to Δf . Our goal is to generalise this idea to functions of two variables.

Suppose that a surface given by $z = F(x, y)$ has a tangent plane at the point (x_0, y_0, z_0) . The equation of the tangent is given by

$$z = z_0 + F_x(x_0, y_0)(x - x_0) + F_y(x_0, y_0)(y - y_0).$$

If (x, y) is near (x_0, y_0) then

$$F(x, y) \approx z_0 + F_x(x_0, y_0)(x - x_0) + F_y(x_0, y_0)(y - y_0).$$

Since (x_0, y_0, z_0) lies on the surface, $z_0 = F(x_0, y_0)$. Thus

$$F(x, y) - F(x_0, y_0) \approx F_x(x_0, y_0)(x - x_0) + F_y(x_0, y_0)(y - y_0). \quad (1.3)$$

Let Δx denote the difference $x - x_0$ between x and x_0 and let Δy denote the difference $y - y_0$ between y and y_0 . Then (1.3) becomes

$$F(x_0 + \Delta x, y_0 + \Delta y) - F(x_0, y_0) \approx F_x(x_0, y_0)\Delta x + F_y(x_0, y_0)\Delta y.$$

The left-hand side of this approximation represents the change in F when x_0 and y_0 are changed by Δx and Δy respectively. We call this the *increment* in F and denote it by ΔF ; that is,

$$\Delta F = F(x_0 + \Delta x, y_0 + \Delta y) - F(x_0, y_0).$$

Hence

$$\Delta F \approx F_x(x_0, y_0)(x - x_0) + F_y(x_0, y_0)(y - y_0).$$

This formula is called the *total differential approximation to ΔF* . By supressing the point of evaluation, the total differential approximation may be written as

$$\Delta F \approx \frac{\partial F}{\partial x} \Delta x + \frac{\partial F}{\partial y} \Delta y.$$

The approximation improves when Δx and Δy are smaller.

The total differential approximation can be used to estimate the change in the output of a function given changes to each of the inputs.

Example 1.4.1. The ideal gas law asserts that the pressure P , volume V and temperature T of an ideal gas are related by the formula

$$PV = kT,$$

where k is a constant. If the temperature is increased by 4% and the volume is decreased by 5%, estimate the percentage increase in pressure.

Solution. We have

$$P = \frac{kT}{V}, \quad \frac{\partial P}{\partial T} = \frac{k}{V} \quad \text{and} \quad \frac{\partial P}{\partial V} = -\frac{kT}{V^2}.$$

Now the temperature is increased by 4%, so $\Delta T = 0.04T$. Similarly, the volume is decreased by 5% and so $\Delta V = -0.05V$. By the total differential approximation,

$$\begin{aligned} \Delta P &\approx \frac{\partial P}{\partial T} \Delta T + \frac{\partial P}{\partial V} \Delta V \\ &= \frac{k}{V} \times 0.04T - \frac{kT}{V^2} \times (-0.05V) \\ &= 0.04P + 0.05P && (\text{since } P = \frac{kT}{V}) \\ &= 0.09P. \end{aligned}$$

Hence the pressure increases by approximately 9%. □

In science, engineering, psychology, economics and so on, measurements are often made that are not exact. Any quantities calculated from these measurements will also contain errors. The total differential approximation can give us an idea how bad such errors can get. Given a function F of two variables x and y , one can interpret ΔF as the error in the output given errors Δx and Δy in the inputs. Typically, one does not know the precise value of Δx and Δy , but sometimes one can

find an upper bound for the absolute errors $|\Delta x|$ and $|\Delta y|$. The total differential approximation then gives an approximate upper bound for the absolute error $|\Delta F|$ in F :

$$\begin{aligned} |\Delta F| &\approx \left| \frac{\partial F}{\partial x} \Delta x + \frac{\partial F}{\partial y} \Delta y \right| \\ &\leq \left| \frac{\partial F}{\partial x} \right| |\Delta x| + \left| \frac{\partial F}{\partial y} \right| |\Delta y|, \end{aligned}$$

(where we have used the triangle inequality in the last step). The next example illustrates these ideas.

Example 1.4.2. The dimensions of a cylinder are measured to the nearest millimeter using a measuring tape. The circumference is measured to be 22.0 cm and height is measured to be 15.0 cm. Use these measurements to (a) estimate the volume of the cylinder, and (b) estimate an upper bound for the percentage error in your answer to part (a).

Solution. (a) Let r , C and h denote the radius, circumference and height respectively. Then $C = 2\pi r$ and so

$$V = \pi r^2 h = \pi \left(\frac{C}{2\pi} \right)^2 h = \frac{C^2 h}{4\pi}.$$

By using the measurements $C = 22$ and $h = 15$, one finds that

$$V = \frac{1815}{\pi}.$$

And so the volume is estimated to be $\frac{1815}{\pi} \text{ cm}^3$, which is approximately 577.73 cm^3 .

(b) The absolute error in each measurement is at most 0.5 mm, which is 0.05 cm. Let ΔC and Δh denote the error in each measurement. Then

$$|\Delta C| \leq 0.05 \quad \text{and} \quad |\Delta h| \leq 0.05.$$

The increment ΔV is the error in our calculation for the volume. Now

$$V = \frac{C^2 h}{4\pi}, \quad \frac{\partial V}{\partial C} = \frac{Ch}{2\pi} \quad \text{and} \quad \frac{\partial V}{\partial h} = \frac{C^2}{4\pi}.$$

So the total differential approximation (when $C = 22$ and $h = 15$) is

$$\begin{aligned} \Delta V &\approx \frac{\partial V}{\partial C} \Delta C + \frac{\partial V}{\partial h} \Delta h \\ &= \frac{Ch}{2\pi} \Delta C + \frac{C^2}{4\pi} \Delta h \\ &= \frac{165}{\pi} \Delta C + \frac{121}{\pi} \Delta h \end{aligned}$$

If we take absolute values of both sides then

$$\begin{aligned} |\Delta V| &\approx \left| \frac{165}{\pi} \Delta C + \frac{121}{\pi} \Delta h \right| \\ &\leq \frac{165}{\pi} |\Delta C| + \frac{121}{\pi} |\Delta h| && \text{(by the triangle inequality)} \\ &\leq \frac{165}{\pi} \times 0.05 + \frac{121}{\pi} \times 0.05 \\ &= \frac{286}{20\pi}. \end{aligned}$$

So an upper bound for the absolute error in V is approximately $\frac{286}{20\pi}$ (that is, approximately 4.55 cm^3). An upper bound for the percentage error is given by

$$\begin{aligned}\frac{\max |\Delta V|}{V} \times 100\% &\approx \frac{286}{20\pi} \cdot \frac{\pi}{1815} \times 100\% \\ &= \frac{26}{33} \%. \end{aligned}$$

Hence the percentage error is no more than about 0.79%. $\square$

1.5 Chain rules

If f and g are functions of one variable, then the derivative of $f \circ g$ may be calculated using the chain rule for functions of one variable. In this section, we study compositions of functions of more than one variable. To calculate their partial derivatives, we use a chain rule for functions of more than one variable.

Suppose that F is a function of x and y and that x and y are each functions of t . A small change Δt in t produces a corresponding change Δx and Δy in x and y . These changes in turn produce a corresponding change ΔF in F . By the total differential approximation,

$$\Delta F \approx \frac{\partial F}{\partial x} \Delta x + \frac{\partial F}{\partial y} \Delta y,$$

and this approximation gets better as Δx and Δy approach zero. If we divide through by Δt then

$$\frac{\Delta F}{\Delta t} \approx \frac{\partial F}{\partial x} \frac{\Delta x}{\Delta t} + \frac{\partial F}{\partial y} \frac{\Delta y}{\Delta t}.$$

As $\Delta t \rightarrow 0$,

$$\frac{\Delta x}{\Delta t} = \frac{x(t + \Delta t) - x(t)}{\Delta t} \rightarrow \frac{dx}{dt}$$

and

$$\frac{\Delta y}{\Delta t} = \frac{y(t + \Delta t) - y(t)}{\Delta t} \rightarrow \frac{dy}{dt}.$$

Finally, if we view F as a function of t then, by a similar argument, $\frac{\Delta F}{\Delta t} \rightarrow \frac{dF}{dt}$. So in the limit, the total differential approximation becomes

$$\boxed{\frac{dF}{dt} = \frac{\partial F}{\partial x} \frac{dx}{dt} + \frac{\partial F}{\partial y} \frac{dy}{dt}}. \quad (1.4)$$

This is an example of a chain rule for a function of two variables.

The above chain rule must be interpreted properly. First, each derivative must be evaluated at a correct point. Second, the F appearing on the left-hand side is a function of one variable t , while the F that appears on the right-hand side is a function of two variables x and y . Technically, these are two different functions. The next theorem expresses chain rule (1.4) without these ambiguities.


Theorem 1.5.1. *Suppose that F is a function of two variables and that x and y are both functions of one variable. Define the function ϕ by $\phi(t) = F(x(t), y(t))$ and the point (x_0, y_0) by $(x_0, y_0) = (x(t_0), y(t_0))$. If x and y are both differentiable at t_0 and the partial derivatives of F exist and are continuous at (x_0, y_0) , then ϕ is differentiable at t_0 and*

$$\phi'(t_0) = D_1 F(x_0, y_0) x'(t_0) + D_2 F(x_0, y_0) y'(t_0). \quad (1.5)$$

1.5. CHAIN RULES

17

Remark 1.5.2. Formulae (1.4) and (1.5) are equivalent. The former is easier to remember while the latter is more precise. To remember the rule, consider the following *chain diagram*.



To construct the diagram, draw an arrow from each function to each of its variables. Then $\frac{dF}{dt}$ is the sum of all paths (left to right) from F to t , where the derivatives are multiplied across each path.

Example 1.5.3. The potential energy E of a particle at point (x, y) is given by $E(x, y) = \sin(\pi x^2 y)$. If the x -ordinate of the particle is increasing at a rate of 3 units per second, and the y -ordinate of the particle is decreasing at a rate of 2 units per second, find the rate of change of potential energy when the particle has coordinate $(-1, 2)$.

Solution. Since the ordinates x and y of the particle change with time t , we may view x and y as functions of t . We are told that

$$\frac{dx}{dt} = 3 \quad \text{and} \quad \frac{dy}{dt} = -2.$$

By the chain rule,

$$\begin{aligned} \frac{dE}{dt} &= \frac{\partial E}{\partial x} \frac{dx}{dt} + \frac{\partial E}{\partial y} \frac{dy}{dt} \\ &= 2\pi xy \cos(\pi x^2 y) \times 3 + \pi x^2 \cos(\pi x^2 y) \times (-2) \\ &= 2\pi x(3y - x) \cos(\pi x^2 y). \end{aligned}$$

When $(x, y) = (-1, 2)$,

$$\frac{dE}{dt} = -14\pi.$$

So the rate of change of E at $(-1, 2)$ is -14π . □

We now examine the case when F is a function of x and y , where each of x and y is a function of both s and t . This situation is sometimes written as

$$F = F(x, y), \quad x = x(s, t) \quad \text{and} \quad y = y(s, t).$$


If we treat s as a constant and differentiate F with respect to t , then chain rule (1.4) gives

$$\boxed{\frac{\partial F}{\partial t} = \frac{\partial F}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial t}.}$$

Similarly, if we treat t as a constant and differentiate F with respect to s , then chain rule (1.4) gives

$$\boxed{\frac{\partial F}{\partial s} = \frac{\partial F}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial s}.}$$

Each of these new chain rules may be remembered using the following chain diagram.



For example, to remember the rule for $\frac{\partial F}{\partial s}$, simply sum all paths (left to right) from F to s , where the derivatives are multiplied across each path.

Example 1.5.4. Suppose that $z = F(x, y)$. Express the point (x, y) in terms of polar coordinates (r, θ) . Hence express $\frac{\partial z}{\partial r}$ and $\frac{\partial z}{\partial \theta}$ in terms of x, y, F_x and F_y . Finally, show that the partial derivatives satisfy the equation

$$r \frac{\partial z}{\partial r} + \frac{\partial z}{\partial \theta} = (x - y) \frac{\partial F}{\partial x} + (x + y) \frac{\partial F}{\partial y}.$$

Solution. We have

$$z = F(x, y), \quad x = r \cos \theta, \quad y = r \sin \theta \quad \text{and} \quad r^2 = x^2 + y^2.$$

So the chain rule gives

$$\begin{aligned} \frac{\partial z}{\partial r} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial r} \\ &= F_x(x, y) \cos \theta + F_y(x, y) \sin \theta \\ &= \frac{x}{r} F_x(x, y) + \frac{y}{r} F_y(x, y) \\ &= \frac{x}{\sqrt{x^2 + y^2}} F_x(x, y) + \frac{y}{\sqrt{x^2 + y^2}} F_y(x, y) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial z}{\partial \theta} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial \theta} \\ &= -F_x(x, y) r \sin \theta + F_y(x, y) r \cos \theta \\ &= -y F_x(x, y) + x F_y(x, y). \end{aligned}$$

Finally,

$$\begin{aligned} r \frac{\partial z}{\partial r} + \frac{\partial z}{\partial \theta} &= r \left(\frac{x}{\sqrt{x^2 + y^2}} \frac{\partial F}{\partial x} + \frac{y}{\sqrt{x^2 + y^2}} \frac{\partial F}{\partial y} \right) + \left(-y \frac{\partial F}{\partial x} + x \frac{\partial F}{\partial y} \right) \\ &= x \frac{\partial F}{\partial x} + y \frac{\partial F}{\partial y} - y \frac{\partial F}{\partial x} + x \frac{\partial F}{\partial y} \\ &= (x - y) \frac{\partial F}{\partial x} + (x + y) \frac{\partial F}{\partial y}, \end{aligned}$$

as required. □

1.6. FUNCTIONS OF MORE THAN TWO VARIABLES

19

We present one more useful version of the chain rule. Suppose that F is a function of u and that u is a function of both x and y . This is sometimes written as

$$F = F(u) \quad \text{and} \quad u = u(x, y).$$


The corresponding chain rules are

$$\boxed{\frac{\partial F}{\partial x} = \frac{dF}{du} \frac{\partial u}{\partial x}}$$

and

$$\boxed{\frac{\partial F}{\partial y} = \frac{dF}{du} \frac{\partial u}{\partial y}}.$$

The chain diagram corresponding to this situation is illustrated below.



These chain rules may be easily written down after sketching the chain diagram.

1.6 Functions of more than two variables

Until now we have only discussed functions of two variables. In this section, the ideas met in this chapter are generalised to functions of three variables. We present a summary only.

Suppose that F is a function of three variables x , y and z . The partial derivatives of F are defined by

$$\begin{aligned} F_x(x, y, z) &= \lim_{h \rightarrow 0} \frac{F(x+h, y, z) - F(x, y, z)}{h} \\ F_y(x, y, z) &= \lim_{h \rightarrow 0} \frac{F(x, y+h, z) - F(x, y, z)}{h} \\ F_z(x, y, z) &= \lim_{h \rightarrow 0} \frac{F(x, y, z+h) - F(x, y, z)}{h} \end{aligned}$$

wherever these limits exist. Equivalent notation for each of these partial derivatives is given below:

$$F_x = \frac{\partial F}{\partial x} = D_1 F, \quad F_y = \frac{\partial F}{\partial y} = D_2 F, \quad \text{and} \quad F_z = \frac{\partial F}{\partial z} = D_3 F.$$

If (a, b, c) is a point in $\mathbb{R}^3$ then $F_x(a, b, c)$ is the rate of change of F in the x -direction at (a, b, c) . Similarly, $F_z(a, b, c)$ is the rate of change of F in the z -direction at (a, b, c) .

The partial derivatives of F are calculated by differentiating F with respect to one variable and treating the other variables as constants. For example, if

$$F(x, y, z) = e^{2x} z \cos y$$

then

$$F_x(x, y, z) = 2e^{2x} z \cos y, \quad F_y(x, y, z) = -e^{2x} z \sin y \quad \text{and} \quad F_z(x, y, z) = e^{2x} \cos y.$$

A ‘hyper-surface’ in $\mathbb{R}^4$ is the natural generalisation of a surface in $\mathbb{R}^3$. A function F of three variables can be used to define a hyper-surface $w = F(x, y, z)$ in $\mathbb{R}^4$. Given a point (x_0, y_0, z_0, w_0) on the surface, the equation of the ‘tangent plane’ to the surface at this point is given by


$$w = w_0 + F_x(x_0, y_0, z_0)(x - x_0) + F_y(x_0, y_0, z_0)(y - y_0) + F_z(x_0, y_0, z_0)(z - z_0)$$

(assuming, of course, that that a ‘tangent plane’ to the surface exists at this point).

The *total differential approximation* ΔF is given by

$$\Delta F \approx \frac{\partial F}{\partial x} \Delta x + \frac{\partial F}{\partial y} \Delta y + \frac{\partial F}{\partial z} \Delta z.$$

The chain rules are easily written down using chain diagrams. For example, suppose that F is a function of x , y and z and that x , y and z are each functions of both u and v . The corresponding chain diagram is shown below.



So the chain rule for $\frac{\partial F}{\partial u}$ is given by

$$\frac{\partial F}{\partial u} = \frac{\partial F}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial u} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial u},$$

while the chain rule for $\frac{\partial F}{\partial v}$ is given by

$$\frac{\partial F}{\partial v} = \frac{\partial F}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial v} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial v}.$$

The generalisation of each of these formulae to a function of four (or more) variables should be obvious.

1.7 Maple notes

The `plot3d` command is useful for visualizing the graphs of functions of several variables.

The MAPLE `diff` command carries out partial differentiation: `diff(f(x,y), x);` computes $\frac{\partial f}{\partial x}$, and `diff(f(x,y), x,y);` calculates $\frac{\partial^2 f}{\partial y \partial x}$. For example,

```
> diff(x^3*y-sin(y^2),x);
```

$$3x^2y$$

```
> diff(x^3*y-sin(y^2),y$2);
```

$$4 \sin(y^2)y^2 - 2 \cos(y^2)$$

Questions marked with [R] are routine, with [H] are harder and with [X] are for MATH1241 only. You should make sure that you can do the easier questions before you tackle the more difficult questions.

Problems 1.1 : Sketching simple surfaces in $\mathbb{R}^3$

1. [R] For each of the following surfaces, sketch some level curves and sketch the yz -profile (which is found by intersecting the surface with the plane $x = 0$). Hence sketch the surface.
 - a) $z = x^2 + y^2$
 - b) $x^2 + y^2 + z^2 = 1$
 - c) $z^2 = x^2 + y^2 - 1$
 - d) $z^2 = x^2 + y^2$
 - e) [H] $z = x^2 - y^2$

Problems 1.2 : Partial differentiation

2. [R] Given that $z = e^{x^2y}$, find $\frac{\partial z}{\partial x}$, $\frac{\partial z}{\partial y}$ and $\frac{\partial^2 z}{\partial y \partial x}$.
3. [R] In each case, find all first and second order partial derivatives and verify that $\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$.
 - a) $z = x^2y + y^2$
 - b) $z = \tan^{-1}(y/x)$
 - c) $z = \sin(x - cy)$

Problems 1.3 : Tangent planes to surfaces

4. [R] Find a normal vector $\mathbf{n}$ and the equation of the tangent plane to the surface $\mathcal{S}$ at the point $\mathbf{x}_0$.
 - a) $\mathcal{S} : z = x^2 + y^2, \quad \mathbf{x}_0 = (3, 5, 34).$
 - b) $\mathcal{S} : z = 4x^2y, \quad \mathbf{x}_0 = (2, -1, -16).$
 - c) $\mathcal{S} : z = \ln(x^2 + 3y^2), \quad \mathbf{x}_0 = (2, -1, \ln 7).$
 - d) $\mathcal{S} : z^2 + x^2 + y^2 = 1, \quad \mathbf{x}_0 = \left(\frac{1}{3}, \frac{1}{2}, \frac{\sqrt{23}}{6}\right).$

Problems 1.4 : The total differential approximation

5. [R] The volume V of a football in the shape of an ellipsoid of revolution with semi-axes of length a , b and b is given by

$$V = \frac{4}{3}\pi ab^2.$$

The values of a and b are measured to be 12.0 cm and 7.0 cm respectively, each to the nearest millimetre.

- a) Use these measurements to calculate the volume of the football.
 - b) Use the total differential to estimate the maximum absolute error in the calculated value of V .
 - c) Hence estimate the percentage error in your answer of (a).
6. [R] Suppose that $z = \frac{x+1}{y^2+1}$. The measured values of x and y are 3 and 1 respectively and each of the measurements is made with an error whose absolute value is at most 0.02. Use the total differential approximation of z to estimate the maximum error in the calculated value of z .
7. [R] Use the total differential approximation of $f(x, y) = \sqrt{x^2 + y^2}$ to estimate $\sqrt{2.98^2 + 4.03^2}$.
8. [R] The specific gravity S of a solid is given by

$$S = \frac{A}{A - W},$$

where A and W are its weights in air and water respectively.

- a) Find $\frac{\partial S}{\partial A}$ and $\frac{\partial S}{\partial W}$.
 - b) If A and W are measured to be 15.1 gm and 5.1 gm respectively, and if each of these measurements is made with an error whose absolute value is at most 0.2 gm, then use the total differential approximation of S to estimate the maximum error in the calculated value of S .
9. [R] The specific volume v of a compressible fluid flowing through a section of area A with mean velocity V is given by
- $$v = kAV$$
- where k is a constant. If v decreases by 5% and A increases by 4%, then estimate the percentage change in V .
10. [R] A triangle has two sides of length a and b with an included angle measuring $\pi/3$ radians. Given that a increases by 5%, b decreases by 6% and the included angle increases by 2%, estimate the percentage increase of area of the triangle.

Problems 1.5 : Chain rules

11. [R] Use a chain rule to calculate $\frac{dw}{dt}$ (as a function of t) when
- a) $w = xy$, $x = e^t$, $y = t^2$;
 - b) $w = x^2 + y^2 + z^2$, $x = \cos t$, $y = \sin t$, $z = t$.
12. [R] A cylindrical metallic solid is expanding under heat in such a way that its height is increasing at the rate of 0.1 cm/sec and its radius is increasing at the rate of 0.05 cm/sec. Find the rate of increase of its volume at the instant when the height is 10 cm and the radius is 5 cm.

13. [H] Suppose that f is a differentiable function of a single variable and $F(x, y)$ is defined by $F(x, y) = f(x^2 - y)$.

a) Show that F satisfies the partial differential equation

$$\frac{\partial F}{\partial x} + 2x \frac{\partial F}{\partial y} = 0.$$

b) Given that $F(0, y) = \sin y$ for all y , find a formula for $F(x, y)$.

14. [H] Consider the differential equation

$$\frac{\partial^2 u}{\partial t^2} - 16 \frac{\partial^2 u}{\partial x^2} = 0.$$

(This is an example of the one dimensional wave equation, which can be used to model, for example, the displacement $u(x, t)$ of a particle at position x along a vibrating guitar string at time t .)

- a) Suppose that g is an arbitrary twice-differentiable function of one variable and that $u(x, t) = g(x + \lambda t)$, where λ is a constant. Calculate u_{xx} and u_{tt} .
 b) Given that $u(x, t) = g(x + \lambda t)$, find all values of λ such that u satisfies the differential equation.

15. [X] A function f of two variables is said to be homogeneous of degree n if

$$f(tx, ty) = t^n f(x, y)$$

whenever $t > 0$. Show that such a function f satisfies the equation

$$x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = n f.$$

16. [X]

a) Suppose that a and b real numbers. Consider the partial differential equation

$$a \frac{\partial u}{\partial x} + b \frac{\partial u}{\partial y} = 0. \quad (1)$$

Use the chain rule to show that for a suitable choice of constants α and β , every function of the form $u(x, y) = f(\alpha x + \beta y)$, where f is differentiable, satisfies (1).

b) Generalise this remark to find solutions $u : \mathbb{R}^n \rightarrow \mathbb{R}$ to the differential equation

$$a_1 \frac{\partial u}{\partial x_1} + a_2 \frac{\partial u}{\partial x_2} + \cdots + a_n \frac{\partial u}{\partial x_n} = 0.$$

17. [X] A function u of two variables is defined implicitly by

$$u(x, t) = f(x - t u(x, t)),$$

where f is a given bounded, differentiable function of one variable, $f : \mathbb{R} \rightarrow \mathbb{R}$.

- a) Calculate $\frac{\partial u}{\partial t}$ and $\frac{\partial u}{\partial x}$.
- b) Show that $\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0$.
- c) Given that $f(s) = 1 - \tanh s$, find the smallest positive number t_m such that $\frac{\partial u}{\partial x}$ is undefined for precisely one value of x .
- d) Sketch $u(x, t)$ as a function of x for several fixed values of t (taking t to be positive), and hence interpret your result in (c). (Maple would be useful for the plots.)
- e) For fixed $t > t_m$ is $u(x, t)$ a function of x ?
- f) Generalise these results and find a way to predict t_m for arbitrary differentiable, bounded, monotonic decreasing functions f .

18. [H] A point sits on the hyperboloid

$$F(x, y, z) = x^2 + y^2 - z^2 = 1.$$

The position of the point moves with respect to time as $x(t) = z(t) = t$. The parameterisation of y is unspecified aside from $y(t) \neq 0$.

- a) Find the parameterisation of the y co-ordinate and then find $\frac{dy}{dt}$.
- b) Calculate a normal to the hyperboloid at the point (x, y, z) by calculating the vector

$$\nabla F = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}, \frac{\partial F}{\partial z} \right)^T.$$

- c) The chain rule states that

$$\frac{dF}{dt} = (\nabla F) \cdot \left(\frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right)^T.$$

Deduce that $(\nabla F) \cdot \left(\frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right)^T = 0$ and interpret this equation geometrically. Use this equation to find $\frac{dy}{dt}$ (without first finding y as a function of t as in part a).

Chapter 2

Integration techniques

Many real world problems, such as

- calculating the area of a region
- locating the centre of a region,
- calculating the volume, surface area and centre of mass of a solid,
- calculating the length of a curve,
- determining the probability that a certain event occurs,
- analysing the harmonics of a musical instrument,
- determining the solution to models of various physical phenomena, and
- calculating the work done by a force,

boil down to evaluating an appropriate integral. Some of these applications were explored in last semester's course while others will be discussed later in this course or in second year. But what these applications demand is mastery of integration. In this chapter we work towards that goal by examining techniques for integrating various types of integrals that arise when solving real world problems. The hard work done here will pay off when applications of such integrals are studied.

2.1 Trigonometric integrals

(Ref: SH10 §8.3)

In this section we focus specifically on integrals involving the trigonometric functions.

2.1.1 Integrating powers of sine and cosine

The first class of trigonometric integrals considered consist of integrals of the form

$$\int \cos^m x \sin^n x \, dx, \tag{2.1}$$

where m and n are non-negative integers. There are essentially two cases: (i) either m or n (or both) are odd; or (ii) both m and n are even. We'll begin with the first case.

Case (i). Suppose that m is odd in (2.1). Then we use the substitution $u = \sin x$ along with the identity

$$\sin^2 x + \cos^2 x = 1$$

to evaluate the integral.

Example 2.1.1. Evaluate the integral $\int \cos^3 x \sin^4 x \, dx$.

Solution. The substitution

$$u = \sin x, \quad du = \cos x \, dx$$

yields

$$\begin{aligned} \int \cos^3 x \sin^4 x \, dx &= \int \cos^2 x \sin^4 x \cos x \, dx \\ &= \int (1 - \sin^2 x) \sin^4 x \cos x \, dx \\ &= \int (1 - u^2) u^4 \, du \\ &= \int u^4 - u^6 \, du \\ &= \frac{u^5}{5} - \frac{u^7}{7} + C \\ &= \frac{\sin^5 x}{5} - \frac{\sin^7 x}{7} + C. \end{aligned}$$

□

If n is odd in (2.1) then we use the substitution $u = \cos x$ and follow the same strategy. (If both m and n are odd, then either of the substitutions $u = \sin x$ or $u = \cos x$ will work.)

Example 2.1.2. Evaluate the integral $\int \cos^6 x \sin^5 x \, dx$.

Solution. This time we use the substitution

$$u = \cos x, \quad du = -\sin x \, dx$$

to evaluate the integral:

$$\begin{aligned} \int \cos^6 x \sin^5 x \, dx &= - \int \cos^6 x (\sin^2 x)^2 (-\sin x) \, dx \\ &= - \int \cos^6 x (1 - \cos^2 x)^2 (-\sin x) \, dx \\ &= - \int u^6 (1 - u^2)^2 \, du \\ &= - \int u^6 - 2u^8 + u^{10} \, du \\ &= - \left(\frac{u^7}{7} - \frac{2u^9}{9} + \frac{u^{11}}{11} \right) + C \\ &= -\frac{\cos^7 x}{7} + \frac{2\cos^9 x}{9} - \frac{\cos^{11} x}{11} + C. \end{aligned}$$

□

Case (ii). The case where *both* m and n are even in (2.1) requires an entirely different approach. This time we use the identities

$$\cos^2 x = \frac{1 + \cos 2x}{2} \quad \text{and} \quad \sin^2 x = \frac{1 - \cos 2x}{2} \quad (2.2)$$

to change integral (2.1) into a sum of integrals of the form

$$\int \cos^k(2x) dx.$$

We then repeat the methods of Case (i) or Case (ii) until each integral in the sum is easy to compute.

Example 2.1.3. Evaluate $\int \sin^2 x dx$.

Solution. The second identity in (2.2) gives

$$\begin{aligned} \int \sin^2 x dx &= \frac{1}{2} \int 1 - \cos 2x dx \\ &= \frac{1}{2} \left(x - \frac{\sin 2x}{2} \right) + C. \end{aligned}$$

□

The next example is much harder.

Example 2.1.4. Evaluate $\int \sin^2 x \cos^4 x dx$.

Solution. The identities (2.2) give

$$\begin{aligned} \int \sin^2 x \cos^4 x dx &= \int \left(\frac{1 - \cos 2x}{2} \right) \left(\frac{1 + \cos 2x}{2} \right)^2 dx \\ &= \frac{1}{8} \int (1 - \cos 2x)(1 + \cos 2x)^2 dx \\ &= \frac{1}{8} \int 1 + \cos 2x - \cos^2 2x - \cos^3 2x dx \\ &= \frac{x}{8} + \frac{\sin 2x}{16} - \frac{1}{8} \int \cos^2 2x dx - \frac{1}{8} \int \cos^3 2x dx. \end{aligned} \quad (2.3)$$

The first integrand of (2.3) is an even power of $\cos 2x$ and is evaluated using the first identity in (2.2):

$$\begin{aligned} \int \cos^2 2x dx &= \frac{1}{2} \int 1 + \cos 4x dx \\ &= \frac{x}{2} + \frac{\sin 4x}{8} + C_1. \end{aligned}$$

The second integrand of (2.3) is an odd power of $\cos 2x$. The substitution

$$u = \sin 2x, \quad du = 2 \cos 2x dx$$

$$\begin{aligned}
\int \cos^3 2x \, dx &= \frac{1}{2} \int (1 - \sin^2 2x) 2 \cos 2x \, dx \\
&= \frac{1}{2} \int 1 - u^2 \, du \\
&= \frac{u}{2} - \frac{u^3}{6} + C_2 \\
&= \frac{\sin 2x}{2} - \frac{\sin^3 2x}{6} + C_2.
\end{aligned}$$

Following from (2.3) we obtain

$$\begin{aligned}
\int \sin^2 x \cos^4 x \, dx &= \frac{x}{8} + \frac{\sin 2x}{16} - \frac{x}{16} - \frac{\sin 4x}{64} - \frac{\sin 2x}{16} + \frac{\sin^3 2x}{48} + C \\
&= \frac{x}{16} - \frac{\sin 4x}{64} + \frac{\sin^3 2x}{48} + C.
\end{aligned}$$

□

2.1.2 Integrating multiple angles of sine and cosine

The next class of trigonometric integrals consists of integrals of the form

$$\int \cos mx \sin nx \, dx, \quad \int \cos mx \cos nx \, dx \quad \text{or} \quad \int \sin mx \sin nx \, dx, \quad (2.4)$$

where m and n are real numbers. These have many applications, including analysis of waves, musical harmonics and distribution of heat in solids. Such applications are discussed in some second year courses.

To evaluate the integrals in (2.4) we need the following trigonometric identities.

Lemma 2.1.5. *Suppose that A and B are real numbers. Then*

$$\sin A \cos B = \frac{1}{2} (\sin(A + B) + \sin(A - B)) \quad (2.5)$$

$$\cos A \cos B = \frac{1}{2} (\cos(A - B) + \cos(A + B)) \quad (2.6)$$

$$\sin A \sin B = \frac{1}{2} (\cos(A - B) - \cos(A + B)) \quad (2.7)$$

Proof. We only prove the first identity; the other proofs are similar. We begin with the sum and difference formulae

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B.$$

Adding the two identities gives

$$\sin(A + B) + \sin(A - B) = 2 \sin A \cos B,$$

whereupon dividing by 2 establishes (2.5). □

Example 2.1.6. Evaluate $\int \cos 5x \cos 3x \, dx$.

$$\begin{aligned}\int \cos 5x \cos 3x \, dx &= \frac{1}{2} \int \cos(5x - 3x) + \cos(5x + 3x) \, dx \\ &= \frac{1}{2} \int \cos(2x) + \cos(8x) \, dx \\ &= \frac{\sin 2x}{4} + \frac{\sin 8x}{16} + C.\end{aligned}$$

□

2.1.3 Integrating powers of tan and sec

Students will not be expected to evaluate difficult integrals involving powers of tan and sec. However, it is expected that they will be able to use the facts that

$$\tan^2 x + 1 = \sec^2 x, \quad \frac{d}{dx} \tan x = \sec^2 x \quad \text{and} \quad \frac{d}{dx} \sec x = \tan x \sec x$$

to find suitable substitutions or strategies.

Example 2.1.7. Evaluate $\int \tan^2 x \, dx$.

Proof. The Pythagorean identity gives

$$\begin{aligned}\int \tan^2 x \, dx &= \int \sec^2 x - 1 \, dx \\ &= \tan x - x + C.\end{aligned}$$

□

Example 2.1.8. Evaluate $\int \sec^4 x \tan x \, dx$.

Proof. The substitution

$$u = \sec x, \quad du = \sec x \tan x \, dx$$

yields

$$\begin{aligned}\int \sec^4 x \tan x \, dx &= \int u^3 \, du \\ &= \frac{\sec^4 x}{4} + C.\end{aligned}$$

(Of course, one can always by-pass the substitution and integrate by inspection.)

□

2.2 Reduction formulae

We begin with an example.

Example 2.2.1. Suppose that I_n is defined by

$$I_n = \int_0^{\pi/4} \tan^n x \, dx$$

whenever $n \geq 0$. Show that

$$I_n = \frac{1}{n-1} - I_{n-2} \quad \forall n \geq 2. \quad (2.8)$$

Hence evaluate

$$\int_0^{\pi/4} \tan^6 x \, dx.$$

Proof. By the identity $\tan^2 x = \sec^2 x - 1$,

$$\begin{aligned} I_n &= \int_0^{\pi/4} \tan^{n-2} x \tan^2 x \, dx \\ &= \int_0^{\pi/4} \tan^{n-2} x (\sec^2 x - 1) \, dx \\ &= \int_0^{\pi/4} \tan^{n-2} x \sec^2 x \, dx - \int_0^{\pi/4} \tan^{n-2} x \, dx \\ &= \left[\frac{\tan^{n-1} x}{n-1} \right]_0^{\pi/4} - I_{n-2} \\ &= \frac{1}{n-1} - I_{n-2} \end{aligned}$$

as required.

Using (2.8), we see that

$$\begin{aligned} \int_0^{\pi/2} \tan^6 x \, dx &= I_6 \\ &= \frac{1}{5} - I_4 \\ &= \frac{1}{5} - \left(\frac{1}{3} - I_2 \right) \\ &= \frac{1}{5} - \frac{1}{3} + \left(\frac{1}{1} - I_0 \right) \\ &= \frac{1}{5} - \frac{1}{3} + \frac{1}{1} - \int_0^{\pi/4} dx \\ &= \frac{1}{5} - \frac{1}{3} + \frac{1}{1} - \frac{\pi}{4} \\ &= \frac{13}{15} - \frac{\pi}{4}. \end{aligned}$$

(Note that I_0 must be evaluated directly, since formula (2.8) is only valid when $n \geq 2$.) □

Formula (2.8) is an example of a *reduction formula*, since it expresses an integral in terms of a ‘smaller’ integral of the same type. As illustrated above, once a reduction formula is known, integrals of that type may be evaluated rapidly. Although not the case with the previous example, most reduction formulae are proved using integration by parts.

Example 2.2.2. Suppose that

$$I_n = \int \sin^n x \, dx$$

whenever $n \geq 0$. Show that

$$I_n = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} I_{n-2} \quad \forall n \geq 2.$$

Solution. If we apply integration by parts with

$$\begin{aligned} u &= \sin^{n-1} x & v &= -\cos x \\ u' &= (n-1) \sin^{n-2} x \cos x & v' &= \sin x \end{aligned}$$

then

$$\begin{aligned} I_n &= \int \sin^{n-1} x \sin x \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos x \cos x \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x (1 - \sin^2 x) \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \, dx - (n-1) \int \sin^n x \, dx \\ &= -\sin^{n-1} x \cos x + (n-1) I_{n-2} - (n-1) I_n. \end{aligned}$$

If we gather the I_n terms to the left-hand side then

$$n I_n = -\sin^{n-1} x \cos x + (n-1) I_{n-2}.$$

Dividing both sides by n gives the result. □

In the final example, the reduction formula has two parameters (m and n) instead of one.

Example 2.2.3. Suppose that

$$I_{m,n} = \int_0^{\pi/2} \cos^m x \sin^n x \, dx \tag{2.9}$$

whenever m and n are nonnegative integers.

(a) [X] Show that

$$I_{m,n} = \begin{cases} \left(\frac{m-1}{m+n} \right) I_{m-2,n} & \text{provided that } m \geq 2 \\ \left(\frac{n-1}{m+n} \right) I_{m,n-2} & \text{provided that } n \geq 2. \end{cases} \tag{2.10}$$

(b) [R] Using the result of (a), evaluate $\int_0^{\pi/2} \cos^4 x \sin^6 x \, dx$.

Solution. (a) Integration by parts with

$$\begin{aligned} u &= \cos^{m-1} x & v &= \frac{\sin^{n+1} x}{n+1} \\ u' &= -(m-1) \cos^{m-2} x \sin x & v' &= \sin^n x \cos x \end{aligned}$$

gives

$$\begin{aligned} I_{m,n} &= \left[\frac{\cos^{m-1} x \sin^{n+1} x}{n+1} \right]_0^{\pi/2} + \frac{m-1}{n+1} \int_0^{\pi/2} \cos^{m-2} x \sin^{n+2} x \, dx \\ &= \frac{m-1}{n+1} \int_0^{\pi/2} \sin^n x \cos^{m-2} x (1 - \cos^2 x) \, dx \\ &= \frac{m-1}{n+1} I_{m-2,n} - \frac{m-1}{n+1} I_{m,n}. \end{aligned}$$

By bringing the $I_{m,n}$ terms to the left-hand side and rearranging, the first formula is obtained. The second formula is proved similarly.

(b) The first formula in (2.10) allows us to reduce the first parameter:

$$\begin{aligned} \int_0^{\pi/2} \cos^4 x \sin^6 x \, dx &= I_{4,6} \\ &= \frac{3}{10} I_{2,6} \\ &= \frac{3}{10} \cdot \frac{1}{8} I_{0,6}. \end{aligned}$$

The second formula in (2.10) allows us to reduce the second parameter:

$$\begin{aligned} \int_0^{\pi/2} \cos^4 x \sin^6 x \, dx &= \frac{3}{10} \cdot \frac{1}{8} I_{0,6} \\ &= \frac{3}{10} \cdot \frac{1}{8} \cdot \frac{5}{6} I_{0,4} \\ &= \frac{3}{10} \cdot \frac{1}{8} \cdot \frac{5}{6} \cdot \frac{3}{4} I_{0,2} \\ &= \frac{3}{10} \cdot \frac{1}{8} \cdot \frac{5}{6} \cdot \frac{3}{4} \cdot \frac{1}{2} I_{0,0}. \end{aligned}$$

Finally, we note from (2.9) that $I_{0,0} = \frac{\pi}{2}$. Hence

$$\int_0^{\pi/2} \cos^4 x \sin^6 x \, dx = \frac{3}{10} \cdot \frac{1}{8} \cdot \frac{5}{6} \cdot \frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \frac{3\pi}{512},$$

completing the problem. □

2.2.1 [X] Application: the irrationality of π

In this subsection, we use a reduction formula to prove that π is an irrational number. Students studying MATH1231 may want to skip to the next section if this does not interest them.

Although the number π has been studied for over 2000 years, it was only in 1770 that it was shown (by Johann Heinrich Lambert) that π is an irrational number. The proof we give is simpler than Lambert's proof and is similar to a proof discovered in the twentieth century. The main

2.2. REDUCTION FORMULAE

33

idea (as with many other irrationality proofs) is to assume that π is a rational number and find a contradiction. In our proof, a contradiction arises by showing that a certain definite integral, which is known to lie in the interval $(0, 1)$, is an integer if one assumes that π is a rational number.

Suppose that q is a positive integer. For each natural number n , define the integral I_n by

$$I_n = \frac{q^{2n}}{n!} \int_{-\pi/2}^{\pi/2} \left(\frac{\pi^2}{4} - x^2 \right)^n \cos x \, dx. \quad (2.11)$$

Lemma 2.2.4. *Suppose that q is a positive integer and I_n is defined as above. If $n \geq 2$ then*

$$I_n = (4n - 2)q^2 I_{n-1} - q^4 \pi^2 I_{n-2}. \quad (2.12)$$

Moreover,

$$I_1 = 4q^2 \quad \text{and} \quad I_0 = 2.$$

Proof. We give an outline proof only; students should be able to fill in the details. Suppose that $n \geq 2$. Integration by parts with

$$\begin{aligned} u &= \left(\frac{\pi^2}{4} - x^2 \right)^n & v &= \sin x \\ u' &= -2nx \left(\frac{\pi^2}{4} - x^2 \right)^{n-1} & v' &= \cos x \end{aligned}$$

yields

$$I_n = \frac{2nq^{2n}}{n!} \int_{-\pi/2}^{\pi/2} x \left(\frac{\pi^2}{4} - x^2 \right)^{n-1} \sin x \, dx.$$

A second application of integration by parts with

$$u = x \left(\frac{\pi^2}{4} - x^2 \right)^{n-1} \quad v' = \sin x$$

gives

$$\begin{aligned} I_n &= \frac{2q^{2n}}{(n-1)!} \int_{-\pi/2}^{\pi/2} \left(\frac{\pi^2}{4} - x^2 \right)^{n-1} \cos x \, dx - \frac{4q^{2n}}{(n-2)!} \int_{-\pi/2}^{\pi/2} x^2 \left(\frac{\pi^2}{4} - x^2 \right)^{n-2} \cos x \, dx \\ &= 2q^2 I_{n-1} - \frac{4q^{2n}}{(n-2)!} \int_{-\pi/2}^{\pi/2} x^2 \left(\frac{\pi^2}{4} - x^2 \right)^{n-2} \cos x \, dx. \end{aligned}$$

In the right-most integrand, write x^2 as $\frac{\pi^2}{4} - \left(\frac{\pi^2}{4} - x^2 \right)$. Hence

$$\begin{aligned} I_n &= 2q^2 I_{n-1} - \frac{4q^{2n}}{(n-2)!} \left(\frac{\pi^2}{4} \int_{-\pi/2}^{\pi/2} \left(\frac{\pi^2}{4} - x^2 \right)^{n-2} \cos x \, dx - \int_{-\pi/2}^{\pi/2} \left(\frac{\pi^2}{4} - x^2 \right)^{n-1} \cos x \, dx \right) \\ &= 2q^2 I_{n-1} - q^4 \pi^2 I_{n-2} + 4(n-1)q^2 I_{n-1}. \end{aligned}$$

If we gather both I_{n-1} terms then we obtain (2.12) as required. The proof that $I_1 = 4q^2$ and $I_0 = 2$ is straightforward and is left as an exercise. $\square$

The next lemma will be used to show that $0 < I_n < 1$ for sufficiently large n .

Lemma 2.2.5. *If $a > 0$ then $\lim_{n \rightarrow \infty} \frac{a^n}{n!} = 0$.*

Proof. Take N to be any integer greater than $2a$. If $n > N$ then

$$\begin{aligned} \frac{a^n}{n!} &= \frac{a^N}{N!} \cdot \frac{a^{n-N}}{(N+1)(N+2)\dots n} \\ &= \frac{a^N}{N!} \cdot \frac{a}{N+1} \cdot \frac{a}{N+2} \cdots \frac{a}{n}. \end{aligned}$$

Now $\frac{a^N}{N!}$ is some fixed number and

$$\frac{a}{N+1} < \frac{1}{2}, \quad \frac{a}{N+2} < \frac{1}{2}, \quad \dots, \quad \frac{a}{n} < \frac{1}{2},$$

so

$$0 < \frac{a^n}{n!} < \frac{a^N}{N!} \left(\frac{1}{2}\right)^{n-N}.$$

As $n \rightarrow \infty$, the right hand side approaches 0 and hence $\lim_{n \rightarrow \infty} \frac{a^n}{n!} = 0$ by a sequence version of the pinching theorem. □

Theorem 2.2.6. *The number π is irrational.*

Proof. Suppose that $\pi = \frac{p}{q}$ where p and q are positive integers and consider the integral I_n defined by (2.11) whenever $n \geq 0$.

First we argue by mathematical induction that I_n is an integer for every value of n . By Lemma 2.2.4, I_0 and I_1 are integers. Suppose inductively that I_{k-2} and I_{k-1} are integers whenever $k \geq 2$. By (2.12) and the assumption that $\pi = p/q$,

$$I_k = (4n-2)q^2 I_{k-1} - p^2 q^2 I_{k-2}$$

and so I_k is also an integer. Hence I_n is an integer whenever $n \geq 0$.

On the other hand, it is not hard to see that

$$0 < \left(\frac{\pi^2}{4} - x^2\right)^n \cos x \leq \left(\frac{\pi^2}{4}\right)^n$$

whenever $n \geq 0$ and $-\frac{\pi}{2} < x < \frac{\pi}{2}$. Hence

$$\begin{aligned} 0 < I_n &< \frac{q^{2n}}{n!} \int_{-\pi/2}^{\pi/2} \left(\frac{\pi^2}{4}\right)^n dx \\ &= \frac{q^{2n}}{n!} \left(\frac{\pi^2}{4}\right)^n \pi \\ &= \frac{p}{q} \cdot \frac{(p^2/4)^n}{n!}. \end{aligned} \tag{2.13}$$

As $n \rightarrow \infty$, the expression in (2.13) approaches 0 by Lemma 2.2.5. Hence $0 < I_n < 1$ whenever n is sufficiently large. In particular, there is a large value of n for which I_n is not an integer, giving a contradiction.

Hence we conclude that π is an irrational number. □

(Ref: SH10 §8.4)

Many integrals can be evaluated by finding the right substitution, but unfortunately there is no general systematic way to do this. Integrals involving square roots of quadratics often yield to trigonometric or hyperbolic substitutions.

The following table indicates which substitution can be tried for integrals containing an expression of the form $\sqrt{\pm x^2 \pm a^2}$.

Expression in integrand	Trigonometric substitution	Hyperbolic substitution
$\sqrt{a^2 - x^2}$	$x = a \sin \theta$	$x = a \tanh \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta$	$x = a \sinh \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta$	$x = a \cosh \theta$

Whether or not a trigonometric substitution is more efficient than a hyperbolic substitution depends on the particular integral. In general, trigonometric substitutions are favoured because once integration is completed in the variable θ , it is easier to restate the result in terms of x .

Example 2.3.1. Evaluate $\int \sqrt{1 - x^2} dx$.


Solution. The substitution

$$x = \sin \theta \qquad dx = \cos \theta d\theta$$

yields

$$\begin{aligned}
 \int \sqrt{1 - x^2} dx &= \int \sqrt{1 - \sin^2 \theta} \cos \theta d\theta \\
 &= \int \sqrt{\cos^2 \theta} \cos \theta d\theta && \text{(since } \sin^2 \theta + \cos^2 \theta = 1) \\
 &= \int \cos^2 \theta d\theta \\
 &= \frac{1}{2} \int 1 + \cos 2\theta d\theta && \text{(by the double-angle formula for cosine)} \\
 &= \frac{1}{2} \left(\theta + \frac{\sin 2\theta}{2} \right) + C \\
 &= \frac{1}{2} (\theta + \sin \theta \cos \theta) + C && \text{(by the double-angle formula for sin).}
 \end{aligned}$$

To state our answer in terms of x , it is easiest to draw a triangle.



We see that $\theta = \sin^{-1} x$ and $\cos \theta = \frac{\sqrt{1-x^2}}{1}$. Hence

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \left(\sin^{-1} x + x\sqrt{1-x^2} \right) + C.$$

□

Example 2.3.2. Evaluate $\int \frac{dx}{(4+x^2)^{3/2}}$.


Solution. By using the substitution

$$x = 2 \tan \theta \qquad dx = 2 \sec^2 \theta d\theta$$

and the identity $\tan^2 \theta + 1 = \sec^2 \theta$, we have

$$\begin{aligned} \int \frac{dx}{(4+x^2)^{3/2}} &= \int \frac{2 \sec^2 \theta d\theta}{(\sqrt{4 \tan^2 \theta + 4})^3} \\ &= \int \frac{2 \sec^2 \theta d\theta}{(2\sqrt{\tan^2 \theta + 1})^3} \\ &= \int \frac{2 \sec^2 \theta d\theta}{(2 \sec \theta)^3} \\ &= \frac{1}{4} \int \frac{d\theta}{\sec \theta} \\ &= \frac{1}{4} \int \cos \theta d\theta \\ &= \frac{\sin \theta}{4} + C. \end{aligned}$$

To write the answer in terms of x , consider the following triangle.



Thus $\sin \theta = \frac{x}{\sqrt{x^2+4}}$ and hence

$$\int \frac{dx}{(4+x^2)^{3/2}} = \frac{x}{4\sqrt{x^2+4}} + C.$$

□

Example 2.3.3. Use the substitution $x = 3 \cosh \theta$ to evaluate $\int \frac{x^3 dx}{\sqrt{x^2-9}}$.

Solution. The substitution

$$x = 3 \cosh \theta \qquad dx = 3 \sinh \theta d\theta$$

and the identity $\cosh^2 \theta - \sinh^2 \theta = 1$ gives $\sqrt{x^2 - 9} = 3 \sinh \theta$. Hence

$$\begin{aligned}
 \int \frac{x^3 dx}{\sqrt{x^2 - 9}} &= \int \frac{3^4 \cosh^3 \theta \sinh \theta d\theta}{\sqrt{9 \cosh^2 \theta - 9}} \\
 &= \int \frac{3^4 \cosh^3 \theta \sinh \theta d\theta}{3 \sinh \theta} && (\text{since } \cosh^2 \theta - \sinh^2 \theta = 1) \\
 &= 27 \int \cosh^3 \theta d\theta \\
 &= 27 \int \cosh \theta \cosh^2 \theta d\theta \\
 &= 27 \int \cosh \theta (1 + \sinh^2 \theta) d\theta && (\text{since } \cosh^2 \theta - \sinh^2 \theta = 1) \\
 &= 27 \int 1 + u^2 dx && (\text{using the substitution } u = \sinh \theta) \\
 &= 27(u + \frac{1}{3}u^3) + C.
 \end{aligned}$$

As was observed above, $\sqrt{x^2 - 9} = 3 \sinh \theta$ and hence

$$u = \sinh \theta = \frac{1}{3}\sqrt{x^2 - 9}.$$

Therefore

$$\begin{aligned}
 \int \frac{x^3 dx}{\sqrt{x^2 - 9}} &= 27(u + \frac{1}{3}u^3) + C \\
 &= 9\sqrt{x^2 - 9} + \frac{1}{3}(\sqrt{x^2 - 9})^3 + C.
 \end{aligned}$$

□

Exercise: Evaluate the integral in Example 2.3.3 by using an appropriate trigonometric substitution.

2.4 Integrating rational functions

(Ref: SH10 §8.5)

The main result of this section is that every rational function has an antiderivative among the elementary functions. Moreover, there is a systematic way of finding this antiderivative.

Before we begin, we remind the reader that a rational function f is of the form

$$f(x) = \frac{p(x)}{q(x)},$$

where p and q are polynomials. We say that f is *proper* if the **degree** of the denominator q is greater than the **degree** of the numerator p . We say that f is *improper* if the degree of the denominator q is less than or equal to the degree of the numerator p . We say that a quadratic polynomial is *irreducible* if it has no real linear factors. (Equivalently, a quadratic $ax^2 + bx + c$ is irreducible if its discriminant $b^2 - 4ac$ is negative.)

Before articulating the general strategy for integrating a rational function, we revise some known tactics for integrating simpler examples.

Example 2.4.1. Evaluate $\int \frac{x}{x^2 + 2x + 10} dx$.

Solution. The first tactic is to rewrite integrand so that the derivative of the denominator is sitting on the numerator:

$$\begin{aligned} \int \frac{x}{x^2 + 2x + 10} dx &= \frac{1}{2} \int \frac{2x}{x^2 + 2x + 10} dx \\ &= \frac{1}{2} \int \frac{(2x + 2) - 2}{x^2 + 2x + 10} dx \\ &= \frac{1}{2} \int \frac{2x + 2}{x^2 + 2x + 10} - \frac{2}{x^2 + 2x + 10} dx. \end{aligned}$$

The first term can now be integrated using the $\ln$ function. To integrate the second term, we complete the square in the denominator:

$$\begin{aligned} x^2 + 2x + 10 &= x^2 + 2x + 1 + 9 \\ &= (x + 1)^2 + 3^2. \end{aligned}$$

Hence

$$\begin{aligned} \int \frac{x}{x^2 + 2x + 10} dx &= \frac{1}{2} \int \frac{2x + 2}{x^2 + 2x + 10} dx - \int \frac{1}{(x + 1)^2 + 3^2} dx \\ &= \frac{1}{2} \ln |x^2 + 2x + 10| - \frac{1}{3} \tan^{-1} \left(\frac{x + 1}{3} \right) + C. \end{aligned}$$

□

The integrand of Example 2.4.1 is a proper rational function whose denominator is an irreducible quadratic. Any such function can be integrated using the techniques illustrated in that example. We turn now to study a general strategy for integrating *any* rational function.

2.4.1 The overall strategy

In this subsection we give an overview of the approach to integrating rational functions. The basic procedure is summarised below, afterwards illustrated with an example.

1. If the rational function is improper, then use polynomial division to write f as the sum of a polynomial and a proper rational function. Since the polynomial is easy to integrate, we need only focus on integrating a proper rational function.
2. It can be shown using algebra that every proper rational function f can be written as a unique sum of functions of the form

$$\frac{A}{(x - a)^k} \quad \text{and} \quad \frac{Bx + C}{(x^2 + bx + c)^k}, \quad (2.14)$$

where the quadratic $x^2 + bx + c$ is *irreducible*. This sum is called the *partial fractions decomposition* of f . We discuss how to find the partial fractions decomposition in the next subsection.

2.4. INTEGRATING RATIONAL FUNCTIONS

39

3. Now we only need to integrate functions of the form given by (2.14). By completing the square, using a substitution or performing simple algebraic manipulation, these can be integrated by the standard formulae

$$\begin{aligned}\int x^k dx &= \frac{x^{k+1}}{k+1} + C, & k \neq -1 \\ \int \frac{g'(x)}{g(x)} dx &= \ln |g(x)| + C \\ \int \frac{dx}{a^2 + x^2} dx &= \frac{1}{a} \tan^{-1} \frac{x}{a} + C.\end{aligned}$$

Example 2.4.2. Find $\int \frac{x^4 - 5x^3 + 12x^2 - 21x + 35}{x^3 - 3x^2 + 4x - 12} dx$.

Solution. Denote the integrand by $f(x)$.

Step 1. Note that f is improper. So polynomial division gives

$$\begin{array}{r} x^3 - 3x^2 + 4x - 12 \overline{) x^4 - 5x^3 + 12x^2 - 21x + 35} \\ \underline{x^4 - 3x^3 + 4x^2 - 12x} \\ -2x^3 + 8x^2 - 9x + 35 \\ \underline{-2x^3 + 6x^2 - 8x + 24} \\ 2x^2 - x + 11 \end{array}$$

and hence

$$f(x) = x + 2 + \frac{2x^2 - x + 11}{x^3 - 3x^2 + 4x - 12}.$$

Note that rational expression on the far right-hand side is proper.

Step 2. The partial fractions decomposition of $\frac{2x^2 - x + 11}{x^3 - 3x^2 + 4x - 12}$ is given by

$$\frac{2x^2 - x + 11}{x^3 - 3x^2 + 4x - 12} = \frac{2}{x - 3} - \frac{1}{x^2 + 4}. \quad (2.15)$$

(Note that the quadratic $x^2 + 4$ is irreducible.) It is not hard to verify that (2.15) is true; the question is, How does one find such a decomposition? We answer this question in Subsection 2.4.2.

Step 3. The results of Steps 1 and 2 give

$$f(x) = x - 2 + \frac{2}{x - 3} - \frac{1}{x^2 + 4}.$$

To integrate f , we need only integrate each term in the sum. Hence

$$\begin{aligned}\int \frac{x^4 - 5x^3 + 12x^2 - 21x + 35}{x^3 - 3x^2 + 4x - 12} dx &= \int \left(x - 2 + \frac{2}{x - 3} - \frac{1}{x^2 + 4} \right) dx \\ &= \frac{1}{2}x^2 - 2x + 2 \ln |x - 3| - \frac{1}{2} \tan^{-1} \frac{x}{2} + C,\end{aligned}$$

completing the problem. □

In the next subsection, we focus on finding the partial fractions decomposition of a proper rational function.

2.4.2 Partial fractions decompositions

To find the partial fractions decomposition of a proper rational function $\frac{p}{q}$, we factorise the denominator q as much as possible; that is, we express q as a product of real linear factors and real irreducible quadratic factors. The form of the partial fractions decomposition is determined by this factorisation. There are several cases, depending on the type of factorisation.

Case 1: The denominator splits into distinct linear factors. Examples of two such rational functions and the form of their partial fractions decompositions are given below:

$$\frac{x-3}{(x-1)(x-2)} = \frac{A}{x-1} + \frac{B}{x-2}$$

$$\frac{x^2-x+7}{x(2x+1)(x-3)} = \frac{A}{x} + \frac{B}{2x+1} + \frac{C}{x-3}.$$

The constants A , B and C in each case can be determined using the following method.

Example 2.4.3. Find the partial fractions decomposition of $\frac{7x-1}{x^2-2x-3}$.

Solution. By factorising we find that $x^2-2x-3 = (x-3)(x+1)$. So the partial fractions decomposition takes the form

$$\frac{7x-1}{(x-3)(x+1)} = \frac{A}{x-3} + \frac{B}{x+1},$$

where A and B are constants to be determined. To find A and B , multiply through by $(x-3)(x+1)$ to obtain the polynomial equation

$$7x-1 = A(x+1) + B(x-3) \quad \forall x \in \mathbb{R}.$$

Since this identity is true for all values x , the values of A and B are easily determined by choosing suitable values of x :

$$\begin{array}{llll} x=3 & \Rightarrow & 7 \times 3 - 1 = A(3+1) & \Rightarrow & A=5 \\ x=-1 & \Rightarrow & 7 \times (-1) - 1 = B(-1-3) & \Rightarrow & B=2. \end{array}$$

Hence the partial fractions decomposition is given by

$$\frac{7x-1}{x^2-2x-3} = \frac{7x-1}{(x-3)(x+1)} = \frac{5}{x-3} + \frac{2}{x+1}.$$

This may be easily verified by rewriting the right-hand side over a common denominator and simplifying. $\square$

Case 2: The denominator has a repeated linear factor. Examples of two such rational functions and the form of their partial fractions decompositions are given below:

$$\frac{x^2+1}{(x+4)^3} = \frac{A}{x+4} + \frac{B}{(x+4)^2} + \frac{C}{(x+4)^3}$$

$$\frac{x^2-2}{(x-1)(x-2)^2} = \frac{A}{x-1} + \frac{B}{x-2} + \frac{C}{(x-2)^2}.$$

Note carefully how the repeated factors appear on the right-hand side. The constants A , B and C in each case can be determined using the following method.

Example 2.4.4. Find the partial fractions decomposition of $\frac{x^2 - 3x + 8}{x(x-2)^2}$.

Solution. The partial fractions decomposition takes the form

$$\frac{x^2 - 3x + 8}{x(x-2)^2} = \frac{A}{x} + \frac{B}{x-2} + \frac{C}{(x-2)^2},$$

where A , B and C are constants. To find these constants, we multiply through by $x(x-2)^2$ to obtain

$$x^2 - 3x + 8 = A(x-2)^2 + Bx(x-2) + Cx \quad \forall x \in \mathbb{R}.$$

Now substitute the obvious values for x to determine the values of A and C :

$$\begin{array}{llll} x = 2 & \Rightarrow & 6 = 2C & \Rightarrow & C = 3 \\ x = 0 & \Rightarrow & 8 = 4A & \Rightarrow & A = 2. \end{array}$$

To determine B , we can substitute any other value for x . However, it is best to choose a small integer to keep the arithmetic simple:

$$x = 1 \quad \Rightarrow \quad 6 = A - B + C \quad \Rightarrow \quad B = A + C - 6 = -1.$$

(Alternately, one can find B by noting that

$$x^2 - 3x + 8 = 2(x-2)^2 + Bx(x-2) + 3x \quad \forall x \in \mathbb{R}$$

and comparing coefficients for x^2 .) Hence we obtain the partial fractions decomposition

$$\frac{x^2 - 3x + 8}{x(x-2)^2} = \frac{2}{x} - \frac{1}{x-2} + \frac{3}{(x-2)^2}.$$

□

Case 3: The denominator has an irreducible quadratic factor. Examples of two such rational functions and the form of their partial fractions decompositions are given below:

$$\begin{aligned} \frac{x^2 + x}{(x-1)(x^2 + 9)} &= \frac{A}{x-1} + \frac{Bx + C}{x^2 + 9} \\ \frac{x^3 - 2x + 4}{(x^2 + 5)(x^2 + x + 1)} &= \frac{Ax + B}{x^2 + 5} + \frac{Cx + D}{x^2 + x + 1} \end{aligned}$$

Note carefully how the irreducible quadratic appears on the right-hand side. As before, the constants A , B , C and D in each case can be determined by algebra.

Example 2.4.5. Find the partial fractions decomposition of $\frac{4x^2 + 2x + 1}{(x+1)(x^2 + x + 1)}$.

Solution. The partial fractions decomposition takes the form

$$\frac{4x^2 + 2x + 1}{(x+1)(x^2 + x + 1)} = \frac{A}{x+1} + \frac{Bx + C}{x^2 + x + 1},$$

where A , B and C are constants. Multiplying through by $(x+1)(x^2 + x + 1)$ gives

$$4x^2 + 2x + 1 = A(x^2 + x + 1) + (Bx + C)(x+1) \quad \forall x \in \mathbb{R}.$$

Now substitute suitable values for x :

$$\begin{array}{llll} x = -1 & \Rightarrow & 3 = A & \Rightarrow & A = 3 \\ x = 0 & \Rightarrow & 1 = A + C & \Rightarrow & C = 1 - A = -2 \\ x = 1 & \Rightarrow & 7 = 3A + 2(B + C) & \Rightarrow & B = 1. \end{array}$$

(Alternatively, after finding A , we could compare the coefficients of the x^2 terms on both sides to deduce that $B = 1$, and compare the constant terms to deduce that $C = -2$.) Hence

$$\frac{4x^2 + 2x + 1}{(x+1)(x^2 + x + 1)} = \frac{3}{x+1} + \frac{x-2}{x^2 + x + 1}$$

is the partial fractions decomposition. □

Case 4: The denominator has repeated irreducible quadratic factor. This case rarely appears in first year mathematics courses because it is more computationally intensive. Nevertheless, for completeness the basic form of decomposition is illustrated below:

$$\begin{aligned} \frac{x^2 + x}{(x^2 + 9)^3} &= \frac{Ax + B}{x^2 + 9} + \frac{Cx + D}{(x^2 + 9)^2} + \frac{Ex + F}{(x^2 + 9)^3} \\ \frac{x^3 - 2x + 4}{(x-2)(x^2 + x + 1)^2} &= \frac{A}{x-2} + \frac{Bx + C}{x^2 + x + 1} + \frac{Dx + E}{(x^2 + x + 1)^2}. \end{aligned}$$

As before, the constants appearing in each example can be determined by algebra.

The final example tests our ability to generalise each of these cases to rational functions whose denominators have many factors of different types.

Example 2.4.6. Write down the *form* of partial fractions decomposition for the rational function given by

$$\frac{4x^4 - 3x^2 + x - 9}{x^3(x-7)(x^2+3)^2(x^2+x+2)}.$$

(You are not required to evaluate the constant coefficients.)

Solution. The partial fractions decomposition is given by

$$\frac{4x^4 - 3x^2 + x - 9}{x^3(x-7)(x^2+3)^2(x^2+x+2)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x^3} + \frac{D}{x-7} + \frac{Ex+F}{x^2+3} + \frac{Gx+H}{(x^2+3)^2} + \frac{Ix+J}{x^2+x+2}$$

where $A, B, \dots, J$ are real constants. □

Remark 2.4.7. It is important to check that the denominator of the rational function has been completely factorised before writing down the form of partial fractions decomposition. In particular, one should check that every quadratic factor appearing in the factorisation is irreducible.

2.4.3 Integrating rational functions: two examples

In this subsection we illustrate how techniques discussed in the previous subsections are applied.

Example 2.4.8. Find $\int \frac{8x^3 - 12x^2 - 13x - 5}{2x^2 - 3x - 2} dx$.

2.4. INTEGRATING RATIONAL FUNCTIONS

43

Solution. We follow the steps outlined in Subsection 2.4.1. Denote the integrand by $f(x)$.

Step 1. Since f is improper, we begin with polynomial division. This gives

$$\begin{array}{r} 4x \\ 2x^2 - 3x - 2 \overline{) 8x^3 - 12x^2 - 13x - 5} \\ \underline{8x^3 - 12x^2 - 8x} \\ - 5x - 5 \end{array}$$

whence

$$\begin{aligned} f(x) &= 4x + \frac{-5x - 5}{2x^2 - 3x - 2} \\ &= 4x - \frac{5x + 5}{2x^2 - 3x - 2}. \end{aligned}$$

Step 2. To find the partial fractions decomposition of

$$\frac{5x + 5}{2x^2 - 3x - 2}$$

we factorise the denominator:

$$2x^2 - 3x - 2 = (2x + 1)(x - 2).$$

Hence the decomposition is given by

$$\frac{5x + 5}{(2x + 1)(x - 2)} = \frac{A}{2x + 1} + \frac{B}{x - 2},$$

where A and B are constants. Multiplying through by $(2x + 1)(x - 2)$ gives

$$5x + 5 = A(x - 2) + B(2x + 1)$$

and by using the substitution $x = 2$ we deduce that $B = 3$. By comparing coefficients of x on both sides it is easy to see that $A = -1$. Hence

$$\frac{5x + 5}{(2x + 1)(x - 2)} = \frac{-1}{2x + 1} + \frac{3}{x - 2}.$$

Step 3. By the previous two steps,

$$f(x) = 4x + \frac{1}{2x + 1} - \frac{3}{x - 2}.$$

Integrating gives

$$\int f(x) dx = 2x^2 + \frac{1}{2} \ln |2x + 1| - 3 \ln |x - 2| + C,$$

completing our answer. □

Example 2.4.9. Find $\int \frac{4x^2 - 15x + 29}{(x - 5)(x^2 - 4x + 13)} dx$.

Solution. The integrand is a proper rational function and its denominator completely factorised in the real numbers. So we immediately look for its partial fractions decomposition, which is of the form

$$\frac{4x^2 - 15x + 29}{(x-5)(x^2 - 4x + 13)} = \frac{A}{x-5} + \frac{Bx+C}{x^2 - 4x + 13}$$

for some real constants A , B and C . Hence

$$4x^2 - 15x + 29 = A(x^2 - 4x + 13) + (Bx + C)(x - 5),$$

from which appropriate substitutions allow the evaluation of the unknown constants:

$$\begin{array}{llll} x = 5 & \Rightarrow & 54 = 18A & \Rightarrow & A = 3 \\ x = 0 & \Rightarrow & 29 = 13A - 5C & \Rightarrow & C = 2 \\ x = 1 & \Rightarrow & 18 = 10A - 4(B + C) & \Rightarrow & B = 1. \end{array}$$

Hence

$$\frac{4x^2 - 15x + 29}{(x-5)(x^2 - 4x + 13)} = \frac{3}{x-5} + \frac{x+2}{x^2 - 4x + 13}.$$

The first term of the decomposition is easy to integrate. We therefore focus on the second term:

$$\begin{aligned} \int \frac{x+2}{x^2 - 4x + 13} dx &= \frac{1}{2} \int \frac{2x+4}{x^2 - 4x + 13} dx \\ &= \frac{1}{2} \left[\int \frac{2x-4}{x^2 - 4x + 13} dx + \int \frac{8}{x^2 - 4x + 13} dx \right] \\ &= \frac{1}{2} \int \frac{2x-4}{x^2 - 4x + 13} dx - 4 \int \frac{1}{x^2 - 4x + 13} dx \\ &= \frac{1}{2} \ln |x^2 - 4x + 13| - 4 \int \frac{1}{(x-2)^2 + 9} dx \\ &= \frac{1}{2} \ln |x^2 - 4x + 13| - \frac{4}{3} \tan^{-1} \left(\frac{x-2}{3} \right) + C. \end{aligned}$$

Putting everything together gives

$$\int \frac{4x^2 - 15x + 29}{(x-5)(x^2 - 4x + 13)} dx = 3 \ln |x-5| + \frac{1}{2} \ln |x^2 - 4x + 13| - \frac{4}{3} \tan^{-1} \left(\frac{x-2}{3} \right) + C.$$

□

2.5 Other substitutions

(Ref: SH10 §8.6)

The method of partial fractions allows us, in principle, to find an antiderivative, among the elementary functions, for any given rational function. So given a ‘non-standard’ integral, a sound technique for integration is to look for a substitution that will convert the given integral into the integral of rational function. Choosing a good substitution is often a matter of experience and a little inspiration.

Example 2.5.1. Evaluate the following integrals.

$$(a) \int \frac{dx}{1+x^{1/4}}$$

$$(b) \int \frac{x^{1/2}}{x^{1/3}+x^{1/4}} dx$$

$$(c) \int \frac{dx}{\sqrt{e^{2x}-1}}$$

Proof. (a) The aim is to replace the fractional power $x^{1/4}$ with something more convenient. The obvious substitution to use is $x = u^4$, which leads to the substitution $dx = 4u^3 du$. Hence

$$\int \frac{dx}{1+x^{1/4}} = \int \frac{4u^3 du}{1+u}$$

and we now have the integral of a rational function. Polynomial division gives

$$\frac{4u^3}{1+u} = 4 \left(u^2 - u + 1 - \frac{1}{1+u} \right).$$

(This result may also be obtained by writing

$$\begin{aligned} u^3 &= (u^3 + 1) - 1 \\ &= (u+1)(u^2 - u + 1) - 1, \end{aligned}$$

thus avoiding the use of polynomial long division.) Consequently,

$$\begin{aligned} \int \frac{dx}{1+x^{1/4}} &= 4 \left(\frac{u^3}{3} - \frac{u^2}{2} + u - \ln|1+u| \right) + C \\ &= \frac{4x^{3/4}}{3} - 2x^{1/2} + 4x^{1/4} - 4 \ln|1+x^{1/4}| + C. \end{aligned}$$

(b) We aim to remove the fractional powers $x^{1/2}$, $x^{1/3}$ and $x^{1/4}$. The lowest common multiple of 2, 3 and 4 is 12, so we choose the substitution $x = u^{12}$. Hence

$$\begin{aligned} \int \frac{x^{1/2}}{x^{1/3}+x^{1/4}} dx &= \int \frac{u^6}{u^4+u^3} 12u^{11} du \\ &= 12 \int \frac{u^{14}}{u+1} du. \end{aligned}$$

From here we either use polynomial division, or we can observe the factorisation

$$u^n - 1 = (u+1)(u^{n-1} - u^{n-2} + \dots + u - 1) \quad \text{when } n \text{ is even,}$$

to obtain

$$\int \frac{x^{1/2}}{x^{1/3}+x^{1/4}} dx = 12 \int \frac{u^{14}}{u+1} du = 12 \int (u^{13} - u^{12} + u^{11} - \dots + u - 1 + \frac{1}{u+1}) du.$$

It is easy to evaluate the integral from here.

(c) One option is to use the substitution

$$u = e^x \quad du = e^x dx$$

so that

$$\begin{aligned}\int \frac{dx}{\sqrt{e^{2x}-1}} &= \int \frac{e^x dx}{e^x \sqrt{e^{2x}-1}} \\ &= \int \frac{du}{u\sqrt{u^2-1}}.\end{aligned}$$

From here the integral can be evaluated using the substitution $u = \sec \theta$ or $u = \cosh \theta$. This is left as an exercise.

A better approach is to remove the square root from the very first substitution. The substitution $u^2 = e^{2x} - 1$ implies that

$$2u \frac{du}{dx} = 2e^{2x} = 2(u^2 + 1),$$

which leads to the substitution

$$dx = \frac{u du}{u^2 + 1}.$$

Hence

$$\begin{aligned}\int \frac{dx}{\sqrt{e^{2x}-1}} &= \int \frac{u du}{u(u^2 + 1)} \\ &= \int \frac{du}{u^2 + 1} \\ &= \tan^{-1} u + C \\ &= \tan^{-1} \sqrt{e^{2x}-1} + C.\end{aligned}$$

Thus the second method is more efficient than the first. □

As seen in the last example, there may more than one method to evaluate a given integral. Choosing the most efficient substitution to use is not always easy, but intuition can be developed with time, experience and practice. The next example is therefore left to the student as an exercise.

Example 2.5.2. Evaluate $\int_0^1 \frac{x^3}{(4+x^2)^{5/2}} dx$ by

- (i) using the substitution $x = 2 \tan \theta$ (since the integrand involves $\sqrt{4+x^2}$);
- (ii) using the substitution $x = 2 \sinh \theta$ (since the integrand involves $\sqrt{4+x^2}$);
- (iii) using the substitution $u^2 = 4+x^2$ (aiming for a rational function);
- (iv) using the substitution $u = 4+x^2$.

Which method works best?

The following MAPLE command is relevant to the material of this chapter:

`convert(f, parfrac, x);` performs a partial fraction decomposition of the rational function `f` in the variable `x`. For example,

`> convert(x^2/(x+2), parfrac, x);`

$$x - 2 + \frac{4}{x + 2}$$

`> convert(x/(x-b)^2, parfrac, x);`

$$\frac{b}{(x - b)^2} + \frac{1}{x - b}$$

Revision problems

1. [R] Evaluate each of the following integrals by inspection. Do not use substitution.

a) $\int x e^{2x^2} dx$	b) $\int x \sin(x^2) dx$	c) $\int x^2 \cos(2x^3) dx$
d) $\int \frac{x}{5x^2 - 11} dx$	e) $\int \sin x \cos^3 x dx$	f) $\int \frac{dx}{x \ln x}$
g) $\int \frac{x+2}{\sqrt{x^2+4x+7}} dx$	h) $\int x \sqrt{1+x^2} dx$	i) $\int x^2 \sqrt{9-4x^3} dx$
j) $\int \frac{x^2}{\sqrt{9-4x^3}} dx$	k) $\int \frac{x^3}{(1+x^4)^3} dx$	l) $\int \frac{\sec^2 x}{\tan^4 x} dx$
m) $\int \frac{\cos x}{\sin^3 x} dx$	n) $\int e^{2x} (4 + 3e^{2x})^{\frac{1}{3}} dx$	o) $\int \frac{1}{x (\ln x)^5} dx$

2. [R] Integrate the following by parts.

a) $\int x^2 e^{-x} dx$	b) $\int x^3 \ln x dx$	c) $\int \frac{x}{\cos^2 x} dx$	d) $\int \frac{(\ln x)^2}{x^2} dx$
e) $\int e^x \cos x dx$	f) $\int \ln x dx$	g) $\int \tan^{-1} x dx$	

Problems 2.1 : Trigonometric integrals

3. [R] Evaluate the following integrals.

a) $\int_0^{\pi/2} \sin^7 x \cos x dx$	b) $\int_0^{\pi} \sin^3 x \cos^2 x dx$
c) $\int \sec^3 x \tan x dx$	d) $\int \cos^2 \theta d\theta$
e) $\int \cos x \cos 10x dx$	f) $\int \sin 2x \cos 3x dx$

Problems 2.2 : Reduction formulae

4. [R]

- a) By multiplying the integrand by $\frac{\tan x + \sec x}{\tan x + \sec x}$, find $\int \sec x dx$.
- b) The reduction formula

$$\int \sec^n x dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx$$

is valid whenever $n \geq 2$. Use it to find the following integrals.

i) $\int \sec^4 x \, dx$ ii) $\int \sec^5 x \, dx$

c) [X] Prove the reduction formula given above.

5. [R] Suppose that

$$I_{m,n} = \int_0^{\pi/2} \cos^m x \sin^n x \, dx$$

whenever m and n are nonnegative integers. Use the reduction formula

$$I_{m,n} = \begin{cases} \left(\frac{m-1}{m+n}\right) I_{m-2,n} & \text{provided that } m \geq 2 \\ \left(\frac{n-1}{m+n}\right) I_{m,n-2} & \text{provided that } n \geq 2 \end{cases}$$

to evaluate the following integrals.

a) $\int_0^{\pi/2} \cos^6 x \sin^4 x \, dx$ b) $\int_0^{\pi/2} \cos^5 x \sin^5 x \, dx$ c) $\int_0^{\pi/2} \cos^3 x \sin^4 x \, dx$

6. [R] Suppose that $I_n = \int_0^1 x^n e^{-x} \, dx$. Prove that

$$I_n = n I_{n-1} - \frac{1}{e}$$

whenever $n > 0$. Hence evaluate $\int_0^1 x^3 e^{-x} \, dx$.

7. [R] It was proven in the notes that if $I_n = \int_0^{\pi/4} \tan^n x \, dx$ then

$$I_n = \frac{1}{n-1} - I_{n-2}$$

whenever $n > 1$. Use this to evaluate I_7 and I_8 .

8. [R] Suppose that $I_n = \int_1^e x(\ln x)^n \, dx$. Show that

$$I_n = \frac{1}{2} (e^2 - n I_{n-1})$$

whenever $n \geq 1$. Hence evaluate I_3 .

9. [R] By writing $\cos^n x$ as $\cos^{n-1} x \cos x$ and integrating by parts, show that

$$\int_0^{\pi/2} \cos^n x \, dx = \frac{n-1}{n} \int_0^{\pi/2} \cos^{n-2} x \, dx$$

whenever $n \geq 2$. Hence find $\int_0^{\pi/2} \cos^8 x \, dx$ and $\int_0^{\pi/2} \cos^7 x \, dx$.

10. [R] Suppose that $I_n = \int_0^1 \frac{x^n}{\sqrt{1+x}} dx$. Find a reduction formula for I_n .
11. [H] Show that $\int_0^1 x^m(1-x)^n dx = \frac{m!n!}{(m+n+1)!}$ for all nonnegative integers m and n .
12. [H] Suppose that $I_n = \int_0^{\pi/2} \cos^n x dx$.

- a) By writing the reduction formula of Question 9 as $I_n = \left(1 - \frac{1}{n}\right) I_{n-2}$, show that

$$I_{2m} = \left(1 - \frac{1}{2m}\right) \left(1 - \frac{1}{2m-2}\right) \cdots \left(1 - \frac{1}{2}\right) \frac{\pi}{2} = \frac{\pi}{2} \prod_{k=1}^m \left(1 - \frac{1}{2k}\right)$$

and

$$I_{2m+1} = \prod_{k=1}^m \left(1 - \frac{1}{2k+1}\right).$$

- b) Deduce that

$$\frac{2}{\pi} \frac{I_{2m}}{I_{2m+1}} = \prod_{k=1}^m \left(1 - \frac{1}{(2k)^2}\right).$$

- c) By considering $\cos x$ on $(0, \frac{\pi}{2})$, show that

$$I_{2m+2} \leq I_{2m+1} \leq I_{2m}$$

whenever $m \geq 1$.

- d) Use the result of (a) and (c) and the pinching theorem to deduce that

$$\lim_{m \rightarrow \infty} \frac{I_{2m}}{I_{2m+1}} = 1.$$

- e) Conclude that

$$\lim_{m \rightarrow \infty} \prod_{k=1}^m \left(1 - \frac{1}{(2k)^2}\right) = \frac{2}{\pi}.$$

This limit is called *Wallis' product*.

- f) Show that

$$\prod_{k=1}^m \left(1 - \frac{1}{(2k)^2}\right) = \prod_{k=1}^m \frac{(2k-1)(2k+1)}{(2k)^2} = \frac{(2m+1)((2m)!)^2}{2^{4m}(m!)^4}$$

and deduce that Wallis' product may be written as

$$\frac{\pi}{2} = \lim_{m \rightarrow \infty} \frac{2^{4m}(m!)^4}{(2m+1)((2m)!)^2}.$$

13. [X]

a) Show that

$$\int_m^{2m} \ln x \, dx = m \ln m + 2m \ln 2 - m.$$

b) Show that the trapezoidal rule (see SH6 page 457, SH7 page 537 or SH8 page 484) with m subintervals gives the approximate value of this integral as

$$\ln \left(\frac{(2m)!}{m!} \right) - \frac{1}{2} \ln 2.$$

c) It can be shown that if the trapezoidal rule with m intervals of equal width is used to approximate $\int_a^b f(x) \, dx$, where f is a twice differentiable function, then the absolute error in the approximation is no greater than $\frac{(b-a)^3 M}{12m^2}$, where M is the maximum of $|f''(x)|$ on $[a, b]$. Use this error bound to show that the error in the approximation in part (b) approaches 0 as $m \rightarrow \infty$.

d) Conclude that

$$\frac{(2m)!}{m!} \bigg/ \left\{ \sqrt{2} 2^{2m} m^m e^{-m} \right\} \rightarrow 1$$

as $m \rightarrow \infty$.

14. [X] Using the previous two questions (or otherwise), show that

$$\frac{\pi}{2} = \lim_{m \rightarrow \infty} \frac{(m!)^2}{4m m^{2m} e^{-2m}},$$

so that

$$\frac{m!}{\sqrt{2\pi} m^{m+\frac{1}{2}} e^{-m}} \rightarrow 1$$

as $m \rightarrow \infty$. This is sometimes called Stirling's formula (or Stirling's approximation for $m!$).

Problems 2.3 : Trigonometric and hyperbolic substitutions

15. [R] Evaluate the following integrals.

$$\text{a) } \int_0^1 \frac{x^2}{\sqrt{4-x^2}} \, dx \quad \text{b) } \int \frac{dx}{\sqrt{x^2-6x+13}}$$

$$\text{c) } \int_0^3 \sqrt{9-x^2} \, dx \quad \text{d) } \int \frac{dx}{x^2 \sqrt{x^2+16}}$$

$$\text{e) } \int (1-x^2)^{-\frac{3}{2}} \, dx \quad \text{f) } \int_{-1}^1 \frac{dx}{x^2+2x+2}$$

16. [R] Evaluate $\int \frac{x}{\sqrt{x^2-4}} \, dx$ by making an appropriate substitution. Are there any other methods or substitutions that could be used? Which one is most efficient?

Problems 2.4 : Integrating rational functions

17. [R] Evaluate the following integrals.

- a) $\int \frac{1}{x^2 + 4x + 3} dx$ b) $\int \frac{5x - 7}{x^2 - 3x + 2} dx$
 c) $\int \frac{(x + 1)}{x^2(x - 1)} dx$ d) $\int \frac{1}{(x^2 - 1)^2} dx$
 e) $\int \frac{x^2 + 1}{x^2 - 1} dx$ f) $\int \frac{18}{(x^2 + 9)(x - 3)} dx$
 g) $\int \frac{x^2 + x + 2}{(x + 1)(x + 2)^2} dx$ h) $\int \frac{1 - x}{(1 + x)^3} dx$

Problems 2.5 : Other substitutions and miscellaneous integrals

18. [R] Evaluate the following integrals.

- a) $\int \frac{x}{x^2 + 2x + 10} dx$ b) $\int \frac{x}{\sqrt{x^2 + 2x + 10}} dx$
 c) $\int \frac{dx}{1 + \sqrt{x}}$ d) [H] $\int_1^{64} \frac{1}{x^{1/2} + x^{1/3}} dx$

19. [R] Evaluate $\int_0^1 \frac{x^3}{(4 + x^2)^{5/2}} dx$ by

- a) using the substitution $x = 2 \tan \theta$ (since the integrand involves $\sqrt{4 + x^2}$);
 b) using the substitution $x = 2 \sinh \theta$ (since the integrand involves $\sqrt{4 + x^2}$);
 c) using the substitution $u^2 = 4 + x^2$ (aiming for a rational function);
 d) using the substitution $u = 4 + x^2$.

Which method works best?

20. [X] *Caution: for each integral below, it is unlikely to be profitable to seek an indefinite integral.*

- a) Use the substitution $x = \pi - u$ to show that $\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx = \frac{\pi^2}{4}$.
 b) Show that $\int_0^1 \frac{\ln(1 + x)}{1 + x^2} dx = \frac{\pi}{8} \ln 2$.

21. [X] Find $\int \frac{x^2 - 1}{x^2 + 1} \frac{1}{\sqrt{1 + x^4}} dx$.

The substitution $u^2 = x^2 + \frac{1}{x^2}$ may be useful, but there are other possible approaches.
 [Taken from Spivak's *Calculus*.]

22. [R] The following integrals were selected from past papers. Evaluate each one.

a) $\int \frac{dx}{x(x^2 + x + 1)}$

c) $\int \frac{3x + 5}{x^2 + 4x + 8} dx$

e) $\int \frac{3x^2 - 5x + 3}{(x - 1)(x^2 - 2x + 2)} dx$

g) $\int \frac{dx}{(x^2 + 3)^{3/2}}$

i) $\int_1^\infty \frac{e^{-\sqrt{x}}}{\sqrt{x}} dx$. If convergent evaluate the integral.

b) $\int 8 \sinh x \cosh^4 x dx$

d) $\int \sqrt{25 - x^2} dx$

f) $\int \frac{1}{x^2 \sqrt{1 + x^2}} dx$

h) $\int \cos(4x) \sin(3x) dx$

Chapter 3

Ordinary differential equations

In many practical applications (in physics, economics, social sciences, engineering, applied science, mathematics and so on), information is known about the relationship between a quantity and its rates of change, but one may not have an exact formula for the quantity itself. For example, a simple population model states that the rate of change of a population, at any given time, is proportional to size of the population itself. If we write $P(t)$ for the population at time t , then we arrive at the equation

$$\frac{dP}{dt} = kP,$$

where k is the constant of proportionality. An equation, such as the one given above, which involves one (or more) of the derivatives of a function, is called a *differential equation*. Some other simple examples include

- $\frac{d^2x}{dx^2} = -k^2x$, which is used to describe the displacement x from the origin of a particle undergoing simple harmonic motion;
- $\frac{dT}{dt} = k(T - 20)$, which is used to describe how the temperature T of an object changes in room temperature; and
- $\frac{dx}{dy} - 0.08y = 0.05(60000 + 1000t)$, which is used to describe how the amount y (in dollars) of a particular investment changes in time (see Example 3.4.3).

If possible, the aim from here is to find an explicit formula (or formulae) describing the unknown function (respectively P , x , T and y in the examples above) appearing in each differential equation. The primary goal of this chapter is to examine some techniques for obtaining a formula for a function given a differential equation for that function.

We begin with a definition.

Definition 3.1.1. An *ordinary differential equation* is an equation expressed in terms of exactly one independent variable and one (or more) of the derivatives of a function of this variable. The *order* of an ordinary differential equation is the order of the highest derivative present.

For example, the equation

$$\frac{d^3y}{dx^3} + \sin x \frac{dy}{dx} = 3x^2y$$

is an ordinary differential equation of order 3; the independent variable is x and y is assumed to be a function of x . The equation

$$\left(\frac{d^2x}{dt^2}\right)^{3/2} + \frac{dx}{dt} - tx = 0$$

is an ordinary differential equation of order 2; the independent variable is t and x is assumed to be a function of t .

In these notes, the term ‘ordinary differential equation’ will often be abbreviated as ODE. Such equations are called ‘ordinary’ because they involve ordinary derivatives. This is to distinguish them from differential equations that involve partial derivatives. (The study of ‘partial differential equations’ will be introduced in some second year courses.)

Ordinary differential equations can be written in several ways using a variety of notations. For example, each of the equations

$$\begin{aligned}\frac{d^2y}{dx^2} + 4x \frac{dy}{dx} &= e^x \\ f''(x) + 4xf'(x) &= e^x \\ y'' + 4xy' &= e^x\end{aligned}$$

represent the same ODE.

Definition 3.1.2. A *solution* to an n th order ordinary differential equation is a function which is n -times differentiable and satisfies the given equation.

The next example illustrates this definition as well as introducing the terms ‘particular solution’ and ‘general solution.’

Example 3.1.3. Consider the ODE

$$\frac{dy}{dx} = x^2 + 5.$$

Then the function y , given by

$$y(x) = \frac{x^3}{3} + 5x,$$

3.1. AN INTRODUCTION

is a solution to the ODE. Note that if $y(x) = \frac{x^3}{3} + 5x + 6$ or if $y(x) = \frac{x^3}{3} + 5x - 45$, then y is also a solution to the ODE. Each of these solutions is called a *particular solution* to the ODE. Using the mean value theorem (see Section 5.9 of the MATH1131 calculus notes), it is easily shown that *every* particular solution y to the ODE can be written in the form

$$y(x) = \frac{x^3}{3} + 5x + C, \quad (3.1)$$

where $C \in \mathbb{R}$. The family of solutions given by (3.1), where $C \in \mathbb{R}$, is called the *general solution* to the ODE.

In the above example, solution y could be expressed explicitly as a function of the independent variable x . Hence we obtained an *explicit solution* to the ODE. However, this cannot always be done, as the following example illustrates. Sometimes we must settle for an *implicit solution* to the ODE.

Example 3.1.4. Show that y , given implicitly by the equation

$$y^2 = \cos(x^2 + y^2), \quad (3.2)$$

is a particular solution to the ODE

$$2x \sin(x^2 + y^2) + (2y \sin(x^2 + y^2) + 2y) \frac{dy}{dx} = 0.$$

Proof. To verify that y solves the ODE, we first need to calculate $\frac{dy}{dx}$. Implicit differentiation of (3.2) with respect to x gives

$$2y \frac{dy}{dx} = -\sin(x^2 + y^2) \times \left(2x + 2y \frac{dy}{dx} \right)$$

(where we have used the chain rule to obtain the right-hand side). Hence

$$(2y + 2y \sin(x^2 + y^2)) \frac{dy}{dx} = -2x \sin(x^2 + y^2).$$

By simple rearrangement it is easily seen that

$$2x \sin(x^2 + y^2) + (2y \sin(x^2 + y^2) + 2y) \frac{dy}{dx} = 0,$$

and hence the ODE is satisfied. (This ODE shall be revisited again in Section 3.5, where we shall *find* the general solution, instead of merely verifying that a given function is a solution.) $\square$

For some differential equations, it may not even be possible to find an implicit solution. *If* it is possible to prove that an solution exists, then mathematicians and scientists must often settle for working with an approximate solution to the ODE. However, such issues will not concern us in this course.

3.2 Initial value problems

In most practical applications where ODEs are used, information is also known about the value of the unknown function and its derivatives at a particular point. This information, together with an ODE, forms an *initial value problem*.

Definition 3.2.1. An *initial value problem* is an n th order ODE together with a set of values of the solution and its first $(n - 1)$ derivatives at some fixed point x_0 . These values are called the *initial conditions* of the initial value problem.

For example,

- $\frac{dy^2}{dx^2} + 5x \frac{dy}{dx} + y = \sin x, \quad y(0) = 2, \quad \left. \frac{dy}{dx} \right|_{x=0} = 7;$
- $f'(t) - e^{2t} f(t) = 3t^2, \quad f(1) = 5;$ and
- $y''' + 3y'' + 4y = \cosh x, \quad y''(\pi) = 2, \quad y'(\pi) = 0, \quad y(\pi) = -1$

are all initial value problems. The term ‘initial value problem’ is often abbreviated as IVP.

To solve an initial value problem, we usually try to find a general solution to the ODE (which is expressed using unspecified constants) and then determine the values of these constants by imposing the initial conditions.

Example 3.2.2. Solve the IVP

$$\frac{d^2 y}{dx^2} = 6x, \quad y'(0) = 2, \quad y(0) = -1.$$

Solution. Integrating the ODE once gives

$$\frac{dy}{dx} = 3x^2 + C$$

where $C \in \mathbb{R}$. By imposing the initial condition $y'(0) = 2$, we deduce that $C = 2$. Hence

$$\frac{dy}{dx} = 3x^2 + 2.$$

Integrating again gives

$$y = x^3 + 2x + D,$$

where $D \in \mathbb{R}$. The initial condition $y(0) = -1$ implies that $D = -1$. Hence the solution y to the IVP is given by

$$y = x^3 + 2x - 1.$$

Note that this solution is valid for all x in $\mathbb{R}$; that is, the solution y is defined on $\mathbb{R}$. □

Not every initial value problem is as straightforward to solve as the example above. Solving an IVP is, in general, very difficult, and the following questions arise.

- (a) Does the IVP have a solution?

(b) Does it have a unique solution?

(c) If initial values are given at the point a , then how far on either side of a does the solution extend?

The following two examples show that care must be taken in answering such questions, even for IVPs that appear to be ‘simple.’

Example 3.2.3. Solve the initial values problem

$$\frac{dy}{dx} = \sqrt{y}, \quad y(0) = 0.$$

Solution. If we assume that $y(x) \neq 0$ then the ODE can be written as

$$\frac{dx}{dy} = \frac{1}{\sqrt{y}}, \quad (3.3)$$

from which we find that $x = 2\sqrt{y} + C$, where C is a real number. When $x = 0$ we have that $y = 0$ and hence $C = 0$. Rearranging gives

$$y(x) = \frac{x^2}{4}.$$

However, note that $y(x) = 0$ is also a solution to the IVP. Hence the IVP does not have a unique solution. $\square$

Example 3.2.4. Solve the initial values problem

$$\frac{dy}{dx} = \frac{1}{x}, \quad y(1) = 2.$$

How far does the solution extend on either side of the point 1?

Solution. First, the ODE implies that $\frac{dy}{dx}$ does not exist at 0. However, on the interval $(0, \infty)$, we obtain the general solution

$$y(x) = \ln x + C,$$

where $C \in \mathbb{R}$. The initial condition implies that $C = 2$ and so

$$y(x) = \ln x + 2$$

whenever $x > 0$. Hence we have found solution that extends to the interval $(0, \infty)$. $\square$

Remark 3.2.5. Note that in example 3.2.4 we could give a *family* of solutions defined on the set $\{x \in \mathbb{R} : x \neq 0\}$ by

$$y(x) = \begin{cases} \ln |x| + D & \text{if } x < 0 \\ \ln |x| + 2 & \text{if } x > 0, \end{cases}$$

where $D \in \mathbb{R}$. However, for most practical applications such a solution would not be used on $(-\infty, 0)$ because of the break in the domain of y at 0.

(Ref: SH10 §9.2)

A separable ODE is a differential equation where the two variables involved (say x and y) can be separated so that all the y 's are on one side of the equation and all x 's are on the other. We give an example and then state the general form.

Example 3.3.1. Solve the initial value problem

$$\frac{dy}{dx} = y^2(1 + x^2), \quad y(0) = 1.$$

Solution. First we separate the variables x and y to obtain

$$\frac{1}{y^2} dy = (1 + x^2) dx. \quad (3.4)$$

Integrating gives

$$\int \frac{1}{y^2} dy = \int (1 + x^2) dx, \quad (3.5)$$

whence

$$-\frac{1}{y} = x + \frac{x^3}{3} + C,$$

where $C \in \mathbb{R}$. This gives an *implicit* solution to the ODE.

We now impose the initial condition to evaluate the constant C . When $x = 0$ and $y = 1$ we obtain

$$-\frac{1}{1} = 0 + \frac{0^3}{3} + C,$$

whence $C = -1$ and

$$-\frac{1}{y} = x + \frac{x^3}{3} - 1.$$

Finally, in this particular example, one can easily make y the subject to obtain the *explicit* solution

$$\begin{aligned} y &= \frac{-1}{x + \frac{x^3}{3} - 1} \\ &= \frac{-3}{3x + x^3 - 3}, \end{aligned}$$

thus completing the problem. □

Remark 3.3.2. Equation (3.4) makes sense within the context of integration (see equation (3.5)). The fact that *this* kind of symbolic manipulation with dy and dx ‘works’ can be traced back to the chain rule, which is the basis for implicit differentiation. (To make this clear, note in the example above that

$$\frac{1}{y^2} \frac{dy}{dx} = 1 + x^2$$

and so integrating both sides with respect to x gives

$$\int \frac{1}{y^2} \frac{dy}{dx} dx = \int (1 + x^2) dx.$$

Hence

$$-\frac{1}{y} = x + \frac{x^3}{3} + C, \quad (3.6)$$

which may be easily verified by (implicitly) differentiating both sides of (3.6) respect to x .) Students should not think that they can manipulate the symbols dy and dx in other ways and still obtain valid results.

In general, a separable ODE is one that can be written in the form

$$\frac{dy}{dx} = \frac{g(x)}{h(y)}. \quad (3.7)$$

To solve this equation, write

$$h(y) dy = g(x) dx,$$

and then integrate both sides to obtain an implicit solution

$$H(y) = G(x) + C,$$

where C is the constant of integration. Whenever possible, isolate y on the left-hand side to find the explicit solution. If initial conditions are given, then the constant C can be determined.

Example 3.3.3. Solve the equation

$$\sinh y \cos^2 x \frac{dy}{dx} = \tan x + 4.$$

Proof. We separate the variables to obtain

$$\sinh y dy = \frac{\tan x + 4}{\cos^2 x} dx.$$

Simplification followed by integration gives

$$\begin{aligned} \sinh y dy &= (\tan x \sec^2 x + 4 \sec^2 x) dx \\ \int \sinh y dy &= \int (\tan x \sec^2 x + 4 \sec^2 x) dx \\ \cosh y &= \frac{1}{2} \tan^2 x + 4 \tan x + C, \end{aligned} \quad (3.8)$$

where $C \in \mathbb{R}$. In this case it is best to leave the solution in implicit form (3.8), since $\cosh$ is not a one-to-one function. $\square$

We end with an application of this method to the real world.

Example 3.3.4 (Newton's law of cooling). (a) Newton's law of cooling states that the rate of heat loss of a body is proportional to the difference in temperatures between the body and its surroundings. Set up an ODE to model this law and solve it.

- (b) A hot object is placed into a room of temperature 20°C . Unfortunately the object is too hot for the thermometer to measure its initial temperature. However, after 6 minutes, the temperature of the object was measured as 80°C , and after eight minutes as 50°C . What was the original temperature of the object?

Proof. (a) Suppose that

- T is the temperature of the object at time t ,
- A is the ambient temperature (that is, the temperature of the surroundings), and
- k is the constant of proportionality.

Newton's law of cooling implies that

$$\frac{dT}{dt} = k(T - A).$$

To solve this equation, we separate variables:

$$\begin{aligned}\frac{1}{T - A} dT &= k dt \\ \int \frac{1}{T - A} dT &= \int k dt \\ \ln(T - A) &= kt + C,\end{aligned}$$

where C is the constant of integration. By taking exponentials of both sides we obtain

$$T - A = e^{kt+C}.$$

Hence

$$T = A + Ke^{kt}, \tag{3.9}$$

where $K = e^C > 0$.

(b) We have

$$A = 20, \quad T(6) = 80, \quad \text{and} \quad T(8) = 50.$$

Hence (3.9) implies that

$$\begin{cases} 80 = 20 + Ke^{6k} \\ 50 = 20 + Ke^{8k}, \end{cases}$$

which can be solved to give

$$k = -\frac{\ln 2}{2} \quad \text{and} \quad K = 480.$$

Hence

$$T(0) = 20 + 480e^0 = 500.$$

So the initial temperature of the object was $500^\circ C$. □

3.4 First order linear ODEs

(Ref: SH10 §9.1)

A first order linear ODE can be written in the form

$$\frac{dy}{dx} + f(x)y = g(x), \tag{3.10}$$

where f and g are given functions of a single variable x . The ODE is called *linear* since there are no non-linear terms (such as y^2 , $\sin y$ or $\sqrt{y'}$) involving y or its derivative y' .

A very slick method for solving first order linear ODEs is summarised in the steps below.

1. Write the ODE in the form (3.10).
2. Calculate $e^{\int f(x) dx}$ (ignoring the constant of integration). We denote this by $h(x)$ and call it the *integrating factor*.
3. Multiply (3.10) by the integrating factor $h(x)$ to obtain

$$h(x) \frac{dy}{dx} + h(x)f(x)y = g(x)h(x).$$

By using the product rule for differentiation, the left-hand side can now be rewritten so that

$$\frac{d}{dx}(h(x)y) = g(x)h(x)$$

(this is easily seen in the examples that follow).

4. Integrate both sides and then rearrange for y to solve the ODE. Don't forget the constant of integration!

Example 3.4.1. Solve $\frac{dy}{dx} + 3y = e^{-x}$.

Solution. The ODE is already in the form (3.10) with $f(x)$ equal to 3. The integrating factor h is therefore given by

$$h(x) = e^{\int 3 dx} = e^{3x}.$$

Multiplying the ODE by the integrating factor e^{3x} gives

$$e^{3x} \frac{dy}{dx} + 3e^{3x}y = e^{2x}.$$

By the product rule, we can contract the left hand side to obtain

$$\frac{d}{dx}(e^{3x}y) = e^{2x}$$

(this step is easy to check by working backwards). Integrating gives

$$e^{3x}y = \frac{1}{2}e^{2x} + C,$$

where $C \in \mathbb{R}$. Now divide by e^{3x} to obtain the explicit solution

$$y = \frac{1}{2}e^{-x} + Ce^{-3x},$$

where $C \in \mathbb{R}$. (Note that if the constant of integration is accidentally omitted then we lose half the solution!) $\square$

Example 3.4.2. Solve the IVP

$$(x-1)^3 \frac{dy}{dx} + 4(x-1)^2 y = x+1, \quad y(0) = 2. \quad (3.11)$$

Solution. First rewrite the ODE into the standard form (3.10) to obtain

$$\frac{dy}{dx} + 4(x-1)^{-1}y = \frac{x+1}{(x-1)^3}. \quad (3.12)$$

The integrating factor h is given by

$$h(x) = e^{\int 4(x-1)^{-1} dx} = e^{4 \ln(x-1)} = e^{\ln(x-1)^4} = (x-1)^4.$$

(Note that it is important to simplify $h(x)$ before proceeding with the method.)

Multiplying (3.12) by the integrating factor gives

$$(x-1)^4 \frac{dy}{dx} + 4(x-1)^3 y = x^2 - 1,$$

from which we obtain

$$\frac{d}{dx}((x-1)^4 y) = x^2 - 1$$

by the product rule. Integrating gives

$$(x-1)^4 y = \frac{1}{3}x^3 - x + C,$$

where $C \in \mathbb{R}$. To evaluate C , we impose the initial condition $y(0) = 2$ and find that

$$(0-1)^4 2 = \frac{1}{3}0^3 - 0 + C.$$

Hence $C = 2$. Therefore the solution of the IVP is given by

$$y = \frac{\frac{1}{3}x^3 - x + 2}{(x-1)^4} = \frac{x^3 - 3x + 6}{3(x-1)^4}.$$

(Note that the solution is valid when $x \in (-\infty, 1)$ or $x \in (1, \infty)$ but not when $x = 1$. In fact, it is not hard to see from (3.11) that there is no real-valued function y satisfying the ODE with 1 in its domain.) $\square$

The following example shows an application of a first order linear ODE to a real world problem.

Example 3.4.3. An investor has a salary of \$60,000 per year which is expected to increase at a rate of \$1000 per annum. Suppose that an initial deposit of \$1000 is invested in a program that pays 8% per annum, and that the investor deposits 5% of their salary each year. Find the amount invested after t years.

Solution. We will approximate the situation by assuming that interest is calculated continuously and that deposits are made continuously.

Let $y(t)$ denote the dollars invested after t years. Then

$$\frac{dy}{dt} = 0.08y + 0.05(60000 + 1000t)$$

$$(\text{rate of increase of investment} = 8\% \text{ of investment} + 5\% \text{ of salary}).$$

This is a first order linear ODE. If we rewrite the ODE as

$$\frac{dy}{dt} - 0.08y = 0.05(60000 + 1000t),$$

then we see that the integrating factor h is given by

$$h(t) = e^{\int -0.08 dt} = e^{-0.08t}.$$

After multiplying the ODE by the integrating factor and contracting the left-hand side by the product rule, one obtains

$$\frac{d}{dt}(e^{-0.08t}y) = 0.05(60000 + 1000t)e^{-0.08t}.$$

Integration (where we use integration by parts for the right-hand side) and rearrangement gives

$$y(t) = -625t - 46312.5 + Ce^{0.08t},$$

where C is the constant of integration. Imposing the initial condition $y(0) = 1000$ yields the final solution

$$y(t) = 45312.5 e^{0.08t} - 625t - 46312.5,$$

where $t \geq 0$.

To illustrate, note that after 10 years the investment totals about $y(10) \approx 51507.86$ dollars. $\square$

3.5 Exact ODEs

(Ref: SH10 §19.2)

In this section we examine another approach to solving (some) first order ODEs.

To begin, suppose that H is a function of two variables x and y satisfying the equation

$$H(x, y) = C,$$

where C is a real constant. If we consider y as a function of x and differentiate both sides with respect to x , then the chain rule gives

$$\frac{\partial H}{\partial x} + \frac{\partial H}{\partial y} \frac{dy}{dx} = 0.$$

If $\frac{\partial H}{\partial x}$ and $\frac{\partial H}{\partial y}$ are denoted by F and G respectively, then we obtain the differential equation

$$F(x, y) + G(x, y) \frac{dy}{dx} = 0. \quad (3.13)$$

Of course, if F and G are defined as above, then $H(x, y) = C$ is a solution to (3.13).

Conversely, suppose that we want to solve a differential equation of the form (3.13). The above discussion shows that *if* there exists a function H of two variables such that $F = \frac{\partial H}{\partial x}$ and $G = \frac{\partial H}{\partial y}$ *then* the solution is given by $H(x, y) = C$, where $C \in \mathbb{R}$. The difficulty is, this condition on F and G is not so easy to verify. Fortunately, there is an easier condition. Recall, by the theorem on mixed partial derivatives, that if H is a ‘nice’ function then

$$\frac{\partial^2 H}{\partial y \partial x} = \frac{\partial^2 H}{\partial x \partial y},$$

or in other words,

$$\frac{\partial F}{\partial y} = \frac{\partial G}{\partial x}.$$

This second condition on F and G is known as the condition for *exactness* and is much easier to verify.

We summarise these observations in the next definition and theorem.

Definition 3.5.1. An ordinary differential equation of the form

$$F(x, y) + G(x, y) \frac{dy}{dx} = 0$$

is called *exact* if

$$\frac{\partial F}{\partial y} = \frac{\partial G}{\partial x}.$$

Theorem 3.5.2. Suppose that an ordinary differential equation of the form (3.13) is exact. Then the solution to (3.13) is given by $H(x, y) = C$, where C is a constant and where H is a function satisfying the equations

$$\frac{\partial H}{\partial x} = F \quad \text{and} \quad \frac{\partial H}{\partial y} = G.$$

Remark 3.5.3. The differential equation (3.13) is equivalent to

$$\frac{dy}{dx} = -\frac{F(x, y)}{G(x, y)}$$

or

$$F(x, y)dx + G(x, y)dy = 0.$$

The left-hand side of the second expression is an example of a *differential form*.

Example 3.5.4. Show that the differential equation

$$\frac{dy}{dx} = -\frac{2x + y + 1}{2y + x + 1}$$

is exact, and hence find its solution.

Solution. First we rewrite the differential equation to obtain

$$(2x + y + 1) + (2y + x + 1) \frac{dy}{dx} = 0.$$

Write $F = 2x + y + 1$ and $G = 2y + x + 1$. Then

$$\frac{\partial F}{\partial y} = 1 = \frac{\partial G}{\partial x},$$

so the differential equation is exact. Hence there exists a function H satisfying

$$\frac{\partial H}{\partial x} = F(x, y) = 2x + y + 1, \tag{3.14}$$

$$\frac{\partial H}{\partial y} = G(x, y) = 2y + x + 1. \tag{3.15}$$

To find H , we begin by integrating (3.14) with respect to x (and treating y as a constant), so that

$$H(x, y) = x^2 + xy + x + C_1(y), \quad (3.16)$$

where the ‘constant of integration’ $C_1(y)$ is a function of y . Similarly, integrating (3.15) with respect to y (and treating x as a constant) gives

$$H(x, y) = y^2 + xy + y + C_2(x), \quad (3.17)$$

where the ‘constant of integration’ $C_2(x)$ is a function of x . A comparison of (3.16) and (3.17) shows that

$$H(x, y) = x^2 + xy + y^2 + x + y. \quad (3.18)$$

Hence the solution to the differential equation is given by

$$x^2 + xy + y^2 + x + y = C, \quad (3.19)$$

where C is a real constant.

Note that, since (3.19) is a quadratic in y , we could rewrite the solution explicitly for y to give

$$y = \frac{-(x+1) \pm \sqrt{(x+1)^2 - 4(x^2 + x - C)}}{2}, \quad C \in \mathbb{R}$$

(this gives *two* functions for every value of C). However, in this case the solution is probably better left in implicit form, as in (3.19). $\square$

Remark 3.5.5. Technically, (3.18) should read

$$H(x, y) = x^2 + xy + y^2 + x + y + K,$$

where K is an arbitrary constant. However, then the solution to the ODE is given by

$$x^2 + xy + y^2 + x + y + K = C_0,$$

where C_0 is yet another constant. By combining the constants K and C_0 on the right-hand side, this solution is equivalent to (3.19). Hence it is customary to ignore the constant K .

The next example follows the same overall strategy, but illustrates a slightly different approach to finding the function H .

Example 3.5.6. Solve the differential equation

$$2x \sin(x^2 + y^2) + (2y \sin(x^2 + y^2) + 2y) \frac{dy}{dx} = 0.$$

Solution. Write $F(x, y) = 2x \sin(x^2 + y^2)$ and $G(x, y) = 2y \sin(x^2 + y^2) + 2y$. It is easily seen that

$$\frac{\partial F}{\partial y} = 4xy \cos(x^2 + y^2) = \frac{\partial G}{\partial x},$$

and so the equation is exact. Hence we look for a function H satisfying

$$\frac{\partial H}{\partial x} = F(x, y) = 2x \sin(x^2 + y^2), \quad (3.20)$$

$$\frac{\partial H}{\partial y} = G(x, y) = 2y \sin(x^2 + y^2) + 2y. \quad (3.21)$$

As with the previous example, we integrate (3.20) with respect to x (and treating y as a constant) to obtain

$$H(x, y) = -\cos(x^2 + y^2) + C_1(y), \quad (3.22)$$

where the ‘constant of integration’ $C_1(y)$ is a function of y . So now we only need to determine $C_1(y)$. To do so, differentiating (3.22) with respect to y (and treating x as a constant) gives

$$\frac{\partial H}{\partial y} = 2y \sin(x^2 + y^2) + C_1'(y).$$

Comparing this with (3.21) shows that $C_1'(y) = 2y$, whence $C_1(y) = y^2$. (Here we omit the constant of integration for reasons given in Remark 3.5.5.) Hence

$$H(x, y) = -\cos(x^2 + y^2) + y^2$$

and the solution to the differential equation is given by

$$-\cos(x^2 + y^2) + y^2 = C,$$

where $C \in \mathbb{R}$. (Note that in this example it is not possible to give an explicit expression for y in terms of x .) $\square$

The final example gives an interesting variation on this theme.

Example 3.5.7. Solve the differential equation

$$(e^x - \sin y) dx + \cos y dy = 0. \quad (3.23)$$

Solution. Suppose that $F(x, y) = e^x - \sin y$ and $G(x, y) = \cos y$. Since

$$\frac{\partial F}{\partial y} = -\cos y \quad \text{and} \quad \frac{\partial G}{\partial x} = 0,$$

the differential equation is *not* exact. What happens if we use the method of the last two examples regardless?

Suppose that there is a function H such that

$$\frac{\partial H}{\partial x} = F(x, y) = e^x - \sin y, \quad (3.24)$$

$$\frac{\partial H}{\partial y} = G(x, y) = \cos y. \quad (3.25)$$

Integrating (3.24) with respect to x gives

$$H(x, y) = e^x - x \sin y + C_1(y),$$

where $C_1(y)$ is *independent of* x . Partial differentiation with respect to y yields

$$\frac{\partial H}{\partial y} = -x \cos y + C_1'(y).$$

If we compare this with (3.25), then we conclude that $C_1'(y) = (1 + x) \cos y$, which contradicts the fact that $C_1(y)$ is independent of x . Hence no such function H exists.

Fortunately, not all is lost. If we multiply (3.23) through by the function e^{-x} , then we obtain

$$(1 - e^{-x} \sin y) dx + e^{-x} \cos y dy = 0. \quad (3.26)$$

Since the integrating factor e^{-x} is never zero, solutions to (3.26) will also be solutions to (3.23). Moreover, it is easily verified that (3.26) is exact. Hence we can now (successfully) use the method for solving exact differential equations to obtain the solution

$$x + e^{-x} \sin y = C,$$

where $C \in \mathbb{R}$. Details are left to the reader. $\square$

Remark 3.5.8. As illustrated in the previous example, if an ODE of the form

$$F(x, y) dx + G(x, y) dy = 0$$

is not exact, then it may be possible to transform it into an exact ODE by multiplying through by a suitable function. In general, finding such a function is difficult and lies beyond the scope of this course.

3.6 Solving ODEs by using a change of variable [X]

(Ref: SH10 §19.1)

This section is for MATH1241 students only. We have seen in previous sections how to solve separable, linear and exact first order ODEs. While not all first order ODEs are among these types, some can be transformed into one of these types by a suitable change of variables. We illustrate the principle with two examples.

Example 3.6.1. Use the substitution $y(x) = x \cdot v(x)$ to solve the differential equation

$$\frac{dy}{dx} = \frac{xy - y^2}{x^2}. \quad (3.27)$$

Solution. The idea is to transform (3.27) into a separable ODE involving v and x . Once the general solution for v is found, then it is easy to write down the solution for y .

Using the substitution $y(x) = x \cdot v(x)$ and the product rule for differentiation, we see that

$$\frac{dy}{dx} = \frac{d}{dx}(xv) = v \frac{dx}{dx} + x \frac{dv}{dx} = v + x \frac{dv}{dx}.$$

Hence (3.27) becomes

$$v + x \frac{dv}{dx} = \frac{x(xv) - (xv)^2}{x^2}.$$

If we simplify the right-hand side then

$$v + x \frac{dv}{dx} = v - v^2,$$

and hence we obtain the separable ODE

$$-\frac{dv}{v^2} = \frac{dx}{x}.$$

Integrating both sides gives

$$\frac{1}{v} = \ln |x| + C$$

which implies that

$$v = \frac{1}{\ln |x| + C}$$

where $C \in \mathbb{R}$. Now $v = \frac{y}{x}$ and so

$$y = \frac{x}{\ln |x| + C}$$

gives the general solution to (3.27). □

Example 3.6.2. Solve the equation

$$\frac{dy}{dt} + 2y + y^2 t^2 e^{2t} = 0 \quad (3.28)$$

by using the substitution $z = 1/y$.

Solution. Note that (3.28) is not a first order linear ODE because of the term involving y^2 . In this example we will see that the nonlinear equation in y and t becomes a linear ODE in z and t under the transformation $z = 1/y$.

To make use of the substitution $z = 1/y$, we need to express $\frac{dy}{dx}$ in terms of z . To do so, differentiate both sides of the equation

$$y = \frac{1}{z}$$

with respect to t to obtain

$$\frac{dy}{dt} = -\frac{1}{z^2} \frac{dz}{dt}.$$

With this substitution, (3.28) becomes

$$-\frac{1}{z^2} \frac{dz}{dt} + \frac{2}{z} + \frac{t^2 e^{2t}}{z^2} = 0.$$

Rearranging gives

$$\frac{dz}{dt} - 2z = t^2 e^{2t},$$

which is a first order linear ODE in z . As usual, multiply through by the integrating factor e^{-2t} to obtain

$$\frac{d}{dt}(e^{-2t} z) = t^2.$$

Integrating gives

$$e^{-2t} z = \frac{t^3}{3} + C_0,$$

whereupon

$$z = e^{2t} \left(\frac{t^3}{3} + C_0 \right),$$

where $C_0 \in \mathbb{R}$. Since $y = 1/z$, the solution to (3.28) is given by

$$y = \frac{3}{e^{2t}(t^3 + C)},$$

where $C = 3C_0 \in \mathbb{R}$. □

(Ref: SH10 §9.1, 9.2)

Many real-life problems can be analysed and solved by attempting to convert them into mathematics. In doing so, a number of assumptions have to be made and a theoretical framework set up which attempts to reflect what is happening in the real world. Such a framework is called a *mathematical model*. The reliability of that model can be judged by how well it predicts what actually happens in the real world.

To construct a mathematical model, one should

1. describe accurately the data we have,
2. decide exactly what information we want to extract from the model,
3. decide which variables in the model are dependent and which are independent, and
4. describe how the dependent variables change as the independent ones vary (which may lead to a differential equation).

Examples of mathematical modelling with differential equations have already been encountered (see Examples 3.3.4 and 3.4.3). The following two subsections provide further examples and discussion.


3.7.1 Mixing problems

In this subsection we illustrate how to construct a mathematical model that leads to a differential equation.

Example 3.7.1. A martini drink is, in essence, a mixture of the two liquids *gin* and *vermouth*. James Blond insists that his martinis be prepared as follows. Initially, 40 cc of gin are placed in a large container. Then gin is poured into the container at a rate of 2 cc/sec and at the same time vermouth is poured in at a rate of 6 cc/sec. The mixture is constantly shaken (not stirred) and flows out at a rate of 4 cc/sec.

- (a) Find an expression for the volume of vermouth in the container t seconds after the pouring commences.
- (b) James likes his martini to have roughly two parts gin to three of vermouth. How many seconds should elapse before he stops pouring and inserts a cocktail glass in the outflow from the container?

Solution. (a) To begin, it is recommended that we place the relevant information on a diagram.



Next, it is important to identify what we want to find. In this case, we want a formula for the volume of Vermouth. So let $V(t)$ denote the volume of vermouth (in cc) in the container at time t .

Now we write down as much information about V as we can. We know that $V(0) = 0$. The other information given tells us how V changes with time. In particular,

$$\begin{aligned}\frac{dV}{dt} &= \text{rate of change of } V \\ &= (\text{rate of inflow}) - (\text{rate of outflow}).\end{aligned}\tag{3.29}$$

Now the rate of inflow of vermouth is 6 cc/sec. To calculate the the rate of outflow, we note that the total volume of liquid in the container at time t is given by

$$40 + 2t + 6t - 4t = 40 + 4t.$$

Hence the *proportion* by volume of vermouth in the container at time t is

$$\frac{V(t)}{40 + 4t}.\tag{3.30}$$

Since the rate of outflow of liquid is 4 cc/sec, it follows that the rate of outflow of vermouth is

$$\frac{V(t)}{40 + 4t} \times 4 \text{ cc/sec}.$$

Following from (3.29), we obtain the initial value problem

$$\frac{dV}{dt} = 6 - \frac{V(t)}{10 + t}, \quad V(0) = 0.\tag{3.31}$$

Finally, we solve the IVP. The ODE is first order linear equation, whose solution (when $V(0) = 0$) is given by

$$V(t) = \frac{3t^2 + 60t}{10 + t}.\tag{3.32}$$

The details for finding this solution are left to the reader as an exercise.

(b) The liquid will be two parts gin to three of vermouth exactly when three-fifths of the liquid is vermouth. Hence, using (3.30), we require that

$$\frac{V(t)}{40 + 4t} = \frac{3}{5}.$$

But $V(t)$ is given by (3.32), so we require that

$$\frac{3t^2 + 60t}{4(10 + t)^2} = \frac{3}{5}.$$

By rearranging this equation, we obtain the quadratic equation

$$t^2 + 20t - 400 = 0.$$

The quadratic formula shows that

$$t = \sqrt{500} - 10 \approx 12.36,$$

where we have chosen the positive solution to the quadratic equation. Hence James should insert the cocktail glass 12.36 seconds after the mixing process begins. $\square$

3.7.2 Population models

In this subsection we compare three different mathematical models for population growth.

Suppose that the city of Mathopolis initially has a population of 3,000,000 inhabitants and (initially) grows at a rate of 2% per annum. Can we predict what the population of the city will be in 10, 20 or 100 years time? In the next three examples, we examine different population models, discuss their accuracy and see what growth forecast each model gives for Mathopolis.

Example 3.7.2 (Population model 1). In this model, we assume that the growth rate of the population remains constant. That is, we assume that the rate of change of population is proportional to the population, where the constant of proportionality r is the growth rate. In other words, we have the IVP

$$\frac{dP}{dt} = rP, \quad P(0) = P_0,$$

where P is the population at time t , r is the growth rate and P_0 is the initial population. Solving this (separable) ODE gives the solution

$$P(t) = P_0 e^{rt}.$$

(This model goes back to Thomas Malthus' book *An Essay on the Principle of Population*, published in 1798.)


Application. For Mathopolis, $P_0 = 3,000,000$ and $r = 0.02$. Thus the population $P(t)$ of the city at time t is given by

$$P(t) = 3000000e^{0.02t}.$$

The table in Figure 3.1 shows the predicted population when t is 10, 20 and 100. The graph of P against t is also shown.

	Initially	After 10 years	After 20 years	After 100 years
Model 1	3.00 million	3.66 million	4.48 million	22.17 million
Model 2	3.00 million	3.56 million	4.04 million	6.11 million
Model 3	3.00 million	3.61 million	4.21 million	6.73 million

(a) Table showing population forecast according to various models



(b) Graphs showing how models compare

Figure 3.1: Population growth for a city as projected by different models.

Criticisms of this model.

1. The model predicts that population will grow indefinitely. This ignores the fact that the resources and space needed to support such a population are finite.
2. The model ignores external factors (such as disease, natural disasters and wars) that have an effect on population size.

In light of the first criticism, we introduce a second model.

Example 3.7.3 (Population model 2). In this model the rate of population growth is not proportional to the population. Instead, there is a critical population P_c which when exceeded causes the population P to decrease; otherwise the population increases. We try the IVP

$$\frac{dP}{dt} = k(P_c - P), \quad P(0) = P_0, \quad (3.33)$$

where k is a positive constant. Hence the rate of change of P is positive if $P < P_c$ and negative if $P > P_c$.

To solve (3.33), we separate the variables and integrate:

$$\int \frac{dP}{P_c - P} = \int k dt.$$

In the case when $P < P_c$ we obtain

$$-\ln(P_c - P) = kt + C$$

and so

$$P = P_c - Ae^{-kt},$$

where $A = e^{-C}$. Now $P = P_0$ when $t = 0$ and so $A = P_c - P_0$. Hence

$$P(t) = P_c - (P_c - P_0)e^{-kt}.$$

(The case when $P > P_c$ gives the same solution, as can be easily verified by carefully working through the details.) Note that $P(t) \rightarrow P_c$ as $t \rightarrow \infty$.

Application. For Mathopolis, $P_0 = 3,000,000$ and $\frac{dP}{dt} = 0.02P_0 = 60,000$ when $t = 0$. Assume also that $P_c = 7,000,000$. (That is, due to available land, resources and other factors, one expects the city's maximum sustainable population size is 7 million.) By considering the differential equation (3.33) when $t = 0$, we conclude that

$$60000 = \frac{dP}{dt} = k(P_c - P_0) = k(7000000 - 3000000)$$

and hence that $k = 0.015$. Therefore the population $P(t)$ of Mathopolis at time t is given by

$$P(t) = 7000000 - 4000000e^{-0.015t}.$$

See Figure 3.1 for specific population projections under this model and a corresponding graph.

Criticisms of this model.

1. Observe that if P is close to 0 then $\frac{dP}{dt} \approx kP_c$, which means that when the population is very small the growth rate may be a large positive number. In fact, the rate of increase is most rapid for tiny populations!
2. As with the first population model, external factors are ignored.

In light of the first criticism we introduce a third model.

Example 3.7.4 (Population model 3). To overcome the first criticism of the previous model, we instead try the IVP

$$\frac{dP}{dt} = kP(P_c - P), \quad P(0) = P_0, \quad (3.34)$$

where k is a positive constant and P_c is the critical population. Note that $\frac{dP}{dt}$ is small when P is small. (This model was first published by Pierre Verhulst in 1838 after he had read Thomas Malthus' *An Essay on the Principle of Population*.)

To solve (3.34), separation of variables and integration gives

$$\int \frac{dP}{P(P_c - P)} = \int k dt.$$

The integral on the left-hand side is evaluated by the method of partial fractions:

$$\frac{1}{P_c} \int \frac{1}{P} + \frac{1}{P_c - P} dP = \int k dt.$$

From here it is not difficult to show that

$$P(t) = \frac{P_c P_0}{P_0 + (P_m - P_0)e^{-kP_c t}}$$

(try this as an exercise). Note once again that $P(t) \rightarrow P_c$ as $t \rightarrow \infty$. The resulting curve (see, for example, the solid gray curve in Figure 3.1 (b)) is called a *logistic curve*. The initial stage of growth is approximately exponential; then, as P approaches P_c the growth slows and approaches P_c asymptotically.

Application. Once again, for Mathopolis we have the data

$$P_0 = 3,000,000, \quad P_c = 7,000,000 \quad \text{and} \quad \frac{dP}{dt} = 0.02P_0 = 60,000 \quad \text{when} \quad t = 0.$$

By considering the differential equation (3.34) when $t = 0$, we conclude that

$$60000 = \frac{dP}{dt} = kP_0(P_c - P_0)$$

and hence that $k = 5 \times 10^{-9}$. Therefore the population $P(t)$ of the city at time t is given by

$$P(t) = \frac{21000000}{3 + 4e^{-0.035t}}.$$

See Figure 3.1 for specific population projections under this model.

Criticisms of this model. As with the other population models, external factors are ignored. In particular, P_c may change due to factors such as technological advances or climate change.

Remark 3.7.5. Note that as the models introduced become more realistic, the mathematics needed to solve the corresponding differential equations is more sophisticated. In the case of modelling fluid flow (such as water flow in the pipes or air flow around an aeroplane wing), the set of (partial) differential equations which must be solved (known as the *Navier–Stokes equations*) raises difficulties that are beyond the grasp of current mathematical knowledge. For example, it is as yet unknown whether a solution to these equations always exists and whether (in the case that a solution exists) it is a ‘smooth’ solution. The Clay Mathematics Institute has listed these problems as one of the seven *Millennium problems* and carries prize money of US\$1,000,000 for a correct solution.

3.8 Second order linear ODEs with constant coefficients

(Ref: SH10 §9.3, §19.4)

In this section we consider a special class of second order equations, known as *second order linear ODEs with constant coefficients*. An equation of this class has the form

$$\frac{d^2 y}{dx^2} + a \frac{dy}{dx} + by = f(x), \tag{3.35}$$

where a and b are real numbers. Such equations naturally arise in modelling wave mechanics and prey-predator interaction. You will have seen in earlier calculus courses (and in Physics) that the second order equation $\frac{d^2 x}{dt^2} + n^2 x = 0$ is used to model simple harmonic motion.

To simplify our treatment of the second order ODE (3.35), we first look at the case when $f(x) \equiv 0$ (which means that $f(x) = 0$ for all x).

Definition 3.8.1. A second order linear ODE with constant coefficients is said to be *homogeneous* if it is of the form

$$\frac{d^2y}{dx^2} + a\frac{dy}{dx} + by = 0, \quad (3.36)$$

where a and b are real numbers.

It turns out that we can always solve a homogeneous second order ODE with real coefficients. The first important observation towards proving this fact is given by the following lemma.

Lemma 3.8.2. If y_1 and y_2 are two solutions to the differential equation (3.36) then any linear combination $Ay_1 + By_2$, where A and B are real numbers, is also a solution to (3.36).

Proof. Suppose that y_1 and y_2 are two solutions to the differential equation (3.36). If $y = Ay_1 + By_2$, where A and B are real numbers, then

$$\begin{aligned} y'' + ay' + by &= (Ay_1 + By_2)'' + a(Ay_1 + By_2)' + b(Ay_1 + By_2) \\ &= Ay_1'' + Ay_2'' + Aay_1' + Bay_2' + Aby_1 + Bby_2 \\ &= A(y_1'' + ay_1' + by_1) + B(y_2'' + ay_2' + by_2) \\ &= 0 + 0 \quad (\text{since } y_1 \text{ and } y_2 \text{ are solutions}) \\ &= 0. \end{aligned}$$

Hence y is also a solution to (3.36). □

Remark 3.8.3. It can also be shown that every second order ODE has at most two linearly independent solutions (this will be demonstrated in second year linear algebra courses). Hence if y_1 and y_2 are two *linearly independent* solutions of (3.36) then every solution y to (3.36) is of the form $y = Ay_1 + By_2$. In this context, y_1 and y_2 are linearly independent if and only if they are not constant multiples of each other.

In view of the above lemma and remark, to find a complete solution to (3.36), one only needs to find two linearly independent solutions. To look for a solution to (3.36), we try a function y that does not change too much when differentiated. (The idea is that, upon substitution, the terms on the left-hand side need to cancel each other out to give zero.) If $y = e^{\lambda x}$, where λ is a constant, then $y' = \lambda e^{\lambda x}$ and $y'' = \lambda^2 e^{\lambda x}$. When these are substituted into (3.36) we obtain

$$\lambda^2 e^{\lambda x} + a\lambda e^{\lambda x} + be^{\lambda x} = 0.$$

Dividing by $e^{\lambda x}$ gives

$$\lambda^2 + a\lambda + b = 0, \quad (3.37)$$

Hence we have shown that

$$y = e^{\lambda x} \text{ is a solution to (3.36) if and only if } \lambda \text{ is a root of (3.37).}$$

We give the quadratic equation (3.37) a special name.

Definition 3.8.4. The *characteristic equation* of the second order linear ODE

$$\frac{d^2y}{dx^2} + a\frac{dy}{dx} + by = 0,$$

is given by

$$\lambda^2 + a\lambda + b = 0. \quad (3.38)$$

The following example illustrates what we have learnt so far.

Example 3.8.5. Solve the second order homogeneous linear ODE

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = 0. \quad (3.39)$$

Solution. The characteristic equation associated to (3.39) is given by

$$\lambda^2 - 5\lambda + 6 = 0.$$

By solving the quadratic equation we find that $\lambda = 2, 3$. Hence $y_1 = e^{2x}$ and $y_2 = e^{3x}$ are solutions to (3.39). By Lemma 3.8.2 the linear combination y , given by

$$y = Ay_1 + By_2 = Ae^{2x} + Be^{3x}$$

where A and B are real numbers, is also a solution to (3.39). Moreover, since y_1 and y_2 are linearly independent, *every* solution is of this form (see Remark 3.8.3). $\square$

Note that in the last example, the characteristic equation had two distinct real roots, thus leading to two linearly independent solutions to the homogeneous ODE. In general, there are three possibilities since all the coefficients we consider are *real* numbers. Either (i) the characteristic equation has two distinct real roots, (ii) the characteristic equation has a repeated real root, or (iii) the characteristic equation has two distinct complex roots (which are complex conjugates of each other).

Case (i): The characteristic equation has two distinct real roots λ_1 and λ_2 . Then, as seen above, we obtain two linearly independent solutions $y_1 = e^{\lambda_1 x}$ and $y_2 = e^{\lambda_2 x}$. Hence the general solution is given by

$$y = Ae^{\lambda_1 x} + Be^{\lambda_2 x},$$

where A and B are real numbers.

Case (ii): The characteristic equation has a repeated real root λ_1 . Then one solution is given by $y_1 = e^{\lambda_1 x}$. Is there another independent solution? It turns out in this case that $y_2 = xe^{\lambda_1 x}$ also solves the homogeneous equation (3.36), as can be easily verified by substituting this into the left-hand side of (3.36). Hence the general solution is given by

$$y = Ae^{\lambda_1 x} + Bxe^{\lambda_1 x},$$

where A and B are real numbers.

Case (iii): The characteristic equation has two distinct complex roots $\alpha + \beta i$ and $\alpha - \beta i$, where α and β are real numbers and $\beta \neq 0$. Then we obtain two solutions

$$y_1 = e^{(\alpha + \beta i)x} \quad \text{and} \quad y_2 = e^{(\alpha - \beta i)x}.$$

$$y = Ce^{(\alpha+\beta i)x} + De^{(\alpha-\beta i)x},$$

and C and D are (complex) constants. The goal is to choose C and D so that the solution is real. Now Euler's formula

$$e^{i\theta} = \cos \theta + i \sin \theta$$

gives

$$\begin{aligned} y &= Ce^{(\alpha+\beta i)x} + De^{(\alpha-\beta i)x} \\ &= e^{\alpha x}(Ce^{\beta i x} + De^{-\beta i x}) \\ &= e^{\alpha x}(C(\cos \beta x + i \sin \beta x) + D(\cos \beta x - i \sin \beta x)) \\ &= e^{\alpha x}((C + D) \cos \beta x + i(C - D) \sin \beta x) \\ &= e^{\alpha x}(A \cos \beta x + B \sin \beta x), \end{aligned}$$

where $A = C + D$, $B = i(C - D)$. If we choose C and D to be complex conjugates of each other, but otherwise with arbitrary real and imaginary parts, then A and B will be real. It is easy to see that $e^{\alpha x} \cos(\beta x)$ and $e^{\alpha x} \sin(\beta x)$ are independent solutions; hence the general solution in this case is given by

$$y = e^{\alpha x}(A \cos \beta x + B \sin \beta x)$$

where A and B are real numbers.

We summarise our findings in the following theorem.

Theorem 3.8.6. *Consider the second order homogeneous ODE given by (3.36) and let λ_1 and λ_2 denote the roots of the corresponding characteristic equation (3.38).*

(i) *If λ_1 and λ_2 are different real numbers then the solution to (3.36) is given by*

$$y = Ae^{\lambda_1 x} + Be^{\lambda_2 x},$$

where $A, B \in \mathbb{R}$.

(ii) *If $\lambda_1 = \lambda_2$ then the solution to (3.36) is given by*

$$y = Ae^{\lambda_1 x} + Bxe^{\lambda_1 x},$$

where $A, B \in \mathbb{R}$.

(iii) *If $\lambda_1 = \alpha + \beta i$ and $\lambda_2 = \alpha - \beta i$, where $\alpha, \beta \in \mathbb{R}$ and $\beta \neq 0$, then the solution to (3.36) is given by*

$$y = e^{\alpha x}(A \cos(\beta x) + B \sin(\beta x)),$$

where $A, B \in \mathbb{R}$.

Example 3.8.7. Solve the following differential equations:

(a) $y'' - 6y' + 25 = 0$,

(b) $y'' + 4y' + 4y = 0$, with initial conditions $y(0) = 1$ and $y'(0) = 0$.

Solution. (a) The characteristic equation

$$\lambda^2 - 6\lambda + 25 = 0$$

has roots $3+4i$ and $3-4i$ (as determined by completing the square or using the quadratic formula). Hence the solution y to the ODE is given by

$$y = e^{3x}(A \cos 4x + B \sin 4x),$$

where A and B are real numbers.

(b) The left-hand side of the characteristic equation

$$\lambda^2 + 4\lambda + 4 = 0$$

is easily factorised to give

$$(\lambda + 2)^2 = 0.$$

Hence -2 is a repeated root and the solution y to the IVP is given by

$$y = Ae^{-2x} + Bxe^{-2x}, \quad (3.40)$$

where the constants A and B are to be determined by imposing initial conditions. Differentiation shows that

$$y' = -2Ae^{-2x} - 2Bxe^{-2x} + Be^{-2x}. \quad (3.41)$$

When $x = 0$, (3.40), (3.41) and the initial conditions imply that $A = 1$ and $B = 2$. Hence

$$y = e^{-2x} + 2xe^{-2x}$$

describes the solution y to the IVP. □

3.8.2 The non-homogeneous case

We return to solving the second order linear ODE (3.36) in the case when f is not identically zero. The main idea will be illustrated in the following example.

Example 3.8.8. Solve the equation

$$y'' - 5y' + 6y = 12x - 4. \quad (3.42)$$

Solution. Since the first and second derivatives of a polynomial are also polynomials, it seems likely that at least one particular solution y_P to the ODE is a polynomial. A little more thought shows that if y_P is polynomial that satisfies (3.42), then the degree of y_P is no greater than one. So we look for a particular solution y_P of the form

$$y_P = ax + b,$$

where a and b are real numbers whose values are to be determined. Now $y'_P = a$ and $y''_P = 0$, so substituting y_P into (3.42) gives

$$0 - 5a + 6(ax + b) = 12x - 4.$$

By equating coefficients we find that $a = 2$ and $b = 1$. Hence one particular solution y_P to (3.39) is given by $y_P = 2x + 1$.

Are there any other solutions? The answer to this question is ‘yes’. To find them, we consider the associated homogeneous ODE

$$y'' - 5y' + 6y = 0, \quad (3.43)$$

whose solution y_H is given by

$$y_H = Ae^{2x} + Be^{3x},$$

where A and B are real numbers (see Example 3.8.5). Now we will show that y , where $y = y_H + y_P$, is a solution to (3.42):

$$\begin{aligned} y'' - 5y' + 6y &= (y_H + y_P)'' - 5(y_H + y_P)' + 6(y_H + y_P) \\ &= (y_H'' - 5y_H' + 6y_H) + (y_P'' - 5y_P' + 6y_P) && \text{(by linearity of differentiation)} \\ &= 0 + (12x - 4) && \text{(by the properties of } y_H \text{ and } y_P) \\ &= 12x - 4. \end{aligned}$$

Hence y , given by

$$y = y_H + y_P = Ae^{2x} + Be^{3x} + 2x - 1$$

where A and B are real constants, also solves the ODE. In fact, this gives the general solution (the discussion in Subsection 3.8.4 explains why). $\square$

Bearing in mind the above example, we now detail an algorithm for solving a second order ODE of the form (3.35).

1. Find the solution y_H to the corresponding homogeneous equation (3.36) (by first identifying the roots of the characteristic equation, as in Subsection 3.8.1).
2. Find a particular solution y_P to (3.35).
3. The general solution y to (3.35) is then given by $y = y_H + y_P$.

In general, it is best to perform Step 1 before Step 2; the reason for this will soon become obvious.

Step 1 has already been discussed in some detail, and Step 3 is easy. We need only devote some discussion to performing Step 2.

Example 3.8.9. Solve the ODE

$$y'' - 4y' + 5y = 20e^{-x}. \quad (3.44)$$

Proof. First, we solve the homogeneous equation

$$y'' - 4y' + 5y = 0.$$

The roots of the characteristic equation $\lambda^2 - 4\lambda + 5 = 0$ are $2 - i$ and $2 + i$ (these can be found using the quadratic formula). Hence the solution y_H of the homogeneous equation is given by

$$y_H = e^{2x}(A \cos x + B \sin x),$$

where A and B are real numbers.

Second, we find a particular solution y_P . Since the right hand side is an exponential, we try

$$y_P = ae^{-x},$$

where a is a constant to be determined. Note that $y_P' = -ae^{-x}$ and $y_P'' = ae^{-x}$. So substituting y_P into (3.44) gives

$$ae^{-x} + 4ae^{-x} + 5ae^{-x} = 20e^{-x}.$$

By comparing coefficients on each side we conclude that y_P is a particular solution if and only if $a = 2$. Hence our particular solution is given by $y_P = 2e^{-x}$.

Third, the general solution y to (3.44) is given by

$$\begin{aligned} y &= y_H + y_P \\ &= e^{2x}(A \cos x + B \sin x) + 2e^{-x}, \end{aligned}$$

where A and B are real numbers. □

Example 3.8.10. Solve the ODE

$$y'' - 3y' + 2y = 5e^{2x}. \quad (3.45)$$

Solution. First we solve the corresponding homogeneous ODE

$$y'' - 3y' + 2y = 0.$$

The characteristic equation $\lambda^2 - 3\lambda + 2 = 0$ has the solutions $\lambda = 1$ and $\lambda = 2$. Hence the solution y_H to the homogeneous equation is given by

$$y_H = Ae^{2x} + Be^x, \quad (3.46)$$

where A and B are real constants.

Second, we look for a particular solution y_P . Since the right-hand side of (3.45) is a multiple of e^{2x} , it seems natural to try $y_P = ae^{2x}$. However, note that ae^{2x} is a particular solution to the homogeneous equation. (To see this, simply set A as a and B as 0 in (3.46)). Hence substituting $y_P = ae^{2x}$ into (3.45) will produce 0 on the left-hand side, and consequently this guess for y_P will *not* work.

In this circumstance, the ‘trick’ for finding a particular solution y_P is to *multiply the old guess* ae^{2x} by x . That is, we now try

$$y_P = axe^{2x}.$$

(This guess for a particular solution is certainly not in the solution space for the homogeneous equation.) Now we find that

$$y'_P = ae^{2x}(2x + 1) \quad \text{and} \quad y''_P = 4ae^{2x}(x + 1),$$

and hence substituting $y_P = axe^{2x}$ into (3.45) gives

$$4ae^{2x}(x + 1) - 3ae^{2x}(2x + 1) + 2axe^{2x} = 5e^{2x}.$$

This simplifies to

$$4ae^{2x} - 3ae^{2x} = 5e^{2x},$$

and we thereby deduce that $a = 5$. Therefore a particular solution y_P is given by $y_P = 5xe^{2x}$.

Finally, the general solution y to (3.45) is given by

$$\begin{aligned} y &= y_H + y_P \\ &= Ae^{2x} + Be^x + 5xe^{2x} \end{aligned}$$

where A and B are real constants. □

Given a function f , the following table indicates which guess for y_P will always yield a particular solution for the nonhomogeneous ODE (3.35).

$f(x)$	Guess for particular solution y_P
$P(x)$ (a polynomial of degree n)	$Q(x)$ (a polynomial of degree n)
$P(x)e^{sx}$	$Q(x)e^{sx}$
$P(x)\cos(sx)$	$Q_1(x)\cos(sx) + Q_2(x)\sin(sx)$
$P(x)\sin(sx)$	$Q_1(x)\cos(sx) + Q_2(x)\sin(sx)$
$P(x)e^{sx}\cos(tx)$ or $P(x)e^{sx}\sin(tx)$	$Q_1(x)e^{sx}\cos(tx) + Q_2(x)e^{sx}\sin(tx)$
If any term of the guess for y_P is a solution to the homogeneous ODE, then multiply it by x. If any term of the new guess is still a solution to the homogeneous ODE, then multiply by x again.	

The next example illustrates the directive given in the last two rows of the table.

Example 3.8.11. Solve the ODE

$$y'' - 6y' + 9y = 8e^{3x}. \quad (3.47)$$

Solution. First, the characteristic equation

$$\lambda^2 - 6\lambda + 9 = 0$$

factorises as $(\lambda - 3)^2 = 0$ and thus has the repeated root 3. So the solution y_H to the corresponding homogeneous equation is given by

$$y_H = Ae^{3x} + Bxe^{3x}, \quad (3.48)$$

where A and B are real numbers.

Second, we search for a particular solution y_P . Our first guess $y_P = ae^{3x}$ will not work, as it is a solution to the corresponding homogeneous equation (to see why, simply set A as a and B as 0 in (3.48)). So we multiply by x to obtain a new guess $y_P = axe^{3x}$. However, this guess also solves the homogeneous equation. So once again, multiply by x to obtain a new guess $y_P = ax^2e^{3x}$. It is easy to see that y_P no longer lies in the solution space to the homogeneous equation. So this guess will work.

To determine the value of a , substitute $y_P = ax^2e^{3x}$ into (3.47). We leave it to the reader to verify that $a = 4$. Hence a particular solution y_P is given by $y_P = 4x^2e^{3x}$.

Finally, the general solution y is the sum of y_H and y_P , namely

$$y = Ae^{3x} + Bxe^{3x} + 4x^2e^{3x},$$

where A and B are real numbers. □

Remark 3.8.12. The previous example illustrates the importance of finding y_H before making a guess for y_P ; without knowing y_H it is not possible to make a suitable guess for y_P . The next example emphasises this point.

Example 3.8.13. Consider the nonhomogeneous ODE

$$\frac{d^2y}{dt^2} + 6\frac{dy}{dt} + 13y = 5e^{3t} \cos(2t). \quad (3.49)$$

Write down the *form* of a particular solution y_P to this ODE. (You are *not* required to evaluate the undetermined coefficients appearing in the the form of y_P .)

Solution. The characteristic equation $\lambda^2 + 6\lambda + 13 = 0$ has roots $3 + 2i$ and $3 - 2i$ (these can be found by completing the square or using the quadratic formula). Hence the solution y_H to the corresponding homogeneous equation is given by

$$y_H = e^{3t}(A \cos 2t + B \sin 2t), \quad (3.50)$$

where $A, B \in \mathbb{R}$.

Since the right-hand side of (3.49) is a product of e^{3t} and $\cos 2t$, our initial guess for the particular solution y_P is given by

$$y_P = ae^{3t} \cos 2t + be^{3t} \sin 2t.$$

However, (3.50) shows that this guess solves the homogeneous equation. Instead, try

$$y_P = ate^{3t} \cos 2t + bte^{3t} \sin 2t.$$

This new guess for y_P does not lie in the solution space for the homogeneous equation; hence this gives the form of particular solution that we seek. $\square$

Remark 3.8.14. One must take care in the use of the method of undetermined coefficients, and in particular the two rules at the end of the table given above.

Consider, for example, the differential equation

$$y'' - 6y' + 9y = x^2e^{3x}.$$


The homogeneous solution, from above, is $y_H = Ae^{3x} + Bxe^{3x}$. Now since x^2e^{3x} is **not** one of the homogeneous solutions, one would try the particular solution $y_P = (Cx^2 + Dx + E)e^{3x}$. However, since this contains terms which are part of the homogeneous solution, we need to multiply by x twice - thus $y_P = (Cx^4 + Dx^3 + Ex^2)e^{3x}$ - so that no term in this expression is a homogeneous solution. Substitution will reveal that D and E are zero, while $C = \frac{1}{12}$.

There is an easier method to solve such a problem using differential operators. This is left to more advanced courses in differential equations.

3.8.3 An application: vibrations and resonance

Many structures have a natural frequency of vibration. If an external agent causes them to vibrate at or near one of these frequencies then large oscillations build up and *resonance* occurs. This can cause disasters such as the collapse of bridges. The ‘trick’ used by singers breaking wine glasses is another example of this phenomenon. The first example of this subsection examines the behaviour of a natural oscillating system; the second example illustrates how introducing external vibrations into such a system causes resonance.

Example 3.8.15. Suppose that a spring is mounted to a (fixed) point P and that an object of mass m is suspended from the spring. Let x denote the (vertical) displacement from the equilibrium position (or resting position) of the object, taking x to be positive if it lies above the equilibrium position.



By using Newton's second law of motion, Hooke's law and making some simple assumptions, one can show that the system satisfies the differential equation

$$\frac{d^2x}{dt^2} + \omega^2 x = 0, \quad (3.51)$$

where ω is a positive constant that depends only on the mass of the object and the stiffness of the spring.

The object is pulled downwards from its equilibrium position by a distance of 4 units and then released from rest. Find $x(t)$ when $t \geq 0$.

Solution. We need to solve the differential equation (3.51) subject to the initial conditions $x(0) = -4$ and $x'(0) = 0$. The characteristic equation $\lambda^2 + \omega^2 = 0$ is easily solved, leading to the solution

$$x(t) = A \cos \omega t + B \sin \omega t$$

of (3.51), where A and B are constants. Before imposing initial conditions, we calculate $x'(t)$. This is given by

$$x'(t) = -A\omega \sin \omega t + B\omega \cos \omega t.$$

Now $x'(0) = 0$ implies that $0 = B\omega$, from which we conclude that $B = 0$ (since ω is positive). Finally, $x(0) = -4$ implies that $A = -4$. Hence the solution is given by

$$x(t) = -4 \cos \omega t.$$

This type of motion is known as *simple harmonic motion* and is graphed in Figure 3.2 (a). $\square$

Example 3.8.16. Consider the same scenario as Example 3.8.15, except that now the point P vibrates up and down, such that its vertical displacement y is given by $y = 2 \sin \Omega t$. In these circumstances, a simple physical argument shows that x obeys the differential equation


$$\frac{d^2x}{dt^2} + \omega^2 x = 2 \sin \Omega t. \quad (3.52)$$

Describe the motion of the object, given that $x(0) = -4$ and $x'(0) = 0$.

Solution. We have already solved the homogeneous equation in Example 3.8.15. So we look for a particular solution x_P .

Case 1: Suppose that $\Omega \neq \omega$. Then we look for a particular solution of the form

$$x_P = C \cos \Omega t + D \sin \Omega t.$$



(a) Simple harmonic motion



(b) Stable oscillation: $\omega \neq \Omega$ (c) Resonance: $\omega = \Omega$

Figure 3.2: Different oscillating systems.

$$x_P = \frac{2}{\omega^2 - \Omega^2} \sin \Omega t.$$

By using the fact that $x = x_H + x_P$ and imposing initial conditions we find that

$$x(t) = -4 \cos \omega t - \frac{\Omega}{\omega} \frac{2}{\omega^2 - \Omega^2} \sin \omega t + \frac{2}{\omega^2 - \Omega^2} \sin \Omega t.$$

This is simply another oscillating system. Its graph is shown in Figure 3.2 (b) (in the case when $\Omega = \frac{3}{2}\omega$).

Case 2: Suppose that $\Omega = \omega$. This time we look for a particular solution of the form

$$x_P = Ct \cos \omega t + Dt \sin \omega t.$$

By substituting this into (3.52) we obtain

$$x_P = -\frac{t}{\omega} \cos \omega t.$$

Using the fact that $x = x_H + x_P$ and imposing initial conditions we find that

$$x(t) = -4 \cos \omega t + \frac{1}{\omega^2} \sin \omega t - \frac{t}{\omega} \cos \omega t.$$

Hence as t increases, the amplitude of the $\cos \omega t$ term grows without bound and the system becomes unstable. Its graph is shown in Figure 3.2 (c). $\square$

3.8.4 A connection with linear algebra

In Section 3.8, we have made two claims whose proofs have not been given yet; namely (A) that every second order homogeneous ODE has at most two linearly independent solutions and (B) that every solution y to a nonhomogeneous second order ODE can be written in the form $y_H + y_P$. Both of these assertions can be proved by using linear algebra. In this subsection we highlight the connection between second order linear ODEs and linear algebra, and explain why assertion (B) is true. The proof of assertion (A) is given in MATH2501. Students should not continue reading this subsection until Chapter 7 from MATH1231/1241 algebra has been completed.

Consider the second order ODE

$$y'' + ay' + by = f, \tag{3.53}$$

where a and b are real numbers and $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function. Let V denote the vector space of all (infinitely) differentiable functions $y : \mathbb{R} \rightarrow \mathbb{R}$. We define the linear transformation $T : V \rightarrow V$ by the formula

$$T(y) = y'' + ay' + by.$$

Observe that

$$\begin{aligned} y_H \text{ is a solution to the homogeneous equation} &\iff y_H'' + ay_H' + by_H = 0 \\ &\iff T(y_H) = 0 \\ &\iff y_H \in \ker(T). \end{aligned}$$

Hence the general solution to the homogeneous equation *is* the kernel of T . Also observe that

$$\begin{aligned} y_P \text{ is a particular solution to the nonhomogeneous equation} &\iff y_P'' + ay_P' + by_P = f \\ &\iff T(y_P) = f. \end{aligned}$$

Hence the ODE (3.53) has a particular solution if and only if f is in the image of T . Moreover, since

$$\begin{aligned} T(y_H + y_P) &= T(y_H) + T(y_P) && \text{(by the linearity of } T) \\ &= 0 + f \\ &= f, \end{aligned}$$


we conclude that $y_H + y_P$ is also a solution to (3.53).

Finally, we prove assertion (B). Suppose that y is a solution to (3.53) and that y_P is some particular solution to (3.53). Then $T(y) = f$ and $T(y_P) = f$. Hence

$$\begin{aligned} T(y - y_P) &= T(y) - T(y_P) && \text{(by the linearity of } T) \\ &= f - f \\ &= 0. \end{aligned}$$

Therefore $y - y_P$ is in the kernel of T . But, as observed above, every function in the kernel of T is a solution y_H to the homogeneous equation. Hence $y - y_P = y_H$, where y_H is some solution to the homogeneous equation. It follows that $y = y_H + y_P$, thus completing the proof of assertion (B).

The kernel and image of T , and their connection to the homogeneous solution space and a particular solution for the differential equation, may be represented pictorially as shown below.



3.9 Maple notes

The following MAPLE command is relevant to the material of this chapter:

`dsolve(deqn, y(x));` solves the ordinary differential equation (or IVP) `deqn` for the function `y(x)`. For example,

```
> dsolve(diff(y(x), x$2) - y(x) = 1, y(x));
```

$$y(x) = -1 + _C1 \exp(x) + _C2 \exp(-x)$$

```
> dsolve({diff(v(t), t) + 2*t = 0, v(1)=5}, v(t));
```

$$v(t) = -t^2 + 6$$

Problems 3.3 : Separable ODEs

1. [R] Solve the following differential equations.

$$\begin{array}{ll}
 \text{a) } \frac{dy}{dt} = t^2(1 + y^2) & \text{b) } \frac{dy}{dx} = xy^2 \\
 \text{c) } \frac{dy}{dx} = \frac{\sin x}{y^2} & \text{d) } \frac{dy}{dx} = \frac{y}{x(x-1)} \\
 \text{e) } \frac{dy}{dx} = e^{x+y} & \text{f) } y \cos^2 x \frac{dy}{dx} = \tan x + 2, \quad y = 2 \text{ when } x = \pi/4 \\
 \text{g) } x \frac{dy}{dx} = y \ln x & \text{h) } \frac{dy}{dx} = 3x^2y^2, \quad \text{given that } y = 1 \text{ when } x = 0.
 \end{array}$$

2. [H] Try to find the general solution for $\frac{dy}{dx} = 3y^{2/3}$.

Your answer will probably be $y = (x + C)^3$.

Observe $y = 0$ is also a solution and it cannot be expressed as $y = (x + C)^3$ for any value of C .

How do you account for this? Are there any other solutions?

3. [R] Oil is leaking out of a tank in such a way that the depth of oil h in the tank at time t satisfies $\frac{dh}{dt} = -\sqrt{2h}$. If the initial height is 4 cm, then find the time taken for the tank to empty.

Problems 3.4 : First order linear ODEs

4. [R] Solve the following linear ODEs.

$$\begin{array}{ll}
 \text{a) } \frac{dy}{dx} - 2y = x^2e^{2x} & \text{b) } \frac{dy}{dx} + 3y = \frac{e^{-3x}}{1+x^2} \\
 \text{c) } x \frac{dy}{dx} + (1+x)y = 2 & \text{d) } x \frac{dy}{dx} - 2y = 6x^5 \\
 \text{e) } \cos^2 x \frac{dy}{dx} + y = \tan x & \text{f) } \frac{dy}{dx} = x + 2y \tan 2x
 \end{array}$$

5. [R] Solve $\frac{dx}{dt} = t - x$. Sketch the solution curves passing through $(0, 1)$, $(0, 0)$, $(0, -1)$ and $(0, -2)$.

6. [R] An object falling vertically experiences a resistance which is proportional to its velocity.

- Explain why its acceleration is given by $\frac{dv}{dt} = g - kv$, where g is the acceleration due to gravity and k is a positive constant.
- Solve this as a linear equation for the initial condition $v = 0$.

- c) Solve this as a separable equation.
 d) What is the terminal (or limiting) velocity of the object?

Problems 3.5 : Exact ODEs and miscellaneous first order ODEs

7. [R] Solve the following exact ODEs.
 a) $2xy + (x^2 + y^2)\frac{dy}{dx} = 0$ b) $(\sin y - xy^2)dx + (x \cos y - x^2y)dy = 0$
8. [R] Determine which of the following differential equations are exact and solve those which are.
 a) $(2xe^y + e^x) + (x^2 + 1)e^y \frac{dy}{dx} = 0$
 b) $y(x^2 + \ln y) + x \frac{dy}{dx} = 0$
 c) $\left(\frac{y}{1+x^2} + e^y\right)dx + (\tan^{-1}x + xe^y)dy = 0$
 d) $e^{xy}(y \cos x - \sin x) + xe^{xy} \cos x \frac{dy}{dx} = 0$
9. [R] (*Miscellaneous first order ODEs*)
 Solve the following ODEs.
 a) $(2xy - 3 \tan x) \frac{dy}{dx} = 3y \sec^2 x - y^2$
 b) $x - (x^2y + y) \frac{dy}{dx} = 0$
 c) $x^2 \frac{dy}{dx} + xy = 1, x > 0$
 d) $x^2 \frac{dy}{dx} - xy = y$
 e) $x \frac{dy}{dx} = \frac{1+x^2}{y^2}, y(1) = 3.$
10. [X] Solve the differential equation

$$(2x - 10y^3) \frac{dy}{dx} + y = 0$$

by first multiplying it through by some function $\mu(y)$ to make it exact.

Problems 3.6 : Solving ODEs by using a change of variable [X]

11. [X] Solve the following ODEs by using the substitution $y(x) = x \cdot v(x)$.
 a) $\frac{dy}{dx} = \frac{xy - y^2}{x^2}$ b) $\frac{dy}{dx} = \frac{y}{x} + \cos\left(\frac{y-x}{x}\right)$
 c) $\frac{dy}{dx} + \frac{2xy}{x^2 + y^2} = 0$ d) $\frac{dy}{dx} + \frac{x^2 + y^2}{2xy} = 0$
 e) $x \frac{dy}{dx} = \sqrt{x^2 + y^2} + y, y(1) = 0.$

12. [X] Use the substitution $y = x \cdot v(x)$ to find the solution to the differential equation

$$\frac{dy}{dx} = \frac{y-x}{y+x}, \quad y(1) = 0.$$

Use polar coordinates to sketch the corresponding curve.

13. [X] Solve $\frac{dy}{dx} = (y-x)^2$ using the substitution $u = y-x$.

14. [X]

- a) Solve $\frac{dy}{dt} = y - 2y^2$ by making the substitution $z = \frac{1}{y}$ to obtain a linear differential equation in z and t .
 b) Use the same trick to solve

$$\frac{dy}{dt} = 5y - ty^2 \quad \text{given that } y = 1 \quad \text{when } t = 0.$$

Find the maximum value of y when $t \geq 0$.

15. [X] Solve the following differential equations by using the given substitution.

- a) $y \frac{dy}{dx} - 2y^2 = x; \quad u = y^2$.
 b) $x \frac{dy}{dx} + y(x^2 + \ln y) = 0, \quad x > 0, \quad y > 0; \quad u = \ln y$.
 c) $xyy'' = yy' + x(y')^2; \quad u = y'/y$ or $u = \ln y$.

16. [X] In each case, use the substitution $v = \frac{dy}{dt}$ to solve the given ODE.

- a) $\frac{d^2y}{dt^2} + 2 \frac{dy}{dt} = 6$
 b) $t^2 \frac{d^2y}{dt^2} - \left(\frac{dy}{dt} \right)^2 = 0$

17. [X] Suppose that $\alpha(x)$ is a solution to the equation

$$\frac{d^2u}{dx^2} + b(x) \frac{du}{dx} + c(x)u = 0.$$

- a) Use the substitution $u(x) = \alpha(x)v(x)$ to write down a first order equation for $\frac{dv}{dx}$. What method can you see to solve this equation?
 b) Verify that $u(x) = x^3$ is one solution to the differential equation

$$x^2 \frac{d^2u}{dx^2} - 4x \frac{du}{dx} + 6u = 0.$$

Hence use the technique of (a) to find the general solution.

Problems 3.7 : Modelling with first order ODEs

18. [R] The simple population growth model, $\frac{dy}{dt} = ky$, is (because of limitations on resources, pollution, ...) unsatisfactory over a 'long' period. We might look at

$$\frac{dy}{dt} = k(y)y \quad (1)$$

(which of course ignores seasonal and other variations with time). Mathematically one of the simplest assumptions we can make (Verhulst 1839) is that $k(y)$ decreases *linearly* as y increases. In this case (1) may be written in the form

$$\frac{dy}{dt} = k \left(1 - \frac{y}{K}\right) y, \quad (2)$$

where k and K are constants.

- a)
 - i) Equation (2) has two constant (stationary) solutions. What are they?
 - ii) Solve (2), given that $y = y_0$ when $t = 0$ and $0 < y_0 < K$.
 - iii) State what happens to y as $t \rightarrow \infty$.
 - iv) For what value of y is the rate of increase of y a maximum?
(Caution — there is a simple, two line, method.)
 - v) If $y_0 > K$ then what happens to y as t increases?
- b) [H] It has been found empirically that for certain bodies $k(y)$ decreases linearly with $\ln y$. (Here y could be the volume, mass or number of cells of the body.) In this case (1) may be written as

$$\frac{dy}{dt} = \alpha \ln \left(\frac{K}{y} \right) y. \quad (3)$$

Solve this differential equation, given that $y = y_0$ when $t = 0$.

19. [R] An initially unpolluted lake of 10^9 litres has a river flowing through it at 1,000,000 litres per day. A factory is built which discharges 10,000 litres per day of pollutant into the lake. Assume that the total volume of liquid in the lake remains constant at 10^9 litres.
- a) What will be the eventual level of pollution in the lake?
 - b) How long will it take to reach half this level?
 - c) Is there anything unrealistic about your model?
20. [R] A tank can hold 100 litres. Initially it holds 50 litres of pure water. Brine, which contains 2 grams of salt per litre, is run in at the rate of 3 litres per minute. The mixture, which is stirred continuously, is run off at 1 litre per minute. Let $x(t)$ denote the mass of salt (in grams) present in the tank after t minutes.
- a) Set up a differential equation in x and t to model the system.
 - b) Show that when the tank is at the point of overflowing it contains $50(4 - \sqrt{2})$ grams of salt.
21. [R] A population of size P is subject to seasonal variation. The population has a growth rate given by

$$\frac{dP}{dt} + P = 100 + 50 \sin t.$$

- a) Solve the differential equation given that $P(0) = 20$.
- b) Find the average population size over a long period of time.

22. [R] The equation

$$\frac{dy}{dt} = k(1 + a \cos(2\pi t))y.$$

is an attempt to model seasonal variation. Solve it.

23. [R] It is estimated that the population growth rate of a certain developing country will fall linearly from 2% per year to 1.5% per year over the next decade.

- a) Express the population growth rate as a function of time.
- b) Given that the present population is 10 million, find the estimated population in 10 years time.

24. [R] An object falling in a resisting medium has a constant acceleration due to gravity of 9.8 m/sec^2 and also a drag force, which is approximately proportional to the speed (see Question 6). For a stone falling in water with velocity $v \text{ m/sec.}$, the acceleration from this drag force is approximately $10v \text{ m/sec}^2$.

- a) Given that the stone is dropped from rest at the water surface, write a differential equation describing this situation, and solve it to find v .
- b) Determine the terminal velocity, $\lim_{t \rightarrow \infty} v(t)$.
- c) How long does it take before the stone is travelling with 95% of its terminal velocity?
- d) What if it starts at a velocity higher than the terminal velocity?

25. [R] The differential equation $\frac{dy}{dx} = \lambda \frac{y}{x}$ is employed as a simple model for comparative growth.

- a) Solve this differential equation.
- b) The mass y of the large claw of a fiddler crab is compared to the mass x of its body (without claw) over a period of time. The measurements taken (in grams) are recorded in the following table.


x	55	300	536	1080	1449	2233
y	5	72	175	522	773	1498

By graphing $f(x)$ against $f(y)$ for a suitable function f , show that the above model appears reasonable in this example. Determine approximately the value of λ and the constant of integration.

26. [R] An investor puts \$500,000 into Hitek Bonds, which pay a profit of 20% a year, compounded daily (i.e., if P is the amount of money in the bonds at any time, $\frac{dP}{dt} = 0.2P$ approximately).

- a) How much money would he own at the end of a year?

- b) [H] His tax accountant advises that this is overprofitable, and suggests that he use \$200 a week of the interest towards the hire of a Volvo, which would be a tax write-off under the government's Small Business Incentive Scheme, and that at the end of each six months, all the remaining interest should be taken out and invested in the Lake Eyre Ricegrowers' Cooperative, which loses money continuously at a rate of 10% a year (so that $\frac{dP}{dt} = -0.1P$). This would qualify the investor for various advantages under the Rural Rorts Scheme. If the investor takes this advice, what would the total of both his investments be after one year? [Take it that there are 52 weeks in a year.]
27. [H] There are $n + 1$ tanks each containing 100 litres and connected as shown.



Throughout, the liquid in every tank is kept well-mixed. The 0-th tank contains 100 litres of pure water with 50 grams of salt dissolved in it. The remainder contain pure water. Pure water is pumped into tank 0 at 3 litres per minute and liquid leaves the system from tank n at 3 litres per minute. Let m_k denote the mass (in grams) of salt in tank k .

- a) Show that

$$\frac{dm_k}{dt} = 0.03(m_{k-1} - m_k), \quad k = 1, 2, \dots, n$$

and

$$\frac{dm_0}{dt} = -0.03m_0.$$

- b) Show that

$$m_n = \frac{50(0.03)^n}{n!} t^n e^{-0.03t}.$$

28. [H] The decay of one atom of radioactive element A yields one atom of B which is itself radioactive. The decay constants for A and B are 0.25 and 2 (per day) respectively. A pure sample of K atoms of A is placed in a closed container. Let $y_1(t)$ and $y_2(t)$ denote the numbers of atoms of A and B that are present at time t . Write down two differential equations to describe this situation. Solve to obtain a formula for y_2 . When is the amount of B in the container a maximum?
29. [X] A cyclist freewheeling on a level road experiences a retardation (i.e., a negative acceleration) that is proportional to the square of his speed. His speed is reduced from 20 metres per second to 10 metres per second in a distance of 100 metres. Find his average (with respect to time) speed during this period.

30. [R] Find the general solutions of the following second order ODEs.

- a) $y'' + 3y' + 2y = 0$ b) $y'' + 2y' + 10y = 0$
 c) $y' + 3y = 0$ d) $y'' + 4y' + 4y = 0$

31. [R] Find the solutions of the following initial value problems.

- a) $y'' - 6y' + 5y = 0$; $y = 1$, $y' = 0$, when $x = 0$.
 b) $y'' + 2y' + 2y = 0$; $y = 1$, $y' = 0$, when $x = 0$.

32. [R] Solve the following differential equations.

- a) $y'' + 4y' + 3y = x$ b) $y'' - 6y' + 9y = 5e^{2x}$
 c) $y'' + 2y' + 2y = 10 \cos 2x$ d) $y'' - y = e^{-x}$
 e) $y'' + 4y = \sin 2x$ f) $y' + 3y = 10e^{2x}$
 g) $y'' + 4y = \sin x$ $y = 0$, $y' = 1$ when $x = 0$
 h) $y'' - 5y' + 4y = 2e^{2x}$ $y = 1$, $y' = 3$ when $x = 0$

33. [R] For each of the following differential equations, find the general solution of the associated homogeneous equation and write down the form of the particular solution you would seek. (Do not evaluate the unknown coefficients.)

- a) $2y'' - 3y' - 5y = (x + 4)e^{5x/2}$
 b) $y'' + 2y' - 24y = e^{4x} \sin 6x$
 c) $y'' + 6y' + 9y = e^{-3x}$

34. [H] Find a particular solution for the differential equation $y'' - 4y' + 4y = 6x^2e^{2x}$.

35. [X] By making the substitution $x = e^t$, reduce the linear differential equation

$$x^2 \frac{d^2 y}{dx^2} - 4x \frac{dy}{dx} + 6y = x^5, \quad x > 0,$$

to one with constant coefficients, and hence solve it.

36. [X]

- a) Show that the second order differential equation

$$y'' + (m_1 + m_2)y' + m_1m_2y = 0,$$

where m_1 and m_2 are constants, can be written in the form

$$\frac{d}{dx}(y' + m_1y) + m_2(y' + m_1y) = 0.$$


- b) Suppose that $m_1 = m_2 = m$. Using the result of (a), solve the first differential equation by solving successively two first order linear equations.

37. [R] A cylindrical buoy of 80 cm in diameter floats in water with its axis vertical. When depressed slightly and released, it bobs up and down according to the differential equation

$$m \frac{d^2 x}{dt^2} = -\pi 40^2 g x,$$

where m is the mass (in grams) of the buoy and x is the displacement (in centimetres) from the equilibrium position. Take the acceleration g due to gravity to be 980 cm/sec^2 . The period of oscillation is observed to be 2.5 seconds. What is the mass of the buoy?

38. [R] A block of wood is attached to a spring and slides across a surface as shown.



Let $x(t)$ denote the horizontal distance of the block from its 'resting' position at time t . The motion of the block is modelled by the initial value problem

$$\frac{d^2 x}{dt^2} + c \frac{dx}{dt} + 4x = 0, \quad x'(0) = 0, \quad x(0) = 1,$$

where c is the 'coefficient of friction' between the block and the surface.

- a) By solving the differential equation, describe the motion of the block in the 'ideal' situation when there is no friction between the block and the surface (that is, when $c = 0$).
 - b) Solve the initial value problem when $c = 2$ and when $c = 5$ and explain why the block oscillates if $c = 2$ but does not if $c = 5$.
 - c) [H] Find the smallest positive c such that the system does not oscillate. (This value of c corresponds to what is known as 'critical damping'.)
39. [R] A circuit consists of an inductor and capacitor connected in series with a sinusoidal power source. The charge q (in coulombs) stored in a capacitor is given by the differential equation

$$\frac{d^2 q}{dt^2} + 10\,000q = 1000 \sin \Omega t,$$

where $\Omega/(2\pi)$ is the frequency of the power source.

- a) Find the solution to the corresponding homogeneous equation.
- b) Write down the form of particular solution you would seek for the differential equation. (Warning: there are two cases, depending on the value of Ω .)
- c) What sinusoidal frequency leads to unbounded oscillatory behaviour (known as 'resonance')?

40. [R]

- a) Find the general solution of the vibrating system modelled by the equation

$$\frac{d^2 y}{dt^2} + 3 \frac{dy}{dt} + 2y = 20 \sin t.$$

- b) The long term behaviour of this system is independent of the initial conditions. What is this ‘steady state solution’?

41. [X]

- a) Solve the previous question by first finding a particular complex-valued function
- $z(t)$
- satisfying

$$\frac{d^2 z}{dt^2} + 3 \frac{dz}{dt} + 2z = 20e^{it}.$$

- b) Similarly, find a particular solution of

$$\frac{d^2 y}{dt^2} + 4y = \sin 2t$$

by considering

$$\frac{d^2 z}{dt^2} + 4z = e^{2it}.$$

42. [R] A stationary wave on a guitar string of length
- L
- can be (partially) modelled by the boundary value problem

$$y'' = ky; \quad y(0) = 0, \quad y(L) = 0, \quad L > 0,$$

where k is a constant. Assume that y is not identically zero.

- a) Suppose that k is positive and write k as μ^2 , where $\mu \geq 0$. Show that the boundary value problem has no nonzero solutions.
- b) Are there any nonzero solutions when $k = 0$?
- c) Suppose now that k is negative and write k as $-\mu^2$, where $\mu > 0$. Find all possible values of μ such that the boundary value problem has a nonzero solution. Give the corresponding solutions.
(Each such solution corresponds to a natural harmonic of the string.)

43. [X]

- a) If
- g
- is a given function which is continuous and
- positive*
- on the interval
- $[0, L]$
- , show that the only solution of the boundary value problem

$$y'' - g(x)y = 0; \quad y(0) = 0, \quad y(L) = 0, \quad L > 0,$$

is $y = 0$.

(Hint: Give a proof by contradiction using the maximum-minimum theorem for continuous functions.)

- b) Find all possible values for λ such that there is a solution to the differential equation

$$y'' + 2y' + 2\lambda y = 0$$

satisfying $y(0) = y(L) = 0$ and y not identically zero.

44. [H] A simple linear predator-prey model:

Let X denote the number of predators at time t and Y the number of prey. Experience and theoretical considerations suggest that

- (α) there is an equilibrium state $(X, Y) = (X_0, Y_0)$ with $X_0, Y_0 \neq 0$;
 (β) an increase in the number of prey, from Y_0 , results in an increase in the number of predators at a rate (approximately) proportional to the increase in prey, so that

$$\frac{dX}{dt} = a(Y - Y_0), \quad a > 0; \quad (1)$$

and

- (γ) an increase in the number of predators, from X_0 , results in a decrease in the number of prey at a rate (approximately) proportional to the increase in predators, so that

$$\frac{dY}{dt} = -b(X - X_0), \quad b > 0. \quad (2)$$

[In fact the ‘linearisation’ of a number of complicated models near an equilibrium state leads to (1) and (2).]

If $x = X - X_0$ and $y = Y - Y_0$ then we obtain

$$\frac{dx}{dt} = ay, \quad (3)$$

and

$$\frac{dy}{dt} = -bx. \quad (4)$$

- a) Eliminate dt between (3) and (4) and solve to obtain a relation between x and y (the ‘phase trajectories’ of the system).
 b) Eliminate x by differentiating (4) and substituting from (3). Hence solve for y and then use (4) again to obtain x .
 c) Suppose that $a = 0.8$ and $b = 3.2$. If $x = -1.2$ and $y = 3.2$ when $t = 0$, then find x and y in terms of t and also find the phase trajectory.
45. [X] (*Note: do not attempt this question until Chapter 7 of MATH1241 Algebra has been completed.*)

Let V denote the vector space of twice differentiable functions on $\mathbb{R}$. Define a linear map L on V by the formula

$$Lu = a \frac{d^2 u}{dx^2} + b \frac{du}{dx} + cu, \quad \text{where } a, b \text{ and } c \text{ are real numbers.}$$

Suppose that u_1, u_2 is a basis for the solution space of $L(u) = 0$. Find a basis for the solution space of the fourth order equation $L(L(u)) = 0$. What can you say about the kernels of L and L^2 ?

Chapter 4

Taylor series

Polynomials are nice functions to work with. Their values can be easily evaluated using a finite number of additions and multiplications. They are easy to differentiate and integrate; moreover their derivatives and antiderivatives are also polynomials and thus have these properties. Some (but not all) of these properties are shared by a few other classes of functions. For example, the derivatives of exponential functions are exponentials, but in general exponentials cannot be evaluated using a finite number additions and multiplications. Many other useful functions share none of these properties.

If a function can be accurately approximated by a polynomial, then we can use the polynomial to approximate the values, derivatives and antiderivative of the function. This generalises an idea met in MATH1131, where we saw that a differentiable function can be locally approximated by a linear function (which is a degree 1 polynomial). In this chapter we will see that many n -times differentiable functions can be locally approximated by a polynomial of degree n . We will devote considerable time discussing how accurate such approximations are. Finally, we shall prove that some functions can not only be approximated by polynomials but are also equal to series consisting of infinitely many polynomial terms. Such series are known as *Taylor series*.

The ideas presented in this chapter have a long history, going back to Archimedes' *method of exhaustion*, which he used to approximate π . Beginning in the fourteenth century, a school of Indian mathematicians based in Kerala found accurate polynomial approximations to trigonometric functions. This allowed them to solve problems in astronomy. In the seventeenth century, the Scottish astronomer James Gregory independently employed similar techniques. However, it was not until 1715 that the English mathematician Brook Taylor published a theorem which gave a general method for polynomial approximation, and which described precisely the errors involved. In the twenty-first century, computers and calculators regularly use algorithms to find the approximate value of functions; many of these modern computation techniques trace their roots back to Taylor's method.

4.1 Taylor polynomials

(Ref: SH10 §12.6, 12.7)

Recall from MATH1131 that the function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = e^x$, is defined as the inverse of the function $\ln : (0, \infty) \rightarrow \mathbb{R}$, and that $\ln$ is defined in terms of an integral. How, then, does one evaluate $e^{0.1}$? (If the answer given is 'use a calculator,' then we merely ask the question, How does a calculator evaluate $e^{0.1}$?)

One approach is to suppose that $y = e^{0.1}$. Then we need to solve the equation $\ln y = 0.1$. By the definition of $\ln$, this boils down to solving the integral equation

$$\int_1^y \frac{dt}{t} = 0.1.$$

From here we could guess an approximate value for y and use Riemann sums to check whether our guess is reasonable. Clearly this is an unsatisfactory approach to the problem.

Another method is to locally approximate the function f with a linear function. This technique was discussed in MATH1131 (see Chapter 4 of the MATH1131 calculus notes) and is based on the idea that the tangent lies close to the graph of f near the point of contact. Suppose, once again, that $f(x) = e^x$. To find an approximate value for $e^{0.1}$, we will approximate f using the tangent to the graph of f at 0. The tangent function p_1 at 0 is a polynomial of degree 1 and is given by

$$p_1(x) = 1 + x.$$

So when x is close to 0,

$$e^x \approx 1 + x$$

and hence $e^{0.1} \approx 1.1$.

The polynomial p_1 has the property that its value at 0 and gradient at 0 agree with the value and gradient of f at 0. That is,

$$p_1(0) = f(0) \quad \text{and} \quad p_1'(0) = f'(0)$$

If we want to improve our approximation, we could generalise this idea and look for a degree two polynomial p_2 such that the value, gradient and concavity of f and p_2 agree at 0. That is, if

$$p_2(x) = b_0 + b_1x + b_2x^2,$$

then we choose the coefficients b_0 , b_1 and b_2 such that

$$p_2(0) = f(0), \quad p_2'(0) = f'(0) \quad \text{and} \quad p_2''(0) = f''(0).$$

By calculating the first and second derivatives of f and p , we find that

$$b_0 = 1, \quad b_1 = 1 \quad \text{and} \quad 2b_2 = 1.$$

Hence

$$p_2(x) = 1 + x + \frac{x^2}{2}.$$

One can see in Figure 4.1 that p_2 gives a better approximation to f near 0 than does p_1 . Using p_2 , we obtain the approximation

$$e^{0.1} = f(0.1) \approx p_2(0.1) = 1.105.$$

One can again improve this approximation by using a polynomial of degree 3. If

$$p_3(x) = c_0 + c_1x + c_2x^2 + c_3x^3$$

then we can determine the unknown coefficients by solving the equations

$$p_3(0) = f(0), \quad p_3'(0) = f'(0), \quad p_3''(0) = f''(0) \quad \text{and} \quad p_3'''(0) = f'''(0).$$




Figure 4.1: Polynomial approximations (in gray) for the exponential function (in black) about 0.

These equations imply that

$$c_0 = 1, \quad c_1 = 1, \quad 2c_2 = 1 \quad \text{and} \quad 6c_3 = 1$$

and hence

$$p_3(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$$

(see Figure 4.1). Thus we have the approximation

$$e^{0.1} = f(0.1) \approx p_3(0.1) = 1.10516.$$

The table below compares the approximations to $e^{0.1}$ given by these first, second and third degree polynomials.

n	$p_n(x)$	$p_n(0.1)$	Error in approximation (3 s.f.)
1	$1 + x$	1.1	5.17×10^{-3}
2	$1 + x + \frac{x^2}{2}$	1.105	1.71×10^{-4}
3	$1 + x + \frac{x^2}{2} + \frac{x^3}{6}$	1.10516	4.25×10^{-6}

In fact, $e^{0.1}$ (rounded to five decimal places) is equal to 1.10517 and so the approximation $p_3(0.1)$ is accurate to four decimal places. Each of the polynomials p_1 , p_2 and p_3 are called *Taylor polynomials* for f about 0.

Using the same technique, one can attempt to approximate any function f at 0 with an n degree polynomial p_n , provided that f is n -times differentiable at 0. Suppose that

$$p_n(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \cdots + a_nx^n.$$

We require that

$$p_n(0) = f(0), \quad p'_n(0) = f'(0), \quad p''_n(0) = f''(0), \quad p_n^{(3)}(0) = f^{(3)}(0), \quad \dots, \quad p_n^{(n)}(0) = f^{(n)}(0),$$

where $f^{(j)}$ denotes the j th derivative of f . By calculating the derivatives of p , one finds that

$$a_0 = f(0), \quad a_1 = f'(0), \quad 2!a_2 = f''(0), \quad 3!a_3 = f^{(3)}(0), \quad \dots, \quad n!a_n = f^{(n)}(0).$$

Hence

$$p_n(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \cdots + \frac{f^{(n)}(0)}{n!}x^n.$$

Definition 4.1.1. Suppose that f is n -times differentiable at 0. Then the *Taylor polynomial* p_n of degree n for f about 0 is given by

$$p_n(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \cdots + \frac{f^{(n)}(0)}{n!}x^n.$$

We also call p_n the *n th Taylor polynomial* for f about 0.

Example 4.1.2. Suppose that $f(x) = e^x$ and $n \geq 0$. Find the Taylor polynomial of degree n for f about 0.

Solution. It is clear that

$$f'(x) = e^x, \quad f''(x) = e^x, \quad f'''(x) = e^x, \quad \dots, \quad f^{(n)}(x) = e^x.$$

Hence $f(0) = f'(0) = f''(0) = \cdots = f^{(n)}(0) = 1$ and

$$p_n(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!}$$

gives the Taylor polynomial of degree n about 0. □


Example 4.1.3. Suppose that $f(x) = \sin x$. Find the Taylor polynomials for f up to degree seven about the point 0.

Solution. We have

$f(x) = \sin x$	$f(0) = 0$
$f'(x) = \cos x$	$f'(0) = 1$
$f''(x) = -\sin x$	$f''(0) = 0$
$f^{(3)}(x) = -\cos x$	$f^{(3)}(0) = -1$
$f^{(4)}(x) = \sin x$	$f^{(4)}(0) = 0$
$\vdots$	$\vdots$

$$\begin{aligned} p_1(x) &= x \\ p_3(x) &= x - \frac{x^3}{3!} \\ p_5(x) &= x - \frac{x^3}{3!} + \frac{x^5}{5!} \\ p_7(x) &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \end{aligned}$$

are the Taylor polynomials for f up to degree seven about the point 0. Their graphs are compared to the sine function in the following diagram.



Note that the sine function is odd and that its Taylor polynomials about 0 are also odd. $\square$

The diagram in Example 4.1.3 shows that the Taylor polynomials about 0 are good approximations for f near 0. However, the approximations get worse farther away from 0. If one wants to approximate a function f near a point a , where a is not close to 0, then one can try (i) using Taylor polynomials about 0 of higher degree (which may or may not give satisfactory results) or (ii) approximating f with *Taylor polynomials about a* . Method (ii) is particularly pertinent in the case when $f(x) = \ln x$; since f is not defined at 0, the Taylor polynomials for f about 0 do not exist.

Suppose f is n -times differentiable at a . To approximate f about a , we try a degree n polynomial p_n , given by

$$p_n(x) = a_0 + a_1(x - a) + a_2(x - a)^2 + \cdots + a_n(x - a)^n,$$

where

$$p_n(a) = f(a), \quad p'_n(a) = f'(a), \quad p''_n(a) = f''(a), \quad \dots, \quad p_n^{(n)}(a) = f^{(n)}(a).$$

By calculating the derivatives of p_n at a , the unknown coefficients can be determined. Hence

$$p_n(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f^{(3)}(a)}{3!}(x - a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n.$$

This leads to the following definition.

Definition 4.1.4. Suppose that f is n -times differentiable at a . Then the *Taylor polynomial* p_n of degree n for f about a is given by

$$p_n(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n.$$

We also call p_n the *n th Taylor polynomial* for f about a .

The n th Taylor polynomial for f about a can be expressed using summation notation as

$$p_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k.$$

Here $0! = 1$ (by definition) and we use the convention of writing $f(a)$ as $f^{(0)}(a)$.

Example 4.1.5. Suppose that $f(x) = \ln x$. Find the Taylor polynomial of degree 5 for f about 1.

Solution. Since

$$\begin{array}{ll} f(x) = \ln x & f(1) = 0 \\ f'(x) = \frac{1}{x} & f'(1) = 1 \\ f''(x) = -\frac{1}{x^2} & f''(1) = -1 \\ f^{(3)}(x) = \frac{2!}{x^3} & f^{(3)}(1) = 2! \\ f^{(4)}(x) = -\frac{3!}{x^4} & f^{(4)}(1) = -3! \\ f^{(5)}(x) = \frac{4!}{x^5} & f^{(5)}(1) = 4!, \end{array}$$

we see that

$$\begin{aligned} p_5(x) &= (x-1) + \frac{-1}{2!}(x-1)^2 + \frac{2!}{3!}(x-1)^3 + \frac{-3!}{4!}(x-1)^4 + \frac{4!}{5!}(x-1)^5 \\ &= (x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \frac{(x-1)^5}{5}. \end{aligned}$$

It is preferable to express such a polynomial in powers of $(x-1)$ rather than in powers of x . There is usually no need to expand each of the terms $(x-1)^k$. $\square$

4.2 Taylor's theorem

(Ref: SH10 §12.6, 12.7)

We saw in the last section that $e^{0.1}$ can be approximated using Taylor polynomials. Suppose that a calculator (with a 10 digit display) uses Taylor polynomials to calculate the (approximate) value of $e^{0.1}$. What degree Taylor polynomial should the calculator use so that the displayed value

4.2. TAYLOR'S THEOREM


107

is accurate to 10 digits? Importantly, is there a way of answering this question without knowing in advance the decimal expansion of $e^{0.1}$?

In other words, we seek a method for determining how bad the error can get when approximating a function by one of its Taylor polynomials without knowing the precise values of the function. The functions f and g , given in the table below, further highlight the need for such a method.

Function	n th Taylor polynomial about 0
$f(x) = \sin x$	$p_n(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots \frac{x^n}{n!} \quad (n \text{ odd})$
$g(x) = \ln(1+x)$	$q_n(x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^5}{5} + \cdots \frac{x^n}{n}$

Suppose that we want to approximate $f(7)$. Although 7 is not close to 0, the graph below shows that the approximation $f(7) \approx p_n(7)$ is reasonable if $n = 19$ or $n = 21$.



By plotting more graphs, one would discover that the approximation $f(7) \approx p_n(7)$ seems to improve as n increases. We could conjecture that as $n \rightarrow \infty$ the error in the approximation approaches 0.

On the other hand, suppose that we want to approximate $g(1.5)$. Although 1.5 is closer to 0 than is 7, the Taylor polynomials do not provide a good approximation to the true value of $g(1.5)$, as Figure 4.2 shows. The Taylor polynomials approximate g very well on the interval $(-0.7, 0.7)$ and are not too bad on $(0.7, 1)$. However, to the right of 1 they should not be used to approximate the function. Moreover, while the higher degree Taylor polynomials give better approximations near 0, they give larger errors than lower degree polynomials to the right of 1. This surprising observation highlights the need for establishing a rigorous basis Taylor polynomial approximation. In particular, we must have some idea whether the error involved in each approximation is going to be large or small.

In light of the above discussion, we want an exact expression for the difference between a function and one of its Taylor polynomials. To obtain this expression, we use integration by parts.

Suppose that f has $n + 1$ continuous derivatives on an open interval I containing 0. We are about to compute the n th Taylor polynomial of f about 0 in such a way that we keep track of the *difference* (also known as the *error* or *remainder*) between $f(x)$ and $p_n(x)$. Fix a number x in the interval I and note that

$$\int_0^x f'(t) dt = f(x) - f(0). \quad (4.1)$$




Figure 4.2: Taylor polynomials for $\ln(1+x)$ about 0.

On the other hand, we can evaluate the same integral using integration by parts. If we set

$$\begin{aligned} u &= f'(t) & v &= -(x-t) \\ \frac{du}{dt} &= f''(t) & \frac{dv}{dt} &= 1 \end{aligned}$$

then integration by parts gives

$$\begin{aligned} \int_0^x f'(t) dt &= \left[-f'(t)(x-t) \right]_0^x + \int_0^x f''(t)(x-t) dt \\ &= f'(0)x + \int_0^x f''(t)(x-t) dt. \end{aligned} \quad (4.2)$$

Thus, from equations (4.1) and (4.2) we see that

$$f(x) = f(0) + f'(0)x + \int_0^x f''(t)(x-t) dt.$$

(Note that the first two terms on the right-hand side gives the Taylor polynomial of degree 1, and that the third term on the right-hand side is the error in the approximation $f(x) \approx p_1(x)$.) By applying integration by parts with

$$\begin{aligned} u &= f''(t) & v &= -\frac{1}{2}(x-t)^2 \\ \frac{du}{dt} &= f'''(t) & \frac{dv}{dt} &= x-t, \end{aligned}$$

we obtain

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{1}{2!} \int_0^x f'''(t)(x-t)^2 dt.$$

(The right-hand side is now the second Taylor polynomial of f with the associated error (also known as remainder)). If we continue integrating by parts then we obtain after n steps

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \frac{1}{n!} \int_0^x f^{(n+1)}(t)(x-t)^n dt.$$

That is,

$$f(x) = p_n(x) + R_{n+1}(x)$$

where p_n is the n th Taylor polynomial for f about 0 and

$$R_{n+1}(x) = \frac{1}{n!} \int_0^x f^{(n+1)}(t)(x-t)^n dt.$$

We call $R_{n+1}(x)$ the *remainder* term.

This argument may be easily adapted for Taylor polynomials about a point a , rather than 0. We thus have the following theorem.

Theorem 4.2.1 (Taylor's theorem). *Suppose that f has $n+1$ continuous derivatives on an open interval I containing a . Then for each x in I ,*

$$f(x) = p_n(x) + R_{n+1}(x),$$

where the n th Taylor polynomial p_n about a is given by

$$p_n(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n$$

and the remainder $R_{n+1}(x)$ is given by

$$R_{n+1}(x) = \frac{1}{n!} \int_a^x f^{(n+1)}(t)(x-t)^n dt. \quad (4.3)$$

Taylor's theorem tells us that the error in the approximation $f(x) \approx p_n(x)$ is *exactly* equal to $R_{n+1}(x)$. However, the remainder $R_{n+1}(x)$, as given in the integral form (4.3), is usually very difficult to compute. A more convenient form is known as the *Lagrange formula for the remainder*.

Corollary 4.2.2 (Lagrange formula for the remainder). *Suppose that f has $n+1$ continuous derivatives on an open interval I containing a . Then for each x in I ,*

$$f(x) = p_n(x) + R_{n+1}(x),$$

where p_n is the n th Taylor polynomial for f about a and

$$R_{n+1}(x) = \frac{f^{(n+1)}(c)}{(n+1)!}(x-a)^{n+1} \quad (4.4)$$

for some real number c between a and x .

The proof of Corollary 4.2.2 is left as an exercise in the tutorial problems.

Remark 4.2.3. Taylor's theorem with the Lagrange formula for the remainder is a generalisation of the mean value theorem. To see this, suppose that f is once differentiable. Then

$$f(x) - p_0(x) = R_1(x),$$

which means that

$$f(x) - f(a) = f'(c)(x-a).$$

for some c between x and a . In other words,

$$\frac{f(x) - f(a)}{x-a} = f'(c)$$

for some c between x and a .

In most examples, it is difficult to find the exact value of the number c that appears in the Lagrange formula for the remainder. Instead, one uses the fact that c lies between x and a to find an upper bound for the remainder term. That is while we may not know *what* c is, we do know *where* c is.

Example 4.2.4. Suppose that $f(x) = \cos x$. By considering the second Taylor polynomial for f about 0, estimate $f(1/5)$ and find an upper bound for the error.

4.2. TAYLOR'S THEOREM

111

Solution. The second Taylor polynomial p_2 is given by

$$\begin{aligned} p_2(x) &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 \\ &= \cos(0) - \sin(0)x - \frac{\cos(0)}{2!}x^2 \\ &= 1 - \frac{x^2}{2!}. \end{aligned}$$

Hence we use the estimate $\cos(\frac{1}{5}) \approx p_2(\frac{1}{5}) = \frac{49}{50}$. Using the Lagrange formula for the remainder, we calculate an upper bound for absolute error:

$$\begin{aligned} |\text{error}| &= |f(1/5) - p_2(1/5)| \\ &= |R_3(1/5)| && \text{(by Taylor's theorem)} \\ &= \left| \frac{f^{(3)}(c)}{3!} (1/5)^3 \right| && \text{(for some } c \text{ in } [0, 1/5]) \\ &= \frac{\sin c}{6} \times \frac{1}{125} && \text{(for some } c \text{ in } [0, 1/5]) \\ &\leq \frac{1}{6} \times \frac{1}{125} && \text{(since } \sin c \leq 1) \\ &= \frac{1}{750}. \end{aligned}$$

So an upper bound for the error is $\frac{1}{750}$, which is approximately 0.001333.

This upper bound is actually quite crude due to the estimate $\sin c \leq 1$. If instead we use the inequality

$$\sin t < t \quad \text{whenever } t > 0$$

(see Chapter 5 of MATH1131), then $\sin c < c < \frac{1}{5}$ and we obtain the upper bound

$$|\text{error}| < \frac{1}{3750}.$$

In summary, $\cos(\frac{1}{5}) \approx \frac{49}{50}$ and the absolute error in this estimate is less than $\frac{1}{3750}$ (that is, less than 0.0002667). $\square$

Example 4.2.5. A calculator with a 10 digit display uses a Taylor polynomial about 0 to estimate $e^{0.1}$. What degree polynomial should be used to guarantee that $e^{0.1}$ is displayed accurately?

Solution. We consider the function f given by $f(x) = e^x$. By Taylor's theorem,

$$e^x = p_n(x) + R_{n+1}(x)$$

where

$$p_n(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}$$

and

$$R_{n+1}(x) = \frac{e^c}{(n+1)!} x^{n+1}$$

for some c between 0 and x . Since the calculator will display the first ten digits of $p_n(0.1)$, it suffices if the error in the approximation $e^{0.1} \approx p_n(0.1)$ is less than 10^{-10} . That is, we need to find n large enough so that

$$|R_{n+1}(0.1)| < 10^{-10}. \quad (4.5)$$

$$\begin{aligned}
|R_{n+1}(0.1)| &= \frac{e^c}{(n+1)!} (0.1)^{n+1} && (\text{for some } c \text{ in } [0, 0.1]) \\
&\leq \frac{e^{0.1}}{(n+1)!} (0.1)^{n+1} && (\text{since exp is an increasing function}) \\
&< \frac{2}{(n+1)!} (0.1)^{n+1} && (\text{since } e^{0.1} < 3^{0.1} < 3^{1/2} < 2) \\
&= \frac{2}{(n+1)!} 10^{-(n+1)} && (\text{since } 0.1 = 10^{-1}).
\end{aligned}$$

Now $\frac{2}{(n+1)!} < 1$ whenever $n \geq 1$, so we obtain the (very crude) estimate

$$|R_{n+1}(0.1)| < 10^{-(n+1)}$$

provided that $n \geq 1$. Thus if $n = 9$ then (4.5) is satisfied. That is, if the calculator displays the first 10 digits appearing in the decimal expansion of $p_9(0.1)$, then the number appearing on the display will be accurate.

However, we can do better than this. By trial and error we find that the first (positive) integer n such that

$$\frac{2}{(n+1)!} 10^{-(n+1)} < 10^{-10}$$

is 6. That is, the calculator should display the first 10 digits appearing in the decimal expansion of $p_6(0.1)$.

(Note that we did not use the decimal expansion of $e^{0.1}$ to estimate the error.) $\square$

4.2.1 Classifying stationary points

Taylor's theorem can also be applied to the problem of classifying the stationary points of differentiable functions. For example, the function f given by $f(x) = (x-3)^4$ has a stationary point at 3 (since $f'(3) = 0$). However, since $f''(3) = 0$, the second derivative test cannot be used to determine whether the stationary point is a maximum, minimum or horizontal point of inflexion. By using Taylor's theorem, one can deduce the following improvement on the second derivative test.

Corollary 4.2.6. *Suppose that f is n times differentiable at a and that $f'(a) = 0$. If*

$$f''(a) = f'''(a) = \dots = f^{(k-1)}(a) = 0$$

but $f^{(k)}(a) \neq 0$, where $k \leq n$, then

- (i) a is a local minimum point if k is even and $f^{(k)}(a) > 0$;*
- (ii) a is a local maximum point if k is even and $f^{(k)}(a) < 0$;*
- (iii) a is an horizontal point of inflexion if k is odd.*

Sketch proof. Suppose that f is n times differentiable at a , and that

$$f'(a) = 0 = f''(a) = f'''(a) = \dots = f^{(k-1)}(a) = 0 \quad \text{and} \quad f^{(k)}(a) \neq 0.$$

We will make the additional assumptions that $f^{(k)}(x)$ exists for all x sufficiently close to a and that $f^{(k)}$ is continuous at a . (A proof of the general case is more complicated and will be omitted.)

Taylor's theorem (with Lagrange's remainder) implies that

$$\begin{aligned}
 f(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^{(k-1)}(a)}{(k-1)!}(x-a)^{k-1} + \frac{f^{(k)}(c)}{k!}(x-a)^k \\
 &= f(a) + 0 + 0 + \cdots + 0 + \frac{f^{(k)}(c)}{k!}(x-a)^k \\
 &= f(a) + \frac{f^{(k)}(c)}{k!}(x-a)^k
 \end{aligned} \tag{4.6}$$

for some c between x and a .

We now prove case (i). If k is even then $(x-a)^k > 0$ whenever $x \neq a$. Since $f^{(k)}(a) > 0$ and $f^{(k)}$ is continuous at a , we conclude that $f^{(k)}(c) > 0$ whenever x (and hence c) is sufficiently close to a . By combining these inequalities with (4.6), we conclude that $f(x) \geq f(a)$ for all x sufficiently close to a . Hence a is a local minimum point for f .

Cases (ii) and (iii) are proved in a similar way. $\square$

Example 4.2.7. Suppose that

$$f(x) = x^7 - 17x^6 + 101x^5 - 229x^4 + 3x^3 + 621x^2 - 297x - 567.$$

You are given that 3 is a stationary point of f . Classify this stationary point.

Solution. It is easy to check that

$$f'(3) = f''(3) = f'''(3) = 0 \quad \text{and} \quad f^{(4)}(3) = -1536 < 0.$$

By applying Corollary 4.2.6 (with k equal to 4), we conclude that 3 is a local maximum point for f . $\square$

4.2.2 Some questions arising from Taylor's theorem

To set the agenda for the rest of the chapter, we return now to our solution of Example 4.2.5, where Taylor polynomials were used to estimate $e^{0.1}$. We saw that

$$e^{0.1} = 1 + (0.1) + \frac{(0.1)^2}{2!} + \frac{(0.1)^3}{3!} + \cdots + \frac{(0.1)^n}{n!} + R_{n+1}(0.1), \tag{4.7}$$

where

$$0 < |R_{n+1}(0.1)| < \frac{2}{10^{n+1}(n+1)!}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{2}{10^{n+1}(n+1)!} = 0,$$

it seems reasonable to surmise that $|R_{n+1}(0.1)|$ (and hence $R_{n+1}(0.1)$) approaches 0 as $n \rightarrow \infty$. By letting n approach infinity in (4.7), this suggests that

$$e^{0.1} = 1 + (0.1) + \frac{(0.1)^2}{2!} + \frac{(0.1)^3}{3!} + \frac{(0.1)^4}{4!} + \frac{(0.1)^5}{5!} + \frac{(0.1)^6}{6!} + \cdots.$$

We therefore ask the following questions.

- What do we mean by $R_{n+1}(0.1) \rightarrow 0$ as $n \rightarrow \infty$? More generally, given a sequence of numbers $\{a_1, a_2, a_3, \dots\}$, how do we determine the limiting behaviour of a_n as $n \rightarrow \infty$?

- What is meant by the infinite sum

$$1 + (0.1) + \frac{(0.1)^2}{2!} + \frac{(0.1)^3}{3!} + \frac{(0.1)^4}{4!} + \frac{(0.1)^5}{5!} + \frac{(0.1)^6}{6!} + \dots?$$

Does the sum converge to a real number or (since we are adding infinitely many positive numbers) does the sum diverge to infinity? If the sum does converge to a real number then is that number $e^{0.1}$? More generally, given an infinite sum (also called a *series*) $a_1 + a_2 + a_3 + \dots$, how can we determine whether the series converges to a real number or else diverges?

- Each of these questions may be framed in a much larger context. Given a function f which is infinitely differentiable at a , Taylor's theorem gives

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_{n+1}(x).$$

For what values of x will $\lim_{n \rightarrow \infty} R_{n+1}(x) = 0$? For what values of x will the infinite series

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots$$

converge to a finite number? If the series does converge for some x , is it always true that

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots?$$

The example of $\ln(x+1)$ given at the beginning of this section suggests that we cannot always answer 'yes' to each of these questions. To appreciate a comprehensive answer to the questions posed, we must spend some time studying the limiting behaviour of sequences and series. We do so in the next three sections.

4.3 Sequences

(Ref: SH10 §11.6–11.4)

A *sequence* is a real-valued function defined on (a subset of) the natural numbers. For example, the function $f : \mathbb{N} \rightarrow \mathbb{R}$, given by $f(n) = n^2$, is a sequence. Sequences are usually denoted using a and b , rather than f and g . In the case of sequences, subscript notation is traditionally preferred over function notation. Thus

$$a(n) = n^2 \quad \forall n \in \mathbb{N}$$

and

$$a_n = n^2 \quad \text{whenever } n = 0, 1, 2, 3, \dots$$

describe the same sequence, but the second notation is used more frequently. Another way of describing the same sequence is

$$\{a_n\} = \{0, 1, 4, 16, 25, \dots\},$$

or more precisely,

$$\{a_n\}_{n=0}^{\infty} = \{n^2\}_{n=0}^{\infty}.$$

In each case, the number a_n is called the *n th term of the sequence*.

Example 4.3.1. The sequence

$$\left\{ \frac{1}{2}, \frac{2}{5}, \frac{3}{10}, \frac{4}{17}, \frac{5}{26}, \dots \right\}$$

is described by the rule $a_n = \frac{n}{n^2 + 1}$ whenever $n \geq 1$.

Example 4.3.2. The Fibonacci sequence

$$\{1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots\}$$

is described *recursively* by the rule

$$a_n = \begin{cases} 1 & \text{if } n = 1 \text{ or } n = 2 \\ a_{n-1} + a_{n-2} & \text{if } n \geq 3. \end{cases}$$

Example 4.3.3. Suppose that f is $m + 1$ times differentiable at a . Then for each x in $\mathbb{R}$, the remainder term $R_{n+1}(x)$ from Taylor's theorem gives rise to the (finite) sequence $\{R_{n+1}(x)\}_{n=1}^m$.

Since sequences are a class of functions, we can add, subtract, multiply and divide any two sequences that share a common domain. For example,

$$\{n^2\}_{n \in \mathbb{N}} \{\sqrt{n}\}_{n \in \mathbb{N}} = \{n^2 \sqrt{n}\}_{n \in \mathbb{N}}.$$

If the domain (in this case $\mathbb{N}$) is understood, then one can simply write $\{n^2\}\{\sqrt{n}\} = \{n^2 \sqrt{n}\}$.

4.3.1 Describing the limiting behaviour of sequences

Suppose that $\{a_n\}$ is a sequence. Our primary objective is to describe the behaviour of a_n as $n \rightarrow \infty$. There are two main types of behaviour. Either

- (a) a_n approaches some finite number L , in which case we say that the sequence $\{a_n\}$ is *convergent* and write $\lim_{n \rightarrow \infty} a_n = L$; or
- (b) the sequence $\{a_n\}$ is not convergent, in which case we say that $\{a_n\}$ is *divergent*.

Divergent sequences can be further classified according to the list below.

- (i) If $a_n \rightarrow \infty$ as $n \rightarrow \infty$ (that is, a_n grows without bound) then we say that the sequence $\{a_n\}$ *diverges to infinity*.
- (ii) If $a_n \rightarrow -\infty$ as $n \rightarrow \infty$ then we say that the sequence $\{a_n\}$ *diverges to negative infinity*.
- (iii) If $\{a_n\}$ has no limit as $n \rightarrow \infty$ but remains bounded then we say that $\{a_n\}$ *boundedly divergent*.
- (iv) If $\{a_n\}$ exhibits none of the above behaviour then we say that $\{a_n\}$ *unboundedly divergent*.

There are rigorous definitions for each of these cases. We will see one of them later in Remark 4.3.5.

Example 4.3.4. Describe the behaviour of each sequence $\{a_n\}$ as $n \rightarrow \infty$.

- (a) $a_n = n^3$

(b) $a_n = \sin(n\pi/2)$

(c) $a_n = \frac{3n^2}{n^2+4n+3}$

(d) $a_n = (-1)^n 2^n$

Solution. (a) Since $a_n \rightarrow \infty$ as $n \rightarrow \infty$, the sequence $\{a_n\}$ diverges to infinity.

(b) Since

$$\{a_n\}_{n=0}^{\infty} = \{0, 1, 0, -1, 0, 1, 0, -1, 0, \dots\}$$

we see that $\{a_n\}$ is bounded but does not have a limit. Hence $\{a_n\}$ is boundedly divergent.

(c) Since

$$\frac{3n^2}{n^2 + 4n + 3} = \frac{3}{1 + 4/n + 3/n^2} \rightarrow \frac{3}{1 + 0 + 0}$$

as $n \rightarrow \infty$, we conclude that $\{a_n\}$ converges to 3.

(d) Since

$$\{a_n\}_{n \in \mathbb{N}} = \{1, -2, 4, -8, 16, -32, 64, -128, \dots\}$$

we see that $\{a_n\}$ is not bounded. Moreover, since the even terms of the sequence approach infinity while the odd terms approach negative infinity, we conclude that the sequence $\{a_n\}$ is unboundedly divergent. $\square$


Remark 4.3.5. The formal definition for $\lim_{n \rightarrow \infty} a_n = L$ is similar to that for $\lim_{x \rightarrow \infty} f(x) = L$ (see the MATH1131 calculus notes) and is given as follows.

Suppose that $\{a_n\}_{n=0}^{\infty}$ is a sequence of real numbers and that $L \in \mathbb{R}$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if, for every positive number ϵ , there is a number M such that $|a_n - L| < \epsilon$ whenever $n > M$.

This definition may be interpreted geometrically.



Given *any* small number ϵ , there is a point M such that the distance between the a_n and the limit L is less than ϵ for every n past M . In other words, for every small band about the limit L , there is a point M such the sequence always lies in the band past M .

4.3.2 Techniques for calculating limits of sequences

Many of the rules and techniques given in MATH1131 for calculating limits of functions also apply for limits of sequences.

The first proposition of this subsection shows that limits behave well under the standard arithmetic operations.

Proposition 4.3.6. *Suppose that $\lim_{n \rightarrow \infty} a_n$ and $\lim_{n \rightarrow \infty} b_n$ exist. Then*

- (i) $\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$;
- (ii) $\lim_{n \rightarrow \infty} (a_n b_n) = \lim_{n \rightarrow \infty} a_n \times \lim_{n \rightarrow \infty} b_n$;
- (iii) $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$, provided that $\lim_{n \rightarrow \infty} b_n \neq 0$ and $b_n \neq 0$ for any n ; and
- (iv) $\lim_{n \rightarrow \infty} (\alpha a_n) = \alpha \lim_{n \rightarrow \infty} a_n$ for every real number α .

Example 4.3.7. Suppose that $a_n = \sqrt{n^2 + 4n} - n$. Determine the limiting behaviour of a_n as $n \rightarrow \infty$.

Solution. We proceed in the same manner as if we were asked to calculate $\lim_{x \rightarrow \infty} (\sqrt{x^2 + 4x} - x)$. Now

$$\begin{aligned}
 a_n &= \sqrt{n^2 + 4n} - n \\
 &= \frac{(\sqrt{n^2 + 4n} - n)(\sqrt{n^2 + 4n} + n)}{(\sqrt{n^2 + 4n} + n)} && \text{(multiplying top and bottom by the 'conjugate')} \\
 &= \frac{n^2 + 4n - n^2}{\sqrt{n^2 + 4n} + n} && \text{(difference of two squares)} \\
 &= \frac{4n}{\sqrt{n^2 + 4n} + n} \\
 &= \frac{4}{\sqrt{1 + 4/n} + 1} && \text{(dividing top and bottom by } n\text{).}
 \end{aligned}$$

Clearly $4/n \rightarrow 0$ as $n \rightarrow \infty$ and so by applying the limit rules of the above proposition we find that $a_n \rightarrow 2$ as $n \rightarrow \infty$. $\square$

The next proposition can be used when a function is composed with a sequence.

Proposition 4.3.8. *Suppose that $\lim_{n \rightarrow \infty} a_n = a$ and that f is continuous at a . Then*

$$\lim_{n \rightarrow \infty} f(a_n) = f(a).$$

This proposition is easy to remember if f is continuous everywhere; it amounts to saying that the function and limit can be swapped, as shown below:

$$\lim_{n \rightarrow \infty} f(a_n) = f\left(\lim_{n \rightarrow \infty} a_n\right).$$

Example 4.3.9. Find $\lim_{n \rightarrow \infty} \sin\left(\frac{\pi n^2}{4n^2 + 1}\right)$, if it exists.

Solution. Note that the sine function is continuous everywhere. Therefore

$$\begin{aligned}\lim_{n \rightarrow \infty} \sin\left(\frac{\pi n^2}{4n^2 + 1}\right) &= \sin\left(\lim_{n \rightarrow \infty} \frac{\pi n^2}{4n^2 + 1}\right) \\ &= \sin\left(\lim_{n \rightarrow \infty} \frac{\pi}{4 + 1/n^2}\right) \\ &= \sin\left(\frac{\pi}{4}\right) \\ &= \frac{1}{\sqrt{2}}.\end{aligned}$$

□

The following rule allows the use of l'Hôpital's rule when calculating limits of sequences.

Proposition 4.3.10. *Suppose that $\{a_n\}$ is a sequence and f is a function defined on some interval (b, ∞) . If $a_n = f(n)$ for all n sufficiently large and $\lim_{x \rightarrow \infty} f(x)$ exists then*

$$\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} f(x).$$

Note that this proposition only works for limits which are finite. If $a_n = f(n)$ and $f(x)$ diverges as $x \rightarrow \infty$, one cannot say that a_n diverges. (Consider, for example, the case when $a_n = 0$ for all n and $f(x) = x \sin(\pi x)$ for all x .)

The next example shows how Proposition 4.3.10 is applied in conjunction with l'Hôpital's rule.

Example 4.3.11. Suppose that $a_n = (1 + 1/n)^n$. Determine the limiting behaviour of a_n as $n \rightarrow \infty$.

Solution. Suppose that $f(x) = (1 + 1/x)^x$ whenever $x > 0$. Then $a_n = f(n)$ whenever $n > 0$. Hence

$$\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} f(x).$$

The method for calculating the limit of f is standard: we first take the logarithm of f (to remove the power) and then rearrange the resulting function so that we can apply l'Hôpital's rule. That

$$\begin{aligned}
 \lim_{n \rightarrow \infty} a_n &= \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x \\
 &= \lim_{x \rightarrow \infty} \exp \left\{ \ln \left(1 + \frac{1}{x}\right)^x \right\} && \text{(since ln and exp are inverses)} \\
 &= \exp \left\{ \lim_{x \rightarrow \infty} \ln \left(1 + \frac{1}{x}\right)^x \right\} && \text{(since exp is continuous everywhere)} \\
 &= \exp \left\{ \lim_{x \rightarrow \infty} x \ln \left(1 + \frac{1}{x}\right) \right\} && \text{(by the log law)} \\
 &= \exp \left\{ \lim_{x \rightarrow \infty} \frac{\ln(1 + 1/x)}{1/x} \right\} && \text{(to prepare for l'Hôpital's rule)} \\
 &= \exp \left\{ \lim_{x \rightarrow \infty} \frac{\left(\frac{-1/x^2}{1+1/x}\right)}{-1/x^2} \right\} && \text{(by l'Hôpital's rule)} \\
 &= \exp \left\{ \lim_{x \rightarrow \infty} \frac{1}{1 + 1/x} \right\} && \text{(simplifying the fraction)} \\
 &= \exp\{1\} \\
 &= e.
 \end{aligned}$$

In summary, a_n converges to e as $n \rightarrow \infty$. □

Remark 4.3.12. The limit $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$ is a standard result and should be familiar to students.

Finally, we present a version of the pinching theorem for sequences.

Proposition 4.3.13 (The pinching theorem for sequences). *Suppose that $\{a_n\}$, $\{b_n\}$ and $\{c_n\}$ are sequences and that for some positive integer N the inequality*

$$a_n \leq b_n \leq c_n$$

is satisfied whenever $n > N$. If $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$ then $\lim_{n \rightarrow \infty} b_n = L$.

The following example is important because sequences involving both factorials and powers arise frequently in applications of Taylor's theorem.

Example 4.3.14. Suppose that $a_n = \frac{n!}{n^n}$. Discuss the limiting behaviour of a_n as $n \rightarrow \infty$.

Solution. Note that

$$\begin{aligned}
 a_n &= \frac{1}{n} \cdot \frac{2}{n} \cdot \frac{3}{n} \cdots \frac{n}{n} \\
 &\leq \frac{1}{n} \cdot \frac{n}{n} \cdot \frac{n}{n} \cdots \frac{n}{n} \\
 &= \frac{1}{n}
 \end{aligned}$$

whenever $n \geq 1$. On the other hand, a_n is always positive. Thus

$$0 \leq a_n \leq \frac{1}{n}.$$

As $n \rightarrow \infty$ we conclude that $a_n \rightarrow 0$ by the pinching theorem. □

A similar technique was used in Lemma 2.2.5 to prove that

$$\lim_{n \rightarrow \infty} \frac{c^n}{n!} = 0$$

whenever $c > 0$.

Remark 4.3.15. It is helpful to have a good intuition of the *order of growth of sequences*. The following table compares the growth of various sequences as $n \rightarrow \infty$. The lower down on the table the sequence appears, the faster it grows at infinity.

a_n	growth rate as $n \rightarrow \infty$
1	constant: does not grow
$\ln n$	grows slowly
n^k , where $k > 0$	growth rate is faster for larger k
c^n , where $c > 1$	growth rate is faster for larger c
$n!$	grows rapidly
n^n	grows very rapidly

For example, the ordering in the table reflects the fact that

$$\lim_{n \rightarrow \infty} \frac{e^n}{n!} = 0 \quad \text{while} \quad \lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0.$$

The final theorem is of great theoretical importance and will be used in later sections. We begin with a definition.

Definition 4.3.16. A sequence $\{a_n\}_{n=0}^{\infty}$ of real numbers is said to be

- (a) *increasing* if $a_n < a_{n+1}$ for each natural number n ,
- (b) *nondecreasing* if $a_n \leq a_{n+1}$ for each natural number n ,
- (c) *decreasing* if $a_n > a_{n+1}$ for each natural number n , and
- (d) *nonincreasing* if $a_n \geq a_{n+1}$ for each natural number n .

If any of these four properties holds then the sequence is said to be *monotonic*.

Theorem 4.3.17. If $\{a_n\}_{n=0}^{\infty}$ is a bounded monotonic sequence of real numbers then it converges to some real number L .

This theorem is proved using a property that distinguishes the real numbers from the rational numbers. Given a bounded monotonic sequence of rational numbers, it is not true, in general, that the sequence converges to a rational number.

4.3.3 Suprema and infima [X]

In this subsection we summarise some more advanced ideas for students studying MATH1241.

Definition 4.3.18. Suppose that $\{a_n\}_{n=0}^{\infty}$ is a sequence of real numbers.

- (a) We say that M is an *upper bound* for $\{a_n\}_{n=0}^{\infty}$ if $a_n \leq M$ for every natural number n .
- (b) We say that M is a *lower bound* for $\{a_n\}_{n=0}^{\infty}$ if $a_n \geq M$ for every natural number n .
- (c) We say that K is the *least upper bound* for $\{a_n\}_{n=0}^{\infty}$ if K is an upper bound for $\{a_n\}_{n=0}^{\infty}$ and $K \leq M$ whenever M is an upper bound for $\{a_n\}_{n=0}^{\infty}$.
- (d) We say that K is the *greatest lower bound* for $\{a_n\}_{n=0}^{\infty}$ if K is a lower bound for $\{a_n\}_{n=0}^{\infty}$ and $K \geq M$ whenever M is a lower bound for $\{a_n\}_{n=0}^{\infty}$.

Example 4.3.19. Find the greatest lower bound and least upper bound for the sequence $\{a_n\}_{n=1}^{\infty}$, where $a_n = \frac{(-1)^n n}{n+1}$. Prove your answer.

Solution. We write out the first few terms of the sequence to get a feel for what is happening:

$$\left\{ -\frac{1}{2}, \frac{2}{3}, -\frac{3}{4}, \frac{4}{5}, -\frac{5}{6}, \frac{6}{7}, -\frac{7}{8}, \dots \right\}.$$

It is clear that the odd terms approach -1 (from above) while the even terms approach 1 (from below).

We will prove that 1 is the least upper bound. First, it is clear that $|a_n| = \frac{n}{n+1} < 1$ for every positive integer n . So 1 is an upper bound. Suppose now that K is also an upper bound but that $K < 1$. Hence $K = 1 - \epsilon$ for some positive number ϵ , while

$$\frac{n}{n+1} < K$$

for every positive integer n . Therefore

$$\frac{n}{n+1} < 1 - \epsilon$$

and hence

$$1 - \frac{1}{n+1} < 1 - \epsilon$$

for every positive integer n . But rearranging this inequality gives

$$n < \frac{1}{\epsilon} - 1$$

for every positive integer n , which gives a contradiction since the set of positive integers has no upper bound. Hence no such K exists. We conclude that 1 is the least upper bound.

Using a similar technique, one can show that -1 is the greatest lower bound for the sequence.

Note that the sequence has neither a maximum nor minimum value. $\square$

The fact that every bounded monotonic sequence of real numbers has a limit in $\mathbb{R}$ (see Theorem 4.3.17) follows from one of the axioms of the real number system. This axiom is called the *least upper bound axiom* and may be stated as

‘Every nonempty set of real numbers that has an upper bound has a *least* upper bound.’

Note that this axiom is not true for the rational number system.

To prove Theorem 4.3.17 in the case when $\{a_n\}_{n=0}^{\infty}$ is a bounded increasing sequence of real numbers, we note that the values of the sequence forms a bounded nonempty set of real numbers. By the least upper bound axiom, it therefore has an upper bound, which we denote by L . Using the definition of the limit (see Remark 4.3.5), one can now show that $\lim_{n \rightarrow \infty} a_n = L$. The proof of the other cases is similar.

We now introduce some alternate terminology and new notation for least upper bound and greatest lower bound.

Definition 4.3.20. Suppose that $\{a_n\}_{n=0}^{\infty}$ is a sequence of real numbers.

- (a) If $\{a_n\}_{n=0}^{\infty}$ has a least upper bound M , then M is also called the *supremum* of $\{a_n\}_{n=0}^{\infty}$ and is denoted by

$$\sup_{n \geq 0} a_n \quad \text{or} \quad \sup\{a_n : n \geq 0\}.$$

- (b) If $\{a_n\}_{n=0}^{\infty}$ has a greatest lower bound M , then M is also called the *infimum* of $\{a_n\}_{n=0}^{\infty}$ and is denoted by

$$\inf_{n \geq 0} a_n \quad \text{or} \quad \inf\{a_n : n \geq 0\}.$$

The plural for supremum and infimum is *suprema* and *infima*.

4.4 Infinite series

(Ref: SH10 §12.1, 12.2)

At the end of Section 4.2, we asked the question What is meant by the infinite sum

$$1 + (0.1) + \frac{(0.1)^2}{2!} + \frac{(0.1)^3}{3!} + \frac{(0.1)^4}{4!} + \frac{(0.1)^5}{5!} + \frac{(0.1)^6}{6!} + \dots?$$

In this section, we will develop a framework that gives meaning to this infinite sum by using existing notions for convergence (and divergence) of sequences. The key is to recognise that

- the terms of the above series form a sequence $\{a_k\}_{k=0}^{\infty}$, where

$$a_k = \frac{(0.1)^k}{k!};$$

and

- if s_n denotes the sum of the first n terms in the series (so that

$$s_n = 1 + (0.1) + \frac{(0.1)^2}{2!} + \frac{(0.1)^3}{3!} + \dots + \frac{(0.1)^n}{n!}$$

whenever $n \geq 0$) then $\{s_n\}_{n=0}^{\infty}$ is also a sequence.

Thus questions concerning the meaning of an infinite series can be reduced to studying the limiting behaviour of $\{s_n\}_{n=0}^{\infty}$. (Naturally, the limiting behaviour of $\{s_n\}$ depends on limiting properties of the sequence $\{a_k\}$; we will pay more attention to this aspect of the theory in Section 4.5.)

Definition 4.4.1. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of real numbers. For each natural number n , let s_n denote the n th partial sum given by

$$s_n = a_0 + a_1 + a_2 + \cdots + a_n = \sum_{k=0}^n a_k.$$

If the sequence $\{s_n\}_{n=0}^{\infty}$ of partial sums converges to a number L then we say that the infinite series $\sum_{k=0}^{\infty} a_k$ *converges to L* and we write

$$\sum_{k=0}^{\infty} a_k = L.$$

In this case we also say that the series is *summable*. If the sequence $\{s_n\}_{n=0}^{\infty}$ of partial sums diverges then we say that the infinite series $\sum_{k=0}^{\infty} a_k$ *diverges*.

The following example illustrates this definition.

Example 4.4.2 (Geometric series). Suppose that $r \in \mathbb{R}$ and consider the *geometric series*

$$\sum_{k=0}^{\infty} r^k = 1 + r + r^2 + r^3 + r^4 + \cdots.$$

Determine the values of r for which the series (a) converges and (b) diverges.

Solution. If $r \neq 1$ then s_n is given by the formula

$$s_n = \frac{1 - r^{n+1}}{1 - r}$$

(as is taught in high school). In the case when $r = 1$, we simply have

$$s_n = \underbrace{1 + 1 + 1 + \cdots + 1}_{n \text{ times}} = n.$$

In summary,

$$s_n = \begin{cases} \frac{1 - r^{n+1}}{1 - r} & \text{if } r \neq 1 \\ n & \text{if } r = 1. \end{cases}$$

To determine the convergence (or otherwise) of the infinite series $\sum_{k=0}^{\infty} r^k$, we simply determine the convergence of $\{s_n\}_{n=0}^{\infty}$. We break this up into four cases.

- If $|r| < 1$ then $r^{n+1} \rightarrow 0$ as $n \rightarrow \infty$ and so

$$\lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} \frac{1 - r^{n+1}}{1 - r} = \frac{1}{1 - r}.$$

Since the sequence of partial sums converges, so does the series and thus

$$\sum_{k=0}^{\infty} r^k = \frac{1}{1 - r}.$$

- If $|r| > 1$ then r^{n+1} diverges (either to infinity if $r > 1$ or unboundedly if $r < -1$). The sequence $\{s_n\}$ of partial sums therefore diverges and consequently the series $\sum_{k=0}^{\infty} r^k$ also diverges.
- If $r = 1$ then $s_n = n \rightarrow \infty$ as $n \rightarrow \infty$. Consequently the series diverges.
- If $r = -1$ then it is easily seen that $\{s_n\}_{n=0}^{\infty} = \{1, 0, 1, 0, 1, \dots\}$. Hence $\{s_n\}$ is boundedly divergent and the series

$$\sum_{k=0}^{\infty} r^k$$


also diverges.

□

The next example illustrates the technique of comparing a series with an integral.

Example 4.4.3 (The harmonic series). Show that the *harmonic series* $\sum_{k=1}^{\infty} \frac{1}{k}$ diverges.

Solution. This proof uses results from integration. Consider the diagram below.



It is clear that the area of the k th rectangle is $1/k$, which is also equal to the k th term of the series. Moreover, the area under the rectangles on the interval $[1, n+1]$ is greater than the area under the curve under the same interval. From these two observations it follows that

$$\begin{aligned} s_n &= \sum_{k=1}^n \frac{1}{k} \\ &\geq \int_1^{n+1} \frac{1}{x} dx \\ &= [\ln x]_1^{n+1} \\ &= \ln(n+1). \end{aligned}$$

4.4. INFINITE SERIES

125

Now $\ln(n+1) \rightarrow \infty$ as $n \rightarrow \infty$, so we conclude that $s_n \rightarrow \infty$ as $n \rightarrow \infty$. Hence the series diverges. $\square$

Using the technique illustrated in the previous example, one can show that

- $\sum_{k=1}^{\infty} \frac{1}{k^2}$ converges,
- $\sum_{k=1}^{\infty} \frac{1}{\sqrt{k}}$ diverges, and
- $\sum_{k=2}^{\infty} \frac{1}{k \ln k}$ diverges.

All one needs to do is draw a diagram and compare the series with an appropriate improper integral.

Since a convergent infinite series is the limit of a sequence (the sequence of partial sums), many results for sequences can be interpreted as results for series. The following proposition illustrates this point.

Proposition 4.4.4. Suppose that $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$ are two summable series. Then

$$(i) \sum_{k=0}^{\infty} (a_k + b_k) = \sum_{k=0}^{\infty} a_k + \sum_{k=0}^{\infty} b_k; \text{ and}$$

$$(ii) \sum_{k=0}^{\infty} (\alpha a_k) = \alpha \sum_{k=0}^{\infty} a_k \text{ for every real number } \alpha.$$

Proof. Let s_n and t_n denote the partial sums of $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$ respectively. Now apply Proposition 4.3.6 (i) and (iv) to the sequences $\{s_n\}$ and $\{t_n\}$. $\square$

Remark 4.4.5. While all the terms of a convergent series contribute to the value of the series, the convergence (or otherwise) of any series only depends on the ‘tail’ of the series. That is, the first hundred, thousand or even billion terms of the series are irrelevant to the question of whether the series converges. More precisely, given any positive integer N ,

$$\sum_{k=0}^{\infty} a_k \text{ converges} \quad \text{if and only if} \quad \sum_{k=N}^{\infty} a_k \text{ converges.}$$

With this in mind, all of the theorems presented in the next section are just as true for series of the form

$$\sum_{k=1}^{\infty} a_k, \quad \sum_{k=50}^{\infty} a_k \quad \text{or} \quad \sum_{k=2000}^{\infty} a_k$$

as they are for series of the form $\sum_{k=0}^{\infty} a_k$. When the starting point for the series does not matter, one sometimes simply writes

$$\sum a_k \quad \text{or} \quad \sum a_k.$$

(Ref: SH10 §12.3–12.5)

To determine whether a series

$$\sum_{k=0}^{\infty} a_k$$

converges or diverges, mathematicians have developed some simple tests. Typically, these tests examine the behaviour of the *sequence* $\{a_k\}$ and thereby deduce the convergence (or otherwise) of the corresponding *series*. In this section we introduce three such tests: the *kth term test*, the *ratio test* and the *alternating series test*.

Remark 4.5.1 (Warning). Care must be taken through this section not to confuse sequences and series. For example, suppose that $a_k = \frac{1}{k}$. While the *sequence* $\{a_k\}$ converges, the infinite *series* $\sum a_k$ does not.

4.5.1 Some preliminary results on series summation

The next two results are fundamental to the study of infinite series. They will later be used to establish some simple tests for convergence or divergence.

Lemma 4.5.2. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of positive numbers and let s_n denote the partial sum given by

$$s_n = \sum_{k=0}^n a_k.$$

If $\{s_n\}_{n=0}^{\infty}$ is a bounded sequence then the infinite series $\sum_{k=0}^{\infty} a_k$ is convergent.

Proof. For any natural number n ,

$$s_{n+1} = s_n + a_{n+1} > s_n,$$

since a_{n+1} is positive. Hence $\{s_n\}_{n=0}^{\infty}$ is a bounded increasing sequence and hence has a limit L (see Theorem 4.3.17). Therefore

$$\sum_{k=0}^{\infty} a_k = L$$

and the series converges. □

4.5.2 The *kth term* divergence test

The next test we introduce is a simple test for divergence. *One should always use this test first when trying to decide whether a series converges.*

Theorem 4.5.3 (The *kth term* test for divergence.). If $a_k \not\rightarrow 0$ as $k \rightarrow \infty$ then $\sum_{k=0}^{\infty} a_k$ diverges.

Before proving the theorem, we give an example.

Example 4.5.4. Determine whether the series $\sum_{k=1}^{\infty} \frac{k^2 + 2k}{\sqrt{k^4 + 2}}$ converges.

Solution. If $a_k = \frac{k^2 + 2k}{\sqrt{k^4 + 2}}$ then

$$\begin{aligned}\lim_{k \rightarrow \infty} a_k &= \lim_{k \rightarrow \infty} \frac{k^2 + 2k}{\sqrt{k^4 + 2}} \\ &= \lim_{k \rightarrow \infty} \frac{1 + 2/k}{\sqrt{1 + 2/k^4}} \quad (\text{by dividing top and bottom by } k^2) \\ &= 1.\end{aligned}$$

Since $a_k \not\rightarrow 0$ as $k \rightarrow \infty$, the series diverges by the k th term test. $\square$

Remark 4.5.5. The k th term test is not a test for convergence. For example, consider the series $\sum_{k=1}^{\infty} \frac{1}{k}$. In this case, $\lim_{k \rightarrow \infty} 1/k = 0$ but the series diverges (see Example 4.4.3).

The k th term divergence test is equivalent to the following theorem, whose proof we give below.

Theorem 4.5.6. *If the series $\sum_{k=0}^{\infty} a_k$ converges then $a_k \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. [H] Suppose that $\sum_{k=0}^{\infty} a_k$ converges to the real number L and let s_n denote the n th partial sum of the series. Then

$$\begin{aligned}s_n - s_{n-1} &= (a_0 + a_1 + \cdots + a_{n-1} + a_n) - (a_0 + a_1 + \cdots + a_{n-1}) \\ &= a_n.\end{aligned} \tag{4.8}$$

Now $\lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} s_{n-1} = L$ and so

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} (s_n - s_{n-1}) \quad (\text{by (4.8)}) \\ &= \lim_{n \rightarrow \infty} s_n - \lim_{n \rightarrow \infty} s_{n-1} \\ &= L - L \\ &= 0,\end{aligned}$$

thus completing the proof. $\square$

4.5.3 The integral test

The idea from Example 4.4.3, where we bounded a sum by an integral, can be applied more generally to produce a test for either convergence or divergence.

Theorem 4.5.7 (The integral test). *Suppose that $\sum a_k$ is an infinite series with positive terms. Suppose $f(x)$ is a positive integrable function decreasing on $[1, \infty)$ such that for each positive integer k , $f(k) = a_k$.*

(i) *If $\int_1^{\infty} f(x) dx$ converges then so does $\sum_{k=1}^{\infty} a_k$.*

(ii) *If $\int_1^{\infty} f(x) dx$ diverges then so does $\sum_{k=1}^{\infty} a_k$.*

The proof is similar to that given in Example 4.4.3.

Example 4.5.8. Determine whether or not the following series converge.

$$(a) \sum_{k=1}^{\infty} \frac{1}{k^2} \quad (b) \sum_{k=1}^{\infty} \frac{k}{2k^2 + 1} \quad (c) \sum_{k=2}^{\infty} \frac{1}{k(\log k)^2}$$

Proof. (a) Consider the improper integral, $\int_1^{\infty} \frac{1}{x^2} dx$.

$$\int_1^{\infty} \frac{1}{x^2} dx = \lim_{N \rightarrow \infty} \int_1^N \frac{1}{x^2} dx = \lim_{N \rightarrow \infty} \left[-\frac{1}{x} \right]_1^N = 1.$$

Since the improper integral converges, so does the series $\sum_{k=1}^{\infty} \frac{1}{k^2}$.

(Note: This is a famous series, first summed by Euler. Its value is remarkably $\frac{\pi^2}{6}$. This will be proven in later courses, but a proof of it appeared in the NSW Extension 2 paper, 2010.)

(b) Consider the improper integral, $\int_1^{\infty} \frac{x}{2x^2 + 1} dx$.

$$\int_1^{\infty} \frac{x}{2x^2 + 1} dx = \lim_{N \rightarrow \infty} \int_1^N \frac{x}{2x^2 + 1} dx = \lim_{N \rightarrow \infty} \left[\frac{1}{4} \log(2x^2 + 1) \right]_1^N \rightarrow \infty.$$

Since the improper integral diverges, so does the series $\sum_{k=1}^{\infty} \frac{k}{2k^2 + 1}$.

(c) Consider the improper integral, $\int_2^{\infty} \frac{1}{x(\log x)^2} dx$.

$$\int_2^{\infty} \frac{1}{x(\log x)^2} dx = \lim_{N \rightarrow \infty} \int_2^N \frac{1}{x(\log x)^2} dx = \lim_{N \rightarrow \infty} \left[-\frac{1}{\log x} \right]_2^N = \frac{1}{\log 2}.$$

Since the improper integral converges, so does the series $\sum_{k=1}^{\infty} \frac{1}{k(\log k)^2}$. □

4.5.4 The comparison test

The integral test was basically a comparison between each term of a given series and the area of a corresponding rectangle. This idea can also be applied to the terms of two series. If each term of a given series is less than each term of another series - whose convergence is easy to determine, then we can conclude the given series also converges. A similar test can be found for divergence.

Theorem 4.5.9 (The comparison test). *Suppose that $\{a_k\}_{k=0}^{\infty}$ and $\{b_k\}_{k=0}^{\infty}$ are two positive sequences such that $a_k \leq b_k$ for every natural number k .*

(i) *If $\sum_{k=0}^{\infty} b_k$ converges then $\sum_{k=0}^{\infty} a_k$ also converges.*

(ii) *If $\sum_{k=0}^{\infty} a_k$ diverges then $\sum_{k=0}^{\infty} b_k$ also diverges.*

The comparison test is often used in conjunction with series of the following type.

Proposition 4.5.10 (Convergence and divergence of p -series). *The series*

$$\sum_{k=1}^{\infty} \frac{1}{k^p}$$

converges if $p > 1$ and diverges if $p \leq 1$.

Proof. This theorem may be easily proved by the integral test, or by adapting the proof in Example 4.4.3. The details are left to the reader. $\square$

Example 4.5.11. Determine whether or not the following series converge.

$$(a) \sum_{k=1}^{\infty} \frac{k}{k^3 + 1} \quad (b) \sum_{k=2}^{\infty} \frac{1}{\sqrt{k^2 - 1}} \quad (c) \sum_{k=1}^{\infty} \frac{1}{\sqrt{k^2 + 1}}$$

Typically, one needs some intuition as to whether the series will converge or diverge before the comparison test is used to construct a rigorous solution.

Solution. (a) By considering the dominant term (as $k \rightarrow \infty$), we see that

$$\frac{k}{k^3 + 1} \approx \frac{k}{k^3} = \frac{1}{k^2}$$

whenever k is a large positive integer. Since $\sum \frac{1}{k^2}$ converges (p -series when $p = 2$), this suggests that $\sum_{k=1}^{\infty} \frac{k}{k^3 + 1}$ also converges.

To prove that $\sum_{k=1}^{\infty} \frac{k}{k^3 + 1}$ converges, we use part (i) of the comparison test. Note that

$$0 \leq \frac{k}{k^3 + 1} \leq \frac{k}{k^3} = \frac{1}{k^2}$$

whenever $k \geq 1$. So suppose that $a_k = \frac{k}{k^3 + 1}$ and $b_k = \frac{1}{k^2}$. We have shown that $0 \leq a_k \leq b_k$. Since $\sum b_k$ converges, it follows from the comparison test that $\sum a_k$ converges.

$$(b) \text{ Looking at the dominant terms, } \sum_{k=2}^{\infty} \frac{1}{\sqrt{k^2 - 1}} \approx \sum_{k=2}^{\infty} \frac{1}{\sqrt{k^2}} = \sum_{k=2}^{\infty} \frac{1}{k}.$$

Since this series diverges (p -series when $p = 1$), this suggests that $\sum_{k=2}^{\infty} \frac{1}{\sqrt{k^2 - 1}}$ also diverges.

To prove that $\sum_{k=2}^{\infty} \frac{1}{\sqrt{k^2 - 1}}$ diverges, we use part (ii) of the comparison test.

Note that $\frac{1}{\sqrt{k^2 - 1}} \geq \frac{1}{\sqrt{k^2}} = \frac{1}{k}$, whenever $k \geq 2$. So suppose that $a_k = \frac{1}{\sqrt{k^2 - 1}}$ and $b_k = \frac{1}{k}$. We have shown that $a_k \geq b_k$.

Since $\sum b_k$ diverges, it follows from the comparison test that $\sum a_k$ diverges.

(c) The same analysis as in (b) suggests that this series also diverges. However, the inequality $\frac{1}{\sqrt{k^2 + 1}} \geq \frac{1}{\sqrt{k^2}} = \frac{1}{k}$ is *false*. To overcome this hurdle, we introduce a ‘fudge factor’ to obtain the inequality we want. Thus, $\frac{1}{\sqrt{k^2 + 1}} \geq \frac{1}{2\sqrt{k^2}} = \frac{1}{2k}$ for all sufficiently large k (in fact $k \geq 1$ will do – you should check this!). So suppose that $a_k = \frac{1}{\sqrt{k^2 + 1}}$ and $b_k = \frac{1}{2k}$. We have shown that $a_k \geq b_k$.

Since $\sum b_k$ diverges, it follows from the comparison test that $\sum a_k$ diverges. $\square$

4.5.5 [X] The limit form of the comparison test

This form of the comparison test is extremely useful and allows us to rely on our intuition without having to work with inequalities. The disadvantage is that it does not always work in quite the same way as the straight comparison test does.

Proposition 4.5.12. Suppose a_n, b_n are sequences with positive terms and suppose $\lim_{n \rightarrow \infty} \frac{a_n}{b_n}$ is finite and **not** zero, then $\sum a_n$ converges if and only if $\sum b_n$ converges.

Proof. Suppose $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = K > 0$. For any given $\epsilon > 0$, we have

$$\left| \frac{a_k}{b_k} - K \right| < \epsilon$$

for all sufficiently large k . For such k we have

$$K - \epsilon < \frac{a_k}{b_k} < K + \epsilon \Rightarrow (K - \epsilon)b_k < a_k < (K + \epsilon)b_k.$$

Thus, from the last inequality, if $\sum a_k$ converges then $\sum (K - \epsilon)b_k$ converges and hence $\sum b_k$ does also; if $\sum b_k$ converges then $\sum (K + \epsilon)b_k$ converges and hence $\sum a_k$ does also. $\square$

Example 4.5.13. Discuss the convergence of $\sum_{k=5}^{\infty} \frac{k^2}{k^4 + 3}$

Proof. For large k the summand $a_k = \frac{k^2}{k^4 + 3}$ is roughly $b_k = \frac{1}{k^2}$.

Now $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$. Hence, since $\sum_{k=5}^{\infty} \frac{1}{k^2}$ converges (by p -series with $p = 2$), so does $\sum_{k=5}^{\infty} \frac{k^2}{k^4 + 3}$. $\square$

Example 4.5.14. Discuss the convergence of $\sum_{k=1}^{\infty} \sin\left(\frac{1}{k}\right)$

Proof. Since $\sin x \approx x$ for small x , we try comparing $a_k = \sin\left(\frac{1}{k}\right)$, (whose terms are positive), with $b_k = \frac{1}{k}$.

Now $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\sin\left(\frac{1}{k}\right)}{\frac{1}{k}} = 1$. Hence, since $\sum_{k=1}^{\infty} \frac{1}{k}$ diverges (by p -series with $p = 1$), so does $\sum_{k=1}^{\infty} \sin\left(\frac{1}{k}\right)$. $\square$

Remark 4.5.15. The series $\sum \frac{1}{\log n}$ diverges, since, for $n \geq 1$, we have $\log n < n$ giving $\frac{1}{\log n} > \frac{1}{n}$ and so the comparison test may be applied. On other hand, if we try to use the limit comparison test with $a_n = \frac{1}{\log n}$ and $b_n = \frac{1}{n}$, then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{n}{\log n} \rightarrow \infty$ and so the test fails.

We now introduce a simple convergence and divergence test known as the ratio test. It is important to note that this test can only be applied to series whose terms are positive.

Theorem 4.5.16 (The ratio test). *Suppose that $\sum a_k$ is an infinite series with positive terms and that*

$$\lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} = r.$$

(i) *If $r < 1$ then $\sum a_k$ converges.*

(ii) *If $r > 1$ then $\sum a_k$ diverges.*

Remark 4.5.17. The ratio test does not specify what happens if $r = 1$. In this case, the test is inconclusive; the series may converge or diverge.

The reason the ratio test works is that the tail of any series $\sum a_k$ with ‘ratio’ r given by

$$r = \lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k}$$

behaves like a geometric series with common ratio r . When $r < 1$, the geometric series is convergent and hence $\sum a_k$ also converges. Similarly, When $r > 1$, the geometric series is divergent and therefore $\sum a_k$ also diverges. Of course, these assertions need to be proved.

Before we see a proof of the ratio test, we shall see how it is applied. As seen in the examples below, the ratio test is particularly useful when $k!$ or k th powers appear in each term a_k .

Example 4.5.18. Determine whether or not the following series converge.

$$(a) \sum_{k=1}^{\infty} \frac{1}{k!} \quad (b) \sum_{k=1}^{\infty} \frac{k^k}{k!} \quad (c) \sum_{k=1}^{\infty} \frac{1}{k}$$

Solution. (a) Suppose that $a_k = \frac{1}{k!}$. Then

$$r = \lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} = \lim_{k \rightarrow \infty} \frac{k!}{(k+1)!} = \lim_{k \rightarrow \infty} \frac{1}{k+1} = 0.$$

Since $r < 1$, the series converges by the ratio test.

(b) Suppose that $a_k = \frac{k^k}{k!}$. Then

$$\begin{aligned} r &= \lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} \\ &= \lim_{k \rightarrow \infty} \frac{(k+1)^{k+1}}{(k+1)!} \cdot \frac{k!}{k^k} \\ &= \lim_{k \rightarrow \infty} \frac{(k+1)^k (k+1)}{k! (k+1)} \cdot \frac{k!}{k^k} \\ &= \lim_{k \rightarrow \infty} \frac{(k+1)^k}{k^k} \\ &= \lim_{k \rightarrow \infty} \left(\frac{k+1}{k} \right)^k \\ &= \lim_{k \rightarrow \infty} \left(1 + \frac{1}{k} \right)^k \\ &= e \end{aligned}$$

(see Example 4.3.11 for a calculation of this well known limit). Since $r = e > 1$, the series diverges by the ratio test.

(c) Since $r = \lim_{k \rightarrow \infty} \frac{k}{k+1} = 1$, we cannot say from the ratio test whether or not the series converges. It can be shown using another method that the series diverges (see Example 4.4.3). $\square$

Sketch proof of the ratio test. [X] (i) Suppose that $r < 1$ and choose R such that $r < R < 1$. Since $\lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} = r$, the terms of the sequence $\left\{ \frac{a_{k+1}}{a_k} \right\}$ eventually gets so close to r that they must also be less than R . More precisely, there is an integer N such that

$$\frac{a_{k+1}}{a_k} < R \quad \text{whenever } k \geq N. \quad (4.9)$$

Our goal from here is to show that the tail $\sum_{k=N}^{\infty} a_k$ of the series can be bounded above by a convergent geometric series with common ratio R . From (4.9) we see that

$$a_{N+1} < Ra_N, \quad a_{N+2} < Ra_{N+1} < R^2 a_N, \quad a_{N+3} < Ra_{N+2} < R^2 a_{N+1} < R^3 a_N,$$

and more generally that

$$a_{N+j} < R^j a_N \quad \text{whenever } j \geq 0. \quad (4.10)$$

Hence

$$\begin{aligned} \sum_{k=N}^{\infty} a_k &= \sum_{j=0}^{\infty} a_{N+j} \\ &< \sum_{j=0}^{\infty} R^j a_N && \text{(by inequality (4.10))} \\ &= a_N \sum_{j=0}^{\infty} R^j && \text{(which is a geometric series)} \\ &= \frac{a_N}{1-R} && \text{(since } R < 1). \end{aligned}$$

Since the tail of the series $\sum a_k$ converges, the series itself must converge.

(ii) If $r > 1$, then

$$\lim_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} > 1$$

and so

$$\frac{a_{k+1}}{a_k} > 1$$

for all k sufficiently large. That is, $a_{k+1} > a_k$ for all k sufficiently large, which means that the positive sequence $\{a_k\}$ eventually becomes an increasing sequence. Hence $a_k \not\rightarrow 0$ as $k \rightarrow \infty$. It follows from the k th term test that the series is divergent. $\square$

4.5.7 Leibniz' test for alternating series

So far we have given a convergence test for series all of whose terms are positive. If the terms are all negative, then we simply multiply the series by -1 to obtain series whose terms are positive. However, if the series has a mixture of positive and negative terms, we cannot apply this trick. In the next two subsections, we deal with series whose terms have mixed signs. The simplest case is when the sign alternates from term to term.

Definition 4.5.19. If $\{a_k\}_{k=0}^{\infty}$ is a sequence of positive real numbers, then the series

$$a_0 - a_1 + a_2 - a_3 + a_4 - a_5 + a_6 - a_7 + a_8 - a_9 + \cdots$$

is called an *alternating series*.

An alternating series is often written in the form

$$\sum_{k=0}^{\infty} (-1)^k a_k.$$

The following theorem, proved by Leibniz in the early eighteenth century, is a simple test for the convergence of alternating series.

Theorem 4.5.20 (Alternating series test). *Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of real numbers satisfying the following properties:*

- (a) $a_k \geq 0$;
- (b) $a_k \geq a_{k+1}$ for all k (that is, the sequence is nonincreasing); and
- (c) $\lim_{k \rightarrow \infty} a_k = 0$.

Then the alternating series $\sum_{k=0}^{\infty} (-1)^k a_k$ converges.

Before proving the theorem, we give an example and state a corollary.

Example 4.5.21. Determine whether the series $\sum_{k=2}^{\infty} \frac{(-1)^k k}{k^2 + 1}$ is summable.

Solution. Since this is an alternating series, one naturally tries the alternating series test. Suppose that $a_k = \frac{k}{k^2 + 1}$. We need to check that $\{a_k\}$ satisfies hypotheses (a), (b) and (c) of the alternating series test.

It is clear that (a) and (c) hold. To prove (b), consider the function f given by

$$f(x) = \frac{x}{x^2 + 1}.$$

Now

$$f'(x) = \frac{1 - x^2}{(1 + x^2)^2}$$

and hence $f'(x) < 0$ whenever $x > 1$. That is, f is decreasing on the interval $(1, \infty)$. Since $f(k) = a_k$ whenever $k \geq 2$, it follows that $\{a_k\}_{k=2}^{\infty}$ is a decreasing sequence.

We now apply the alternating series test and deduce that the series converges. $\square$

If the hypotheses of the alternating sequence test are satisfied, then not only do we know that the series converges to some limit L , but we can also approximate L with any partial sum s_n and obtain an upper bound for the corresponding error.

Corollary 4.5.22. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of numbers satisfying properties (a), (b) and (c) of the alternating series test. Denote the value of the convergent series $\sum_{k=0}^{\infty} (-1)^k a_k$ by L and the n th partial sum of the same series by s_n . Then

$$|s_n - L| \leq a_{n+1} \quad (4.11)$$

for every natural number n .

In effect, the corollary says that if you chop the series off after the n th term, the error in approximation will be less than the $(n+1)$ st term. The proof is given at the end of this subsection.

Example 4.5.23. Estimate the value of the series $\sum_{k=0}^{\infty} \frac{(-1)^k}{k^2 + 1}$ such that the error is less than $\frac{1}{100}$.

Solution. It is easy to verify that the sequence $\{a_k\}$, where $a_k = \frac{1}{k^2 + 1}$, satisfies the hypotheses of the alternating series test. Hence the infinite series converges. Denote the value of the series by L . We will use the estimate $s_n \approx L$, where n is chosen such that the absolute error is less than $\frac{1}{100}$. Now

$$\begin{aligned} \text{absolute error} &= |s_n - L| \\ &\leq a_{n+1} && \text{(by Corollary 4.5.22)} \\ &= \frac{1}{(n+1)^2 + 1}. \end{aligned}$$

So it is enough to guarantee that


$$\frac{1}{(n+1)^2 + 1} < \frac{1}{100}.$$

Clearly the smallest positive integer n that satisfies this inequality is 9. So

$$L \approx s_9 = \sum_{k=0}^9 \frac{(-1)^k}{k^2 + 1} = 1 - \frac{1}{5} + \frac{1}{10} - \frac{1}{17} + \cdots - \frac{1}{82}$$

and the error in this approximation is less than $\frac{1}{100}$. (Using Maple, one finds that $s_9 = 0.6305785114$ (correct to 10 decimal places) and so $L \approx 0.63$.) $\square$

We conclude this subsection by proving the alternating series test and its corollary. The following diagram illustrates the typical behaviour of the partial sums of a series that satisfies hypotheses (a), (b) and (c) of the alternating series test. It will be helpful to bear this diagram in mind when reading the proofs.



4.5. TESTS FOR SERIES CONVERGENCE

135

Proof of Theorem 4.5.20. [X] Let s_n denote the n th partial sum of the series and suppose that properties (a), (b) and (c) hold. The proof proceeds in three steps.

Step 1. We will prove that the sequence $\{s_{2n}\}_{n=0}^{\infty}$ of even partial sums is bounded above by 0. Now

$$s_{2n} = (a_0 - a_1) + (a_2 - a_3) + (a_4 - a_5) + \cdots + (a_{2n-2} - a_{2n-1}) + a_{2n}. \quad (4.12)$$

By property (b) we see that

$$a_0 - a_1 \geq 0, \quad a_2 - a_3 \geq 0, \quad \dots, \quad a_{2n-2} - a_{2n-1} \geq 0$$

and by property (a) it is evident that $a_{2n} \geq 0$. It follows from (4.12) that $s_{2n} \geq 0$ for every n .

Step 2. We will prove that the sequence $\{s_{2n}\}_{n=0}^{\infty}$ of even partial sums is nonincreasing. Now

$$\begin{aligned} s_{2n} - s_{2n+2} &= (a_0 - a_1 + \cdots + a_{2n}) - (a_0 - a_1 + \cdots + a_{2n} - a_{2n+1} + a_{2n+2}) \\ &= a_{2n+1} - a_{2n+2} \\ &\geq 0 \end{aligned}$$

since $a_{2n+1} \geq a_{2n+2}$ by property (b). Thus $s_{2n} \geq s_{2n+2}$ for all n , which means that $\{s_{2n}\}_{n=0}^{\infty}$ is nonincreasing.

Step 3. From Steps 1 and 2, we conclude that $\{s_{2n}\}_{n=0}^{\infty}$ is a bounded monotonic sequence and hence convergent. Call the limit of this sequence L . If we can show that the sequence $\{s_{2n+1}\}_{n=0}^{\infty}$ of odd partial sums also converges to L , then we can conclude that $\{s_n\}$ converges. Now

$$s_{2n+1} = s_{2n} + a_{2n+1} \rightarrow L + 0$$

as $n \rightarrow \infty$ by property (c). Hence $\{s_n\}$ converges. □

Proof of Corollary 4.5.22. [X] Suppose that $\{a_k\}$ satisfies the hypotheses (a), (b) and (c) of the alternating series test. Since the infinite series converges, $\lim_{n \rightarrow \infty} s_n = L$ for some real number L . If n is odd then

$$s_{n+2} = s_n + a_{n+1} - a_{n+2} \geq s_n,$$

and so the odd partial sums increase towards L from below. If n is even then

$$s_{n+2} = s_n - a_{n+1} + a_{n+2} \leq s_n,$$

and so the even partial sums decrease towards L from above. Hence if n is odd then

$$s_n \leq L \leq s_{n+1} = s_n + a_{n+1},$$

while if n is even then

$$s_n - a_{n+1} = s_{n+1} \leq L \leq s_n.$$

That is,

$$\text{either } s_n \leq L \leq s_n + a_{n+1} \quad \text{or} \quad s_n - a_{n+1} \leq L \leq s_n.$$

Both cases imply (4.11). □

Consider the series

$$1 + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} + \frac{1}{5!} - \frac{1}{6!} + \frac{1}{7!} + \frac{1}{8!} - \frac{1}{9!} + \cdots. \quad (4.13)$$

Since

$$\lim_{k \rightarrow \infty} |a_k| = \lim_{k \rightarrow \infty} \frac{1}{k!} = 0,$$

the k th term test for divergence does not apply. One cannot apply the ratio test (since not all the terms positive) and clearly this is not an alternating series. Is there a way of determining whether the series is summable? The theorem given below proves very helpful in this instance. First we give a definition.

Definition 4.5.24. A series $\sum_{k=0}^{\infty} a_k$ is said to be *absolutely convergent* if the series

$$\sum_{k=0}^{\infty} |a_k|$$

is convergent.

Theorem 4.5.25. *If a series is absolutely convergent then it converges.*

Proof. [H] Suppose that the series $\sum_{k=0}^{\infty} a_k$ converges absolutely. For each natural number k ,

$$-|a_k| \leq a_k \leq |a_k|$$

and hence

$$0 \leq a_k + |a_k| \leq 2|a_k|. \quad (4.14)$$

Since $\sum |a_k|$ converges, it follows that $2\sum |a_k|$ converges and hence $\sum 2|a_k|$ converges. By the comparison test, we deduce from (4.14) that $\sum (a_k + |a_k|)$ converges. Now

$$a_k = (a_k + |a_k|) - |a_k|.$$

Since the sum of the terms on the right-hand side converges (by Proposition 4.4.4), the sum of the terms on the left must also converge. That is, $\sum_{k=0}^{\infty} a_k$ converges, thus completing the proof. $\square$

Example 4.5.26. Determine whether or not the series given by (4.13) is convergent.

Solution. Let a_k denote the k th term of the series. By Theorem 4.5.25, it is enough to show that the series converges absolutely. Now

$$\sum_{k=1}^{\infty} |a_k| = \sum_{k=1}^{\infty} \frac{1}{k!}$$

and this series converges by the ratio test (see Example 4.5.18 (a)). $\square$

Not every convergent series is absolutely convergent. For example, the alternating series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots \quad (4.15)$$

converges (by the alternating series test) but the corresponding absolute series

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \cdots$$

diverges (see Example 4.4.3). In this situation, we say that the series (4.15) is *conditionally convergent*.

Definition 4.5.27. A series $\sum_{k=0}^{\infty} a_k$ is said to be *conditionally convergent* if it converges but the series

$$\sum_{k=0}^{\infty} |a_k|$$

diverges.

The distinction between conditionally and absolutely convergent series is brought into bold relief when considering rearrangements of series.

Definition 4.5.28. A *rearrangement* of a series $\sum a_k$ is a series that has exactly the same terms but that is summed in a different order.

For example,

$$1 + \frac{1}{3} + \frac{1}{5} - \frac{1}{2} + \frac{1}{7} + \frac{1}{9} - \frac{1}{4} + \frac{1}{11} + \cdots$$

and

$$-\frac{1}{2} + 1 - \frac{1}{4} + \frac{1}{3} - \frac{1}{6} + \frac{1}{5} - \frac{1}{8} + \frac{1}{7} - \cdots$$

are both rearrangements of

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \frac{1}{8} + \cdots.$$

If the series is finite, then every rearrangement has the same value (since addition of real numbers is commutative). However, if the series is infinite, then the value (if it exists) of each rearrangement is determined by a limit of partial sums, and one cannot appeal to commutativity of addition, as in the finite case. In fact, some rather surprising phenomena occur with rearrangements of conditionally convergent series.

Theorem 4.5.29. Suppose that $\sum_{k=0}^{\infty} a_k$ is an infinite series.

(i) If $\sum a_k$ converges absolutely, then every rearrangement of the series converges absolutely and all rearrangements have the same limit as $\sum a_k$.

(ii) If $\sum a_k$ converges conditionally, then given any real number L , the series has a rearrangement that converges to L . Moreover, every conditionally convergent series has a rearrangement that diverges to ∞ , and another rearrangement that diverges to $-\infty$.

This theorem was published in 1867 by Riemann. One of the tutorial problems illustrates that the conditionally convergent series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \cdots$$

can be rearranged to sum to different real numbers. The moral of the story is that one should not rearrange a conditionally convergent series to determine its value.

4.6 Taylor series

(Ref: SH10 §12.7)

At the end of Section 4.2 we posed the question, When is it true that

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots?$$

Having studied sequences and series of real numbers, we now have the tools to deal with this and related questions. First, we give the above series expansion a special name.

Definition 4.6.1. Suppose that a function f has derivatives of all orders at a . Then the series

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots,$$

which may also be written as

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k,$$

is called the *Taylor series for f about a* . In the case when $a = 0$, the series is also called the *Maclaurin series for f* .

Next, we need to define what we mean by the convergence (or divergence) of a Taylor series.

Definition 4.6.2. Suppose that I is an interval and that f has derivatives of all orders at some point a . We say that

- (a) the Taylor series for f about a *converges on I* if the series

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$$

converges for each point x in I ;

- (b) the Taylor series for f about a *converges to f on I* if for each x in I , x lies in the domain of f and

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k;$$

and

- (c) the Taylor series for f at a *diverges on I* if the series

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$$

diverges for each point x in I .

Thus the question asked at the beginning of this section may be rephrased as,

For what intervals I will the Taylor series of a function f converge to f ?

The following corollary to Taylor's theorem helps answer this question.

Corollary 4.6.3. Suppose that f has derivatives of all orders at a and that x lies in the domain of f . Let $R_{n+1}(x)$ denote the remainder term of Theorem 4.2.1 (or its equivalent form as given in Corollary 4.2.2). If

$$\lim_{n \rightarrow \infty} R_{n+1}(x) = 0 \quad (4.16)$$

then

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n.$$

Proof. Let p_n denote the n th Taylor polynomial for f about a . Taylor's theorem implies that

$$f(x) = p_n(x) + R_{n+1}(x). \quad (4.17)$$

Note that $p_n(x)$ is the n th partial sum of the series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n.$$

So we only have to show that

$$\lim_{n \rightarrow \infty} p_n(x) = f(x). \quad (4.18)$$

$$\begin{aligned} |p_n(x) - f(x)| &= |R_{n+1}(x)| && \text{(by (4.17))} \\ &\rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$ by (4.16). That is, the distance between $p_n(x)$ and $f(x)$ can be made as small as we like. Hence (4.18) follows. $\square$

Example 4.6.4. Suppose that $x \in \mathbb{R}$. Show that

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots .$$

Solution. Suppose that $f(t) = e^t$ and fix x in $\mathbb{R}$. The sum of the terms up to (and including) $\frac{x^n}{n!}$ is equal to $p_n(x)$, where p_n is the n th Taylor polynomial for f about 0. So by Corollary 4.6.3, we only need to show that $\lim_{n \rightarrow \infty} R_{n+1}(x) = 0$. Now

$$f^{(n)}(t) = e^t \quad \forall t \in \mathbb{R},$$

and so by the Lagrange formula for the remainder,

$$R_{n+1}(x) = \frac{e^c}{(n+1)!} x^{n+1}$$

for some c between 0 and x . Now $e^c \leq e^{|c|} \leq e^{|x|}$. If $M = e^{|x|}$ then

$$\begin{aligned} 0 &\leq |R_{n+1}(x)| \\ &= \frac{e^c}{(n+1)!} |x|^{n+1} \\ &\leq \frac{M|x|^{n+1}}{(n+1)!} \\ &\rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$ by Lemma 2.2.5. Therefore

$$\lim_{n \rightarrow \infty} R_{n+1}(x) = 0$$

by the pinching theorem for sequences. $\square$

We have just proved that the Taylor series about 0 for the exponential function converges to the exponential function on $\mathbb{R}$. We say that the exponential function is *represented* by its Taylor series about 0 on $\mathbb{R}$. Some other convergent Taylor series representations are given in the next theorem.

Theorem 4.6.5. *The following formulae hold whenever x lies in the given interval.*

$$\begin{aligned}
 \frac{1}{1-x} &= 1 + x + x^2 + x^3 + x^4 + \dots & x &\in (-1, 1) \\
 e^x &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots & x &\in \mathbb{R} \\
 \sin x &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots & x &\in \mathbb{R} \\
 \cos x &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots & x &\in \mathbb{R} \\
 \sinh x &= x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots & x &\in \mathbb{R} \\
 \cosh x &= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots & x &\in \mathbb{R} \\
 \ln(1+x) &= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots & x &\in (-1, 1] \\
 \tan^{-1} x &= x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots & x &\in [-1, 1]
 \end{aligned}$$

Moreover, if x lies outside the given interval then the corresponding Maclaurin series diverges.

Most of these formulae can be proved by showing that the Lagrange formula for the remainder tends to 0 as $n \rightarrow \infty$. However, sometimes one must resort to using the integral form of the remainder (as given by Theorem 4.2.1). One of the tutorial problems illustrates its use. In Section 4.8, we introduce tools that provide an alternate approach to deriving some of these expansions.

Remark 4.6.6. The Taylor series expansions for $\sin x$ and $\ln(1+x)$ given by Theorem 4.6.5 explain the phenomena discussed at the beginning of Section 4.2. In particular, the Taylor series for $\sin x$ converges for all x in $\mathbb{R}$, which explains why $\sin(7)$ could be approximated by Taylor polynomials of sufficiently high degree. On the other hand, we cannot use Taylor polynomials to approximate $\ln(1+x)$ when $x > 1$ (as suggested by Figure 4.2) because the Taylor series *diverges* when $x > 1$. This explains why higher order Taylor polynomials give worse approximations for $\ln(1+x)$ when $x > 1$.

Remark 4.6.7. The Maclaurin series given by Theorem 4.6.5 can be used to obtain beautiful series expansions for some irrational numbers. By substituting particular values for x into an appropriate Maclaurin series, one finds that

$$\begin{aligned}
 e &= 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots \\
 \ln 2 &= 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots \\
 \frac{\pi}{4} &= 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots
 \end{aligned}$$

Unfortunately, the last two series converge too slowly to be of high computational value.

Remark 4.6.8. If f equals its Taylor series on an interval I , then the corresponding Taylor polynomial p_n can be used to approximate f on I . However, it is important to appreciate that some Taylor series (such as that for e^x) converge much more quickly to the function than do others (such as that for $\ln(1+x)$, which converges slowly). If the series converges very slowly then the approximation $f \approx p_n$ is only accurate when n is very large.

Remark 4.6.9. If a Taylor series converges to a function f on an interval I , then (obviously) the Taylor series converges on I . However, the converse is not true. That is, if the Taylor series converges on I , then one cannot conclude that the Taylor series converges to f on I . In the tutorial problems we give one example of a function f whose Maclaurin series converges on the entire real line but only converges to f at the origin.

4.7 Power series

(Ref: SH10 §12.8)

A Maclaurin series is a series of the form

$$a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + \cdots, \quad (4.19)$$

where each coefficient a_k is given by

$$a_k = \frac{f^{(k)}(0)}{k!}$$

for some function f that is infinitely differentiable at 0. For the remainder of this chapter, we study more general series of the form (4.19), where each coefficient a_k is not necessarily a Taylor coefficient. Such a series is called a *power series*.

Definition 4.7.1. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of real numbers and that $a \in \mathbb{R}$. A series of the form

$$\sum_{k=0}^{\infty} a_k x^k$$

is called a *power series in powers of x* . A series of the form

$$\sum_{k=0}^{\infty} a_k (x - a)^k$$

is called a *power series in powers of $x - a$* .

Thus a Maclaurin series is a power series in powers of x , while a Taylor series about a is a power series in powers of $x - a$. For the last two sections of this chapter, we discuss the convergence, addition, multiplication, integration and differentiation of power series. Hence whatever is said about power series also applies to Maclaurin and Taylor series.

In this section, we focus on the convergence and divergence of power series.

Definition 4.7.2. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of real numbers, I is an interval and a is a real number. We say that a power series $\sum_{k=0}^{\infty} a_k(x-a)^k$ converges

(a) at a real number c if the series $\sum_{k=0}^{\infty} a_k(c-a)^k$ converges;

(b) on the interval I if the series $\sum_{k=0}^{\infty} a_k(x-a)^k$ converges for each x in I .

We say that a power series $\sum_{k=0}^{\infty} a_k(x-a)^k$ diverges

(a) at a real number c if the series $\sum_{k=0}^{\infty} a_k(c-a)^k$ diverges;

(b) on the interval I if the series $\sum_{k=0}^{\infty} a_k(x-a)^k$ diverges for each x in I .

Using the ratio test, one can often determine for what values of x a power series converges absolutely.

Example 4.7.3. Find an interval I such that the power series $\sum_{k=0}^{\infty} \frac{kx^k}{3^k}$ converges on I .

Solution. We first find an interval I on which the series converges absolutely. To do so, we apply the ratio test to the (absolute) series

$$\sum_{k=0}^{\infty} \left| \frac{kx^k}{3^k} \right|.$$

Now

$$\begin{aligned} r &= \lim_{k \rightarrow \infty} \left| \frac{(k+1)x^{k+1}}{3^{k+1}} \right| \cdot \left| \frac{3^k}{kx^k} \right| \\ &= \lim_{k \rightarrow \infty} \frac{(k+1)|x|}{3k} \\ &= \frac{|x|}{3}. \end{aligned}$$

To conclude from the ratio test that the series converges, we require that $r < 1$, which means that $|x| < 3$. So the series converges absolutely whenever $-3 < x < 3$. Hence the series $\sum_{k=0}^{\infty} \frac{kx^k}{3^k}$ converges on the interval $(-3, 3)$. $\square$

Remark 4.7.4. In the previous example, one can also conclude that the series diverges when $|x| > 3$. To see this, fix x in $\mathbb{R}$ such that $|x| > 3$. Let b_k denote the k th term

$$\frac{kx^k}{3^k}$$

of the series. Now, by the same calculation as before,

$$\lim_{k \rightarrow \infty} \left| \frac{b_{k+1}}{b_k} \right| = \frac{|x|}{3}.$$

Since $|x| > 3$ we deduce that $\lim_{k \rightarrow \infty} \frac{|b_{k+1}|}{|b_k|} > 1$. This shows that $\frac{|b_{k+1}|}{|b_k|} > 1$ whenever k is sufficiently large. Rearranging implies that

$$|b_{k+1}| > |b_k|$$

for all sufficiently large k and hence the tail of the sequence $\{|b_k|\}$ is increasing. Thus $|b_k| \not\rightarrow 0$ as $k \rightarrow \infty$. We conclude that $b_k \not\rightarrow 0$ as $k \rightarrow \infty$ and hence $\sum b_k$ diverges by the k th term test for divergence.

4.7.1 Radius of Convergence

In the previous example, the power series, in powers of x , converged for $|x| < 3$. We call the number 3 the **radius of convergence** of the power series. It is half the length of the interval of convergence.

Definition 4.7.5. If a power series of the form $\sum_{k=0}^{\infty} a_k(x-a)^k$ converges at all points in some interval $(-R+a, R+a)$, or equivalently, for $|z-a| < R$, then the number R is called the *radius of convergence* for the power series. The corresponding interval $(-R+a, R+a)$ is called the *open interval of convergence* for the power series. If the power series converges for all real x , we say that the radius of convergence is infinite.

Notes: 1. The term ‘radius’ is used since, when x is replaced by the complex variable z , the open interval is replaced by an open disc, $|z-a| < R$, in the Argand plane. The number R then is the radius of this open disc.

2. It is easiest to find the interval of convergence first, using the ratio test, and then write down the radius of convergence. It can be shown that $R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$, provided this limit exists. There are, however, power series which have a radius of convergence, but for which this limit does not exist.

By generalising the solution to Example 4.7.3 and the argument in Remark 4.7.4, one obtains the following theorem.

Theorem 4.7.6. Suppose that $\{a_k\}_{k=0}^{\infty}$ is a sequence of real numbers such that

$$\lim_{k \rightarrow \infty} \left| \frac{a_k}{a_{k+1}} \right| = R$$

for some real number R . Then the power series $\sum_{k=0}^{\infty} a_k(x-a)^k$

4.7. POWER SERIES

145

(i) converges absolutely whenever $|x - a| < R$, and

(ii) diverges whenever $|x - a| > R$.

Proof of Theorem 4.7.6. [H] The proof of (i) is similar to the solution of Example 4.7.3. We apply the ratio test to the series

$$\sum_{k=0}^{\infty} |a_k(x - a)^k|.$$

Now

$$r = \lim_{k \rightarrow \infty} \frac{|a_{k+1}(x - a)^{k+1}|}{|a_k(x - a)^k|} = \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| |x - a|.$$

We have convergence whenever $r < 1$, which corresponds to the condition that

$$\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| |x - a| < 1.$$

By rearranging we find that the series converges absolutely whenever

$$|x - a| < \frac{1}{\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right|} = \lim_{k \rightarrow \infty} \left| \frac{a_k}{a_{k+1}} \right| = R,$$

where R is the limit given in the theorem.

The proof of (ii) is a simple modification of the argument given in Remark 4.7.4. □

Note that we cannot tell from the theorem whether or not the power series converges at the endpoints $a + R$ or $a - R$. Sometimes the power series will converge at one endpoint but not at the other. Other times it will converge at both endpoints or diverge at both endpoints.

Example 4.7.7. Find the largest open interval on which the power series

$$\sum_{k=0}^{\infty} \frac{(5x + 2)^k}{k^2 + 1}$$

will converge.

Solution. We apply the ratio test to the series $\sum_{k=0}^{\infty} \left| \frac{(5x+2)^k}{k^2+1} \right|$. Now

$$\begin{aligned} r &= \lim_{k \rightarrow \infty} \left| \frac{(5x+2)^{k+1}}{(k+1)^2+1} \right| \left| \frac{k^2+1}{(5x+2)^k} \right| \\ &= \lim_{k \rightarrow \infty} \frac{k^2+1}{(k+1)^2+1} |5x+2| \\ &= |5x+2|. \end{aligned}$$

We require that $r < 1$ and so that $|5x + 2| < 1$. Hence

$$-1 < 5x + 2 < 1$$

or in other words,

$$-\frac{3}{5} < x < -\frac{1}{5}.$$

So the largest open interval of convergence is $(-\frac{3}{5}, -\frac{1}{5})$.

Hence the radius of convergence is $\frac{1}{5}$. □

Students studying MATH1241 are also required to determine whether a power series converges at the endpoints of its interval of convergence. As the next example illustrates, one deduces the convergence at each endpoint by substituting the endpoint into the power series and determining whether the resulting series of real numbers converges.

Example 4.7.8. Find the interval of convergence (including endpoints, if appropriate) for the power series $\sum_{k=2}^{\infty} \frac{x^k}{\ln k}$.

Solution. First we find the open interval of convergence. Now

$$\begin{aligned} r &= \lim_{k \rightarrow \infty} \left| \frac{x^{k+1}}{\ln(k+1)} \cdot \frac{\ln k}{x^k} \right| \\ &= \lim_{k \rightarrow \infty} \frac{\ln k}{\ln(k+1)} |x| \\ &= \lim_{k \rightarrow \infty} \frac{1/k}{1/(k+1)} |x| && \text{(by l'Hôpital's rule)} \\ &= \lim_{k \rightarrow \infty} \frac{k+1}{k} |x| \\ &= |x|. \end{aligned}$$

The series converges absolutely whenever $r = |x| < 1$. Hence the largest open interval of convergence is $(-1, 1)$ and the series diverges whenever $|x| > 1$.

Now we determine whether the series converges at the endpoints 1 and -1 . When $x = 1$, the series becomes

$$\sum_{k=2}^{\infty} \frac{1}{\ln k},$$

which diverges by comparison with the harmonic series $\sum \frac{1}{k}$. When $x = -1$, the series becomes

$$\sum_{k=2}^{\infty} \frac{(-1)^k}{\ln k},$$

which is alternating and converges (conditionally) by the alternating series test.

Hence the interval of convergence for the power series is $[-1, 1)$. (Note that the largest interval on which the power series is *absolutely* convergent is $(-1, 1)$.) □

4.8 Manipulation of power series

(Ref: SH10 §12.9)

In this section we investigate what sense (if any) can be made of adding, multiplying, differentiating and integrating power series. Since differentiation and integration are operations applied to functions, it is most natural to approach this investigation by viewing a power series as a function.

Suppose that a power series $\sum_{n=0}^{\infty} a_n x^n$ converges in the interval $(-R, R)$, where R is its radius of convergence. Then one can define a function $f : (-R, R) \rightarrow \mathbb{R}$ by the formula

$$f(x) = \sum_{k=0}^{\infty} a_k x^k \quad \text{whenever } |x| < R.$$

Thus the value of f at each point x is a convergent sum of real numbers. Sometimes it is possible to find a *closed* form for f , but other times we must approximate each value $f(x)$ by using partial sums.

Example 4.8.1. Suppose that f is given by the rule

$$f(x) = \sum_{k=0}^{\infty} x^k.$$

By using, say, the ratio test, we see that the series converges whenever $|x| < 1$ and diverges when $|x| > 1$. Hence natural domain for f is $(-1, 1)$.

In fact, by summing the geometric series, we find that $f(x) = \frac{1}{1-x}$ whenever $|x| < 1$. This is the *closed form* of $f(x)$.

Example 4.8.2. Suppose that f is defined by the rule

$$f(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

We instantly recognise this series as the Maclaurin series for the exponential function. Since this series converges on $\mathbb{R}$, the maximal domain of f is $\mathbb{R}$. The closed form of $f(x)$ is given by $f(x) = e^x$ whenever $x \in \mathbb{R}$.

Example 4.8.3. Suppose that f is defined by the rule

$$f(x) = \sum_{k=1}^{\infty} \frac{x^k}{k^2}.$$

By the ratio test, we find that the series converges whenever $|x| < 1$ and diverges whenever $|x| > 1$. Therefore we take the domain of f to be $(-1, 1)$. (Students in MATH1141 will note that the power series also converges when $x = 1$ and $x = -1$. So the domain could be extended to $[-1, 1]$.)

There seems to be no obvious closed form for $f(x)$ whenever $|x| < 1$. How, then, does one evaluate $f(-\frac{1}{2})$? Note that

$$\begin{aligned} f(-\tfrac{1}{2}) &= \sum_{k=1}^{\infty} \frac{(-1)^k}{2^k k^2} \\ &= -\frac{1}{2^1 1^2} + \frac{1}{2^2 2^2} - \frac{1}{2^3 3^2} + \frac{1}{2^4 4^2} - \frac{1}{2^5 5^2} + \cdots \end{aligned}$$

and that the right-hand side is an alternating series. Thus

$$f(-\tfrac{1}{2}) \approx -\frac{1}{2^1 1^2} + \frac{1}{2^2 2^2} - \frac{1}{2^3 3^2} + \frac{1}{2^4 4^2} - \frac{1}{2^5 5^2} + \cdots + \frac{1}{2^{10} 10^2},$$

where the absolute error in this approximation is less than $\frac{1}{2^{11} 11^2}$ by Corollary 4.5.22.

It turns out that power series are very well behaved as functions defined on their interval of convergence. Given two power series with the same interval of convergence, you can add, subtract and multiply them together in the ‘natural’ way. Power series are also differentiable and integrable, and their derivatives and antiderivatives can also be expressed as power series in the ‘natural’ way. The following theorems articulate the precise details. Their proofs are given later in Subsection 4.8.1.

Theorem 4.8.4. *Suppose that the functions $f : I \rightarrow \mathbb{R}$ and $g : I \rightarrow \mathbb{R}$ are defined by*

$$f(x) = \sum_{k=0}^{\infty} a_k(x-a)^k \quad \text{and} \quad g(x) = \sum_{k=0}^{\infty} b_k(x-a)^k,$$

where both power series converge on the interval I . Then, whenever $x \in I$,

$$(f+g)(x) = \sum_{k=0}^{\infty} (a_k + b_k)(x-a)^k$$

and

$$(fg)(x) = \sum_{k=0}^{\infty} c_k(x-a)^k, \tag{4.20}$$

where

$$c_k = \sum_{j=0}^k a_j b_{k-j}.$$

Remark 4.8.5. The product formula (4.20) says that

$$\begin{aligned} (fg)(x) = & a_0 b_0 + (a_0 b_1 + a_1 b_0)(x-a) + (a_0 b_2 + a_1 b_1 + a_2 b_0)(x-a)^2 \\ & + (a_0 b_3 + a_1 b_2 + a_2 b_1 + a_3 b_0)(x-a)^3 + \cdots \end{aligned}$$

This is the natural generalisation of polynomial multiplication.

Theorem 4.8.6. *Suppose that $f : I \rightarrow \mathbb{R}$ is defined by*

$$f(x) = \sum_{k=0}^{\infty} a_k(x-a)^k$$

whenever $x \in I$, where I denotes the open interval of convergence for the power series. Then

(i) f is differentiable on I and

$$f'(x) = \sum_{k=1}^{\infty} k a_k (x-a)^{k-1}$$

whenever $x \in I$; and

(ii) f is integrable on I and an antiderivative F for f is given by

$$F(x) = \sum_{k=0}^{\infty} \frac{a_k}{k+1} (x-a)^{k+1} + C$$

whenever $x \in I$, where C is a constant.

Remark 4.8.7. This theorem says that a power series can be differentiated and integrated ‘term by term’ inside its open interval of convergence. Another way of saying this is that

$$\frac{d}{dx} \left(\sum_{k=1}^{\infty} a_k (x-a)^k \right) = \sum_{k=1}^{\infty} \left(\frac{d}{dx} a_k (x-a)^k \right)$$

and

$$\int \left(\sum_{k=1}^{\infty} a_k (x-a)^k \right) dx = \sum_{k=1}^{\infty} \left(\int a_k (x-a)^k dx \right).$$

Students should note that one cannot always swap infinite summation with differentiation (or with integration). For example, if

$$f(x) = \sum_{k=1}^{\infty} \frac{\sin(2^k x)}{2^k}$$

then f is a sum of differentiable functions but f is not differentiable anywhere!

The following corollary follows from Theorem 4.8.6.

Corollary 4.8.8. Suppose that $f : I \rightarrow \mathbb{R}$ is defined by

$$f(x) = \sum_{k=0}^{\infty} a_k (x-a)^k$$

whenever $x \in I$, where I denotes the open interval of convergence for the power series. Then f is continuous on I and has derivatives of all orders on I .

Proof. By Theorem 4.8.6 (i), f is differentiable on I and is therefore continuous on I .

We now prove that f has derivatives of all orders on I . Suppose that n is any natural number. It suffices to show that f is n -times differentiable on I . By Theorem 4.8.6 (i), f is once differentiable on I and its derivative f' has a power series expansion that converges on I . Now apply Theorem 4.8.6 (i) to f' . We conclude that f' is differentiable on I and its derivative f'' has a power series expansion that converges on I . Now apply Theorem 4.8.6 (i) to f'' . Continuing in this way, after n steps we conclude that $f^{(n-1)}$ is differentiable on I with derivative $f^{(n)}$. Hence f is n -times differentiable on I , thus completing the proof. $\square$

Remark 4.8.9. Suppose that a function $f : I \rightarrow \mathbb{R}$ is defined by

$$f(x) = \sum_{k=0}^{\infty} a_k (x-a)^k,$$

where I is the open interval of convergence for the power series. By the corollary, f has derivatives of all orders at a and therefore has a Taylor series about a . One can easily show that the Taylor series for f about a converges on I to f . Thus the Taylor series for f about a is equal to the power series that defines f .

Theorems 4.8.4 and 4.8.6 allow us to find valid Taylor series expansions of functions without having to derive the Taylor coefficients and verify that the remainder term from Taylor’s theorem vanishes.

Example 4.8.10. Given the Taylor expansions

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \cdots \quad (x \in \mathbb{R}) \quad (4.21)$$

and

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + \cdots \quad (|x| < 1), \quad (4.22)$$

find Taylor expansions for each function f , making sure that you state the interval of convergence.

$$(a) f(x) = \cosh x \quad (b) f(x) = \sinh x \quad (c) f(x) = \tan^{-1}(x)$$

Solution. (a) First note that $\cosh x = \frac{1}{2}(e^x + e^{-x})$. So we aim to add the Maclaurin series for e^x and e^{-x} . By replacing x with $-x$ in (4.21), we find that

$$e^{-x} = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \frac{x^5}{5!} + \frac{x^6}{6!} - \cdots \quad (x \in \mathbb{R}).$$

So

$$e^x + e^{-x} = 2 + 2\frac{x^2}{2!} + 2\frac{x^4}{4!} + 2\frac{x^6}{6!} + \cdots \quad (x \in \mathbb{R})$$

by Theorem 4.8.4. Hence

$$\begin{aligned} \cosh x &= \frac{1}{2}(e^x + e^{-x}) \\ &= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \cdots \end{aligned}$$

whenever $x \in \mathbb{R}$.

(b) By differentiating both sides of the expansion

$$\cosh x = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \frac{x^8}{8!} + \cdots \quad (x \in \mathbb{R})$$

we find that

$$\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \cdots$$

whenever $x \in \mathbb{R}$.

(c) If $|x| < 1$ then $|-x^2| < 1$. So we can replace x with $-x^2$ in (4.22) to obtain the convergent expansion

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + x^8 - x^{10} + \cdots$$

whenever $|x| < 1$. By integrating both sides this of identity, we find that

$$\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \frac{x^9}{9} - \frac{x^{11}}{11} + \cdots$$

whenever $|x| < 1$. □

The function f , given by $f(x) = e^{-x^2}$, has no antiderivative among the elementary functions. In MATH1131, we used Riemann sums to estimate the area underneath the graph of f . As the next example shows, the use of Taylor series provides a more efficient approach to the same problem.

Example 4.8.11. Suppose that $f(x) = e^{-x^2}$. By using the Maclaurin expansion for f , estimate $\int_0^1 f(x) dx$ and give an upper bound for the absolute error.

Solution. We begin with the Maclaurin expansion

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \cdots,$$

which is valid for all x in $\mathbb{R}$. By replacing x with $-x^2$ we find that

$$e^{-x^2} = 1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \frac{x^8}{4!} - \frac{x^{10}}{5!} + \cdots$$

for all real numbers x . By integrating both sides of this equation on the interval $[0, 1]$, we find that

$$\begin{aligned} \int_0^1 e^{-x^2} dx &= \left[x - \frac{x^3}{3} + \frac{x^5}{5(2!)} - \frac{x^7}{7(3!)} + \frac{x^9}{9(4!)} - \frac{x^{11}}{11(5!)} + \cdots \right]_0^1 \\ &= 1 - \frac{1}{3} + \frac{1}{5(2!)} - \frac{1}{7(3!)} + \frac{1}{9(4!)} - \frac{1}{11(5!)} + \cdots. \end{aligned}$$

Hence the integral is expressed as alternating series of the form $\sum(-1)^k a_k$, where $\{a_k\}$ is a positive decreasing sequence. If we estimate the series using a partial sum, then Corollary 4.5.22 gives an upper bound for the absolute error. For example, the absolute error in the approximation

$$\int_0^1 e^{-x^2} dx \approx 1 - \frac{1}{3} + \frac{1}{5(2!)} - \frac{1}{7(3!)}$$

is no greater than $\frac{1}{9(4!)}$. One can evaluate this partial sum numerically to obtain

$$\int_0^1 e^{-x^2} dx \approx 0.7429$$

with an error no greater than 0.005. □

4.8.1 Proof of theorems in Section 4.8 [X]

In this subsection, we prove Theorems 4.8.4 for the case where $a = 0$. It is not hard to adapt the presented proofs to the general case.

Proof of Theorem 4.8.4. We shall only prove the product formula (4.20) when $a = 0$. Fix x in I and define the partial sums $s_n(x)$ and $t_n(x)$ by

$$s_n(x) = \sum_{k=0}^n a_k x^k \quad \text{and} \quad t_n(x) = \sum_{k=0}^n b_k x^k.$$

Then $f(x) = \lim_{n \rightarrow \infty} s_n(x)$ and $g(x) = \lim_{n \rightarrow \infty} t_n(x)$. So using Proposition 4.3.6, we find that the sequence $\{s_n(x)t_n(x)\}$ converges and

$$(f.g)(x) = f(x).g(x) = \lim_{n \rightarrow \infty} s_n(x) \times \lim_{n \rightarrow \infty} t_n(x) = \lim_{n \rightarrow \infty} (s_n(x)t_n(x)).$$

$$\begin{aligned}
s_n(x)t_n(x) &= (a_0 + a_1x + a_2x^2 + \cdots + a_nx^n) \times (b_0 + a_1x + b_2x^2 + \cdots + b_nx^n) \\
&= a_0b_0 + (a_0b_1 + a_1b_0)x + (a_0b_2 + a_1b_1 + a_2b_0)x^2 \\
&\quad + \cdots + (a_0b_n + a_1b_{n-1} + \cdots + a_{n-1}b_1 + a_nb_0)x^n.
\end{aligned}$$

As $n \rightarrow \infty$, one obtains (4.20). $\square$

We move now to the proof of Theorem 4.8.6, which shall be broken into two parts. First we prove the differentiation result, which is difficult and uses the mean value theorem. After this, the integration result can be easily deduced from the first part.

Proof of Theorem 4.8.6 (i). Suppose that $f : (-R, R) \rightarrow \mathbb{R}$ is defined by

$$f(x) = \sum_{k=0}^{\infty} a_k x^k$$

whenever $x \in (-R, R)$, where R is the radius of convergence for the power series. It can be easily shown (see the tutorial problems) that the radius of convergence for the power series

$$\sum_{k=1}^{\infty} k a_k x^{k-1}$$

is also R . So define the function $g : (-R, R) \rightarrow \mathbb{R}$ by the formula

$$g(x) = \sum_{k=1}^{\infty} k a_k x^{k-1}.$$

Fix, now, a number x in $(-R, R)$. Our task is to show that $f'(x) = g(x)$, or in other words, that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = g(x).$$

Now if $x+h \in (-R, R)$ and $h \neq 0$ then

$$\begin{aligned}
\left| g(x) - \frac{f(x+h) - f(x)}{h} \right| &= \left| \sum_{k=1}^{\infty} k a_k x^{k-1} - \sum_{k=0}^{\infty} \frac{a_k (x+h)^k - a_k x^k}{h} \right| \\
&= \left| \sum_{k=1}^{\infty} k a_k x^{k-1} - \sum_{k=1}^{\infty} a_k \left(\frac{(x+h)^k - x^k}{h} \right) \right|.
\end{aligned}$$

By the mean value theorem,

$$\frac{(x+h)^k - x^k}{h} = k c_k^{k-1}$$

for some real number c_k between x and $x+h$. Hence

$$\begin{aligned}
\left| g(x) - \frac{f(x+h) - f(x)}{h} \right| &= \left| \sum_{k=1}^{\infty} k a_k x^{k-1} - \sum_{k=1}^{\infty} k a_k c_k^{k-1} \right| \\
&= \left| \sum_{k=1}^{\infty} k a_k (x^{k-1} - c_k^{k-1}) \right| \\
&= \left| \sum_{k=2}^{\infty} k a_k (x^{k-1} - c_k^{k-1}) \right|.
\end{aligned}$$

$$\frac{x^{k-1} - c_k^{k-1}}{x - c_k} = (k-1)d_{k-1}^{k-2}$$

for some real number d_{k-1} between x and c_k . Hence

$$\left| x^{k-1} - c_k^{k-1} \right| = |x - c_k| \left| (k-1)d_{k-1}^{k-2} \right|.$$

Now $|x - c_k| < |h|$ and $|d_{k-1}| < M$, where $M = \max\{|x|, |x+h|\}$. So

$$\left| x^{k-1} - c_k^{k-1} \right| \leq |h| \left| (k-1)M^{k-2} \right|.$$

Thus

$$\left| g(x) - \frac{f(x+h) - f(x)}{h} \right| \leq |h| \sum_{k=2}^{\infty} \left| k(k-1)a_k M^{k-2} \right|.$$

One can show using the ratio test that the series on the right-hand side converges and hence

$$\lim_{h \rightarrow 0} |h| \sum_{k=2}^{\infty} \left| k(k-1)a_k M^{k-2} \right| = 0.$$

Therefore

$$\lim_{h \rightarrow 0} \left| g(x) - \frac{f(x+h) - f(x)}{h} \right| = 0$$

(by the pinching theorem for limits) and we conclude that

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = g(x).$$

Hence $f'(x) = g(x)$ for all x in $(-R, R)$. □

Now that we have proved that a power series is differentiable inside its open interval of convergence, it is relatively easy to prove that it is integrable inside this interval.

Proof of Theorem 4.8.6 (ii). Suppose that $f : (-R, R) \rightarrow \mathbb{R}$ is defined by

$$f(x) = \sum_{k=0}^{\infty} a_k x^k$$

whenever $x \in (-R, R)$, where R is the radius of convergence for the power series. It can be easily shown (see the tutorial problems) that the radius of convergence for the power series

$$\sum_{k=0}^{\infty} \frac{a_k}{k+1} x^{k+1}$$

is also R . So define the function $F : (-R, R) \rightarrow \mathbb{R}$ by the formula

$$F(x) = \sum_{k=0}^{\infty} \frac{a_k}{k+1} x^{k+1}.$$

We now apply the differentiability theorem (Theorem 4.8.6 (i)) to F . In particular,

$$\begin{aligned} F'(x) &= \sum_{k=0}^{\infty} \frac{d}{dx} \left(\frac{a_k}{k+1} x^{k+1} \right) \\ &= \sum_{k=0}^{\infty} a_k x^k \\ &= f(x) \end{aligned}$$

whenever $x \in (-R, R)$. Hence F is an antiderivative for f on $(-R, R)$ and hence f is integrable on $(-R, R)$. $\square$

4.9 Maple notes

The following MAPLE command is relevant to the material of this chapter:

`sum(f(k) , k=m..n);` computes the sum of $f(k)$ as k runs from m to n . For example,

`> sum(k^2, k=1..4);`

30

`> sum(k^2, k=1..n);`

$$\frac{1}{3}(n+1)^3 - \frac{1}{2}(n+1)^2 + \frac{1}{6}n + \frac{1}{6}$$

`> sum(1/k^2, k=1..infinity);`

$$\frac{1}{6}\pi^2$$

`?powseries` will give information about the MAPLE package for manipulating formal power series.

`taylor(expr, x=a, k);` computes the Taylor series for expr about $x=a$, up to the term of order k .

`convert(taylor(expr, x=a, k), polynom);` computes the Taylor polynomial of order $k-1$ for expr about $x=a$.

`coeftayl(expr, x=a, k);` computes the k th coefficient in the Taylor series expansion of expr about $x=a$.

For example,

`> taylor(sin(x) ,x=0,8);`

$$x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + O(x^8)$$

`> convert(%,polynom);`

$$x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7$$

`> coeftayl(sin(x),x=0,11);`

$$-\frac{1}{39916800}$$

Problems 4.1 : Taylor polynomials

1. [R] For each function f , find the Taylor polynomial of degree 9 for f about 0.
 - a) $f(x) = e^x$
 - b) $f(x) = \sin x$
 - c) $f(x) = \sinh x$
2. [R] Suppose that $f(x) = \sin x$ and $m \geq 0$. Using summation notation, find a formula for the Taylor polynomial p_{2m+1} of degree $2m + 1$ for f about 0.
3. [R]
 - a) Suppose that $f(x) = \sqrt{x}$. Find the Taylor polynomial of degree 3 for f about 4.
 - b) Suppose that $g(x) = \cos x$. Find the Taylor polynomial of degree 4 for g about $\pi/4$.
4. [R] Let $f(x) = 1 + x + x^2$. Find the Taylor polynomial $p_n(x)$
 - a) of degree $n = 1$ about 1.
 - b) of degree $n = 2$ about 1.
 - c) of degree $n = 2$ about 2.

Problems 4.2 : Taylor's theorem

5. [R] Suppose that $f(x) = \ln(1 + x)$.
 - a) Express $f(x)$ in the form $p_1(x) + R_2(x)$, where p_1 is the first Taylor polynomial for f about 0 and R_2 is the Lagrange formula for the remainder.
 - b) Suppose that $x \in [-0.1, 0.1]$ and consider the approximation $\ln(1 + x) \approx x$. Use your answer to (a) to show that an upper bound for the absolute error in this approximation is $1/162$.
6. [R] Suppose that $f(x) = \sqrt{1 + x}$ and let p_2 denote the second Taylor polynomial for f about 0. If $x \in [0, 1]$ then show that the absolute error in the approximation $f(x) \approx p_2(x)$ does not exceed $\frac{1}{16}$.
7. [R] Suppose that $f(x) = \cos x$ and that n is a positive even integer.
 - a) Find the n th Taylor polynomial p_n for f about 0 and the Lagrange formula for the remainder R_{n+1} .
 - b) Use the mean value theorem to prove that

$$\sin x < x \quad \text{whenever } x > 0.$$
 - c) Use parts (a) and (b) to find an upper bound for the absolute error in the approximation $f(1/10) \approx p_n(1/10)$.

- d) Hence find a value for n such that the absolute error in the approximation of (c) is less than 10^{-6} .
- e) The value of $\cos 2$ is estimated using $p_n(2)$ for some n . Explain why it is better *not* to use the inequality of (b) to find an upper bound for $|R_{n+1}(2)|$.
- f) Find a value for n such that the absolute error in the approximation $f(2) \approx p_n(2)$ is less than 10^{-6} .
- g) The Taylor polynomial p_{10} is used to approximate f on the interval $[-a, a]$, where a is a positive real number. Find a value for a such that the absolute error in the approximation $f(x) \approx p_{10}(x)$ is less than 10^{-6} whenever $x \in [-a, a]$.

8. [X] Below is a statement of the **mean value theorem for integrals**:

If f and g are continuous on $[a, b]$ and g is nonnegative on $[a, b]$, then

$$\int_a^b f(t)g(t) dt = f(c) \int_a^b g(t) dt$$

for some real number c in $[a, b]$.

This theorem is used in several of the problems below. We give a proof of it in this question.

- a) Suppose that f is continuous on $[a, b]$ and let m and M denote the minimum and maximum values of f on $[a, b]$ respectively. If $m \leq z \leq M$, then explain why there exists a real number c in $[a, b]$ such that $f(c) = z$.
 - b) By considering a lower and upper bound for the integral $\int_a^b f(t)g(t) dt$ in terms of m and M , prove the mean value theorem for integrals.
9. [X] In this question, we will prove the Lagrange formula for the remainder under some simple assumptions. Suppose that f is $(n+1)$ -times differentiable on some open interval I containing 0 and that $f^{(n+1)}$ is continuous on I . Suppose that $x \in I$ and recall that the remainder term in Taylor's theorem is given by

$$R_{n+1}(x) = \frac{1}{n!} \int_0^x f^{(n+1)}(t)(x-t)^n dt.$$

Use the mean value theorem for integrals (see Question 8) to deduce that

$$R_{n+1}(x) = \frac{f^{(n+1)}(c)}{(n+1)!} x^{n+1},$$

where c is some real number between 0 and x .

10. [R]

- a) Suppose that

$$f(x) = x^7 + 5x^6 + 3x^5 - 17x^4 - 16x^3 + 24x^2 + 16x - 11.$$

Verify that 1 and -2 are stationary points for f and use the corollary to Taylor's theorem to classify each of these stationary points.

b) Suppose that

$$f(x) = x^7 - 7x^6 + 10x^5 + 22x^4 - 43x^3 - 35x^2 + 48x + 40.$$

Verify that -1 , 2 and 3 are stationary points for f and classify each of these stationary points.

11. [H] Use the following outline to show that e is irrational.

a) If e were rational, it would be of the form $e = \frac{p}{q}$, where p and q are positive integers. Select an integer k such that $k \geq 3$ and $k \geq q$. Use Taylor's Theorem to show that

$$\frac{p}{q} = e = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots + \frac{1}{k!} + \frac{e^z}{(k+1)!}$$

for some z in $[0, 1]$.

b) Suppose that

$$s_k = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots + \frac{1}{k!}.$$

Show that $k!(e - s_k)$ is an integer.

c) Show that $0 < k!(e - s_k) < 1$.

d) Conclude that e is irrational.

Problems 4.3 : Sequences

12. [R] Describe the limiting behaviour of the following sequences. If the sequence converges, then state its limit.

a) $\frac{n^2 - 2n + 1}{2n^2 + 4n - 1}$

b) $\frac{\ln n}{n^a}$, where $a > 0$.

c) $\frac{n \sin(n\pi/4)}{\sqrt{n^2 + 1}}$

d) $\frac{n + \cos n\pi}{n - \cos n\pi}$

e) $\frac{n!}{n^n}$

f) $\frac{(2n)!}{(n!)^2}$

g) $(a^n + b^n)^{1/n}$, where $a \geq b > 0$.

13. [H] Suppose that $a_n = \frac{n^2 - 2n + 1}{2n^2 + 4n - 1}$.

a) Find L , where $L = \lim_{n \rightarrow \infty} a_n$.

b) For each positive number ϵ , find a number N such that

$$|a_n - L| < \epsilon \quad \text{whenever } n > N.$$

14. [R] Use the result $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$ and standard results about limits to evaluate the following limits.

a) $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^{4n}$

b) $\lim_{n \rightarrow \infty} \left(\frac{n}{n+1}\right)^n$

15. [X] *The limit of a recursively defined sequence.*

Suppose that $a_1 = 1$ and $a_{n+1} = \sqrt{1 + a_n}$ whenever $n \geq 1$.

- Show that $\sqrt{1+x} \in [1, 2]$ whenever $x \in [1, 2]$.
- Use induction to show that the sequence $\{a_n\}$ is bounded.
- Use induction to show that $\{a_n\}$ is an increasing sequence.
- Explain why $\lim_{n \rightarrow \infty} a_n$ exists.
- Find $\lim_{n \rightarrow \infty} a_n$. (That is, find $\sqrt{1 + \sqrt{1 + \sqrt{1 + \cdots}}}$.)

16. [X] Find the supremum and infimum of each of the following sets.

- $\left\{ \frac{n}{1+n^2} : n = 1, 2, \dots \right\}$
- $\left\{ \frac{n}{1+n^2} : n \in \mathbb{Z} \right\}$
- $\{x \in \mathbb{Q} : x^2 < 2\}$
- $\left\{ \frac{(-1)^n}{n} + \sin n : n = 1, 2, \dots \right\}$
- $\{x \in (0, \infty) : \sin x < 0\}$
- $\{1 + \tan^{-1} x : x \in \mathbb{R}\}$

17. [X] Suppose that A and B are nonempty subsets of $\mathbb{R}^2$. Define the distance $d(A, B)$ between A and B by the formula

$$d(A, B) = \inf\{|\mathbf{a} - \mathbf{b}| : \mathbf{a} \in A, \mathbf{b} \in B\}.$$

- Explain why this infimum always exists.
 - Suppose that $A = \{(x, y) : x^2 + y^2 < 1\}$ and $B = \{(x, y) : x^2 - y^2 > 9\}$. Find $d(A, B)$.
 - Find two disjoint sets A and B such that $d(A, B) = 0$.
18. [X] Suppose that $\{a_n\}_{n=1}^\infty$ is a bounded sequence. Given a positive integer m , define K_m by the formula

$$K_m = \sup\{a_m, a_{m+1}, a_{m+2}, \dots\}.$$

- Explain why $K_m \geq \inf_{n \geq 1} a_n$ for all m .
- Explain why $K_{m+1} \leq K_m$ for all m .
- Explain why the limit $\lim_{m \rightarrow \infty} K_m$ exists.

Let K denote the limit of part (c). We call K the **limit superior** of the sequence and write $K = \limsup_n a_n$. It is the largest number to which any subsequence of $\{a_n\}$ converges. The smallest number to which any subsequence converges is called the **limit inferior**, $\liminf_n a_n = \lim_{m \rightarrow \infty} \left(\inf_{n \geq m} a_n \right)$.

- Prove that $\liminf_n a_n = \limsup_n a_n$ if and only if $\{a_n\}$ converges.

19. [R] *Examples of telescoping series.*

- a) Consider the series $\sum_{k=1}^{\infty} a_k$, where $a_k = \frac{1}{k(k+1)}$.
- Find the partial fractions decomposition of a_k .
 - Find a simple formula for the partial sum s_n , where $s_n = \sum_{k=1}^n a_k$.
 - Hence find $\sum_{k=1}^{\infty} a_k$.
- b) Repeat the question for the series $\sum_{k=2}^{\infty} a_k$, where $a_k = \frac{1}{k^2 - 1}$.

20. [R] Let s_n denote the n th partial sum of the series

$$1 + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \cdots.$$

- Show that $s_n > \sqrt{n}$ whenever $n > 1$.
- Hence explain why the series diverges.

21. [R] Let s_n denote the n th partial sum of the harmonic series series $\sum_{k=1}^{\infty} \frac{1}{k}$. When $n = 2^{k-1}$, the terms of s_n may be bracketed as shown:

$$s_n = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \cdots + \frac{1}{8}\right) + \cdots + \left(\frac{1}{2^{k-2}+1} + \frac{1}{2^{k-2}+2} + \cdots + \frac{1}{2^{k-1}}\right).$$

Hence use an argument similar to that in Question 20 to show that the harmonic series diverges.

22. [R] The following three questions are similar to Example 4.4.3.

- By drawing a diagram and interpreting each term in sum as the area of rectangle, show that

$$\sum_{k=2}^n \frac{1}{k^2} \leq \int_1^n \frac{dx}{x^2}.$$

- Deduce that $\sum_{k=2}^{\infty} \frac{1}{k^2}$ converges.

23. [R]

- By drawing a diagram and interpreting each term in sum as the area of rectangle, show that

$$\sum_{k=1}^n \frac{1}{\sqrt{k}} \geq \int_1^{n+1} \frac{dx}{\sqrt{x}}.$$

- b) Deduce that $\sum_{k=1}^{\infty} \frac{1}{\sqrt{k}}$ diverges.
24. [R] By using the technique of the previous two questions, determine whether or not the sum $\sum_{k=2}^{\infty} \frac{1}{k \ln k}$ converges.

Problems 4.5 : Tests for series convergence

25. [R] Use the integral test to examine the convergence of
- a) $\sum_{k=1}^{\infty} \frac{k}{k^2 + 4}$ b) $\sum_{k=3}^{\infty} \frac{1}{k(\ln k)}$ c) $\sum_{k=1}^{\infty} \frac{1}{(k+9)^3}$
26. [R] Use a comparison test to determine whether or not each series converges.
- a) $\sum_{k=1}^{\infty} \frac{1}{(k^3 + 3)^{1/2}}$ b) $\sum_{k=2}^{\infty} \frac{1}{(k^2 - 1)^{1/3}}$ c) $\sum_{k=1}^{\infty} \frac{k}{k^2 - 6}$
- d) $\sum_{k=2}^{\infty} \frac{1}{\ln k}$ e) $\sum_{k=2}^{\infty} \frac{1}{(\ln k)^9}$ f) $\sum_{k=1}^{\infty} \sin^2 \frac{1}{k}$
27. [R] Use the ratio test to examine the convergence of each series.
- a) $\sum_{k=1}^{\infty} \frac{k^2}{2^k}$ b) $\sum_{k=1}^{\infty} \frac{3^k}{k!}$ c) $\sum_{k=1}^{\infty} \frac{k!}{k^k}$ d) $\sum_{k=1}^{\infty} \frac{5^k}{2^k + 4^k}$
28. [R] Determine which of the following alternating series converge. Which are absolutely convergent?
- a) $\sum_{k=3}^{\infty} \frac{(-1)^k}{\ln k}$ b) $\sum_{k=1}^{\infty} \frac{(-1)^k k^k}{(k+1)^k}$
- c) $\sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{\sqrt{k(k^2 - 2)}}$ d) [X] $\sum_{k=2}^{\infty} \frac{(-1)^k}{\sqrt{k} + (-1)^k}$
29. [R] Consider the alternating series $\sum_{k=0}^{\infty} \frac{(-1)^k}{k^3 + 1}$. Let L denote the value of the series and let s_n denote the n th partial sum of the series whenever $n \geq 0$.
- a) Verify that the series is convergent.
- b) Calculate s_4 and give an upper bound for the absolute error in the approximation $L \approx s_4$.
- c) Find a value for n such that the absolute error in the approximation $L \approx s_n$ is less than 10^{-6} .

30. [R] By using an appropriate test, determine whether or not each series converges.

$$\begin{array}{lll} \text{a)} \sum_{k=2}^{\infty} \frac{(-1)^k}{\sqrt{k}} & \text{b)} \sum_{k=1}^{\infty} \frac{3^k}{k^3} & \text{c)} \sum_{k=1}^{\infty} \frac{3^k}{1+7^k} \\ \text{d)} \sum_{k=2}^{\infty} \frac{2^k k!}{k^k} & \text{e)} \sum_{k=2}^{\infty} \frac{(-1)^{k+1} k^2}{\sqrt{4k^4+1}} & \text{f)} \sum_{k=1}^{\infty} \frac{\sin((2k-1)\pi/4)}{2^k} \end{array}$$

31. [X] Determine the convergence or divergence of $\sum a_k$, for each a_k given below.

$$\begin{array}{llll} \text{a)} \frac{\sin k}{k^2} & \text{b)} \frac{\sqrt{k}}{k^2+1} & \text{c)} \frac{k}{(\ln k)^k} & \text{d)} \frac{(\ln k)^3}{\sqrt{k^3-3k^2+1}} \\ \text{e)} \frac{(-1)^k 2^k}{k^3} & \text{f)} \frac{1}{k^{1+\frac{1}{k}}} & \text{g)} \frac{\ln(k!)}{k^3} & \end{array}$$

32. [H] Discuss the convergence of the series

$$\sum_{k=1}^{\infty} \frac{1}{a_1 a_2 \dots a_k} = \frac{1}{a_1} + \frac{1}{a_1 a_2} + \frac{1}{a_1 a_2 a_3} + \dots,$$

where $\{a_k\}$ is a strictly increasing sequence and $a_1 > 0$.

33. [X]

- a) Prove that if $\sum_k a_k^2$ and $\sum_k b_k^2$ converge, then $\sum_k a_k b_k$ converges.
- b) Prove that if $\sum_k a_k^2$ converges, then $\sum_k \frac{a_k}{k}$ converges.

Problems 4.6 : Taylor series

34. [R] Find, in each case, the Maclaurin series for f . Express your answer using summation notation. (You may find your answers to some questions of Problems 4.1 : Taylor polynomials helpful.)

$$\begin{array}{lll} \text{a)} f(x) = e^x & \text{b)} f(x) = \sin x & \text{c)} f(x) = \sinh x \\ \text{d)} f(x) = \ln(1+x) & \text{e)} f(x) = \frac{1}{x-1} & \end{array}$$

35. [R] Suppose that $f(x) = e^x$.

- a) Express f in the form $f(x) = p_n(x) + R_{n+1}(x)$, where p_n is the n th Taylor polynomial for f about 0 and R_{n+1} is the Lagrange formula for the remainder.
- b) Fix x in $\mathbb{R}$ and show that $R_{n+1}(x) \rightarrow 0$ as $n \rightarrow \infty$.
- c) Hence write down the Taylor series expansion for $f(x)$, stating clearly where the expansion is valid.

36. [H] Repeat the previous question in the case when $f(x) = \sinh x$.

37. [H] Suppose that $f(x) = \ln(1+x)$.

- a) Express f in the form $f(x) = p_n(x) + R_{n+1}(x)$, where p_n is the n th Taylor polynomial for f about 0 and R_{n+1} is the Lagrange formula for the remainder.
- b) Suppose that $0 \leq x \leq 1$ and show that $R_{n+1}(x) \rightarrow 0$ as $n \rightarrow \infty$.
- c) Hence write down the Taylor series expansion for $f(x)$, when $0 \leq x \leq 1$.
- d) [X] We will show that this Taylor series expansion is also valid on $(-1, 0)$. Suppose that $-1 < x < 0$.
- i) By using the integral form for the remainder, show that

$$R_{n+1}(x) = \int_0^x \left(\frac{t-x}{1+t} \right)^n \frac{1}{1+t} dt.$$

- ii) Use the mean value theorem for integrals (when $g(t) = 1$; see Question 8) to show that

$$|R_{n+1}(x)| < \left(\frac{c_n + |x|}{1 + c_n} \right)^n \left(\frac{|x|}{1 + x} \right)$$

for some number c_n between x and 0.

- iii) Deduce that

$$|R_{n+1}(x)| < |x|^n \left(\frac{|x|}{1 + x} \right)$$

and hence that $R_{n+1}(x) \rightarrow 0$ as $n \rightarrow \infty$.

38. [H] Let I denote the interval $(x_0 - R, x_0 + R)$, where x_0 and R are real numbers. Suppose that a function f has derivatives of all orders on I and that all these derivatives have a common bound (that is, $|f^{(n)}(x)| \leq M$ for all x in I and for all positive integers n). Show that f is represented by its Taylor series about x_0 on I .

39. [X] Suppose that

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0. \end{cases}$$

- a) Show that f is differentiable everywhere and find a formula for f' .
- b) Show that f' is differentiable everywhere and find a formula for f'' .
- c) Suppose that k is a positive integer. Prove that $\frac{d}{dx} \left(\frac{e^{-1/x^2}}{x^k} \right)$ is a linear combination of functions of the form $\frac{e^{-1/x^2}}{x^m}$, where m is a positive integer.
- d) Hence deduce that $f^{(n)}(0) = 0$ for every natural number n .
- e) Write down the Maclaurin series for f . Where does the Maclaurin series converge? Where does it converge to f ?

40. [H] (This exercise illustrates that a conditionally convergent series, when rearranged, can have a different sum.)

Consider the series s and t , given by

$$s = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots$$

$$t = 1 - \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{6} - \frac{1}{8} + \frac{1}{5} - \frac{1}{10} - \frac{1}{12} + \cdots.$$

Note that second series is a rearrangement of the first.

- Explain why s is conditionally convergent.
- By considering an appropriate Maclaurin series (see Theorem 4.6.5), find the exact value of s .
- Denote the sum of the first n terms of s by s_n and of t by t_n . By using induction, show that $t_{3n} = \frac{1}{2}s_{2n}$ whenever $n \geq 1$.
- Hence find the value of $\lim_{n \rightarrow \infty} t_{3n}$.
- Explain why $\lim_{n \rightarrow \infty} t_{3n+1} = \lim_{n \rightarrow \infty} t_{3n+2} = \lim_{n \rightarrow \infty} t_{3n}$.
- Hence write down the value of t .

Problems 4.7 : Power series

41. [R] Determine the open interval of convergence for each of the following power series. (Students studying MATH1241 should also examine the behaviour of the power series at the endpoints.)

$$\begin{array}{lll} \text{a) } \sum_{k=0}^{\infty} \left(\frac{x}{6}\right)^k & \text{b) } \sum_{k=0}^{\infty} \frac{x^k}{k^2 + 1} & \text{c) } \sum_{k=0}^{\infty} \frac{kx^k}{2^k} \\ \text{d) } \sum_{k=1}^{\infty} \frac{(x-2)^k}{k^3} & \text{e) } \sum_{k=2}^{\infty} \frac{(3x-2)^k}{k \ln k} & \text{f) } \sum_{k=1}^{\infty} \frac{(-1)^k x^k}{(k+1)3^k} \\ \text{g) } [\mathbf{X}] \sum_{k=1}^{\infty} \frac{(\ln k)^k x^k}{k^k} & & \end{array}$$

42. [R] Consider the power series

$$(i) \quad \sum \frac{n!}{n^n} x^n \quad \text{and} \quad (ii) \quad \sum \frac{n^n}{n!} x^n.$$

- Show that their radii of convergence are e and e^{-1} respectively.
- [X] Show, by any method, that $a(n) = \left(1 + \frac{1}{n}\right)^n$ is a strictly increasing sequence (whose limit is e).
- [X] Deduce from your working in (a) and (b) that (i) diverges when $x = \pm e$.
- [X] Here you may assume Stirling's formula:

$$n! \Big/ \left\{ \sqrt{2\pi} n^{n+1/2} e^{-n} \right\} \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

Show that series (ii) diverges when $x = e^{-1}$. Using Stirling's formula and (b), show that (ii) is conditionally convergent when $x = -e^{-1}$.

Problems 4.8 : Manipulation of power series

43. [R] Use your answers to Question 34 to deduce the Maclaurin series for each function g .

a) $g(x) = (x+1)e^x$ b) $g(x) = \sin(x^2)$ c) $g(x) = (x-1)^{-2}$

44. [R] Suppose that $f(x) = (1-x)^{-1}$. Write down the Maclaurin series for f and hence find the Maclaurin series for each function g given below. On what open interval is each function represented by its Maclaurin series?

a) $g(x) = (1+x)^{-1}$ b) $g(x) = (1+x^2)^{-1}$ c) $g(x) = \tan^{-1} x$

45. [R] Suppose that $f(x) = x^2 \sin(x^3)$.

- By using the Maclaurin series for sine, find the Maclaurin series for f .
- Hence show that 0 is a stationary point for f .
- Is 0 a local maximum point, local minimum point or a horizontal point of inflexion? Explain.

46. [R] Consider the Maclaurin series representation

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \cdots,$$

which is valid whenever $|x| < 1$.

- a) By first integrating the above Maclaurin series, deduce that

$$\ln\left(\frac{1+x}{1-x}\right) = 2\left(x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \cdots\right)$$

whenever $|x| < R$, for some real number R .

- What is the largest possible value of R ?
 - Use the first two terms of the series of (a) to find a rational number that approximates $\ln 2$.
47. [R] The function $\text{Si} : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$\text{Si}(x) = \int_0^x f(t) dt,$$

where

$$f(t) = \begin{cases} \frac{\sin t}{t} & \text{if } t \neq 0 \\ 1 & \text{if } t = 0. \end{cases}$$

The Si function is used in signal processing and by surveyors for GPS.

- Show that Si has a stationary point at π and classify this stationary point.
- By first writing down the Maclaurin series for the sine function, find the Maclaurin series for Si.
- Hence find an estimate for the value of $\text{Si}(\pi)$ such that the absolute error is less than $1/100$.

48. [H] A function f is defined by the rule

$$f(x) = \sum_{k=1}^{\infty} kx^k.$$

- a) What is the largest open interval I on which the function well-defined?
- b) By manipulating the power series expansion

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k, \quad |x| < 1,$$

find a closed formula for $f(x)$.

49. [H] Consider the differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = e^x - 1.$$

Suppose that a solution y has a power series representation given by

$$y = \sum_{k=0}^{\infty} a_k x^k,$$

where the coefficients a_k are to be determined.

- a) Write down, in summation notation, the Maclaurin series representation of $e^x - 1$.
- b) Write down the power series representations of $\frac{dy}{dx}$ and $\frac{d^2 y}{dx^2}$.
- c) By expressing both the left- and right-hand sides of the differential equation as a power series, determine the value of each coefficient a_k .
- d) Hence write down a solution to the differential equation, stating the values of x for which the solution is valid.

Note: This method for solving differential equations will be developed in some second year courses.

Chapter 5

Averages, arc length, speed and surface area

In this chapter we look at the application of calculus to the following problems:


- finding the average height of a cable, or the average temperature over a certain time interval;
- finding the length of a curve;
- finding the speed of a particle that travels along a curve in the plane; and
- finding the surface area of certain solids.

Each of these applications involves integration, either through approximating a quantity with a Riemann sum, or through approximating the rate of change of a quantity and thereafter applying the fundamental theorem of calculus.

5.1 The average value of a function

(Ref: SH10 §5.9)

A cable is suspended between two poles as shown.




What is the average height of the cable above the ground? Clearly it must be somewhere between 2 and 3 metres. To obtain a precise answer, recall from MATH1131 that any suspended cable is the graph of a function f of the form


$$f(x) = \frac{1}{c} \cosh(cx)$$

over some interval $[a, b]$, where c is a constant that depends on the tension in and mass of the cable, and where the coordinate system is suitably chosen. Thus we rephrase the question as, ‘What is the average value of f on the interval $[a, b]$?’

To answer this question, we need a suitable definition for the average value of a function f over an interval $[a, b]$. To motivate such a definition, consider the function f whose graph is shown below.



One way to proceed is the following. Divide the interval $[a, b]$ into n subintervals of equal length. We sample the height of the graph in the k th subinterval by choosing a point c_k in that subinterval and calculating $f(c_k)$.



The average (that is, arithmetic mean) a_n of these sampled heights is given by

$$a_n = \frac{1}{n} (f(c_1) + f(c_2) + f(c_3) + \dots + f(c_n)).$$

As the number n of subintervals increases, a_n should get closer to what we intuitively understand by ‘the average height of the graph.’


5.1. THE AVERAGE VALUE OF A FUNCTION

169

Now, by multiplying and dividing a_n by $(b - a)$, we find that

$$\begin{aligned} a_n &= \frac{1}{n} \sum_{k=1}^n f(c_k) \\ &= \frac{1}{n} \sum_{k=1}^n f(c_k) \left(\frac{b-a}{b-a} \right) \\ &= \frac{1}{b-a} \sum_{k=1}^n f(c_k) \left(\frac{b-a}{n} \right). \end{aligned} \quad (5.1)$$

Note that $f(c_k)$ and $\frac{b-a}{n}$ is the height and width of the k th rectangle in the following diagram.



So the sum in (5.1) is a Riemann sum. If f is Riemann integrable then

$$\lim_{n \rightarrow \infty} a_n = \frac{1}{b-a} \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \left(\frac{b-a}{n} \right) = \frac{1}{b-a} \int_a^b f(x) dx.$$

This leads to the following definition.


Definition 5.1.1. Suppose that f is integrable on a closed interval $[a, b]$. Then the *average value* $\bar{f}$ of f on $[a, b]$ is defined by the formula

$$\bar{f} = \frac{1}{b-a} \int_a^b f(x) dx.$$

Remark 5.1.2. By rearranging this formula, we see that $\bar{f}$ is the unique constant such that

$$(b-a)\bar{f} = \int_a^b f(x) dx.$$

Interpreted geometrically, $\bar{f}$ is the unique y -value such that the area of the shaded rectangle is equal to the area under the graph of f .



We now answer the question posed at the beginning of this section.

Example 5.1.3 (The average height of a suspended cable). The suspended cable illustrated at the beginning of this section is a curve given by the equation

$$y = 2 \cosh(x/2), \quad -a \leq x \leq a,$$

where the x -axis runs along the ground, the y -axis passes through the vertex of the curve and $a = 2 \cosh^{-1}(3/2)$. Find, to the nearest centimetre, the average height of the cable above the ground.

Note: The fact that the x -axis runs along the ground is special to this example. This will not necessarily be the case for any given suspended cable.

Solution. (a) Suppose that $f(x) = 2 \cosh(x/2)$. The average value $\bar{f}$ of f on the interval $[-a, a]$ is given by

$$\begin{aligned} \bar{f} &= \frac{1}{a - (-a)} \int_{-a}^a 2 \cosh(x/2) dx \\ &= \frac{2}{2a} \int_0^a 2 \cosh(x/2) dx && \text{(since cosh is even)} \\ &= \frac{1}{a} \left[4 \sinh(x/2) \right]_0^a \\ &= \frac{4}{a} \sinh(a/2) \\ &= \frac{2}{\cosh^{-1}(3/2)} \sinh(\cosh^{-1}(3/2)). \end{aligned}$$

Now $\cosh^2(t) - \sinh^2(t) = 1$ and so

$$\sinh(\cosh^{-1}(3/2)) = \sqrt{\cosh^2(\cosh^{-1}(3/2)) - 1} = \frac{\sqrt{5}}{2}.$$

Hence

$$\bar{f} = \frac{2}{\cosh^{-1}(3/2)} \times \frac{\sqrt{5}}{2}.$$

By using a calculator, we find that $\bar{f} \approx 2.32$. So the average height of the cable above the ground is approximately 2.32 metres. $\square$

5.2. THE ARC LENGTH OF A CURVE

171

We saw in MATH1131 that every continuous function f defined on a closed interval $[a, b]$ attains its maximum and minimum values. (This result is called the maximum-minimum theorem; see Chapter 2 in the MATH1131 calculus notes). The next theorem says that such a function also attains its average value.

Theorem 5.1.4 (The mean value theorem for integrals). *Suppose that f is continuous on $[a, b]$. Then there is a number c in (a, b) such that*

$$\int_a^b f(t) dt = f(c)(b - a).$$

Restated, the conclusion of the mean value theorem for integrals says that there exists a point c in $[a, b]$ such that $f(c) = \bar{f}$, where $\bar{f}$ is the average value of f on $[a, b]$.

Proof. Define $F : [a, b] \rightarrow \mathbb{R}$ by the formula

$$F(x) = \int_a^x f(t) dt.$$

By the fundamental theorem of calculus, F is continuous on $[a, b]$, differentiable on (a, b) and $F'(x) = f(x)$. By the mean value theorem, there exists $c \in (a, b)$ such that

$$\frac{F(b) - F(a)}{b - a} = F'(c). \quad (5.2)$$

But

$$F(a) = 0, \quad F(b) = \int_a^b f(t) dt \quad \text{and} \quad F'(c) = f(c).$$

Hence (5.2) implies that

$$\frac{1}{b - a} \int_a^b f(t) dt = f(c)$$

as required. $\square$

Remark 5.1.5. A more general version of the mean value theorem for integrals is given in the tutorial problems for Chapter 4 and is used to prove the Lagrange formula for the remainder in Taylor's theorem.


5.2 The arc length of a curve

(Ref: SH10 §10.7)

Suppose that $P_0(x_0, y_0)$ and $P_1(x_1, y_1)$ are two points in $\mathbb{R}^2$. The distance between P_0 and P_1 is given by

$$\text{dist}(P_0, P_1) = \sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2}.$$

Suppose that the points P_0 and P_1 are the endpoints of a straight line segment P_0P_1 , as shown.



Then we define the *length* of the line segment P_0P_1 to be the distance between P_0 and P_1 . In other words,

$$\text{length}(P_0P_1) = \sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2}.$$


The line segment P_0P_1 is a special example of a curve in $\mathbb{R}^2$. In this section we study the lengths of curves that are not necessarily straight line segments. We begin by presenting an intuitive derivation of a formula that gives the arc length of a curve. (A rigorous approach to proving the validity of such a formula would involve giving a formal definition for arc length and considering the limits of some technically difficult Riemann sums; we shall not delve into this here.) The subsections following this derivation present variations of this formula and some examples.

5.2.1 An intuitive derivation of the arc length formula

Suppose that $\mathcal{C}$ is a curve in $\mathbb{R}^2$. The goal is to give an heuristic derivation of a formula for the arc length of $\mathcal{C}$. We make the assumption that $\mathcal{C}$ can be expressed in parametric form as

$$\mathcal{C} = \{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq b\},$$

where x and y are differentiable functions of t .




We also assume that the parametrisation is chosen so that the path traversed by the moving point $(x(t), y(t))$ does not retrace its steps (either forwards or backwards).

When $a \leq t \leq b$, let $\ell(s)$ denote the arc length of the curve $\mathcal{C}_s$, given by

$$\mathcal{C}_s = \{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq s\}.$$


The segment $\mathcal{C}_s$ corresponding to the arc length $\ell(s)$ is illustrated in black in the diagram below.



5.2. THE ARC LENGTH OF A CURVE

173

The idea is to take a small segment of the curve and approximate its length with the length of a secant. Suppose that $a < t < b$ and that h is a small real nonzero number. Consider the points $P(x(t), y(t))$ and $Q(x(t+h), y(t+h))$.



The length of the arc from P to Q is approximately equal to the length of secant PQ . That is

$$\ell(t+h) - \ell(t) \approx \sqrt{[x(t+h) - x(t)]^2 + [y(t+h) - y(t)]^2},$$

where the length of the secant is calculated using the distance formula. If we divide both sides by h then

$$\frac{\ell(t+h) - \ell(t)}{h} \approx \sqrt{\left[\frac{x(t+h) - x(t)}{h}\right]^2 + \left[\frac{y(t+h) - y(t)}{h}\right]^2}.$$

This approximation gets better as h gets smaller. If we make the assumption that ℓ is a differentiable function of s then, by taking the limit as h approaches zero, one obtains

$$\ell'(t) = \sqrt{[x'(t)]^2 + [y'(t)]^2}.$$

Hence

$$\ell(s) = \int_a^s \sqrt{[x'(t)]^2 + [y'(t)]^2} dt + K$$

for some constant of integration K , by the fundamental theorem of calculus. To evaluate K , note that $\ell(a) = 0$. So if $s = a$ then

$$0 = \ell(a) = \int_a^a \sqrt{[x'(t)]^2 + [y'(t)]^2} dt + K = 0 + K = K.$$

Hence $K = 0$ and thus

$$\ell(s) = \int_a^s \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Finally, the length of the entire curve is $\ell(b)$. So the arc length of $\mathcal{C}$ is

$$\int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Our findings are summarised at the beginning of the next subsection.

5.2.2 Arc length for a parametrised curve

Suppose that a curve $\mathcal{C}$ can be expressed in parametric form as

$$\mathcal{C} = \{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq b\},$$


where x and y are differentiable functions of t . Then its arc length ℓ is given by the formula

$$\ell = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2} dt \quad (5.3)$$

The next example shows how this formula is applied. We consider the cycloid, which is closely related to the so-called ‘curve of fastest descent’ (see Section 7.2 of the MATH1131 calculus course notes).

Example 5.2.1 (The arc length of a cycloid). Find the arc length of one arch of the cycloid

$$x(t) = r(t - \sin t), \quad y(t) = r(1 - \cos t), \quad 0 \leq t \leq 2\pi.$$



Solution. We begin by calculating the derivatives:

$$x'(t) = r(1 - \cos t), \quad y'(t) = r \sin t.$$

Hence

$$\begin{aligned} [x'(t)]^2 + [y'(t)]^2 &= r^2(1 - 2\cos t + \cos^2 t) + r^2 \sin^2 t \\ &= 2r^2(1 - \cos t), \end{aligned}$$

since $\cos^2 t + \sin^2 t = 1$. We want to substitute this into formula (5.3). Before doing so, it is best to express $1 - \cos t$ as a square. Now $1 - \cos 2\theta = 2\sin^2 \theta$ and so

$$\begin{aligned} [x'(t)]^2 + [y'(t)]^2 &= 2r^2(1 - \cos t) \\ &= 4r^2 \sin^2(t/2). \end{aligned}$$

Hence (5.3) gives

$$\begin{aligned} \ell &= \int_0^{2\pi} \sqrt{[x'(t)]^2 + [y'(t)]^2} dt \\ &= \int_0^{2\pi} \sqrt{4r^2 \sin^2(t/2)} dt. \end{aligned}$$

5.2. THE ARC LENGTH OF A CURVE

At this point we should be careful with taking the square root, since $\sqrt{a^2} = a$ only when $a \geq 0$. Now $\sin(t/2)$ is positive whenever $0 < t < 2\pi$, and so taking the squareroot in the ‘naïve’ way causes no problems. Therefore

$$\begin{aligned}\ell &= \int_0^{2\pi} 2r \sin(t/2) dt \\ &= 2r \left[-2 \cos(t/2) \right]_0^{2\pi} \\ &= 8r.\end{aligned}$$

So the arc length of one arch of the cycloid is $8r$ units. $\square$

Remark 5.2.2. For parametrisations of closed curves, one should be careful with the limits of integration. For example, a circle of radius r and centre $(0, 0)$ may be parametrised as

$$x(t) = r \cos t, \quad y(t) = r \sin t, \quad 0 \leq t \leq 2\pi.$$

Hence

$$\ell = \int_0^{2\pi} \sqrt{[-r \sin t]^2 + [r \cos t]^2} dt = \int_0^{2\pi} r dt = 2\pi r,$$

which shows that the circumference of the circle is $2\pi r$, as expected.

On the other hand, if we use the parametrisation

$$x(t) = r \cos 2t, \quad y(t) = r \sin 2t$$

then

$$\int_0^{2\pi} \sqrt{[-2r \sin 2t]^2 + [2r \cos 2t]^2} dt = \int_0^{2\pi} 2r dt = 4\pi r,$$

which is *not* the circumference of the circle. The reason for this is that, as t varies from 0 to 2π , the point $(x(t), y(t))$ moves around the circle twice! To find the arc length using this parametrisation, one should instead integrate from 0 to π .

5.2.3 Arc length for the graph of a function

Suppose that f is a function of one variable. To find the arc length of the graph of f on the interval $[a, b]$, we parametrise the curve

$$y = f(x)$$

by

$$x(t) = t, \quad y(t) = f(t), \quad a \leq t \leq b.$$

Now

$$x'(t) = 1 \quad \text{and} \quad y'(t) = f'(t).$$

By using the arc length formula (5.3), one finds that

$$\ell = \int_a^b \sqrt{1 + [f'(t)]^2} dt$$

We usually write the variable of integration as x .


In summary, the arc length ℓ of the graph of a function f on the interval $[a, b]$ is given by

$$\ell = \int_a^b \sqrt{1 + [f'(x)]^2} dx. \quad (5.4)$$

The use of this formula is illustrated below. We remind readers that a catenary is the shape of a hanging cable and is described using the hyperbolic cosine function. See Chapter 10 of the MATH1131 calculus notes for further details.

Example 5.2.3 (The arc length of a catenary). Find the arc length of a catenary whose graph is given by

$$y = \frac{1}{a} \cosh(ax), \quad x \in [-b, b].$$



Solution. Suppose that $f(x) = \frac{1}{a} \cosh(ax)$. By formula (5.4),

$$\begin{aligned} \ell &= \int_{-b}^b \sqrt{1 + [f'(x)]^2} dx \\ &= \int_{-b}^b \sqrt{1 + \sinh^2(ax)} dx \\ &= \int_{-b}^b \sqrt{\cosh^2(ax)} dx && (\text{since } \cosh^2 t - \sinh^2 t = 1) \\ &= \int_{-b}^b \cosh(ax) dx && (\text{since } \cosh \text{ is always positive}) \\ &= 2 \int_0^b \cosh(ax) dx && (\text{since } \cosh \text{ is even}) \\ &= 2 \left[\frac{1}{a} \sinh ax \right]_0^b \\ &= \frac{2}{a} \sinh(ab) \\ &= \frac{1}{a} (e^{ab} - e^{-ab}). \end{aligned}$$

So the arc length of the catenary is $\frac{1}{a} (e^{ab} - e^{-ab})$ units. □

Suppose that a curve is described using polar coordinates by

$$r = f(\theta), \quad \theta_0 \leq \theta \leq \theta_1.$$

Since

$$x = r \cos \theta = f(\theta) \cos \theta \quad \text{and} \quad y = r \sin \theta = f(\theta) \sin \theta,$$

we have a parametrisation for the curve in terms of θ . Now

$$x'(\theta) = -f(\theta) \sin \theta + f'(\theta) \cos \theta, \quad \text{while} \quad y'(\theta) = f(\theta) \cos \theta + f'(\theta) \sin \theta.$$

Hence

$$\begin{aligned} [x'(\theta)]^2 + [y'(\theta)]^2 &= [f(\theta)]^2 \sin^2 \theta - 2f(\theta)f'(\theta) \sin \theta \cos \theta + [f'(\theta)]^2 \cos^2 \theta \\ &\quad + [f(\theta)]^2 \cos^2 \theta + 2f(\theta)f'(\theta) \sin \theta \cos \theta + [f'(\theta)]^2 \sin^2 \theta \\ &= [f(\theta)]^2 + [f'(\theta)]^2, \end{aligned}$$

where we have used the fact that $\sin^2 \theta + \cos^2 \theta = 1$. So by using the parametric form (5.3) for arc length, we find that the arc length ℓ of the polar curve is given by


$$\ell = \int_{\theta_0}^{\theta_1} \sqrt{[f(\theta)]^2 + [f'(\theta)]^2} d\theta.$$

We usually write $f(\theta)$ as r and $f'(\theta)$ as $\frac{dr}{d\theta}$.

In summary, the arc length ℓ of a polar curve is given by

$$\ell = \int_{\theta_0}^{\theta_1} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$

Example 5.2.4. The spiral graphed below is given by the polar equation $r = e^{-\theta/10}$, where $\theta \geq 0$. Is the total arc length finite? Explain.



Solution. We begin by calculating the arc length for the segment of the curve when $0 \leq \theta \leq \theta_1$. Now

$$r = e^{-\theta/10} \quad \text{and} \quad \frac{dr}{d\theta} = -\frac{1}{10}e^{-\theta/10}.$$

$$\begin{aligned}
\ell &= \int_0^{\theta_1} \sqrt{(e^{-\theta/10})^2 + (-\frac{1}{10}e^{-\theta/10})^2} d\theta \\
&= \int_0^{\theta_1} \sqrt{(1 + \frac{1}{100}) e^{-2\theta/10}} d\theta \\
&= \frac{\sqrt{101}}{10} \int_0^{\theta_1} e^{-\theta/10} d\theta \\
&= \sqrt{101} (1 - e^{-\theta_1/10}).
\end{aligned}$$

Now as $\theta_1 \rightarrow \infty$, $\ell \rightarrow \sqrt{101}$. Hence the total arc length is finite and equals $\sqrt{101}$ units. $\square$

5.3 The speed of a moving particle

(Ref: SH10 §10.7)

In Chapter 4 of the MATH1131 calculus notes, we discussed the speed of a particle that moves along a straight line. Now we consider the speed of a particle that moves along a curve in the plane.

Suppose that a particle P is moving in the plane and that its position at time t is given by $(x(t), y(t))$. The distance $s(t)$ that the particle has travelled from time zero to any later time t is given by the formula

$$s(t) = \int_0^t \sqrt{[x'(u)]^2 + [y'(u)]^2} du$$

(which is simply the arc length formula for the path that P traverses in this time interval). By definition, the speed of P is the rate of change of its distance with respect to time. So if $v(t)$ denotes the speed of P at time t then

$$v(t) = s'(t) = \sqrt{[x'(t)]^2 + [y'(t)]^2}$$

by the fundamental theorem of calculus.

In summary, the speed $v(t)$ of a particle P at time t is given by

$$\boxed{v(t) = \sqrt{[x'(t)]^2 + [y'(t)]^2}},$$

where the functions x and y give the position $(x(t), y(t))$ of P at time t .

Example 5.3.1. A stone is thrown horizontally from the deck of the Sydney Harbour Bridge at 20 metres per second. Its position $(x(t), y(t))$ exactly t seconds after the stone is thrown is given by

$$x(t) = 20t, \quad y(t) = 50 - 5t^2, \quad 0 \leq t \leq \sqrt{10},$$

where $y(t)$ is the height above the water (see Example 7.2.1 in the MATH1131 calculus notes). Find the speed of the stone an instant before it hits the water.

Solution. We have

$$x'(t) = 20 \quad \text{and} \quad y'(t) = -10t.$$

So the speed $v(t)$ of the stone at time t is given by

$$v(t) = \sqrt{20^2 + 100t^2}$$

5.4. SURFACE AREA

179

whenever $0 < t < \sqrt{10}$. Now the stone hits the water when $y(t) = 0$, which is precisely when $t = \sqrt{10}$. Hence the speed of the stone *an instant before* it hits the water is given by

$$\lim_{t \rightarrow (\sqrt{10})^-} v(t) = \sqrt{20^2 + 100(\sqrt{10})^2} = 10\sqrt{14} \approx 37.42.$$


So the speed of the stone just before it hits the water is approximately 37.42 metres per second. $\square$

5.4 Surface area

(Ref: SH10 §10.8)

The problem of finding the surface area of a surface or solid in $\mathbb{R}^3$ is not easy. In this section, we focus on finding the surface area for a surface (or solid) that is formed by rotating a curve about one of the axes. In the first subsection we derive, intuitively, a formula for the area of such a surface. In the second subsection, the relevant formulae are summarised and examples given.

The formulae presented all rely on the formula for the surface area of the frustum of a right circular cone.



Given a frustum of slant height s and radii r and R , the surface area A of the ‘curved surface’ is given by

$$A = \pi(r + R)s. \quad (5.5)$$

This formula may be proved using elementary methods and is left as an exercise in the tutorial problems.

5.4.1 An heuristic derivation for the surface area of a surface of revolution


Suppose that a curve $\mathcal{C}$ has parametrisation given by

$$\mathcal{C} = \{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq b\}.$$


We will assume that

- the curve lies in the upper half-plane (more precisely, $x(t) \geq 0$ and if the curve meets the x -axis then it does so at only a finite number of points); and
- the curve $\mathcal{C}$ is *simple*: if $(x(t_0), y(t_0)) = (x(t_1), y(t_1))$ then $t_0 = t_1$ (that is, the curve does not intersect itself).

An example of such a curve is shown below.



If we rotate the curve about the axis, then a *surface of revolution* is formed.




The goal of this subsection is to derive a formula for the area of this surface.

Let $A(s)$ denote the area of the surface formed when the curve segment


$$\{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq s\}$$

is rotated about the x -axis. We make the assumption that A is a differentiable function.

Our immediate goal is to compute the derivative of A . Fix t in (a, b) and suppose that h is a small nonzero real number. Consider the points $P(x(t), y(t))$ and $Q(x(t+h), y(t+h))$ and note that $A(t+h) - A(t)$ is the area of the surface formed by rotating the curve segment from P to Q about the x -axis.



Since h is small, this surface area is approximately equal to the area of the surface formed when the secant PQ is rotated about the x axis.



This area may be calculated using formula (5.5), where

$$\text{slant height} = \sqrt{[x(t+h) - x(t)]^2 + [y(t+h) - y(t)]^2}.$$

Hence

$$A(t+h) - A(h) \approx \pi(y(t+h) + y(t)) \sqrt{[x(t+h) - x(t)]^2 + [y(t+h) - y(t)]^2},$$

and dividing both sides by h gives

$$\frac{A(t+h) - A(h)}{h} \approx \pi(y(t+h) + y(t)) \sqrt{\left[\frac{x(t+h) - x(t)}{h}\right]^2 + \left[\frac{y(t+h) - y(t)}{h}\right]^2}.$$

Note that approximation improves as h gets smaller. Moreover, since y is differentiable at t , it follows that y is continuous at t and so $y(t+h) \rightarrow y(t)$ as $h \rightarrow 0$. By taking the limit as h

approaches zero, we obtain

$$\begin{aligned} A'(t) &= \pi(y(t) + y(t))\sqrt{[x'(t)]^2 + [y'(t)]^2} \\ &= 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2}. \end{aligned}$$

This gives an expression for the derivative of A . By applying the fundamental theorem of calculus, we find that

$$A(s) = \int_a^s 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2} dt + K$$

for some constant of integration K . To evaluate K , note that $A(a) = 0$. So the substitution $s = a$ yields

$$0 = A(a) = \int_a^a 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2} dt + K = 0 + K = K.$$

Thus $K = 0$ and hence

$$A(s) = \int_a^s 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Finally, the substitution $s = b$ yields

$$A(b) = \int_a^b 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2} dt,$$

which is a formula for the area of the surface of revolution formed by rotating $\mathcal{C}$ about the x -axis.

5.4.2 Surface area formulae and examples

Assume that a curve $\mathcal{C}$ lies in the upper-half plane and is simple (see the assumptions stated at the beginning of Subsection 5.4.1). We present formulae for the area of the surface of revolution about the x -axis when $\mathcal{C}$ is described either parametrically, as the graph of a function or using polar coordinates. In each case we assume that the appropriate derivatives exist.

If $\mathcal{C}$ is described parametrically by

$$\mathcal{C} = \{(x(t), y(t)) \in \mathbb{R}^2 : a \leq t \leq b\},$$

then the area A of the surface of revolution about the x -axis is given by

$$A = \int_a^b 2\pi y(t)\sqrt{[x'(t)]^2 + [y'(t)]^2} dt. \quad (5.6)$$

If $\mathcal{C}$ is the graph

$$y = f(x), \quad x \in [a, b],$$

of a function f on $[a, b]$ then the area A of the surface of revolution about the x -axis is given by

$$A = \int_a^b 2\pi f(x)\sqrt{1 + [f'(x)]^2} dx. \quad (5.7)$$

If $\mathcal{C}$ is described using polar coordinates by

$$r = f(\theta), \quad \theta_0 \leq \theta \leq \theta_1,$$

then the area A of the surface of revolution about the x -axis is given by

$$A = \int_{\theta_0}^{\theta_1} 2\pi r \sin \theta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta. \quad (5.8)$$

Remark 5.4.1. Formula (5.6) was derived heuristically in Subsection 5.4.1. Formula (5.7) may be easily deduced from (5.6) by using the following parameterisation of the graph of f :

$$x(t) = t, \quad y(t) = f(t), \quad a \leq t \leq b.$$

Formula (5.8) may be easily deduced from (5.6) by using the following parameterisation of the polar curve:

$$x(\theta) = f(\theta) \cos \theta, \quad y(\theta) = f(\theta) \sin \theta, \quad \theta_0 \leq \theta \leq \theta_1.$$

Remark 5.4.2. In parametric form, the formula for the area A of surface of revolution about the y -axis is given by

$$A = \int_a^b 2\pi x(t) \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Other versions of this formula may be easily deduced using the parametrisations given in the previous remark.


Remark 5.4.3. Each of these formulae only give the area of the *surface* of revolution. To find the surface area of the *solid* of revolution, one must also add the surface area contributed by any circular ‘caps’ appearing at each end of the surface of revolution. See, for example, Example 5.4.5.

Example 5.4.4. Find the surface area of a sphere of radius r .

Solution. A sphere of radius r is formed by rotating the curve

$$x(t) = r \cos t, \quad y(t) = r \sin t, \quad 0 \leq t \leq \pi$$

about the x -axis.



By using formula (5.6), we find that the surface area A is given by

$$\begin{aligned} A &= \int_0^\pi 2\pi r \sin t \sqrt{[-r \sin t]^2 + [r \cos t]^2} dt \\ &= \int_0^\pi 2\pi r \sin t \sqrt{r^2(\sin^2 t + \cos^2 t)} dt \\ &= \int_0^\pi 2\pi r^2 \sin t dt \\ &= 2\pi r^2 \left[-\cos t \right]_0^\pi \\ &= 4\pi r^2. \end{aligned}$$


So the surface area of a sphere of radius r is $4\pi r^2$. □

Example 5.4.5. A solid $\mathcal{S}$ is formed by rotating the curve given by

$$y = \sqrt{2-x}, \quad x \in [0, 2],$$

about the x -axis. Find the surface area of the solid, making sure that the every face of the solid is accounted for.

Solution. The solid $\mathcal{S}$ is drawn below.



It has two faces: the truncated paraboloid (shaded in lighter gray) and the circular cap (shaded in darker gray). The area A_1 of the truncated paraboloid is given by

$$\begin{aligned} A_1 &= \int_0^2 2\pi f(x) \sqrt{1 + [f'(x)]^2} dx && (\text{where } f(x) = \sqrt{2-x}) \\ &= \int_0^2 2\pi \sqrt{2-x} \left(1 + \left(\frac{-1}{2\sqrt{2-x}} \right)^2 \right)^{1/2} dx \\ &= \int_0^2 2\pi \sqrt{2-x} \left(1 + \frac{1}{4(2-x)} \right)^{1/2} dx \\ &= \pi \int_0^2 \sqrt{4(2-x) + 1} dx \\ &= \pi \int_0^2 \sqrt{9-4x} dx \\ &= \pi \left[\frac{(9-4x)^{3/2}}{-6} \right]_0^2 \\ &= \frac{13\pi}{3}. \end{aligned}$$

The area A_2 of the circular cap is given by

$$A_2 = \pi r^2 = \pi(\sqrt{2})^2 = 2\pi.$$

Hence the total surface area of the solid $\mathcal{S}$ is given by

$$A_1 + A_2 = \frac{19\pi}{3},$$


which is approximately 19.9 square units. □

We end with a simple but interesting example. Recall from high school that the volume V of the solid formed when the graph of a function $f : [a, b] \rightarrow [0, \infty)$ is rotated about the x -axis is given by

$$V = \int_a^b \pi [f(x)]^2 dx. \tag{5.9}$$


5.4. SURFACE AREA

Example 5.4.6 (Gabriel's horn). Suppose that the function $f : [1, \infty) \rightarrow \mathbb{R}$ is defined by $f(x) = \frac{1}{x}$. Rotate the graph of f about the x -axis, as shown.



The surface of revolution formed is known as *Gabriel's horn* (after the biblical figure Gabriel) or *Torricelli's trumpet* (after the mathematician and philosopher Evangelista Torricelli, who was a pupil of Galileo). Even though the length of the horn is infinite, it may still be possible to calculate its surface area and the volume of the corresponding solid.

We begin with the volume. First, consider the volume V_R of the truncated solid shown below.



Using formula (5.9),

$$\begin{aligned} V_R &= \int_1^R \frac{\pi}{x^2} dx \\ &= \pi \left[-\frac{1}{x} \right]_1^R \\ &= \pi \left(1 - \frac{1}{R} \right). \end{aligned}$$

Now $\lim_{R \rightarrow \infty} V_R = \pi$, and so the volume of the solid of revolution is finite and equal to π cubic units.

We now examine the surface area. The area A of the surface of revolution for the truncated curve is given by

$$A = 2\pi \int_1^R \frac{1}{x} \sqrt{1 + \frac{1}{x^4}} dx.$$

Finding an antiderivative for the integrand looks difficult. However, we are mainly interested in whether or not the improper integral

$$\int_1^\infty \frac{1}{x} \sqrt{1 + \frac{1}{x^4}} dx$$

converges. Note that

$$\frac{1}{x} \sqrt{1 + \frac{1}{x^4}} > \frac{1}{x} \sqrt{1 + 0} = \frac{1}{x}$$

whenever $x \geq 1$. Since the improper integral

$$\int_1^\infty \frac{1}{x} dx$$

diverges, so too does the integral

$$\int_1^\infty \frac{1}{x} \sqrt{1 + \frac{1}{x^4}} dx$$

by the comparison test for integrals (see Chapter 8 in the MATH1131 calculus notes). Hence the surface area of Gabriel's horn is infinite.

The fact that Gabriel's horn has finite volume (π cubic units) and infinite surface area leads to the following paradox. To paint the outside surface of the horn, one requires an infinite amount of paint since the surface area is infinite. However, to paint the inside surface of the horn, one only needs at most π cubic units of paint. Simply fill the horn with paint and then remove whatever paint is not touching the surface. This is sometimes called the *painter's paradox*.

Question: How is the paradox resolved?

Problems 5.1 : The average value of a function

1. [R] It can be shown that when a cable is hanging between two poles the curve that it forms is always the graph of a hyperbolic cosine function. Suppose that the height h (in metres) of a cable above the ground is given by

$$h(x) = 4 \cosh \left(\frac{x - 10}{20} \right),$$

where $0 \leq x \leq 20$. Find the average height of the cable above the ground.

2. [R] Suppose that the air temperature $T(t)$, measured in degrees Celsius t hours after noon, is given by

$$T(t) = 25 + 2t - \frac{t^2}{3}.$$

Find the average temperature between noon and 5 p.m.

Problems 5.2 : The arc length of a curve

3. [R] Calculate the lengths of the given arcs given by

- a) $y = x^{3/2}$, where $0 \leq x \leq 1$;
- b) $x = t - \sin t$, $y = 1 - \cos t$, where $0 \leq t \leq 2\pi$; and
- c) $x = t^3$, $y = t^2$ from $(0, 0)$ to $(8, 4)$.

4. [R] The astroid

$$x^{2/3} + y^{2/3} = a^{2/3} \quad (1)$$

has a parametrisation given by

$$x(\theta) = a \cos^3 \theta, \quad y(\theta) = a \sin^3 \theta.$$

Its graph was sketched, in the case when $a = 1$, in one of the problems from Chapter 7 of MATH1131.

- a) Use the parametric form to calculate the arc length of the astroid.

- b) [H]

- i) Show that the improper integral $\int_0^1 x^{-1/3} dx$ converges to $3/2$ by calculating the limit

$$\lim_{h \rightarrow 0^+} \int_h^1 x^{-1/3} dx.$$

- ii) Hence calculate the arc length of the astroid by using the implicit equation (1).

- 5. [R] Find the length of the curve $r = e^\theta$, where $0 \leq \theta \leq 2\pi$.
- 6. [R] Find the length of the cardioid $r = 1 + \cos \theta$.

CHAPTER 5. AVERAGES, ARC LENGTH, SPEED AND SURFACE AREA
Problems 5.3 : The speed of a moving particle

7. [R] A projectile is fired from an elevated cannon. Its horizontal distance x (in metres) from the cannon and height y (in metres) above the ground, exactly t seconds after it is fired, is given by

$$x(t) = 40t, \quad y(t) = -5t^2 + 40t + 45, \quad 0 \leq t \leq t_1,$$

where t_1 is the time the projectile hits the ground.

- Find t_1 .
 - Find the speed of the projectile immediately prior to impact.
 - What was the average height of the projectile above the ground during the period after it was fired and before impact?
 - [X] What distance did the projectile travel during this period?
8. [H] The position $(x(t), y(t))$ of a particle P at time t is given by


$$x(t) = \cos\left(\frac{\pi}{2}(\cos \pi t - 1)\right), \quad y(t) = -\sin\left(\frac{\pi}{2}(\cos \pi t - 1)\right),$$

where $t \geq 0$.

- Find a formula for the speed $v(t)$ of the particle at time t .
- For what values of t is the speed of the particle (i) a maximum and (ii) a minimum?
- What curve does the trajectory of the particle trace out?
- What is the length of the curve of (c)?
- Find the distance that the particle travels during the time interval $[0, 3]$.

Problems 5.4 : Surface area

9. [R] In this question we will show that the surface area of the frustum of a right circular cone is given by $\pi(R + r)s$. (For a diagram, see the beginning of Section 5.4.) Consider a truncated right circular cone with slant height s and base radius R . Cut a line from the vertex Q to the base and flatten the cone as shown below.



- Explain why the flattened surface is the sector of a circle.

- b) Find, in terms of R and s , the area of the sector and hence the surface area of the cone.
- c) Hence show that the surface area of a frustum of radii r and R and of slant height s is given by $\pi(R + r)s$.
10. [R] Find the area of the surface of revolution formed when the given curve is rotated about the x -axis.
- a) $y = x^3$, where $0 \leq x \leq 2$. b) $x = t - \sin t$, $y = 1 - \cos t$, where $0 \leq t \leq 2\pi$.
11. [R] Show that if $-a \leq b < c \leq a$ then the surface area of the sphere $x^2 + y^2 + z^2 = a^2$ between the planes $x = b$ and $x = c$ is $2\pi a(c - b)$.
12. [R] Suppose that $0 < r < R$. A surface (doughnut) is formed by rotating a circle of radius r and centre $(0, R)$ about the x -axis. By finding a suitable parametrisation for the circle, show that the area of the surface is $(2\pi R)(2\pi r)$.
13. [R] Find the area of the surface formed when the polar curve $r = 1 + \cos \theta$, where $0 \leq \theta \leq \pi$, is rotated about the x -axis.

Answers to selected problems

Chapter 1

1. a) cup shaped b) unit sphere c) umbrella stand
d) cone with semi-vertical angle $\pi/4$ e) saddle

2. $2xye^{x^2y}$, $x^2e^{x^2y}$, $(2x + 2x^3y)e^{x^2y}$.

	$\frac{\partial z}{\partial x}$	$\frac{\partial z}{\partial y}$	$\frac{\partial^2 z}{\partial x^2}$	$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$	$\frac{\partial^2 z}{\partial y^2}$
3. a)	$2xy$	$x^2 + 2y$	$2y$	$2x$	2
b)	$\frac{-y}{(x^2 + y^2)}$	$\frac{x}{(x^2 + y^2)}$	$\frac{2xy}{(x^2 + y^2)^2}$	$\frac{(y^2 - x^2)}{(x^2 + y^2)^2}$	$\frac{-2xy}{(x^2 + y^2)^2}$
c)	$\cos(x - cy)$	$-c \cos(x - cy)$	$-\sin(x - cy)$	$c \sin(x - cy)$	$-c^2 \sin(x - cy)$

4. a) $z = 6x + 10y - 34$, $\mathbf{n} = (6, 10, -1)^T$
b) $z = 32 - 16x + 16y$, $\mathbf{n} = (-16, 16, -1)^T$
c) $4x - 6y - 7z - 14 + 7 \ln 7 = 0$, $\mathbf{n} = (4, -6, -7)^T$
d) $2x + 3y + \sqrt{23}z - 6 = 0$, $\mathbf{n} = (2, 3, \sqrt{23})^T$

5. a) $784\pi \text{ cm}^3$ b) $\frac{217\pi}{15} \text{ cm}^3$ c) 1.85%

6. 0.05

7. 5.012 (calculator gives 5.012115)

8. $|\Delta S| \leq 0.0404$

9. 9% decrease.

10. 0.21%

11. a) $e^t(t^2 + 2t)$ b) $2t$

12. 7.5π cubic centimetres per second

13. b) $F(x, y) = \sin(y - x^2)$

14. a) $u_{xx}(x, t) = g''(x + \lambda t)$, $u_{tt}(x, t) = \lambda^2 g''(x + \lambda t)$
b) 4, -4

17. a) $u_t(x, t) = \frac{-u(x, t) f'(x - tu(x, t))}{1 + t f'(x - tu(x, t))}$, $u_x(x, t) = \frac{f'(x - tu(x, t))}{1 + t f'(x - tu(x, t))}$

- c) $t_m = 1$
- d) Hint: Try the Maple commands
`with (plots):`
`implicitplot(y-1+tanh(x-t*y),x=-5..7, y=-1..3, gridrefine = 2);`
 for a few values of t . Can you animate this in time t ?
- e) No
18. a) $y = \pm 1, \frac{dy}{dt} = 0$.
- b) $\nabla F = (2x, 2y, -2z)^T$.
- c) The vector $\left(\frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt}\right)^T$ represents the velocity of a particle on the hyperboloid. So the equation states that the velocity is perpendicular to the surface's normal.

Chapter 2

1. a) $\frac{1}{4}e^{2x^2}$ b) $-\frac{1}{2}\cos(x^2)$ c) $\frac{1}{6}\sin(2x^3)$
 d) $\frac{1}{10}\ln|5x^2 - 11|$ e) $-\frac{1}{4}\cos^4 x$ f) $\ln(\ln x)$
 g) $\sqrt{x^2 + 4x + 7}$ h) $\frac{1}{3}(1 + x^2)^{\frac{3}{2}}$ i) $-\frac{1}{18}(9 - 4x^3)^{\frac{3}{2}}$
 j) $-\frac{1}{6}(9 - 4x^3)^{\frac{1}{2}}$ k) $-\frac{1}{8(1 + x^4)^2}$ l) $\frac{-1}{3\tan^3 x}$
 m) $\frac{-1}{2\sin^2 x}$ n) $\frac{1}{8}(4 + 3e^{2x})^{\frac{4}{3}}$ o) $\frac{-1}{4(\ln x)^4}$
2. a) $-e^{-x}(x^2 + 2x + 2)$ b) $\frac{1}{4}x^4 \ln x - \frac{1}{16}x^4$ c) $x \tan x + \ln(\cos x)$
 d) $-\frac{(\ln x)^2}{x} - 2\frac{\ln x}{x} - \frac{2}{x}$ e) $\frac{1}{2}e^x(\sin x + \cos x)$ f) $x \ln x - x$
 g) $x \tan^{-1} x - \frac{1}{2}\ln(x^2 + 1)$
3. a) $1/8$ b) $4/15$
 c) $\frac{1}{3}\sec^3 x + C$ d) $\frac{1}{4}\sin 2\theta + \frac{1}{2}\theta + C$
 e) $-\frac{1}{99}(\sin x \cos 10x - 10 \cos x \sin 10x) + C$ or $\frac{1}{2}\left(\frac{1}{9}\sin 9x + \frac{1}{11}\sin 11x\right) + C$ f) $-\frac{1}{10}\cos 5x + \frac{1}{2}\cos x + C$
4. a) $\ln|\tan x + \sec x| + C$
 b) i) $\frac{1}{3}\sec^2 x \tan x + \frac{2}{3}\tan x + C$
 ii) $\frac{1}{4}\sec^3 x \tan x + \frac{3}{8}\sec x \tan x + \frac{3}{8}\ln|\tan x + \sec x| + C$
5. a) $3\pi/512$ b) $1/60$ c) $2/35$
6. $6 - 16e^{-1}$
7. $5/12 - (\ln 2)/2, \quad \pi/4 - 76/105$
8. $(e^2 + 3)/8$
9. $35\pi/256, \quad 16/35$
10. $I_n = (2\sqrt{2} - 2nI_{n-1})/(1 + 2n)$

15. a) $\pi/3 - \sqrt{3}/2$; $x = 2 \sin \theta$
 b) $\sinh^{-1} \frac{x-3}{2} + C$ or $\ln(x-3 + \sqrt{x^2 - 6x + 13}) + K$; complete the square.
 c) $9\pi/4$
 d) $-\frac{\sqrt{x^2+16}}{16x} + C$; $x = 4 \tan \theta$ to obtain $\int \frac{\cos \theta}{\sin^2 \theta} d\theta$.
 e) $x/\sqrt{1-x^2} + C$
 f) $\tan^{-1} 2$
16. $\sqrt{x^2-4} + C$
17. a) $\frac{1}{2} \ln \left| \frac{x+1}{x+3} \right| + C$ b) $3 \log |x-2| + 2 \log |x-1| + C$
 c) $2 \log |x| + \frac{1}{x} + 2 \log |x-1| + C$ d) $-\frac{1}{4} \left(\frac{2x}{x^2-1} - \log \left| \frac{x+1}{x-1} \right| \right) + C$
 e) $x + \ln \left| \frac{x-1}{x+1} \right| + C$ f) $\ln |x-3| - \frac{1}{2} \ln(x^2+9) - \tan^{-1} \frac{x}{3} + C$
 g) $\ln \frac{(x+1)^2}{|x+2|} + \frac{4}{x+2} + C$ h) $\frac{-1}{(1+x)^2} + \frac{1}{1+x} = \frac{x}{(1+x)^2} + C$
18. a) $\frac{1}{2} \ln(x^2+2x+10) - \frac{1}{3} \tan^{-1} \frac{x+1}{3} + C$ b) $\sqrt{x^2+2x+10} - \sinh^{-1} \frac{x+1}{3} + C$
 c) $2\sqrt{x} - 2 \ln(1 + \sqrt{x}) + C$ d) $11 - 6 \ln(3/2)$; $x = u^6$ first.
22. a) $\ln x - \frac{1}{2} \ln(x^2+x+1) - \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2x+1}{\sqrt{3}} \right) + C$
 b) $\frac{8}{5} \cosh^5(x) + C$
 c) $\frac{3}{2} \ln(x^2+4x+8) - \frac{1}{2} \tan^{-1} \left(\frac{x+2}{2} \right) + C$
 d) $\frac{1}{2} x \sqrt{25-x^2} + \frac{25}{2} \sin^{-1} \left(\frac{x}{5} \right) + C$
 e) $\ln |x-1| + \ln(x^2-2x+2) + \tan^{-1}(x-1) + C$
 f) $-\frac{\sqrt{1+x^2}}{x} + C$
 g) $\frac{1}{3} \frac{x}{\sqrt{x^2+3}} + C$
 h) $-\frac{1}{14} \cos(7x) + \frac{1}{2} \cos(x) + C$
 i) $2/e$

Chapter 3

1. a) $y = \tan(t^3/3 + C)$ b) $y = -2/(x^2 + A)$
 c) $y = \sqrt[3]{A - 3 \cos x}$ d) $y = C(x-1)/x$
 e) $e^{x+y} + Ce^y + 1 = 0$ f) $y^2 = \tan^2 x + 4 \tan x - 1$, where $y > 0$
 g) $(\ln x)^2 = 2 \ln y + C$ h) $y = 1/(1-x^3)$
2. Let a and b be arbitrary real numbers with $a < b$. Then there are solutions of the form

$$y = \begin{cases} (x-a)^3 & \text{if } x < a \\ 0 & \text{if } a \leq x \leq b \\ (x-b)^3 & \text{if } x > b \end{cases}.$$

3. $2\sqrt{2}$

4. a) $y = (x^3/3 + C)e^{2x}$ b) $y = e^{-3x}(\tan^{-1} x + C)$
 c) $y = (2 + Ce^{-x})/x$ d) $y = 2x^5 + Ax^2$
 e) $y = \tan x - 1 + Ae^{-\tan x}$ f) $y = \frac{x}{2} \tan 2x + \frac{1}{4} + A \sec 2x$
5. $x = t - 1 + Ce^{-t}$
7. a) $3x^2y + y^3 = A$ b) $x \sin y - \frac{1}{2}x^2y^2 = C$
8. a) $(x^2 + 1)e^y + e^x = C$ b) Not exact
 c) $y \tan^{-1} x + xe^y = C$ d) $e^{xy} \cos x = C$
9. a) $xy^2 - 3y \tan x = C$ b) $y^2 = \ln(x^2 + 1) + C$
 c) $y = \frac{\ln x + C}{x}$ d) $y = Axe^{-1/x}$
 e) $y = \left[3 \left(\ln|x| + \frac{x^2}{2} + \frac{17}{2} \right) \right]^{1/3}$
10. $xy^2 = 2y^5 + C$
11. a) $x/y = \ln|x| + C$ b) $\sec \frac{y-x}{x} + \tan \frac{y-x}{x} = Ax$
 c) $3x^2y + y^3 = A$ d) $3xy^2 + x^3 = A$
 e) $\sinh^{-1}(y/x) = \ln x$; that is, $y = \frac{1}{2}(x^2 - 1)$
12. $\ln \sqrt{x^2 + y^2} = -\tan^{-1}(y/x)$. Logarithmic or equiangular spiral, $r = e^{-\theta}$.
13. $2x + C = \ln \left| \frac{y-x-1}{y-x+1} \right|$ or $y = x - 1 + \frac{2}{1 + De^{2x}}$
14. a) $y = \frac{1}{2 + Ae^{-t}}$
 b) $y = 25/(5t - 1 + 26e^{-5t})$; $y_{\max} = \frac{25}{\ln 26}$ when $t = \frac{\ln 26}{5}$
15. a) $y^2 = Ce^{4x} - \frac{1}{2}x - \frac{1}{8}$ b) $x \ln y + x^3/3 = C$ c) $y = Ae^{Bx^2}$
16. a) $y = Ae^{-2t} + B + 3t$ b) $y = \frac{t}{c} - \frac{\ln(1+ct)}{c^2} + d$
17. a) Let $z = \frac{dv}{dx}$. We obtain $\alpha z' + (2\alpha' + b\alpha)z = 0$, which is first order homogeneous linear.
 b) $u = Ax^2 + Bx^3$
18. a) i) $y = 0$, $y = K$
 ii) $y = \frac{y_0 K}{y_0 + (K - y_0)e^{-kt}}$
 iii) y is strictly increasing and approaches K .
 iv) $y = K/2$; work from the differential equation.
 v) The solution is the same as for ii); y is strictly decreasing (and concave upwards) with $y = K$ as a horizontal asymptote.
 b) $y = K \exp \{-\ln(K/y_0)e^{-\alpha t}\}$; one way, let $z = \ln y$.
19. Let y litres of pollutant be present in the lake after t days.

$$\frac{dy}{dt} = 10^4 - \frac{y(t)}{10^9} \times (10^6 + 10^4), \quad \text{giving} \quad y = \frac{10^9}{101} \left(1 - e^{-1.01t/10^3} \right)$$

- a) $y \rightarrow 10^9/101$ litres or just under 1%
- b) $\frac{10^5 \ln 2}{101} \approx 686.3$ days ≈ 1.88 years
- c) That there is perfect mixing, that the pollutant does not precipitate or dissolve, that the pollutant does not itself create more pollution, that
20. With an inflow of 3 litres per minute and an outflow of 1 litre per minute, the volume of liquid in the tank at time t is $50 + 2t$ litres. The inflow of salt is 3×2 grams per minute. In running off 1 litre per minute with a concentration of $x/(50 + 2t)$ grams of salt per litre, the rate of removal of salt is $1 \times x/(50 + 2t)$ grams per minute. So the net rate of increase of x is given by $\frac{dx}{dt} = 6 - x/(50 + 2t)$ or $\frac{dx}{dt} + \frac{1}{50 + 2t}x = 6$. This is a first order linear ODE. (You may find it helpful to consider the outflow over a small time interval $[t, t + \Delta t]$.)
21. a) $P(t) = 100 - 25 \cos t + 25 \sin t - 55e^{-t}$
b) 100
22. $y = A \exp \left\{ k \left(t + \frac{a}{2\pi} \sin(2\pi t) \right) \right\}$
23. a) $r(t) = 2\% + (1.5\% - 2\%)/10t = 0.02 - 0.0005t$, where t is measured in years.
b) The differential equation is $\frac{dy}{dt} = (0.02 - 0.0005t)y$
 $y = 10^7 \exp(0.02t - 0.00025t^2)$. When $t = 10$, $y = 10^7 e^{0.175} \approx 10^7 \times 1.19$
24. a) $\frac{dv}{dt} = g - kv$; with $g = 9.8$, $k = 10$. So $v = Ae^{-kt} + g/k$. With $v = 0$ at $t = 0$,
 $v = \frac{g}{k}(1 - e^{-kt}) = \frac{49}{50}(1 - e^{-10t})$
b) $g/k = 49/50$
c) $t \geq \frac{\ln 20}{k}$ (≈ 0.3 seconds)
d) $A > 0$ and v decreases (rapidly) towards g/k .
25. a) $y = Ax^\lambda$
b) Graph $\ln y$ against $\ln x$. From the graph, $\lambda \approx 1.54$ and $\ln A \approx -4.55$ (so that $A \approx 0.0106$).
26. a) \$610701.38
b) Treating the car payments as continuous at the rate of \$10400 a year, $\frac{dP}{dt} = 0.2P - 10400$. Then $P = 52000 + 448000e^{0.2t}$, for $0 \leq t \leq 1/2$. At $t = 1/2$, $P = 547116.57$. The capital remaining in the cooperative after 6 months is $47116.57 e^{-0.05} = 44818.67$. At the end of one year, the total is 547116.57 (capital plus new interest in Hitek) + 44818.67 (in the co-op) = 591935.24 dollars.
28. $\frac{dy_1}{dt} = -0.25y_1$, $\frac{dy_2}{dt} = 0.25y_1 - 2y_2$ with $y_1(0) = K$, $y_2(0) = 0$. Thus $y_1 = Ke^{-0.25t}$ and hence $y_2 = \frac{K}{7}(e^{-0.25t} - e^{-2t})$. The maximum value of y_2 occurs for $t = \frac{12}{7} \ln 2$. That is, after about 1.188 days.
29. $20 \ln 2 \approx 13.8$ m/sec
30. a) $y = Ae^{-2x} + Be^{-x}$
b) $y = e^{-x}(C \cos 3x + D \sin 3x)$
c) $y = Ae^{-3x}$
d) $y = (Ax + B)e^{-2x}$

31. a) $\frac{1}{4}(5e^x - e^{5x})$ b) $y = e^{-x}(\cos x + \sin x)$
32. a) $y = Ae^{-3x} + Be^{-x} + \frac{1}{3}x - \frac{4}{9}$ b) $y = (Ax + B)e^{3x} + 5e^{2x}$
 c) $y = e^{-x}(A \sin x + B \cos x) + 2 \sin 2x - \cos 2x$ d) $y = (A - x/2)e^{-x} + Be^x$
 e) $y = A \cos 2x + B \sin 2x - \frac{1}{4}x \cos 2x$ f) $y = Ae^{-3x} + 2e^{2x}$
 g) $y = \frac{1}{3}(\sin x + \sin 2x)$ h) $y = e^x + e^{4x} - e^{2x}$
33. a) $Ae^{-x} + Be^{5x/2}$; seek $y_P = x(ax + b)e^{5x/2}$
 b) $Ae^{4x} + Be^{-6x}$; seek $y_P = e^{4x}(a \cos 6x + b \sin 6x)$
 c) $(Ax + B)e^{-3x}$; seek $y_P = x^2 ae^{-3x}$
34. $y = \frac{1}{2}x^4 e^{2x}$.
35. You should obtain $\frac{d^2y}{dt^2} - 5 \frac{dy}{dt} + 6y = e^{5t}$. $y = Ax^3 + Bx^2 + \frac{1}{6}x^5$
37. $50^2 g/\pi$ grams; about 780 kilograms
38. a) $x(t) = \cos 2t$, so the block oscillates with fixed amplitude.
 b) If $c = 2$ then $x(t) = e^{-t}(\cos \sqrt{3}t + \frac{1}{\sqrt{3}} \sin \sqrt{3}t)$, so the system has damped oscillations.
 If $c = 5$ then $x(t) = \frac{1}{3}(e^{-t} - e^{-4t})$, so the system does not oscillate.
 c) If the characteristic equation has real roots then the solution has no oscillating terms. This happens whenever $c \geq 4$. So the smallest value of c is 4.
39. a) $q(t) = A \cos 100t + B \sin 100t$
 b) $x_P = a \cos \Omega t + b \sin \Omega t$ if $\Omega \neq 100$; $x_P = t(a \cos \Omega t + b \sin \Omega t)$ if $\Omega = 100$.
 c) $50/\pi$ (which corresponds to when $\Omega = 100$).
40. a) $y = Ae^{-t} + Be^{-2t} + 2 \sin t - 6 \cos t$
 b) $y = -6 \cos t + 2 \sin t$
41. b) $y_p = -\frac{1}{4}t \cos t$
42. b) No
 c) $\mu = n\pi/L$, where $n = 1, 2, 3, \dots$. The corresponding solutions are $y_n(x) = B_n \sin(n\pi x/L)$.
43. b) $2\lambda = 1 + \frac{k^2\pi^2}{L^2}$, $k = 1, 2, 3, \dots$ with $y_k = B_k e^{-x} \sin(k\pi x/L)$
44. a) $\frac{x^2}{a} + \frac{y^2}{b} = C$, a family of ellipses.
 b) $\frac{d^2y}{dx^2} + aby = 0$, $y = A \cos \omega t + B \sin \omega t$, where $\omega^2 = ab$
 and $x = (A\omega/b) \sin \omega t - (B\omega/b) \cos \omega t$.
 c) $\omega = 1.6$, $A = 3.2$, $B = 2.4$
 $x = 1.6 \sin \omega t - 1.2 \cos \omega t$, $y = 3.2 \cos \omega t + 2.4 \sin \omega t$ or
 $x = -2 \cos(\omega t + \phi) = 2 \sin(\omega t + \phi - \pi/2)$ and $y = 4 \sin(\omega t + \phi)$
 where $\phi = \sin^{-1} \frac{4}{5}$, and $\frac{x^2}{4} + \frac{y^2}{16} = 1$. The peaks in the predator population lag behind the peaks in the prey population by a quarter of the period.
45. Care is required if the characteristic equation of $Lu = 0$ has a double root. Otherwise a basis for $\ker(L^2)$ is $\{u_1, xu_1, u_2, xu_2\}$.
 The two dimensional kernel of L is a subspace of the four dimensional kernel of L^2 .

1. a) $1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^9}{9!}$
 b) $x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!}$
 c) $x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \frac{x^9}{9!}$
2. $p_{2m+1}(x) = \sum_{k=0}^m \frac{(-1)^k x^{2k+1}}{(2k+1)!}$
3. a) $2 + \frac{1}{4}(x-4) - \frac{1}{64}(x-4)^2 + \frac{1}{512}(x-4)^3$
 b) $\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}\left(x - \frac{\pi}{4}\right) - \frac{1}{2\sqrt{2}}\left(x - \frac{\pi}{4}\right)^2 + \frac{1}{6\sqrt{2}}\left(x - \frac{\pi}{4}\right)^3 + \frac{1}{24\sqrt{2}}\left(x - \frac{\pi}{4}\right)^4$
4. a) $3 + 3(x-1)$
 b) $3 + 3(x-1) + (x-1)^2$
 c) $7 + 5(x-2) + (x-2)^2$
5. a) $p_1(x) = x$ and $R_2(x) = -\frac{x^2}{2(1+c)^2}$ for some c between 0 and x .
7. a) $p_n(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots + (-1)^k \frac{x^n}{n!}$ where $n = 2k$;
 $R_{n+1}(x) = \frac{(-1)^{k+1} \sin c}{(n+1)!} x^{n+1}$ for some c between 0 and x .
 c) $|R_{n+1}(x)| < \frac{1}{10^{n+2}(n+1)!}$
 d) $n = 4$
 e) The estimate $\sin x \leq 1$ (whenever $x \in \mathbb{R}$) is less crude in the case when $x \geq 1$.
 f) $n = 14$
 g) Rearranging the inequality $|\text{error}| \leq \frac{a^{11}}{11!} < 10^{-6}$ gives $a < \sqrt[11]{11!10^{-6}}$. So a less than 1.398 will do.
10. a) Horizontal point of inflexion at 1; local maximum at -2 .
 b) Horizontal point of inflexion at -1 ; local minima at 2 and 3.
12. a) $1/2$ b) 0 c) boundedly divergent d) 1
 e) 0 f) diverges to ∞ g) a
13. a) $1/2$ b) $N = 2/\epsilon$ works
14. a) e^4 b) $1/e$
15. e) $(1 + \sqrt{5})/2$
16. a) $1/2, 0$ b) $1/2, -1/2$ c) $\sqrt{2}, -\sqrt{2}$
 d) $1/2 + \sin 2, -1/5 + \sin 5$ e) ∞, π f) $1 + \pi/2, 1 - \pi/2$
17. b) 2

19. a) iii) 1 b) iii) $3/4$
24. diverges
25. a) divergent b) divergent c) convergent
26. a) convergent b) divergent c) divergent
d) divergent e) divergent f) convergent
27. a) convergent b) convergent c) convergent d) divergent
28. a) conditionally convergent
b) divergent (by the k th term test)
c) absolutely convergent
d) diverges to ∞ ; $\frac{(-1)^k}{\sqrt{k} + (-1)^k} = \frac{(-1)^k \sqrt{k}}{k - 1} - \frac{1}{k - 1}$
29. b) $s_4 = 9677/16380 \approx 0.59078$; $|\text{error}| \leq 1/126$
c) n equal to 99 will do.
30. a) convergent b) divergent c) convergent
d) convergent e) divergent f) convergent
31. Only e) and f) are divergent.
32. Converges if $\lim_{k \rightarrow \infty} a_k > 1$ and diverges otherwise.
34. a) $\sum_{k=0}^{\infty} \frac{x^k}{k!}$ b) $\sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!}$ c) $\sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!}$
d) $\sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^k}{k}$ e) $-\sum_{k=0}^{\infty} x^k$
35. a) $e^x = 1 + x + \frac{x^2}{2!} \cdots + \frac{x^n}{n!} + \frac{e^{c_n} x^{n+1}}{(n+1)!}$, for some c_n between 0 and x .
36. a) $R_{n+1}(x) = \frac{\sinh c_n}{(n+1)!} x^{n+1}$ if n is odd and $R_{n+1}(x) = \frac{\cosh c_n}{(n+1)!} x^{n+1}$ if n is even. In each case, c_n lies between 0 and x .
37. a) $R_{n+1}(x) = \frac{(-1)^n x^{n+1}}{(n+1)(1+c_n)^{n+1}}$ for some c_n between 0 and x .
39. a) $f'(x) = 2e^{-1/x^2}/x^3$ if $x \neq 0$; $f'(0) = 0$.
b) $f''(x) = 4e^{-1/x^2}/x^6 - 6e^{-1/x^2}/x^4$ if $x \neq 0$; $f''(0) = 0$.
e) The Maclaurin series is $0 + 0x + 0x^2 + 0x^3 + \cdots$, and hence converges everywhere. It only converges to f at 0.
40. b) $\ln 2$
f) $\frac{1}{2} \ln 2$

41. Students studying MATH1231 only need to state the corresponding open interval in each case.

- a) $(-6, 6)$ b) $[-1, 1]$ c) $(-2, 2)$
 d) $[1, 3]$ e) $[1/3, 1]$ f) $(-3, 3]$
 g) $(-\infty, \infty)$

43. a) $\sum_{k=0}^{\infty} \frac{(k+1)x^k}{k!}$ b) $\sum_{k=0}^{\infty} \frac{(-1)^k x^{4k+2}}{(2k+1)!}$

c) Differentiate 34(e): $\sum_{k=1}^{\infty} kx^{k-1}$

44. All series are valid on $(-1, 1)$.

a) $\sum_{k=0}^{\infty} (-1)^k x^k$ b) $\sum_{k=0}^{\infty} (-1)^k x^{2k}$ c) $\sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1}$

45. a) $f(x) = x^5 - \frac{x^{11}}{3!} + \frac{x^{17}}{5!} - \frac{x^{23}}{7!} + \cdots = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} x^{6k-1}}{(2k-1)!}$

c) A horizontal point of inflexion.

46. $R = 1$, $\ln 2 \approx \frac{56}{81}$.

47. a) A local maximum

b) $\text{Si}(x) = x - \frac{x^3}{3!3} + \frac{x^5}{5!5} - \frac{x^7}{7!7} + \cdots$

c) $\text{Si}(\pi) \approx \pi - \frac{\pi^3}{3!3} + \frac{\pi^5}{5!5} - \frac{\pi^7}{7!7} \approx 1.84$

48. a) $(-1, 1)$

b) $f(x) = \frac{x}{(1-x)^2}$

49. a) $\sum_{k=1}^{\infty} \frac{x^k}{k!}$

b) $y' = \sum_{k=1}^{\infty} k a_k x^{k-1}$, $y'' = \sum_{k=2}^{\infty} k(k-1) a_k x^{k-2}$

c) $a_k = \frac{1}{k^2 k!}$ whenever $k \geq 1$

d) $y = \sum_{k=1}^{\infty} \frac{x^k}{k^2 k!}$ whenever $x \in \mathbb{R}$.

Chapter 5

1. $8 \sinh(1/2) \approx 4.17$ metres

2. $27\frac{2}{9}^\circ\text{C}$

3. a) $(13^{3/2} - 8)/27$ b) 8 c) $\frac{8}{27}(10\sqrt{10} - 1)$

4. $6a$

5. $\sqrt{2}(e^{2\pi} - 1)$

6. 8

7. a) 9 seconds

- b) $10\sqrt{41} \approx 64.03$ metres per second
 - c) 90 metres
 - d) $80\sqrt{2} - 80 \ln(\sqrt{2} - 1) + 25\sqrt{41} + 160 \ln 2 - 80 \ln(\sqrt{41} - 5) \approx 427.53$ metres
8. a) $v(t) = \frac{\pi^2}{2} |\sin(\pi t)|$
- b) (i) $t = \frac{1}{2} + k$, where k is a positive integer. (ii) $t = n$, where n is a positive integer.
- c) A semicircle of centre $(0, 0)$ and radius 1 in the upper half-plane.
- d) π
- e) 3π
10. a) $\frac{\pi}{27}(145^{3/2} - 1)$ b) $64\pi/3$
13. $32\pi/5$

Index

- absolutely convergent, 136
- alternating series, 133
- alternating series test, 133
- answers, 191
- arc length, 174
 - of a catenary, 176
 - of a cycloid, 174
 - of a polar curve, 177
 - of the graph of a function, 176
- Archimedes, 101
- area of the surface of revolution, 182
- average height of a suspended cable, 170
- average value of f , 169
- bounded monotonic sequence, 120
- chain rule(s), 16, 17, 19, 20
 - chain diagram(s), 17, 20
- characteristic equation, 78
- classifying stationary points, 112
 - second derivative test, 112
- closed form, 147
- comparison test, 128
- conditionally convergent, 137
- contour, 1
 - contour lines, 1
- convergence and divergence of p -series, 129
- convergent sequence, 115
- decreasing sequence, 120
- differential approximation to Δf , 13
- differential equations, 55
 - applications of, 71
 - exact equations, 66
 - explicit solution, 57
 - general solution, 56
 - implicit solution, 57
 - initial value problem, 58
 - integrating factor, 63
 - order, 56
 - ordinary
 - particular solution, 56, 83
 - second order, 76
 - separable equations, 60
 - solution, 56
 - differential form, 66
 - divergent sequence, 115
 - diverges to negative infinity, 115
- e , 157
 - is irrational, 157
- exact differential equations, 66
- first order linear differential equations, 62
- frustum, 179
- Gabriel's horn, 185
- general solution, 56
- greatest lower bound, 121
- Gregory, James, 101
- growth of sequences, 120
- harmonic series, 124
- increasing, 120
- increment, 14
- Indian mathematicians, 101
- infima, 122
- infimum, 122
- infinite series, 122
 - absolutely convergent, 136
 - alternating series, 133
 - comparison test, 128
 - conditionally convergent, 137
 - converges to L , 123
 - diverges, 123
 - integral test, 127
 - k th term test for divergence, 126
 - n th partial sum, 123
 - p -series, 129
 - ratio test, 131
 - summable, 123
 - tail of the series, 125

202

- initial conditions, 58
- initial value problem, 58
- integral test, 127
- integrating factor, 63
- integration
 - powers of sin and cos, 25
 - by hyperbolic substitutions, 35
 - by trigonometric substitutions, 35
 - multiple angles of sin and cos, 28
 - of rational functions, 37
 - of trigonometric functions, 25
 - powers of tan and sec, 29
 - reduction formula, 30
- irreducible, 38
- Lambert, Johann Heinrich, 32
- least upper bound, 121, 122
- least upper bound axiom, 121
- Leibniz, 133
- length of a curve, 172
- level curve, 1
- limit inferior, 158
- limit of sequence, 116
- limit superior, 158
- linear combination, 77
- logistic curve, 76
- lower bound, 121
- Maclaurin series for f , 138
- Malthus, Thomas, 73
- mathematical model, 71
- mean value theorem for integrals, 156, 171
- Millennium problems, 76
- monotonic, 120
- Navier–Stokes equations, 76
- Newton’s law of cooling, 61
- non-homogeneous differential equations, 80, 83
- nondecreasing, 120
- nonincreasing, 120
- normal vector, 10, 11
- open interval of convergence, 144
- order of growth of sequences, 120
- π , 32, 101
 - is irrational, 34
- painter’s paradox, 186
- partial derivative(s), 7, 19
 - mixed derivative theorem, 9
 - of F with respect to x , 5
 - of F with respect to y , 5
 - second order, 8
- particular solution, 56, 83
- pinching theorem for sequences, 119
- population growth, 73
- power series, 148
 - closed form of, 147
 - differentiable, 148
 - in powers of x , 142
 - in powers of $x - a$, 142
 - integrable, 148
 - open interval of convergence, 144
 - radius of convergence, 144
- radius of convergence, 144
- ratio test, 131
- rational function(s), 37
 - distinct linear factors, 40
 - improper, 37
 - irreducible, 37
 - irreducible quadratic factor, 41
 - partial fractions decomposition, 38, 40
 - proper, 37
 - repeated irreducible quadratic factor, 42
 - repeated linear factor, 40
- rearrangement of a series, 137
- reduction formula, 30
- remainder term, 109
 - Lagrange formula, 110, 156
- resonance, 84
- Riemann sum, 169
- right circular cone, 179
- second derivative test, 112
- second order linear differential equations
 - characteristic equation, 78
 - homogeneous equations, 77
 - linearly independent solutions, 77
 - non-homogeneous equations, 80
 - particular solution, 56, 83
 - resonance, 84
 - with constant coefficients, 76
- separable differential equations, 60
- sequence(s), 114
 - bounded monotonic, 120
 - boundedly divergent, 115
 - convergent, 115

INDEX

- decreasing, 120
- definition of limit, 116
- divergent, 115
- diverges to $-\infty$, 115
- diverges to ∞ , 115
- greatest lower bound, 121
- infimum, 122
- least upper bound, 121
- lower bound, 121
- n th term of, 114
- order of growth, 120
- pinching theorem, 119
- supremum, 122
- unboundedly divergent, 115
- upper bound, 121
- series, *see* infinite series
- simple curve, 179
- sine integral $\text{Si}(x)$, 164
- speed $v(t)$ of, 178
- summable, 123
- suprema, 122
- supremum, 122
- surface of revolution, 180
- suspended cable, 167
- tail of the series, 125
- tangent plane(s), 10, 11, 20
- Taylor polynomial
 - about a , 105
 - for f about a , 106
 - of degree n , 104
- Taylor series
 - converges on I , 139
 - converges to f on I , 139
 - diverges on I , 139
 - for f about a , 138
 - function represented by, 140
- Taylor's theorem, 109
 - remainder term, 109
 - Lagrange formula, 110, 156
- Taylor, Brook, 101
- Torricelli's trumpet, 185
- Torricelli, Evangelista, 185
- total differential, 13
 - approximation ΔF , 20
 - approximation to ΔF , 14
 - use with measurement errors, 14
- upper bound, 121
- Verhulst, Pierre, 75
- Wallis' product, 50