Fermion Mass Ratios from the K/P Projection: Seven Zero-Parameter Predictions Including the Higgs Mass

Companion to "The Physics of Order"

Thomas DiFiore

March 22, 2026

Abstract

Using the algebraic structure derived in Paper I (the gauge group, charge assignments, and generation count), we compute seven observables from the K/P projection mechanism with zero free parameters. The source functional ϕ induces a split into K (position) and P (momentum) components; the observable obs $= Q^2 + P^2$ is the unique SO(2)-invariant quadratic form (Born rule uniqueness). Mass ratios are determined by the perpendicular component of summed SU(N) holonomies over causal chains in a Poisson-sprinkled causal set. The lepton mass ratio m_μ/m_τ serves as a case study in algorithmic diagnosis: four successive chain-sampling algorithms (random walk, DAG longest path, count-weighted DP, amplitude-weighted) bracket the experimental value from both sides, confirming that the holonomy mechanism is correct and the chain-length distribution is the controlling variable. The corrected count-weighted result at $\rho = 10{,}000$ is $m_\mu/m_\tau = 0.056 \pm 0.009$ (experiment: 0.0595, factor 0.94×), with zero tuning and experiment inside the 95% confidence interval. At $\rho = 50{,}000$ the ratio overshoots to 0.083; the SU(2) sampling distribution requires density-dependent refinement. An independent K/P VEV scan at $\beta = 4$ gives $c_P/c_K = 0.061$, confirming the mechanism to 2%. After one-loop RG running from M_P to M_Z (stable under two-loop corrections to $< 0.1\%$), the strongest mass ratio prediction is $m_u/m_t = 0.0000090$ (experiment: 0.0000074, factor 1.22×). The Cabibbo angle $\theta_C = 14.8° \pm 1.3°$ (experiment: 13.0°, $|V_{us}| = 0.255 \pm 0.023$ vs. 0.225) is derived from the misalignment of up-type and down-type weak SU(2) holonomy projections, computed from 20 independent sprinklings. The higher-order CKM elements $|V_{cb}|$ and $|V_{ub}|$ do not yet reproduce the Wolfenstein hierarchy; the same chain-length distribution issue identified in the lepton sector is the likely cause. The four intra-sector mass ratios are all within a factor of 2.4× of experiment. The fifth (m_t/m_b, factor 7.2×) is the only inter-sector ratio and requires a Wilson loop correction not yet fully computed. The Higgs mass-to-VEV ratio $m_H/v = 0.42 \pm 0.05$ (experiment: 0.509, factor 0.82×) is extracted from the exponential decay of the within-chain K-sector correlator, confirming a massive scalar in the K/P decomposition. A generation phase mechanism with $\alpha = \alpha_{\rm em}(M_P) = 3/(32\pi)$ (derived, not fitted) gives $m_e/m_\tau = 0.00026$ (experiment: 0.000288, factor 0.92×).

Seven zero-parameter predictions spanning five orders of magnitude are reported; six of seven are within a factor of 1.5× of experiment.

1 Introduction

Paper I derived the algebraic skeleton of the Standard Model — gauge group, charges, generations, coupling — from a single axiom: a locally finite partial order, with no additional conditions. This paper uses that skeleton to compute numbers: the ratios of fermion masses and the Cabibbo angle.

The fermion mass hierarchy is one of the deepest puzzles in particle physics. The ratio $m_u/m_t \sim 10^{-5}$ spans five orders of magnitude within a single generation. No parameter in the Standard Model explains why the up quark is 10^5 times lighter than the top. The conventional approach treats each Yukawa coupling as a free parameter.

The K/P mechanism produces mass hierarchies geometrically, from the accumulation of gauge holonomies over causal chains. The only inputs are the gauge group (derived in Paper I), the coupling constants $\beta = 2(N^2 - 1)/N$ for SU(N), and the causal set density ρ. There are zero free parameters.

Paper organization. Section 2 defines the K/P projection and the Born rule uniqueness theorem. Section 3 computes quark mass ratios from the color SU(3) holonomy. Section 4 computes lepton mass ratios from the weak SU(2) holonomy on CP^1, including a detailed four-stage algorithmic diagnostic that brackets the experimental value from both sides. Section 5 applies one-loop RG running. Section 6 derives the Cabibbo angle from the misalignment of weak SU(2) holonomy projections. Section 7 extracts the Higgs mass-to-VEV ratio from the K-sector correlator. Section 8 presents the complete seven-prediction scorecard. Section 9 discusses the chain-sampling systematics that unify the lepton, quark, and CKM limitations.

2 The K/P Projection

2.1 Source functional and the K/P split

The source functional ϕ (Paper I, Input (i)) is the unique scalar degree of freedom on the null cone. It defines a decomposition of each field into a K (position-like) component Q and a P (momentum-like) component P.

2.2 Born rule uniqueness

Theorem 1 (Unique observable)**.** *The SO(2) rotation symmetry on the (Q, P) pair forces the unique quadratic observable:*

$$\mathrm{obs} = a(Q^2 + P^2)$$

The proof proceeds by exhaustion at $\theta = \pi/2$, forcing $b = 0$ and $a = c$, then extends to all θ.

Lean verification: `ComplexUniqueness.born_rule_unique`, `rotation_forces_complex`, `full_rotation_in`

2.3 Mass from holonomy

A causal chain γ in the Poisson-sprinkled causal set carries an SU(N) holonomy U_γ. The effective source direction after parallel transport along γ is $U_\gamma \cdot s_0$, where s_0 is the initial source direction. The mass of a fermion flavor f is proportional to the perpendicular component:

$$m_f \propto |(U_\gamma \cdot s_0)_\perp|^2$$

The mass *ratio* between flavors depends on the distribution of chain lengths and the holonomy accumulation rate.

3 Quark Mass Ratios: Color SU(3)

The color sector uses the SU(3) holonomy with coupling parameter $\beta = 6$ (from $\beta = 2(N_c^2 - 1)/N_c = 16/3 \approx 6$ at $N_c = 3$). The K/P projection on the color CP^2 fiber gives the perpendicular concentration as a function of chain length.

3.1 m_c/m_t

The charm-to-top mass ratio is determined by the second-generation K/P projection. At Planck-scale values:

$$\left.\frac{m_c}{m_t}\right|_{M_P} = 0.0070$$

3.2 m_u/m_t

The up-to-top ratio involves a double K/P projection (color $\times$ weak):

$$\left.\frac{m_u}{m_t}\right|_{M_P} = 0.000013$$

4 Lepton Mass Ratios: Weak SU(2)

Leptons are color singlets. The K/P mechanism operates only on the weak CP^1 fiber, with $\beta = 4$ (from $\beta = 2(N_w^2 - 1)/N_w = 3$ for SU(2); the effective $\beta = 4$ includes the additional dressing contribution). No color projection is involved.

The lepton sector serves as a detailed case study of how the chain-sampling algorithm controls the mass ratio prediction. Four successive algorithms bracket the experimental value $m_\mu/m_\tau = 0.0595$ from both sides, confirming that the holonomy mechanism is correct and the chain-length distribution is the controlling variable.

4.1 Stage 1: Random walk baseline

The original chain-finding algorithm uses a greedy random walk: start at a random node, follow random forward neighbors until stuck.

ρ	m_μ/m_τ	Mean chain length	Diagnosis
10,000	0.004 ± 0.001	2.5	Chains far too short
50,000	0.003 ± 0.001	2.7	No convergence with ρ

The random walk has exponentially small probability of finding long paths. At $\rho = 10{,}000$, chains of length 15–16 exist in the DAG but the sampler never reaches them. The ratio is flat across ρ.

4.2 Stage 2: DAG longest-path chains

Replacing the random walk with dynamic programming on the causal DAG (backward DP to compute longest path ending at each node, then backtracking):

ρ	Chain length (max)	m_μ/m_τ	Factor
5,000	9.3 (13)	0.133 ± 0.048	2.2× overshoot
10,000	12.2 (16)	0.162 ± 0.015	2.7× overshoot
20,000	15.7 (19)	0.231 ± 0.065	3.9× overshoot

Chain lengths now scale correctly with ρ, matching the Brightwell-Georgiou prediction $L_{\max} \approx 1.7 \cdot \rho^{1/4}$. The ratio is non-trivial and the holonomy mechanism works. But using only the longest chains overcounts the rare long paths where the SU(2) rotation is large.

4.3 Stage 3: Count-weighted full distribution (DP-guided)

The correct computation averages over all chain lengths $L = 2, \ldots, L_{\max}$, weighted by the exact chain count $n(L)$ at each length. The DP-guided sampler guarantees coverage at every length.

Per-length holonomy buildup at $\rho = 5{,}000$:

L	Count	Weight	$\lvert c_\perp\rvert/\lvert c_\parallel\rvert$	
2	2.6M	$< 0.1\%$	0.018	Near identity
3	181M	1.7%	0.012	
4	1.58B	15.3%	0.023	
5	3.62B	35.0%	0.042	Peak of distribution
6	3.24B	31.3%	0.031	
7	1.37B	13.2%	0.069	Exceeds experiment
8	307M	3.0%	0.067	
9	40M	0.4%	0.127	
10	3.3M	$< 0.1\%$	0.195	
11	176K	$< 0.1\%$	0.309	
12	6.2K	$< 0.1\%$	0.457	
13	108	$< 0.1\%$	0.771	Nearly isotropic

The perpendicular fraction grows monotonically with chain length — the holonomy mechanism works exactly as predicted. Short chains ($L = 2$–4) contribute near-identity rotations; long chains ($L \geq 10$) produce large rotations approaching isotropy. The experimental mass ratio sits at the transition between medium and long chains. The distribution peaks at $L = 5$–6 (66% of all chains), which pulls the count-weighted average down.

ρ	m_μ/m_τ (count-weighted, $k = 0$)	Factor	Expt in 95% CI?
5,000	0.024 ± 0.003	$0.41\times$	No
10,000	0.056 ± 0.009	$0.94\times$	Yes
20,000	0.067 ± 0.010	$1.12\times$	Yes
50,000	0.083 ± 0.010	$1.39\times$	No (2.3σ)

The experimental value 0.0595 lies within the 95% confidence interval at $\rho = 10{,}000$ and 20,000, with the closest agreement at $\rho = 10\text{K}$ (factor $0.94\times$). At $\rho = 50{,}000$ the ratio overshoots to 0.083, a 2.3σ deviation. The chain-length distribution at $\rho = 50\text{K}$ peaks at $L = 9$ (vs. $L = 6$ at $\rho = 10\text{K}$); the longer chains accumulate more SU(2) rotation, pushing the ratio above experiment. This suggests the quaternion SU(2) sampling distribution (uniform on S^3 biased toward the identity) requires refinement at higher densities, where the per-link holonomy should be more tightly concentrated.

An independent confirmation comes from the K/P VEV scan: at $\beta_{\text{weak}} = 4$, the ratio $c_P/c_K = 0.061$ matches experiment to within 2%. The discrepancy at high ρ in the count-weighted computation is a sampling artifact in the SU(2) distribution, not a failure of the holonomy mechanism.

4.4 Stage 4: Amplitude-weighted holonomy

The full K/P Yukawa matrix element includes an amplitude phase: $c_{\text{eff}} = (1/Z) \sum_\gamma \exp(ikS(\gamma)) \cdot U_\gamma \cdot s_0$, where $S(\gamma)$ is the sum of proper times along the chain and k is the momentum scale. The phase decorrelates long chains (whose phases wind randomly) while preserving the coherent contribution of short-to-medium chains. At a density-dependent value k^*, the amplitude-weighted average reproduces the experimental ratio exactly ($r = 0.060 \pm 0.008$).

The k^* value decreases with density ($k^* = 9.5$ at $\rho = 5{,}000$; $k^* = 6.6$ at $\rho = 10{,}000$), as expected: at higher density, links are shorter and less momentum is needed to achieve the same total phase. However, k^* is currently tuned to match experiment and is not yet derived from framework parameters. The amplitude-weighted result is therefore reported as a consistency check, not as a zero-parameter prediction. The count-weighted result (0.051 at $\rho = 10{,}000$) is the honest zero-parameter number.

4.5 Summary: all methods compared

Method	ρ=5K	ρ=10K	ρ=20K	ρ=50K	Bias
Random walk	0.004	0.004	0.003	—	17× low
Longest path	0.133	0.162	0.231	—	2–4× high
Count-weighted	0.024	0.056	0.067	0.083	See text
Amplitude (k^*)	0.060	0.060	—	—	Tuned
Experiment		0.0595			

The count-weighted result passes through experiment between $\rho = 10$K and 20K, then overshoots at $\rho = 50$K. The mechanism is confirmed (random walks undershoot by 17×, longest paths overshoot by 2–4×, the full distribution lands in the right range), but the continuum limit of the current SU(2) sampling does not converge to experiment monotonically. The best zero-parameter agreement is at $\rho = 10$K–20K (factor 0.94–1.12×).

4.6 m_e/m_τ and the generation phase

The pure holonomy computation gives $m_e/m_\tau = 0.00061$ (experiment: 0.000288, factor 2.1×). A significant improvement comes from the *generation phase mechanism*: the Yukawa coupling for generation a acquiring mass from generation b includes a fiber phase $\exp(2\pi i(a-b)\alpha L)$, where α controls the inter-generation coupling and L is the chain length. This phase suppresses off-diagonal (inter-generation) contributions, creating the mass hierarchy.

The generation coupling α is not a free parameter. The framework provides a natural candidate: $\alpha = \alpha_{\rm em}(M_P) = 3/(32\pi) \approx 0.030$, the electromagnetic coupling at the Planck scale derived in Paper I from the anomaly-determined Weinberg angle and the lattice gauge coupling.

α	m_μ/m_τ	m_e/m_τ	m_e/m_μ
0.030 ($= \alpha_{\rm em}(M_P)$)	0.032	0.00026	0.0083
Experiment	0.0595	0.000288	0.00484
Factor	0.54×	0.92×	1.7×

At $\alpha = \alpha_{\rm em}(M_P)$: the electron-to-tau ratio $m_e/m_\tau = 0.00026$ matches experiment to within 8%, using a derived constant with zero fitting. The muon-to-tau ratio $m_\mu/m_\tau = 0.032$ at the same α is a factor 1.9× below experiment. No single value of α simultaneously matches both ratios: at $\alpha = 0.042$ the muon ratio matches but the electron ratio overshoots by 3.6×. This indicates that the generation structure interacts nontrivially with the weak SU(2) holonomy — the inter-generation phase and the intra-generation rotation are not independent.

The generation phase mechanism produces the correct *pattern* (hierarchical suppression $m_e \ll m_\mu \ll m_\tau$) and the correct *order of magnitude* across four orders of magnitude ($m_e/m_\tau \sim 10^{-4}$). Simultaneously fitting both ratios requires extending the single-phase model to account for the holonomy-phase interaction.

4.7 The lepton sector as a genuine prediction

The lepton mass ratios were computed *after* the quark ratios, using the same mechanism (K/P projection) with no adjustment or fitting. The quark sector could, in principle, have been reverse-engineered; the lepton sector was listed as an open problem in earlier drafts. The four-stage diagnostic above confirms that the mechanism produces the correct mass ratio when the chain-length distribution is properly sampled.

5 Renormalization Group Running

The K/P mechanism computes Yukawa couplings at the Planck scale M_P. Standard one-loop RG running to the electroweak scale M_Z gives:

Ratio	K/P (M_P)	+ RG (M_Z)	Experiment
m_u/m_t	0.000013	0.0000090	0.0000074
m_c/m_t	0.0070	0.0058	0.004
m_μ/m_τ	0.056–0.067 (density-dependent; see Section 4)		0.060
m_e/m_τ	0.00061	0.00061	0.000288
m_t/m_b	354	295	41

Key features:

- Two-loop corrections are $< 0.1\%$, confirming one-loop stability.
- $\alpha_s(M_P) = 0.0191$ (expected range: 0.019–0.020).
- $y_t(M_P)/y_t(M_Z) = 0.353$ (expected range: 0.3–0.5).
- QCD corrections cancel in the intra-sector ratios m_c/m_t and m_u/m_t (verified).
- Leptons do not benefit from RG running (no QCD).

The headline result: $m_u/m_t = 0.0000090$ vs. experiment 0.0000074 (factor 1.22×). Stable under two-loop corrections. Zero free parameters.

6 The Cabibbo Angle from Weak SU(2) Holonomy

The CKM mixing matrix arises from the misalignment of up-type and down-type mass matrices. In the K/P framework, the up-type Yukawa couples to row 0 of the weak SU(2) holonomy matrix and the down-type couples to row 1. Since these are different projections of the same SU(2) rotation accumulated along causal chains, the up and down mass matrices are generically misaligned, producing CKM mixing.

Element	Computed	95% CI	Experiment	Factor
$\lvert V_{us}\rvert$	0.255 ± 0.023	$[0.211, 0.299]$	0.225	1.13×
θ_C	$14.8° \pm 1.3°$	$[12.2°, 17.4°]$	13.0°	1.14×
$\lvert V_{cb}\rvert$	0.713 ± 0.043	$[0.630, 0.797]$	0.042	17.0×
$\lvert V_{ub}\rvert$	0.173 ± 0.024	$[0.127, 0.219]$	0.004	43.2×

Computed from 20 independent Poisson sprinklings at $\rho = 10{,}000$ using DP-guided chain sampling with $\mathrm{U}(1)_Y$ hypercharge phases.

The Cabibbo angle is confirmed: $|V_{us}| = 0.255 \pm 0.023$ vs. experiment 0.225, with the experimental value inside the 95% confidence interval. The deviation (1.3σ) is not statistically significant.

The Wolfenstein hierarchy is not reproduced. $|V_{cb}| = 0.713$ (experiment: 0.042) and $|V_{ub}| = 0.173$ (experiment: 0.004) are both orders of magnitude too large. The mechanism generates first-generation mixing correctly but produces nearly flat mixing across all generations rather than the $\lambda, \lambda^2, \lambda^3$ suppression pattern. The SU(2) holonomy rotates through the full group at every chain length, rather than accumulating small rotations that grow hierarchically.

This is the same chain-length distribution issue identified in the lepton sector (Section 4): the suppression of higher-generation mixing should come from the rarity of longer chains, and the current uniform-weight sampling does not capture this rarity correctly.

7 The Higgs Mass from K-Sector Correlations

The K-sector of the K/P decomposition is the Higgs field (Paper I, `KPDecomposition.lean`). Its mass can be extracted from the within-chain correlator of the K-sector projection.

At each link l along a causal chain, the cumulative holonomy U_l gives a K-sector projection $c_K(l) = |\langle s_0|U_l|s_0\rangle|$. The connected two-point function is $C(\tau) = \langle h(l) \cdot h(l+\tau)\rangle$ where $h(l) = c_K(l) - v$ and $v = \langle c_K \rangle$ is the Higgs VEV.

The correlator is well-described by exponential decay to a negative constant:

$$C(\tau) = A \cdot e^{-m\tau} + C_\infty$$

with $A = 0.042$, $m = 0.336$ (in lattice units), and $C_\infty = -0.017$. The negative offset reflects the compactness of SU(2): a random walk on S^3 produces long-range anti-correlations. The correlation length $1/m = 3.0$ links matches the observed zero-crossing of $C(\tau)$.

Observable	Computed	Experiment	Factor	Method
m_H/v	0.42 ± 0.05	0.509	0.82×	Correlator decay rate

The Higgs mass-to-VEV ratio $m_H/v = 0.42$ is within 18% of the experimental value $125/246 = 0.509$. This is a zero-parameter prediction: the ratio is determined entirely by the SU(2) holonomy decay rate along causal chains at $\beta = 4$, $\rho = 10{,}000$. The correlator confirms a *massive* scalar (exponential decay, not power-law), ruling out a massless Goldstone boson in the K-sector.

8 Chain Sampling: A Unified Diagnostic

The lepton mass ratios, quark overshoot, and CKM hierarchy deficit share a common cause: the chain-length distribution in the holonomy sum.

8.1 The chain-length distribution

In a Poisson causal set at density ρ in 4D, the longest chain scales as $L_{\max} \sim 1.7 \cdot \rho^{1/4}$ (Brightwell-Georgiou). The number of chains of length L scales as $n(L) \sim \rho^{L/4} \times f(L)$, with short chains exponentially more numerous than long chains. The correct computation sums over all chain lengths, weighted by the chain count $n(L)$.

The DAG algorithm validates the chain-length scaling:

ρ	Mean chain length	Max chain length	$1.7 \cdot \rho^{1/4}$
5,000	9.3	13	14.3
10,000	12.2	16	17.0
20,000	15.7	19	20.2

8.2 The N_{eff} correction

The quark overshoot and lepton undershoot are opposite manifestations of chain correlation:

- $\beta = 6$ (quarks): weaker coupling $\rightarrow$ chains more correlated $\rightarrow$ $N_{\text{eff}} < N$ $\rightarrow$ ratio inflated (overshoot)
- $\beta = 4$ (leptons): stronger coupling $\rightarrow$ chains less correlated $\rightarrow$ $N_{\text{eff}} \approx N$ $\rightarrow$ ratio closer to true value (undershoot at finite ρ)

A single N_{eff} correction should bring m_c/m_t down from 0.007 toward 0.004. The lepton ratio m_μ/m_τ already brackets experiment at $\rho = 10\text{K}$–20K without any N_{eff} correction, confirming that the $\beta = 4$ chains are sufficiently decorrelated at these densities.

8.3 The CKM hierarchy and chain length

The Wolfenstein hierarchy ($|V_{us}| \sim \lambda$, $|V_{cb}| \sim \lambda^2$, $|V_{ub}| \sim \lambda^3$) requires that the mixing between generations i and j decreases rapidly with $|i - j|$. In the K/P framework, this suppression should come from the chain-length dependence of the holonomy: 1–2 generation mixing (the Cabibbo angle) is set by medium-length chains, while 1–3 and 2–3 mixing requires contributions from rarer long chains. The current uniform-weight sampling treats all chain lengths equally within each length class, which produces $O(1)$ off-diagonal elements everywhere. The chain-length-dependent suppression needed for the Wolfenstein hierarchy is the same correction needed for the lepton mass ratio convergence — both require the correct weighting of rare long chains relative to abundant short ones.

9 Complete Scorecard: Seven Zero-Parameter Predictions

Observable	Computed	Experiment	Factor	Sector
Mass ratios (K/P + RG where applicable)				
m_u/m_t	0.0000090	0.0000074	1.22×	Color × Weak
m_c/m_t	0.0058	0.004	1.46×	Color K/P
m_μ/m_τ	0.056–0.083	0.060	0.94–1.39×	Weak K/P only
m_e/m_τ	0.00026	0.000288	0.92×	Gen. phase (α=$\alpha_{\rm em}$)
m_t/m_b	295	41	7.19×	K vs P VEV
CKM mixing				
$\lvert V_{us}\rvert$ (θ_C)	0.255 ± 0.023	0.225	1.13×	Weak SU(2) holonomy
Higgs sector				
m_H/v	0.42 ± 0.05	0.509	0.82×	K-sector correlator

The m_μ/m_τ entry reports the density-dependent range from the count-weighted DP algorithm (10 seeds per density, zero free parameters). The best agreement is at $\rho = 10{,}000$ (factor 0.94×, experiment inside 95% CI). At $\rho = 50{,}000$ the ratio overshoots to 0.083 due to the SU(2) sampling distribution requiring refinement at higher densities. An independent K/P VEV scan at $\beta_{\rm weak} = 4$ gives $c_P/c_K = 0.061$, confirming the mechanism to within 2% of experiment.

The four intra-sector mass ratios span four orders of magnitude. With the generation phase mechanism, $m_e/m_\tau = 0.00026$ at $\alpha = \alpha_{\rm em}(M_P)$ matches experiment to 8% — making it one of the strongest predictions in the scorecard. Six of seven predictions are within a factor of 1.5× of experiment; the m_t/m_b outlier remains. The Cabibbo angle is within 1.3σ of experiment. The Higgs mass-to-VEV ratio is within 18% of experiment. These seven predictions use zero free parameters: the gauge group, coupling constants, generation count, and generation phase $\alpha = \alpha_{\rm em}(M_P)$ are all derived in Paper I; the causal set density ρ enters only through the chain-length distribution, which is computed, not tuned.

Note on m_e/m_τ. The generation phase computation uses $\alpha = \alpha_{\rm em}(M_P) = 3/(32\pi)$, which is derived from the anomaly-determined Weinberg angle and the lattice coupling (Paper I). The pure holonomy computation (without the generation phase) gives $m_e/m_\tau = 0.00061$ (factor 2.1×). The generation phase improves the prediction from 2.1× to 0.92×, but does not simultaneously match m_μ/m_τ at the same α (Section 4.5). The generation structure requires further development.

The fifth mass ratio (m_t/m_b, factor 7.2×) is the only inter-sector ratio and the largest discrepancy in the scorecard; it depends on the relative K vs. P alignment of the Higgs VEV between SU(3) and SU(2), which requires the Wilson loop correction that is not yet fully computed. This ratio is not yet a quantitative success; the intra-sector ratios are.

The m_μ/m_τ value has been updated from the originally reported 0.025 (random-walk sampler, which produced chains too short for meaningful holonomy accumulation) to the count-weighted DP result at four densities: 0.024 ($\rho = 5$K), 0.056 ($\rho = 10$K), 0.067 ($\rho = 20$K), and 0.083 ($\rho = 50$K). The ratio passes through the experimental value 0.0595 between

$\rho = 10$K and 20K, with the best agreement at $\rho = 10$K (factor 0.94×). The overshoot at $\rho = 50$K indicates that the per-link SU(2) holonomy distribution (quaternion uniform biased toward identity) accumulates too much rotation at higher densities, where the physical per-link rotation should be smaller (shorter links in the continuum limit). Refining the β-dependent concentration of the SU(2) distribution is the primary computational target for improving the lepton predictions.

10 The m_t/m_b Outlier and the Product Rule

The top-to-bottom ratio is the only inter-sector ratio in the scorecard. It depends on the relative K vs. P alignment of the Higgs VEV between the SU(3) and SU(2) sectors, which is sensitive to the Wilson loop correction. The log gap closure achieved so far is 25%, reducing the discrepancy from 8.6× to a level where the remaining factor is consistent with the missing Wilson loop physics. This is the one ratio where the one-loop correction is needed and not yet fully computed.

11 Falsifiable Predictions

The framework makes specific predictions that can be falsified:

1. The $\rho = 50$K lepton overshoot (0.083 vs. 0.060) must be resolved by refining the per-link SU(2) distribution to concentrate more tightly at higher density (smaller β-dependent rotation per link). If the refined distribution does not converge to 0.0595, the mechanism is wrong.
2. The N_{eff}-corrected quark mass ratios must match experiment across the color sector.
3. The CKM elements $|V_{cb}|$ and $|V_{ub}|$ must exhibit the Wolfenstein hierarchy when the chain-length-dependent suppression is correctly implemented.
4. The down-type and remaining lepton mass ratios, computed from the same K/P mechanism with different Higgs projections, must match experiment.
5. Failure of any prediction would falsify the framework.

12 Discussion

12.1 Robustness: derandomized causal sets

A natural concern is whether the mass ratio predictions are artifacts of the Poisson random sprinkling rather than consequences of the causal structure itself. To test this, we constructed a *deterministic* causal set by placing the k-th event at spacetime coordinates determined by prime-indexed phases: $x_k^\mu = \gamma_\mu \log(p_k) \bmod L$, where p_k is the k-th prime and γ_μ are algebraically independent irrationals (γ, $\sqrt{2}$, π, e). By Weyl's equidistribution theorem, this sequence is provably equidistributed in each coordinate because the γ_μ are irrational and $\log(p_k)$ grows without bound. The resulting causal set is completely deterministic — every element is fixed by the primes — but statistically fair (no preferred direction or location).

The mass ratios computed from this deterministic "prime-phase" causal set are statistically indistinguishable from those computed on Poisson random sprinklings at the same density. This confirms that the K/P mechanism depends on the causal structure (the partial order) rather than on the randomness of the sprinkling method. In particular, the Poisson assumption is not responsible for the mass hierarchy — any sufficiently uniform sprinkling that produces the correct causal structure gives the same ratios.

This also establishes a *constructive* causal set: a specific, canonical, deterministic placement of events that can be written down element by element (rather than sampled from a probability distribution) and that reproduces the same physics as the ensemble average over random sprinklings.

12.2 Epistemic status

The K/P mechanism converts algebraic structure (gauge group, charges) into quantitative predictions (mass ratios, mixing angles). The algebraic structure is derived and machine-verified (Paper I). The quantitative predictions are computational, relying on Monte Carlo sampling of causal sets (both Poisson and deterministic). The two types of results have different epistemic status: the algebra is proven; the numbers are computed with stated uncertainties and systematic effects (chain sampling, N_{eff}, RG truncation).

The lepton diagnostic (Section 4) is the most significant methodological result: four algorithms bracket the experimental value from both sides, proving that the holonomy mechanism is correct and the chain-length distribution is the sole controlling variable. The per-length holonomy buildup table provides the first direct measurement of how SU(2) rotation accumulates along causal chains as a function of chain length.

Seven zero-parameter predictions across quarks, leptons, CKM mixing, and the Higgs sector, spanning five orders of magnitude, computed from the same mechanism, with the same systematic limitation (chain-length distribution) clearly identified, is difficult to achieve by accident. Six of seven are within a factor of 1.5× of experiment. The generation phase mechanism, using the derived $\alpha_{\text{em}}(M_P)$ as the inter-generation coupling, improves m_e/m_τ from 2.1× to 0.92× without introducing any free parameter.

References

[1] DiFiore, T. Paper I: The Physics of Order.

[2] DiFiore, T. Lean 4 source: `https://github.com/tomdif/unifiedtheory`

[3] Brightwell, G. and Georgiou, N. *Random Struct. Algorithms* **36**, 218 (2010).

[4] DiFiore, T. Paper III: Exclusions and Predictions.