


Domain Theory in Constructive and Predicative Univalent Foundations

Tom de Jong 

University of Birmingham, United Kingdom

<https://www.cs.bham.ac.uk/~txd880>

t.dejong@pgr.bham.ac.uk

Martín Hötzel Escardó 

University of Birmingham, United Kingdom

<https://www.cs.bham.ac.uk/~mhe>

m.escardo@cs.bham.ac.uk

Abstract

We develop domain theory in constructive univalent foundations without Voevodsky’s resizing axioms. In previous work in this direction, we constructed the Scott model of PCF and proved its computational adequacy, based on directed complete posets (dcpo). Here we further consider algebraic and continuous dcpo, and construct Scott’s D_∞ model of the untyped λ -calculus. A common approach to deal with size issues in a predicative foundation is to work with *information systems* or *abstract bases* or *formal topologies* rather than dcpo, and *approximable relations* rather than Scott continuous functions. Here we instead accept that dcpo may be large and work with type universes to account for this. For instance, in the Scott model of PCF, the dcpo have carriers in the second universe \mathcal{U}_1 and suprema of directed families with indexing type in the first universe \mathcal{U}_0 . Seeing a poset as a category in the usual way, we can say that these dcpo are large, but locally small, and have small filtered colimits. In the case of algebraic dcpo, in order to deal with size issues, we proceed mimicking the definition of accessible category. With such a definition, our construction of Scott’s D_∞ again gives a large, locally small, algebraic dcpo with small directed suprema.

2012 ACM Subject Classification Theory of computation \rightarrow Constructive mathematics; Theory of computation \rightarrow Type theory

Keywords and phrases domain theory, constructivity, predicativity, univalent foundations

Digital Object Identifier 10.4230/LIPIcs.CVIT.2016.23

1 Introduction

In domain theory [1] one considers posets with suitable completeness properties, possibly generated by certain elements called *compact*, or more generally generated by a certain *way below* relation, giving rise to algebraic and continuous domains. As is well known, domain theory has applications to programming language semantics [40, 38, 31], higher-type computability [25], topology, topological algebra and more [17, 16].

In this work we explore the development of domain theory from the univalent point of view [43, 46]. Additionally, we work constructively (we don’t assume excluded middle or choice axioms) and predicatively (we don’t assume Voevodsky’s resizing principles [44, 45, 46], and so, in particular, powersets are large). Most of the work presented here has been formalized in the proof assistant Agda [6, 15, 10] (see Section 7 for details). In our predicative setting, it is extremely important to check universe levels carefully, and the use of a proof assistant such as Agda has been invaluable for this purpose.

In previous work in this direction [8] (extended by Brendan Hart [18]), we constructed the Scott model of PCF and proved its computational adequacy, based on directed complete posets (dcpo). Here we further consider algebraic and continuous dcpo, and construct Scott’s D_∞ model of the untyped λ -calculus [38].



© Tom de Jong and Martín H. Escardó;
licensed under Creative Commons License CC-BY

42nd Conference on Very Important Topics (CVIT 2016).

Editors: John Q. Open and Joan R. Access; Article No. 23; pp. 23:1–23:25

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

A common approach to deal with size issues in a predicative foundation is to work with *information systems* [39], *abstract bases* [1] or *formal topologies* [36, 7] rather than dcpos, and *approximable relations* rather than (Scott) continuous functions. Here we instead accept that dcpos may be large and work with type universes to account for this. For instance, in our development of the Scott model of PCF [40, 31], the dcpos have carriers in the second universe \mathcal{U}_1 and suprema of directed families with indexing type in the first universe \mathcal{U}_0 . Seeing a poset as a category in the usual way, we can say that these dcpos are large, but locally small, and have small filtered colimits. In the case of algebraic dcpos, in order to deal with size issues, we proceed mimicking the definition of accessible category [27]. With such a definition, our construction of Scott's D_∞ again gives a large, locally small, algebraic dcpo with small directed suprema.

Organization

Section 2: Foundations. *Section 3:* (Im)predicativity. *Section 4:* Basic domain theory, including directed complete posets, continuous functions, lifting, Ω -completeness, exponentials, powersets as dcpos. *Section 5:* Limit and colimits of dcpos, Scott's D_∞ . *Section 6:* Way below relation, bases, compact element, continuous and algebraic dcpos, ideal completion, retracts, examples. *Section 7:* Conclusion and future work.

Related Work

Domain theory has been studied predicatively in the setting of *formal topology* [36, 7] in [37, 28, 29, 26] and the more recent categorical paper [22]. In this predicative setting, one avoids size issues by working with abstract bases or formal topologies rather than dcpos, and approximable relations rather than Scott continuous functions. Hedberg [19] presented these ideas in Martin-Löf Type Theory and formalized them in the proof assistant ALF. A modern formalization in Agda based on Hedberg's work was recently carried out in Lidell's master thesis [24].

Our development differs from the above line of work in that it studies dcpos directly and uses type universes to account for the fact that dcpos may be large. There are two Coq formalizations of domain theory in this direction [5, 11]. Both formalizations study ω -chain complete preorders, work with setoids, and make use of Coq's impredicative sort **Prop**. In our development we avoid the use of setoids thanks to the adoption of the univalent point of view. Moreover, we work predicatively and we work with directed sets rather than ω -chains, as we intend our theory to be also applicable to topology and algebra [17, 16].

There are also constructive accounts of domain theory aimed at program extraction [4, 30]. Both [4] and [30] study ω -chain complete posets (ω -cpo) and define notions of ω -continuity for them. Interestingly, Bauer and Kavkler [4] note that there can only be non-trivial examples of ω -continuous ω -cpo when Markov's Principle holds [4, Proposition 6.2]. This leads the authors of [30] to weaken the definition of ω -continuous ω -cpo by using the double negation of existential quantification in the definition of the way below relation [30, Remark 3.2]. In light of this, it is interesting to observe that when we study directed complete posets (dcpos) rather than ω -cpo, and continuous dcpos rather than ω -continuous ω -cpo, we can avoid Markov's Principle or a weakened notion of the way below relation to obtain non-trivial continuous dcpos (see for instance Examples 58, 59 and 82).

Another approach is the field of *synthetic domain theory* [35, 34, 20, 32, 33]. Although the work in this area is constructive, it is still impredicative, based on topos logic, but more importantly it has a focus different from that of regular domain theory: the aim is to isolate

a few basic axioms and find models in (realizability) toposes where “every object is a domain and every morphism is continuous”. These models often validate additional axioms, such as Markov’s Principle and countable choice, and moreover falsify excluded middle. Our development has a different goal, namely to develop regular domain theory constructively and predicatively, but in a foundation compatible with excluded middle and choice, while not relying on them or Markov’s Principle or countable choice.

2 Foundations

We work in intensional Martin-Löf Type Theory with type formers $+$ (binary sum), Π (dependent products), Σ (dependent sum), Id (identity type), and inductive types, including $\mathbf{0}$ (empty type), $\mathbf{1}$ (type with exactly one element $\star : \mathbf{1}$), \mathbf{N} (natural numbers). Moreover, we have type universes (for which we typically write \mathcal{U} , \mathcal{V} , \mathcal{W} or \mathcal{T}) with the following closure conditions. We assume a universe \mathcal{U}_0 and two operations: for every universe \mathcal{U} a successor universe \mathcal{U}^+ with $\mathcal{U} : \mathcal{U}^+$, and for every two universes \mathcal{U} and \mathcal{V} another universe $\mathcal{U} \sqcup \mathcal{V}$ such that for any universe \mathcal{U} , we have $\mathcal{U}_0 \sqcup \mathcal{U} \equiv \mathcal{U}$ and $\mathcal{U} \sqcup \mathcal{U}^+ \equiv \mathcal{U}^+$. Moreover, $(-) \sqcup (-)$ is idempotent, commutative, associative, and $(-)^+$ distributes over $(-) \sqcup (-)$. We write $\mathcal{U}_1 \equiv \mathcal{U}_0^+$, $\mathcal{U}_2 \equiv \mathcal{U}_1^+$, \dots and so on. If $X : \mathcal{U}$ and $Y : \mathcal{V}$, then $X + Y : \mathcal{U} \sqcup \mathcal{V}$ and if $X : \mathcal{U}$ and $Y : X \rightarrow \mathcal{V}$, then the types $\Sigma_{x:X} Y(x)$ and $\Pi_{x:X} Y(x)$ live in the universe $\mathcal{U} \sqcup \mathcal{V}$; finally, if $X : \mathcal{U}$ and $x, y : X$, then $\text{Id}_X(x, y) : \mathcal{U}$. The type of natural numbers \mathbf{N} is assumed to be in \mathcal{U}_0 and we postulate that we have copies $\mathbf{0}_{\mathcal{U}}$ and $\mathbf{1}_{\mathcal{U}}$ in every universe \mathcal{U} . All our examples go through with just two universes \mathcal{U}_0 and \mathcal{U}_1 , but the theory is more easily developed in a general setting.

In general we adopt the same conventions of [43]. In particular, we simply write $x = y$ for the identity type $\text{Id}_X(x, y)$ and use \equiv for the judgemental equality, and for dependent functions $f, g : \Pi_{x:X} A(x)$, we write $f \sim g$ for the pointwise equality $\Pi_{x:X} f(x) = g(x)$.

Within this type theory, we adopt the univalent point of view [43]. A type X is a *proposition* (or *truth value* or *subsingleton*) if it has at most one element, i.e. the type $\text{is-prop}(X) \equiv \prod_{x,y:X} x = y$ is inhabited. A major difference between univalent foundations and other foundational systems is that we *prove* that types are propositions or properties. For instance, we can show (using function extensionality) that the axioms of directed complete poset form a proposition. A type X is a *set* if any two elements can be identified in at most one way, i.e. the type $\prod_{x,y:X} \text{is-prop}(x = y)$ is inhabited.

We will assume two extensionality principles:

- (i) *Propositional extensionality*: if P and Q are two propositions, then we postulate that $P = Q$ exactly when both $P \rightarrow Q$ and $Q \rightarrow P$ are inhabited.
- (ii) *Function extensionality*: if $f, g : \prod_{x:X} A(x)$ are two (dependent) functions, then we postulate that $f = g$ exactly when $f \sim g$.

Function extensionality has the important consequence that the propositions form an exponential ideal, i.e. if X is a type and $Y : X \rightarrow \mathcal{U}$ is such that every $Y(x)$ is a proposition, then so is $\Pi_{x:X} Y(x)$. In light of this, universal quantification is given by Π -types in our type theory.

In Martin-Löf Type Theory, an element of $\prod_{x:X} \sum_{y:Y} \phi(x, y)$, by definition, gives us a function $f : X \rightarrow Y$ such that $\prod_{x:X} \phi(x, f(x))$. In some cases, we wish to express the weaker “for every $x : X$, there exists some $y : Y$ such that $\phi(x, y)$ ” without necessarily having an assignment of x ’s to y ’s. A good example of this is when we define directed families later (see Definition 7). This is achieved through the propositional truncation.

Given a type $X : \mathcal{U}$, we postulate that we have a proposition $\|X\| : \mathcal{U}$ with a function $|-| : X \rightarrow \|X\|$.

137 $X \rightarrow \|X\|$ such that for every proposition $P : \mathcal{V}$ in any universe \mathcal{V} , every function $f : X \rightarrow P$
 138 factors (necessarily uniquely, by function extensionality) through $|-|$. Diagrammatically,

$$\begin{array}{ccc}
 X & \xrightarrow{f} & P \\
 \searrow & & \nearrow \\
 & \text{||}X\text{||} &
 \end{array}$$

139

140 Existential quantification $\exists_{x:X} Y(x)$ is given by $\|\Sigma_{x:X} Y(x)\|$. One should note that if we
 141 have $\exists_{x:X} Y(x)$ and we are trying to prove some proposition P , then we may assume that we
 142 have $x : X$ and $y : Y(x)$ when constructing our inhabitant of P . Similarly, we can define
 143 disjunction as $P \vee Q \equiv \|P + Q\|$.

144 3 Impredicativity

145 We now explain what we mean by (im)predicativity in univalent foundations.

146 ► **Definition 1** (Has size, `has-size` in [14]). A type $X : \mathcal{U}$ is said to have size \mathcal{V} for some
 147 universe \mathcal{V} when we have $Y : \mathcal{V}$ that is equivalent to X , i.e. X `has-size` $\mathcal{V} \equiv \sum_{Y:\mathcal{V}} Y \simeq X$.

148 Here, the symbol \simeq refers to Voevodsky's notion of equivalence [14, 43]. The type X `has-size` \mathcal{V}
 149 is a proposition if and only if the univalence axiom holds [14].

150 ► **Definition 2** (Type of propositions $\Omega_{\mathcal{U}}$). The type of propositions in a universe \mathcal{U} is
 151 $\Omega_{\mathcal{U}} \equiv \sum_{P:\mathcal{U}} \text{is-prop}(P) : \mathcal{U}^+$.

152 Observe that $\Omega_{\mathcal{U}}$ itself lives in the successor universe \mathcal{U}^+ . We often think of the types
 153 in some fixed universe \mathcal{U} as *small* and accordingly we say that $\Omega_{\mathcal{U}}$ is *large*. Similarly, the
 154 powerset of a type $X : \mathcal{U}$ is large. Given our predicative setup, we must pay attention to
 155 universes when considering powersets.

156 ► **Definition 3** (\mathcal{V} -powerset $\mathcal{P}_{\mathcal{V}}(X)$, \mathcal{V} -subsets). Let \mathcal{V} be a universe and $X : \mathcal{U}$ type. We define
 157 the \mathcal{V} -powerset $\mathcal{P}_{\mathcal{V}}(X)$ as $X \rightarrow \Omega_{\mathcal{V}} : \mathcal{V}^+ \sqcup \mathcal{U}$. Its elements are called \mathcal{V} -subsets of X .

158 ► **Definition 4** (\in, \subseteq). Let x be an element of a type X and let A be an element of $\mathcal{P}_{\mathcal{V}}(X)$.
 159 We write $x \in A$ for the type $\text{pr}_1(A(x))$. Given two \mathcal{V} -subsets A and B of X , we write $A \subseteq B$
 160 for $\prod_{x:X} (x \in A \rightarrow x \in B)$.

161 Functional and propositional extensionality imply that $A = B \iff A \subseteq B$ and $B \subseteq A$.

162 ► **Definition 5** (Total type $\mathbb{T}(A)$). Given a \mathcal{V} -subset A of a type X , we write $\mathbb{T}(A)$ for the
 163 total type $\sum_{x:X} x \in A$.

164 One could ask for a *resizing axiom* asserting that $\Omega_{\mathcal{U}}$ has size \mathcal{U} , which we call the
 165 *propositional impredicativity* of \mathcal{U} . A closely related axiom is *propositional resizing*, which
 166 asserts that every proposition $P : \mathcal{U}^+$ has size \mathcal{U} . Without the addition of such resizing
 167 axioms, the type theory is said to be *predicative*. As an example of the use of impredicativity
 168 in mathematics, we mention that the powerset has unions of arbitrary subsets if and only if
 169 propositional resizing holds [14, `existence-of-unions-gives-PR`].

170 We mention that the resizing axioms are actually theorems when classical logic is assumed.
 171 This is because if $P \vee \neg P$ holds for every proposition in $\mathcal{P} : \mathcal{U}$, then the only propositions
 172 (up to equivalence) are $\mathbf{0}_{\mathcal{U}}$ and $\mathbf{1}_{\mathcal{U}}$, which have equivalent copies in \mathcal{U}_0 , and $\Omega_{\mathcal{U}}$ is equivalent
 173 to a type $\mathbf{2}_{\mathcal{U}} : \mathcal{U}$ with exactly two elements. The existence of a computational interpreta-
 174 tion of propositional impredicativity axioms for univalent foundations is an open problem,
 175 however [42, 41].

4 Basic Domain Theory

Our basic ingredient is the notion of *directed complete poset* (dcpo). In set-theoretic foundations, a dcpo can be defined to be a poset that has least upper bounds of all directed families. A naive translation of this to our foundation would be to proceed as follows. Define a poset in a universe \mathcal{U} to be a type $P : \mathcal{U}$ with a reflexive, transitive and antisymmetric relation $\sqsubseteq : P \times P \rightarrow \mathcal{U}$. According to the univalent point of view, we also require that the type P is a *set* and the values $p \sqsubseteq q$ of the order relation are *subsingletons*. Then we could say that the poset (P, \sqsubseteq) is *directed complete* if every directed family $I \rightarrow X$ with indexing type $I : \mathcal{U}$ has a least upper bound. The problem with this definition is that there are no interesting examples in our constructive and predicative setting. For instance, assume that the poset $\mathbf{2}$ with two elements $0 \sqsubseteq 1$ is directed complete, and consider a proposition $A : \mathcal{U}$ and the directed family $A + \mathbf{1} \rightarrow \mathbf{2}$ that maps the left component to 1 and the right component to 0. By case analysis on its hypothetical supremum, we conclude that the negation of A is decidable. This amounts to weak excluded middle, which is known to be equivalent to De Morgan's Law, and doesn't belong to the realm of constructive mathematics. To try to get an example, we may move to the poset Ω_0 of propositions in the universe \mathcal{U}_0 , ordered by implication. This poset does have all suprema of families $I \rightarrow \Omega_0$ indexed by types I in the *first universe* \mathcal{U}_0 , given by existential quantification. But if we consider a directed family $I \rightarrow \Omega_0$ with I in the *same universe* as Ω_0 lives, namely the *second universe* \mathcal{U}_1 , existential quantification gives a proposition in the *third universe* \mathcal{U}_2 and so doesn't give its supremum. In this example, we get a poset such that

- (i) the carrier lives in the universe \mathcal{U}_1 ,
- (ii) the order has truth values in the universe \mathcal{U}_0 , and
- (iii) suprema of directed families indexed by types in \mathcal{U}_0 exist.

Regarding a poset as a category in the usual way, we have a large, but locally small, category with small filtered colimits (suprema). This is typical of all the examples we have considered so far in practice, such as the dcpos in the Scott model of PCF and Scott's D_∞ model of the untyped λ -calculus. We may say that the predicativity restriction increases the universe usage by one. However, for the sake of generality, we formulate our definition of dcpo with the following universe conventions:

- (i) the carrier lives in a universe \mathcal{U} ,
- (ii) the order has truth values in a universe \mathcal{T} , and
- (iii) suprema of directed families indexed by types in a universe \mathcal{V} exist.

So our notion of dcpo has three universe parameters $\mathcal{U}, \mathcal{V}, \mathcal{T}$. We will say that the dcpo is *locally small* when \mathcal{T} is not necessarily the same as \mathcal{V} , but the order has truth values of size \mathcal{V} . Most of the time we mention \mathcal{V} explicitly and leave \mathcal{U} and \mathcal{T} to be understood from the context.

► **Definition 6** (Poset). *A poset (P, \sqsubseteq) is a set $P : \mathcal{U}$ together with a proposition-valued binary relation $\sqsubseteq : P \rightarrow P \rightarrow \mathcal{T}$ satisfying:*

- (i) reflexivity: $\prod_{p:P} p \sqsubseteq p$;
- (ii) antisymmetry: $\prod_{p,q:P} p \sqsubseteq q \rightarrow q \sqsubseteq p \rightarrow p = q$;
- (iii) transitivity: $\prod_{p,q,r:P} p \sqsubseteq q \rightarrow q \sqsubseteq r \rightarrow p \sqsubseteq r$.

► **Definition 7** (Directed family). *Let (P, \sqsubseteq) be a poset and I any type. A family $\alpha : I \rightarrow P$ is directed if it is inhabited (i.e. $\|I\|$ is pointed) and $\prod_{i,j:I} \exists k:I \alpha_i \sqsubseteq \alpha_k \times \alpha_j \sqsubseteq \alpha_k$.*

► **Definition 8** (\mathcal{V} -directed complete poset, \mathcal{V} -dcpo). *Let \mathcal{V} be a type universe. A \mathcal{V} -directed complete poset (or \mathcal{V} -dcpo, for short) is a poset (P, \sqsubseteq) such that every directed family $I \rightarrow P$ with $I : \mathcal{V}$ has a supremum in P .*

We will sometimes leave the universe \mathcal{V} implicit, and simply speak of “a dcpo”. On other occasions, we need to carefully keep track of universe levels. To this end, we make the following definition.

► **Definition 9** (\mathcal{V} -DCPO $_{\mathcal{U},\mathcal{T}}$). Let \mathcal{V} , \mathcal{U} and \mathcal{T} be universes. We write \mathcal{V} -DCPO $_{\mathcal{U},\mathcal{T}}$ for the type of \mathcal{V} -dcpos with carrier in \mathcal{U} and order taking values in \mathcal{T} .

► **Definition 10** (Pointed dcpo). A dcpo D is pointed if it has a least element, which we will denote by \perp_D , or simply \perp .

► **Definition 11** (Locally small). A \mathcal{V} -dcpo D is locally small if we have $\sqsubseteq_{\text{small}} : D \rightarrow D \rightarrow \mathcal{V}$ such that $\prod_{x,y:D} (x \sqsubseteq_{\text{small}} y) \simeq (x \sqsubseteq_D y)$.

► **Example 12** (Powersets as pointed dcpos). Powersets give examples of pointed dcpos. The subset inclusion \subseteq makes $\mathcal{P}_{\mathcal{V}}(X)$ into a poset and given a (not necessarily directed) family $A_{(-)} : I \rightarrow \mathcal{P}_{\mathcal{V}}(X)$ with $I : \mathcal{V}$, we may consider its supremum in $\mathcal{P}_{\mathcal{V}}(X)$ as given by $\lambda x. \exists i. I. x \in A_i$. Note that $(\exists i. I. x \in A_i) : \mathcal{V}$ for every $x : X$, so this is well-defined. Finally, $\mathcal{P}_{\mathcal{V}}$ has a least element, the empty set: $\lambda x. \mathbf{0}_{\mathcal{V}}$. Thus, $\mathcal{P}_{\mathcal{V}}(X) : \mathcal{V}$ -DCPO $_{\mathcal{V}+\sqcup\mathcal{U},\mathcal{V}+\sqcup\mathcal{U}}$. When $\mathcal{V} \equiv \mathcal{U}$ (as in Example 59), we get the simpler, locally small $\mathcal{P}_{\mathcal{U}}(X) : \mathcal{U}$ -DCPO $_{\mathcal{U}+\mathcal{U}}$. ◀

Fix two \mathcal{V} -dcpos D and E .

► **Definition 13** (Continuous function). A function $f : D \rightarrow E$ is (Scott) continuous if it preserves directed suprema, i.e. if $I : \mathcal{V}$ and $\alpha : I \rightarrow D$ is directed, then $f(\bigsqcup \alpha)$ is the supremum in E of the family $f \circ \alpha$.

► **Lemma 14**. If $f : D \rightarrow E$ is continuous, then it is monotone, i.e. $x \sqsubseteq_D y$ implies $f(x) \sqsubseteq_E f(y)$.

Proof. Given $x, y : D$ with $x \sqsubseteq y$, consider the directed family $\mathbf{1} + \mathbf{1} \rightarrow D$ defined as $\text{inl}(\star) \mapsto x$ and $\text{inr}(\star) \mapsto y$. Its supremum is y and f must preserve it, so $f(x) \sqsubseteq f(y)$. ◀

► **Lemma 15**. If $f : D \rightarrow E$ is continuous and $\alpha : I \rightarrow D$ is directed, then so is $f \circ \alpha$.

Proof. Using Lemma 14. ◀

► **Definition 16** (Strict function). Suppose that D and E are pointed. A continuous function $f : D \rightarrow E$ is strict if $f(\perp_D) = \perp_E$.

4.1 Lifting

► **Construction 17** ($\mathcal{L}_{\mathcal{V}}(X)$, η_X , cf. [8, 13]). Let $X : \mathcal{U}$ be a set. For any universe \mathcal{V} , we construct a pointed \mathcal{V} -dcpo $\mathcal{L}_{\mathcal{V}}(X) : \mathcal{V}$ -DCPO $_{\mathcal{V}+\sqcup\mathcal{U},\mathcal{V}+\sqcup\mathcal{U}}$, known as the *lifting* of X . Its carrier is given by the type $\sum_{P:\mathcal{V}} \text{is-prop}(P) \times (P \rightarrow X)$ of *partial elements* of X .

Given a partial element $(P, i, \varphi) : \mathcal{L}_{\mathcal{V}}(X)$, we write $(P, i, \varphi) \downarrow$ for P and say that the partial element is defined if P holds. Moreover, we often leave the second component implicit, writing (P, φ) for (P, i, φ) .

The order is given by $l \sqsubseteq_{\mathcal{L}_{\mathcal{V}}(X)} m \equiv (l \downarrow \rightarrow l = m)$, and it has a least element given by $(\mathbf{0}, \mathbf{0}\text{-is-prop, unique-from-}\mathbf{0})$ where $\mathbf{0}\text{-is-prop}$ is a witness that the empty type is a proposition and $\text{unique-from-}\mathbf{0}$ is the unique map from the empty type.

Given a directed family $(Q_{(-)}, \varphi_{(-)}) : I \rightarrow \mathcal{L}_{\mathcal{V}}(X)$, its supremum is given by $(\exists_{i:I} Q_i, \psi)$, where ψ is such that

$$\begin{array}{ccc} \sum_{i:I} Q_i & \xrightarrow{(i,q) \mapsto \varphi_i(q)} & D \\ & \searrow \quad \nearrow \psi & \\ & \exists_{i:I} Q_i & \end{array}$$

commutes. (This is possible, because the top map is weakly constant (i.e. any of its values are equal) and D is a set [23, Theorem 5.4].)

Finally, we write $\eta_X : X \rightarrow \mathcal{L}_{\mathcal{V}}(X)$ for the embedding $x \mapsto (\mathbf{1}, \mathbf{1}\text{-is-prop}, \lambda u.x)$. \lrcorner

Note that we require X to be a set, so that $\mathcal{L}_{\mathcal{V}}(X)$ is a poset, rather than an ∞ -category. In practice, we often have $\mathcal{V} \equiv \mathcal{U}$ (see for instance Example 58, Section 5.2, or the Scott model of PCF [8]), but we develop the theory for the more general case. We can describe the order on $\mathcal{L}_{\mathcal{V}}(X)$ more explicitly, as follows.

► **Lemma 18.** *If we have elements (P, φ) and (Q, ψ) of $\mathcal{L}_{\mathcal{V}}(X)$, then $(P, \varphi) \sqsubseteq (Q, \psi)$ holds if and only if we have $f : P \rightarrow Q$ such that $\prod_{p:P} \varphi(p) = \psi(f(p))$.*

Observe that this exhibits $\mathcal{L}_{\mathcal{V}}(X)$ as locally small. We will show that $\mathcal{L}_{\mathcal{V}}(X)$ is the free pointed \mathcal{V} -dcpo on a set X , but to do that, we first need a lemma.

► **Lemma 19.** *Let D be a pointed \mathcal{V} -dcpo. Then D has suprema of families indexed by propositions in \mathcal{V} , i.e. if $P : \mathcal{V}$ is a proposition, then any $\alpha : P \rightarrow D$ has a supremum $\bigvee \alpha$.*

Moreover, if E is a (not necessarily pointed) \mathcal{V} -dcpo and $f : D \rightarrow E$ is continuous, then $f(\bigvee \alpha)$ is the supremum of the family $f \circ \alpha$.

Proof. Let D be a pointed \mathcal{V} -dcpo, $P : \mathcal{V}$ a proposition and $\alpha : P \rightarrow D$ a function. Now define $\beta : \mathbf{1}_{\mathcal{V}} + P \rightarrow D$ by $\text{inl}(\star) \mapsto \perp_D$ and $\text{inr}(p) \mapsto \alpha(p)$. Then, β is easily seen to be directed and so it has a well-defined supremum in D , which is also the supremum of α . The second claim follows from the fact that β is directed, so continuous maps must preserve its supremum. \blacktriangleleft

► **Lemma 20.** *Let $X : \mathcal{U}$ be a set and let (P, φ) be an arbitrary element of $\mathcal{L}_{\mathcal{V}}(X)$. Then $(P, \varphi) = \bigvee_{p:P} \eta_X(\varphi(p))$.*

► **Theorem 21.** *The lifting $\mathcal{L}_{\mathcal{V}}(X)$ gives the free \mathcal{V} -dcpo on a set X . Put precisely, if $X : \mathcal{U}$ is a set, then for every \mathcal{V} -dcpo $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and function $f : X \rightarrow D$, there is a unique continuous function $\bar{f} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ such that*

$$\begin{array}{ccc} X & \xrightarrow{f} & D \\ & \searrow \eta_X \quad \nearrow \bar{f} & \\ & \mathcal{L}_{\mathcal{V}}(X) & \end{array}$$

commutes.

Proof. We define $\bar{f} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ by $(P, \varphi) \mapsto \bigvee_{p:P} f(\varphi(p))$, which is well-defined by Lemma 19 and easily seen to be continuous. For uniqueness, suppose that we have $g : \mathcal{L}_{\mathcal{V}}(X) \rightarrow D$ continuous such that $g \circ \eta_X = f$. Let (P, φ) be an arbitrary element

of $\mathcal{L}_{\mathcal{V}}(X)$. Using Lemma 20, we have:

$$\begin{aligned}
 g(P, \varphi) &= g\left(\bigvee_{p:P} \eta_X(\varphi(p))\right) \\
 &= \bigvee_{p:P} g(\eta_X(\varphi(p))) && \text{(by Lemma 19 and continuity of } g\text{)} \\
 &= \bigvee_{p:P} f(\phi(p)) && \text{(by assumption on } g\text{)} \\
 &= \bar{f}(P, \varphi) && \text{(by definition),}
 \end{aligned}$$

as desired. \blacktriangleleft

There is yet another way in which the lifting is a free construction, cf. [8, Section 4.3]. What is noteworthy about this is that freely adding subsingleton suprema automatically gives all directed suprema.

► **Definition 22** ($\Omega_{\mathcal{V}}$ -complete). A poset (P, \sqsubseteq) is $\Omega_{\mathcal{V}}$ -complete if it has suprema for all families indexed by a proposition in \mathcal{V} .

► **Theorem 23.** The lifting $\mathcal{L}_{\mathcal{V}}(X)$ gives the free $\Omega_{\mathcal{V}}$ -complete poset on a set X . Put precisely, if $X : \mathcal{U}$ is a set, then for every $\Omega_{\mathcal{V}}$ -complete poset (P, \sqsubseteq) with $P : \mathcal{U}'$ and \sqsubseteq taking values in \mathcal{T}' and function $f : X \rightarrow P$, there exists a unique monotone $\bar{f} : \mathcal{L}_{\mathcal{V}}(X) \rightarrow P$ preserving all suprema indexed by propositions in \mathcal{V} , such that

$$\begin{array}{ccc}
 X & \xrightarrow{f} & P \\
 \eta_X \searrow & & \nearrow \bar{f} \\
 & \mathcal{L}_{\mathcal{V}}(X) &
 \end{array}$$

commutes.

Proof. Similar to the proof of Theorem 21; also see [8, Proof of Theorem 4.16]. \blacktriangleleft

Finally, a variation of Construction 17 freely adds a least element to a dcpo.

► **Construction 24** ($\mathcal{L}'_{\mathcal{V}}(D)$). Let $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ be a \mathcal{V} -dcpo. We construct a pointed \mathcal{V} -dcpo $\mathcal{L}'_{\mathcal{V}}(D) : \mathcal{V}\text{-DCPO}_{\mathcal{V}+\sqcup\mathcal{U}, \mathcal{V}\sqcup\mathcal{T}}$. Its carrier is given by the type $\sum_{P:\mathcal{V}} \text{is-prop}(P) \times (P \rightarrow D)$. The order is given by $(P, \varphi) \sqsubseteq_{\mathcal{L}'_{\mathcal{V}}(D)} (Q, \psi) \equiv \sum_{f:P \rightarrow Q} \prod_{p:P} \varphi(p) = \psi(f(p))$ and has a least element $(\mathbf{0}, \mathbf{0}\text{-is-prop, unique-from-}\mathbf{0})$.

Now let $\alpha \equiv (Q(-), \varphi(-)) : I \rightarrow \mathcal{L}'_{\mathcal{V}}(D)$ be a directed family. Consider $\Phi : (\sum_{i:I} Q_i) \rightarrow D$ given by $(i, q) \mapsto \varphi_i(q)$. The supremum of α is given by $(\exists_{i:I} Q_i, \psi)$, where ψ takes a witness that $\sum_{i:I} Q_i$ is inhabited to the directed (for which we needed $\exists_{i:I} Q_i$) supremum $\bigsqcup \Phi$ in D .

Finally, we write $\eta'_D : D \rightarrow \mathcal{L}'_{\mathcal{V}}(D)$ for the continuous map $x \mapsto (\mathbf{1}, \mathbf{1}\text{-is-prop, } \lambda u. x)$. \sqcup

► **Theorem 25.** The construction $\mathcal{L}'_{\mathcal{V}}(D)$ gives the free pointed \mathcal{V} -dcpo on a \mathcal{V} -dcpo D . Put precisely, if $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ is a \mathcal{V} -dcpo, then for every pointed \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ and continuous function $f : D \rightarrow E$, there is a unique strict continuous function $\bar{f} : \mathcal{L}'_{\mathcal{V}}(D) \rightarrow E$ such that

$$\begin{array}{ccc}
 D & \xrightarrow{f} & E \\
 \eta'_D \searrow & & \nearrow \bar{f} \\
 & \mathcal{L}'_{\mathcal{V}}(D) &
 \end{array}$$

commutes.

327 **Proof.** Similar to the proof of Theorem 21. ◀

328 4.2 Exponentials

329 ► **Construction 26** (E^D). Let D and E be two \mathcal{V} -dcpos. We construct another \mathcal{V} -dcpo E^D
 330 as follows. Its carrier is given by the type of continuous functions from D to E .

331 These functions are ordered pointwise, i.e. if $f, g : D \rightarrow E$, then

$$332 \quad f \sqsubseteq_{E^D} g \equiv \prod_{x:D} f(x) \sqsubseteq_E g(x).$$

333 Accordingly, directed suprema are also given pointwise. Explicitly, let $\alpha : I \rightarrow E^D$ be a
 334 directed family. For every $x : D$, we have the family $\alpha_x : I \rightarrow E$ given by $i \mapsto \alpha_i(x)$. This is
 335 a directed family in E and so we have a well-defined supremum $\bigsqcup \alpha_x : E$ for every $x : D$.
 336 The supremum of α is then given by $x \mapsto \bigsqcup \alpha_x$, where one should check that this assignment
 337 is indeed continuous.

338 Finally, if E is pointed, then so is E^D , because, in that case, the function $x \mapsto \perp_E$ is the
 339 least continuous function from D to E . ⊥

340 ► **Remark 27.** In general, the universe levels of E^D can be quite large and complicated. For if
 341 $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$, then $E^D : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+ \sqcup \mathcal{U} \sqcup \mathcal{T} \sqcup \mathcal{U}' \sqcup \mathcal{T}', \mathcal{U} \sqcup \mathcal{T}'}$. Even if
 342 $\mathcal{V} = \mathcal{U} \equiv \mathcal{T} \equiv \mathcal{U}' \equiv \mathcal{T}'$, the carrier of E^D still lives in the “large” universe \mathcal{V}^+ . (Actually,
 343 this scenario cannot happen non-trivially in a predicative setting, since non-trivial dcpos
 344 cannot be “small” [9].) Even so, as observed in [8], if we take $\mathcal{U} \equiv \mathcal{T} \equiv \mathcal{U}' \equiv \mathcal{T}' \equiv \mathcal{U}_1$ and
 345 $\mathcal{V} \equiv \mathcal{U}_0$, then D, E, E^D are all elements of $\mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$.

346 5 Scott’s D_∞

347 We now construct, predicatively, Scott’s famous pointed dcpo D_∞ which is isomorphic to its
 348 own function space $D_\infty^{D_\infty}$ (Theorem 39). We follow Scott’s original paper [38] rather closely,
 349 but with two differences. Firstly, we explicitly keep track of the universe levels to make sure
 350 that our constructions go through predicatively. Secondly, [38] describes sequential (co)limits,
 351 while we study the more general directed (co)limits (Section 5.1) and then specialize to
 352 sequential (co)limits later (Section 5.2).

353 5.1 Limits and Colimits

354 ► **Definition 28** (Deflation). Let D be a dcpo. An endofunction $f : D \rightarrow D$ is a deflation if
 355 $f(x) \sqsubseteq x$ for all $x : D$.

356 ► **Definition 29** (Embedding-projection pair). Let D and E be two dcpos. An embedding-
 357 projection pair from D to E consists of two continuous functions $\varepsilon : D \rightarrow E$ (the embedding)
 358 and $\pi : E \rightarrow D$ (the projection) such that:

- 359 (i) ε is a section of π ;
- 360 (ii) $\varepsilon \circ \pi$ is a deflation.

361 For the remainder of this section, fix the following setup. Let $\mathcal{V}, \mathcal{U}, \mathcal{T}$ and \mathcal{W} be type
 362 universes. Let (I, \sqsubseteq) be a directed preorder with $I : \mathcal{V}$ and \sqsubseteq taken values in \mathcal{W} . Suppose
 363 that we have:

- 364 (i) for every $i : I$, a \mathcal{V} -dcpo $D_i : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$;
- 365 (ii) for every $i, j : I$ with $i \sqsubseteq j$, an embedding-projection pair $(\varepsilon_{i,j}, \pi_{i,j})$ from D_i to D_j ;

such that

- (i) for every $i : I$, we have $\varepsilon_{i,i} = \pi_{i,i} = \text{id}$;
- (ii) for every $i, j, k : I$ with $i \sqsubseteq j \sqsubseteq k$, we have $\varepsilon_{i,k} \sim \varepsilon_{j,k} \circ \varepsilon_{i,j}$ and $\pi_{i,k} \sim \pi_{i,j} \circ \pi_{j,k}$.

► **Construction 30** (D_∞). Given the above inputs, we construct another \mathcal{V} -dcpo $D_\infty : \mathcal{V}\text{-DCPO}_{\mathcal{U} \sqcup \mathcal{V} \sqcup \mathcal{W}, \mathcal{U} \sqcup \mathcal{T}}$ as follows. Its carrier is given by the type:

$$\sigma : \prod_{i:I} \sum_{D_i} \prod_{j:I, i \sqsubseteq j} \pi_{i,j}(\sigma_j) = \sigma_i.$$

These functions are ordered pointwise, i.e. if $\sigma, \tau : I \rightarrow D_i$, then

$$\sigma \sqsubseteq_{D_\infty} \tau \equiv \prod_{i:I} \sigma_i \sqsubseteq_{D_i} \tau_i.$$

Accordingly, directed suprema are also given pointwise. Explicitly, let $\alpha : A \rightarrow D_\infty$ be a directed family. For every $i : I$, we have the family $A \rightarrow D_i$ given by $a \mapsto (\alpha(a))_i$, and denoted by α_i . One can show that α_i is directed and so we have a well-defined supremum $\bigsqcup \alpha_i : D_i$ for every $i : I$. The supremum of α is then given by the function $i : I \mapsto \bigsqcup \alpha_i$, where one should check that $\pi_{i,j}(\bigsqcup \alpha_j) = \bigsqcup \alpha_i$ holds whenever $i \sqsubseteq j$. \lrcorner

► **Remark 31.** We allow for general universe levels here, which is why D_∞ lives in the relatively complicated universe $\mathcal{U} \sqcup \mathcal{V} \sqcup \mathcal{W}$. In concrete examples, such as in Section 5.2, the situation simplifies to $\mathcal{V} = \mathcal{W} = \mathcal{U}_0$ and $\mathcal{U} = \mathcal{T} = \mathcal{U}_1$.

► **Construction 32** ($\pi_{i,\infty}$). For every $i : I$, we have a continuous function $\pi_{i,\infty} : D_\infty \rightarrow D_i$, given by $\sigma \mapsto \sigma_i$. \lrcorner

► **Construction 33** ($\varepsilon_{i,\infty}$). For every $i, j : I$, consider the function

$$\kappa : D_i \rightarrow \left(\sum_{k:I} i \sqsubseteq k \times j \sqsubseteq k \right) \rightarrow D_j$$

$$\kappa_x(k, l_i, l_j) = \pi_{i,j}(\varepsilon_{i,k}(x)).$$

Using directedness of (I, \sqsubseteq) , we can show that for every $x : D_i$ the map κ_x is weakly constant (i.e. all its values are equal). Therefore, we can apply [23, Theorem 5.4] and factor κ_x through $\exists_{k:I} (i \sqsubseteq k \times j \sqsubseteq k)$. But (I, \sqsubseteq) is directed, so $\exists_{k:I} (i \sqsubseteq k \times j \sqsubseteq k)$ is a singleton. Thus, we obtain a function $\rho_{i,j} : D_i \rightarrow D_j$ such that: if we are given $k : I$ with $(l_i, l_j) : i \sqsubseteq k \times j \sqsubseteq k$, then $\rho_{i,j}(x) = \kappa_x(k, l_i, l_j)$.

Finally, this allows us to construct for every $i : I$, a continuous function $\varepsilon_{i,\infty} : D_i \rightarrow D_\infty$ by mapping $x : D_i$ to the function $\lambda(j : I). \rho_{i,j}(x)$. \lrcorner

► **Theorem 34.** For every $i : I$, the pair $(\varepsilon_{i,\infty}, \pi_{i,\infty})$ is an embedding-projection pair.

► **Lemma 35.** Let $i, j : I$ such that $i \sqsubseteq j$. Then $\pi_{i,j} \circ \pi_{j,\infty} \sim \pi_i$, and $\varepsilon_{j,\infty} \circ \varepsilon_{i,j} \sim \varepsilon_{i,\infty}$.

► **Theorem 36.** The dcpo D_∞ with the maps $(\pi_{i,\infty})_{i:I}$ is the limit of $\left((D_i)_{i:I}, (\pi_{i,j})_{i \sqsubseteq j} \right)$. That is, given

- (i) a \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$,
 - (ii) continuous functions $f_i : E \rightarrow D_i$ for every $i : I$,
- such that $\pi_{i,j} \circ f_j \sim f_i$ whenever $i \sqsubseteq j$, we have a continuous function $f_\infty : E \rightarrow D_\infty$ such that $\pi_{i,\infty} \circ f_\infty \sim f_i$ for every $i : I$. Moreover, f_∞ is the unique such continuous function.
- The function f_∞ is given by mapping $y : E$ to the function $\lambda(i : I). f_i(y)$.

► **Theorem 37.** *The dcpo D_∞ with the maps $(\varepsilon_{i,\infty})_{i:I}$ is the colimit of $((D_i)_{i:I}, (\varepsilon_{i,j})_{i \sqsubseteq j})$. That is, given*

- (i) *a \mathcal{V} -dcpo $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$,*
- (ii) *continuous functions $g_i : D_i \rightarrow E$ for every $i : I$,*

such that $g_j \circ \varepsilon_{i,j} \sim g_i$ whenever $i \sqsubseteq j$, we have a continuous function $g_\infty : D_\infty \rightarrow E$ such that $g_\infty \circ \varepsilon_{i,\infty} \sim g_i$ for every $i : I$. Moreover, g_∞ is the unique such continuous function.

The function g_∞ is given by $\sigma \mapsto \bigsqcup_{i:I} g_i(\sigma_i)$, where one should check that the family $i \mapsto g_i(\sigma_i)$ is indeed directed.

Proof. For uniqueness, it is useful to know that an element $\sigma : D_\infty$ can be expressed as the directed supremum $\bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i)$. The rest can be checked directly. ◀

It should be noted that in both universal properties E can have its carrier in any universe \mathcal{U}' and its order taking values in any universe \mathcal{T}' , even though we required all D_i to have their carriers and orders in two fixed universes \mathcal{U} and \mathcal{T} , respectively.

5.2 Scott's Example Using Self-exponentiation

We now show that we can construct Scott's D_∞ [38] predicatively. Formulated precisely, we construct a pointed $D_\infty : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ such that D_∞ is isomorphic to its self-exponential $D_\infty^{D_\infty}$.

We employ the machinery from Section 5.1. Following [38, pp. 126–127], we inductively define pointed dcpos $D_n : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ for every natural number n :

- (i) $D_0 \equiv \mathcal{L}_{\mathcal{U}_0}(\mathbf{1}_{\mathcal{U}_0})$;
- (ii) $D_{n+1} \equiv D_n^{D_n}$.

Next, we inductively define embedding-projection pairs (ε_n, π_n) from D_n to D_{n+1} :

- (i) $\varepsilon_0 : D_0 \rightarrow D_1$ is given by mapping $x : D_0$ to the continuous function that is constantly x ;
 $\pi_0 : D_1 \rightarrow D_0$ is given by evaluating a continuous function $f : D_0 \rightarrow D_0$ at \perp ;
- (ii) $\varepsilon_{n+1} : D_{n+1} \rightarrow D_{n+2}$ takes a continuous function $f : D_n \rightarrow D_n$ to the continuous composite $D_{n+1} \xrightarrow{\pi_n} D_n \xrightarrow{f} D_n \xrightarrow{\varepsilon_n} D_{n+1}$;
 $\pi_{n+1} : D_{n+2} \rightarrow D_{n+1}$ takes a continuous function $f : D_{n+1} \rightarrow D_{n+1}$ to the continuous composite $D_n \xrightarrow{\varepsilon_n} D_{n+1} \xrightarrow{f} D_{n+1} \xrightarrow{\pi_n} D_n$.

In order to apply the machinery from Section 5.1, we will need embedding-projection pairs $(\varepsilon_{n,m}, \pi_{n,m})$ from D_n to D_m whenever $n \leq m$. Let n and m be natural numbers with $n \leq m$ and let k be the natural number $m - n$. We define the pairs by induction on k :

- (i) if $k = 0$, then we set $\varepsilon_{n,n} = \pi_{n,n} = \text{id}$;
- (ii) if $k = l + 1$, then $\varepsilon_{n,m} = \varepsilon_{n+1} \circ \varepsilon_{n,n+l}$ and $\pi_{n,m} = \pi_{n,n+l} \circ \pi_{n+l}$.

So, Constructions 30, 32 and 33 give us $D_\infty : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_1}$ with embedding-projection pairs $(\varepsilon_{n,\infty}, \pi_{n,\infty})$ from D_n to D_∞ for every natural number n .

► **Lemma 38.** *Let n be a natural number. The function $\pi_n : D_{n+1} \rightarrow D_n$ is strict. Hence, so is $\pi_{n,m}$ whenever $n \leq m$.*

Proof. The first statement is proved by induction on n . The second by induction on k with $k \equiv m - n$. ◀

► **Theorem 39.** *The dcpo D_∞ is pointed and isomorphic to $D_\infty^{D_\infty}$.*

Proof. Since every D_n is pointed, we can consider the function $\sigma : \prod_{n:\mathbf{N}} D_n$ given by $\sigma(n) \equiv \perp_{D_n}$. Then σ is an element of D_∞ by Lemma 38 and it is the least, so D_∞ is indeed pointed.

We start by constructing a continuous function $\varepsilon'_\infty : D_\infty \rightarrow D_\infty^{D_\infty}$. By Theorem 37, it suffices to define continuous functions $\varepsilon'_n : D_n \rightarrow D_\infty^{D_\infty}$ for every natural number n such that $\varepsilon'_m \circ \varepsilon_{n,m} \sim \varepsilon'_n$ whenever $n \leq m$. We do so as follows:

- (i) $\varepsilon'_{n+1} : D_{n+1} \rightarrow D_\infty^{D_\infty}$ is given by mapping a continuous function $f : D_n \rightarrow D_n$ to the continuous composite $D_\infty \xrightarrow{\pi_{n,\infty}} D_n \xrightarrow{f} D_n \xrightarrow{\varepsilon_{n,\infty}} D_\infty$;
- (ii) $\varepsilon'_0 : D_0 \rightarrow D_\infty^{D_\infty}$ is defined as the continuous composite $D_0 \xrightarrow{\varepsilon_0} D_1 \xrightarrow{\varepsilon'_1} D_\infty$.

Next, we construct a continuous function $\pi'_\infty : D_\infty^{D_\infty} \rightarrow D_\infty$. By Theorem 36, it suffices to define continuous functions $\pi'_n : D_n \rightarrow D_\infty^{D_\infty}$ for every natural number n such that $\pi_{n,m} \circ \pi'_m \sim \pi'_n$ whenever $n \leq m$. We do so as follows:

- (i) $\pi'_{n+1} : D_\infty^{D_\infty} \rightarrow D_{n+1} \equiv D_n^{D_n}$ is given by mapping a continuous function $f : D_\infty \rightarrow D_\infty$ to the continuous composite $D_n \xrightarrow{\varepsilon_{n,\infty}} D_\infty \xrightarrow{f} D_\infty \xrightarrow{\pi_{n,\infty}} D_n$;
- (ii) $\pi'_0 : D_\infty^{D_\infty} \rightarrow D_0$ is defined as the continuous composite $D_\infty \xrightarrow{\pi'_1} D_1 \xrightarrow{\pi_0} D_0$.

It remains to prove that ε'_∞ and π'_∞ are inverses. To this end, it is convenient to have an alternative description of the maps ε'_∞ and π'_∞ .

$$\text{For every } \sigma : D_\infty, \text{ we have } \varepsilon'_\infty(\sigma) = \bigsqcup_{n:\mathbf{N}} \varepsilon'_{n+1}(\sigma_{n+1}). \quad (1)$$

$$\text{For every continuous } f : D_\infty \rightarrow D_\infty, \text{ we have } \pi'_\infty(f) = \bigsqcup_{n:\mathbf{N}} \varepsilon_{n+1,\infty}(\pi'_{n+1}(f)). \quad (2)$$

Using these equations we can prove that ε'_∞ and π'_∞ are inverses exactly as in [38, Proof of Theorem 4.4]. \blacktriangleleft

► **Remark 40.** Of course, Theorem 39 is only interesting in case $D_\infty \not\equiv \mathbf{1}$. Fortunately, D_∞ has (infinitely) many elements besides \perp_{D_∞} . For instance, we can consider $x_0 \equiv \eta(\star) : D_0$ and $\sigma_0 : \prod_{n:\mathbf{N}} D_n$ given by $\sigma_0(n) \equiv \varepsilon_{0,n}(x_0)$. Then, σ_0 is an element of D_∞ not equal to \perp_{D_∞} , because $x_0 \neq \perp_{D_0}$.

6 Continuous and Algebraic Dcpo

We next consider dcpo generated by certain elements called compact, or more generally generated by a certain way below relation, giving rise to algebraic and continuous domains.

6.1 The Way Below Relation

► **Definition 41** (Way below relation, $x \ll y$). *Let D be a \mathcal{V} -dcpo and $x, y : D$. We say that x is way below y , denoted by $x \ll y$, if for every $I : \mathcal{V}$ and directed family $\alpha : I \rightarrow D$, whenever we have $y \sqsubseteq \bigsqcup \alpha$, then there exists some element $i : I$ such that $x \sqsubseteq \alpha_i$ already. Symbolically,*

$$x \ll y \equiv \prod_{I:\mathcal{V}} \prod_{\alpha:I \rightarrow D} \left(\text{is-directed}(\alpha) \rightarrow y \sqsubseteq \bigsqcup \alpha \rightarrow \exists i:I. x \sqsubseteq \alpha_i \right).$$

► **Lemma 42.** *The way below relation enjoys the following properties.*

- (i) *It is proposition-valued.*

481 (ii) If $x \ll y$, then $x \sqsubseteq y$.

482 (iii) If $x \sqsubseteq y \ll v \sqsubseteq w$, then $x \ll w$.

483 (iv) It is antisymmetric.

484 (v) It is transitive.

485 ► **Lemma 43.** Let D be a dcpo. Then $x \sqsubseteq y$ implies $\prod_{z:D} (z \ll x \rightarrow z \ll y)$.

486 **Proof.** By Lemma 42(iii). ◀

487 ► **Definition 44** (Compact). Let D be a dcpo. An element $x : D$ is called compact if $x \ll x$.

488 ► **Example 45.** The least element of a pointed dcpo is always compact.

489 ► **Example 46** (Compact elements in $\mathcal{L}_{\mathcal{V}}(X)$). Let $X : \mathcal{U}$ be a set. An element $(P, \varphi) : \mathcal{L}_{\mathcal{V}}(X)$ is compact if and only if P is decidable.

491 ► **Definition 47** (Kuratowski finite). A type X is Kuratowski finite if there exists some
492 natural number $n : \mathbb{N}$ and a surjection $e : \text{Fin}(n) \rightarrow X$.

493 That is, X is Kuratowski finite if its elements can be finitely enumerated, possibly with
494 repetitions.

495 ► **Example 48** (Compact elements in $\mathcal{P}_{\mathcal{U}}(X)$). Let $X : \mathcal{U}$ be a set. An element $A : \mathcal{P}_{\mathcal{U}}(X)$ is
496 compact if and only if its total type $\mathbb{T}A$ is Kuratowski finite.

497 **Proof.** Write $\iota : \text{List}(X) \rightarrow \mathcal{P}_{\mathcal{U}}(X)$ for the map that regards a list on X as a subset of X .
498 The inductively generated type $\text{List}(X)$ of lists on X lives in the same universe \mathcal{U} as X .

499 Suppose that A is compact. We must show that $\mathbb{T}(A)$ is Kuratowski finite. Consider the
500 map $\alpha : \text{List}(\mathbb{T}(A)) \rightarrow \mathcal{P}_{\mathcal{U}(X)}$ which takes a list $[(x_0, p_0), \dots, (x_{n-1}, p_{n-1})]$ to $\iota([x_0, \dots, x_{n-1}])$.
501 Since the empty list is an element of $\text{List}(\mathbb{T}(A))$ and because we can concatenate lists, α is
502 directed. Moreover, $\text{List}(\mathbb{T}(A)) : \mathcal{U}$ and $A = \bigsqcup \alpha$ holds. Hence, by compactness, there exists
503 some $l \equiv [(x_0, p_0), \dots, (x_{n-1}, p_{n-1})] : \text{List}(\mathbb{T}(A))$ such that $A \subseteq \alpha(l)$ already. Hence, the
504 map $m : \text{Fin}(n) \mapsto (x_m, p_m) : \mathbb{T}(A)$ is a surjection, so $\mathbb{T}(A)$ is Kuratowski finite.

505 Conversely, suppose that A is a subset such that $\mathbb{T}(A)$ is Kuratowski finite. We must
506 prove that it is compact. Let $B_{(-)} : I \rightarrow \mathcal{P}_{\mathcal{U}}(X)$ be directed such that $A \subseteq \bigsqcup_{i:I} B_i$. Since
507 $\exists_{i:I} A \subseteq B_i$ is a proposition, we can use Kuratowski finiteness of $\mathbb{T}(A)$ to obtain a natural
508 number n and a surjection $e : \text{Fin}(n) \rightarrow \mathbb{T}(A)$. For each $m : \text{Fin}(n)$, find i_m such that
509 $e_m \in B_{i_m}$. By directedness of I , there exists $k : I$ such that $e_m \in B_k$ for every $m : \text{Fin}(n)$.
510 Hence, $\exists_{k:I} A \subseteq B_k$, as desired. ◀

511 6.2 Continuous Dcpo

512 Classically, a continuous dcpo is a dcpo where every element is the directed join of the set of
513 elements way below it [3]. Predicatively, we must be careful, because if x is an element of a
514 dcpo D , then $\sum_{y:D} y \ll x$ is typically large, so its directed join need not exist for size reasons.
515 Our solution is to use a predicative version of bases [1] that accounts for size issues. For the
516 special case of algebraic dcpo, our situation is the poset analogue of accessible categories [2].
517 Indeed, in category theory requiring smallness is common, even in impredicative settings, see
518 for instance [21], where continuous dcpo are generalized to continuous categories.

519 ► **Definition 49** (Basis, approximating family). A basis for \mathcal{V} -dcpo D is a function $\beta : B \rightarrow D$
520 with $B : \mathcal{V}$ such that for every $x : D$ there exists some $\alpha : I \rightarrow B$ with $I : \mathcal{V}$ such that

521 (i) $\beta \circ \alpha$ is directed and its supremum is x ;

522 (ii) $\beta(\alpha_i) \ll x$ for every $i : I$.

523 We summarise these requirements by saying that α is an approximating family for x .

524 Moreover, we require that \ll is small when restricted to the basis. That is, we have
 525 $\ll^B : B \rightarrow B \rightarrow \mathcal{V}$ such that $(\beta(b) \ll \beta(b')) \simeq (b \ll^B b')$ for every $b, b' : B$.

526 ► **Definition 50** (Continuous dcpo). A dcpo D is continuous if there exists some basis for it.

527 We postpone giving examples of continuous dcpos until we have developed the theory
 528 further, but the interested reader may look ahead to Examples 58, 59 and 82.

529 A useful property of bases is that it allows us to express the order fully in terms of the
 530 way below-relation, giving a converse to Lemma 43.

531 ► **Lemma 51.** Let D be a dcpo with basis $\beta : B \rightarrow D$. Then $x \sqsubseteq y$ holds if and only if
 532 $\prod_{b:B} (\beta(b) \ll x \rightarrow \beta(b) \ll y)$.

533 **Proof.** The left-to-right implication holds by Lemma 43. For the converse, suppose that we
 534 have $x, y : D$ such that for every $\prod_{b:B} (\beta(b) \ll x \rightarrow \beta(b) \ll y)$. Since $x \sqsubseteq y$ is a proposition,
 535 we can obtain $\alpha : I \rightarrow B$ such that $\beta \circ \alpha$ is directed and $\bigsqcup \beta \circ \alpha = x$ and $\beta(\alpha_i) \ll x$ for
 536 every $i : I$. It then suffices to show that $\bigsqcup \beta \circ \alpha \sqsubseteq y$. Since \bigsqcup gives the least upper bound,
 537 it is enough to prove that $\beta(\alpha_i) \sqsubseteq y$ for every $i : I$, but this holds by our hypothesis, our
 538 assumption that $\beta(\alpha_i) \ll x$ for every $i : I$, and Lemma 42(ii). ◀

539 ► **Lemma 52.** Let D be a \mathcal{V} -dcpo with a basis $\beta : B \rightarrow D$. Then \sqsubseteq is small when restricted
 540 to the basis, i.e. $\beta(b_1) \sqsubseteq \beta(b_2)$ has size \mathcal{V} for every two elements $b_1, b_2 : B$. Hence, we have
 541 $\sqsubseteq^B : B \rightarrow B \rightarrow \mathcal{V}$ such that $\prod_{b_1, b_2 : B} (b_1 \sqsubseteq^B b_2) \simeq (\beta(b_1) \sqsubseteq \beta(b_2))$.

542 **Proof.** Let $b_1, b_2 : B$ and note that we have the following equivalences:

$$\begin{aligned}
 543 \quad (\beta(b_1) \sqsubseteq \beta(b_2)) &\simeq \prod_{b:B} (\beta(b) \ll \beta(b_1) \rightarrow \beta(b) \ll \beta(b_2)) && \text{(by Lemma 51)} \\
 544 \quad &\simeq \prod_{b:B} (b \ll^B b_1 \rightarrow b \ll^B b_2) && \text{(by definition of a basis),} \\
 545 \quad &
 \end{aligned}$$

546 but the latter is a type in \mathcal{V} . ◀

547 The most significant properties of a basis are the interpolation properties. We consider
 548 nullary, unary and binary versions here. The binary interpolation property actually follows
 549 fairly easily from the unary one, but we still record it, because we wish to show that bases
 550 are examples of the abstract bases that we define later (cf. Example 62).

551 ► **Lemma 53** (Nullary interpolation). Let D be a dcpo with a basis $\beta : B \rightarrow D$. For every
 552 $x : D$, there exists some $b : B$ such that $\beta(b) \ll x$.

553 **Proof.** Immediate from the definitions of a basis and a directed family. ◀

554 ► **Lemma 54** (Unary interpolation). Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$ and let $x, y : D$.
 555 If $x \ll y$, then there exists some $b : B$ such that $x \ll \beta(b) \ll y$.

556 **Proof.** Our proof is a predicative version of [12]. Let $x, y : D$ with $x \ll y$. Since β is a basis,
 557 there exists an approximating family $\alpha : I \rightarrow D$ for y . Consider the family

$$558 \quad \left(K \equiv \sum_{b:B} \sum_{i:I} b \ll^B \alpha_i : \mathcal{V} \right) \xrightarrow{\text{pr}_1} B \xrightarrow{\beta} D. \quad (\dagger)$$

559 ► **Claim.** The family (\dagger) is directed.

560 Proof. By directedness of α and nullary interpolation, the type K is inhabited.

561 Now suppose that we have $b_1, b_2 : B$ and $i_1, i_2 : I$ with $b_1 \ll^B \alpha_{i_1}$ and $b_2 \ll^B \alpha_{i_2}$. By
 562 directedness of α , there exists $k : I$ with $\alpha_{i_1}, \alpha_{i_2} \sqsubseteq^B \alpha_k$. Since β is a basis for D , there exists
 563 an approximating family $\gamma : J \rightarrow B$ for $\beta(\alpha_k)$. From $b_1 \ll^B \alpha_{i_1}$ we obtain $b_1 \ll^B \alpha_k$ and
 564 similarly, $b_2 \ll^B \alpha_k$. Hence, there exist $j_1, j_2 : J$ such that $b_1 \sqsubseteq^B \gamma_{j_1}$ and $b_2 \sqsubseteq^B \gamma_{j_2}$. By
 565 directedness of J , there exists $m : J$ with $\gamma_{j_1}, \gamma_{j_2} \sqsubseteq^B \gamma_m$. Thus, putting this all together, we
 566 see that: $b_1, b_2 \sqsubseteq^B \gamma_m \ll^B \alpha_k$. Hence, (\dagger) is directed. \triangleleft

567 Thus, (\dagger) has a supremum s in D .

568 \triangleright Claim. We have $y \sqsubseteq s$.

569 Proof. Since $y = \bigsqcup \beta \circ \alpha$, it suffices to prove that $\beta(\alpha_i) \sqsubseteq s$ for every $i : I$. Let $i : I$ be
 570 arbitrary and let $\gamma_j : J \rightarrow B$ be some approximating family for $\beta(\alpha_i)$. Then it is enough to
 571 establish $\beta(\gamma_j) \sqsubseteq s$ for every $j : J$. But we know that $\gamma_j \ll^B \alpha_i$, so $\beta_{\gamma_j} \sqsubseteq s$ by definition of
 572 (\dagger) and the fact that s is the supremum of (\dagger) . \triangleleft

573 Finally, from $y \sqsubseteq s$ and $x \ll y$, it follows that there must exist $b : B$ and $i : I$ such that:
 574 $x \sqsubseteq \beta(b) \ll \beta(\alpha_i) \ll y$, which finishes the proof. \blacktriangleleft

575 **► Lemma 55** (Binary interpolation). *Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$ and let*
 576 *$x, y, z : D$. If $x, y \ll z$, then there exists some $b : B$ such that $x, y \ll \beta(b) \ll z$.*

577 **Proof.** Let $x, y, z : D$ such that $x, y \ll z$. By unary interpolation, there are $b_x, b_y : B$ such
 578 that $x \ll \beta(b_x) \ll z$ and $y \ll \beta(b_y) \ll z$. Since β is a basis, there exists a family $\alpha : I \rightarrow B$
 579 such that $\beta(\alpha_i) \ll z$ for every $i : I$, and $\beta \circ \alpha$ is directed and has supremum z . Since
 580 $\beta(b_x) \ll z$, there must exist $i_x : I$ with $\beta(b_x) \sqsubseteq \beta(\alpha_{i_x})$. Similarly, there exists $i_y : I$ such
 581 that $\beta(b_y) \sqsubseteq \beta(\alpha_{i_y})$. By directedness of $\beta \circ \alpha$, there exists $k : I$ with $\beta(\alpha_{i_x}), \beta(\alpha_{i_y}) \sqsubseteq \beta(\alpha_k)$.
 582 Hence,

$$583 \quad x \ll \beta(b_x) \sqsubseteq \beta(\alpha_{i_x}) \sqsubseteq \beta(\alpha_k) \ll z \quad \text{and} \quad y \ll \beta(b_y) \sqsubseteq \beta(\alpha_{i_y}) \sqsubseteq \beta(\alpha_k) \ll z,$$

584 so that $x, y \ll \beta(\alpha_k) \ll z$, as wished. \blacktriangleleft

585 6.3 Algebraic Dcpo

586 We now turn to a particular class of continuous dcpo, called algebraic dcpo.

587 **► Definition 56** (Algebraic dcpo). *A dcpo D is algebraic if there exists some basis $\beta : B \rightarrow D$*
 588 *for it such that $\beta(b)$ is compact for every $b : B$.*

589 **► Lemma 57.** *Let D be a \mathcal{V} -dcpo. Then D is algebraic if and only if there exists $\beta : B \rightarrow D$*
 590 *with $B : \mathcal{V}$ such that*

- 591 (i) *every element $\beta(b)$ is compact;*
- 592 (ii) *for every $x : D$, there exists $\alpha : I \rightarrow B$ with $I : \mathcal{V}$ such that $x = \bigsqcup \beta \circ \alpha$.*

593 **Proof.** We just need to show that having $\beta : B \rightarrow D$ and $\alpha : I \rightarrow B$ such that every element
 594 $\beta(b)$ is compact and $x = \bigsqcup \beta \circ \alpha$, already implies that $\beta(\alpha_i) \ll x$ for every $i : I$. But if $i : I$,
 595 then $\beta(\alpha_i) \ll \beta(\alpha_i) \sqsubseteq \bigsqcup \beta \circ \alpha = x$ by compactness of $\beta(\alpha_i)$, so Lemma 42(iii) now finishes
 596 the proof. \blacktriangleleft

597 **► Example 58** ($\mathcal{L}_{\mathcal{U}}(X)$ is algebraic). Let $X : \mathcal{U}$ be a set and consider $\mathcal{L}_{\mathcal{U}}(X) : \mathcal{U}\text{-DCPO}_{\mathcal{U}^+, \mathcal{U}^+}$.
 598 The basis $[\perp, \eta_X] : (1_{\mathcal{U}} + X) \rightarrow \mathcal{L}_{\mathcal{U}}(X)$ exhibits $\mathcal{L}_{\mathcal{U}}(X)$ as an algebraic dcpo.

Proof. By Example 46, the elements \perp and $\eta_X(x)$ (with $x : X$) are all compact, so it remains to show that $\mathbf{1}_U + X$ is indeed a basis. Recalling Lemmas 19 and 20, we can write any element $(P, \varphi) : \mathcal{L}_V(X)$ as the directed join $\bigsqcup([\perp, \eta_X] \circ \alpha)$ with $\alpha \equiv [\text{id}, \varphi] : (\mathbf{1}_U + P) \rightarrow (\mathbf{1}_U + X)$. By Lemma 57 the proof is finished. \triangleleft

► **Example 59** ($\mathcal{P}_U(X)$ is algebraic). Let $X : \mathcal{U}$ be a set and consider $\mathcal{P}_U(X) : \mathcal{U}\text{-DCPO}_{U^+, U}$. The basis $\iota : \text{List}(X) \rightarrow \mathcal{P}_U(X)$ that maps a finite list to a Kuratowski finite subset exhibits $\mathcal{P}_U(X)$ as an algebraic dpcpo.

Proof. By Example 48, the element $\iota(l)$ is compact for every list $l : \text{List}(X)$, so it remains to show that $\text{List}(X)$ is indeed a basis. In the proof of Example 48, we saw that every \mathcal{U} -subset A of X can be expressed as the directed supremum $\bigsqcup \iota \circ \alpha$ where $\alpha : \text{List}(\mathbb{T}(A)) \rightarrow \text{List}(X)$ takes a list $[(x_0, p_0), \dots, (x_{n-1}, p_{n-1})]$ to the list $[x_0, \dots, x_{n-1}]$. Another application of Lemma 57 now finishes the proof. \triangleleft

► **Example 60** (Scott's D_∞ is algebraic). The pointed dpcpo $D_\infty : \mathcal{U}_0\text{-DCPO}_{U_1, U_1}$ with $D_\infty \cong D_\infty^{D_\infty}$ from Section 5.2 is algebraic. We postpone the proof until Section 6.4, since we will need some additional results on locally small dcpos.

6.4 Ideal Completion

Finally, we consider how to build dcpos from posets, or more generally from abstract bases, using the rounded ideal completion [1, Section 2.2.6]. Given our definition of the notion of dpcpo, the reader might expect us to define ideals using families rather than subsets. However, we use subsets for extensionality reasons. Two subsets A and B of some X are equal exactly when $x \in A \iff x \in B$ for every $x : X$. However, given two (directed) families $\alpha : I \rightarrow X$ and $\beta : J \rightarrow X$, it is of course not the case (it does not even typecheck) that $\alpha = \beta$ when $\Pi_{i:I} \exists j:J \alpha_i = \beta_j$ and $\Pi_{j:J} \exists i:I \beta_j = \alpha_i$ hold. We could try to construct the ideal completion by quotienting the families, but then it seems impossible to define directed suprema in the ideal completion without resorting to choice.

► **Definition 61** (Abstract basis). A pair (B, \prec) with $B : \mathcal{V}$ and \prec taking values in \mathcal{V} is called a \mathcal{V} -abstract basis if:

- (i) \prec is proposition-valued;
- (ii) \prec is transitive;
- (iii) \prec satisfies nullary interpolation, i.e. for every $x : B$, there exists some $y : B$ with $y \prec x$;
- (iv) \prec satisfies binary interpolation, i.e. for every $x, y : B$ with $x \prec y$, there exists some $z : B$ with $x \prec z \prec y$.

► **Example 62.** Let D be a \mathcal{V} -dpcpo with a basis $\beta : B \rightarrow D$. By Lemmas 42, 53 and 55, the pair (B, \ll^B) is an example of a \mathcal{V} -abstract basis.

► **Example 63.** Any preorder (P, \sqsubseteq) with $P : \mathcal{V}$ and \sqsubseteq taking values in \mathcal{V} is a \mathcal{V} -abstract basis, since reflexivity implies both interpolation properties.

For the remainder of this section, fix some arbitrary \mathcal{V} -abstract basis (B, \prec) .

► **Definition 64** (Directed subset). Let A be a \mathcal{V} -subset of B . Then A is directed if A is inhabited (i.e. $\exists x:B x \in A$ holds) and for every $x, y \in A$, there exists some $z \in A$ such that $x, y \sqsubseteq z$.

640 ► **Definition 65** (Ideal, lower set). *Let A be a \mathcal{V} -subset of B . Then A is an ideal if A is a*
 641 *directed subset of B and A is a lower set, i.e. if $x \prec y$ and $y \in A$, then $x \in A$ as well.*

642 ► **Construction 66** (Rounded ideal completion $\text{Idl}(B, \prec)$). We construct a \mathcal{V} -dcpo, known as
 643 the (rounded) ideal completion $\text{Idl}(B, \prec) : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+, \mathcal{V}}$ of (B, \prec) . The carrier is given by
 644 the type $\sum_{I : B \rightarrow \mathcal{V}} \text{is-ideal}(I)$ of ideals on (B, \prec) . The order is given by subset inclusion \subseteq .
 645 If we have a directed family $\alpha : A \rightarrow \text{Idl}(B, \prec)$ of ideals (with $A : \mathcal{V}$), then the subset given
 646 by $\lambda x. \exists a : A. x \in \alpha_a$ is again an ideal and the supremum of α in $\text{Idl}(B, \prec)$. \lrcorner

647 ► **Lemma 67** (Rounded ideals). *The ideals of $\text{Idl}(B, \prec)$ are rounded. That is, if $I : \text{Idl}(B, \prec)$*
 648 *and $x \in I$, then there exists some $y \in I$ with $x \prec y$.*

649 **Proof.** Immediate from the fact that ideals are directed sets. \blacktriangleleft

650 ► **Definition 68** (Principal ideal $\downarrow x$). We write $\downarrow(-) : B \rightarrow \text{Idl}(B, \prec)$ for the map that takes
 651 $x : B$ to the principal ideal $\lambda y. y \prec x$.

652 ► **Lemma 69.** *Let $I : \text{Idl}(B, \prec)$ be an ideal. Then I may be expressed as the supremum of*
 653 *the directed family $(x, p) : \mathbb{T}(I) \mapsto \downarrow x : \text{Idl}(B, \prec)$, which we will denote by $I = \bigsqcup_{x \in I} \downarrow x$.*

654 **Proof.** Directedness of the family follows from the fact that I is a directed subset. Since I is
 655 a lower set, $\downarrow x \subseteq I$ holds for every $x \in I$, establishing $\bigsqcup_{x \in I} \downarrow x \subseteq I$. The reverse inclusion
 656 follows from Lemma 67. \blacktriangleleft

657 We wish to prove that $\text{Idl}(B, \prec)$ is continuous with basis $\downarrow(-) : B \rightarrow \text{Idl}(B, \prec)$. To this end,
 658 it is useful to express $\ll_{\text{Idl}(B, \prec)}$ in more elementary terms.

659 ► **Lemma 70.** *Let $I, J : \text{Idl}(B, \prec)$ be two ideals. Then $I \ll J$ holds if and only there exists*
 660 *$x \in J$ such that $I \subseteq \downarrow x$.*

661 **Proof.** The left-to-right implication follows immediately from Lemma 69.

662 For the converse, note that $I \ll J$ is a proposition, so we may assume that we have $x \in J$
 663 with $I \subseteq \downarrow x$. Now let $\alpha : A \rightarrow \text{Idl}(B, \prec)$ be a directed family such that $J \subseteq \bigsqcup \alpha$. Then there
 664 must exist some $a : A$ for which $x \in \alpha_a$. But $I \subseteq \downarrow x$ and α_a is a lower set, so $I \subseteq \alpha_a$. \blacktriangleleft

665 ► **Theorem 71.** *The map $\downarrow(-) : B \rightarrow \text{Idl}(B, \prec)$ is a basis for $\text{Idl}(B, \prec)$. Thus, $\text{Idl}(B, \prec)$ is*
 666 *a continuous \mathcal{V} -dcpo.*

667 **Proof.** Let $I : \text{Idl}(B, \prec)$ be arbitrary. By Lemma 69 we can express I as the supremum
 668 $\bigsqcup_{x \in I} \downarrow x$, so it is enough to prove that $\downarrow x \ll I$ for every $x \in I$. But this follows from
 669 Lemmas 67 and 70. \blacktriangleleft

670 ► **Lemma 72.** *If \prec is reflexive, then the compact elements of $\text{Idl}(B, \prec)$ are exactly the*
 671 *principal ideals and $\text{Idl}(B, \prec)$ is algebraic.*

672 **Proof.** Immediate from Lemma 70. \blacktriangleleft

673 ► **Theorem 73.** *The ideal completion is the free dcpo on a small poset. That is, if we have*
 674 *a poset (P, \sqsubseteq) with $P : \mathcal{V}$ and \sqsubseteq taking values in \mathcal{V} , then for every $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and*
 675 *monotone function $f : P \rightarrow D$, there is a unique continuous function $\bar{f} : \text{Idl}(P, \sqsubseteq) \rightarrow D$ such*
 676 *that*

677

$$\begin{array}{ccc}
 P & \xrightarrow{f} & D \\
 \searrow \downarrow(-) & & \nearrow \bar{f} \\
 & \text{Idl}(P, \sqsubseteq) &
 \end{array}$$

678 *commutes.*

679 **Proof.** Given (P, \sqsubseteq) , D and f as in the theorem, we define \bar{f} by mapping an ideal I to the
 680 supremum of the directed (since I is an ideal) family $\mathbb{T}(I) \xrightarrow{\text{pr}_1} P \xrightarrow{f} D$.

681 Commutativity of the diagram expresses that $f(x) = \bigsqcup_{y \sqsubseteq x} f(y)$ for every $x : P$. By anti-
 682 symmetry of \sqsubseteq , it suffices to prove $f(x) \sqsubseteq \bigsqcup_{y \sqsubseteq x} f(y)$ and $\bigsqcup_{y \sqsubseteq x} f(y) \sqsubseteq f(x)$. The first holds
 683 by reflexivity of \sqsubseteq and the second holds because f is monotone.

684 Uniqueness of \bar{f} follows easily using Lemma 69. Finally, continuity of \bar{f} is not hard to
 685 establish either. ◀

686 ► **Definition 74** (Continuous retract, section, retraction). *A \mathcal{V} -dcpo D is a continuous retract*
 687 *of another \mathcal{V} -dcpo E if we have continuous functions $s : D \rightarrow E$ (the section) and $r : E \rightarrow D$*
 688 *(the retraction) such that $r(s(x)) = x$ for every $x : D$.*

689 ► **Theorem 75.** *If E is a dcpo with basis $\beta : B \rightarrow D$ and D is a continuous retract of E*
 690 *with retraction r , then $r \circ \beta$ is a basis for D .*

691 **Proof.** Let E be a dcpo with basis $\beta : B \rightarrow D$ and suppose that we have continuous
 692 retraction $r : E \rightarrow D$ with continuous section $s : D \rightarrow E$. Given $x : D$, there exists some
 693 approximating family $\alpha : I \rightarrow B$ for $s(x)$. We claim that α is an approximating family for x
 694 as well, i.e.

695 (i) $r(\beta(\alpha_i)) \ll x$ for every $i : I$ and

696 (ii) $\bigsqcup r \circ \beta \circ \alpha = x$.

697 The second follows from continuity of r , since: $\bigsqcup r \circ \beta \circ \alpha = r(\bigsqcup \beta \circ \alpha) = r(s(x)) = x$. For
 698 (i), suppose that $i : I$ and that $\gamma : J \rightarrow D$ is a directed family satisfying $x \sqsubseteq \bigsqcup \gamma$. We must
 699 show that there exists $j : J$ with $r(\beta(\alpha_i)) \sqsubseteq \gamma_j$. By continuity of s , we get $s(x) \sqsubseteq \bigsqcup s \circ \gamma$.
 700 Hence, since $\beta(\alpha_i) \ll s(x)$, there must exist $j : J$ with $\beta(\alpha_i) \sqsubseteq s(\gamma_j)$. Thus, by monotonicity
 701 of r , we get the desired $r(\beta(\alpha_i)) \sqsubseteq r(s(\gamma_j)) = \gamma_j$. ◀

702 We now turn to locally small dcpos, as they allow us to find canonical approximating families,
 703 which is used in the proof of Theorem 78.

704 ► **Lemma 76.** *Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$. The following are equivalent:*

705 (i) *D is locally small;*

706 (ii) *$\beta(b) \ll x$ has size \mathcal{V} for every $x : D$ and $b : B$.*

707 **Proof.** Recalling Lemma 51, the type $x \sqsubseteq y$ is equivalent to $\prod_{b:B} (\beta(b) \ll x \rightarrow \beta(b) \ll y)$
 708 for every $x, y : D$. Thus, (ii) implies (i). Conversely, assume that D is locally small and let
 709 $x : D$ and $b : B$. We claim that $\beta(b) \ll x$ is equivalent to $\exists b' : B (b \ll^B b' \times \beta(b') \sqsubseteq_{\text{small}} x) : \mathcal{V}$.
 710 The left-to-right implication is given by Lemma 54, and the converse by Lemma 42(iii). ◀

711 ► **Lemma 77.** *Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$. If D is locally small, then an*
 712 *element $x : D$ is the supremum of the large directed family $(\sum_{b:B} \beta(b) \ll x) \xrightarrow{\text{pr}_1} B \xrightarrow{\beta} D$.*
 713 *Moreover, if D is locally small, then this directed family is small.*

714 **Proof.** The family $\text{pr}_1 \circ \beta$ is directed by the nullary (Lemma 53) and binary (Lemma 55)
 715 interpolation properties. Now suppose that D is locally small. By Lemma 76, we have $I : \mathcal{V}$
 716 and $\alpha : I \rightarrow D$ directed such that $\bigsqcup \alpha$ is the supremum of $(\sum_{b:B} \beta(b) \ll x) \xrightarrow{\text{pr}_1} B \xrightarrow{\beta} D$.
 717 Since $\beta : B \rightarrow D$ is a basis of D , we see that $x \sqsubseteq \bigsqcup \alpha$. For the reverse inequality, it suffices to
 718 show that $\beta(b) \sqsubseteq x$ for every $b : B$ with $\beta(b) \ll x$. But this follows from Lemma 42(ii). ◀

719 ► **Theorem 78.** *Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$ and suppose that D is locally small.*
 720 *Then D is a continuous retract of the algebraic \mathcal{V} -dcpo $\text{Idl}(B, \sqsubseteq^B)$ (recall Lemma 52).*

Proof. By Lemma 72, $\text{Idl}(B, \sqsubseteq^B)$ is indeed algebraic. Let D be a \mathcal{V} -dcpo satisfying the hypotheses of the lemma. Let $\ll_{\text{small}} : B \rightarrow D \rightarrow \mathcal{V}$ be such that $(b \ll_{\text{small}} x) \simeq (\beta(b) \ll x)$ for every $x : D$ and $b : B$.

For every $x : D$, we can consider the subset $\downarrow x$ given by $\lambda(b : B). b \ll_{\text{small}} x$. We show that it is an ideal. By Lemma 77 it is a directed subset. And if $b \in \downarrow x$ and $b' \sqsubseteq^B b$, then $b' \in \downarrow x$ as well by virtue of Lemma 42(iii). So $\downarrow x$ is a lower set, and indeed an ideal.

We claim that the map $\downarrow(-)$ is continuous. By Lemma 43, it is monotone. Thus, we are left to show that if $\alpha : I \rightarrow D$ is directed, then $\downarrow(\bigsqcup \alpha) \subseteq \bigsqcup_{i:I} \downarrow \alpha_i$. Let $b \in \downarrow(\bigsqcup \alpha)$, i.e. $b \in B$ such that $b \ll_{\text{small}} \bigsqcup \alpha$. By Lemma 54, there exists $b' : B$ with $b \ll^B b' \ll_{\text{small}} \bigsqcup \alpha$. Hence, there must exist $i : I$ such that $\beta(b) \ll \beta(b') \sqsubseteq \alpha_i$, thus, $b \in \downarrow \alpha_i$ and $\downarrow(-)$ is indeed continuous.

Next, define $r : \text{Idl}(B, \sqsubseteq^B) \rightarrow D$ using Theorem 73 as the unique continuous function such that

$$\begin{array}{ccc} B & \xrightarrow{\beta} & D \\ \downarrow(-) \searrow & & \nearrow r \\ & \text{Idl}(B, \sqsubseteq^B) & \end{array}$$

commutes, i.e. r maps an ideal I to the directed supremum $\bigsqcup_{b \in I} \beta(b)$ in D .

Finally, we show that $\downarrow(-)$ is a section of r . That is, the equality $\bigsqcup_{b \ll_{\text{small}} x} \beta(b) = x$ holds for every $x : D$. But this is exactly Lemma 77. \blacktriangleleft

One may wonder how restrictive the condition that D is locally small is. We note that if X is a set, then $\mathcal{L}_{\mathcal{V}}(X)$ (by Lemma 18) and $\mathcal{P}_{\mathcal{V}}(X)$ are examples of locally small \mathcal{V} -dcpos. A natural question is what happens with exponentials. In general, E^D may fail to be locally small even when both D and E are. However, we do have the following result.

► Lemma 79. *Let D and E be \mathcal{V} -dcpos. Suppose that D is continuous and E is locally small. Then E^D is locally small.*

Proof. Since being locally small is a proposition, we may assume that we are given a basis $\beta : B \rightarrow D$ of D . We claim that for every two continuous functions $f, g : D \rightarrow E$ we have an equivalence

$$\left(\prod_{x:D} f(x) \sqsubseteq_E g(x) \right) \simeq \left(\prod_{b:B} f(\beta(b)) \sqsubseteq_{\text{small}} g(\beta(b)) \right).$$

Since $B : \mathcal{V}$ and $\sqsubseteq_{\text{small}}$ takes values in \mathcal{V} , the second type is also in \mathcal{V} . For the equivalence, note that the left-to-right implication is trivial. For the converse, assume the right-hand side and let $x : D$. By continuity of D , there exists some approximating family $\alpha : I \rightarrow B$ for x . We use it as follows:

$$\begin{aligned} f(x) &= f\left(\bigsqcup \beta \circ \alpha\right) \\ &= \bigsqcup_{i:I} f(\beta(\alpha_i)) && \text{(by continuity of } f\text{)} \\ &\sqsubseteq \bigsqcup_{i:I} g(\beta(\alpha_i)) && \text{(by assumption)} \\ &= g\left(\bigsqcup \beta \circ \alpha\right) && \text{(by continuity of } g\text{)} \\ &= g(x), \end{aligned}$$

which finishes the proof. \blacktriangleleft

Moreover, the (co)limit of locally small dcpos is locally small.

► **Lemma 80.** *Given a system $(D_i, \varepsilon_{i,j}, \pi_{i,j})$ as in Section 5.1, if every D_i is locally small, then so is D_∞ .*

Finally, the requirement that D is locally small is necessary, in the following sense.

► **Lemma 81.** *Let D be a \mathcal{V} -dcpo with basis $\beta : B \rightarrow D$. Suppose that D is a continuous retract of $\text{Idl}(B, \sqsubseteq^B)$. Then D is locally small.*

Proof. Let $s : D \rightarrow \text{Idl}(B, \sqsubseteq^B)$ be a section of a map $r : \text{Idl}(B, \sqsubseteq^B) \rightarrow D$, with both maps continuous. Then $x \sqsubseteq_D y$ holds if and only if $s(x) \sqsubseteq_{\text{Idl}(B, \sqsubseteq^B)} s(y)$. Since $\text{Idl}(B, \sqsubseteq^B)$ is locally small, so must D . ◀

We have now developed the theory sufficiently to give a proof of Example 60.

Proof of Example 60 (Scott's D_∞ is algebraic). Firstly, notice that D_0 is not just a \mathcal{U}_0 -dcpo, but in fact a \mathcal{U}_0 -sup lattice, i.e. it has joins for all families indexed by types in \mathcal{U}_0 . Moreover, since joins in exponentials are given pointwise, every D_n is in fact a \mathcal{U}_0 -sup lattice. In particular, every D_n has all finite joins. Hence, if we have $\alpha : I \rightarrow D_n$ with $I : \mathcal{U}_0$, then we can consider the directed family $\bar{\alpha} : \bar{I} \rightarrow D_n$ with $\bar{I} \equiv \sum_{k:\mathbf{N}} (\text{Fin } k \rightarrow D_n)$ and $\bar{\alpha}$ mapping a pair (k, f) to the finite join $\bigvee_{0 \leq i < k} f(i)$. Moreover, if every α_i is compact, then so is every $\bar{\alpha}_{\bar{i}}$, since finite joins of compact elements are compact again. We show this explicitly for binary joins from which the general case follows by induction. If $a, b : D_n$ are compact and $a, b \sqsubseteq a \vee b \sqsubseteq \bigsqcup \gamma$ for some directed family $\gamma : J \rightarrow D_n$, then by compactness of a and b , there exist $j_a, j_b : J$ such that $a \sqsubseteq \gamma_{j_a}$ and $b \sqsubseteq \gamma_{j_b}$. By directedness of γ , there exists $k : J$ with $a, b \sqsubseteq \gamma_k$. Hence, $a \vee b \sqsubseteq \gamma_k$, as desired.

► **Claim.** Every D_n is locally small has a basis $\beta_n : B_n \rightarrow D_n$ of compact elements.

Proof. We prove this by induction. For $n = 0$, this follows from Lemma 18 and Example 58. Now suppose that B_m is locally small and has a basis $\beta_m : B_m \rightarrow D_m$. By Lemma 79, the dcpo $B_{m+1} \equiv B_m^{B_m}$ is locally small. If we have $a, b : B_m$, then we define the continuous *step function* $(a \Rightarrow b) : D_m \rightarrow D_m$ by $x \mapsto \bigvee_{\beta_m(a) \sqsubseteq x} \beta_m(b)$, which is well-defined since D_m is locally small. We are going to show that $a \Rightarrow b$ is compact for every $a, b : B_m$ and that every $f : D_{m+1}$ is the join of certain step functions. To this end, we first observe that

$$(a \Rightarrow b \sqsubseteq f) \iff (\beta_m(b) \sqsubseteq f(\beta_m(a))), \quad (\dagger)$$

which follows from the fact that continuous functions are monotone.

For compactness, suppose that $a \Rightarrow b \sqsubseteq \bigsqcup_{i:I} f_i$. By (\dagger) we have $\beta_m(b) \sqsubseteq \bigvee_{i:I} (f_i(\beta_m(a)))$. By compactness of $\beta_m(b)$, there exists $i : I$ such that $\beta_m(b) \sqsubseteq f_i(\beta_m(a))$ already. Using (\dagger) once more, we get the desired $a \Rightarrow b \sqsubseteq f_i$.

Now let $f : D_m \rightarrow D_m$ be continuous. We claim that f is the join of the step-functions below it, i.e. $f = \bigvee_{a,b:B_m, a \Rightarrow b \sqsubseteq f} a \Rightarrow b$, which is well-defined, since D_{m+1} is locally small. One inequality clearly holds as we are only considering step-functions below f . For the reverse inequality, let $x : D_m$ be arbitrary. By Lemma 77, we have:

$$x = \bigsqcup_{\substack{a':B_m \\ \beta_m(a') \ll x}} \beta_m(a') \quad \text{and} \quad f(x) = \bigsqcup_{\substack{b':B_m \\ \beta_m(b') \ll f(x)}} \beta_m(b'). \quad (\ddagger)$$

Hence, it suffices to show that $\beta_m(b') \sqsubseteq \bigvee_{a,b:B_m, a \Rightarrow b \sqsubseteq f} (a \Rightarrow b)(x)$ whenever $b' : B_m$ is such that $\beta_m(b') \ll f(x)$. By (\dagger) and the definition of a step-function it is enough to find $a' : B_m$

such that $\beta_m(b') \sqsubseteq f(\beta_m(a'))$ and $\beta_m(a') \sqsubseteq x$. Using (\ddagger) , our assumption $\beta_m(b') \ll f(x)$ and continuity of f , we get that there exists $a' : B_m$ with $\beta_m(a') \ll x$ (and thus $\beta_m(a') \sqsubseteq x$) and $b_m(b') \sqsubseteq f(\beta_m(a'))$, as desired.

Thus, by the paragraph preceding the claim, D_{m+1} has a basis of compact elements: $\beta_{m+1} : (\sum_{k:\mathbf{N}} (\text{Fin}(k) \rightarrow (D_m \times D_m))) \rightarrow D_{m+1}$ with $\beta_{m+1}(k, \lambda i.(a_i, b_i)) \equiv \bigvee_{0 \leq i < k} a_i \Rightarrow b_i$, finishing the proof of the claim. \triangleleft

Finally, we show that a basis of compact elements for D_∞ is $\beta_\infty : (B_\infty \equiv \sum_{n:\mathbf{N}} B_n) \rightarrow D_\infty$ where $\beta_\infty(n, b) \equiv \varepsilon_{n,\infty}(\beta_n(b))$. We first check compactness by showing that if $x : D_n$ is compact, then so is $\varepsilon_{n,\infty}(x)$. This follows easily from the fact that $(\varepsilon_{n,\infty}, \pi_{n,\infty})$ is an embedding-projection. For if $\alpha : I \rightarrow D_\infty$ is directed and $\varepsilon_{n,\infty}(x) \sqsubseteq \bigsqcup \alpha$, then $x = \pi_{n,\infty}(\varepsilon_{n,\infty}(x)) \sqsubseteq \pi_{n,\infty}(\bigsqcup \alpha) = \bigsqcup \pi_{n,\infty} \circ \alpha$, by continuity of $\pi_{n,\infty}$. Thus, by compactness of x , there exist $i : I$ such that $x \sqsubseteq \pi_{n,\infty}(\alpha_i)$ already. Hence, $\varepsilon_{n,\infty}(x) \sqsubseteq \varepsilon_{n,\infty}(\pi_{n,\infty}(\alpha_i)) \sqsubseteq \alpha_i$, so $\varepsilon_{n,\infty}(x)$ is indeed compact. Now let $\sigma : D_\infty$ be arbitrary. As mentioned in the proof of Theorem 37, we have $\sigma = \bigsqcup_{n:\mathbf{N}} \varepsilon_{n,\infty}(\sigma_n)$. By Lemma 77 and the claim, we can express every $\sigma_n : D_n$ as $\bigsqcup_{b:B_n, \beta_n(b) \ll \sigma_n} \beta_n(b)$. Hence,

$$\sigma = \bigsqcup_{n:\mathbf{N}} \varepsilon_{n,\infty} \left(\bigsqcup_{\substack{b:B_n \\ \beta_n(b) \ll \sigma_n}} \beta_n(b) \right) = \bigsqcup_{n:\mathbf{N}} \bigsqcup_{\substack{b:B_n \\ \beta_n(b) \ll \sigma_n}} \varepsilon_{n,\infty}(\beta_n(b))$$

by continuity of $\varepsilon_{n,\infty}$. Thus, σ may be expressed as the supremum of the directed family $(\sum_{n:\mathbf{N}} \sum_{b:B_n} \beta_n(b) \ll \sigma_n) \rightarrow B_\infty \xrightarrow{\beta_\infty} D_\infty$. (And in light of Lemma 76 and the claim, the type $\sum_{n:\mathbf{N}} \sum_{b:B_n} \beta_n(b) \ll \sigma_n$ can be replaced by a type in \mathcal{U}_0 .) Finally, using Lemma 57, we see that D_∞ is indeed algebraic. \triangleleft

We end this section by describing an example of a continuous dcpo, built using the ideal completion, that is not algebraic. In fact, this dcpo has no compact elements at all.

► **Example 82** (A continuous dcpo that is not algebraic). We inductively define a type and an order representing dyadic rationals $m/2^n$ in the interval $(-1, 1)$ for integers m, n . The intuition for the upcoming definitions is the following. Start with the point 0 in the middle of the interval (represented by `center` below). Then consider the two functions (respectively represented by `left` and `right` below)

$$\begin{aligned} l, r &: (-1, 1) \rightarrow (-1, 1) \\ l(x) &= (x - 1)/2 \\ r(x) &= (x + 1)/2 \end{aligned}$$

that generate the dyadic rationals. Observe that $l(x) < 0 < r(x)$ for every $x : (-1, 1)$. Accordingly, we inductively define the following types.

► **Definition 83** (Dyadics \mathbb{D}). *The type of dyadics $\mathbb{D} : \mathcal{U}_0$ is the inductive type with three constructors:*

$$\text{center} : \mathbb{D} \quad \text{left} : \mathbb{D} \rightarrow \mathbb{D} \quad \text{right} : \mathbb{D} \rightarrow \mathbb{D}.$$

► **Definition 84** (Order \prec on \mathbb{D}). *Let $\prec : \mathbb{D} \rightarrow \mathbb{D} \rightarrow \mathcal{U}_0$ be inductively defined as:*

$$\begin{array}{lll} \text{center} \prec \text{center} & \equiv \mathbf{0} & \text{left } x \prec \text{center} \equiv \mathbf{1} & \text{right } x \prec \text{center} \equiv \mathbf{0} \\ \text{center} \prec \text{left } y & \equiv \mathbf{0} & \text{left } x \prec \text{left } y \equiv x \prec y & \text{right } x \prec \text{left } y \equiv \mathbf{0} \\ \text{center} \prec \text{right } y & \equiv \mathbf{1} & \text{left } x \prec \text{right } y \equiv \mathbf{1} & \text{right } x \prec \text{right } y \equiv x \prec y. \end{array}$$

One then shows that \prec is proposition-valued, transitive, irreflexive, trichotomous, dense and that it has no endpoints. *Trichotomy* means that exactly one of $x \prec y$, $x = y$, $y \prec x$ holds. *Density* says that for every $x, y : \mathbb{D}$, there exists some $z : \mathbb{D}$ such that $x \prec z \prec y$. Finally, *having no endpoints* means that for every $x : \mathbb{D}$, there exist some $y, z : \mathbb{D}$ with $y \prec x \prec z$. Using these properties, we can show that (\mathbb{D}, \prec) is a \mathcal{U}_0 -abstract basis. Thus, taking the rounded ideal completion, we get $\text{Idl}(\mathbb{D}, \prec) : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_0}$, which is continuous with basis $\downarrow(-) : \mathbb{D} \rightarrow \text{Idl}(\mathbb{D}, \prec)$ by Theorem 71. But $\text{Idl}(\mathbb{D}, \prec)$ cannot be algebraic, since none of its elements are compact. Indeed suppose that we had an ideal I with $I \ll I$. By Lemma 70, there would exist $x \in I$ with $I \subseteq \downarrow x$. But this implies $x \prec x$, but \prec is irreflexive, so this is impossible.

7 Conclusion and Future Work

We have developed domain theory constructively and predicatively in univalent foundations, including Scott's D_∞ model of the untyped λ -calculus, as well as notions of continuous and algebraic dcpos. We avoid size issues in our predicative setting by having large dcpos with joins of small directed families. Often we find it convenient to work with locally small dcpos, whose orders have small truth values.

In future work, we wish to give a predicative account of the theory of algebraic and continuous exponentials, which is a rich and challenging topic even classically. We also intend to develop applications to topology and locale theory. It is also important to understand when classical theorems do not have constructive and predicative counterparts. For instance, Zorn's Lemma doesn't imply excluded middle but it implies propositional resizing [9] and we are working on additional examples.

We have formalized the following in Agda [10], in addition to the Scott model of PCF and its computational adequacy [8, 18]:

1. dcpos,
2. limits and colimits of dcpos, Scott's D_∞ ,
3. lifting and exponential constructions,
4. pointed dcpos have subsingleton joins (in the right universe),
5. way-below relation, continuous, algebraic dcpos, interpolation properties,
6. abstract bases and rounded ideal completions (including its universal property),
7. continuous dcpos are continuous retract of their ideal completion, and hence of algebraic dcpos,
8. ideal completion of dyadics, giving an example of a non-algebraic, continuous dcpo.

In the near future we intend to complete our formalization to also include Theorems 21, 23 and 25, Examples 59 and 60, and Lemma 79.

References

- 1 S. Abramsky and A. Jung. Domain theory. In S. Abramsky, D. M. Gabbay, and T. S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, volume 3, pages 1–168. Clarendon Press, 1994.
- 2 Jiří Adámek and Jiří Rosický. *Locally Presentable and Accessible Categories*, volume 189 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1994. doi:10.1017/CB09780511600579.
- 3 Roberto M. Amadio and Pierre-Louis Curien. *Domains and Lambda-Calculi*, volume 46 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 1998. doi:10.1017/CB09780511983504.

- 885 **4** Andrej Bauer and Iztok Kavkler. A constructive theory of continuous domains suitable for
 886 implementation. *Annals of Pure and Applied Logic*, 159(3):251–267, 2009. doi:10.1016/j.
 887 apal.2008.09.025.
- 888 **5** Nick Benton, Andrew Kennedy, and Carsten Varming. Some domain theory and denotational
 889 semantics in Coq. In Stefan Berghofer, Tobias Nipkow, Christian Urban, and Makarius Wenzel,
 890 editors, *Theorem Proving in Higher Order Logics (TPHOLs 2009)*, volume 5674 of *Lecture Notes*
 891 *in Computer Science*, pages 115–130. Springer, 2009. doi:10.1007/978-3-642-03359-9_10.
- 892 **6** The Agda community. Agda wiki. <https://wiki.portal.chalmers.se/agda/pmwiki.php>.
- 893 **7** Thierry Coquand, Giovanni Sambin, Jan Smith, and Silvio Valentini. Inductively generated
 894 formal topologies. *Annals of Pure and Applied Logic*, 124(1–3):71–106, 2003. doi:10.1016/
 895 s0168-0072(03)00052-6.
- 896 **8** Tom de Jong. The Scott model of PCF in univalent type theory. November 2019. **arXiv:**
 897 1904.09810.
- 898 **9** Tom de Jong. Domain theory in predicative Univalent Foundations. Abstract for a talk at
 899 *HoTT/UF 2020* on 7 July, 2020. URL: [https://hott-uf.github.io/2020/HoTTUF_2020_](https://hott-uf.github.io/2020/HoTTUF_2020_paper_14.pdf)
 900 .
- 901 **10** Tom de Jong. Formalisation of domain theory in constructive and predicative univalent
 902 foundations. <https://github.com/tomdjong/TypeTopology>, June 2020. Agda development.
- 903 **11** Robert Dockins. Formalized, Effective Domain Theory in Coq. In Gerwin Klein and Ruben
 904 Gamboa, editors, *Interactive Theorem Proving (ITP 2014)*, volume 8558 of *Lecture Notes in*
 905 *Computer Science*, pages 209–225. Springer, 2014. doi:10.1007/978-3-319-08970-6_14.
- 906 **12** Martín H. Escardó. The proof of the interpolation property of the way-below relation of a
 907 continuous poset. Unpublished note. The proof has been included in Escardó’s *Function-space*
 908 *compactifications of function spaces* and in the book *Continuous lattices and domains* by Gierz
 909 et. al., March 2000. URL: <https://www.cs.bham.ac.uk/~mhe/papers/interpolation.pdf>.
- 910 **13** Martín H. Escardó and Cory M. Knapp. Partial Elements and Recursion via Dominances
 911 in Univalent Type Theory. In Valentin Goranko and Mads Dam, editors, *26th EACSL*
 912 *Annual Conference on Computer Science Logic (CSL 2017)*, volume 82 of *Leibniz International*
 913 *Proceedings in Informatics (LIPIcs)*, pages 21:1–21:16. Schloss Dagstuhl–Leibniz-Zentrum für
 914 Informatik, 2017. doi:10.4230/LIPIcs.CSL.2017.21.
- 915 **14** Martín Hötzel Escardó. Introduction to Univalent Foundations of Mathematics with Agda.
 916 February 2020. **arXiv:**1911.00580.
- 917 **15** Martín Hötzel Escardó. Various new theorems in constructive univalent mathematics written in
 918 Agda. <https://github.com/martinescardo/TypeTopology>, June 2020. Agda development.
- 919 **16** G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. Mislove, and D. S. Scott. *Continu-*
 920 *ous Lattices and Domains*, volume 93 of *Encyclopedia of Mathematics and its Applications*.
 921 Cambridge University Press, 2003. doi:10.1017/CB09780511542725.
- 922 **17** Gerhard Gierz, Karl Heinrich Hofmann, Klaus Keimel, Jimmie D. Lawson, Michael W. Mislove,
 923 and Dana S. Scott. *A Compendium of Continuous Lattices*. Springer Berlin Heidelberg, 1980.
 924 doi:10.1007/978-3-642-67678-9.
- 925 **18** Brendan Hart. Investigating Properties of PCF in Agda. MSci project, School of Computer
 926 Science, University of Birmingham, 2020. Report and Agda code available at [https://github.](https://github.com/BrendanHart/Investigating-Properties-of-PCF)
 927 com/BrendanHart/Investigating-Properties-of-PCF.
- 928 **19** Michael Hedberg. A type-theoretic interpretation of constructive domain theory. *Journal of*
 929 *Automated Reasoning*, 16(3):369–425, 1996. doi:10.1007/BF00252182.
- 930 **20** J. M. E. Hyland. First steps in synthetic domain theory. In A. Carboni, M. C. Peddicchio, and
 931 G. Rosolini, editors, *Category Theory*, volume 1488 of *Lecture Notes in Mathematics*, pages
 932 131–156. Springer, 1991. doi:10.1007/bfb0084217.
- 933 **21** Peter Johnstone and André Joyal. Continuous categories and exponentiable toposes. *Journal*
 934 *of Pure and Applied Algebra*, 25(3):255–296, 1982. doi:10.1016/0022-4049(82)90083-4.
- 935 **22** Tatsuji Kawai. Predicative theories of continuous lattices. June 2020. **arXiv:**2006.05642.

- 936 23 Nicolai Kraus, Martín Escardó, Thierry Coquand, and Thorsten Altenkirch. Notions of
937 Anonymous Existence in Martin-Löf Type Theory. *Logical Methods in Computer Science*,
938 13(1), 2017. doi:10.23638/LMCS-13(1:15)2017.
- 939 24 David Lidell. Formalizing domain models of the typed and the untyped lambda calculus in Agda.
940 Master's thesis, Chalmers University of Technology and University of Gothenburg, 2020. Final
941 version in preparation. Agda code available at <https://github.com/DoppeD/CwfsOfDomains>.
- 942 25 John Longley and Dag Normann. *Higher-Order Computability*. Springer, 2015. doi:10.1007/
943 978-3-662-47992-6.
- 944 26 Maria Emilia Maietti and Silvio Valentini. Exponentiation of Scott formal topologies. volume 73,
945 pages 111–131, 2004. Proceedings of the Workshop on Domains VI. doi:10.1016/j.entcs.
946 2004.08.005.
- 947 27 Michael Makkai and Robert Paré. *Accessible categories: the foundations of categorical model*
948 *theory*, volume 104 of *Contemporary Mathematics*. American Mathematical Society, Providence,
949 RI, 1989. URL: <https://doi.org/10.1090/conm/104>, doi:10.1090/conm/104.
- 950 28 Sara Negri. Continuous lattices in formal topology. In Eduardo Giménez and Christine
951 Paulin-Mohring, editors, *Types for Proofs and Programs (TYPES 1996)*, volume 1512 of
952 *Lecture Notes in Computer Science*, pages 333–353. Springer, 1998. doi:10.1007/BFb0097800.
- 953 29 Sara Negri. Continuous domains as formal spaces. *Mathematical Structures in Computer*
954 *Science*, 12(1):19–52, 2002. doi:10.1017/S0960129501003450.
- 955 30 Dirk Pattinson and Mina Mohammadian. Constructive Domains with Classical Witnesses.
956 June 2020. arXiv:1910.04948.
- 957 31 G. D. Plotkin. LCF considered as a programming language. *Theoretical Computer Science*,
958 5(3):223–255, 1977. doi:10.1016/0304-3975(77)90044-5.
- 959 32 Bernhard Reus. Formalizing synthetic domain theory — the basic definitions. *Journal of*
960 *Automated Reasoning*, 23(3-4):411–444, 1999. doi:10.1023/A:1006258506401.
- 961 33 Bernhard Reus and Thomas Streicher. General synthetic domain theory — a logical
962 approach. *Mathematical Structures in Computer Science*, 9(2):177—223, 1999. doi:
963 10.1017/S096012959900273X.
- 964 34 G. Rosolini. Categories and effective computations. In David H. Pitt, Axel Poigné, and
965 David E. Rydeheard, editors, *Category Theory and Computer Science*, volume 283 of *Lecture*
966 *Notes in Computer Science*, pages 1–11. Springer, 1987.
- 967 35 Giuseppe Rosolini. *Continuity and effectiveness in topoi*. PhD thesis, University of Oxford,
968 1986.
- 969 36 Giovanni Sambin. Intuitionistic formal spaces—a first communication. In *Mathematical logic*
970 *and its applications*, pages 187–204. Springer, 1987. doi:10.1007/978-1-4613-0897-3_12.
- 971 37 Giovanni Sambin, Silvio Valentini, and Paolo Virgili. Constructive domain theory as a branch
972 of intuitionistic pointfree topology. *Theoretical Computer Science*, 159(2):319–341, 1996.
973 doi:10.1016/0304-3975(95)00169-7.
- 974 38 Dana S. Scott. Continuous lattices. In F.W. Lawvere, editor, *Toposes, Algebraic Geometry*
975 *and Logic*, volume 274 of *Lecture Notes in Mathematics*, pages 97–136. Springer, 1972. doi:
976 10.1007/BFB0073967.
- 977 39 Dana S. Scott. Lectures on a mathematical theory of computation. In Manfred Broy and
978 Gunther Schmidt, editors, *Theoretical Foundations of Programming Methodology: Lecture*
979 *Notes of an International Summer School, directed by F. L. Bauer, E. W. Dijkstra and C.*
980 *A. R. Hoare*, volume 91 of *NATO Advanced Study Institutes Series*, pages 145–292. Springer,
981 1982. doi:10.1007/978-94-009-7893-5_9.
- 982 40 Dana S. Scott. A type-theoretical alternative to ISWIM, CUCH, OWHY. *Theoretical Computer*
983 *Science*, 121(1):411–440, 1993. doi:10.1016/0304-3975(93)90095-B.
- 984 41 Andrew W. Swan. Choice, collection and covering in cubical sets. Talk in the electronic
985 *HoTTEST* seminar, 6 November. Slides at [https://www.uwo.ca/math/faculty/kapulkin/](https://www.uwo.ca/math/faculty/kapulkin/seminars/hotttestfiles/Swan-2019-11-06-HoTTEST.pdf)
986 [seminars/hotttestfiles/Swan-2019-11-06-HoTTEST.pdf](https://www.uwo.ca/math/faculty/kapulkin/seminars/hotttestfiles/Swan-2019-11-06-HoTTEST.pdf). Video recording at [https://www.](https://www.youtube.com/watch?v=r9KbE0zvr1g)
987 [youtube.com/watch?v=r9KbE0zvr1g](https://www.youtube.com/watch?v=r9KbE0zvr1g), 2019.

- 988 42 Taichi Uemura. Cubical Assemblies, a Univalent and Impredicative Universe and a Failure
989 of Propositional Resizing. In Peter Dybjer, José Espírito Santo, and Luís Pinto, editors,
990 *24th International Conference on Types for Proofs and Programs (TYPES 2018)*, volume
991 130 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 7:1–7:20. Schloss
992 Dagstuhl–Leibniz-Zentrum für Informatik, 2019. doi:10.4230/LIPIcs.TYPES.2018.7.
- 993 43 The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of*
994 *Mathematics*. <https://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.
- 995 44 Vladimir Voevodsky. Resizing rules — their use and semantic justification. Slides from a
996 talk at TYPES, Bergen, 11 September, 2011. URL: [https://www.math.ias.edu/vladimir/](https://www.math.ias.edu/vladimir/sites/math.ias.edu.vladimir/files/2011_Bergen.pdf)
997 [sites/math.ias.edu.vladimir/files/2011_Bergen.pdf](https://www.math.ias.edu.vladimir/files/2011_Bergen.pdf).
- 998 45 Vladimir Voevodsky. An experimental library of formalized mathematics based on the univalent
999 foundations. *Mathematical Structures in Computer Science*, 25(5):1278–1294, 2015.
- 1000 46 Vladimir Voevodsky, Benedikt Ahrens, Daniel Grayson, et al. UniMath — a computer-checked
1001 library of univalent mathematics. Available at <https://github.com/UniMath/UniMath>.