

Random Walks on Simple Two-Dimensional Manifolds

Tom Eichlersmith

Hamline University

`teichlersmith01@hamline.edu`

April 9, 2018

Overview

Introduction

Background

Method

Results

Questions

Random Walks

Tom Eichlersmith

Introduction

Background

Method

Results

Questions

- ▶ Random
- ▶ Walk
- ▶ Simple
- ▶ Two-Dimensional
- ▶ Manifolds

Coordinate Patch $\mu : U \rightarrow V$: continuous functions mapping from $U \subseteq \mathbb{R}^2$ to a subset of the surface V

Chart : covers entire surface

Regular Surfaces:

- ▶ Differentiable — the coordinate functions of μ in \mathbb{R}^3 have continuous partial derivatives for all orders
- ▶ Homeomorphic — μ and its inverse are continuous
- ▶ Satisfies the Regularity Condition — The differential of μ is a one-to-one linear transformation

$$\begin{aligned}\phi : \mathbb{R}^2 &\rightarrow P \\ \phi(u, v) &= (u, v, 0)\end{aligned}$$

$$\sigma : \mathbb{R}^2 \rightarrow S$$

$$\sigma(u, v) = \left(\frac{2u}{1 + u^2 + v^2}, \frac{2v}{1 + u^2 + v^2}, \frac{-1 + u^2 + v^2}{1 + u^2 + v^2} \right)$$

$$\begin{aligned}\tau : [0, 1) \times [0, 1) &\rightarrow T(R, r) \\ \tau(u, v) &= \left((R + r \cos(2\pi v)) \cos(2\pi u), \right. \\ &\quad (R + r \cos(2\pi v)) \sin(2\pi u), \\ &\quad \left. r \sin(2\pi v) \right)\end{aligned}$$

1. Extend definition of line to other surfaces
2. Assume a path is a geodesic contained in a coordinate patch
3. Derive geodesic equations for coordinate functions of path

Geodesic Equations

$$u'' + \frac{\mu_{uu} \cdot \mu_u}{\mu_u \cdot \mu_u} (u')^2 + \frac{\mu_{vv} \cdot \mu_u}{\mu_u \cdot \mu_u} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_u}{\mu_u \cdot \mu_u} u' v' = 0$$
$$v'' + \frac{\mu_{uu} \cdot \mu_v}{\mu_v \cdot \mu_v} (u')^2 + \frac{\mu_{vv} \cdot \mu_v}{\mu_v \cdot \mu_v} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_v}{\mu_v \cdot \mu_v} u' v' = 0$$

$$\Gamma_{jk}^i = \frac{\mu_{jk} \cdot \mu_i}{\mu_i \cdot \mu_i} \quad \text{where } i, j, k \in \{u, v\}$$

↓ abuse of symbols

$$\frac{d^2 x^i}{dt^2} + \sum_{j,k \in \{1,2\}} \Gamma_{jk}^i \frac{dx^j}{dt} \frac{dx^k}{dt} = 0$$

Stepping Method

Runge-Kutta 4th Order Method (RK4)

$$\frac{dy}{dt} = F(y) \quad y_0 = y(0)$$

Numerically solve up to $t = h$ with N iterations.

$$\delta \leftarrow h/N$$

$$y \leftarrow y_0$$

loop N times:

$$k_1 \leftarrow F(y)$$

$$k_2 \leftarrow F(y + (\delta/2)k_1)$$

$$k_3 \leftarrow F(y + (\delta/2)k_2)$$

$$k_4 \leftarrow F(y + \delta k_3)$$

$$y \leftarrow y + (\delta/6)(k_1 + 2k_2 + 2k_3 + k_4)$$

Tom Eichlersmith



- ▶ Collection of every step point
- ▶ Number of steps in RK4
- ▶ Simplifications due to symmetry
 - ▶ Plane with radius representation
 - ▶ Sphere with polar angle representation
- ▶ Method of “compressing” the data

Plane

Random Walks

Tom Eichlersmith


Introduction

Background


Method

Results

Questions



Plane



Sphere

Random Walks

Tom Eichlersmith


Introduction

Background


Method

Results

Questions



Sphere



Random Walks

Tom Eichlersmith


Introduction


Background

Method

Results

Questions





Overall Package

Package Attributes

- ▶ Versatility
- ▶ Flexibility
- ▶ Speed

Specific Parts

- ▶ Stepper
- ▶ Coordinate Wrappers
- ▶ Escape Checks

Questions?