

Random Walks on Simple Two-Dimensional Manifolds

Tom Eichlersmith

Hamline University

`teichlersmith01@hamline.edu`

April 19, 2018

Introduction

Random Walks

Tom Eichlersmith

Introduction

Background

Method

Results

Questions

- ▶ Random
- ▶ Walk
- ▶ Simple
- ▶ Two-Dimensional
- ▶ Manifolds

Regular Surfaces

Random Walks

Tom Eichlersmith

Introduction

Background

Method

Results

Questions




Figure: By Leonid_2 - Own work, CC BY-SA 3.0,
<https://commons.wikimedia.org/w/index.php?curid=8643414>

1. Extend definition of line to other surfaces
2. Assume a path is a geodesic contained in a coordinate patch
3. Derive geodesic equations for coordinate functions of path

Geodesic Equations

$$u'' + \frac{\mu_{uu} \cdot \mu_u}{\mu_u \cdot \mu_u} (u')^2 + \frac{\mu_{vv} \cdot \mu_u}{\mu_u \cdot \mu_u} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_u}{\mu_u \cdot \mu_u} u' v' = 0$$
$$v'' + \frac{\mu_{uu} \cdot \mu_v}{\mu_v \cdot \mu_v} (u')^2 + \frac{\mu_{vv} \cdot \mu_v}{\mu_v \cdot \mu_v} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_v}{\mu_v \cdot \mu_v} u' v' = 0$$

$$\Gamma_{jk}^i = \frac{\mu_{jk} \cdot \mu_i}{\mu_i \cdot \mu_i} \quad \text{where } i, j, k \in \{u, v\}$$

↓ abuse of symbols

$$\frac{d^2 x^i}{dt^2} + \sum_{j,k \in \{1,2\}} \Gamma_{jk}^i \frac{dx^j}{dt} \frac{dx^k}{dt} = 0$$

Stepping Method

Runge-Kutta 4th Order Method (RK4)

$$\frac{dy}{dt} = F(y) \quad y_0 = y(0)$$

Numerically solve up to $t = h$ with N iterations.

$$\delta \leftarrow h/N$$

$$y \leftarrow y_0$$

loop N times:

$$k_1 \leftarrow F(y)$$

$$k_2 \leftarrow F(y + (\delta/2)k_1)$$

$$k_3 \leftarrow F(y + (\delta/2)k_2)$$

$$k_4 \leftarrow F(y + \delta k_3)$$

$$y \leftarrow y + (\delta/6)(k_1 + 2k_2 + 2k_3 + k_4)$$

Stepping Method

Define

$$p = \frac{du}{dt} \quad \text{and} \quad q = \frac{dv}{dt}$$

Then the geodesic equations become

$$\frac{du}{dt} = p$$

$$\frac{dv}{dt} = q$$

$$\frac{dp}{dt} = -\Gamma_{uu}^u p^2 - 2\Gamma_{uv}^u pq - \Gamma_{vv}^u q^2$$


$$\frac{dq}{dt} = -\Gamma_{uu}^v p^2 - 2\Gamma_{uv}^v pq - \Gamma_{vv}^v q^2$$

Random Walks

Method



- ▶ Collection of every step point
- ▶ Number of steps in RK4
- ▶ Simplifications due to symmetry
 - ▶ Plane with radius representation
 - ▶ Sphere with polar angle representation
- ▶ Method of “compressing” the data



Plane

Random Walks

Tom Eichlersmith


Introduction

Background

Method

Results

Questions



Plane

Random Walks

Tom Eichlersmith


Introduction

Background


Method

Results


Questions



Plane



Sphere



Random Walks

Tom Eichlersmith

Introduction

Background

Method

Results

Questions

Sphere

Random Walks

Tom Eichlersmith


Introduction

Background


Method

Results

Questions



Sphere



Random Walks

Tom Eichlersmith

Introduction

Background

Method

Results

Questions

Questions?