

Random Walks on Simple Two-Dimensional Manifolds

Tom Eichlersmith

Hamline University

`teichlersmith01@hamline.edu`

April 19, 2018

- ▶ Random
- ▶ Walk
- ▶ Simple
- ▶ Two-Dimensional
- ▶ Manifolds
- ▶ Smooth Surfaces
- ▶ Geodesic Equations
- ▶ Christoffel Symbols

“Smooth” Surfaces

Random Walks

Tom Eichlersmith

Background

Method

Results

Questions




Figure: By Leonid_2 - Own work, CC BY-SA 3.0,
<https://commons.wikimedia.org/w/index.php?curid=8643414>

Geodesic Equations

Random Walks

Tom Eichlersmith

Background

Method

Results

Questions

1. Extend definition of line to other surfaces
2. Assume a path is a geodesic contained in a coordinate patch
3. Derive geodesic equations for coordinate functions of path

Christoffel Symbols

Random Walks

Tom Eichlersmith

Background


Method

Results

Questions

- ▶ Represent surface in geodesic equations
- ▶ Characterize properties of surface

- ▶ Runge-Kutta 4th Order Method
- ▶ Stack Linked List
- ▶ Function Pointers



Plane

Random Walks


Tom Eichlersmith

Background

Method

Results

Questions



Plane

Random Walks


Tom Eichlersmith

Background

Method

Results

Questions



Plane

Random Walks


Tom Eichlersmith

Background


Method

Results

Questions



Sphere



Random Walks

Tom Eichlersmith

Background

Method

Results

Questions

Sphere

Random Walks


Tom Eichlersmith

Background


Method

Results

Questions



Sphere



Random Walks

Tom Eichlersmith

Background

Method

Results

Questions

Questions?

Coordinate Patch $\mu : U \rightarrow V$: continuous functions mapping from $U \subseteq \mathbb{R}^2$ to a subset of the surface V

Chart : covers entire surface

Regular Surfaces:

- ▶ Differentiable — the coordinate functions of μ in \mathbb{R}^3 have continuous partial derivatives for all orders
- ▶ Homeomorphic — μ and its inverse are continuous
- ▶ Satisfies the Regularity Condition — The differential of μ is a one-to-one linear transformation

Overall Package

Random Walks

Tom Eichlersmith

Background

Method

Results

Questions

Package Attributes

- ▶ Versatility
- ▶ Flexibility
- ▶ Speed

Specific Parts

- ▶ Stepper
- ▶ Coordinate Wrappers
- ▶ Escape Checks

Geodesic Equations

Random Walks

Tom Eichlersmith

Background

Method

Results

Questions

$$u'' + \frac{\mu_{uu} \cdot \mu_u}{\mu_u \cdot \mu_u} (u')^2 + \frac{\mu_{vv} \cdot \mu_u}{\mu_u \cdot \mu_u} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_u}{\mu_u \cdot \mu_u} u' v' = 0$$
$$v'' + \frac{\mu_{uu} \cdot \mu_v}{\mu_v \cdot \mu_v} (u')^2 + \frac{\mu_{vv} \cdot \mu_v}{\mu_v \cdot \mu_v} (v')^2 + 2 \frac{\mu_{uv} \cdot \mu_v}{\mu_v \cdot \mu_v} u' v' = 0$$

$$\Gamma_{jk}^i = \frac{\mu_{jk} \cdot \mu_i}{\mu_i \cdot \mu_i} \quad \text{where } i, j, k \in \{u, v\}$$

↓ abuse of symbols

$$\frac{d^2 x^i}{dt^2} + \sum_{j,k \in \{1,2\}} \Gamma_{jk}^i \frac{dx^j}{dt} \frac{dx^k}{dt} = 0$$

Stepping Method

Runge-Kutta 4th Order Method (RK4)

$$\frac{dy}{dt} = F(y) \quad y_0 = y(0)$$

Numerically solve up to $t = h$ with N iterations.

$$\delta \leftarrow h/N$$

$$y \leftarrow y_0$$

loop N times:

$$k_1 \leftarrow F(y)$$

$$k_2 \leftarrow F(y + (\delta/2)k_1)$$

$$k_3 \leftarrow F(y + (\delta/2)k_2)$$

$$k_4 \leftarrow F(y + \delta k_3)$$

$$y \leftarrow y + (\delta/6)(k_1 + 2k_2 + 2k_3 + k_4)$$

Stepping Method

Define

$$p = \frac{du}{dt} \quad \text{and} \quad q = \frac{dv}{dt}$$

Then the geodesic equations become

$$\frac{du}{dt} = p$$

$$\frac{dv}{dt} = q$$

$$\frac{dp}{dt} = -\Gamma_{uu}^u p^2 - 2\Gamma_{uv}^u pq - \Gamma_{vv}^u q^2$$

$$\frac{dq}{dt} = -\Gamma_{uu}^v p^2 - 2\Gamma_{uv}^v pq - \Gamma_{vv}^v q^2$$

Random Walks

Questions



- ▶ Collection of every step point
- ▶ Number of steps in RK4
- ▶ Simplifications due to symmetry
 - ▶ Plane with radius representation
 - ▶ Sphere with polar angle representation
- ▶ Method of “compressing” the data

