#### The z-Transform

Digital Signal Processing

October 2, 2025



### **Motivation for the** *z***-Transform**

- Until now, we've used the Fourier transform to analyze frequency content of signals.
- With the z-transform, we'll analyze systems (specifically, LTI systems).
- Helps determine properties of a system, such as stability, causality, frequency response, etc.
- Used to design LTI systems (filters).

## **Review: Discrete Fourier Transform**

#### **Definition**

Let x[n] be a complex-valued, periodic signal with period L. The **discrete Fourier transform (DFT)** of x[n] is given by

DFT analysis:

$$X[k] = \frac{1}{\sqrt{L}} \sum_{n=0}^{L-1} e^{-i\omega_0 k n} x[n]$$

DFT synthesis:

$$x[n] = \frac{1}{\sqrt{L}} \sum_{k=0}^{L-1} e^{i\omega_0 k n} X[k]$$

### **Review: Discrete Fourier Transform**

#### **Definition**

Let x[n] be a complex-valued, periodic signal with period L. The **discrete Fourier transform (DFT)** of x[n] is given by


DFT analysis:

$$X[k] = \frac{1}{\sqrt{L}} \sum_{n=0}^{L-1} e^{-i\omega_0 kn} x[n]$$

DFT synthesis:

$$x[n] = \frac{1}{\sqrt{L}} \sum_{k=0}^{L-1} e^{i\omega_0 k n} X[k]$$

# **Transform Domains**



#### **Definition (DTFT Analysis)**

$$X(e^{i\omega}) = \sum_{n=-\infty}^{\infty} e^{-i\omega n} x[n], \quad \text{for } \omega \in [-\pi, \pi).$$

- Note range of n is all of  $\mathbb{Z}$ .
- Note  $X(e^{i\omega})$  is defined on the **continuous** unit circle in  $\mathbb{C}$ .
- Can also think of X as a function of angular frequency,  $\omega$ .

#### **Definition (DTFT Analysis)**

$$X(e^{i\omega}) = \sum_{n=-\infty}^{\infty} e^{-i\omega n} x[n], \quad \text{for } \omega \in [-\pi, \pi).$$

- Note range of n is all of  $\mathbb{Z}$ .
- Note  $X(e^{i\omega})$  is defined on the **continuous** unit circle in  $\mathbb{C}$ .
- Can also think of X as a function of angular frequency,  $\omega$ .

#### **Definition (DTFT Analysis)**

$$X(e^{i\omega}) = \sum_{n=-\infty}^{\infty} e^{-i\omega n} x[n], \quad \text{for } \omega \in [-\pi, \pi).$$

- Note range of n is all of  $\mathbb{Z}$ .
- Note  $X(e^{i\omega})$  is defined on the **continuous** unit circle in  $\mathbb{C}$ .
- Can also think of X as a function of angular frequency,  $\omega$ .

#### **Definition (DTFT Analysis)**

$$X(e^{i\omega}) = \sum_{n=-\infty}^{\infty} e^{-i\omega n} x[n], \quad \text{for } \omega \in [-\pi, \pi).$$


- Note range of n is all of  $\mathbb{Z}$ .
- Note  $X(e^{i\omega})$  is defined on the **continuous** unit circle in  $\mathbb{C}$ .
- Can also think of X as a function of angular frequency,  $\omega$ .

#### **Definition (DTFT Analysis)**

$$X(e^{i\omega}) = \sum_{n=-\infty}^{\infty} e^{-i\omega n} x[n], \quad \text{for } \omega \in [-\pi, \pi).$$

- Note range of n is all of  $\mathbb{Z}$ .
- Note  $X(e^{i\omega})$  is defined on the **continuous** unit circle in  $\mathbb{C}$ .
- Can also think of X as a function of angular frequency,  $\omega$ .

## **Transform Domains**



### **Inverse DTFT**

#### **Definition (DTFT Synthesis)**

Let  $X(e^{i\omega})$  be the DTFT of a signal x[n]. The **inverse** discrete-time Fourier transform is given by

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{i\omega}) e^{i\omega n} d\omega.$$

• Note this is an integral around the unit circle in  $\mathbb{C}$ .

#### The *z*-Transform

#### **Definition (**z**-Transform Analysis**)

Given a complex discrete signal x[n], its z-transform is given by

$$X(z) = \sum_{n = -\infty}^{\infty} z^{-n} x[n].$$


### The *z*-Transform

#### **Definition (**z**-Transform Analysis**)

Given a complex discrete signal x[n], its z-transform is given by

$$X(z) = \sum_{n = -\infty}^{\infty} \overline{z^{-n}} x[n].$$

## **Transform Domains**



### **Overloaded Notation!**

Notice we reused the notation "X" for the DFT, DTFT, and z-transform.

The context is clear from the input to X:

DFT	X[k]
DTFT	$X(e^{i\omega})$
z-transform	X(z)

## **Quick Side Note: Geometric Series**

A geometric series looks like:

$$s = 1 + r + r^2 + r^3 + \cdots$$
$$= \sum_{n=0}^{\infty} r^n$$

The infinite sum evaluates to

$$s = \frac{1}{1 - r},$$
 for  $|r| < 1$ .

This holds for r real or complex!

## **DTFT Example: Right-Sided Exponential**

Right-sided exponential signal, for some constant  $a \in \mathbb{C}$ , is:

$$x[n] = \begin{cases} a^n & \text{for } n \ge 0, \\ 0 & \text{for } n < 0, \end{cases}$$
$$= a^n u[n].$$


The DTFT is


$$\begin{split} X(e^{i\omega}) &= \sum_{n=-\infty}^{\infty} a^n u[n] e^{-i\omega n} = \sum_{n=0}^{\infty} (ae^{-i\omega})^n \\ &= \frac{1}{1-ae^{-i\omega}}, \qquad \text{for } |ae^{-i\omega}| < 1, \quad \text{or} \quad |a| < 1. \end{split}$$

Diverges for  $|a| \ge 1$ .

## **Plotting the DTFT**

$$x[n] = 0.5^n u[n]$$
  $\leftrightarrow$   $X(e^{i\omega}) = \frac{1}{1 - 0.5e^{-i\omega}}$ 





## z-Transform of Right-Sided Exponential

The *z*-transform of  $x[n] = a^n u[n]$  is

$$\begin{split} X(z) &= \sum_{n=-\infty}^{\infty} a^n u[n] z^{-n} = \sum_{n=0}^{\infty} (az^{-1})^n \\ &= \frac{1}{1-az^{-1}}, \qquad \text{for } |az^{-1}| < 1, \quad \text{or} \quad |z| > |a| \\ &= \frac{z}{z-a}. \end{split}$$

## **Region of Convergence (ROC)**

#### **Definition (ROC)**


The **region of convergence** for the z-transform of a signal x[n] is defined as all points  $z\in\mathbb{C}$  where

$$|X(z)| = \left| \sum_{n=-\infty}^{\infty} x[n]z^{-n} \right| < \infty.$$

## **ROC for Right-Sided Exponential**

For  $x[n] = u[n]a^n$ , we had

$$X(z) = \frac{z}{z - a}, \qquad \text{for } |z| > |a|.$$



## **Left-Sided Exponential**

Left-sided exponential for a constant  $a \in \mathbb{C}$  is

$$x[n] = \begin{cases} -a^n & \text{for } n < 0\\ 0 & \text{for } n \ge 0 \end{cases}$$
$$= -a^n u[-n-1]$$

The z-transform is

$$\begin{split} X(z) &= \sum_{n=-\infty}^{\infty} -a^n u[-n-1] z^{-n} = -\sum_{n=-\infty}^{-1} (az^{-1})^n \\ &= -\sum_{m=1}^{\infty} a^{-m} z^m = 1 - \sum_{m=0}^{\infty} (a^{-1}z)^m \\ &= 1 - \frac{1}{1 - a^{-1}z}, \qquad \text{for } |a^{-1}z| < 1, \quad \text{or } |z| < |a| \end{split}$$

# **Left-Sided Exponential**

#### Further simplifying:

$$\begin{split} X(z) &= 1 - \frac{1}{1 - a^{-1}z} \\ &= \frac{1 - a^{-1}z}{1 - a^{-1}z} - \frac{1}{1 - a^{-1}z} \\ &= \frac{-a^{-1}z}{1 - a^{-1}z} \\ &= \frac{z}{z - a}, \qquad \text{for } |z| < |a|. \end{split}$$


Same as right-sided exponential, but different ROC!

Right-sided ROC: |z| > |a|, Left-sided ROC: |z| < |a|


## **ROC for Left-Sided Exponential**

For  $x[n] = -a^n u[-n-1]$ , we had

$$X(z) = \frac{z}{z - a}, \qquad \text{for } |z| < |a|.$$



# **Quick Side Note: Triangle Inequality**



Complex Addition

Triangle inequality tells us:

$$|z_1 + z_2| \le |z_1| + |z_2|$$

Do this iteratively to get:

$$\left| \sum_{n=-\infty}^{\infty} z_n \right| \le \sum_{n=-\infty}^{\infty} |z_n|.$$

### **General ROC of** *z***-Transform**

In general, we have:

$$|X(z)| = \left| \sum_{n = -\infty}^{\infty} x[n]z^{-n} \right|$$

$$\leq \sum_{n = -\infty}^{\infty} |x[n]z^{-n}|$$

$$= \sum_{n = -\infty}^{\infty} |x[n]||z|^{-n}$$

Note this only depends on the magnitude  $r=\vert z\vert$ , not the angle of z.

### General ROC of z-Transform

#### ROC is an annulus:

$$0 \le r_R < |z| < r_L \le \infty$$

 $r_R$ : right-side radius

 $\sum_{n=0}^{\infty}|x[n]|r^{-n}$  diverges when  $r < r_R.$ 

 $r_L$ : left-side radius

$$\sum_{n=-\infty}^{-1}|x[n]|r^{-n}$$
 diverges when  $r>r_L$ .

