



ETT

mm	8th	9th
2.5	<1000 g <28 wk	≤1200 g ≤28 wk*
3.0	1000–2000 g 28–34 wk	1201–2200 g 29–34 wk
3.5	>2000 g >34 wk	>2200 g >34 wk

*2.0 mm optional <800 g (22–25 wk)

GA wk	cm
<23	5.0–5.5
23–24	5.5
25–26	6.0
27–29	6.5
30–32	7.0
33–34	7.5
35–37	8.0
38–40	8.5
41–43	9.0

Measure to:

9th upper gum

8th upper lip

Check: NTL + 1 cm

Tip: T1–T2

Flexion ↓ deeper

Extension ↑ higher



RSI + RESUS

RSI order **Unit**

1	Atropine 0.02 mg/kg
2	Fentanyl 1–4 mcg/kg Dilute in 2 mL NS over 3 min, flush 1 mL over 3 min
3	Rocuronium 0.6–1.2 mg/kg

Doses: MacDonald's Atlas. Confirm with order set.

Confirm ETT

HR ↑, chest rise, equal BS, CO₂ color change, CXR

Epi 0.1 mg/mL

IV/IO 0.2 mL/kg (0.1–0.3), flush 3 mL NS

ETT 0.5–1 mL/kg

Repeat every 3–5 min

Volume

NS 10 mL/kg over 5–10 min

Emergency UVC

2–4 cm below skin, to blood return



UMBILICAL LINES

UAC high T6–T9, low L3–L4

$(3 \times \text{kg}) + 9$

UVC IVC–RA junction

$[(3 \times \text{kg}) + 9] \div 2 + 1$

Revised: drop the +1

Add stump length. Confirm on film.

Cath (Atlas): UAC 3.5 Fr <1.2 kg, UVC 3.5 Fr <3.5 kg;
otherwise 5 Fr

kg	UAC cm	UVC cm
0.5	10.5	6.3
0.75	11.3	6.6
1	12	7
1.25	12.8	7.4
1.5	13.5	7.8
2	15	8.5
2.5	16.5	9.3
3	18	10
3.5	19.5	10.8
4	21	11.5



PICC + FILM

PICC measure

Arm/scalp: site → R sternal border, 3rd ICS
(arm 90°)

Leg: site → R of umbilicus → xiphoid

Tip targets on film

ETT	T1–T2
UAC	T6–T9 / L3–L4
UVC	IVC–RA jxn
PICC arm	SVC–RA jxn
PICC leg	IVC–RA jxn + lateral

9th Start FiO₂

≥35 wk	32–34 wk	<32 wk
21%	21–30%	≥30%

SpO₂ targets

2'	3'	4'	5'	10'
65–70	70–75	75–80	80–85	85–95