

The power rule

Given a monomial $f(x) = x^n$, where n is a real number,

$$f'(x) = nx^{n-1}$$

The antiderivative

The antiderivative is the opposite of the derivative.

Infinitely many antiderivatives map to the same function.

For example, since the derivative of x^2 is $2x$, we have x^2 as an antiderivative of $2x$. But $x^2 + 2$, $x^2 + 1$, etc. are also antiderivatives of $2x$.

In general, the antiderivatives of $2x$ follow the pattern $x^2 + C$ where C is a constant.

We can also formulate that using the \int symbol for integration:

$$\int 2x dx = x^2 + C$$

Where C is called the **constant of integration**.

These integrals are **indefinite integrals**.

The function being integrated (here $2x$) is called the **integrand**.

The power rule for integration

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

Exercises

1. Antiderivative of $\frac{1}{x^4}$?

$$\begin{aligned}\int \frac{1}{x^4} dx &= \int x^{-4} dx \\ &= \frac{x^{-4+1}}{-4+1} + C \\ &= \frac{x^{-3}}{-3} + C \\ &= -\frac{1}{3}x^3 + C\end{aligned}$$

2. Antiderivative of $x^{\frac{5}{8}}$

$$\begin{aligned}\int x^{\frac{5}{8}} dx &= \frac{x^{\frac{5}{8}+1}}{\frac{5}{8}+1} + C \\ &= \frac{x^{\frac{13}{8}}}{\frac{13}{8}} + C \\ &= \frac{8x^{\frac{13}{8}}}{13} + C\end{aligned}$$

3. Antiderivative of $\sqrt[5]{x^7}$

$$\begin{aligned}
 \int x^{\frac{7}{5}} dx &= \frac{x^{\frac{7}{5}+1}}{\frac{7}{5}+1} + C \\
 &= \frac{x^{12/5}}{12/5} \\
 &= \frac{5x^{12/5}}{12}
 \end{aligned}$$

The constant factor rule for indefinite integrals

To take the integral of a power function with a constant factor, like $5x^2$, we can take the constant factor out of the integral:

$$\int 5x^2 dx = 5 \cdot \int x^2 dx$$

This is the **constant factor rule**. In general:

$$\int kf(x)dx = k \int f(x)dx$$

Where k is a **constant**. It does not work if k is a variable.

Exercises

1. Calculate $\int \frac{1}{2y^5} dy$

$$\begin{aligned}
 \int \frac{1}{2y^5} dy &= \frac{1}{2} \int y^{-5} \\
 &= \frac{1}{2} \cdot \frac{y^{-5+1}}{-5+1} + C \\
 &= \frac{1}{2} \cdot \frac{y^{-4}}{-4} + C \\
 &= \frac{1}{2} \cdot -\frac{1}{4y^4} + C \\
 &= -\frac{1}{8y^4} + C
 \end{aligned}$$

2. Calculate $\int \frac{1}{3x^6} dx$

$$\begin{aligned}
 \int \frac{1}{3x^6} dx &= \frac{1}{3} \cdot \int x^{-6} \\
 &= \frac{1}{3} \cdot \frac{x^{-5}}{-5} + C \\
 &= \frac{1}{3} \cdot -\frac{1}{5x^5} + C \\
 &= -\frac{1}{15x^5} + C
 \end{aligned}$$

3. Calculate $\int \frac{\pi}{2} dz$

$$\begin{aligned}
\int \frac{\pi}{2} dz &= \frac{\pi}{2} \int z^0 dz \\
&= \frac{\pi}{2} \cdot z + C \\
&= \frac{\pi z}{2} + C
\end{aligned}$$