

NOTES

Contents

Combinatorics	2
Geometry	4
Logarithm	5
Falling objects	5
Geometric Sequences	5
Polynomials	7
Rationalizing the denominator	12
Partial fraction decomposition	16
Functions	17
Statistics	20
Graphing	21
Logarithms	22
Complex Numbers	22
Trigonometry	22
Calculus	29
Vectors	37

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aequale doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguere possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et collaudata est, cum id, quod maxime placeat, facere possimus, omnis voluptas assumenda est, omnis dolor repellendus. Temporibus autem quibusdam et aut officiis debitis aut rerum necessitatibus saepe eveniet, ut et voluptates repudiandae sint et molestiae non recusandae. Itaque earum rerum defuturum, quas natura non depravata desiderat. Et quem ad me accedis, saluto: 'chaere,' inquam, 'Tite!' lictores, turma omnis chorusque: 'chaere, Tite!' hinc hostis mi Albucius, hinc inimicus. Sed iure Mucius. Ego autem mirari satis non queo unde hoc sit tam insolens domesticarum rerum fastidium. Non est omnino hic docendi locus; sed ita prorsus existimo, neque eum Torquatum, qui hoc primus cognomen invenerit, aut torquem illum hosti detraxisse, ut aliquam ex eo est consecutus? – Laudem et caritatem, quae sunt vitae.

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aequale doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguere possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et collaudata est, cum id, quod maxime placeat, facere possimus, omnis voluptas assumenda est, omnis dolor repellendus. Temporibus autem quibusdam et.

abcdefghijklmnopqrstuvwxyz abcdefghijklmnopqrstuvwxyz

Combinatorics

Permutations

Sequences without repetition of n , written ${}_nP_k$, are ordered arrangements of k elements in an n elements set. The formula for the number of permutations is:

$$\frac{n!}{(n-k)!}$$

Combination

A *combination* is a selection of elements of a set that has distinct members, such that the order of the selection does not matter.

$$\binom{n}{k}$$

or

$${}_nC_k$$

The number of k -combinations of a set of cardinality n is given by:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Exercises

- A club has 10 female and 8 male members. Three members are to be chosen at random to represent the club. What is the probability that 2 women and one man will be chosen? Round your answer to the nearest percent.

First let's figure out how many ways to pick there are in total. The order doesn't matter, so we use a permutation:

$$N = {}_3C_{18} = \frac{18!}{(18-3)!3!} = 816$$

And how many ways to pick one male (m) or two women (w):

$$m = {}_1C_8 = \frac{8!}{(8-1)!} = 8$$

$$w = {}_2C_{10} = \frac{10!}{(10-2)!} = 45$$

Multiplying these two, we get the numbers of ways to pick one man and two women.

Our final result is:

$$\frac{mw}{N} = \frac{8 \cdot 45}{816} = 0.4411764...$$

- What is the probability that among 4 randomly chosen people whose birthdays are all in the last week of September, at least two will have birthdays on the same day? Round your answer to 3 decimal places.

It's easier to answer the question "what is the probability that all their birthdays are on a different day?" and take the complement.

We have 7 days to choose from, so there are ${}_7P_4 = 840$ permutations. There are 7^4 possibilities for all the birthday combinations. We have:

$$P(A') = \frac{n}{N} = \frac{840}{7^4} = 0.3498... P(A) = 1 - P(A') = 0.650...$$

- From a bag containing ten names, of which four are boy's names and six are girl's names, a magician must randomly choose two names to perform a magic trick. What is the probability that he will choose two boy's names?

First let's compute how many possibilities in total there are to choose two names from ten, where order does not matter (combinations):

$$\frac{10!}{2!(10-2)!} = 45$$

Now how many ways to pick two boy's names:

$$\frac{4!}{2!(4-2)!} = 6$$

So the probability we are looking for is $\frac{6}{45} = \frac{2}{15}$.

- From a box that contains 20 colored balls, a child randomly draws 2 black balls and one white ball. What is the probability of this event, given that the box contains 10 balls of each color?

First how many ways to pick 3 balls from 20, where order does not matter (combinations):

$$\frac{20!}{3!(20-3)!} = 1140$$

How many to pick 2 black balls out of 10:

$$\frac{10!}{2!(10-2)!} = 45$$

And one white ball out of 10:

$$\frac{10!}{1!(10-1)!} = 10$$

So $\frac{45 \cdot 10}{1140} \approx 0.39$

Geometry

Circles

Circumference: $2\pi r$

Surface: πr^2

Cylinders

Surface: $2\pi r^2 + 2\pi rh$

Volume: $V = \pi r^2 h$

Exercises

- Find an expression for the radius r of the base of a cylinder in terms of its volume V given that the ratio of the radius to the height is 3π .

We have $\frac{r}{3\pi} = h$. Let's substitute that in the volume formula:

$$\begin{aligned}\pi r^2 h &= V \\ \pi r^2 \frac{r}{3\pi} &= V \\ \frac{\pi r^3}{3} \pi &= V \\ \frac{r^3}{3} &= V \\ r^3 &= 3V \\ r &= \sqrt[3]{3V}\end{aligned}$$

- Calculate the circumference of the base, given volume is $625\pi m^2$ and the ratio of its height to its radius is 5.

$$\begin{aligned}\pi r^2 h &= V \\ \pi r^2 \cdot (5r) &= 625\pi \\ 5r^3 &= 625 \\ r^3 &= 125 \\ r &= 5\end{aligned}$$

Now the circumference is $2\pi r = 2\pi(5) = 10\pi$

Spheres

Surface: $4\pi r^2$ **Volume:** $\frac{4}{3}\pi r^3$

Triangles

Surface

The formula for the area of a triangle from one angle C and its two adjacent sides a and b is:

$$\frac{1}{2}ab \sin C$$

C is called the **included angle**.

Polygons

The sum of the interior angles of a polygon is always $180 * (n - 2)$ where n is the number of sides.

Regular polygons

With n as the number of sides, exterior angles are given by:

$$\theta = \frac{360}{n}$$

Logarithm

Domain

$(0, \infty)$

Range

$(-\infty, \infty)$

Falling objects

The *displacement* equation:

$$s = ut + \frac{1}{2}gt^2$$

where:

- s is the **distance** travelled in meters
- u is the **initial velocity** in m/s
- t is the **time** in seconds
- g is the **acceleration due to gravity**, constant equal to $9.8m/s^2$

Geometric Sequences

Geometric sequences are of the form $a_n = a_1 \cdot r^{n-1}$, where r is called the *common ratio*.

Exercises

1. Given that a geometric sequence has the first term $a_1 = -2$ and the 4th term $a_4 = -54$, what is the index of the term -6 ?

The formula for a_n is $a_n = b \cdot r^{n-1}$. Since $a_1 = b$, we have:

$$\begin{aligned}
 a_4 &= -2 \cdot r^3 \\
 -54 &= -2 \cdot r^3 \\
 27 &= r^3 \\
 r &= \sqrt[3]{27} \\
 r &= 3
 \end{aligned}$$

So the formula is $a_n = -2 \cdot 3^{n-1}$. Now we are looking for the index. We substitute -6 for a_n :

$$\begin{aligned}
 -6 &= -2 \cdot 3^{n-1} \\
 3 &= 3^{n-1} \\
 n &= 2
 \end{aligned}$$

2. First term: $a_1 = 81$, common ratio $r = \frac{1}{3}$, and $a_n = 1$. Find n .

$$\begin{aligned}
 1 &= 81 \cdot \left(\frac{1}{3}\right)^{n-1} \\
 \frac{1}{81} &= \left(\frac{1}{3}\right)^{n-1} \\
 \left(\frac{1}{3}\right)^4 &= \left(\frac{1}{3}\right)^{n-1} \\
 4 &= n - 1 \\
 n &= 5
 \end{aligned}$$

3. Given $a_3 = 36$, $r = 3$, what is the index of the term 108?

First let's find the formula:

$$\begin{aligned}
 36 &= a_1 \cdot 3^{3-1} \\
 36 &= a_1 \cdot 9 \\
 a_1 &= 4
 \end{aligned}$$

So $a_n = 4 \cdot 3^{n-1}$. Now,

$$\begin{aligned}
 108 &= 4 \cdot 3^{n-1} \\
 27 &= 3^{n-1} \\
 3^3 &= 3^{n-1} \\
 3 &= n - 1 \\
 n &= 4
 \end{aligned}$$

Polynomials

Forms

- Standard form: $y = ax^2 + bx + c$
- Vertex form: $y = a(x - h)^2 + k$
 - In this form, the coordinates of the vertex are (h, k)
- Factored form: $y = a(x - x_1)(x - x_2)$
- Slope-intercept form: $y = mx + b$
 - In this form, the slope is m and the y-intercept is b

Exercises

- Find the equation of the line that passes through the point $(1, 3)$ and is parallel to the line passing through the points $(1, -3)$ and $(-1, 7)$.

First let's find the slope of the line:

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - (-3)}{-1 - 1} = \frac{10}{-2} = -5$$

Now the equation of the line passing through $(1, 3)$ with slope -5 , using point-slope form:

$$\begin{aligned}y - 3 &= -5(x - 1) \\y &= -5x + 5 + 3 \\y &= -5x + 8\end{aligned}$$

- Find the equation of the line that passes through $(2, 0)$ and is parallel to the line on $(0, 0)$ and $(2, 1)$.

First the slope: $\frac{y_2 - y_1}{x_2 - x_1} = \frac{1}{2}$.

Then, using point-slope form:

$$\begin{aligned}y - 0 &= \frac{1}{2}(x - 2) \\y &= \frac{1}{2}x - 1\end{aligned}$$

- Same. Point: $(0, 0)$. Line: $(0, -2)$, $(-2, 0)$.

Slope: $\frac{2}{-2} = -1$.

$$y = -x$$

- Same. Point: $(6, 4)$. Line: $(2, -2)$, $(4, -1)$.

Slope: $\frac{-1 - (-2)}{4 - 2} = \frac{1}{2}$.

Point-slope form:

$$\begin{aligned}y - 4 &= \frac{1}{2}(x - 6) \\y &= \frac{1}{2}x - 3 + 4 \\y &= \frac{1}{2}x + 1\end{aligned}$$

- Point: $(-3, 2)$. Line: $(5, -3)$, $(6, -5)$.

Slope: $\frac{-5 - (-3)}{6 - (-3)} = -\frac{2}{9} = -\frac{2}{9}$.

$$\begin{aligned}
 y - 2 &= -2(x + 3) \\
 y &= -2x - 6 + 2 \\
 y &= -2x - 4
 \end{aligned}$$

Vertex

If the binomial is given in standard form, the vertex' x coordinate is given by $-\frac{b}{2a}$. Once you have the x coordinate, substitute in the equation to find y .

Domain and range

The **domain** is always $(-\infty, +\infty)$.

The **range** is, for upward opening parabolas, assuming vertex form: $y = a(x - h)^2 + k$, the range is $[k, +\infty)$.

Factor theorem

Theorem. $(x - r)$ is a factor of a polynomial $p(x)$ if and only if the root $x = r$ is also a root of $p(x)$.

Exercises

- The polynomial $-5x^4 + 8x^3 + 29x^2 - 20x - 12$ can be expressed as $-(x - 1)(x - 3) \cdot P(x)$. Find $P(x)$.

First let's divide by $(x - 1)$

$$\begin{array}{r|rrrrrr}
 1 & -5 & 8 & 29 & -20 & -12 \\
 & & -5 & 3 & 32 & 12 \\
 \hline
 & -5 & 3 & 32 & 12 & 0
 \end{array}$$

So we have $(x - 1) \cdot (-5x^3 + 3x^2 + 32x + 12)$

Let's divide by $(x - 3)$

$$\begin{array}{r|rrrr}
 3 & -5 & 3 & 32 & 12 \\
 & & -15 & -36 & -12 \\
 \hline
 & -5 & -12 & -4 & 0
 \end{array}$$

So we have $(x - 1)(x - 3)(-5x^2 - 12x - 4)$

So we have $(x - 1) \cdot (x - 3) \cdot (-5x^2 - 2x + 4)$

Finally, $-(x - 1)(x - 3)(5x^2 + 2x + 4)$. So $P(x) = 5x^2 + 2x + 4$.

Factoring further, $P(x) = (5x + 2)(x + 2)$.

- Find the polynomial $P(x)$ given that

$$4x^4 - 19x^3 - 9x^2 + 19x + 5 = (x - 1)(x - 5)P(x)$$

First let's divide by $(x - 1)$.

$$\begin{array}{r|rrrrr}
 1 & 4 & -19 & -9 & 19 & 5 \\
 & & 4 & -15 & -24 & -5 \\
 \hline
 & 4 & -15 & -24 & -5 & 0
 \end{array}$$

So $(x - 1)(4x^3 - 15x^2 - 24x - 5)$. Now divide by $(x - 5)$.

$$\begin{array}{r|rrrr}
 5 & 4 & -15 & -24 & -5 \\
 & & 20 & 25 & 5 \\
 \hline
 & 4 & 5 & 1 & 0
 \end{array}$$

So $(x - 1)(x - 5)(4x^2 + 5x + 1)$.

- Find $P(x)$ given $4x^4 - 8x^3 - 3x^2 + 7x - 2 = (x - 2)(x + 1)P(x)$.

First divide by $(x + 1)$:

$$\begin{array}{r|rrrrr}
 -1 & 4 & -8 & -3 & 7 & -2 \\
 & & -4 & 12 & -9 & 2 \\
 \hline
 & 4 & -12 & 9 & -2 & 0
 \end{array}$$

So $(x + 1)(4x^3 - 12x^2 + 9x - 2)$. Now divide the remainder by $(x - 2)$:

$$\begin{array}{r|rrrr}
 2 & 4 & -12 & 9 & -2 \\
 & & 8 & -8 & 2 \\
 \hline
 & 4 & -4 & 1 & 0
 \end{array}$$

So $(x + 1)(x - 2)(4x^2 - 4x + 1)$. So $P(x) = (4x^2 - 4x + 1)$. Factoring further,

$$\begin{aligned}
 P(x) &= 4x^2 - 4x + 1 \\
 &= (4x^2 - 2x) + (-2x + 1) \\
 &= 2x(2x - 1) - 1(2x - 1) \\
 &= (2x - 1)^2
 \end{aligned}$$

- Let $p(x) = -6x^4 + 5x^3 + 52x^2 - 19x - 60$. Given that $p(-1) = p(3) = 0$, find the sum of all distinct roots of $p(x)$.

$p(-1) = 0$ means $x = -1$ is a root, so $(x + 1)$ is a factor.

$p(3) = 0$ means $x = 3$ is a root, so $(x - 3)$ is a factor.

Let's first divide by $(x + 1)$.

$$\begin{array}{r|rrrrr}
 -1 & -6 & 5 & 52 & -19 & -60 \\
 & & 6 & -11 & -41 & 60 \\
 \hline
 & -6 & 11 & 41 & -60 & 0
 \end{array}$$

So $p(x) = (x + 1)(-6x^3 + 11x^2 + 41x - 60)$. Now let's divide the remainder by $(x - 3)$.

$$\begin{array}{r|rrrr}
 3 & -6 & 11 & 41 & -60 \\
 & & -18 & -21 & 60 \\
 \hline
 & -6 & -7 & 20 & 0
 \end{array}$$

So $p(x) = (x + 1)(x - 3)(-6x^2 - 7x + 20)$.

Factoring the remainder further:

$$\begin{aligned}
 p(x) &= (x + 1)(x - 3)(-6x^2 - 7x + 20) \\
 &= (x + 1)(x - 3)(-6x^2 + 8x) + (-15x + 20) \\
 &= (x + 1)(x - 3)(2x + 5)(-3x + 4)
 \end{aligned}$$

The roots are $x = -1, 3, -\frac{5}{2}, \frac{4}{3}$.

The number we are looking for is $-1 + 3 - \frac{5}{2} + \frac{4}{3} = 2 - \frac{5}{2} + \frac{4}{3} = -\frac{1}{2} + \frac{4}{3} = \frac{5}{6}$.

- Find the distinct roots of $x^4 - 5x^2 + 4$ given $p(1) = p(-1) = 0$.

$$\begin{array}{c|ccccc} 1 & 1 & 0 & -5 & 0 & 4 \\ & & 1 & 1 & -4 & -4 \\ & 1 & 1 & -4 & -4 & 0 \end{array}$$
$$\begin{array}{c|cccc} -1 & 1 & 1 & -4 & -4 \\ & & -1 & 0 & 4 \\ & 1 & 0 & -4 & 0 \end{array}$$

- Find $P(x)$ given $x^4 - 2x^3 - 7x^2 + 18x - 18 = (x - 3)(x + 3)P(x)$.

$$\begin{array}{c|cccc} 3 & 1 & -2 & -7 & 18 & -18 \\ & & 3 & 3 & -12 & 18 \\ & 1 & 1 & -4 & 6 & 0 \end{array}$$
$$\begin{array}{c|cccc} -3 & 1 & 1 & -4 & 6 \\ & & -3 & 6 & -6 \\ & 1 & -2 & 2 & 0 \end{array}$$

- Find $P(x)$ given that $x^4 + x^3 - 36x^2 + 37x - 35 = (x - 5)(x + 7)P(x)$.

$$\begin{array}{r|rrrrr} 5 & 1 & 1 & -36 & 37 & -35 \\ & & 5 & 30 & -30 & 35 \\ & & 1 & 6 & -6 & 7 & 0 \end{array}$$
$$\begin{array}{c|cccc} -7 & 1 & 6 & -6 & 7 \\ & & -7 & 7 & -7 \\ & 1 & -1 & 1 & 0 \end{array}$$

Polynomial long division

- What is the remainder of $3x^2 - 2x^2 - 18x - 1$ divided by $3x^2 - 8x - 2$?

$$\begin{array}{r}
 3x^2 - 8x - 2 \mid \begin{array}{rrr}
 x & +2 & \\
 3 & -2 & -18 & -1 \\
 3 & -8 & -2 & \\
 \hline
 & 6 & -16 & \\
 & 6 & -16 & -4 \\
 & & & \hline
 & & & 3
 \end{array}
 \end{array}$$

- What is the remainder when $x^4 - 2x^3 - 4x^2 - 10x - 7$ is divided by $x^2 + 2x + 1$?

$$\begin{array}{r}
 x^2 - 4x + 3 \\
 x^2 + 2x + 1 \mid 1 \quad -2 \quad -4 \quad -10 \quad -7 \\
 \underline{1 \quad 2 \quad 1} \\
 -4 \quad -5 \\
 \underline{-4 \quad -8 \quad -4} \\
 3 \quad -6 \\
 \underline{3 \quad 6 \quad 3} \\
 -12x \quad -10
 \end{array}$$

The remainder is $-12x - 10$.

- What is the quotient of $2x^4 - x^2 - 1$ divided by $2x^2 + 1$?

$$\begin{array}{r}
 x^2 - 1 \\
 2x^2 + 1 \mid 2 \quad 0 \quad -1 \quad 0 \quad -1 \\
 \underline{2 \quad 0 \quad 1} \\
 -2 \\
 \underline{-2 \quad 0 \quad 1} \\
 0
 \end{array}$$

So it's $x^2 - 1$.

Rational roots theorem

The rational roots theorem (a.k.a. rational zeros theorem) gives the rational roots of a polynomial with integer coefficients.

Theorem. Suppose that $\frac{p}{q} \in \mathbb{Q}$, in lowest terms, is a root of a polynomial with integer coefficients. Then p must be a factor of the polynomial's constant term, and q must be a factor of its leading coefficient.

That means every root must take the form $\pm \frac{\text{factor of the constant term}}{\text{factor of the leading coefficient}}$.

Exercises

- For the polynomial $f(x) = 3x^8 - 18x^2 - 33x - 18$, how many potential roots are given by the rational roots theorem?

The numerator must be a factor of -18 . The denominator must be a factor of 3. So here are the possibilities:

$$\pm \frac{1}{3}, \pm \frac{2}{3}, \pm 1, \pm 2, \pm 3, \pm 6, \pm 9, \pm 18$$

So 16.

- Given that the polynomial $2x^3 + x^2 + 5x - 3$ has a root in the interval $(0, \frac{3}{2})$, find the sum of the distinct real roots of $f(x)$.

Using the rational roots theorem, we can find the potential roots of the polynomial. The numerator must be a factor of -3 , and the denominator must be a factor of 2. So here are the possibilities in the interval $(0, \frac{3}{2})$:

$$\frac{1}{2}, 1$$

We can quickly find which one of these roots by evaluating the polynomial at these points. It's $\frac{1}{2}$.

Now let's use synthetic division to find the remaining roots:

$$\begin{array}{r|rrrr} 1/2 & 2 & 1 & 5 & -3 \\ & & 1 & 1 & 3 \\ \hline & 2 & 2 & 6 & 0 \end{array}$$

So $2x^3 + x^2 + 5x - 3 = \left(x - \frac{1}{2}\right)(2x^2 + 2x + 6)$. Since the discriminant of $2x^2 + 2x + 6$ is negative, it has no real roots. Therefore, the only real root is $x = \frac{1}{2}$.

Rationalizing the denominator

In general, you want to avoid radicals in the denominator. To rationalize the denominator, multiply both the numerator and the denominator by the conjugate of the denominator.

$$\begin{aligned} & \frac{1}{1 + \sqrt{2}} \\ &= \frac{1}{1 + \sqrt{2}} \cdot \frac{1 - \sqrt{2}}{1 - \sqrt{2}} \\ &= \frac{1 - \sqrt{2}}{(1 + \sqrt{2})(1 - \sqrt{2})} \\ &= \frac{1 - \sqrt{2}}{1^2 - \sqrt{2}^2} \\ &= \frac{1 - \sqrt{2}}{1 - 2} \\ &= \frac{1 - \sqrt{2}}{-1} \\ &= -1 + \sqrt{2} \end{aligned}$$

Exercises

•

$$\begin{aligned}
& \frac{1}{1 + \sqrt{2}} \\
&= \frac{1}{1 + \sqrt{2}} \cdot \frac{1 - \sqrt{2}}{1 - \sqrt{2}} \\
&= \frac{1 - \sqrt{2}}{(1 + \sqrt{2})(1 - \sqrt{2})} \\
&= \frac{1 - \sqrt{2}}{1^2 - \sqrt{2}^2} \\
&= \frac{1 - \sqrt{2}}{1 - 2} \\
&= \frac{1 - \sqrt{2}}{-1} \\
&= -1 + \sqrt{2}
\end{aligned}$$

1.

$$\begin{aligned}
& \frac{1 + \sqrt{8}}{\sqrt{9} - \sqrt{2}} \\
&= \frac{(1 + \sqrt{8})(\sqrt{9} + \sqrt{2})}{7} \\
&= \frac{\sqrt{9} + 3\sqrt{8} + \sqrt{2} + \sqrt{16}}{7} \\
&= \frac{3 + 6\sqrt{2} + \sqrt{2} + 4}{7} \\
&= \frac{7 + 7\sqrt{2}}{7} \\
&= 1 + \sqrt{2}
\end{aligned}$$

2.

$$\begin{aligned}
\frac{x^2 y^2}{\sqrt{x^3 y^3}} &= \frac{x^2 y^2 \sqrt{x^3 y^3}}{x^3 y^3} \\
&= \frac{x^2 y^2 x^{3/2} y^{3/2}}{x^3 y^3} \\
&= \frac{x^{7/2} y^{7/2}}{x^3 y^3} \\
&= \sqrt{xy}
\end{aligned}$$

3.

$$\begin{aligned}
\frac{\sqrt{12x}}{3\sqrt{2x^2y}} &= \frac{\sqrt{12x} \cdot \sqrt{2x^2y}}{3 \cdot 2x^2y} \\
&= \frac{\sqrt{24x^3y}}{6x^2y} \\
&= \frac{\sqrt{4x^2 \cdot 6xy}}{6x^2y} \\
&= \frac{2x \cdot \sqrt{6xy}}{6x^2y} \\
&= \frac{\sqrt{6xy}}{3xy}
\end{aligned}$$

4. Given that $d > 0$.

$$\begin{aligned}
\frac{cd}{\sqrt{cd^3}} &= \frac{cd\sqrt{cd^3}}{cd^3} \\
&= \frac{\sqrt{cd^3}}{d^2} \\
&= \frac{\sqrt{c} \cdot d^{3/2}}{d^2} \\
&= \frac{\sqrt{c} \cdot d^{1/2}}{d} \\
&= \frac{\sqrt{cd}}{d}
\end{aligned}$$

5. Given that $p > 0$

$$\begin{aligned}
\frac{7t}{\sqrt{14tp^2}} &= \frac{7t \cdot \sqrt{14tp^2}}{14tp^2} \\
&= \frac{7tp\sqrt{14t}}{14tp^2} \\
&= \frac{\sqrt{14t}}{2p}
\end{aligned}$$

6. Given that $a > 0$

$$\begin{aligned}
\sqrt{\frac{5b}{a^3}} &= \frac{\sqrt{5b}}{\sqrt{a^3}} \\
&= \frac{\sqrt{5b}}{a\sqrt{a}} \\
&= \frac{\sqrt{5ab}}{a\sqrt{a}^2} \\
&= \frac{\sqrt{5ab}}{a^2}
\end{aligned}$$

7.

$$\begin{aligned}\sqrt{\frac{14r}{p^5}} &= \frac{\sqrt{14r}}{\sqrt{p^5}} \\ &= \frac{\sqrt{14p^5r}}{p^5} \\ &= \frac{\sqrt{14r} \cdot p^{5/2}}{p^5} \\ &= \frac{\sqrt{14pr}}{p^3}\end{aligned}$$

8.

$$\begin{aligned}\sqrt{\frac{9a^2}{b^3}} &= \frac{\sqrt{9a^2}}{\sqrt{b^3}} \\ &= \frac{\sqrt{9a^2}}{b\sqrt{b}} \\ &= \frac{\sqrt{9a^2b}}{b\sqrt{b}^2} \\ &= \frac{3a\sqrt{b}}{b^2}\end{aligned}$$

9. Given that $z > 0$.

$$\begin{aligned}\sqrt{\frac{11}{wz^2}} &= \frac{\sqrt{11}}{\sqrt{wz^2}} \zeta = \frac{\sqrt{11}}{z\sqrt{w}} \\ &= \frac{\sqrt{11w}}{zw}\end{aligned}$$

10.

$$\begin{aligned}\sqrt{\frac{5a}{3b}} &= \frac{\sqrt{5a}}{\sqrt{3b}} \\ &= \frac{\sqrt{15ab}}{3b}\end{aligned}$$

11.

$$\begin{aligned}
\sqrt{\frac{a^2+3}{a^2-3}} &= \frac{\sqrt{a^2+3}}{\sqrt{a^2-3}} \\
&= \frac{\sqrt{(a^2+3)(a^2-3)}}{a^2-3} \\
&= \frac{\sqrt{a^4-9}}{a^2-3} \\
&= \frac{\sqrt{a^4-9}}{a^2-3}
\end{aligned}$$

12. Given $y > 0$

$$\begin{aligned}
\frac{2+3\sqrt{xy}}{3\sqrt{y}} &= \frac{(2+3\sqrt{xy}) \cdot \sqrt{y}}{3y} \\
&= \frac{3y\sqrt{x}+2\sqrt{y}}{3y} \\
&= \sqrt{x} + \frac{2\sqrt{y}}{3y}
\end{aligned}$$

Partial fraction decomposition

We have something of the form:

$$\frac{1}{(x-1)(3x-2)}$$

and we want to decompose it into partial fractions:

$$\frac{1}{(x-1)(3x-2)} = \frac{A}{x-1} + \frac{B}{3x-2}$$

Multiplying both sides by $(x-1)(3x-2)$, we get:

$$1 = A(3x-2) + B(x-1)$$

Setting $x = 1$, we get:

$$1 = A(3(1)-2) + B(1-1) = A$$

Setting $x = \frac{2}{3}$, we get:

$$1 = A\left(3\left(\frac{2}{3}\right)-2\right) + B\left(\frac{2}{3}-1\right) = -\frac{B}{3}$$

Solving for B , we get:

$$B = -3$$

So the partial fraction decomposition is:

$$\frac{1}{(x-1)(3x-2)} = \frac{1}{x-1} - \frac{3}{3x-2}$$

Exercises

1. Decompose $\frac{44x + 48}{(x + 6)(x - 3)(x + 2)}$

$$\frac{44x + 48}{(x + 6)(x - 3)(x + 2)} = \frac{A}{x + 6} + \frac{B}{x - 3} + \frac{C}{x + 2}$$

$$44x + 48 = A(x - 3)(x + 2) + B(x + 6)(x + 2) + C(x + 6)(x - 3)$$

Set $x = 3$:

$$44 \cdot 3 + 48 = B \cdot 9 \cdot 5$$

$$\frac{180}{9 \cdot 5} = B$$

$$B = 4$$

Set $x = -2$:

$$44 \cdot -2 + 48 = C \cdot 4 \cdot -5$$

$$-40 = C \cdot -20$$

$$C = 2$$

Set $x = -6$:

$$44 \cdot -6 + 48 = A \cdot -9 \cdot -4$$

$$A = -6$$

2. Given that

$$\frac{-2x - 9}{x + 3}x = \frac{1}{x + 3} + \frac{B}{x}$$

find B

$$\frac{-2x - 9}{x + 3}x = \frac{1}{x + 3} + \frac{B}{x}$$

$$(-2x - 9) = x + B(x + 3)$$

$$-3x - 9 = B(x + 3)$$

$$\frac{-3(x + 3)}{x + 3} = B$$

$$B = -3$$

3. Given that

$$\frac{-9x + 36}{(x - 3)(x - 6)} = -\frac{3}{x - 3} + \frac{B}{x - 6}$$

$$-9x + 36 = -3(x - 6) + B(x - 3)$$

$$-6x + 18 = B(x - 3)$$

$$\frac{-6(x - 3)}{x - 3} = B$$

$$B = -6$$

Functions

Horizontal asymptotes of rational functions

Rational functions are of the form:

$$f(x) = \frac{ax^n + bx^{n-1} + \dots}{cx^m + dx^{m-1} + \dots}$$

To get the horizontal asymptotes, first take the **dominant term**. It's the leading term of the polynomial in the denominator, *without the coefficient*. Here x^m .

Then, divide every term in both numerator and denominator by the dominant term, and evaluate at $x \rightarrow \infty$.

Exercises

- Determine the horizontal asymptote of $f(x) = \frac{x^3+2x+1}{x^2-x-12}$

The dominant term in x^2 . Let's divide by it.

$$\frac{\frac{x^3+2x+1}{x^2}}{\frac{x^2-x-12}{x^2}} = \frac{\left(x + \frac{2}{x} + \frac{1}{x^2}\right)}{1 - \frac{x+12}{x^2}}$$

As $x \rightarrow \infty$, this becomes $\frac{x}{1} = x$. Since this is not a constant, there is no horizontal asymptote.

- Find the horizontal asymptote of $\frac{3x^2+3}{6x^2+x}$.

The dominant term is x^2 .

$$\frac{\frac{3x^2+3}{x^2}}{\frac{6x^2+x}{x^2}} = \frac{3 + \frac{3}{x^2}}{6 + \frac{1}{x}}$$

As $x \rightarrow \infty$, this becomes $\frac{3}{6} = \frac{1}{2}$. The horizontal asymptote is $y = \frac{1}{2}$.

- Find the horizontal asymptote of $f(x) = \frac{3x^2-2}{2x^2+1}$.

The dominant term is x^2 .

$$\frac{\frac{3x^2-2}{x^2}}{\frac{2x^2+1}{x^2}} = \frac{3 - \frac{2}{x^2}}{2 + \frac{1}{x^2}}$$

As $x \rightarrow \infty$, this becomes $\frac{3}{2}$. The horizontal asymptote is $y = \frac{3}{2}$.

Vertical asymptotes of rational functions

1. Factor numerator and denominator.
2. Cancel out common factors.
3. Set the denominator equal to zero and solve for x .

Exercises

- Vertical asymptotes of $\frac{4x}{2x^3+5x^2+2x}$?

$$\begin{aligned}
\frac{4x}{2x^3 + 5x^2 + 2x} &= \frac{4x}{x(2x^2 + 5x + 2)} \\
&= \frac{4}{2x^2 + 5x + 2} \\
&= \frac{4}{2x^2 + 4x + x + 2} \\
&= \frac{4}{2x(x + 2) + 1(x + 2)} \\
&= \frac{4}{((2x + 1)(x + 2))}
\end{aligned}$$

So the vertical asymptotes are $x = -\frac{1}{2}$ and $x = -2$.

- Vertical asymptotes of $f(x) = \frac{x^2 - x - 2}{x^3 + 7x^2 - x - 7}$.

$$\begin{aligned}
\frac{x^2 - x - 2}{x^3 + 7x^2 - x - 7} &= \frac{x^2 - 2x + x - 2}{x^2(x + 7) - 1(x + 7)} \\
&= \frac{x(x - 2) + 1(x - 2)}{(x - 1)(x + 1)(x + 7)} \\
&= \frac{(x + 1)(x - 2)}{(x + 1)(x - 1)(x + 7)} \\
&= \frac{x - 2}{(x - 1)(x + 7)}
\end{aligned}$$

So the vertical asymptotes are $x = 1$ and $x = -7$.

- Vertical asymptotes of $f(x) = \frac{2x^2 - 10x}{x^2 + 2x - 8}$.

$$\begin{aligned}
\frac{2x^2 - 10x}{x^2 + 2x - 8} &= \frac{2x(x - 5)}{x^2 - 2x + 4x - 8} \\
&= \frac{2x(x - 5)}{(x + 4)(x - 2)}
\end{aligned}$$

Domain of radical functions

All the values where the radicand > 0 for even degrees, $(-\infty, \infty)$ for odd degrees.

Exercises

1. Domain of $d(x) = \sqrt[7]{-2x + \frac{1}{2}} + 1$?

$$-2x + \frac{1}{2} >$$

Roots of rational functions

First factor numerator and denominator, then cancel out common factors, then set the numerator to zero and solve.

Exercises

1. Solve $f(t) = 0$ where $f(t) = \frac{7t^2 + 14}{t^2 - 25}$.

$$\frac{7t^2 + 14}{t^2 - 25} = 0$$

$$\frac{7(t^2 + 2)}{(t + 5)(t - 5)} = 0$$

$$t^2 + 2 = 0$$

$$t^2 = -2$$

2. Solve $f(x) = 0$ where $f(x) = \frac{x^2 + 2x - 15}{x^3 + 5x^2 - 9x - 45}$.

$$\frac{x^2 + 2x - 15}{x^3 + 5x^2 - 9x - 45} = 0$$

$$\frac{x^2 + 5x - 3x - 15}{x^2(x + 5) - 9(x + 5)} = 0$$

$$\frac{(x + 5)(x - 3)}{(x + 5)(x^2 - 9)} = 0$$

$$\frac{(x + 5)(x - 3)}{(x + 5)(x - 3)(x + 3)} = 0$$

$$1 = 0$$

The equation has no solution.

3. Solve $f(t) = 0$ where $f(t) = \frac{t-3}{t^2+6t+5}$

$$\frac{t-3}{t^2+6t+5} = 0$$

$$\frac{t-3}{t^2+t+5t+5} = 0$$

$$\frac{t-3}{(t+5)(t+1)} = 0$$

$$t-3 = 0$$

$$t = 3$$

Invertible functions

A function is invertible on an interval if and only if it is one-to-one on that interval.

To check if a function is one-to-one, we can use the horizontal line test. If any horizontal line intersects the graph of the function at most once, then the function is one-to-one.

Statistics

Variance

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

Covariance

Covariance is the measure of the joint variability of two random variables.

$$\text{Cov}(x, y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$$

Sum of squares

The sum of squares is the sum of the squared differences between each data point and the mean.

$$\sum_{i=1}^n (x_i - \bar{x})^2$$

Denoted S_{xx} . If two variables are involved (covariance), it is denoted as S_{xy} .

Shortcut formula:

$$S_{xx} = \sum_{i=1}^n x_i^2 - \frac{1}{n \left(\sum_{i=1}^n x_i \right)^2}$$
$$S_{xy} = \sum_{i=1}^n x_i y_i - \frac{1}{n \left(\sum_{i=1}^n x_i \right)} \left(\sum_{i=1}^n y_i \right)$$

Mean estimation for continuous grouped data

$$\bar{x} \approx \frac{1}{n} \sum_{i=1}^K f_i \cdot m_i$$

where

- f_i is the frequency of the i th interval
- m_i is the midpoint of the i th interval
- n is the total number of observations

Estimating a variance for continuous grouped data

Using the same notation as above:

$$\sigma_n^2 \approx \frac{1}{n} \sum_{i=1}^K f_i \cdot (m_i - \bar{x})^2$$

Graphing

Graphing cubic curves with one distinct real root

If we are given one root of a cubic equation, we can use synthetic division to factor it.

Example

Given that $(x - 3)$ is a factor of $f(x) = x^3 + x^2 - 4x - 24$, sketch the curve of $y = f(x)$.

First, synthetic division:

	x^3	x^2	x^1	x^0	
3		1	1	-4	-24
			3	12	24
		1	4	8	0


So we have $f(x) = (x - 3)(x^2 + 4x + 8)$

$\therefore x = 3$ is a root.

The discriminant of $x^2 + 4x + 8$ is $4^2 - 4 \cdot 1 \cdot 8 = -16$, which is negative, so $x = 3$ is the only real root.

The y intercept is $f(0) = 0^3 + 0^2 - 4 \cdot 0 - 24 = -24$

Now we can draw the curve:



Logarithms

Domain

$(0, \infty)$

Range

$(-\infty, \infty)$

Complex Numbers

Magnitude

The magnitude of a complex number $z = a + bi$ is defined as $|z| = \sqrt{a^2 + b^2}$.

Argument

For any complex number, like $z = 3 + 4i$, there is $\arg(z)$ that is the angle (in radians) between the x axis and the complex number in the Argand diagram.

So for z :

$$\begin{aligned}\tan \theta &= \frac{4}{3} \\ \theta &= \arctan\left(\frac{4}{3}\right) \\ &\approx 0.93\end{aligned}$$

Trigonometry

Unit circle

Reference angle

The *reference angle* of an angle is the smallest angle between the terminal side of the angle and the x -axis. It is always a positive angle less than or equal to 90 degrees.

Pythagorean identity in the first quadrant

In the first quadrant, any point (x, y) forms a right triangle with the origin, where x and y are the lengths of the legs and $\sqrt{x^2 + y^2}$ is the length of the hypotenuse. Since in the unit circle, the length of the hypotenuse is 1, by definition, the Pythagorean identity states that $x^2 + y^2 = 1$.

Exercises

1. For a point in the first quadrant with $y = \frac{\sqrt{5}}{3}$, find its x coordinate.

$$\sqrt{\left(\frac{\sqrt{5}}{3}\right)^2 + x^2} = 1$$

$$\left(\frac{\sqrt{5}}{3}\right)^2 + x^2 = 1$$

$$x^2 = 1 - \left(\frac{\sqrt{5}}{3}\right)^2$$

$$x^2 = 1 - \frac{5}{9}$$

$$x^2 = \frac{4}{9}$$

$$x = \sqrt{\frac{4}{9}}$$

$$x = \frac{2}{3}$$

2. Point in the first quadrant. y coordinate is $\frac{\sqrt{2}}{3}$. What is the x coordinate?

$$\sqrt{\left(\frac{\sqrt{2}}{3}\right)^2 + x^2} = 1$$

$$\left(\frac{\sqrt{2}}{3}\right)^2 + x^2 = 1$$

$$x^2 = 1 - \left(\frac{\sqrt{2}}{3}\right)^2$$

$$x^2 = 1 - \frac{2}{9}$$

$$x^2 = \frac{7}{9}$$

$$x = \sqrt{\frac{7}{9}}$$

$$x = \frac{\sqrt{7}}{3}$$

Pythagorean trigonometric identity

In the unit circle, since for any central angle $x = \cos \theta$ and $y = \sin \theta$.

That means

$$x^2 + y^2 = 1$$

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

This is the **Pythagorean trigonometric identity**. It works in all quadrants.

Exercises

1. Point P on the unit circle with $x = \frac{2}{5}$. Find $\sin \theta$.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\sin^2 \theta = 1 - \cos^2 \theta$$

$$\sin \theta = \sqrt{1 - \cos^2 \theta}$$

$$\sin \theta = \sqrt{1 - \frac{4}{25}}$$

$$\sin \theta = \sqrt{\frac{21}{25}}$$

$$\sin \theta = \frac{\sqrt{21}}{5}$$

2. y coordinate is $\frac{4}{9}$. Find $\cos \theta$.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\left(\frac{4}{9}\right)^2 + \cos^2 \theta = 1$$

$$\cos^2 \theta = 1 - \left(\frac{4}{9}\right)^2$$

$$\cos^2 \theta = 1 - \frac{16}{81}$$

$$\cos^2 \theta = \frac{65}{81}$$

$$\cos \theta = \sqrt{\frac{65}{81}}$$

$$\cos \theta = \frac{\sqrt{65}}{9}$$

3. x coordinate is $\frac{\sqrt{7}}{4}$. Find $\csc \theta$.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\sin^2 \theta = 1 - \cos^2 \theta$$

$$\sin \theta = \sqrt{1 - \cos^2 \theta}$$

$$\sin \theta = \sqrt{1 - \left(\frac{\sqrt{7}}{4}\right)^2}$$

$$\sin \theta = \sqrt{1 - \frac{7}{16}}$$

$$\sin \theta = \sqrt{\frac{9}{16}}$$

$$\sin \theta = \frac{3}{4}$$

$$\therefore$$

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\csc \theta = \frac{4}{3}$$

4. x coordinate is $\frac{3}{7}$. Find $\csc \theta$.

$$\sin^2 \theta = 1 - \cos^2 \theta$$

$$\sin \theta = \sqrt{1 - \frac{9}{49}}$$

$$\sin \theta = \sqrt{\frac{40}{49}}$$

$$\sin \theta = \frac{\sqrt{40}}{7}$$

$$\therefore$$

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\csc \theta = \frac{7}{\sqrt{40}}$$

$$\csc \theta = \frac{7}{2\sqrt{10}}$$

$$\csc \theta = \frac{7}{2\sqrt{10}}$$

$$\csc \theta = \frac{7\sqrt{10}}{20}$$

5. y coordinate is $\frac{1}{9}$. Find $\tan \theta$.

We have $\sin \theta = \frac{1}{9}$.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\cos \theta = \sqrt{1 - \sin^2 \theta}$$

$$\cos \theta = \sqrt{1 - \left(\frac{1}{9}\right)^2}$$

$$\cos \theta = \sqrt{1 - \frac{1}{81}}$$

$$\cos \theta = \sqrt{\frac{80}{81}}$$

$$\cos \theta = \frac{\sqrt{80}}{9}$$

$$\cos \theta = \frac{4\sqrt{5}}{9}$$

$$\therefore$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\tan \theta = \frac{\frac{1}{9}}{\frac{4\sqrt{5}}{9}}$$

$$\tan \theta = \frac{1}{9} \cdot \frac{9}{4\sqrt{5}}$$

$$\tan \theta = \frac{\sqrt{5}}{20}$$

6. x coordinate is $\frac{\sqrt{11}}{6}$. Find $\cot(\theta)$

$$\text{We have } \cos \theta = \frac{\sqrt{11}}{6} \text{ and } \cot(\theta) = \frac{1}{\tan \theta} = \frac{\cos \theta}{\sin \theta} = \frac{\frac{\sqrt{11}}{6}}{\sin \theta}$$

$$\begin{aligned}
\sin^2 \theta + \cos^2 \theta &= 1 \\
\sin^2 \theta &= 1 - \cos^2 \theta \\
\sin^2 \theta &= 1 - \left(\frac{\sqrt{11}}{6}\right)^2 \\
\sin^2 \theta &= 1 - \frac{11}{36} \\
\sin^2 \theta &= \frac{25}{36} \\
\sin \theta &= \sqrt{\frac{25}{36}} \\
\sin \theta &= \frac{5}{6} \\
\therefore \\
\cot \theta &= \frac{\cos \theta}{\sin \theta} \\
&= \frac{\frac{\sqrt{11}}{6}}{\frac{5}{6}} \\
\cot \theta &= \frac{\sqrt{11}}{5}
\end{aligned}$$

Quadrantal angles

Quadrantal angles are angles that are multiples of 90° . For example, 0° , 90° , 180° , 270° , and 360° are all quadrantal angles. They lie on an axis.

They are useful to find the values of trigonometric functions. Just remember that for points on the unit circle, $\cos \theta = x$ and $\sin \theta = y$.

The corresponding angles on $[0, 360^\circ]$ are $0, \frac{\pi}{2}, \pi, 3\frac{\pi}{2}, 2\pi$.

Example: $\cos 0 + \sin\left(\frac{\pi}{2}\right) = 1 + 1 = 2$

Examples

- Find the value of $\tan(180^\circ) + \cot(270^\circ)$

$$\begin{aligned}
\tan(180^\circ) + \cot(270^\circ) &= \frac{\sin(180^\circ)}{\cos(180^\circ)} + \frac{\cos(270^\circ)}{\sin(270^\circ)} \\
&= \frac{0}{-1} + \frac{0}{-1} \\
&= 0
\end{aligned}$$

- $\sec\left(\frac{\pi}{2}\right)$?

$$\begin{aligned}
\sec\left(\frac{\pi}{2}\right) &= \frac{1}{\cos\left(\frac{\pi}{2}\right)} \\
&= \frac{1}{0}
\end{aligned}$$

So it's undefined.

Law of sines

In any triangle, where a , b , and c are the lengths of the sides opposite angles A , B , and C respectively, the law of sines states that:

$$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$$

For acute angles, the formula can be flipped:

$$\frac{\sin(A)}{a} = \frac{\sin(B)}{b} = \frac{\sin(C)}{c}$$

For obtuse angles, the same formula works, but it will always give acute angles. To find an obtuse angle, subtract the acute angle from 180° .

Exercises

- Find a if $A = 70^\circ$, $B = 35^\circ$, $b = 12$.

$$\begin{aligned}\frac{a}{\sin(A)} &= \frac{b}{\sin(B)} \\ \frac{a}{\sin(70^\circ)} &= \frac{12}{\sin(35^\circ)} \\ a &= 12 \cdot \frac{\sin(70^\circ)}{\sin(35^\circ)} \\ a &\approx 12 \cdot \frac{0.9397}{0.5736} \\ a &\approx 19.66\end{aligned}$$

- R , Q and P form a triangle. $PR = 21m$. $\angle Q = 89^\circ$, $\angle P = 38^\circ$. What is QR ?

$$\begin{aligned}\frac{QR}{\sin(\angle P)} &= \frac{PR}{\sin(\angle Q)} \\ \frac{QR}{\sin(38^\circ)} &= \frac{21}{\sin(89^\circ)} \\ QR &= \sin(38^\circ) \cdot \frac{21}{\sin(89^\circ)}\end{aligned}$$

- $\triangle XYZ$. $\angle Y = 57^\circ$, $XZ = 35$, $XY = 24$. Find $\angle Z$.

$$\begin{aligned}\frac{\sin(\angle Z)}{XY} &= \frac{\sin(\angle Y)}{XZ} \\ \frac{\sin(\angle Z)}{24} &= \frac{\sin(57^\circ)}{35} \\ \sin(\angle Z) &= 24 \cdot \frac{\sin(57^\circ)}{35} \\ \sin(\angle Z) &\approx 0.575 \\ \angle Z &\approx 35^\circ\end{aligned}$$

Law of cosines

In a triangle where a , b and c are the sides opposite angles A , B and C respectively, the law of cosines states that:

$$a^2 = b^2 + c^2 - 2bc \cos(A)$$

$$b^2 = a^2 + c^2 - 2ac \cos(B)$$

$$c^2 = a^2 + b^2 - 2ab \cos(C)$$

Calculus

Limits

Limits of logarithmic functions


For any real n :

$$\lim_{x \rightarrow n} \log(x) = \log(x)$$

For infinity, remember the domain and apply transformations.

Limits of reciprocal functions

Remember, reciprocal functions have the form $f(x) = \frac{1}{x}$. The limit of a reciprocal function as x approaches infinity is always zero.



In general, for any non-constant polynomial $P(x)$, we have $\lim_{x \rightarrow \pm\infty} \frac{1}{P(x)} = 0$

But for $\lim(x \rightarrow 0)$, we have to consider the degree of the leading term of $P(x)$.

- Any function of the form $y = \frac{1}{x^n}$ where n is odd has the same shape as $y = \frac{1}{x}$. There is no limit at $x = 0$ because left and right limit are not the same.
- Any function of the form $y = \frac{1}{x^n}$ where n is even has the same shape as $y = \frac{1}{x^2}$. There is a limit, since the function will approach ∞ or $-\infty$.

Differentiation

Constant multiple rule

$$\frac{d}{dx}(kf(x)) = k \frac{d}{dx}(f(x))$$

Sum rule

$$\frac{d}{dx}(u(x) + v(x)) = \frac{du}{dx} + \frac{dv}{dx}$$

For example for $y = x^2 + x$:

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx}(x^2 + x) \\
 &= \frac{d}{dx}(x^2) + \frac{d}{dx}(x) \\
 &= 2x + 1
 \end{aligned}$$

The power rule

Given a monomial $f(x) = x^n$, where n is a real number,

$$f'(x) = nx^{n-1}$$

Exercises

1. Calculate $\frac{dy}{dx}$ for $y = 2x - \sqrt{x}$

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx}(2x - \sqrt{x}) \\
 &= \frac{d}{dx}2x - \frac{d}{dx}\sqrt{x} \\
 &= 2 - \frac{d}{dx}x^{1/2} \\
 &= 2 - \frac{1}{2} \cdot x^{-1/2} \\
 &= 2 - \frac{1}{2\sqrt{x}}
 \end{aligned}$$

2. Find the derivative of $f(x) = \frac{1}{8}x^8$

$$\begin{aligned}
 f'(x) &= \frac{1}{8} \cdot \frac{d}{dx}x^8 \\
 &= \frac{1}{8} \cdot 8x^7 \\
 &= \frac{8x^7}{8} \\
 &= x^7
 \end{aligned}$$

3. Derivative of $f(x) = 8\sqrt{x}$

$$\begin{aligned}
 f'(x) &= \frac{d}{dx}8\sqrt{x} \\
 &= \frac{d}{dx}8x^{1/2} \\
 &= 8 \cdot \frac{d}{dx}\left(\frac{1}{2} \cdot x^{-1/2}\right) \\
 &= 8 \cdot \frac{1}{2} \cdot \frac{1}{2\sqrt{x}} \\
 &= \frac{4}{\sqrt{x}}
 \end{aligned}$$

Exponential function

When the base is e :

$$\frac{d}{dx}e^x = e^x$$

It is the only function whose derivative is itself.

For an arbitrary base a :

$$\frac{d}{dx}a^x = a^x \ln(a)$$

Exercises

- Find the slope of the tangent to $f(x) = 5^x + 2e^x - \ln 2$ at $x = 1$.

$$\frac{d}{dx} 5^x + 2e^x - \ln 2 = 5^x \ln 5 + 2e^x$$

Evaluating at $x = 1$, we get $5 \ln 5 + 2e$

Rate of change

Average rate of change on an interval $[a, a + h]$:

$$\frac{\delta y}{\delta x} = \frac{f(a + h) - f(a)}{h}$$

The *instantaneous rate of change* at a point $x = a$ is given by:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

Exercises

- Given that $f(x) = x^2$, what is $f'(2)$ according to the definition?

$$\begin{aligned} f'(2) &= \lim_{h \rightarrow 0} \frac{(2 + h)^2 - 2^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{4 + 4h + h^2 - 4}{h} \\ &= \lim_{h \rightarrow 0} \frac{4h + h^2}{h} \\ &= \lim_{h \rightarrow 0} (4 + h) \\ &= 4 \end{aligned}$$

Slope of a tangent line

The derivative of a function $f(x)$ at a point $x = a$ is the slope of the tangent line to the graph of $f(x)$ at that point.

So to find the slope of the tangent line at a point (x_1, y_1) , we evaluate $f'(x_1)$.

Exercises

- Find the slope of the tangent to $y = 5x^7 - 3x^5 + 10x^3$ at the point where $x = 1$.

First find $f'(x)$:

$$f'(x) = \frac{d}{dx} (5x^7 - 3x^5 + 10x^3)$$

$$f'(x) = 5 \cdot \frac{d}{dx} x^7 - 3 \cdot \frac{d}{dx} x^5 + 10 \cdot \frac{d}{dx} x^3$$

$$f'(x) = 5 \cdot 7x^6 - 3 \cdot 5x^4 + 10 \cdot 3x^2$$

$$f'(x) = 35x^6 - 15x^4 + 30x^2$$

$$f'(x) = 5(7x^6 - 3x^4 + 6x^2)$$

- Find the slope of the tangent to $y = 4x^3 + \frac{5}{2}x^2 - 200x$ at $x = 4$.

$$\begin{aligned} f'(x) &= \frac{d}{dx} 4x^3 + \frac{5}{2}x^2 - 200x \\ &= 12x^2 + 5x - 200 \end{aligned}$$

The slope is $f'(4) = 12$

- Find the x -coordinate of the point on the curve $y = 2x^2 - 9x - 6$ whose tangent has a slope of 3.

First let's get the derivative of the function:

$$\begin{aligned} f'(x) &= \frac{d}{dx} (2x^2 - 9x - 6) \\ &= 4x - 9 \end{aligned}$$

Now we set the derivative equal to the slope of the tangent line:

$$\begin{aligned} 4x - 9 &= 3 \\ 4x &= 12 \\ x &= 3 \end{aligned}$$

Finding the equation of a tangent line at a point

- Find the derivative.
- Evaluate the derivative at the given x . That is the slope.
- Replace the slope, the x and y coordinates we are interested in in the point-slope form of the equation: $y - y_1 = m(x - x_1)$.

Exercises

- Find the equation of the tangent to the curve $y = x^2 + 5x$ at the point $(-1, -4)$.

First find the derivative:

$$y' = 2x + 5$$

Find the slope: $2 \cdot -1 + 5 = 3$.

Now substitute the coordinates in the point-slope form:

$$\begin{aligned} y + 4 &= 3(x + 1) \\ y &= 3x + 3 - 4 \\ y &= 3x - 1 \end{aligned}$$

- Find the equation of the tangent line to the curve $y = \frac{2}{x^2} + 6x$ at $(-1, -4)$.

First the derivative: $y' = 2 \cdot -2x^{-1} + 6 = -\frac{4}{x} + 6$

Then the slope: $\frac{-4}{-1} + 6 = 10$.

Finally, substitute in the point-slope form:

$$\begin{aligned}(y + 4) &= 10(x + 1) \\ y &= 10x + 10 - 4 \\ y &= 10x + 6\end{aligned}$$

- Find where the tangent to the curve $y = 4\sqrt{x} - 4x$ at $(1, 0)$ intersects the y -axis.

First, find the derivative: $y' = \frac{d}{dx}(4\sqrt{x} - 4x) = 4 \cdot \frac{1}{2} \cdot x^{-1/2} - 4 = \frac{2}{\sqrt{x}} - 4$

Then, the slope: $\frac{2}{\sqrt{1}} - 4 = 2 - 4 = -2$.

The equation:

$$\begin{aligned}y - 0 &= -2(x - 1) \\ y &= -2x + 2\end{aligned}$$

Then set $x = 0$:

$$\begin{aligned}y &= -2x + 2 \\ y &= 2\end{aligned}$$

- Find where the tangent to the curve $y = \frac{9}{x} - 2x$ at the point $(3, -3)$ intersects the x -axis.

First, find the derivative: $y' = 9 \cdot -1 \cdot x^{-2} - 2 = -\frac{9}{x^2} - 2$.

Then the slope: $-\frac{9}{3^2} - 2 = -\frac{9}{9} - 2 = -1 - 2 = -3$.

Substitute in the point-slope equation:

$$\begin{aligned}y + 3 &= -3(x - 3) \\ y &= -3x + 9 - 3 \\ y &= -3x + 6\end{aligned}$$

Then, set $y = 0$:

$$\begin{aligned}0 &= -3x + 6 \\ -6 &= -3x \\ x &= 2\end{aligned}$$

Integration

The antiderivative

The antiderivative is the opposite of the derivative.

Infinitely many antiderivatives map to the same function.

For example, since the derivative of x^2 is $2x$, we have x^2 as an antiderivative of $2x$. But $x^2 + 2$, $x^2 + 1$, etc. are also antiderivatives of $2x$.

In general, the antiderivatives of $2x$ follow the pattern $x^2 + C$ where C is a constant.

We can also formulate that using the $\int$ symbol for integration:

$$\int 2x \, dx = x^2 + C$$

Where C is called the **constant of integration**.

These integrals are **indefinite integrals**.

The function being integrated (here $2x$) is called the **integrand**.

The power rule for integration

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

Exercises

1. Antiderivative of $\frac{1}{x^4}$?

$$\begin{aligned}\int \frac{1}{x^4} dx &= \int x^{-4} dx \\ &= \frac{x^{-4+1}}{-4+1} + C \\ &= \frac{x^{-3}}{-3} + C \\ &= -\frac{1}{3}x^3 + C\end{aligned}$$

2. Antiderivative of $x^{\frac{5}{8}}$

$$\begin{aligned}\int x^{\frac{5}{8}} dx &= \frac{x^{\frac{5}{8}+1}}{\frac{5}{8}+1} + C \\ &= \frac{x^{\frac{13}{8}}}{\frac{13}{8}} + C \\ &= \frac{8x^{\frac{13}{8}}}{13} + C\end{aligned}$$

3. Antiderivative of $\sqrt[5]{x^7}$

$$\begin{aligned}\int x^{\frac{7}{5}} dx &= \frac{x^{\frac{7}{5}+1}}{\frac{7}{5}+1} + C \\ &= \frac{x^{12/5}}{12/5} \\ &= \frac{5x^{12/5}}{12}\end{aligned}$$

4. Calculate $\int \left(\frac{z}{5} dz\right)$

$$\begin{aligned}\int \left(\frac{z}{5} dz\right) &= \frac{1}{5} \int z dz \\ &= \frac{1}{5} \cdot \left(\frac{z^2}{2} + C\right) \\ &= \frac{z^2}{10} + C\end{aligned}$$

The constant factor rule for indefinite integrals

To take the integral of a power function with a constant factor, like $5x^2$, we can take the constant factor out of the integral:

$$\int 5x^2 dx = 5 \cdot \int x^2 dx$$

This is the **constant factor rule**. In general:

$$\int k f(x) dx = k \int f(x) dx$$

Where k is a **constant**. It does not work if k is a variable.

Exercises

1. Calculate $\int \frac{1}{2y^5} dy$

$$\begin{aligned}\int \frac{1}{2y^5} dy &= \frac{1}{2} \int y^{-5} \\ &= \frac{1}{2} \cdot \frac{y^{-5+1}}{-5+1} + C \\ &= \frac{1}{2} \cdot \frac{y^{-4}}{-4} + C \\ &= \frac{1}{2} \cdot -\frac{1}{4y^4} + C \\ &= -\frac{1}{8y^4} + C\end{aligned}$$

2. Calculate $\int \frac{1}{3x^6} dx$

$$\begin{aligned}\int \frac{1}{3x^6} dx &= \frac{1}{3} \cdot \int x^{-6} \\ &= \frac{1}{3} \cdot \frac{x^{-5}}{-5} + C \\ &= \frac{1}{3} \cdot -\frac{1}{5x^5} + C \\ &= -\frac{1}{15x^5} + C\end{aligned}$$

3. Calculate $\int \frac{\pi}{2} dz$

$$\begin{aligned}\int \frac{\pi}{2} dz &= \frac{\pi}{2} \int z^0 dz \\ &= \frac{\pi}{2} \cdot z + C \\ &= \frac{\pi z}{2} + C\end{aligned}$$

The sum rule for indefinite integrals

To take the integral of a sum, you can take the sum of the integrals of each term:

$$\int (f(x) \pm g(x)) dx = \int f(x) dx \pm \int g(x) dx$$

The sum rule can be combined with the constant factor rule to integrate sums of constant multiples of functions:

$$\int (k_1 f(x) + k_2 g(x)) dx = k_1 \int f(x) dx + k_2 \int g(x) dx$$

Exercises

- Calculate $\int \sqrt{x}(2+x) dx$

$$\begin{aligned} \int \sqrt{x}(2+x) dx &= \int 2\sqrt{x} + x \cdot \sqrt{x} dx \\ &= \int 2\sqrt{x} dx + \int x \cdot \sqrt{x} dx \\ &= 2 \int \sqrt{x} dx + \int x^{3/2} dx \\ &= 2 \cdot \frac{x^{3/2}}{\frac{3}{2}} + \frac{x^{5/2}}{\frac{5}{2}} + C \\ &= \frac{4x^{3/2}}{3} + \frac{2x^{5/2}}{5} + C \\ &= \frac{4\sqrt{x^3}}{3} + \frac{2\sqrt{x^5}}{5} + C \end{aligned}$$

- Calculate $\int \frac{2+x^4}{x^2} dx$

$$\begin{aligned} \int \frac{2+x^4}{x^2} dx &= \int \frac{2}{x^2} dx + \int \frac{x^4}{x^2} dx \\ &= \int 2 \cdot x^{-2} dx + \int x^2 dx \\ &= 2 \cdot \int x^{-2} dx + \frac{x^3}{3} + C \\ &= 2 \cdot -x^{-1} + \frac{x^3}{3} + C \end{aligned}$$

- Calculate:

$$\begin{aligned} \int (3-x^2)^2 dx &= \int (9-6x^2+x^4) dx \\ &= 9 \int 1 dx - 6 \int x^2 dx + \int x^4 dx \\ &= 9x - 6 \frac{x^3}{3} + \frac{x^5}{5} + C \\ &= \frac{x^5}{5} - 2x^3 + 9x + C \end{aligned}$$

- Find the antiderivative of $z^3 + 2z - 1$

$$\begin{aligned}
\int z^3 + 2z - 1 \, dz &= \int z^3 \, dz + \int 2z \, dz - \int 1 \, dz \\
&= \frac{z^4}{4} + 2\frac{z^2}{2} - z + C \\
&= \frac{z^4}{4} + z^2 - z + C
\end{aligned}$$

Integrating reciprocal functions

Since $\frac{d}{dx}(\ln x) = \frac{1}{x}$ for $x > 0$, we can generalize to $x \neq 0$ by using an absolute value:

$$\frac{d}{dx}(\ln|x|) = \frac{1}{x}$$

Why, in the negative case? We have two functions, chained. $|x| = -x$ if x is negative. So we have $\ln|x| = \ln(-x)$. By the chain rule, $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$. Here $(du) \cdot (dx)(-x) = -1$ (constant factor rule), and $\frac{dy}{du} \ln(u) = \frac{1}{u}$. So we substitute x back in, and we get $\frac{dy}{dx} = \frac{1}{-x} \cdot -1 = \frac{1}{x}$

So the integral is:

$$\int \frac{1}{x} \, dx = \ln|x| + C$$

Vectors

Magnitude

For a vector $\vec{AB} = \mathbf{a}$, we denote its magnitude, a.k.a. modulus or length, as $|\vec{AB}|$ or $|\mathbf{a}|$.

When adding two vectors **that are at a right angle**, through the pythagorean identity, we can deduce:

$$|\mathbf{a} + \mathbf{b}|^2 = |\mathbf{a}|^2 + |\mathbf{b}|^2$$

When the angle is not a right angle, we can use the law of cosines instead:

$$|\mathbf{a} + \mathbf{b}|^2 = |\mathbf{a}|^2 + |\mathbf{b}|^2 - 2|\mathbf{a}||\mathbf{b}| \cos \theta$$

Exercises

- In rectangle $ABCD$, $AB = 5$ and $BC = \sqrt{39}$. Find $|\vec{BD}|$.

Since all the angles of the rectangle are right angles, and $|\vec{BD}| = |\vec{AC}|$, we have:

$$\begin{aligned}
|\vec{BD}|^2 &= |\vec{AB}|^2 + |\vec{BC}|^2 \\
&= 25 + 39 \\
&= 64 \\
|\vec{BD}| &= \sqrt{64} \\
&= 8
\end{aligned}$$

Determining whether two vectors are parallel

In general, two vectors $\mathbf{a}$ and $\mathbf{b}$ are **parallel**, a.k.a. **colinear**, if there is a number λ such that $\mathbf{a} = \lambda \mathbf{b}$.

That is also denoted $\mathbf{a} \parallel \mathbf{b}$.

Note that for any vector $\mathbf{a}$, $0 \cdot \mathbf{a}$ is equal to the zero vector, so by definition, the zero vector is parallel to all other vectors.