


This formula was subsequently proved by Mangoldt (1895; Riesel 1994, p. 47; Edwards 2001, pp. 48 and 62-65). The integral on the right-hand side converges only for $x > 1$, but since there are no primes less than 2, the only values of interest are for $x \geq 2$. Since it is monotonic decreasing, the maximum therefore occurs at $x = 2$, which has value

$$\int_2^\infty \frac{dt}{t \ln t (t^2 - 1)} = 0.14001010114328692668 \dots$$

(9)

(OEIS [A096623](#); Derbyshire 2004, p. 329).




Riemann also considered the function

$$R(x) = \sum_{n=1}^\infty \frac{\mu(n)}{n} \operatorname{li}(x^{1/n}),$$

(10)


sometimes also denoted $\operatorname{Ri}(x)$ (Borwein *et al.* 2000), obtained by replacing $f(x^{1/n})$ in the Riemann function with the logarithmic integral $\operatorname{li}(x^{1/n})$, where $\zeta(z)$ is the Riemann zeta function and $\mu(n)$ is the Möbius function (Hardy 1999, pp. 16 and 23; Borwein *et al.* 2000; Havil 2003, p. 198). $R(x)$ is plotted above, including on a semilogarithmic scale (bottom two plots), which illustrate the fact that $R(x)$ has a series of zeros near the origin. These occur at 10^{-x} for $x = 14\,827.7$ (OEIS [A143530](#)), 15300.7, 21381.5, 25461.7, 32711.9, 40219.6, 50689.8, 62979.8, 78890.2, 98357.8, ..., corresponding to $x = 1.829 \times 10^{-14\,828}$ (OEIS [A143531](#)), $2.040 \times 10^{-15\,301}$, $3.289 \times 10^{-21\,382}$, $2.001 \times 10^{-25\,462}$, $1.374 \times 10^{-32\,712}$, $2.378 \times 10^{-40\,220}$, $1.420 \times 10^{-50\,690}$, $1.619 \times 10^{-62\,980}$, $6.835 \times 10^{-78\,891}$, $1.588 \times 10^{-98\,358}$, ...



The quantity $R(x) - \pi(x)$ is plotted above.

This function is implemented in the Wolfram Language as `RiemannR[x]`.

Ramanujan independently derived the formula for $R(n)$, but nonrigorously (Berndt 1994, p. 123; Hardy 1999, p. 23). The following table compares $\pi(10^n)$ and $R(10^n)$ for small n . Riemann conjectured that $R(n) = \pi(n)$ (Knuth 1998, p. 382), but this was disproved by Littlewood in 1914 (Hardy and Littlewood 1918).



The Riemann prime counting function is identical to the Gram series

$$G(x) = 1 + \sum_{k=1}^\infty \frac{(\ln x)^k}{k! \zeta(k+1)},$$

(11)

where $\zeta(x)$ is the [Riemann zeta function](#) (Hardy 1999, pp. 24-25), but the [Gram series](#) is much more tractable for numeric computations. For example, the plots above show the difference $G(x) - R(x)$ where $R(x)$ is computed using the [Wolfram Language's](#) built-in `NSum` command (black) and approximated using the first 10^1 (blue), 10^2 (green), 10^3 (yellow), 10^4 (orange), and 10^5 (red) points.

In the table, $[x]$ denotes the [nearest integer function](#). Note that the values given by Hardy (1999, p. 26) for $x = 10^9$ are incorrect.

n	$\text{nint}(R(10^n))$	$\text{nint}(R(10^n) - \pi(10^n))$
Sloane	A057793	A057794
1	5	1
2	26	1
3	168	0
4	1227	-2
5	9587	-5
6	78527	29
7	664667	88
8	5761552	97
9	50847455	-79
10	455050683	-1828
11	4118052495	-2318
12	37607910542	-1476

Riemann's function is related to the [prime counting function](#) by

$$\pi(x) = R(x) - \sum_{\rho} R(x^{\rho}),$$

(12)

where the [sum](#) is over all complex (nontrivial) zeros ρ of $\zeta(s)$ (Ribenoim 1996), i.e., those in the [critical strip](#) so $0 < \Re[\rho] < 1$, interpreted to mean

$$\sum_{\rho} R(x^{\rho}) = \lim_{t \rightarrow \infty} \sum_{|\Im(\rho)| < t} R(x^{\rho}).$$

(13)

However, no proof of the equality of (12) appears to exist in the literature (Borwein *et al.* 2000).

SEE ALSO:
[Gram Series](#), [Prime Counting Function](#), [Prime Number Theorem](#), [Riemann Function](#), [Riemann Hypothesis](#), [Soldner's Constant](#)

REFERENCES:
Borwein, J. M.; Bradley, D. M.; and Crandall, R. E. "Computational Strategies for the Riemann Zeta Function." *J. Comput. Appl. Math.* **121**, 247-296, 2000.
Berndt, B. C. *Ramanujan's Notebooks, Part IV*. New York: Springer-Verlag, 1994.
Derbyshire, J. *Prime Obsession: Bernhard Riemann and the Greatest Unsolved Problem in Mathematics*. New York: Penguin, 2004.
Edwards, H. M. *Riemann's Zeta Function*. New York: Dover, 2001.
Hardy, G. H. and Littlewood, J. E. *Acta Math.* **41**, 119-196, 1918.
Hardy, G. H. "The Series $R(x)$ " §2.3 in *Ramanujan: Twelve Lectures on Subjects Suggested by His Life and Work, 3rd ed.* New York: Chelsea, 1999.
Havil, J. *Gamma: Exploring Euler's Constant*. Princeton, NJ: Princeton University Press, p. 42, 2003.
Ingham, A. E. *The Distribution of Prime Numbers*. London: Cambridge University Press, p. 83, 1990.
Knuth, D. E. *The Art of Computer Programming, Vol. 2: Seminumerical Algorithms, 3rd ed.* Reading, MA: Addison-Wesley, 1998.
Landau, E. *Handbuch der Lehre von der Verteilung der Primzahlen, 3rd ed.* New York: Chelsea, 1974.
Mangoldt, H. von. "Zu Riemann's Abhandlung 'Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse.'" *J. reine angew. Math.* **114**, 255-305, 1895.
Mathews, G. B. Ch. 10 in *Theory of Numbers*. New York: Chelsea, 1961.
Ribenoim, P. *The New Book of Prime Number Records*. New York: Springer-Verlag, pp. 224-225, 1996.
Riemann, G. F. B. "Über die Anzahl der Primzahlen unter einer gegebenen Grösse." *Monatsber. Königl. Preuss. Akad. Wiss. Berlin*, 671-680, Nov. 1859. Reprinted in *Das Kontinuum und Andere Monographien* (Ed. H. Weyl). New York: Chelsea, 1972. Also reprinted in Edwards, H. M. Appendix. *Riemann's Zeta Function*. New York: Dover, pp. 299-305, 2001.
Riemann, B. "Über die Darstellbarkeit einer Function durch eine trigonometrische Reihe." Reprinted in *Gesammelte math. Abhandlungen*. New York: Dover, pp. 227-264, 1957.
Riesel, H. and Göhl, G. "Some Calculations Related to Riemann's Prime Number Formula." *Math. Comput.* **24**, 969-983, 1970.
Riesel, H. "The Riemann Prime Number Formula." *Prime Numbers and Computer Methods for Factorization, 2nd ed.* Boston, MA: Birkhäuser, pp. 50-52, 1994.
Sloane, N. J. A. Sequences [A057793](#), [A057794](#), [A096623](#), [A096624](#), [A096625](#), [A143530](#), [A143531](#) in "The On-Line Encyclopedia of Integer Sequences."
Wagon, S. *Mathematica in Action*. New York: W. H. Freeman, pp. 28-29 and 362-372, 1991.

Referenced on Wolfram|Alpha: [Riemann Prime Counting Function](#)

CITE THIS AS:
Weisstein, Eric W. "Riemann Prime Counting Function." From *MathWorld*--A Wolfram Web Resource.
<https://mathworld.wolfram.com/RiemannPrimeCountingFunction.html>

The #1 tool for creating Demonstrations and anything technical.

Computerbasedmath.org »
Join the initiative for modernizing math education.

Wolfram Problem Generator »
Unlimited random practice problems and answers with built-in Step-by-step solutions. Practice online or make a printable study sheet.

Explore anything with the first computational knowledge engine.

Online Integral Calculator »
Solve integrals with Wolfram|Alpha.

Wolfram Education Portal »
Collection of teaching and learning tools built by Wolfram education experts: dynamic textbook, lesson plans, widgets, interactive Demonstrations, and more.

Explore thousands of free applications across science, mathematics, engineering, technology, business, art, finance, social sciences, and more.

Step-by-step Solutions »
Walk through homework problems step-by-step from beginning to end. Hints help you try the next step on your own.

Wolfram Language »
Knowledge-based programming for everyone.