

Computing time-dependent reduced models for classical and quantum dynamics

Tommaso Grigoletto^{??}

^a*Department of Information Engineering, University of Padova, Italy*

Abstract

This paper introduces a novel method for approximating the dynamics of a large autonomous system projected onto a fixed subspace. The core contribution is a novel recursive algorithm to construct an effective time-dependent generator that is polynomial in the time variable, ensuring accuracy for short time scales. The derivation is based on the Taylor expansion of the exponential map and a new result for computing the time-ordered exponential of polynomial generators. This work is motivated by the challenge of deriving time-convolutionless master equations in quantum physics and the proposed method offers an alternative to typical derivations based on expansions in the coupling strength. The resulting approximation is accurate for small times, does not require a weak-coupling assumption, performs better than a truncation of the exponential map at low orders, and crucially, guarantees a completely positive and trace-preserving map at the lowest orders. The proposed method is validated against prototypical models: a central spin model and an Ising spin chain.

Key words: Identification and model reduction; Time-varying systems; Time-invariant

1 Introduction

Whenever a dynamical model is too large to be simulated efficiently, one should resort to model reduction techniques to find a smaller model that is simulatable and that retains fundamental characteristics of the original system. A dominant approach in control systems theory – including, for example, minimal realization techniques [?], balanced truncation [?,?], and moment matching [?,?] – consists of finding a low-dimensional subspace where to project the dynamics, obtaining a linear time-invariant (LTI) model. This work considers a different approach: the space onto which the dynamics is projected is assumed to be *given a priori and fixed*, and, instead, the dynamics is approximated by seeking a linear time-varying (LTV) model.

This approach is inspired by quantum physics. To obtain realistic models of open quantum systems, one shall consider the effects of the interaction between the system and its environment [?]. Unfortunately, simulating the system-environment couple is impractical (if not impossible), as the scaling of the Hilbert space makes such a simulation prohibitive even for small environments. This implies that, in order to study the effect of the cou-

pling with the environment, one needs to find a reduced model that (hopefully) captures the interesting features of the interaction.

A common method to reduce the joint system-environment model is known in the physics literature with the name of time-convolutionless master equations (TCL ME) [?,?]. This method aims at obtaining an LTV model of the same size of the system that replicates the effects of the interaction with the environment. Although a formal exact solution to this problem exists, computing this exact LTV generator is generally considered as difficult as solving the original model. The main approach found in the physics literature approximates this formal solution by leveraging an expansion in the strength of the coupling between the system and the environment, which is typically assumed to be small. For more details, see [?] and references therein.

In this work an alternative approach is proposed: instead of an expansion in the coupling strength, an expansion in time is considered. In particular, since the formal solution provides an analytic LTV generator in the time variable t , the exact incomputable generator is approximated with its truncation to a finite order N . Under these assumptions a recursive formula for the coefficients of the polynomial generator is found, and the approximation error of the resulting reduced model scales as

Email address: `tommaso.grigoletto@unipd.it`
(Tommaso Grigoletto).

$O(t^{N+1})$.

Note that while time-dependent dynamical systems are typically more difficult to study than time-independent ones, this approach finds an important application in the simulation of very large systems, as the Runge-Kutta methods allow one to simulate time-dependent models [?]. In this regard, under the assumption of a time-dependent generator that is *polynomial* in t , this paper proposes a novel recursive formula to compute the propagator of this LTV model.

Another benefit of the proposed reduction method consists in the fact that, when computing the reduced LTV generator for unitarily evolving quantum systems, the reduced polynomial generator at first and second order is provably in Lindblad form for all times t thus ensuring that the reduced model at the lowest orders generates a completely positive (CP) and trace preserving (TP) semigroup.

It is important to point out that the method proposed in this work is not intended to substitute the perturbative approach for approximating TCL MEs [?,?] but rather to provide an alternative that can be of use whenever the main assumptions of the standard method do not hold, e.g. in the strong-coupling regime or whenever the time-ordered integrals are difficult to assess.

Structure and paper contributions: For the reader's convenience, the structure of this work and its main contributions are summarized next:

- In Sec. ?? the problem treated in this work is defined rigorously. The known formal solution of the problem is derived in detail using the Nakajima-Schwinger equations [?,?], and the key difficulties associated with computing this LTV model are explained.
- In Sec. ?? the main contribution of the paper is presented: Theorem ?? derives the polynomial approximation of the formal LTV model and the recursive equation that determines its coefficients. This result is based on Theorem ?? where a novel method to compute the time-ordered exponential is derived. The reduction procedure is also validated on a linear model and compared against the truncation of the exponential series.
- Sec. ?? focuses on the application of the proposed reduction procedure to quantum dynamical models. The proposed method is connected and compared with the typical physics approach for the approximation of TCL ME which is based on the expansion of the coupling strength. The connections between the recursive formula obtained here and other similar recursive equations known in the physics literature [?,?,?,?] are discussed, highlighting how the expansion in t greatly simplifies the associated calculations. Most importantly, leveraging a few recent results [?,?] it is proven that when the original generator is purely

Hamiltonian, the proposed approximation at first and second order provides a CPTP two-parameter semigroup.

- In Sec. ?? the proposed procedure is validated against prototypical quantum examples: a central spin model and an Ising spin chain. Note that the majority of the applications presented in this work are in the context of quantum physics since, up to this point, this is where these LTV-based model reduction techniques have found most applications.

Throughout the paper, pedagogical examples are included to intuitively explain the most involved proofs. Furthermore, both the strengths and weaknesses of the proposed approximation procedure are highlighted in detail, namely: the recursive formula used to compute the terms of the generators at different orders makes the procedure numerically easier to access at higher orders than the alternatives; allows for an initial model that is in Lindblad form (not only purely Hamiltonian); does not require the model to be put in rotating frame; and does not rely on a weak coupling assumption, thus allowing to study systems in strong coupling regimes. On the other hand, the proposed reduction procedure relies on an expansion in time and thus provides a good approximation only for small times. Although this aspect is limiting, similar restrictions also hold in the well-known approximations of the TCL ME proposed by [?].

2 Problem setting and connection with the literature

2.1 Problem setting

In this work, we assume to be given an LTI stable autonomous model of the type

$$\dot{x}_t = Lx_t \quad (1)$$

where $x_t \in \mathbb{C}^n$ is the system's state and $L \in \mathbb{C}^{n \times n}$ is the state matrix or generator. Such a generator defines a one-parameter semigroup, i.e. $\{\Lambda_t \equiv e^{Lt}\}_{t \geq 0}$ such that $\Lambda_{t+s} = \Lambda_t \Lambda_s$, for all $t, s \geq 0$.

We next assume that, for some reason dictated by the application at hand, we are not interested in reproducing all the degrees of freedom contained in the system's state x_t , but we are only interested in the degrees of freedom contained on a given subspace $\mathcal{V} \subseteq \mathbb{C}^n$ of dimension $m = \dim(\mathcal{V})$. Importantly, here we assume the subspace to be *given and fixed* $\mathcal{V}$. Let then $P \in \mathbb{C}^{n \times n}$, be a projector $P^2 = P$ onto $\mathcal{V} = \text{Im}P$. Note that P does not need to be orthogonal (with respect to the standard inner product $\langle \cdot, \cdot \rangle$), i.e. $P^\dagger \neq P$. For future convenience, let us define $Q \equiv \mathbb{1} - P$ where $\mathbb{1}$ denotes the identity matrix and factorize P into two linear maps: the *reduction* $R \in \mathbb{C}^{m \times n}$ and the *injection* $J \in \mathbb{C}^{n \times m}$ such that $P = JR$ and $RJ = \mathbb{1}_m$ the identity in $\mathbb{C}^{m \times m}$.




Figure 1. Graphical representation of the approximation problem. The pale blue plane represents the subspace $\mathcal{V}$ onto which P projects. The red curve represents the trajectory of the original model x_t , which starts in $\mathcal{V}$ at $t = 0$. The blue curve represents the projection of the true trajectory Px_t , i.e. the one we want to approximate. Finally, the orange curve represents the approximate evolution we aim to find, i.e. Jz_t .

We further make a simplifying assumption that is quite typical in these settings; see, e.g. [?]: We assume that $x_0 \in \mathcal{V}$, or equivalently $Px_0 = x_0$.

In summary, we have the model

$$\begin{cases} \dot{x}_t = Lx_t \\ y_t = Rx_t \end{cases}, \quad x_0 \in \mathcal{V}$$

where we defined the output signal $y_t \in \mathbb{R}^m$.

In this work, we are interested in deriving a reduced model that is capable of reproducing the trajectories $y_t = Pe^{Lt}x_0$, $\forall x_0 \in \mathcal{V}$, $\forall t \geq 0$ or, equivalently,

$$y_t = Re^{Lt}J\tilde{x}_0, \quad \forall \tilde{x}_0 \in \mathbb{C}^m, \quad \forall t \geq 0. \quad (2)$$

Note that, when $\mathcal{V}$ is L -invariant, the problem has a trivial solution since by projecting the generator L onto $\mathcal{V}$ one obtains the reduced generator RLJ that exactly reproduces all the trajectories of interest. One can obtain an exact and minimal LTI reduced model by computing the orthogonal to the non-observable subspace $\mathcal{N}^\perp$ as the smallest $L^\dagger$ -invariant subspace that contains $\mathcal{V}$ and projecting the dynamics onto it, thus obtaining the smallest observable LTI model [?]. However, the drawback of this approach is the fact that if the couple (L, R) is observable, the model can not be reduced exactly. In those cases, one should recur to approximate model reduction techniques such as balanced truncation [?,?] and

moment matching [?,?]. In those cases, one aims to find a subspace where to project the dynamics in order to obtain an LTI model that approximates the trajectories of interest in a given metric.

In this work, we take a different route: We fix the subspace where the dynamics is projected to $\mathcal{V}$ and lift the time-independence constraint by looking for an LTV model of dimension m that approximates the trajectories of interest. Note that while this approach is not common in the control literature, this has been studied in the physics literature for the derivation of TCL ME as detailed in Sec. ??.

More rigorously, let us define a state $z_t \in \mathbb{C}^m$ whose evolution is given by an LTV model of the type

$$\begin{cases} \dot{z}_t = F_t z_t \\ z_0 = Rx_0 \end{cases} \quad (3)$$

where $F_t \in \mathbb{C}^{n \times n}$ is a time-dependent generator. The evolution of such an LTV model is given by the two-parameter semigroup $\{\Upsilon_t \equiv \mathbb{T}e^{\int_0^t F_s ds}\}_{t \geq 0}$ for which the composition rule $\Upsilon_{t,0} = \Upsilon_{t,s}\Upsilon_{s,0}$ for all $0 \leq s \leq t$ holds. Here $\mathbb{T}$ denotes the time-ordering operator, and $\mathbb{T}e$ denotes the time-ordered exponential or, equivalently, the Volterra series [?,?].

Our aim is then to construct F_t so that $z_t \approx y_t$ or, in other words, we want to determine F_t so that

$$\mathbb{T}e^{\int_0^t F_s ds} \approx Re^{Lt}J.$$

The sense in which the approximation $\approx$ is intended shall be defined more precisely later on in the paper. A graphical representation of the approximation problem is given in Fig. ??.

2.2 Reduced and exact time-dependent dynamics

Before moving on to the method proposed in this work for computing the time-dependent generator F_t we shall discuss the common derivation found in the physics literature to solve the problem at hand [?,?,?,?]. The derivation is particularly elegant hence we report it here for the interested reader.

Dividing the evolution of x_t in the evolution of Px_t and that of Qx_t we have

$$\frac{d}{dt}Px_t = PLPx_t + PLQx_t, \quad (4)$$

$$\frac{d}{dt}Qx_t = QLPx_t + QLQx_t. \quad (5)$$

Solving the second equation for Qx_t with the Lagrange equation, we obtain

$$Qx_t = e^{QLQt}Qx_0 + \int_0^t e^{QLQs}QLPx_{t-s}ds.$$

By substituting this equation into Eq. (??) we obtain the famous Nakajima-Zwanzig equation [?,?]:

$$\begin{aligned} \frac{d}{dt}Px_t &= PLPx_t + PL e^{QLQt}Qx_0 \\ &+ \int_0^t PLQe^{QLQs}QLPx_{t-s}ds. \end{aligned} \quad (6)$$

We then recall that $x_{t-s} = e^{-Ls}x_t$ so that

$$Qx_t = e^{QLQt}Qx_0 + \int_0^t e^{QLQs}QLPe^{-Ls}ds \cdot (P+Q)x_t.$$

Now, defining

$$M_t \equiv \mathbb{1} - \int_0^t e^{QLQs}QLPe^{-Ls}ds \quad (7)$$

we have

$$M_t Qx_t = e^{QLQt}Qx_0 + (\mathbb{1} - M_t)Px_t.$$

If we then assume that M_t is invertible, we obtain

$$Qx_t = M_t^{-1}e^{QLQt}Qx_0 + (M_t^{-1} - \mathbb{1})Px_t.$$

Substituting this into Eq. (??) we obtain

$$\begin{aligned} \frac{d}{dt}Px_t &= PLPx_t + PLM_t^{-1}e^{QLQt}Qx_0 \\ &+ PL(M_t^{-1} - \mathbb{1})Px_t \\ &= PLM_t^{-1}e^{QLQt}Qx_0 + PLM_t^{-1}Px_t. \end{aligned} \quad (8)$$

Note that under the assumption $Px_0 = x_0$ we have $Qx_0 = 0$ and thus Eq. (??) gets simplified to

$$\frac{d}{dt}Px_t = \underbrace{PLM_t^{-1}P}_{\equiv K_t} \rho_t \quad (9)$$

which, separating the projector P into its factors $P = JR$, becomes

$$\frac{d}{dt}\check{x}_t = \underbrace{RLM_t^{-1}J}_{\equiv F_t} \check{x}_t. \quad (10)$$

A few comments on this solution are in order:

i) The invertibility of M_t needs to be verified and typically only holds for small t , (see, e.g. [?]) but for the

times where M_t is invertible, Eq. (??) provides an exact solution to the problem at hand.

ii) Note that, to compute F_t (or equivalently K_t) one needs to first compute the integral that defines M_t and then compute the inverse of M_t . Both these tasks are challenging (numerically and analytically) and it is generally considered that solving this integral and inverse is as difficult as solving the original model (??), see e.g. [?]. However, Eq. (??) provides an excellent starting point for computing an approximation of the reduced evolution.

To avoid these numerical issues, K_t is typically approximated by expanding it as an expansion in the coupling strength ε and by computing the terms of the expansion as ordered cumulants [?, ?, ?]. In many cases of interest, this approximation provides a good approximation of the open quantum system dynamics, both in the long-time and short-time regimes. It is, however, important to note that there is no guarantee that such an approximation holds also for long times. As an example of this, see [?] where it is shown that TCL2 and TCL4 (approximations at orders $N = 2, 4$) do not approximate the true dynamics in the long-time regime. We refer the reader to Sec. ?? for further connections to the physics literature.

3 Polynomial approximations of the time-dependent generator

In this section we derive the key result of this work: a method to approximate the time-dependent generator defined in Eq. (??), F_t . The derivation that follows is based on the following observation.

Proposition 1 *Let M_t be the time-dependent operator defined in Eq. (??) and let F_t be the time-dependent generator defined in Eq. (??). Then, for every t such that M_t is invertible, F_t is analytic in t , i.e.*

$$F_t = \sum_{k=0}^{+\infty} t^k F_{(k+1)}.$$

Proof. We can start by observing that the function $e^{QLQt}QLPe^{-Lt}$ is the product of two analytic functions in t hence it is analytic in t . Furthermore, as the integral of an analytic function is also analytic, see, e.g. [?] we have that M_t is also analytic in t . Finally, as the inverse of an analytic function $f(t)$ is analytic where $f(t) \neq 0$, we have that F_t is also analytic in t whenever M_t is invertible. $\square$ Note

that we label the coefficients in the power series starting from 1 instead of 0 for later convenience. This insightful result allows us to: 1) focus on computing the terms of the power series $\{F_{(k+1)}\}_{k=0}^{+\infty}$ and 2) approximate F_t by truncating the infinite power series to a finite order N .

In what follows, we thus approximate F_t with the *polynomial time-dependent generator*

$$F_{t,N} \equiv \sum_{k=0}^N t^k F_{(k+1)}.$$

Other than an obvious computational benefit introduced by computing the terms $F_{(k)}$ only to order N , the assumption that the generator is polynomial in the time variable greatly simplifies the computation of the time-ordered exponential $\mathbb{T}e^{\int_0^t F_{s,N} ds}$ as shown in the next key result.

Theorem 2 *Let us assume that $F_{t,N} = \sum_{k=0}^N t^k F_{(k+1)}$. Then the time-ordered exponential can be written as*

$$\mathbb{T}e^{\int_0^t F_{s,N} ds} = \sum_{k=0}^{+\infty} t^k E_{(k)} \quad (11)$$

where $E_{(0)} = \mathbb{1}$ and the other terms are defined by the recursion:

$$E_{(k)} = \frac{1}{k} \sum_{s=1}^k F_{(s)} E_{(k-s)} \quad (12)$$

and where we used the convention $F_{(k)} = 0$ for all $k > N$.

Proof of this result and an explanatory example are given in Appendix ??.

Note that Eq. (??) provides a closed formula for the coefficients $E_{(k)}$ of the expansion of $\mathbb{T}e^{\int_0^t F_{s,N} ds}$ as a function of the coefficients $F_{(k)}$. Furthermore, only the terms $\{E_{(\ell)}\}_{\ell=1}^{k-1}$ appear in Eq. (??), which means that this equation can be used iteratively to compute each $E_{(k)}$.

Theorem ?? provides the main intuition on how to compute the coefficients $\{F_{(k)}\}_{k=1}^{N+1}$ of the polynomial time-dependent generator $F_{t,N}$: Eq. (??) provides an expansion for $\mathbb{T}e^{\int_0^t F_{s,N} ds}$ in the time variable t . We can then expand e^{tL} in t and construct $F_{(k)}$ so that the terms of $\mathbb{T}e^{\int_0^t F_{s,N} ds}$ match the terms of $Re^{tL}J$ by orders of t . This approach is similar in principle to the adiabatic elimination techniques [?,?] where, instead of matching orders of ε , we match orders of t . The resulting procedure, which represents the main contribution of this work, is stated next.

Theorem 3 *Let us define*

$$F_{t,N} \equiv \sum_{k=0}^N t^k F_{(k+1)} \quad (13)$$

where

$$F_{(k)} \equiv k \frac{RL^k J}{k!} - \sum_{h=1}^{k-1} F_{(k-h)} \frac{RL^h J}{h!}. \quad (14)$$

Then:

$$Re^{Lt}J - \mathbb{T}e^{\int_0^t F_{s,N} ds} = O(t^{N+1}). \quad (15)$$

Proof. First of all we can use Theorem ?? to write

$$\mathbb{T}e^{\int_0^t F_{s,N} ds} = \sum_{f=0}^{+\infty} t^f E_{(f)}$$

where $E_{(k)} = \frac{1}{k} \sum_{s=1}^k F_{(s)} E_{(k-s)} = \frac{1}{k} \sum_{h=0}^{k-1} F_{(k-h)} E_{(h)}$ and where we reversed the order of the summation by imposing $h = k - s$. We next prove that, for $1 \leq k \leq N$, we have $E_{(k)} = \frac{RL^k J}{k!}$ by induction. The case $k = 1$ is trivial as $E_{(1)} = F_{(1)} E_{(0)} = RLJ$. Let then assume that $E_{(h)} = \frac{RL^h J}{h!}$ for all $h < k$ and let us prove that $E_{(k)} = \frac{RL^k J}{k!}$. Then

$$\begin{aligned} E_{(k)} &= \frac{1}{k} \sum_{h=0}^{k-1} F_{(k-h)} E_{(h)} = \frac{F_{(k)}}{k} + \frac{1}{k} \sum_{h=1}^{k-1} F_{(k-h)} E_{(h)} \\ &= \frac{F_{(k)}}{k} + \frac{1}{k} \sum_{h=1}^{k-1} F_{(k-h)} \frac{RL^h J}{h!} = \frac{RL^k J}{k!} \end{aligned}$$

where we simply substituted the definition of $F_{(k)}$ given in eq. (??).

By leveraging the definition of the exponential map e^{Lt} we can expand the term $Re^{Lt}J$ to obtain

$$Re^{Lt}J = \sum_{k=0}^{+\infty} \frac{t^k}{k!} RL^k J.$$

Combining the two terms we obtain

$$\begin{aligned} Re^{Lt}J - \mathbb{T}e^{\int_0^t F_{s,N} ds} &= \sum_{k=0}^{+\infty} t^k \left(\frac{RL^k J}{k!} - E_{(k)} \right) \\ &= \sum_{k=0}^N t^k \left(\frac{RL^k J}{k!} - E_{(k)} \right) \\ &\quad + O(t^{N+1}) \end{aligned}$$

which concludes the proof. $\square$

We shall observe a few key aspects. First, Eq. (??) finally gives sense to the approximation symbol $\approx$ used in Sec.




Figure 2. Operator and Hilbert-Schmidt norm of the first 20 $F_{(k)}$ terms (left) and of the first 50 $E_{(k)}$ terms when $N = 20$ (right).

?? i.e. an approximation of the type $O(t^{N+1})$. This implies that the approximation holds only for small times. Second, it is important to notice that the right-hand side of equation (??) depends on $RL^k J$ and $\{F_{(s)}\}_{s=1}^{k-1}$. This means that equation (??) can be used iteratively to construct the terms $F_{(k)}$ starting from $k = 1$ and growing up to $k = N$. For the reader's convenience the first few terms of the series are reported next:

$$\begin{aligned} F_{(1)} &= RLJ \\ F_{(2)} &= RL^2 J - F_{(1)} RLJ = RL^2 J - (RLJ)^2 \\ F_{(3)} &= \frac{RL^3 J}{2} - F_{(2)} RLJ - F_{(1)} \frac{RL^2 J}{2} \\ &= \frac{RL^3 J}{2} - \frac{F_{(1)} F_{(2)}}{2} - \frac{F_{(1)}^3}{2} - F_{(2)} F_{(1)} \end{aligned}$$

It is interesting to notice a couple of points about the first few terms: i) As one would expect, the term of order one $F_{(1)}$ coincides with the reduction of the original dynamics which is often used as rough time-independent approximation of the reduced evolution [?]; ii) The second order term does not only contain the reduction of the generator squared $RL^2 J$ but also contains a correction term depends on the first order term $F_{(1)}$.

Lastly, it is important to note that similar (yet distinct) recursive formulas to derive similar approximations have been found in the physics literature [?, ?, ?, ?]. We refer the reader to Sec.?? where we discuss connections and differences between these works.

3.1 Linear testbed

To showcase the capability of the proposed method, we here consider a simple numerical example. Specifically, we consider a random autonomous and stable system of size $n = 20$. Let then A be a random uniform matrix in $\mathbb{C}^{n \times n}$, i.e. $[A]_{i,j} \sim \mathcal{U}([0, 1])$ and let $B = A - \lambda_{\text{Max}} \mathbb{1}_n$ where λ_{Max} is the absolute value of largest eigenvalue of A . The dynamic's generator $L \in \mathbb{C}^{n \times n}$ is then obtained as $L = \frac{1}{2} \frac{B}{\|B\|_{op}}$ where $\|\cdot\|_{op}$ denotes the operator norm.

This ensures that L is simply stable and has operator norm $\|L\|_{op} = \frac{1}{2}$.

The projector P is then constructed through $R = \begin{bmatrix} \mathbb{1}_{m \times m} & 0_{m \times n-m} \end{bmatrix}$ and $J = R^\dagger$ and $P = RJ$.

By using Eq. (??) we can then compute the terms $F_{(k)}$ iteratively for $k = 1, \dots, N$. Fig. ?? (left) shows both the Hilbert-Schmidt and operator norm of the terms $F_{(k)}$ for $k = 1, 20$. Interestingly, the Hilbert-Schmidt norm, $\|F_{(k)}\|_{HS}$, is bounded by $\frac{\|L\|_{HS}^k}{k!}$, indicating that the series $F_{t,N}$ should be converging for $N \rightarrow \infty$. Curiously, the same does not hold for the operator norm. Fig. ?? (right) instead shows the norm of the terms $E_{(k)}$ where we fixed $N = 20$ and $F_{(k)} = 0 \forall k > 20$. Interestingly, for $k \leq N$ $\|E_{(k)}\|$ is bounded by $\frac{\|L\|_{op}^k}{k!}$ while for $k > N$ this is no longer the case.




Figure 3. The continuous lines represent the error norm between the exact reduced evolution Rx_t and the approximated reduced evolution z_t for different values of the approximation order N . The dashed curves represent an approximation obtained by truncating the Taylor expansion of the exponential for different values of the truncation order N .

Fig. ?? shows the error norm $\|Rx_t - z_t\|_2$ versus time. In this simulation Rx_t was simulated by reducing the original model, i.e. $Rx_t = Re^{Lt}x_0$ while z_t is obtained by simulating the time-dependent linear differential equa-

tion $\dot{z}_t = F_{t,N} z_t$, for different values of N , by using an approximation of the equation given in Theorem ??, i.e. $\mathbb{T}e^{\int_0^t F_{s,N} ds} \approx \sum_{k=0}^{100} t^k E_{(k)}$. It can be seen that increasing the degree of approximation N decreases the error committed by the approximated model.

An alternative natural numerical approximation that comes to mind is given by truncating the exponential expansion at a given order N , i.e. $Rx_t \approx \sum_{k=0}^N \frac{t^k}{k!} RL^k J Rx_0$. Note that such an approximation has numerical complexity almost identical to that of the approximation proposed in this work, as in both cases one needs to compute L^N . Furthermore, from a theoretical point of view, both cases commit an error of the order of $O(t^{N+1})$. To test this fact numerically, in Fig. ?? we included the actual error committed by these two approximations at the same orders of approximation: the continuous line denotes the error committed by the procedure proposed here, while the dashed line is the error committed by the truncation of the exponential. One can observe that for lower orders ($N = 1, 2, 5$) the procedure proposed here yields better results, whereas at higher orders this is not the case.

4 Ensuring valid quantum models at lower orders

In this section, we apply the results derived in the Sec. ?? to quantum dynamical models, the application that originally inspired this work.

4.1 The reduction problem for quantum dynamical models

Let us consider a quantum system defined on a Hilbert space $\mathcal{H}$, and let $\mathfrak{B}(\mathcal{H})$ denote the set of bounded operators acting on $\mathcal{H}$. In this work, we only consider finite-dimensional systems, i.e. $\dim(\mathcal{H}) < \infty$, $\mathcal{H} \simeq \mathbb{C}^n$ and $\mathfrak{B}(\mathcal{H}) \simeq \mathbb{C}^{n \times n}$. The state of such a system is described by a density operator $\rho \in \mathfrak{D}(\mathcal{H})$ with $\mathfrak{D}(\mathcal{H}) = \{\rho \in \mathfrak{B}(\mathcal{H}) | \rho = \rho^\dagger \geq 0, \text{tr} \rho = 1\}$. Here, we assume that the state of the system evolves according to the Von-Neumann master equation

$$\dot{\rho}_t = \mathcal{L}(\rho_t) \quad (16)$$

where $\rho_0 \in \mathfrak{D}(\mathcal{H})$ is the system's initial condition, and $\mathcal{L}$ is the celebrated Lindblad (or GKLS) generator [?,?]:

$$\mathcal{L}(\rho) = -i[H, \rho] + \sum_k \mathcal{D}_{L_k}(\rho) \quad (17)$$

$H = H^\dagger$ is the system's Hamiltonian and $\mathcal{D}_L(\rho) = L\rho L^\dagger - \frac{1}{2}\{L^\dagger L, \rho\}$ represents the system's dissipation toward a Markovian environment represented by a set of operators $\{L_k\}$ called noise operators. The evolution

of the quantum system is then described by the exponential map $\rho_t = e^{\mathcal{L}t}(\rho_0)$. Importantly, $e^{\mathcal{L}t}$ generates a quantum dynamical semigroup, i.e. $\{\Lambda_t \equiv e^{\mathcal{L}t}\}_{t \geq 0}$ a one-parameter semigroup of CPTP maps also known as Markovian dynamics. In the following, we will say that a super-operator is of *Lindblad-type* if it can be put in the form of Eq. (??).

Note that in general, in the derivation of TCL ME [?,?], the initial “big” model is considered to be Hamiltonian, i.e. $\mathcal{L}(\cdot) = -i[H, \cdot]$. However, mathematically, there is no reason to restrict to this case; hence, in this work, we start with the general assumption that $\mathcal{L}$ is a generator of Lindblad form and then focus on the Hamiltonian case when necessary.

In many situations of interest in quantum physics, relevant projectors are represented by CPTP ones. We thus assume to be given a CPTP, not necessarily orthogonal (w.r.t. $\langle \cdot, \cdot \rangle_{HS}$), projector $\mathcal{P} : \mathfrak{B}(\mathcal{H}) \rightarrow \mathfrak{B}(\mathcal{H})$, $\mathcal{P}^2 = \mathcal{P}$, whose image will be denoted by $\mathcal{V} \equiv \text{Im} \mathcal{P}$. This provides a generalization to the superoperator projector setting considered by [?,?]. Furthermore, for the sake of simplicity, we assume that $\mathcal{P}$ admits¹ a full rank fixed point, i.e. $\exists \rho \in \mathfrak{D}(\mathcal{H})$, $\rho > 0$ such that $\mathcal{P}(\rho) = \rho$. As we shall see in more detail later (in Sec.??) such a projector is, in fact, a conditional expectation in quantum probability theory [?,?] and enjoys some useful and well known properties [?,?,?,?,?]. However, for now it is sufficient to know that $\mathcal{P}$ can be decomposed into two factors $\mathcal{P} = \mathcal{J}\mathcal{R}$ which take the name of reduction and injection maps $\mathcal{R} : \mathfrak{B}(\mathcal{H}) \rightarrow \mathfrak{B}(\mathcal{K})$ and $\mathcal{J} : \mathfrak{B}(\mathcal{K}) \rightarrow \mathfrak{B}(\mathcal{H})$ such that $\mathcal{R}\mathcal{J} = \mathcal{I}_{\mathcal{K}}$ the identity superoperator, $\text{Im} \mathcal{J} = \mathcal{V}$, $\text{Im} \mathcal{R} = \mathcal{A}$ and where $\mathcal{A} \subseteq \mathfrak{B}(\mathcal{K})$ is a $*$ -subalgebra of $\mathfrak{B}(\mathcal{K})$ for some Hilbert space $\mathcal{K}$, $\dim(\mathcal{K}) \leq \dim(\mathcal{H})$.

To provide some intuition about these mathematical objects, we anticipate the reduction and injection maps we are going to use in the applications of Sec. ??:

- Assume a bipartite case, i.e. $\mathcal{H} = \mathcal{H}_S \otimes \mathcal{H}_B$. We can then consider $\mathcal{P}(\rho) = \text{tr}_B(\rho) \otimes \tau$ for some $\tau \in \mathfrak{D}(\mathcal{H}_B)$, $\tau > 0$ to be our CPTP projector. Then, by choosing $\mathcal{K} = \mathcal{H}_S$ and $\mathcal{A} = \mathfrak{B}(\mathcal{H}_S)$ we have the maps $\mathcal{R}(\rho) = \text{tr}_B(\rho)$ and $\mathcal{J}(\mu) = \mu \otimes \tau$.
- Another CPTP projector we might want to consider is the projector onto the diagonal $\mathcal{P}(\rho) = \sum_{j=0}^{n-1} |j\rangle\langle j| \rho |j\rangle\langle j|$, in which case $\mathcal{K} = \mathcal{H}$, $\mathcal{A}$ is the commutative algebra given by $\mathcal{A} \equiv \text{span}\{|j\rangle\langle j|, j = 0, \dots, n-1\} \simeq \mathbb{C}^n$. $\mathcal{R}(\rho) = \sum_{j=0}^{n-1} |j\rangle\langle j| \rho |j\rangle\langle j|$ so that $\mu = \mathcal{R}(\rho) \in \mathbb{C}^n$, and $\mathcal{J}(\mu) = \sum_{j=0}^{n-1} |j\rangle\langle j| \mu |j\rangle\langle j|$ factorize $\mathcal{P}$.

¹ Note that this assumption is made for simplicity of presentation and can easily be lifted by restricting onto the support of $\text{Im} \mathcal{P}$, see e.g. [?].

The reduction map $\mathcal{R}$ determines the reduced state, i.e. the one we want to keep track of. Let $\eta_t \in \mathfrak{D}(\mathcal{K})$ be defined as $\eta_t = \mathcal{R}(\rho_t)$ for all $t \geq 0$. Under the assumption $\mathcal{P}(\rho_0) = \rho_0$, the evolution of the reduced state is thus given by

$$\eta_t = \mathcal{R}e^{\mathcal{L}t}(\rho_0) = \mathcal{R}e^{\mathcal{L}t}\mathcal{J}(\eta_0).$$

Theorem ?? then offers a method to approximate this evolution with a time-dependent generator $\mathcal{F}_{t,N}$ that is polynomial in time t , i.e. $\mathcal{F}_{t,N} = \sum_{k=0}^N t^k \mathcal{F}_{(k+1)}$ and such that

$$\left\| \mathbb{T}e^{\int_0^t \mathcal{F}_{s,N} ds} - \mathcal{R}e^{\mathcal{L}t}\mathcal{J} \right\| = O(t^{N+1}).$$

Desirable properties for $\mathcal{F}_{t,N}$ certainly include the fact that we would like the semigroup $\{\Upsilon_t \equiv \mathbb{T}e^{\int_0^t \mathcal{F}_s ds}\}_{t \geq 0}$ to be a set of CPTP maps for all $t \geq 0$. Note that necessary conditions to impose on $\mathcal{F}_t$ so that $\mathbb{T}e^{\int_0^t \mathcal{F}_s ds}$ is CPTP for all $t \geq 0$ are not known [?]. Nevertheless, in the following subsections, we shall see that, under mild assumptions, the first two orders of the approximation ensure that this desirable property is preserved.

4.2 Connection with the physics literature

In the typical setting of TCL ME derivation, the system is considered to be bipartite, i.e. $\mathcal{H} = \mathcal{H}_S \otimes \mathcal{H}_B$ and the dynamics purely Hamiltonian, i.e. $\mathcal{L}(\rho) = -i[H, \rho]$ with Hamiltonian $H = H_0 + \varepsilon V$ where H_0 is the free Hamiltonian of the system and bath, V is the interaction Hamiltonian and ε is the coupling strength. Under these assumptions, and assuming that the coupling strength is small, the time-dependent generator $\mathcal{K}_t$ defined in Eq. (??) is computed by performing a perturbative expansion in the coupling parameter. The time-dependent generator is then expressed as

$$\mathcal{K}_t = \sum_{n=1}^{\infty} \mathcal{K}_n(t)$$

where

$$\begin{aligned} \mathcal{K}_n(t) &= \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-2}} dt_{n-1} \\ &\quad \times \sum_q (-1)^q \mathcal{P} \mathcal{L}_t \cdots \mathcal{L}_{t_i} \mathcal{P} \mathcal{L}_{t_j} \cdots \mathcal{L}_{t_k} \mathcal{P} \cdots \mathcal{P} \end{aligned}$$

and where $\mathcal{L}_t = -i[H(t), \cdot]$ and $H(t)$ is the Hamiltonian in rotating frame with respect to H_0 . The series is then approximated by stopping at a given order N , i.e. $\mathcal{K}_t \approx \sum_{n=1}^N \mathcal{K}_n(t)$.

Under these assumptions, a recursive formula similar to Eq. (??) has been derived. Specifically [?, Eq. (38)] (but also [?, Eq. (2)] [?, Eq. (2)], [?, Eq. (16)]) provides a recursive formula for the terms $\mathcal{K}_n(t)$ of the time-dependent generator $\mathcal{K}_t$. However, these recursive formulas were derived in the rotating frame (w.r.t. H_0) under a weak coupling assumption and thus from an initial time-dependent model. This implies that the terms of the recursive formula contain time-ordered integrals that need to be solved to compute the terms of the expansion.

In contrast, in this work, we start from a time-independent model, hence Eq. (??) (which was derived independently from [?, ?, ?, ?]) does not contain integrals, resulting in simpler computations.

Furthermore, the main difference that separates this work from most of the physics literature, see e.g. [?, ?, ?] is the fact that, instead of expanding the time-dependent generator in a power series of a perturbation parameter ε , we expand in the time variable t . This difference allows for a few benefits: It simplifies the calculations of the approximations and lifts the weak-coupling assumption, thus allowing one to explore the effects of strong-coupling regimes.

The main drawback introduced by the expansion in the time variable t is the fact that the approximate model holds only for small times. Nevertheless, as we highlighted above, while this consequence is clearly limiting, this situation is comparable to the typical derivation of TCL ME since no guarantees on the long-time limit are available. In contrast, in this work we were able to characterize how the error between the original and reduced models grows with t .

4.3 Complete positivity of the first order

In this section, we recall relevant known results about CPTP projectors, their images, and reduced dynamics onto distorted algebras from the literature [?, ?, ?, ?, ?, ?]. The consequences of the assumptions we made about the projector $\mathcal{P}$ are summarized in the following proposition.

Proposition 4 *Let $\mathcal{P}$ be a CPTP projector with a full rank fixed point $\bar{\rho} > 0$, $\mathcal{P}(\bar{\rho}) = \bar{\rho}$ and let $\mathcal{V}$ be its image, i.e. $\mathcal{V} \equiv \text{Im} \mathcal{P} \subseteq \mathfrak{B}(\mathcal{H})$. Then there exist a decomposition of the Hilbert space*

$$\mathcal{H} = \bigoplus_k (\mathcal{H}_{F,k} \otimes \mathcal{H}_{G,k}),$$

a unitary operator $U \in \mathfrak{B}(\mathcal{H})$, and a set of density oper-

ators $\tau_k \in \mathfrak{D}(\mathcal{H}_{G,k})$ such that

$$\mathcal{V} = U \left(\bigoplus_k \mathfrak{B}(\mathcal{H}_{F,k}) \otimes \tau_k \right) U^\dagger. \quad (18)$$

Furthermore, $\mathcal{V}$ is a $*$ -algebra closed with respect to the product $X \cdot_\sigma Y \equiv X \sigma Y$ where

$$\sigma = U \left(\bigoplus_k \mathbb{1}_{F,k} \otimes \tau_k^{-1} \right) U^\dagger.$$

Let then W_k be a linear operator from $\mathcal{H}$ to $\mathcal{H}_{F,k} \otimes \mathcal{H}_{G,k}$ such that $W_k W_k^\dagger = \mathbb{1}_{F,k} \otimes \mathbb{1}_{G,k}$ [?]. Then the CPTP projection $\mathcal{P}$ onto $\mathcal{V}$ takes the form

$$\mathcal{P}(X) = U \left[\bigoplus_k \text{tr}_{\mathcal{H}_{G,k}} (W_k X W_k^\dagger) \otimes \tau_k \right] U^\dagger. \quad (19)$$

and is orthogonal with respect to the inner product $\langle X, Y \rangle_{\sigma, \lambda} = \text{tr}[X^\dagger \sigma^\lambda Y \sigma^{1-\lambda}]$ for $\lambda \in [0, 1]$.

Proof of this result can be found in [?] and [?, ?, ?, ?, ?]. Note that projectors of this type take the name of state extensions and are the dual of conditional expectations [?, ?]. Furthermore, note that a $*$ -algebra closed with respect to a modified product often takes the name of *distorted algebra* [?], while the block structure provided in Eq. (??) takes the name of *Wedderburn decomposition*.

The previous result provides us with a structure for the image of a CPTP projection whenever there exists a full rank state $\bar{\rho}$ such that $\mathcal{P}(\bar{\rho}) = \bar{\rho}$. From this structure one can observe that a distorted algebra $\mathcal{V}$ is isomorphic to a smaller representation which we denote $\mathcal{A}$ where the weighted copies are removed, i.e. $\mathcal{A} \equiv \bigoplus_k \mathfrak{B}(\mathcal{H}_{F,k})$. This allows us to find the two factors $\mathcal{R} : \mathfrak{B}(\mathcal{H}) \rightarrow \mathcal{A}$ and $\mathcal{J} : \mathcal{A} \rightarrow \mathfrak{B}(\mathcal{H})$ such that, $\mathcal{R}\mathcal{J} = \mathcal{I}_{\mathcal{A}}$ and $\mathcal{P} = \mathcal{J}\mathcal{R}$. Furthermore, one can prove [?] that these two factors can be chosen to be CPTP, i.e. for all $X \in \mathfrak{B}(\mathcal{H})$

$$\mathcal{R}(X) = \bigoplus_k \text{tr}_{\mathcal{H}_{G,k}} (W_k X W_k^\dagger) = \bigoplus_k X_{F,k} = \check{X}, \quad (20)$$

and for all $\check{X} \in \mathcal{A}$, $\check{X} = \bigoplus_k X_{F,k}$,

$$\mathcal{J}(\check{X}) = U \left(\bigoplus_k X_{F,k} \otimes \tau_k \right) U^\dagger. \quad (21)$$

The reduction and injection maps given in equations (??) and (??) then allow us to compute the reduced dynamics $\mathcal{R}\mathcal{L}\mathcal{J}$ restricted to the algebra $\mathcal{A}$. Then we can prove that the reduced generator $\check{\mathcal{L}}$ is GKLS by leveraging the structure of the reduction and injection maps. Here we recall [?, Theorem 4] for completeness.

Theorem 5 Let $\mathcal{P}$ be a CPTP projection with a full rank fixed point, let $\mathcal{V}$ be its image $\mathcal{V} = \text{Im}\mathcal{P} \subseteq \mathfrak{B}(\mathcal{H})$, and let $\mathcal{R}$ and $\mathcal{J}$ denote the CPTP factorization of $\mathcal{P} = \mathcal{J}\mathcal{R}$, as defined in Eq. (??) and (??). Then for any Lindblad generator $\mathcal{L}$, its restriction to $\mathcal{A}$, $\mathcal{R}\mathcal{L}\mathcal{J}$, is also a Lindblad generator, that is, $\mathcal{R}\mathcal{L}\mathcal{J} : \mathcal{A} \rightarrow \mathcal{A}$ and $\{e^{\check{\mathcal{L}}t}\}_{t \geq 0}$ is a quantum dynamical semigroup.

Note that this theorem only proves that $\mathcal{R}\mathcal{L}\mathcal{J}$ is of Lindblad type. In order to compute the reduced Hamiltonian and noise operators, one can use the procedure described in [?, Appendix]. This result allows us to state that, under the assumptions described above, the approximation at first order $\mathcal{F}_{t,1} = \mathcal{R}\mathcal{L}\mathcal{J}$ is of Lindblad type and thus it is guaranteed to give a CPTP semigroup for all times.

4.4 Second order: The case of purely Hamiltonian evolution

While the results we derived up to this point hold for general Lindblad generators $\mathcal{L}$, we shall now focus on the special case of purely Hamiltonian evolution, i.e. $\mathcal{L}(\cdot) = -i[H, \cdot]$. Other than the fact that this assumption is interesting in practical cases, we shall see that under this simplifying assumption we are able to prove that the second order term $\mathcal{F}_{(2)}$ is also of Lindblad-type.

The next result is based on the fact that the squared of a purely Hamiltonian generator is equal to a dissipator:

$$\begin{aligned} \mathcal{L}^2(\rho) &= -[H[H, \rho]] = -(H^2\rho - 2H\rho H + \rho H^2) \\ &= 2H\rho H - \{H^2, \rho\} = \mathcal{D}_{\sqrt{2}H}(\rho). \end{aligned} \quad (22)$$

This fact allows us to prove the next result.

Theorem 6 Under the assumptions of Theorem ?? we have that $\mathcal{F}_{(2)} = \mathcal{R}\mathcal{L}^2\mathcal{J} - (\mathcal{R}\mathcal{L}\mathcal{J})^2$ is of Lindblad-type and for $\mathcal{F}_{t,2} \equiv \mathcal{F}_{(1)} + t\mathcal{F}_{(2)}$ we have that $\mathbb{V}e^{\int_0^t \mathcal{F}_{s,2} ds}$ forms a CPTP semigroup for all $t \geq 0$.

Proof. By leveraging [?, Proposition 5] we have that $\mathcal{R}\mathcal{L}\mathcal{J}(\check{\rho}) = -i[\check{H}, \check{\rho}]$ where $\check{H} = \mathcal{J}^\dagger(H)$ hence, using equation (??) we obtain $(\mathcal{R}\mathcal{L}\mathcal{J})^2(\check{\rho}) = \mathcal{D}_{\sqrt{2}\check{H}}(\check{\rho})$. On the other hand, by leveraging [?, Proposition 6 and Proposition 7], we have that $\mathcal{R}\mathcal{D}_{\sqrt{2}H}\mathcal{J}(\check{\rho}) = \sum_{\check{C} \in \mathcal{X}(\sqrt{2}H)} \mathcal{D}_{\check{C}}(\check{\rho})$ for a set of operators $\mathcal{X}(\sqrt{2}H) \subset \mathfrak{B}(\mathcal{K})$. Importantly, by [?, Corollary 4], we have that $\mathcal{J}^\dagger(H) \in \mathcal{X}(\sqrt{2}H)$ and thus we can write $\mathcal{R}\mathcal{D}_{\sqrt{2}H}\mathcal{J}(\check{\rho}) = \sum_{\check{C} \in \mathcal{X}(\sqrt{2}H)/\{\sqrt{2}\check{H}\}} \mathcal{D}_{\check{C}}(\check{\rho}) + \mathcal{D}_{\sqrt{2}\check{H}}(\check{\rho})$ since $\sqrt{2}\check{H} = \mathcal{J}^\dagger(\sqrt{2}H)$. Then, by combining these two results we obtain

$$\mathcal{F}_{(2)} = \mathcal{R}\mathcal{L}^2\mathcal{J} - (\mathcal{R}\mathcal{L}\mathcal{J})^2 = \sum_{\check{C} \in \mathcal{X}(\sqrt{2}H)/\{\sqrt{2}\check{H}\}} \mathcal{D}_{\check{C}}(\check{\rho})$$

which is the sum of a set of GKLS, purely dissipative generators and thus of Lindblad type. Combining this with Theorem ?? it trivially follows that $\mathcal{F}_{t,2}$ is of Lindblad type for all $t \geq 0$. $\square$

5 Applications to quantum models

In this section, we consider a few examples of interest:

I) A qubit coupled to a finite bath is studied. Given the fact that the bath is finite-dimensional, this is typically considered a “highly non-Markovian model”. This example allows us to showcase the numerical procedure at higher orders for both a case where the original dynamics is purely Hamiltonian and a case where a dissipative term is included.

III) An Ising spin chain is studied. This example allows us to show that the proposed procedure works in cases that are more general than the bipartite case.

5.1 Dissipative central spin model

In this section we consider a dissipative central spin model to showcase the capabilities of the proposed technique. Similar models have been studied in the literature, see e.g. [?, ?, ?, ?]. The model is composed by a system, i.e. $\mathcal{H}_S \simeq \mathbb{C}^2$ also denoted as spin 1 and a bath composed of N_B other spins, i.e. $\mathcal{H}_B \simeq \mathbb{C}^{2^{N_B}}$ counted from 2 to $N_B + 1$.

The system Hamiltonian is described as $H = H_S + H_B + H_{\text{int}}$ where:

$$\begin{aligned} H_S &= \delta(\sigma_x^{(1)} + \sigma_z^{(1)}) \\ H_B &= \frac{\gamma}{4} \left(2J_x^2 - \frac{N_B}{2} \mathbb{1} \right) \\ H_{\text{int}} &= \frac{\lambda}{2} \left(A_x \sigma_x^{(1)} J_x + A_y \sigma_y^{(1)} J_y + A_z \sigma_z^{(1)} J_z \right) \end{aligned}$$

where $\sigma_u^{(j)}$ denotes the $u \in \{x, y, z\}$ Pauli matrix acting on the j -th spin and $J_u = \frac{1}{2} \sum_{k=2}^{N_B+1} \sigma_u^{(k)}$ denotes the total bath angular momentum operators. The bath is also in contact with a Markovian environment that dissipates information. This interaction is described by the noise operators $L_k = \Lambda \sigma_-^{(k)}$ for all $k = 2, \dots, N_B + 1$ where $\sigma_{\pm}^{(k)} = \frac{1}{2}(\sigma_x^{(k)} \pm \sigma_y^{(k)})$.

For this example we consider as an initial condition the factorized state $\rho_0 = |+\rangle\langle+| \otimes \rho_B$ where

$$\rho_B = \frac{e^{-\beta H_B}}{\text{tr}[e^{-\beta H_B}]}$$

is the thermal state at inverse temperature β . This allows us to define the reduction and injection maps as $\mathcal{R}(\cdot) = \text{tr}_B(\cdot)$ and $\mathcal{J}(\cdot) = \cdot \otimes \rho_B$. Note that with this

choice we have that $\mathcal{P}(\rho_0) = \rho_0$. The reduced state then corresponds to the state of the central spin, i.e.

$$\eta_t = \rho_S(t) \equiv \text{tr}_B(\rho_t).$$

In Fig. ?? the results of the simulation for this dissipative central spin model are shown. The simulation has been run in two settings: a purely Hamiltonian setting ($\Lambda = 0$ on the left) and a dissipative setting ($\Lambda = 0.8$ on the right). The rest of the parameters used were $N_B = 3$, $\delta = 0.3$, $\lambda = 0.1$, $\gamma = 1$, $A_x = 1.2$, $A_y = 1.5$, $A_z = 1.3$ and $\beta = 50$. In the numerical simulations we can observe a few interesting aspects: 1) as expected, the error between the exact solution and the approximate solution $\|\eta_t - \mu_t\|_t$ decreases when increasing the approximation degree N ; 2) the trajectory for approximation orders $N = 1, 2$ and even 3 do not exit the Bloch sphere (note that we have proven the complete positivity for orders $N = 1, 2$ but not for $N = 3$); 3) the simulations for $N > 3$ are displaying Runge’s phenomena at larger times, which is quite typical in these situations; 4) after some time, the evolved state η_t for an approximation degree $N \geq 4$ exits the Bloch sphere, indicating that the dynamics is not CPTP at all times for approximation degrees higher than $N \geq 4$.

5.2 Ising spin chain

Although the idea of approximating a large model with a smaller one that is LTV is common in the setting of a bipartite system-environment couple, this is far from the only case where such an approach is useful. More in general, the same approaches can be adapted to cases where one wants to describe the evolution of only a small set of variables of interest of the original model. For this reason, in this section we showcase the use of the same methods presented before to reproduce the probability distribution in the standard basis.

In order to show that the proposed method works also in cases that are not bi-partite we here consider a spin-chain model. More specifically we consider a model composed of N_B spins, i.e. $\mathcal{H} \simeq \mathbb{C}^n$ where $n = 2^{N_B}$ whose dynamics is driven by the Ising Hamiltonian

$$H = \sum_{j=1}^{N_B} h \sigma_z^{(j)} + A \sum_{j=1}^{N_B-1} \sigma_x^{(j)} \sigma_x^{(j+1)}. \quad (23)$$

In this example, we consider $\mathcal{P}$ to be the CPTP projector on the diagonal $\mathcal{P}(\rho) = \sum_{j=0}^{n-1} |j\rangle\langle j| \rho |j\rangle\langle j|$ which can be factorized into $\mathcal{R}(\rho) = \sum_{j=0}^{n-1} |j\rangle\langle j| \rho |j\rangle\langle j|$ and $\mathcal{J}(\eta) = \sum_{j=0}^{n-1} |j\rangle\langle j| \langle j| \eta$. Note that, in such a case, $\eta = \mathcal{R}(\rho) \in \mathbb{R}^n$ is a probability distribution, i.e. $\eta_i \geq 0$ and $\mathbf{1}^T \eta = 1$ where $\mathbf{1} \in \mathbb{R}^n$ is the vector of all ones. This implies that $\mathcal{F}_t \in \mathbb{R}^{n \times n}$.




Figure 4. Simulations of the dissipative central spin model in the purely Hamiltonian setting ($\Lambda = 0$) on the left and in the dissipative setting ($\Lambda = 0.8$) on the right. The exact evolution and various degrees of approximations ($N = 1, 2, 3, 4, 10, 20$) are shown in each figure. The first row shows the trajectory of the state of the central state ρ_S in the Bloch sphere; the second row shows the magnetization of the central spin in the σ_z direction versus time; the third row shows the trace norm error between the exact trajectory η_t and the reduced one μ_t , i.e. $\|\eta_t - \mu_t\|_{\text{tr}}$ versus time for different values of the approximation N and with dotted curve representing a fitted αt^{N+1} curve. The black crosses denote points where the evolved state μ_t exits the set of density operators.

As an initial condition we consider $\rho_0 = |0\rangle\langle 0|$. Trivially, $\mathcal{P}(\rho_0) = \rho_0$.

Numerically, we verified that $\mathcal{F}_{(1)} = \mathcal{F}_{(3)} = 0$ while $\mathcal{F}_{(2)}$ is graphically represented in Fig. ?? where one can verify that $\mathcal{F}_{(2)}$ is Metzler (a matrix A is Metzler if $A_{i,j} \geq 0$ for all $i \neq j$) and is such that $\mathbf{1}^T \mathcal{F}_{(2)} = 0$. Those two conditions are the classical equivalent to the properties characterizing a Lindblad generator and in fact imply that $e^{\mathcal{F}_{(2)}t}$ is a stochastic matrix for all $t \geq 0$. This directly implies that $\mathbb{T}e^{\int_0^t \mathcal{F}_{s,2} ds}$ is CPTP, or stochastic in this case, for all $t \geq 0$.

Fig. ?? shows the results of the simulations of the Ising spin chain model with $N_B = 4$, $A = 0.3$, $h = 0.36$ where we can observe similar results to the other simulations included in this work: increasing the degree of approximation decreases the error $\|\eta_t - \mu_t\|_2$ at low times; for $N = 2$ the approximate evolved state μ_t remains a probability distribution, while for $N \geq 4$, after a certain time, the state is no longer a probability distribution.




Figure 5. Simulation of the Ising spin chain. Magnetization of the first spin of the chain $\langle \sigma_z^{(1)} \rangle$ versus time on the left and the error norm $\|\eta_t - \mu_t\|_2$ versus time on the right. Various approximation's degrees $N = 2, 4, 10, 20$ are shown in each figure. The black crosses denote the points where the evolved state μ_t exits the n -dimensional simplex, i.e. μ_t is no longer a probability distribution while the dotted curves on the right again represent fitted curves of αt^{N+1} .




Figure 6. Graphical representation of the term $\mathcal{F}_{(2)}$ for the time-dependent generator $\mathcal{F}_t$ of the Ising spin chain model.

6 Conclusions and outlook

In this work, we explored a novel method for approximating the dynamics of a large autonomous system projected onto a given subspace with a time-dependent generator. The problem is inspired by the derivation of TCL ME in quantum physics but has been treated here in the general case of linear autonomous systems.

The main contribution of this work consists of a novel method for computing the coefficients of the time-dependent model that is polynomial in time and that approximates the evolution of the projected dynamics for small times. Importantly, the coefficients of the polynomial can be determined using a recursive formula.

Benefits of the proposed reduction method include: the method does not require to move into a rotating frame, making the calculations simpler; the iterative form of the coefficients allows for easy calculations of the terms; for lower orders the method offers an advantage with respect to truncating the exponential, as shown in Sec. ??; the method does not rely on a perturbation on a small parameter ε allowing for analysis of systems in strong coupling regimes; when the original dynamics is purely Hamiltonian, the reduced time-dependent model at the lowest two orders is ensured to generate a CPTP

two-parameter semigroup at all times. All of this comes at the cost of having a model that approximates the original evolution in the short-time regime.

Future work shall focus on: i) extending Theorem ?? and Theorem ?? to the case of analytic generators; ii) lifting the assumption $Px_0 = x_0$; iii) generalize the results presented here to include control inputs (both linear and bilinear); iv) apply the proposed procedure to other quantum models in the strong coupling regime to study models that were not accessible before and further test the capabilities of the method; v) combine the classical model reduction methods and the time-dependent model reduction proposed here to see if the combination of the two methods can allow to improve the accuracy of the reduced model. Note that a first step in this direction has been taken in the physics literature by [?,?], where, including extra degrees of freedom, has been shown to improve the accuracy of the derived TCL ME.

Given the numerical examples of the central spin model presented in Sec. ?? we do not expect the reduced model to retain the CP property for orders higher than $N = 3$. However, it is possible that, in the Hamiltonian case, for $N = 3$ one can prove that the evolution of the reduced model is necessarily CP at all times, as none of the simulation carried out while preparing this work contradicted this statement.

7 Acknowledgments

The author would like to thank Francesco Ticozzi for his contribution in the initial idea of this work and for his constant support. The author also thanks Lorenza Viola, Michiel Burgelman, Augusto Ferrante, and Giacomo Baggio for useful discussions on the matters of this work.

Code availability: The code necessary to reproduce all the numerical experiments presented in this work is available at [?].

Funding: The author has been supported by the Italian Ministry of University and Research under the PRIN project “Extracting essential information and dynamics from complex networks”, grant no. 2022MBC2EZ.

References

- [1] Athanasios C. Antoulas. *Approximation of Large-Scale Dynamical Systems*. Advances in Design and Control, vol. 6;, Society for Industrial and Applied Mathematics, Philadelphia, 2005.
- [2] Mario Argeri, Stefano Di Vita, Pierpaolo Mastrolia, Edoardo Mirabella, Johannes Schlenk, Ulrich Schubert, and Lorenzo Tancredi. Magnus and dyson series for master integrals. *Journal of High Energy Physics*, 2014(3):1–32, 2014.
- [3] Uri M Ascher, Steven J Ruuth, and Raymond J Spiteri. Implicit-explicit runge-kutta methods for time-dependent partial differential equations. *Applied Numerical Mathematics*, 25(2-3):151–167, 1997.
- [4] Alessandro Astolfi. Model reduction by moment matching for linear and nonlinear systems. *IEEE Transactions on Automatic Control*, 55(10):2321–2336, 2010.
- [5] Rémi Azouit, Francesca Chittaro, Alain Sarlette, and Pierre Rouchon. Towards generic adiabatic elimination for bipartite open quantum systems. *Quantum Science and Technology*, 2(4):044011, 2017.
- [6] Edwin Barnes, Łukasz Cywiński, and S Das Sarma. Nonperturbative master equation solution of central spin dephasing dynamics. *Physical review letters*, 109(14):140403, 2012.
- [7] Fabio Benatti, Dariusz Chruściński, and Giovanni Nichele. Quantum versus classical p-divisibility. *Physical Review A*, 110(5):052212, 2024.
- [8] Michał Białończyk, Andrzej Jamiolkowski, and Karol Życzkowski. Application of shemesh theorem to quantum channels. *Journal of Mathematical Physics*, 59(10), 2018.
- [9] Bruce Blackadar. *Operator algebras: theory of C^* -algebras and von Neumann algebras*, volume 122. Springer Science & Business Media, 2006.
- [10] Robin Blume-Kohout, Hui Khoo Ng, David Poulin, and Lorenza Viola. Information-preserving structures: A general framework for quantum zero-error information. *Physical Review A*, 82(6):062306, 2010.
- [11] Andrei Bourchtein and Ludmila Bourchtein. Analytic functions and their properties. In *Complex Analysis*, pages 61–127. Springer, 2021.
- [12] Heinz-Peter Breuer. Non-markovian generalization of the lindblad theory of open quantum systems. *Phys. Rev. A*, 75:022103, Feb 2007.
- [13] Heinz-Peter Breuer, Jochen Gemmer, and Mathias Michel. Non-markovian quantum dynamics: Correlated projection superoperators and hilbert space averaging. *Phys. Rev. E*, 73:016139, Jan 2006.
- [14] Heinz-Peter Breuer, Francesco Petruccione, et al. *The theory of open quantum systems*. Oxford University Press on Demand, 2002.
- [15] Javier Cerrillo and Jianshu Cao. Non-markovian dynamical maps: Numerical processing of open quantum trajectories. *Phys. Rev. Lett.*, 112:110401, Mar 2014.
- [16] S. Chaturvedi and F. Shibata. Time-convolutionless projection operator formalism for elimination of fast variables. Applications to Brownian motion. *Zeitschrift für Physik B Condensed Matter*, 35(3), September 1979.
- [17] Alessandra Colla, Heinz-Peter Breuer, and Giulio Gasbarri. Recursive perturbation approach to time-convolutionless master equations: Explicit construction of generalized lindblad generators for arbitrary open systems. *arXiv preprint arXiv:2506.04095*, 2025.
- [18] Alessandra Colla, Heinz-Peter Breuer, and Giulio Gasbarri. Unveiling coherent dynamics in non-markovian open quantum systems: exact expression and recursive perturbation expansion. *arXiv preprint arXiv:2506.04097*, 2025.
- [19] Moritz Cygorek and Erik M. Gauger. Time-nonlocal versus time-local long-time extrapolation of non-markovian quantum dynamics, 2025.
- [20] F. J. Dyson. The radiation theories of tomonaga, schwinger, and feynman. *Phys. Rev.*, 75:486–502, Feb 1949.
- [21] E. Ferraro, H.-P. Breuer, A. Napoli, M. Guccione, and A. Messina. Non-markovian master equation for the xx central spin model. In *2008 2nd ICTON Mediterranean Winter*, pages 1–4, 2008.
- [22] G. Gasbarri and L. Ferialdi. Recursive approach for non-markovian time-convolutionless master equations. *Phys. Rev. A*, 97:022114, Feb 2018.
- [23] P.-L. Giscard, K. Lui, S. J. Thwaite, and D. Jaksch. An exact formulation of the time-ordered exponential using path-sums. *Journal of Mathematical Physics*, 56(5), May 2015.
- [24] Vittorio Gorini, Andrzej Kossakowski, and E. C. G. Sudarshan. Completely Positive Dynamical Semigroups of N Level Systems. *J. Math. Phys.*, 17:821, 1976.
- [25] Tommaso Grigoletto. Time ordered expansion – GitHub repository. https://github.com/tommygrigo/Time_ordered_expansion, 2025.
- [26] Tommaso Grigoletto, Clément Pellegrini, and Francesco Ticozzi. Quantum model reduction for continuous-time quantum filters, 2025.
- [27] Tommaso Grigoletto, Yukuan Tao, Francesco Ticozzi, and Lorenza Viola. Exact model reduction for continuous-time open quantum dynamics. *Quantum*, 2025.
- [28] Tommaso Grigoletto and Francesco Ticozzi. Model reduction for quantum systems: Discrete-time quantum walks and open markov dynamics. *IEEE Transactions on Information Theory*, 2025.
- [29] Serkan Gugercin and Athanasios C Antoulas. A survey of model reduction by balanced truncation and some new results. *International Journal of Control*, 77(8):748–766, 2004.
- [30] Jun Jing and Lian-Ao Wu. Decoherence and control of a qubit in spin baths: an exact master equation study. *Scientific reports*, 8(1):1471, 2018.
- [31] Peter D Johnson, Francesco Ticozzi, and Lorenza Viola. General fixed points of quasi-local frustration-free quantum semigroups: from invariance to stabilization. *arXiv preprint arXiv:1506.07756*, 2015.
- [32] P.J. Lawrence. A volterra series approach to the solution of time varying linear differential equations. *International Journal of Control*, 30(4):581–586, 1979.
- [33] Goran Lindblad. On the generators of quantum dynamical semigroups. *Communications in Mathematical Physics*, 48:119–130, 1976.
- [34] Göran Lindblad. A general no-cloning theorem. *Letters in Mathematical Physics*, 47(2):189–196, 1999.
- [35] Sadao Nakajima. On quantum theory of transport phenomena: Steady diffusion. *Progress of Theoretical Physics*, 20(6):948–959, 12 1958.

- [36] Konstantin Nestmann and Carsten Timm. Time-convolutionless master equation: Perturbative expansions to arbitrary order and application to quantum dots. *arXiv preprint arXiv:1903.05132*, 2019.
- [37] Konstantin Nestmann and Carsten Timm. Time-convolutionless master equation: Perturbative expansions to arbitrary order and application to quantum dots, March 2019. arXiv:1903.05132 [cond-mat].
- [38] Alberto Padoan. Model reduction by least squares moment matching for linear and nonlinear systems. *IEEE Transactions on Automatic Control*, 68(12):8267–8274, 2023.
- [39] Dénes Petz. *Quantum information theory and quantum statistics*. Springer Science & Business Media, 2007.
- [40] Howard Harry Rosenbrock. *State-space and multivariable theory*, volume 3. Wiley Interscience Division, 1970.
- [41] Jun John Sakurai and Jim Napolitano. *Modern quantum mechanics*. Cambridge University Press, 2020.
- [42] Fumiaki Shibata, Yoshinori Takahashi, and Natsuki Hashitsume. A generalized stochastic liouville equation. Non-Markovian versus memoryless master equations. *Journal of Statistical Physics*, 17(4), October 1977.
- [43] Francesco Ticozzi, Luca Zuccato, Peter D Johnson, and Lorenza Viola. Alternating projections methods for discrete-time stabilization of quantum states. *IEEE Transactions on Automatic Control*, 63(3):819–826, 2017.
- [44] Devvrat Tiwari, Baibhab Bose, and Subhashish Banerjee. Strong coupling non-markovian quantum thermodynamics of a finite-bath system. *The Journal of Chemical Physics*, 162(11), 2025.
- [45] Masaaki Tokieda and Angela Riva. Time-convolutionless master equation applied to adiabatic elimination. *Physical Review A*, 111(5), May 2025.
- [46] Michael M Wolf. Quantum channels & operations: Guided tour, 2012. Lecture notes available at <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=afa58291b0b8bd47504acb1ab8f553f0b37685cf>.
- [47] Michael M. Wolf and David Perez-Garcia. The inverse eigenvalue problem for quantum channels, 2010.
- [48] Robert Zwanzig. Ensemble method in the theory of irreversibility. *The Journal of Chemical Physics*, 33(5):1338–1341, 11 1960.

A Proof for the time-ordered expansion

Proof.[Theorem ??] We shall start by proving that $\mathbb{T}e^{\int_0^t F_s ds} = \sum_{n=0}^{+\infty} \Phi_{(n),t}$ where $\Phi_{(0),t} = \mathcal{I}$ and, for $n > 0$,

$$\Phi_{(n),t} = \sum_{k_1, \dots, k_n=1}^{N+1} t^{\sum_{j=1}^n k_j} \frac{F_{(k_1)} \cdots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h}.$$

First of all, using the Dyson series expansion for the time-ordered exponential $\mathbb{T}e^{\int_0^t F_s ds}$, see e.g. [?, ?, ?], we have that:

$$\mathbb{T}e^{\int_0^t F_s ds} = \sum_{n=0}^{+\infty} \Phi_{(n),t}$$

with

$$\Phi_{(n),t} = \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-1}} dt_n F_{t_1} F_{t_2} \cdots F_{t_n}.$$

Next, by noticing that $\Phi_{(n),t} = \int_0^t ds F_s \Phi_{(n-1),s}$ we can prove the statement by induction. The case $n = 1$ is proven by simple calculations:

$$\begin{aligned} \Phi_{(1),t} &= \int_0^t ds F_s = \sum_{k_1=0}^N F_{(k_1+1)} \int_0^t ds s^{k_1} \\ &= \sum_{k_1=0}^N F_{(k_1+1)} \frac{t^{k_1+1}}{k_1+1} = \sum_{k_1=1}^{N+1} F_{(k_1)} \frac{t^{k_1}}{k_1}. \end{aligned}$$

Note that $\int_0^t ds \sum_{k=0}^N = \sum_{k=0}^N \int_0^t ds$ under the assumption that N is finite, i.e. $N < \infty$. We next assume that the statement holds for n , i.e.

$$\Phi_{(n),t} = \sum_{k_1, \dots, k_n=1}^{N+1} t^{\sum_{j=1}^n k_j} \frac{F_{(k_1)} \cdots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h}.$$

Then

$$\begin{aligned} \Phi_{(n+1),t} &= \int_0^t ds F_s \Phi_{(n),s} = \\ &= \int_0^t ds F_s \sum_{k_1, \dots, k_n=1}^{N+1} s^{\sum_{j=1}^n k_j} \frac{F_{(k_n)} \cdots F_{(k_1)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h} \\ &= \sum_{k=0}^N \sum_{k_1, \dots, k_n=1}^{N+1} \frac{F_{(k+1)} F_{(k_1)} \cdots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h} \int_0^t s^{k + \sum_{j=1}^n k_j} ds \\ &= \sum_{k=0}^N \sum_{k_1, \dots, k_n=1}^{N+1} \frac{t^{k+1 + \sum_{j=1}^n k_j}}{k+1 + \sum_{j=1}^n k_j} \frac{F_{(k+1)} F_{(k_1)} \cdots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h} \end{aligned}$$

$$\begin{aligned} &= \sum_{k=1}^{N+1} \sum_{k_1, \dots, k_n=1}^{N+1} \frac{t^{k + \sum_{j=1}^n k_j}}{k + \sum_{j=1}^n k_j} \frac{F_{(k)} F_{(k_1)} \cdots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h} \\ &= \sum_{k_1=1}^{N+1} \sum_{k_2, \dots, k_{n+1}=1}^{N+1} \frac{t^{k_1 + \sum_{j=2}^{n+1} k_j}}{k_1 + \sum_{j=2}^{n+1} k_j} \frac{F_{(k_1)} F_{(k_2)} \cdots F_{(k_{n+1})}}{\prod_{j=1}^n \sum_{h=n-j+2}^{n+1} k_h} \\ &= \sum_{k_1, \dots, k_{n+1}=1}^{N+1} \frac{t^{\sum_{j=1}^{n+1} k_j}}{\sum_{j=1}^{n+1} k_j} \frac{F_{(k_1)} \cdots F_{(k_{n+1})}}{\prod_{j=1}^n \sum_{h=n-j+2}^{n+1} k_h} \\ &= \sum_{k_1, \dots, k_{n+1}=1}^{N+1} t^{\sum_{j=1}^{n+1} k_j} \frac{F_{(k_1)} \cdots F_{(k_{n+1})}}{\prod_{j=1}^{n+1} \sum_{h=n-j+2}^{n+1} k_h}. \end{aligned}$$

thus proving the statement.

We next prove that $\mathbb{T}e^{\int_0^t F_s ds} = \sum_{f=0}^{+\infty} t^k E_{(k)}$ where $E_{(0)} = \mathbb{I}$ and, for $k > 0$,

$$E_{(k)} = \sum_{n=1}^k \sum_{\substack{k_1, \dots, k_n=1 \\ \text{s.t. } \sum_{j=1}^n k_j = k}} \frac{F_{(k_1)} \dots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h}.$$

To prove this claim we can start by noticing that each term $\Phi_{(n),t}$ contains powers t^k of order $k \geq n$. Hence, to construct $E_{(k)}$ of order k we need to collect and sum all the coefficients that multiply t^k among the terms $\Phi_{(n),t}$ for $1 \leq n \leq k$. Given $\Phi_{(n),t}$ with $n \leq k$ we have that the coefficient multiplying t^k must satisfy the constraint $\sum_{j=1}^n k_j = k$. Thus, the coefficient of $\Phi_{(n),t}$ that multiply t^k are

$$\sum_{\substack{k_1, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k}} \frac{F_{(k_1)} \dots F_{(k_n)}}{\prod_{j=1}^n \sum_{h=n-j+1}^n k_h}$$

where we used the fact that, if any $k_j > k$ then $\sum_{j=1}^n k_j > k$. Noticing that $\prod_{j=1}^n \sum_{h=n-j+1}^n k_h = k(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)$ and summing over $1 \leq n \leq k$ we obtain, for $k > 0$, the second claim, i.e.

$$E_{(k)} = \sum_{n=1}^k \sum_{\substack{k_1, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k}} \frac{F_{(k_1)} \dots F_{(k_n)}}{k(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)}.$$

We are now finally ready to prove the main claim of the theorem, that is: $E_{(k)} = \frac{1}{k} \sum_{s=1}^k F_{(s)} E_{(k-s)}$. Recalling that $E_{(0)} = \mathbb{I}$ we can write:

$$\begin{aligned} & \frac{1}{k} \sum_{s=1}^k F_{(s)} E_{(k-s)} \\ &= \sum_{s=1}^{k-1} \frac{F_{(s)}}{k} E_{(k-s)} + \frac{F_{(k)}}{k} \\ &= \sum_{s=1}^{k-1} \sum_{n=1}^{k-1} \sum_{\substack{k_1, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k-s}} \frac{F_{(s)} F_{(k_1)} \dots F_{(k_n)}}{k(k-s)(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)} \\ & \quad + \frac{F_{(k)}}{k} \end{aligned}$$

where we used the fact that

$$\sum_{n=k-s+1}^{k-1} \sum_{\substack{k_1, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k-s}} \frac{F_{(s)} F_{(k_1)} \dots F_{(k_n)}}{k(k-s)(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)} = 0$$

since for $n > k-s$ we have that $\sum_{j=1}^n k_j > k-s$ for any set of k_j . This implies that

$$\begin{aligned} & \frac{1}{k} \sum_{s=1}^k F_{(s)} E_{(k-s)} \\ &= \sum_{n=1}^k \sum_{s=1}^{k-1} \sum_{\substack{k_1, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k-s}} \frac{F_{(s)} F_{(k_1)} \dots F_{(k_n)}}{k(k-s)(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)} \\ & \quad + \frac{F_{(k)}}{k} \\ &= \sum_{n=1}^k \sum_{k_1=1}^k \sum_{\substack{k_2, \dots, k_n=1 \\ \sum_{j=1}^n k_j = k}} \frac{F_{(k_1)} F_{(k_2)} \dots F_{(k_n)}}{k(\prod_{j=1}^{n-1} \sum_{h=n-j+1}^n k_h)} = E_{(k)}, \end{aligned}$$

thus concluding the proof. $\square$

Note that the technical reason why we choose to assume F_t to be polynomial ($N < \infty$) instead of analytic is because in this proof we used in a couple of instances the property that $\int_0^t \sum_{k=0}^N s^k F_{(k+1)} ds = \sum_{k=0}^N \int_0^t s^k F_{(k+1)} ds$ which trivially holds if $N < \infty$. To lift this assumption we would need to assume some extra properties about the operators $F_{(k+1)}$ so that the order between the summation and integral can be changed.