

# 1 Introduction

**Definition 1.1.** Let  $\mathbb{Z} \ni p \geq 0$ . A *generalized Hilbert matrix* is a matrix  $H \in M_n(\mathbb{R})$  whose entries are defined:  $H_{ij} = \frac{1}{i+j-1+p}$

Our problem is to find  $a \in \mathbb{R}^N$  such that

$$a^T H a \tag{1.1}$$

is minimized, where  $H$  is a generalized Hilbert matrix (in our case,  $p=2$ ), subject to the constraint:

$$B^T a = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \tag{1.2}$$

where

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix} \in M_{N \times 2}(\mathbb{R})$$

Our goal is to prove that such an  $a$  has coefficients that alternate in sign. Precisely:

**Theorem 1.2.** *Let  $H$  be a generalized Hilbert matrix of size  $N$  and  $B \in M_{N \times 2}$  is as above. Furthermore, let*

$$\hat{a} = \operatorname{argmin}_{a \in \mathbb{R}^N} \quad a^T H a : \quad B^T a = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

*Then for odd  $i$ ,  $\hat{a}(i) > 0$  and for even  $i$   $\hat{a}(i) < 0$ .*

We can obtain expressions for  $\hat{a}$  via the method of Lagrange multipliers. Our problem amounts to minimizing

$$\frac{1}{2} a^T H a - \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} (B^T a - \begin{pmatrix} 1 \\ -1 \end{pmatrix})$$

Taking the derivative with respect to  $a$ , we see that we must have

$$a = H^{-1} B \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} = H^{-1} \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_2 \end{pmatrix}$$

and the constraint of  $B^T a$  results in :

$$\begin{aligned} B^T H^{-1} B \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} &= \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ \implies \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} &= \frac{1}{-} \end{aligned}$$