

# A Saddle Point Paradigm for Finite Element Analysis and Its Role in the DPG Methodology

*Brendan Keith*

Supervisor: Leszek Demkowicz

Collaborators: Marek Behr

Stefanie Elgeti

Federico Fuentes

Jay Gopalakrishnan

Philipp Knechtges

Sriram Nagaraj

Socratis Petrides

Nathan V. Roberts

Committee: George Biros

Thomas J. R. Hughes

J. Tinsley Oden

Nathan V. Roberts

Funding: National Science Foundation

UT Austin Graduate School

Continuing Fellowship

Nathan V. Roberts

Patrick Le Tallec

Ali Vaziri

The Institute for Computational Engineering and Sciences  
The University of Texas at Austin  
August 10, 2018

# Outline

## Introduction

- Babuška to Brezzi
- DPG methods
- DPG\* methods

# Outline

## Introduction

- Babuška to Brezzi
- DPG methods
- DPG\* methods

## DPG\* Analysis

### A Priori Error Estimation

- Role of the Lagrange multiplier
- Duality
- Aubin–Nitsche

# Outline

## Introduction

- Babuška to Brezzi
- DPG methods
- DPG\* methods

## DPG\* Analysis

### A Priori Error Estimation

- Role of the Lagrange multiplier
- Duality
- Aubin–Nitsche

### *A Posteriori* Error Control

- DPG methods
- DPG\* methods
- Crude upper bounds & marking strategies

# Outline

## Introduction

- Babuška to Brezzi
- DPG methods
- DPG\* methods

## DPG\* Analysis

### A Priori Error Estimation

- Role of the Lagrange multiplier
- Duality
- Aubin–Nitsche

### A Posteriori Error Control

- DPG methods
- DPG\* methods
- Crude upper bounds & marking strategies

## Applications

- Poisson
- Viscoelastic fluids

# Prelude: Other Research

# Shape Function Software Library

- Energy Spaces

$$H^1 \xrightarrow{\nabla} H(\text{curl}) \xrightarrow{\nabla \times} H(\text{div}) \xrightarrow{\nabla \cdot} L^2$$

- Hybrid Meshes


1D:



2D:




Segment



Quadrilateral


3D:



Hexahedron



Tetrahedron



Prism



Pyramid

- High order, *anisotropic*, hierarchical shape functions
- Affine invariance to master element geometries
- Sparse stiffness matrices
- Orientation embeddings

---

F. Fuentes, B. Keith, L. Demkowicz, and S. Nagaraj.

Orientation embedded high order shape functions for the exact sequence elements of all shapes.

*Comput. Math. Appl.*, 70(4):353–458, 2015.

# Shape Function Software Library


$$\begin{array}{ccccccc} \text{3D: } & H^1 & \xrightarrow{\nabla} & H(\text{curl}) & \xrightarrow{\nabla \times} & H(\text{div}) & \xrightarrow{\nabla \cdot} L^2 \\ & \text{ext} \uparrow & & \text{ext} \uparrow & & \text{ext} \uparrow & \\ \text{2D: } & H^1 & \xrightarrow{\nabla} & H(\text{curl}) & \xrightarrow{\nabla \times} & L^2 & \\ & \text{ext} \uparrow & & \text{ext} \uparrow & & & \\ \text{1D: } & H^1 & \xrightarrow{\nabla} & L^2, & & & \end{array}$$

Classification into *subtopologies*

1D: vertex and edge shape functions,


2D: vertex, edge, and face shape functions,

3D: vertex, edge, face and interior shape functions.



# Structural Mechanics

## Sheathed hose




$$(E_{\text{steel}} = 200 \text{GPa}, E_{\text{Rubber}} = 0.01 \text{GPa})$$

## Coupled formulations

- All coupled formulations are **mutually compatible** throughout the domain.

Circumferential stress:  $\sigma_{\theta\theta}$



F. Fuentes, B. Keith, L. Demkowicz, and P. L. Tallec.

Coupled variational formulations of linear elasticity and the DPG methodology.  
*J. Comput. Phys.*, 348:715–731, 2017.

B. Keith, F. Fuentes, and L. Demkowicz.

The DPG methodology applied to different variational formulations of linear elasticity.  
*Comput. Methods Appl. Mech. Engrg.*, 309:579–609, 2016.

# Fluid Mechanics

## Oldroyd-B fluid

Conservation of mass and momentum:

$$\begin{aligned} \rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) - \nabla \cdot \boldsymbol{\sigma} &= \rho \mathbf{f} \quad \text{on } \Omega \times (0, T), \\ \nabla \cdot \mathbf{u} &= 0 \quad \text{on } \Omega \times (0, T). \end{aligned}$$

Constitutive law:

$$\boldsymbol{\sigma} = -p \mathbf{I} + 2\mu_S \boldsymbol{\varepsilon}(\mathbf{u}) + \mathbf{T},$$

where

$$\mathbf{T} + \lambda \mathcal{L}_u \mathbf{T} = 2\mu_P \boldsymbol{\varepsilon}(\mathbf{u}).$$

Lie derivative:

$$\mathcal{L}_u \mathbf{T} = \frac{\partial \mathbf{T}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{T} - (\nabla \mathbf{u} \mathbf{T} - \mathbf{T} \nabla^T \mathbf{u}).$$

# Fluid Mechanics

Oldroyd-B fluid

Conservation of mass and momentum:

$$\rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) - \nabla \cdot \boldsymbol{\sigma} = \rho \mathbf{f} \quad \text{on } \Omega \times (0, T).$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{on } \Omega \times (0, T).$$

Constitutive law:

$$\boldsymbol{\sigma} = -p \mathbf{I} + 2\mu_S \boldsymbol{\varepsilon}(\mathbf{u}) + \mathbf{T},$$

where

$$\mathbf{T} + \lambda \mathcal{L}_u \mathbf{T} = 2\mu_P \boldsymbol{\varepsilon}(\mathbf{u}).$$

Lie derivative:

$$\mathcal{L}_u \mathbf{T} = \frac{\partial \mathbf{T}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{T} - (\nabla \mathbf{u} \mathbf{T} - \mathbf{T} \nabla^T \mathbf{u}).$$

Quantity of interest

Drag coefficient


$$c_D(\boldsymbol{\sigma}) = \frac{1}{\mu \bar{u}} \int_{\Gamma_C} (\boldsymbol{\sigma} \vec{n}) \cdot \mathbf{e}_x \, ds.$$

$\Gamma_C$ : boundary of cylinder

$\vec{n}$ : traction

$\mu = \mu_S + \mu_P$ : viscosity

$\bar{u}$ : average inflow velocity.



Confined cylinder domain.


# Wave Mechanics

- Acoustics
- Electromagnetics
- Elastodynamics

$$-\Delta p - \omega^2 p = f$$

$$\frac{1}{\mu} \nabla \times \nabla \times \mathbf{E} - \omega^2 \epsilon \mathbf{E} = i\omega \mathbf{J}^{\text{imp}}$$

$$-\nabla \cdot (\mathbf{C} : \nabla \mathbf{u}) - \rho \omega^2 \mathbf{u} = \mathbf{f}$$



Electromagnetic wave scattering of the discrete electric field  $\mathbf{E}$ .

(a) The  $x$ -component; (b) the  $y$ -component; (c) the  $z$ -component. Only the real part of the solution is visualized.

---

A. Vaziri Astaneh, B. Keith, and L. Demkowicz.

On perfectly matched layers for discontinuous Petrov–Galerkin methods.  
*arXiv:1804.04496 (math.NA)*, 2018.

# Discrete Least Squares


Two solution paradigms

Normal equations

$$\underbrace{B^T G^{-1} B}_A u = B^T G^{-1} f$$

Overdetermined system

$$\underbrace{L^{-1} B}_\sim u = L^{-1} f$$



Acoustics with DPG near resonance

# Discrete Least Squares

## Two solution paradigms


Normal equations

$$\underbrace{B^T G^{-1} B}_A u = B^T G^{-1} f$$

Overdetermined system

$$\underbrace{L^{-1} B}_\sim u = L^{-1} f$$

- ★ Static condensation for  $L^{-1}B$  is explained in the article.
- ★  $L^{-1}B$  has  $O(h^{-1})$  condition number!
- ★ Less round off error when handling  $L^{-1}B$  directly.



Acoustics with DPG near resonance

# Discrete Least Squares

## Two solution paradigms


Normal equations

$$\underbrace{B^T G^{-1} B}_{A} u = B^T G^{-1} f$$

Overdetermined system


$$\underbrace{L^{-1} B}_{\tilde{B}} u = L^{-1} f$$

- ★ Static condensation for  $L^{-1}B$  is explained in the article.
- ★  $L^{-1}B$  has  $O(h^{-1})$  condition number!
- ★ Less round off error when handling  $L^{-1}B$  directly.




Acoustics with DPG near resonance



# Summary



B. Keith, S. Petrides, F. Fuentes, and L. Demkowicz.  
Discrete least-squares finite element methods.  
*Comput. Methods Appl. Mech. Engrg.*, 327:226–255, 2017.



# Summary



F. Fuentes, B. Keith, L. Demkowicz, and S. Nagaraj.

Orientation embedded high order shape functions for the exact sequence elements of all shapes.  
*Comput. Math. Appl.*, 70(4):353–458, 2015.

F. Fuentes, B. Keith, L. Demkowicz, and P. L. Tallec.

Coupled variational formulations of linear elasticity and the DPG methodology.  
*J. Comput. Phys.*, 348:715–731, 2017.

B. Keith, F. Fuentes, and L. Demkowicz.

The DPG methodology applied to different variational formulations of linear elasticity.  
*Comput. Methods Appl. Mech. Engrg.*, 309:579–609, 2016.

B. Keith, P. Knechtges, N. V. Roberts, S. Elgeti, M. Behr, and L. Demkowicz.

An ultraweak DPG method for viscoelastic fluids.  
*J. Non-Newton. Fluid Mech.*, 247:107–122, 2017.

B. Keith, S. Petrides, F. Fuentes, and L. Demkowicz.

Discrete least-squares finite element methods.  
*Comput. Methods Appl. Mech. Engrg.*, 327:226–255, 2017.

A. Vaziri Astaneh, B. Keith, and L. Demkowicz.

On perfectly matched layers for discontinuous Petrov–Galerkin methods.  
*arXiv:1804.04496 (math.NA)*, 2018.

# Outline

## Introduction

- Babuška to Brezzi
- DPG methods
- DPG\* methods

## DPG\* Analysis

### A Priori Error Estimation

- Role of the Lagrange multiplier
- Duality
- Aubin–Nitsche

### A Posteriori Error Control

- DPG methods
- DPG\* methods
- Crude upper bounds & marking strategies

## Applications

- Poisson
- Viscoelastic fluids

# Introduction

# Stability

Babuška to Brezzi

Let  $b : \mathcal{U} \times \mathcal{V} \rightarrow \mathbb{R}$  be a continuous bilinear form.

# Stability

Babuška to Brezzi

Let  $b : \mathcal{U} \times \mathcal{V} \rightarrow \mathbb{R}$  be a continuous bilinear form.

$\mathcal{U}$  and  $\mathcal{V}$  Hilbert

# Stability

Babuška to Brezzi

Let  $b : \mathcal{U} \times \mathcal{V} \rightarrow \mathbb{R}$  be a continuous bilinear form.

$\mathcal{U}$  and  $\mathcal{V}$  Hilbert

$\mathcal{U}_h \subset \mathcal{U}$  and  $\mathcal{V}_h \subset \mathcal{V}$  finite-dimensional

# Stability

Babuška to Brezzi

Let  $b : \mathcal{U} \times \mathcal{V} \rightarrow \mathbb{R}$  be a continuous bilinear form.

$\mathcal{U}$  and  $\mathcal{V}$  Hilbert

$\mathcal{U}_h \subset \mathcal{U}$  and  $\mathcal{V}_h \subset \mathcal{V}$  finite-dimensional

Problem:

Find  $\textcolor{brown}{u}_h \in \mathcal{U}_h$  satisfying  $b(\textcolor{brown}{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$

# Stability

Babuška to Brezzi

Define  $\mathcal{B}(u)(v) = b(u, v)$  for all  $u \in \mathcal{U}$  and  $v \in \mathcal{V}$

# Stability

Babuška to Brezzi

Define  $\mathcal{B}(u)(v) = b(u, v)$  for all  $u \in \mathcal{U}$  and  $v \in \mathcal{V}$

Assume  $F \in (\text{Null } \mathcal{B}')^\perp$

# Stability

Babuška to Brezzi

Define  $\mathcal{B}(u)(v) = b(u, v)$  for all  $u \in \mathcal{U}$  and  $v \in \mathcal{V}$

Assume  $F \in (\text{Null } \mathcal{B}')^\perp$

The following **continuous problem** has a unique solution:

Find  $u \in \mathcal{U}$  satisfying  $b(u, v) = F(v) \quad \forall v \in \mathcal{V},$

$\|u\|_{\mathcal{U}} \leq \gamma^{-1} \|F\|_{\mathcal{V}'},$  if and only if

$$\inf_{u \in \mathcal{U}} \sup_{v \in \mathcal{V}} \frac{b(u, v)}{\|u\|_{\mathcal{U}} \|v\|_{\mathcal{V}}} = \gamma > 0$$

# Stability

Babuška to Brezzi

Define  $\mathcal{B}(u)(v) = b(u, v)$  for all  $u \in \mathcal{U}$  and  $v \in \mathcal{V}$

Assume  $F \in (\text{Null } \mathcal{B}')^\perp$

The following **continuous problem** has a unique solution:

Find  $u \in \mathcal{U}$  satisfying  $b(u, v) = F(v) \quad \forall v \in \mathcal{V},$

$\|u\|_{\mathcal{U}} \leq \gamma^{-1} \|F\|_{\mathcal{V}'},$  if and only if

$$\inf_{u \in \mathcal{U}} \sup_{v \in \mathcal{V}} \frac{b(u, v)}{\|u\|_{\mathcal{U}} \|v\|_{\mathcal{V}}} = \gamma > 0, \quad \gamma = \|\mathcal{B}^{-1}\|^{-1}$$

# Stability

Babuška to Brezzi

## Continuous stability

$$\inf_{u \in \mathcal{U}} \sup_{v \in \mathcal{V}} \frac{b(u, v)}{\|u\|_u \|v\|_v} = \gamma > 0 \quad \not\Rightarrow$$

## Discrete stability

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_u \|v_h\|_v} = \gamma_h > 0$$

# Stability

Babuška to Brezzi

Continuous stability

$$\inf_{u \in \mathcal{U}} \sup_{v \in \mathcal{V}} \frac{b(u, v)}{\|u\|_U \|v\|_V} = \gamma > 0 \quad \not\Rightarrow$$

Discrete stability

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_U \|v_h\|_V} = \gamma_h > 0$$

Q: How to satisfy the discrete inf-sup condition

$$\sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|v_h\|_V} \geq \gamma_h \|u_h\|_U, \quad \forall u_h \in \mathcal{U}_h \quad ?$$

# Stability

Babuška to Brezzi

Continuous stability

$$\inf_{u \in \mathcal{U}} \sup_{v \in \mathcal{V}} \frac{b(u, v)}{\|u\|_u \|v\|_v} = \gamma > 0 \quad \nRightarrow \quad$$

Discrete stability

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_u \|v_h\|_v} = \gamma_h > 0$$

**Q:** How to satisfy the discrete inf-sup condition

$$\sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|v_h\|_v} \geq \gamma_h \|u_h\|_u, \quad \forall u_h \in \mathcal{U}_h \quad ?$$

**A:** Fix  $\mathcal{U}_h$  and increase the dimension of  $\mathcal{V}_h$  until satisfied.

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\textcolor{brown}{u}_h \in \mathcal{U}_h$  :

$$b(\textcolor{brown}{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$b(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

**Brezzi:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\boldsymbol{\varepsilon}_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\boldsymbol{\varepsilon}_h, v_h)_V + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \boldsymbol{\varepsilon}_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

- Requires *two* inf-sup conditions

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$b(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

**Brezzi:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\boldsymbol{\varepsilon}_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\boldsymbol{\varepsilon}_h, v_h)_V + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \boldsymbol{\varepsilon}_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

- Requires *two* inf-sup conditions
- An inf-sup condition relating spaces  $\mathcal{U}_h$  and  $\mathcal{V}_h$

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$b(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

**Brezzi:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\varepsilon_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\varepsilon_h, v_h)_\mathcal{V} + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \varepsilon_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

- Requires *two* inf-sup conditions
- An inf-sup condition relating spaces  $\mathcal{U}_h$  and  $\mathcal{V}_h$
- An inf-sup condition *in the kernel* of the dual operator

# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$b(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

**Brezzi:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\varepsilon_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\varepsilon_h, v_h)_\mathcal{V} + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \varepsilon_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

- Requires *two* inf-sup conditions
- An inf-sup condition relating spaces  $\mathcal{U}_h$  and  $\mathcal{V}_h$
- An inf-sup condition *in the kernel* of the dual operator



# Stability

Babuška to Brezzi

$$\inf_{u_h \in \mathcal{U}_h} \sup_{v_h \in \mathcal{V}_h} \frac{b(u_h, v_h)}{\|u_h\|_{\mathcal{U}} \|v_h\|_{\mathcal{V}}} = \gamma_h > 0 \quad \text{where } \dim \mathcal{V}_h > \dim \mathcal{U}_h$$

**Babuška:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$b(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in \mathcal{V}_h$$

**Brezzi:** Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\varepsilon_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\varepsilon_h, v_h)_\mathcal{V} + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \varepsilon_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

- Solvable only if  
 $\dim \mathcal{V}_h = \dim \mathcal{U}_h$

- Requires *two* inf-sup conditions
- An inf-sup condition relating spaces  $\mathcal{U}_h$  and  $\mathcal{V}_h$  
- An inf-sup condition *in the kernel* of the dual operator 

# Stability

Babuška to Brezzi

$$\dim \mathcal{V}_h > \dim \mathcal{U}_h$$

# Stability

Babuška to Brezzi

$$\dim \mathcal{V}_h > \dim \mathcal{U}_h$$

Dual problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  satisfying  $b(w_h, \mathbf{v}_h) = G(w_h) \quad \forall w_h \in \mathcal{U}_h$

# Stability

Babuška to Brezzi

$$\dim \mathcal{V}_h > \dim \mathcal{U}_h$$

Dual problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  satisfying  $b(w_h, \mathbf{v}_h) = G(w_h) \quad \forall w_h \in \mathcal{U}_h$

**Q:** The dual problem is now *underdetermined*. What to do?

# Stability

Babuška to Brezzi

$$\dim \mathcal{V}_h > \dim \mathcal{U}_h$$

Dual problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  satisfying  $b(w_h, \mathbf{v}_h) = G(w_h) \quad \forall w_h \in \mathcal{U}_h$

**Q:** The dual problem is now *underdetermined*. What to do?

**A:** Consider a different embedding in the saddle point system.

# Stability

Babuška to Brezzi

## Primal:

Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\boldsymbol{\varepsilon}_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\boldsymbol{\varepsilon}_h, v_h)_V + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \boldsymbol{\varepsilon}_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

## Dual:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\boldsymbol{\lambda}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, z_h)_V + b(\boldsymbol{\lambda}_h, z_h) &= 0 \quad \forall z_h \in \mathcal{V}_h \\ b(w_h, \mathbf{v}_h) &= G(w_h) \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

# Stability

Babuška to Brezzi

## Primal:

Find  $\mathbf{u}_h \in \mathcal{U}_h$  and  $\boldsymbol{\varepsilon}_h \in \mathcal{V}_h$  :

$$\begin{aligned} (\boldsymbol{\varepsilon}_h, v_h)_V + b(\mathbf{u}_h, v_h) &= F(v_h) \quad \forall v_h \in \mathcal{V}_h \\ b(w_h, \boldsymbol{\varepsilon}_h) &= 0 \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

## Dual:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\boldsymbol{\lambda}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, z_h)_V - b(\boldsymbol{\lambda}_h, z_h) &= 0 \quad \forall z_h \in \mathcal{V}_h \\ b(w_h, \mathbf{v}_h) &= G(w_h) \quad \forall w_h \in \mathcal{U}_h \end{aligned}$$

# Implementation

General problem:

Find  $v_h \in \mathcal{V}_h$  and  $w_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, \nu)_{\mathcal{V}} + b(\mathbf{w}_h, \nu) &= F(\nu) \quad \forall \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu) \quad \forall \mu \in \mathcal{U}_h \end{aligned}$$

# Implementation

General problem:

Find  $v_h \in \mathcal{V}_h$  and  $w_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, \nu)_h + b(\mathbf{w}_h, \nu) &= F(\nu) \quad \forall \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu) \quad \forall \mu \in \mathcal{U}_h \end{aligned}$$

Discretization:

$$\begin{bmatrix} \mathbf{G} & \mathbf{B} \\ \mathbf{B}^T & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \mathbf{f} \\ \mathbf{g} \end{bmatrix}$$

Schur complement:

$$\begin{aligned} \mathbf{B}^T \mathbf{G}^{-1} \mathbf{B} \mathbf{w} &= \mathbf{B}^T \mathbf{G}^{-1} \mathbf{f} - \mathbf{g} \\ \mathbf{v} &= \mathbf{G}^{-1} (\mathbf{f} - \mathbf{B} \mathbf{w}) \end{aligned}$$

# Implementation

General problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\mathbf{w}_h \in \mathcal{U}_h$  :

$$\begin{aligned}(\mathbf{v}_h, \nu)_{\mathcal{V}} + b(\mathbf{w}_h, \nu) &= F(\nu) \quad \forall \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu) \quad \forall \mu \in \mathcal{U}_h\end{aligned}$$

Discretization:

$$\begin{bmatrix} \mathbf{G} & \mathbf{B} \\ \mathbf{B}^T & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \mathbf{f} \\ \mathbf{g} \end{bmatrix}$$

Schur complement:

$$\begin{aligned}\mathbf{B}^T \mathbf{G}^{-1} \mathbf{B} \mathbf{w} &= \mathbf{B}^T \mathbf{G}^{-1} \mathbf{f} - \mathbf{g} \\ \mathbf{v} &= \mathbf{G}^{-1} (\mathbf{f} - \mathbf{B} \mathbf{w})\end{aligned}$$

DPG:  $\mathbf{G}$  is block-diagonal

# DPG and DPG\* comparison

Minimum norm derivation

Stiffness matrix: Gram matrix: Load vector: Goal vector:

$$B_{ij} = b(u_j, v_i)$$

$$G_{ij} = (v_j, v_i)_V$$

$$f_i = F(v_i)$$

$$g_j = G(u_j)$$

# DPG and DPG\* comparison

Minimum norm derivation

Stiffness matrix: Gram matrix: Load vector: Goal vector:

$$B_{ij} = b(u_j, v_i) \quad G_{ij} = (v_j, v_i)_V \quad f_i = F(v_i) \quad g_j = G(u_j)$$

In DPG, we solve:

$$B^T G^{-1} B u = B^T G^{-1} f$$

# DPG and DPG\* comparison

Minimum norm derivation

Stiffness matrix: Gram matrix: Load vector: Goal vector:

$$B_{ij} = b(u_j, v_i) \quad G_{ij} = (v_j, v_i)_V \quad f_i = F(v_i) \quad g_j = G(u_j)$$

In DPG, we solve:

$$B^T G^{-1} Bu = B^T G^{-1} f$$

In DPG\*, we solve:

$$B^T G^{-1} Bw = g$$

and then post-process  $w$ :

$$v = G^{-1} Bw$$

# Discrete least squares finite element methods

Cholesky factorization

$$G = LL^T.$$

# Discrete least squares finite element methods

Cholesky factorization

$$\mathbf{G} = \mathbf{L}\mathbf{L}^T.$$

Residual minimization problem

$$\mathbf{u} = \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T \mathbf{G}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f})$$

# Discrete least squares finite element methods

Cholesky factorization

$$\mathbf{G} = \mathbf{L}\mathbf{L}^T.$$

Residual minimization problem

$$\begin{aligned}\mathbf{u} &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T \mathbf{G}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T (\mathbf{L}\mathbf{L}^T)^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f})\end{aligned}$$

# Discrete least squares finite element methods

Cholesky factorization

$$\mathbf{G} = \mathbf{L}\mathbf{L}^T.$$

Residual minimization problem

$$\begin{aligned}\mathbf{u} &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T \mathbf{G}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T (\mathbf{L}\mathbf{L}^T)^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} \left( \mathbf{L}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \right)^T \mathbf{L}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f})\end{aligned}$$

# Discrete least squares finite element methods

Cholesky factorization

$$\mathbf{G} = \mathbf{L}\mathbf{L}^T.$$

Residual minimization problem

$$\begin{aligned}\mathbf{u} &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T \mathbf{G}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} (\mathbf{B}\mathbf{w} - \mathbf{f})^T (\mathbf{L}\mathbf{L}^T)^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} \left( \mathbf{L}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \right)^T \mathbf{L}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^N} \left\| \mathbf{L}^{-1} (\mathbf{B}\mathbf{w} - \mathbf{f}) \right\|_2^2.\end{aligned}$$

# Discrete Least Squares

Two solution paradigms


Normal equations

$$\mathbf{B}^T \mathbf{G}^{-1} \mathbf{B} \mathbf{u} = \mathbf{B}^T \mathbf{G}^{-1} \mathbf{f}$$

Overdetermined system

$$\mathbf{L}^{-1} \mathbf{B} \mathbf{u} = \mathbf{L}^{-1} \mathbf{f}$$

- ★ Static condensation for  $\mathbf{L}^{-1} \mathbf{B}$  is explained in the article
- ★  $\mathbf{L}^{-1} \mathbf{B}$  has  $O(h^{-1})$  condition number
- ★ Less round off error when handling  $\mathbf{L}^{-1} \mathbf{B}$  directly



Acoustics with DPG near resonance

# Highlights

- ★ DPG/DPG\* always deliver SPD (HPD) stiffness matrices

## Highlights

- ★ DPG/DPG\* always deliver SPD (HPD) stiffness matrices
- ★ DPG/DPG\* have built-in stability

## Highlights

- ★ DPG/DPG\* always deliver SPD (HPD) stiffness matrices
- ★ DPG/DPG\* have built-in stability
- ★ DPG has a built-in error estimator

## Highlights

- ★ DPG/DPG\* always deliver SPD (HPD) stiffness matrices
- ★ DPG/DPG\* have built-in stability
- ★ DPG has a built-in error estimator
- ★ DPG\* works for some problems without uniqueness

# *A priori* error estimation

Continuous problem:

Find  $v \in \mathcal{V}$  and  $w \in \mathcal{U}$  :

$$(v, \nu)_{\mathcal{V}} + b(w, \nu) = F(\nu), \quad \nu \in \mathcal{V}$$

$$b(\mu, v) = G(\mu), \quad \mu \in \mathcal{U}$$

# *A priori* error estimation

Continuous problem:

Find  $\mathbf{v} \in \mathcal{V}$  and  $\mathbf{w} \in \mathcal{U}$  :

$$\begin{aligned} (\mathbf{v}, \nu)_{\mathcal{V}} + b(\mathbf{w}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \mathbf{v}) &= G(\mu), \quad \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\mathbf{w}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, \nu)_{\mathcal{V}_h} + b(\mathbf{w}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu), \quad \mu \in \mathcal{U}_h \end{aligned}$$

# *A priori* error estimation

Continuous problem:

Find  $\mathbf{v} \in \mathcal{V}$  and  $\mathbf{w} \in \mathcal{U}$  :

$$\begin{aligned} (\mathbf{v}, \nu)_{\mathcal{V}} + b(\mathbf{w}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \mathbf{v}) &= G(\mu), \quad \mu \in \mathcal{U} \end{aligned}$$

Fortin operator:

Continuous operator  $\Pi_h : \mathcal{V} \rightarrow \mathcal{V}_h$  such that

$$b(\mu, \nu - \Pi_h \nu) = 0 \quad \forall \mu \in \mathcal{U}_h, \nu \in \mathcal{V}.$$

Discrete problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\mathbf{w}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, \nu)_{\mathcal{V}_h} + b(\mathbf{w}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu), \quad \mu \in \mathcal{U}_h \end{aligned}$$

# *A priori* error estimation

## Continuous problem:

Find  $\mathbf{v} \in \mathcal{V}$  and  $\mathbf{w} \in \mathcal{U}$  :

$$\begin{aligned} (\mathbf{v}, \nu)_{\mathcal{V}} + b(\mathbf{w}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \mathbf{v}) &= G(\mu), \quad \mu \in \mathcal{U} \end{aligned}$$

## Discrete problem:

Find  $\mathbf{v}_h \in \mathcal{V}_h$  and  $\mathbf{w}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\mathbf{v}_h, \nu)_{\mathcal{V}} + b(\mathbf{w}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \mathbf{v}_h) &= G(\mu), \quad \mu \in \mathcal{U}_h \end{aligned}$$

## Fortin operator:

Continuous operator  $\Pi_h : \mathcal{V} \rightarrow \mathcal{V}_h$  such that

$$b(\mu, \nu - \Pi_h \nu) = 0 \quad \forall \mu \in \mathcal{U}_h, \nu \in \mathcal{V}.$$

## Theorem (Brezzi)

Let  $F \in \mathcal{V}'$  and  $G \in \mathcal{U}'$ . There is a constant  $C$  such that

$$\|\mathbf{v} - \mathbf{v}_h\|_{\mathcal{V}} + \|\mathbf{w} - \mathbf{w}_h\|_{\mathcal{U}} \leq C \left[ \inf_{\nu \in \mathcal{V}_h} \|\mathbf{v} - \nu\|_{\mathcal{V}} + \inf_{\mu \in \mathcal{U}_h} \|\mathbf{w} - \mu\|_{\mathcal{U}} \right].$$

# *A priori* error estimation: DPG

Continuous problem:

Find  $\varepsilon \in \mathcal{V}$  and  $\mathbf{u} \in \mathcal{U}$  :

$$\begin{aligned} (\varepsilon, \nu)_\mathcal{V} + b(\mathbf{u}, \nu) &= F(\nu), & \nu \in \mathcal{V} \\ b(\mu, \varepsilon) &= 0, & \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\varepsilon_h \in \mathcal{V}_h$  and  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\varepsilon_h, \nu)_\mathcal{V} + b(\mathbf{u}_h, \nu) &= F(\nu), & \nu \in \mathcal{V}_h \\ b(\mu, \varepsilon_h) &= 0, & \mu \in \mathcal{U}_h \end{aligned}$$

# *A priori* error estimation: DPG

Continuous problem:

Find  $\varepsilon \in \mathcal{V}$  and  $\mathbf{u} \in \mathcal{U}$  :

$$\begin{aligned} (\varepsilon, \nu)_\mathcal{V} + b(\mathbf{u}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \varepsilon) &= 0, \quad \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\varepsilon_h \in \mathcal{V}_h$  and  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\varepsilon_h, \nu)_\mathcal{V} + b(\mathbf{u}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \varepsilon_h) &= 0, \quad \mu \in \mathcal{U}_h \end{aligned}$$

## Theorem

Let  $F \in (\text{Null } \mathcal{B}')^\perp$ . There is a constant  $C$  such that

$$\|\varepsilon - \varepsilon_h\|_\mathcal{V} + \|\mathbf{u} - \mathbf{u}_h\|_\mathcal{U} \leq C \left[ \inf_{\nu \in \mathcal{V}_h} \|\varepsilon - \nu\|_\mathcal{V} + \inf_{\mu \in \mathcal{U}_h} \|\mathbf{u} - \mu\|_\mathcal{U} \right].$$

# *A priori* error estimation: DPG

Continuous problem:

Find  $\varepsilon \in \mathcal{V}$  and  $\mathbf{u} \in \mathcal{U}$  :

$$\begin{aligned} (\varepsilon, \nu)_\mathcal{V} + b(\mathbf{u}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \varepsilon) &= 0, \quad \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\varepsilon_h \in \mathcal{V}_h$  and  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\varepsilon_h, \nu)_\mathcal{V} + b(\mathbf{u}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \varepsilon_h) &= 0, \quad \mu \in \mathcal{U}_h \end{aligned}$$

Recall:

$$b(\mathbf{u}, \nu) = F(\nu) \quad \forall \nu \in \mathcal{V} \quad \Rightarrow \quad \varepsilon = 0.$$

## Theorem

Let  $F \in (\text{Null } \mathcal{B}')^\perp$ . There is a constant  $C$  such that

$$\|\varepsilon - \varepsilon_h\|_\mathcal{V} + \|\mathbf{u} - \mathbf{u}_h\|_\mathcal{U} \leq C \left[ \inf_{\nu \in \mathcal{V}_h} \|\varepsilon - \nu\|_\mathcal{V} + \inf_{\mu \in \mathcal{U}_h} \|\mathbf{u} - \mu\|_\mathcal{U} \right].$$

# *A priori* error estimation: DPG

Continuous problem:

Find  $\varepsilon \in \mathcal{V}$  and  $\mathbf{u} \in \mathcal{U}$  :

$$\begin{aligned} (\varepsilon, \nu)_\mathcal{V} + b(\mathbf{u}, \nu) &= F(\nu), \quad \nu \in \mathcal{V} \\ b(\mu, \varepsilon) &= 0, \quad \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\varepsilon_h \in \mathcal{V}_h$  and  $\mathbf{u}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\varepsilon_h, \nu)_\mathcal{V} + b(\mathbf{u}_h, \nu) &= F(\nu), \quad \nu \in \mathcal{V}_h \\ b(\mu, \varepsilon_h) &= 0, \quad \mu \in \mathcal{U}_h \end{aligned}$$

Recall:

$$b(\mathbf{u}, \nu) = F(\nu) \quad \forall \nu \in \mathcal{V} \quad \implies \quad \varepsilon = 0.$$

## Theorem

Let  $F \in (\text{Null } \mathcal{B}')^\perp$ . There is a constant  $C$  such that

$$\|\varepsilon_h\|_\mathcal{V} + \|\mathbf{u} - \mathbf{u}_h\|_\mathcal{U} \leq C \inf_{\mu \in \mathcal{U}_h} \|\mathbf{u} - \mu\|_\mathcal{U}.$$

## *A priori* error estimation: DPG\*

Continuous problem:

Find  $\boldsymbol{v} \in \mathcal{V}$  and  $\lambda \in \mathcal{U}$  :

$$\begin{aligned} (\boldsymbol{v}, \nu)_{\mathcal{V}} - b(\lambda, \nu) &= 0, & \nu \in \mathcal{V} \\ b(\mu, \boldsymbol{v}) &= G(\mu), & \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\boldsymbol{v}_h \in \mathcal{V}_h$  and  $\lambda_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\boldsymbol{v}_h, \nu)_{\mathcal{V}} - b(\lambda_h, \nu) &= 0, & \nu \in \mathcal{V}_h \\ b(\mu, \boldsymbol{v}_h) &= G(\mu), & \mu \in \mathcal{U}_h \end{aligned}$$

# *A priori* error estimation: DPG\*

## Continuous problem:

Find  $\boldsymbol{v} \in \mathcal{V}$  and  $\boldsymbol{\lambda} \in \mathcal{U}$  :

$$\begin{aligned} (\boldsymbol{v}, \nu)_\mathcal{V} - b(\boldsymbol{\lambda}, \nu) &= 0, & \nu \in \mathcal{V} \\ b(\mu, \boldsymbol{v}) &= G(\mu), & \mu \in \mathcal{U} \end{aligned}$$

## Discrete problem:

Find  $\boldsymbol{v}_h \in \mathcal{V}_h$  and  $\boldsymbol{\lambda}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\boldsymbol{v}_h, \nu)_\mathcal{V} - b(\boldsymbol{\lambda}_h, \nu) &= 0, & \nu \in \mathcal{V}_h \\ b(\mu, \boldsymbol{v}_h) &= G(\mu), & \mu \in \mathcal{U}_h \end{aligned}$$

## Theorem

Let  $G \in \mathcal{U}'$ . There is a constant  $C$  such that

$$\|\boldsymbol{v} - \boldsymbol{v}_h\|_\mathcal{V} + \|\boldsymbol{\lambda} - \boldsymbol{\lambda}_h\|_\mathcal{U} \leq C \left[ \inf_{\nu \in \mathcal{V}_h} \|\boldsymbol{v} - \nu\|_\mathcal{V} + \inf_{\mu \in \mathcal{U}_h} \|\boldsymbol{\lambda} - \mu\|_\mathcal{U} \right].$$

## *A priori* error estimation: DPG\*

Continuous problem:

Find  $\boldsymbol{v} \in \mathcal{V}$  and  $\boldsymbol{\lambda} \in \mathcal{U}$  :

$$\begin{aligned} (\boldsymbol{v}, \nu)_\mathcal{V} - b(\boldsymbol{\lambda}, \nu) &= 0, & \nu \in \mathcal{V} \\ b(\mu, \boldsymbol{v}) &= G(\mu), & \mu \in \mathcal{U} \end{aligned}$$

Discrete problem:

Find  $\boldsymbol{v}_h \in \mathcal{V}_h$  and  $\boldsymbol{\lambda}_h \in \mathcal{U}_h$  :

$$\begin{aligned} (\boldsymbol{v}_h, \nu)_\mathcal{V} - b(\boldsymbol{\lambda}_h, \nu) &= 0, & \nu \in \mathcal{V}_h \\ b(\mu, \boldsymbol{v}_h) &= G(\mu), & \mu \in \mathcal{U}_h \end{aligned}$$

### Theorem

Let  $G \in \mathcal{U}'$ . There is a constant  $C$  such that

$$\|\boldsymbol{v} - \boldsymbol{v}_h\|_\mathcal{V} + \|\boldsymbol{\lambda} - \boldsymbol{\lambda}_h\|_\mathcal{U} \leq C \left[ \inf_{\nu \in \mathcal{V}_h} \|\boldsymbol{v} - \nu\|_\mathcal{V} + \inf_{\mu \in \mathcal{U}_h} \|\boldsymbol{\lambda} - \mu\|_\mathcal{U} \right].$$

DPG\*: The discretization error is affected by  $\boldsymbol{\lambda}$ !

# *A priori* error estimation: DPG\*

Poisson's equation (ultraweak)

Consider

Broken ultraweak formulation

$$\begin{aligned} -\Delta v = f \quad \implies \quad & b((u, \vec{q}, \hat{u}, \hat{q}_n), (v, \vec{p})) \\ &= (\vec{q}, \vec{p} - \text{grad}_h v)_\Omega - (u, \text{div}_h \vec{p})_\Omega \\ &+ \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_h \end{aligned}$$

# *A priori* error estimation: DPG\*

Poisson's equation (ultraweak)

Consider

Broken ultraweak formulation

$$\begin{aligned} -\Delta v = f \quad \implies \quad & b((u, \vec{q}, \hat{u}, \hat{q}_n), (v, \vec{p})) \\ &= ((u, \vec{q}), \mathcal{L}_h(v, \vec{p}))_{\Omega} \\ &\quad + \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_h \end{aligned}$$

where  $\mathcal{L}_h(v, \vec{p}) = (-\operatorname{div}_h \vec{p}, \vec{p} - \operatorname{grad}_h v)$ .

# *A priori* error estimation: DPG\*

Poisson's equation (ultraweak)

Consider

Broken ultraweak formulation

$$\begin{aligned} -\Delta v = f \quad \implies \quad & b((u, \vec{q}, \hat{u}, \hat{q}_n), (v, \vec{p})) \\ &= ((u, \vec{q}), \mathcal{L}_h(v, \vec{p}))_{\Omega} \\ &\quad + \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_h \end{aligned}$$

where  $\mathcal{L}_h(v, \vec{p}) = (-\operatorname{div}_h \vec{p}, \vec{p} - \operatorname{grad}_h v)$ .

DPG\* formulation:

$$\begin{aligned} ((\boldsymbol{v}, \vec{\boldsymbol{p}}), (\boldsymbol{\nu}, \vec{\boldsymbol{\tau}}))_{\boldsymbol{\nu}} - b((\lambda, \vec{\zeta}, \hat{\lambda}, \hat{\zeta}), (\boldsymbol{\nu}, \vec{\boldsymbol{\tau}})) &= 0 \quad \forall \boldsymbol{\nu}, \vec{\boldsymbol{\tau}} \\ b((\mu, \vec{\sigma}, \hat{\mu}, \hat{\sigma}_n), (\boldsymbol{v}, \vec{\boldsymbol{p}})) &= (f, \mu)_{\Omega} \quad \forall \mu, \vec{\sigma}, \hat{\mu}, \hat{\sigma}_n \end{aligned}$$

# *A priori* error estimation: DPG\*

Poisson's equation (ultraweak)

## Proposition

Let  $\|(\mathbf{v}, \vec{p})\|_{\mathcal{V}}^2 = \|\operatorname{grad}_h \mathbf{v}\|_{\Omega}^2 + \|\mathbf{v}\|_{\Omega}^2 + \|\operatorname{div}_h \vec{p}\|_{\Omega}^2 + \|\vec{p}\|_{\Omega}^2$ . The solution components  $\lambda, \vec{\zeta}, \hat{\lambda}, \hat{\zeta}$  of the system above can be characterized using the remaining solution components  $\mathbf{v}, \vec{p}$  and  $f$  as

$$\begin{aligned}\lambda &= f + \mathbf{e}, & \hat{\lambda} &= \mathbf{e} \\ \vec{\zeta} &= \vec{p} + \vec{r}, & \hat{\zeta} &= 2\vec{p} \cdot \vec{n} + \vec{r} \cdot \vec{n}\end{aligned}$$

where  $\mathbf{e} \in H_0^1(\Omega)$  satisfies the Dirichlet problem  $-\Delta \mathbf{e} = \mathbf{v} + 2f$  and  $\vec{r} = -\operatorname{grad} \mathbf{e}$ .

---

L. Demkowicz, J. Gopalakrishnan, and B. Keith.

The DPG-star method.

*In preparation*, 2018.

T. Führer.

Superconvergence in a DPG method for an ultra-weak formulation.

*Comput. Math. Appl.*, 75(5):1705 – 1718, 2018.

## Sketch of proof

- Use the second line to characterize the solution  $\textcolor{brown}{v}, \vec{p}$ .

Note that  $b((\mu, \vec{\sigma}, \hat{\mu}, \hat{\sigma}_n), (\textcolor{brown}{v}, \vec{p})) = (f, \mu)_\Omega$ .

$$\begin{aligned} \implies ((\mu, \vec{\sigma}), \mathcal{L}_h(\textcolor{brown}{v}, \vec{p}))_\Omega &= (f, \mu)_\Omega & \forall \mu, \vec{\sigma} \\ \langle \hat{\mu}, \vec{p} \cdot \vec{n} \rangle_h &= 0 & \forall \hat{\mu} \\ \langle \hat{\sigma}_n, \textcolor{brown}{v} \rangle_h &= 0 & \forall \hat{\sigma}_n \end{aligned}$$

where  $\mathcal{L}_h(v, \vec{p}) = (-\operatorname{div}_h \vec{p}, \vec{p} - \operatorname{grad}_h v)$ .

# Sketch of proof

- Use the second line to characterize the solution  $\mathbf{v}, \vec{p}$ .

Note that  $b((\mu, \vec{\sigma}, \hat{\mu}, \hat{\sigma}_n), (\mathbf{v}, \vec{p})) = (f, \mu)_\Omega$ .

$$\begin{aligned} \implies ((\mu, \vec{\sigma}), \mathcal{L}_h(\mathbf{v}, \vec{p}))_\Omega &= (f, \mu)_\Omega & \forall \mu, \vec{\sigma} \\ \langle \hat{\mu}, \vec{p} \cdot \vec{n} \rangle_h &= 0 & \forall \hat{\mu} \\ \langle \hat{\sigma}_n, \mathbf{v} \rangle_h &= 0 & \forall \hat{\sigma}_n \end{aligned}$$

where  $\mathcal{L}_h(\mathbf{v}, \vec{p}) = (-\operatorname{div}_h \vec{p}, \vec{p} - \operatorname{grad}_h \mathbf{v})$ .

- Show that this implies

$$\vec{p} - \operatorname{grad} \mathbf{v} = \vec{0}, \quad -\operatorname{div} \vec{p} = f.$$

## Sketch of proof

- Manipulate first term in first line:

$$\begin{aligned} ((\mathbf{v}, \vec{\mathbf{p}}))_{\mathcal{V}} &= (\vec{\mathbf{p}}, \vec{\tau})_{\Omega} + (\operatorname{div} \vec{\mathbf{p}}, \operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} + (\operatorname{grad} \mathbf{v}, \operatorname{grad} \nu)_{\Omega} \\ &= (\vec{\mathbf{p}}, \vec{\tau} - \operatorname{grad} \nu)_{\Omega} + (\operatorname{div} \vec{\mathbf{p}}, \operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} + 2(\operatorname{grad} \mathbf{v}, \operatorname{grad} \nu)_{\Omega} \\ &= (\vec{\mathbf{p}}, \vec{\tau} - \operatorname{grad} \nu)_{\Omega} + (f, -\operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} \\ &\quad + 2 \sum_{K \in \Omega} \left[ \langle \vec{n} \cdot \operatorname{grad} \mathbf{v}, \nu \rangle_{H^{1/2}(\partial K)} - (\Delta \mathbf{v}, \nu)_K \right] \\ &= b((f, \vec{\mathbf{p}}, 0, 2\vec{\mathbf{p}} \cdot \vec{n}), (\nu, \vec{\tau})) + (\mathbf{v} + 2f, \nu)_{\Omega} \end{aligned}$$

## Sketch of proof

- Manipulate first term in first line:

$$\begin{aligned} ((\mathbf{v}, \vec{p}))_{\mathcal{V}} &= (\vec{p}, \vec{\tau})_{\Omega} + (\operatorname{div} \vec{p}, \operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} + (\operatorname{grad} \mathbf{v}, \operatorname{grad} \nu)_{\Omega} \\ &= (\vec{p}, \vec{\tau} - \operatorname{grad} \nu)_{\Omega} + (\operatorname{div} \vec{p}, \operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} + 2(\operatorname{grad} \mathbf{v}, \operatorname{grad} \nu)_{\Omega} \\ &= (\vec{p}, \vec{\tau} - \operatorname{grad} \nu)_{\Omega} + (f, -\operatorname{div} \vec{\tau})_{\Omega} + (\mathbf{v}, \nu)_{\Omega} \\ &\quad + 2 \sum_{K \in \Omega} \left[ \langle \vec{n} \cdot \operatorname{grad} \mathbf{v}, \nu \rangle_{H^{1/2}(\partial K)} - (\Delta \mathbf{v}, \nu)_K \right] \\ &= b((f, \vec{p}, 0, 2\vec{p} \cdot \vec{n}), (\nu, \vec{\tau})) + (\mathbf{v} + 2f, \nu)_{\Omega} \end{aligned}$$

- Define  $\mathbf{e}$ ,  $\vec{r}$ ,  $\hat{\mathbf{e}}$ , and  $\hat{\mathbf{r}}$  such that

$$b((\mathbf{e}, \vec{r}, \hat{\mathbf{e}}, \hat{\mathbf{r}}), (\nu, \vec{\tau})) = (\mathbf{v} + 2f, \nu)_{\Omega} \implies -\Delta \mathbf{e} = \mathbf{v} + 2f$$

# *A priori* error estimation: DPG\*

Poisson's equation

Bramble-Hilbert argument:

## Corollary

Let  $\vec{v}_h = (\mathbf{v}_h, \vec{p}_h) \in \mathcal{V}_h$  and  $\vec{\lambda}_h = (\lambda_h, \vec{\zeta}_h, \hat{\lambda}_h, \hat{\zeta}_h) \in \mathcal{U}_h$  be the DPG\* solutions.

Let  $\mathbf{e} \in H_0^1(\Omega)$  satisfy  $-\Delta \mathbf{e} = \mathbf{v} + 2f$ . Then

$$\|\vec{v} - \vec{v}_h\|_{\mathcal{V}} + \|\vec{\lambda} - \vec{\lambda}_h\|_{\mathcal{U}} \leq Ch^s (\|\mathbf{v}\|_{H^{s+2}(\Omega)} + \|\mathbf{e}\|_{H^{s+2}(\Omega)})$$

for all  $1/2 < s < p + 1$ .

# *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 1:  $v = \sin(\pi x) \sin(\pi y)$ ,  $\Omega = (0, 1)^2$




Figure: Uniform  $h$ -refinements

# *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 2:  $v = 1$




Figure: Uniform  $h$ -refinements

# *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 2:  $v = 1$

$$-\Delta e = 1 \text{ in } \Omega, \quad e|_{\partial\Omega} = 0$$




Figure: Adaptive  $h$ -refinements

## *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 2:  $v = 1$

$$-\Delta e = 1 \text{ in } \Omega, \quad e|_{\partial\Omega} = 0$$

# *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 2:  $v = 1$

$$-\Delta\lambda = 1 \text{ in } \Omega, \quad \lambda|_{\partial\Omega} = 0$$





Figure: Left: The solution component  $\lambda$  when  $v = 1$ . Right: The corresponding mesh (ten refinements).

# Duality

DPG:

$$(1) \quad (\varepsilon, z)_V + b(\textcolor{brown}{u}, z) = F(z)$$

$$(2) \quad b(w, \varepsilon) = 0$$

DPG\*:

$$(3) \quad (\textcolor{brown}{v}, z)_V - b(\lambda, z) = 0$$

$$(4) \quad b(w, \textcolor{brown}{v}) = G(w)$$

DPG:

$$(1) \quad (\varepsilon, z)_V + b(\mathbf{u}, z) = F(z)$$

$$(2) \quad b(w, \varepsilon) = 0$$

DPG\*:

$$(3) \quad (\mathbf{v}, z)_V - b(\lambda, z) = 0$$

$$(4) \quad b(w, \mathbf{v}) = G(w)$$

Observe:

$$G(\mathbf{u} - \mathbf{u}_h) = b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) \quad \text{by (4)}$$

DPG:

$$(1) \quad (\varepsilon, z)_V + b(\mathbf{u}, z) = F(z)$$

$$(2) \quad b(w, \varepsilon) = 0$$

DPG\*:

$$(3) \quad (\mathbf{v}, z)_V - b(\lambda, z) = 0$$

$$(4) \quad b(w, \mathbf{v}) = G(w)$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \end{aligned}$$

**DPG:**

$$(5) \quad (\varepsilon_h, z_h)_V + b(\mathbf{u}_h, z_h) = F(z_h)$$

$$(6) \quad b(w_h, \varepsilon_h) = 0$$

**DPG\*:**

$$(7) \quad (\mathbf{v}_h, z_h)_V - b(\lambda_h, z_h) = 0$$

$$(8) \quad b(w_h, \mathbf{v}_h) = G(w_h)$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \end{aligned}$$

**DPG:**

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_\gamma + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

**DPG\*:**

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_\gamma - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_\gamma - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \end{aligned}$$

DPG:

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_\gamma + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

DPG\*:

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_\gamma - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_\gamma - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_\gamma - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \end{aligned}$$

DPG:

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_\gamma + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

DPG\*:

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_\gamma - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_\gamma - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_\gamma - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_\gamma + b(\lambda - w_h, \varepsilon_h) && \varepsilon = 0 \end{aligned}$$

DPG:

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_V + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

DPG\*:

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_V - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_V - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_V + b(\lambda - w_h, \varepsilon_h) && \varepsilon = 0 \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) - (\mathbf{v} - \mathbf{v}_h, \varepsilon_h)_V + b(\lambda - \lambda_h, \varepsilon_h) \end{aligned}$$

DPG:

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_V + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

DPG\*:

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_V - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_V - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_V + b(\lambda - w_h, \varepsilon_h) && \varepsilon = 0 \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) - (\mathbf{v} - \mathbf{v}_h, \varepsilon_h)_V + b(\lambda - \lambda_h, \varepsilon_h) \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) && \text{by (11)} \end{aligned}$$

DPG:

$$(9) \quad (\varepsilon - \varepsilon_h, z_h)_V + b(\mathbf{u} - \mathbf{u}_h, z_h) = 0$$

$$(10) \quad b(w_h, \varepsilon - \varepsilon_h) = 0$$

DPG\*:

$$(11) \quad (\mathbf{v} - \mathbf{v}_h, z_h)_V - b(\lambda - \lambda_h, z_h) = 0$$

$$(12) \quad b(w_h, \mathbf{v} - \mathbf{v}_h) = 0$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_V - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_V + b(\lambda - w_h, \varepsilon_h) && \varepsilon = 0 \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) - (\mathbf{v} - \mathbf{v}_h, \varepsilon_h)_V + b(\lambda - \lambda_h, \varepsilon_h) \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) && \text{by (11)} \\ &= b(\mathbf{u} - w_h, \mathbf{v} - \mathbf{v}_h) && \text{by (12)} \end{aligned}$$

**DPG:**

$$(1) \quad (\varepsilon, z)_V + b(\mathbf{u}, z) = F(z)$$

$$(2) \quad b(w, \varepsilon) = 0$$

**DPG\*:**

$$(3) \quad (\mathbf{v}, z)_V - b(\lambda, z) = 0$$

$$(4) \quad b(w, \mathbf{v}) = G(w)$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_V - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_V + b(\lambda - w_h, \varepsilon_h) && \varepsilon = 0 \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) - (\mathbf{v} - \mathbf{v}_h, \varepsilon_h)_V + b(\lambda - \lambda_h, \varepsilon_h) \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) && \text{by (11)} \\ &= b(\mathbf{u} - w_h, \mathbf{v} - \mathbf{v}_h) && \text{by (12)} \\ &= F(\mathbf{v} - \mathbf{v}_h) && \text{by (1)} \end{aligned}$$

**DPG:**

$$(1) \quad (\varepsilon, z)_V + b(\mathbf{u}, z) = F(z)$$

$$(2) \quad b(w, \varepsilon) = 0$$

**DPG\*:**

$$(3) \quad (\mathbf{v}, z)_V - b(\lambda, z) = 0$$

$$(4) \quad b(w, \mathbf{v}) = G(w)$$

Observe:

$$\begin{aligned} G(\mathbf{u} - \mathbf{u}_h) &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) && \text{by (4)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + (\mathbf{v}, \varepsilon - \varepsilon_h)_V - b(\lambda, \varepsilon - \varepsilon_h) && \text{by (3)} \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) + (\mathbf{v} - z_h, \varepsilon - \varepsilon_h)_V - b(\lambda - w_h, \varepsilon - \varepsilon_h) && \text{by (9-10)} \\ &= \boxed{b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - z_h) - (\mathbf{v} - z_h, \varepsilon_h)_V + b(\lambda - w_h, \varepsilon_h)} && \varepsilon = 0 \\ &= b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) - (\mathbf{v} - \mathbf{v}_h, \varepsilon_h)_V + b(\lambda - \lambda_h, \varepsilon_h) \\ &= \boxed{b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h)} && \text{by (11)} \\ &= \boxed{b(\mathbf{u} - w_h, \mathbf{v} - \mathbf{v}_h)} && \text{by (12)} \\ &= F(\mathbf{v} - \mathbf{v}_h) && \text{by (1)} \end{aligned}$$

# Aubin-Nitsche

## Corollary

The following crude upper bounds hold:

$$b(\mathbf{u} - \mathbf{u}_h, \mathbf{v} - \mathbf{v}_h) \lesssim \begin{cases} \|\mathbf{u} - \mathbf{u}_h\|_{\mathcal{U}} \inf_{\mathbf{z}_h \in \mathcal{V}_h, \mathbf{w}_h \in \mathcal{U}_h} (\|\mathbf{v} - \mathbf{z}_h\|_{\mathcal{V}} + \|\boldsymbol{\lambda} - \mathbf{w}_h\|_{\mathcal{U}}) \\ \|\mathbf{v} - \mathbf{v}_h\|_{\mathcal{V}} \inf_{\mathbf{w}_h \in \mathcal{U}_h} \|\mathbf{u} - \mathbf{w}_h\|_{\mathcal{U}} \end{cases}$$

# Aubin-Nitsche

## Theorem

Suppose that there is a positive function  $c_1(h)$  that goes to 0 as  $h \rightarrow 0$  satisfying

$$\inf_{w_h \in \mathcal{U}_h} \|\mathbf{u} - w_h\|_{\mathcal{U}} \leq c_1(h) \|F\|_{\mathcal{V}'}$$

Then the error in the DPG\* solution component  $\mathbf{u}_h$  satisfies

$$G(\mathbf{u} - \mathbf{u}_h) \leq c_0(h) \|G\|_{\mathcal{U}'} \|\mathcal{B}\| \inf_{w_h \in \mathcal{U}_h} \|\mathbf{u} - w_h\|_{\mathcal{U}}$$

- Can be used to demonstrate accelerated convergence of DPG\* solution components)

# *A priori* error estimation: DPG\*

Poisson's equation

Manufactured solution 2:  $v = 1$

$$-\Delta\lambda = 1 \text{ in } \Omega, \quad \lambda|_{\partial\Omega} = 0$$




Figure: Adaptive  $h$ -refinements.

# *A posteriori* error estimation

# *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|w\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

## *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|w\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

The energy norm  $\|\cdot\|_{\mathcal{U}}$  is equivalent to the norm  $\|\cdot\|_{\mathcal{U}}$

$$\|\mathcal{B}^{-1}\|^{-1}\|w\|_{\mathcal{U}} \leq \|w\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|w\|_{\mathcal{U}}$$

## *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|\|w\|\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

The energy norm  $\|\| \cdot \| \|_{\mathcal{U}}$  is equivalent to the norm  $\| \cdot \|_{\mathcal{U}}$

$$\|\mathcal{B}^{-1}\|^{-1} \|w\|_{\mathcal{U}} \leq \|\|w\|\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|w\|_{\mathcal{U}}$$

A posteriori error estimator for **broken** test spaces

$$\|\| \mathbf{u} - w \| \|_{\mathcal{U}}^2 = \|\mathcal{B}w - F\|_{\mathcal{V}'}^2$$

## *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|w\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

The energy norm  $\|\cdot\|_{\mathcal{U}}$  is equivalent to the norm  $\|\cdot\|_{\mathcal{U}}$

$$\|\mathcal{B}^{-1}\|^{-1}\|w\|_{\mathcal{U}} \leq \|w\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|w\|_{\mathcal{U}}$$

A posteriori error estimator for **broken test spaces**

$$\|\mathbf{u} - w\|_{\mathcal{U}}^2 = \|\mathcal{B}w - F\|_{\mathcal{V}'}^2 = \sum_{K \in \mathcal{T}} \|\mathcal{B}w - F\|_{\mathcal{V}(K)'}^2$$

# *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|w\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

The energy norm  $\|\cdot\|_{\mathcal{U}}$  is equivalent to the norm  $\|\cdot\|_{\mathcal{U}}$

$$\|\mathcal{B}^{-1}\|^{-1}\|w\|_{\mathcal{U}} \leq \|w\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|w\|_{\mathcal{U}}$$

A posteriori error estimator for **broken test spaces**

$$\|\mathbf{u} - w\|_{\mathcal{U}}^2 = \|\mathcal{B}w - F\|_{\mathcal{V}'}^2 = \sum_{K \in \mathcal{T}} \|\mathcal{B}w - F\|_{\mathcal{V}(K)'}^2 \simeq \sum_{K \in \mathcal{T}} \left( \underbrace{\|\mathcal{B}w - F\|_{\mathcal{V}_{\textcolor{brown}{h}(K)'}}} _{:= \eta_K(w)} \right)^2$$

# *A posteriori* error estimation: DPG

DPG minimizes the error in the **energy norm**

$$\|w\|_{\mathcal{U}} := \|\mathcal{B}w\|_{\mathcal{V}'} \quad \forall w \in \mathcal{U}$$

The energy norm  $\|\cdot\|_{\mathcal{U}}$  is equivalent to the norm  $\|\cdot\|_{\mathcal{U}}$

$$\|\mathcal{B}^{-1}\|^{-1}\|w\|_{\mathcal{U}} \leq \|w\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|w\|_{\mathcal{U}}$$

A posteriori error estimator for **broken test spaces**

$$\|\mathbf{u} - w\|_{\mathcal{U}}^2 = \|\mathcal{B}w - F\|_{\mathcal{V}'}^2 = \sum_{K \in \mathcal{T}} \|\mathcal{B}w - F\|_{\mathcal{V}(K)'}^2 \simeq \sum_{K \in \mathcal{T}} \left( \underbrace{\|\mathcal{B}w - F\|_{\mathcal{V}_{\textcolor{brown}{h}}(K)'}}_{:= \eta_K(w)} \right)^2$$

$$\eta(\mathbf{u}_h)^2 = \sum_{K \in \mathcal{T}} \eta_K(\mathbf{u}_h)^2$$

where

$$\eta_K(\mathbf{u}_h) = \sup_{z_h \in \mathcal{V}_h(K)} \frac{b(\mathbf{u}_h, z_h) - F(z_h)}{\|z_h\|_{\mathcal{V}}}$$

C. Carstensen, L. Demkowicz, and J. Gopalakrishnan.

A posteriori error control for DPG methods.

SIAM J. Numer. Anal., 52(3):1335–1353, 2014.

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{W}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{U}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')_{\perp}$ .

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{U}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')_{\perp}$ .
- $z_0 = \mathcal{P} z \implies \|z\|_{\mathcal{V}}^2 = \|\mathcal{P} z\|_{\mathcal{V}}^2 + \|z_{\perp}\|_{\mathcal{V}}^2$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{U}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}$ .
- $z_0 = \mathcal{P} z \implies \|z\|_{\mathcal{V}}^2 = \|\mathcal{P} z\|_{\mathcal{V}}^2 + \|z_{\perp}\|_{\mathcal{V}}^2$
- By the Closed Range Theorem, we find  $(\text{Null } \mathcal{B}')^{\perp} = \text{Range } \mathcal{B}$ :

$$\|z_{\perp}\|_{\mathcal{V}} = \sup_{\tilde{z}_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{(z, \tilde{z}_{\perp})_{\mathcal{V}}}{\|\tilde{z}_{\perp}\|_{\mathcal{V}}}$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}'z\|_{\mathcal{U}'}^2 + \|\mathcal{P}z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}$ .
- $z_0 = \mathcal{P}z \implies \|z\|_{\mathcal{V}}^2 = \|\mathcal{P}z\|_{\mathcal{V}}^2 + \|z_{\perp}\|_{\mathcal{V}}^2$
- By the Closed Range Theorem, we find  $(\text{Null } \mathcal{B}')^{\perp} = \text{Range } \mathcal{B}$ :

$$\|z_{\perp}\|_{\mathcal{V}} = \sup_{\tilde{z}_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{(z, \tilde{z}_{\perp})_{\mathcal{V}}}{\|\tilde{z}_{\perp}\|_{\mathcal{V}}} = \sup_{E^{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{\langle E^{\perp}, z \rangle_{\mathcal{V}}}{\|E^{\perp}\|_{\mathcal{V}'}}$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}'z\|_{\mathcal{U}'}^2 + \|\mathcal{P}z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}$ .
- $z_0 = \mathcal{P}z \implies \|z\|_{\mathcal{V}}^2 = \|\mathcal{P}z\|_{\mathcal{V}}^2 + \|z_{\perp}\|_{\mathcal{V}}^2$
- By the Closed Range Theorem, we find  $(\text{Null } \mathcal{B}')^{\perp} = \text{Range } \mathcal{B}$ :

$$\|z_{\perp}\|_{\mathcal{V}} = \sup_{\tilde{z}_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{(z, \tilde{z}_{\perp})_{\mathcal{V}}}{\|\tilde{z}_{\perp}\|_{\mathcal{V}}} = \sup_{E^{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{\langle E^{\perp}, z \rangle_{\mathcal{V}}}{\|E^{\perp}\|_{\mathcal{V}'}} = \sup_{\mu \in \mathcal{U}} \frac{\langle \mathcal{B}\mu, z \rangle_{\mathcal{U}}}{\|\mathcal{B}\mu\|_{\mathcal{V}'}}$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}'z\|_{\mathcal{U}'}^2 + \|\mathcal{P}z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V}$$

**Proof:**

- Let  $z = z_0 + z_{\perp}$ , where  $z_0 \in \text{Null } \mathcal{B}'$  and  $z_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}$ .
- $z_0 = \mathcal{P}z \implies \|z\|_{\mathcal{V}}^2 = \|\mathcal{P}z\|_{\mathcal{V}}^2 + \|z_{\perp}\|_{\mathcal{V}}^2$
- By the Closed Range Theorem, we find  $(\text{Null } \mathcal{B}')^{\perp} = \text{Range } \mathcal{B}$ :

$$\|z_{\perp}\|_{\mathcal{V}} = \sup_{\tilde{z}_{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{(z, \tilde{z}_{\perp})_{\mathcal{V}}}{\|\tilde{z}_{\perp}\|_{\mathcal{V}}} = \sup_{E^{\perp} \in (\text{Null } \mathcal{B}')^{\perp}} \frac{\langle E^{\perp}, z \rangle_{\mathcal{V}}}{\|E^{\perp}\|_{\mathcal{V}'}} = \sup_{\mu \in \mathcal{U}} \frac{\langle \mathcal{B}\mu, z \rangle_{\mathcal{U}}}{\|\mathcal{B}\mu\|_{\mathcal{V}'}} = \|\mathcal{B}'z\|_{\mathcal{U}'}.$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{W}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V} \quad \checkmark$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{U}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V} \quad \checkmark$$

Therefore, if  $\mathcal{B}$  is a bijection:

$$\|\boldsymbol{v} - \boldsymbol{v}_h\|_{\mathcal{V}}^2 = \|\mathcal{B}' \boldsymbol{v}_h - G\|_{\mathcal{U}'}^2$$

## *A posteriori* error estimation: DPG\*

Let  $\mathcal{P} : \mathcal{V} \rightarrow \text{Null } \mathcal{B}'$  be the orthogonal projection.

**Claim.** Identity for the error in the dual problem:

$$\|z\|_{\mathcal{V}}^2 = \|\mathcal{B}' z\|_{\mathcal{U}'}^2 + \|\mathcal{P} z\|_{\mathcal{V}}^2 \quad \forall z \in \mathcal{V} \quad \checkmark$$

Therefore, if  $\mathcal{B}$  is a bijection:

$$\|\mathbf{v} - \mathbf{v}_h\|_{\mathcal{V}}^2 = \|\mathcal{B}' \mathbf{v}_h - G\|_{\mathcal{U}'}^2$$

Moreover,  $\|\mathcal{B}^{-1}\|^{-1} \|\cdot\|_{\mathcal{U}} \leq \|\cdot\|_{\mathcal{U}} \leq \|\mathcal{B}\| \|\cdot\|_{\mathcal{U}}$ , so

$$\|\mathcal{B}\|^{-1} \|\mathcal{B}' \mathbf{v}_h - G\|_{\mathcal{U}'}^2 \leq \|\mathbf{v} - \mathbf{v}_h\|_{\mathcal{V}}^2 \leq \|\mathcal{B}^{-1}\| \|\mathcal{B}' \mathbf{v}_h - G\|_{\mathcal{U}'}^2$$

# Residual bounds

## Theorem

Let  $w \in \mathcal{U}$  and  $z \in \mathcal{V}$  be arbitrary. Then the following upper bound holds:

$$|b(\textcolor{brown}{u} - w, \textcolor{brown}{v} - z)| \leq \|\mathcal{B}^{-1}\| \|\mathcal{B}w - F\|_{\mathcal{V}'} \|\mathcal{B}'z - G\|_{\mathcal{U}'}.$$

# Residual bounds

## Theorem

Let  $w \in \mathcal{U}$  and  $z \in \mathcal{V}$  be arbitrary. Then the following upper bound holds:

$$|b(\mathbf{u} - w, \mathbf{v} - z)| \leq \|\mathcal{B}^{-1}\| \|\mathcal{B}w - F\|_{\mathcal{V}'} \|\mathcal{B}'z - G\|_{\mathcal{U}'}.$$

## Proof:

- Let  $z_0 \in \text{Null } \mathcal{B}'$  be arbitrary. Then

$$\begin{aligned}|b(\mathbf{u} - w, \mathbf{v} - z)| &= |b(\mathbf{u} - w, \mathbf{v} - z - z_0)| \\&\leq \|\mathcal{B}w - F\|_{\mathcal{V}'} \|v - z - z_0\| \\&= \|\mathcal{B}w - F\|_{\mathcal{V}'} (\|\mathcal{P}(v - z - z_0)\|_{\mathcal{V}}^2 + \|\mathcal{B}'z - G\|_{\mathcal{U}'}^2)^{1/2}.\end{aligned}$$

- Set  $z_0 = \mathcal{P}(v - z) \in \text{Null } \mathcal{B}'$ .

# Algorithm and Marking Strategies

. . . Solve–Estimate–Estimate–Mark–Refine. . .

## Adaptive mesh refinement

1. Solve for  $u_h$  and/or  $v_h$

- Mark all elements  $K \in \mathcal{T}$  such that  $\theta \cdot \tilde{\eta}_{\max} \leq \tilde{\eta}_K$ ,  $\theta \in (0, 1)$
- Choices for  $\tilde{\eta}$ :  $\tilde{\eta}_K = \eta_K$ ,  $\tilde{\eta}_K = \eta_K^*$ ,  $\tilde{\eta}_K = \eta_K \cdot \eta_K^*$ , . . .

# Algorithm and Marking Strategies

. . . Solve–Estimate–Estimate–Mark–Refine. . .

## Adaptive mesh refinement

1. Solve for  $\mathbf{u}_h$  and/or  $\mathbf{v}_h$
2. Estimate the global error and cease computations if sufficiently small

- Mark all elements  $K \in \mathcal{T}$  such that  $\theta \cdot \tilde{\eta}_{\max} \leq \tilde{\eta}_K$ ,  $\theta \in (0, 1)$
- Choices for  $\tilde{\eta}$ :  $\tilde{\eta}_K = \eta_K$ ,  $\tilde{\eta}_K = \eta_K^*$ ,  $\tilde{\eta}_K = \eta_K \cdot \eta_K^*$ , . . .

# Algorithm and Marking Strategies

. . . Solve–Estimate–Estimate–Mark–Refine. . .

## Adaptive mesh refinement

1. Solve for  $\mathbf{u}_h$  and/or  $\mathbf{v}_h$
2. Estimate the global error and cease computations if sufficiently small
3. Estimate local errors, i.e., compute refinement indicators

- Mark all elements  $K \in \mathcal{T}$  such that  $\theta \cdot \tilde{\eta}_{\max} \leq \tilde{\eta}_K$ ,  $\theta \in (0, 1)$
- Choices for  $\tilde{\eta}$ :  $\tilde{\eta}_K = \eta_K$ ,  $\tilde{\eta}_K = \eta_K^*$ ,  $\tilde{\eta}_K = \eta_K \cdot \eta_K^*$ , . . .

# Algorithm and Marking Strategies

. . . Solve–Estimate–Estimate–Mark–Refine. . .

## Adaptive mesh refinement

1. Solve for  $\mathbf{u}_h$  and/or  $\mathbf{v}_h$
2. Estimate the global error and cease computations if sufficiently small
3. Estimate local errors, i.e., compute refinement indicators
4. Mark elements for refinement as dictated by a marking strategy

e.g., 
$$G(\mathbf{u} - \mathbf{u}_h) \approx \|\mathcal{B}\mathbf{u}_h - F\|_{\mathcal{V}'} \|\mathcal{B}'\mathbf{v}_h - G\|_{\mathcal{W}'} \lesssim \eta(\mathbf{u}_h) \eta^*(\mathbf{v}_h)$$

- Mark all elements  $K \in \mathcal{T}$  such that  $\theta \cdot \tilde{\eta}_{\max} \leq \tilde{\eta}_K$ ,  $\theta \in (0, 1)$
- Choices for  $\tilde{\eta}$ :  $\tilde{\eta}_K = \eta_K$ ,  $\tilde{\eta}_K = \eta_K^*$ ,  $\tilde{\eta}_K = \eta_K \cdot \eta_K^*$ , . . .

# Algorithm and Marking Strategies

. . . Solve–Estimate–Estimate–Mark–Refine. . .

## Adaptive mesh refinement

1. **Solve** for  $\mathbf{u}_h$  and/or  $\mathbf{v}_h$
2. Estimate the global error and cease computations if sufficiently small
3. Estimate local errors, i.e., compute refinement indicators
4. **Mark** elements for refinement as dictated by a marking strategy



$$\text{e.g., } G(\mathbf{u} - \mathbf{u}_h) \approx \|\mathcal{B}\mathbf{u}_h - F\|_{\mathcal{V}'} \|\mathcal{B}'\mathbf{v}_h - G\|_{\mathcal{W}} \lesssim \eta(\mathbf{u}_h) \eta^*(\mathbf{v}_h)$$

5. **Refine** all marked elements and construct a new mesh

- Mark all elements  $K \in \mathcal{T}$  such that  $\theta \cdot \tilde{\eta}_{\max} \leq \tilde{\eta}_K$ ,  $\theta \in (0, 1)$
- Choices for  $\tilde{\eta}$ :  $\tilde{\eta}_K = \eta_K$ ,  $\tilde{\eta}_K = \eta_K^*$ ,  $\tilde{\eta}_K = \eta_K \cdot \eta_K^*$ , . . .


# Applications

# *hp*-adaptive mesh refinement in DPG\* methods




Singular solution and *hp*-adaptive mesh refinements with a DPG\* method for Poisson's equation.


# Goal-Oriented Adaptive Mesh Refinement (AMR) in 3D



The error in the average flux.



Solution-oriented AMR.



Goal-oriented AMR

# Fluid Mechanics

## Oldroyd-B fluid

Conservation of mass and momentum:

$$\begin{aligned} \rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) - \nabla \cdot \boldsymbol{\sigma} &= \rho \mathbf{f} \quad \text{on } \Omega \times (0, T), \\ \nabla \cdot \mathbf{u} &= 0 \quad \text{on } \Omega \times (0, T). \end{aligned}$$

Constitutive law:

$$\boldsymbol{\sigma} = -p \mathbf{I} + 2\mu_S \boldsymbol{\varepsilon}(\mathbf{u}) + \mathbf{T},$$

where

$$\mathbf{T} + \lambda \mathcal{L}_u \mathbf{T} = 2\mu_P \boldsymbol{\varepsilon}(\mathbf{u}).$$

Lie derivative:

$$\mathcal{L}_u \mathbf{T} = \frac{\partial \mathbf{T}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{T} - (\nabla \mathbf{u} \mathbf{T} - \mathbf{T} \nabla^T \mathbf{u}).$$

# Fluid Mechanics

Oldroyd-B fluid

Conservation of mass and momentum:

$$\rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) - \nabla \cdot \boldsymbol{\sigma} = \rho \mathbf{f} \quad \text{on } \Omega \times (0, T).$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{on } \Omega \times (0, T).$$

Constitutive law:

$$\boldsymbol{\sigma} = -p \mathbf{I} + 2\mu_S \boldsymbol{\varepsilon}(\mathbf{u}) + \mathbf{T},$$

where

$$\mathbf{T} + \lambda \mathcal{L}_u \mathbf{T} = 2\mu_P \boldsymbol{\varepsilon}(\mathbf{u}).$$

Lie derivative:

$$\mathcal{L}_u \mathbf{T} = \frac{\partial \mathbf{T}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{T} - (\nabla \mathbf{u} \mathbf{T} - \mathbf{T} \nabla^T \mathbf{u}).$$

Quantity of interest


Drag coefficient

$$c_D(\boldsymbol{\sigma}) = \frac{1}{\mu \bar{u}} \int_{\Gamma_c} (\boldsymbol{\sigma} \vec{n}) \cdot \mathbf{e}_x \, ds.$$

$\Gamma_c$ : boundary of cylinder

$\mu = \mu_S + \mu_P$ : viscosity

$\bar{u}$ : average inflow velocity.



Confined cylinder domain.

# Fluid Mechanics

Oldroyd-B fluid

Conservation of mass and momentum:

$$\rho \left( \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) - \nabla \cdot \boldsymbol{\sigma} = \rho \mathbf{f} \quad \text{on } \Omega \times (0, T).$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{on } \Omega \times (0, T).$$

Constitutive law:

$$\boldsymbol{\sigma} = -p \mathbf{I} + 2\mu_S \boldsymbol{\varepsilon}(\mathbf{u}) + \mathbf{T},$$

where

$$\mathbf{T} + \lambda \mathcal{L}_u \mathbf{T} = 2\mu_P \boldsymbol{\varepsilon}(\mathbf{u}).$$

Lie derivative:

$$\mathcal{L}_u \mathbf{T} = \frac{\partial \mathbf{T}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{T} - (\nabla \mathbf{u} \mathbf{T} - \mathbf{T} \nabla^T \mathbf{u}).$$

Quantity of interest


Drag coefficient

$$c_D(\boldsymbol{\sigma}) = \frac{1}{\mu \bar{u}} \int_{\Gamma_C} (\boldsymbol{\sigma} \vec{n}) \cdot \mathbf{e}_x \, ds.$$

$\Gamma_C$ : boundary of cylinder

$\mu = \mu_S + \mu_P$ : viscosity


$\bar{u}$ : average inflow velocity.



Close-up of rescaled  $T_{11}$ -component from  $\lambda = 0.9$ .

# Fluid Mechanics

## Solution-oriented adaptive mesh refinement



Close-up of rescaled  $T_{11}$ -component from  $\lambda = 0.7$ .


---

B. Keith, P. Knechtges, N. V. Roberts, S. Elgeti, M. Behr, and L. Demkowicz.

An ultraweak DPG method for viscoelastic fluids.

*J. Non-Newton. Fluid Mech.*, 247:107–122, 2017.

# Goal-oriented adaptivity in nonlinear problems



The error in the drag coefficient of the confined cylinder ( $\lambda = 0.4$ ).

# A Posteriori Error Estimation

# DPG\* Methods

- $\mathcal{T}$ : regular subdivision of  $\Omega \subset \mathbb{R}^2$  into triangles

## DPG\* Methods

- $\mathcal{T}$ : regular subdivision of  $\Omega \subset \mathbb{R}^2$  into triangles
- $\mathcal{E}_{\text{int}}$ : the set of all interior edges in  $\mathcal{T}$

## DPG\* Methods

- $\mathcal{T}$ : regular subdivision of  $\Omega \subset \mathbb{R}^2$  into triangles
- $\mathcal{E}_{\text{int}}$ : the set of all interior edges in  $\mathcal{T}$
- $\vec{u} = (u, \vec{q}, \hat{u}, \hat{q}_n) \in L^2(\Omega) \times L^2(\Omega)^d \times H_0^{1/2}(\mathcal{S}) \times H^{-1/2}(\mathcal{S})$
- $\vec{v} = (v, \vec{\tau}) \in H^1(\mathcal{T}) \times H(\text{div}, \mathcal{T})$

# DPG\* Methods

- $\mathcal{T}$ : regular subdivision of  $\Omega \subset \mathbb{R}^2$  into triangles
- $\mathcal{E}_{\text{int}}$ : the set of all interior edges in  $\mathcal{T}$
- $\vec{u} = (u, \vec{q}, \hat{u}, \hat{q}_n) \in L^2(\Omega) \times L^2(\Omega)^d \times H_0^{1/2}(\mathcal{S}) \times H^{-1/2}(\mathcal{S})$
- $\vec{v} = (v, \vec{r}) \in H^1(\mathcal{T}) \times H(\text{div}, \mathcal{T})$
- $b(\vec{u}, \vec{v}) = ((u, \vec{q}), \mathcal{L}_h \vec{v})_{\Omega} + \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_{\partial\mathcal{T}}$

# DPG\* Methods

- $\mathcal{T}$ : regular subdivision of  $\Omega \subset \mathbb{R}^2$  into triangles
- $\mathcal{E}_{\text{int}}$ : the set of all interior edges in  $\mathcal{T}$
- $\vec{u} = (u, \vec{q}, \hat{u}, \hat{q}_n) \in L^2(\Omega) \times L^2(\Omega)^d \times H_0^{1/2}(\mathcal{S}) \times H^{-1/2}(\mathcal{S})$
- $\vec{v} = (v, \vec{r}) \in H^1(\mathcal{T}) \times H(\text{div}, \mathcal{T})$
- $b(\vec{u}, \vec{v}) = ((u, \vec{q}), \mathcal{L}_h \vec{v})_\Omega + \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_{\partial\mathcal{T}}$
- $G(\vec{u}) = (f, u)_\Omega$

# DPG\* Methods

- $\vec{u} = (u, \vec{q}, \hat{u}, \hat{q}_n) \in L^2(\Omega) \times L^2(\Omega)^d \times H_0^{1/2}(\mathcal{S}) \times H^{-1/2}(\mathcal{S})$
- $\vec{v} = (v, \vec{r}) \in H^1(\mathcal{T}) \times H(\text{div}, \mathcal{T})$
- $b(\vec{u}, \vec{v}) = ((u, \vec{q}), \mathcal{L}_h \vec{v})_\Omega + \langle \hat{u}, \vec{p} \cdot \vec{n} \rangle_h + \langle \hat{q}_n, v \rangle_{\partial\mathcal{T}}$
- $G(\vec{u}) = (f, u)_\Omega$

Theorem (Demkowicz, Gopalakrishnan, & K.)

Assume  $\Omega \subset \mathbb{R}^2$  with sufficiently regular boundary. If  $\vec{\mathbf{v}}_h = (\mathbf{v}_h, \vec{\mathbf{p}}_h) \in \mathcal{V}_h$  satisfies

$$b(\vec{w}_h, \vec{\mathbf{v}}_h) = G(\vec{w}_h) \quad \forall \vec{w}_h \in \mathcal{U}_h,$$

then  $\exists C_2 > C_1 > 0$ , independent of the element sizes  $h_K$ , such that

$$C_1 \eta^*(\vec{\mathbf{v}}_h) \leq \|\vec{\mathbf{v}} - \vec{\mathbf{v}}_h\|_{\mathcal{V}} \leq C_2 \eta^*(\vec{\mathbf{v}}_h),$$

where

$$\eta^*(\vec{\mathbf{v}}_h)^2 = \left\| \mathcal{L}_h \vec{\mathbf{v}}_h - f \right\|_\Omega^2 + \sum_{E \in \mathcal{E}_{\text{int}}} h_E \|[\![\vec{\mathbf{p}}_h]\!] \|_{L^2(E)}^2 + \sum_{E \in \mathcal{E}} h_E \|[\![\mathbf{v}_h]\!] \|_{H^1(E)}^2$$

## Sketch of the Proof

- $\mathcal{B}$  is a bijection so

$$\|\textcolor{brown}{v} - \textcolor{brown}{v}_h\|_{\mathcal{V}}^2 \asymp \|\mathcal{B}' \textcolor{brown}{v}_h - G\|_{\mathcal{U}'}^2$$

# Sketch of the Proof

- $\mathcal{B}$  is a bijection so

$$\|\boldsymbol{v} - \boldsymbol{v}_h\|_{\mathcal{V}}^2 \asymp \|\mathcal{B}' \boldsymbol{v}_h - G\|_{\mathcal{U}'}^2$$

- Decompose supremum into three terms:

$$\begin{aligned} \left( \sup_{\vec{u} \in \mathcal{U}} \frac{b(\vec{u}, \vec{v}_h) - G(\vec{u})}{\|\vec{u}\|_{\mathcal{U}}} \right)^2 &= \left( \sup_{(u, \vec{q}) \in L^2(\Omega) \times L^2(\Omega)^d} \frac{((u, \vec{q}), \mathcal{L}_h \vec{v}_h - f)_{\Omega}}{\|(u, \vec{q})\|_{\Omega}} \right)^2 \\ &\quad + \left( \sup_{\hat{u} \in H_0^{1/2}(\mathcal{S})} \frac{\langle \hat{u}, \vec{p}_h \cdot \vec{n} \rangle_h}{\|\hat{u}\|_{H^{1/2}(\partial\mathcal{T})}} \right)^2 \\ &\quad + \left( \sup_{\hat{q}_n \in H^{-1/2}(\mathcal{S})} \frac{\langle \hat{q}_n, v_h \rangle_h}{\|\hat{q}_n\|_{H^{-1/2}(\partial\mathcal{T})}} \right)^2 \end{aligned}$$

# Sketch of the Proof

Term 1:

- $$\sup_{(u, \vec{q}) \in L^2(\Omega) \times L^2(\Omega)^d} \frac{((u, \vec{q}), \mathcal{L}_h \vec{v}_h - f)_\Omega}{\|(u, \vec{q})\|_\Omega} = \|\mathcal{L}_h v_h - f\|_\Omega$$

# Sketch of the Proof

Term 1:

- $$\sup_{(u, \vec{q}) \in L^2(\Omega) \times L^2(\Omega)^d} \frac{((u, \vec{q}), \mathcal{L}_h \vec{v}_h - f)_\Omega}{\|(u, \vec{q})\|_\Omega} = \|\mathcal{L}_h v_h - f\|_\Omega$$

Term 2\*:

- $$\sup_{\hat{u} \in H_0^{1/2}(\mathcal{S})} \frac{\langle \hat{u}, \vec{p}_h \cdot \vec{n} \rangle_h}{\|\hat{u}\|_{H^{1/2}(\partial\mathcal{T})}} = \sup_{u \in H_0^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

# Sketch of the Proof

Term 1:

- $$\sup_{(u, \vec{q}) \in L^2(\Omega) \times L^2(\Omega)^d} \frac{((u, \vec{q}), \mathcal{L}_h \vec{v}_h - f)_\Omega}{\|(u, \vec{q})\|_\Omega} = \|\mathcal{L}_h v_h - f\|_\Omega$$

Term 2\*:

- $$\sup_{\hat{u} \in H_0^{1/2}(\mathcal{S})} \frac{\langle \hat{u}, \vec{p}_h \cdot \vec{n} \rangle_h}{\|\hat{u}\|_{H^{1/2}(\partial\mathcal{T})}} = \sup_{u \in H_0^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

Term 3\*:

- $$\sup_{\hat{q}_n \in H^{-1/2}(\mathcal{S})} \frac{\langle \hat{q}_n, v_h \rangle_h}{\|\hat{q}_n\|_{H^{-1/2}(\partial\mathcal{T})}} = \sup_{\vec{q} \in H(\text{div}, \Omega)} \frac{\langle \text{tr}_n(\vec{q}), v_h \rangle_h}{\|\vec{q}\|_{H(\text{div}, \Omega)}}$$

\*requires a lemma

## Sketch of the Proof

Theorem (Demkowicz, Gopalakrishnan, K.)

If  $\int_{\partial K} \hat{p}_K [\vec{p}_h] = 0$  for all  $\hat{p}_K \in P_C^1(\partial K)$  then

$$\sup_{\mu \in H_0^1(\Omega)} \frac{\langle \operatorname{tr} \mu, \vec{p}_h \cdot \vec{n} \rangle_h^2}{\|\mu\|_{H^1(\Omega)}^2} \asymp \sum_{E \in \mathcal{E}} h_E \|[\vec{p}_h]\|_{L^2(E)}^2.$$

Theorem (Demkowicz, Gopalakrishnan, K.)

If  $\int_E [v_h] = 0$  for all  $E \in \mathcal{E}$  then

$$\sup_{\vec{q} \in H(\operatorname{div}, \Omega)} \frac{\langle \operatorname{tr}_n(\vec{q}), v_h \rangle_h^2}{\|\vec{q}\|_{H(\operatorname{div}, \Omega)}^2} \asymp \sum_{E \in \mathcal{E}} h_E \| [v_h] \|_{H^1(E)}^2 \asymp \sum_{E \in \mathcal{E}} h_E^{-1} \| [v_h] \|_{L^2(E)}^2.$$

# Sketch of the Proof

## Theorem (Regular decompositions)

Given any  $\vec{q} \in H(\text{div}, \Omega)$ , there exist  $\varphi \in H^1(\Omega)$  and  $\vec{\psi} \in H^1(\Omega)$  such that  $\vec{q} = \text{curl}(\varphi) + \vec{\psi}$  and

$$\|\varphi\|_{H^1(\Omega)} + \|\vec{\psi}\|_{H^1(\Omega)} \lesssim \|\vec{q}\|_{H(\text{div}, \Omega)}.$$

In this situation,  $\vec{q} = \text{curl}(\varphi) + \vec{\psi}$  is called a *regular decomposition* of  $\vec{q}$ .

# Sketch of the Proof

## Theorem (Demlow, Hirani, Schöberl)

Let  $E \in \mathcal{E}$ ,  $u \in H^1(\Omega)$ , and  $\vec{q} \in H(\text{div}, \Omega)$  be arbitrary. Let  $h_E$  denote the length of the edge  $E$  and let  $\varphi \in H^1(\Omega)$  and  $\vec{\psi} \in H^1(\Omega)$  belong to a *regular decomposition*  $\vec{q} = \text{curl}(\varphi) + \vec{\psi}$ . There exist *commuting quasi-interpolation operators*  $\mathcal{I} : H^1(\Omega) \rightarrow P_C^1(\mathcal{T})$  and  $\vec{\mathcal{I}} : H(\text{div}, \Omega) \rightarrow \mathcal{RT}^0(\mathcal{T})$ , such that  $\text{curl} \circ \mathcal{I} = \vec{\mathcal{I}} \circ \text{curl}$  and the following inequalities hold:

$$(5) \quad \|u - \mathcal{I}u\|_{L^2(E)}^2 \lesssim h_E \|u\|_{H^1(\Omega_E)}^2,$$

$$(6) \quad \|\varphi - \mathcal{I}\varphi\|_{L^2(E)}^2 + \|(\vec{\psi} - \vec{\mathcal{I}}\vec{\psi}) \cdot \vec{n}\|_{L^2(E)}^2 \lesssim h_E \|\vec{q}\|_{H(\text{div}, \Omega_E)}^2.$$

Each of the statements above hold with  $H^1(\Omega)$  replaced by  $H_0^1(\Omega)$ ,  $H^1(\Omega)$  replaced by  $H_0^1(\Omega)$ , and  $H(\text{div}, \Omega)$  replaced by  $H_0(\text{div}, \Omega)$ . In this case,  $\mathcal{I} : H_0^1(\Omega) \rightarrow P_{C,0}^1(\mathcal{T})$  and  $\vec{\mathcal{I}} : H_0(\text{div}, \Omega) \rightarrow \mathcal{RT}_0^0(\mathcal{T})$ .

---

A. Demlow and A. N. Hirani.

A posteriori error estimates for finite element exterior calculus: The de Rham complex.  
*Found. Computat. Math.*, 14(6):1337–1371, 2014.

J. Schöberl.

A posteriori error estimates for Maxwell equations.  
*Math. Comput.*, 77(262):633–649, 2008.

# Sketch of the Proof

$$\sup_{u \in H^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

## Sketch of the Proof

$$\sup_{u \in H^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

- Let  $u_h = \mathcal{I}u$ . Then

$$\langle u - u_h, \vec{p}_h \cdot \vec{n} \rangle_h = \sum_{K \in \mathcal{T}} (u - u_h, \vec{p}_h \cdot \vec{n}_K)_{\partial K} = \sum_{E \in \mathcal{E}} (u - u_h, [\![\vec{p}_h]\!])_E$$

## Sketch of the Proof

$$\sup_{u \in H^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

- Let  $u_h = \mathcal{I}u$ . Then

$$\langle u - u_h, \vec{p}_h \cdot \vec{n} \rangle_h = \sum_{K \in \mathcal{T}} (u - u_h, \vec{p}_h \cdot \vec{n}_K)_{\partial K} = \sum_{E \in \mathcal{E}} (u - u_h, [\![\vec{p}_h]\!])_E$$

- For each edge  $E$ ,

$$(u - u_h, [\![\vec{p}_h]\!])_E \leq \|u - \mathcal{I}u\|_E \|[\![\vec{p}_h]\!]\|_E \lesssim h_K^{1/2} \|u\|_{H^1(\Omega_E)} \|[\![\vec{p}_h]\!]\|_E$$

where  $\Omega_E$  denotes the patch of elements associated with the edge  $E$ .

## Sketch of the Proof

$$\sup_{u \in H^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

- Let  $u_h = \mathcal{I}u$ . Then

$$\langle u - u_h, \vec{p}_h \cdot \vec{n} \rangle_h = \sum_{K \in \mathcal{T}} (u - u_h, \vec{p}_h \cdot \vec{n}_K)_{\partial K} = \sum_{E \in \mathcal{E}} (u - u_h, [\![\vec{p}_h]\!])_E$$

- For each edge  $E$ ,

$$(u - u_h, [\![\vec{p}_h]\!])_E \leq \|u - \mathcal{I}u\|_E \|[\![\vec{p}_h]\!]\|_E \lesssim h_K^{1/2} \|u\|_{H^1(\Omega_E)} \|[\![\vec{p}_h]\!]\|_E$$

where  $\Omega_E$  denotes the patch of elements associated with the edge  $E$ .

- Upper bound follows from manipulating the supremum

## Sketch of the Proof

$$\sup_{u \in H^1(\Omega)} \frac{\langle \text{tr}(u), \vec{p}_h \cdot \vec{n} \rangle_h}{\|u\|_{H^1(\Omega)}}$$

- Let  $u_h = \mathcal{I}u$ . Then

$$\langle u - u_h, \vec{p}_h \cdot \vec{n} \rangle_h = \sum_{K \in \mathcal{T}} (u - u_h, \vec{p}_h \cdot \vec{n}_K)_{\partial K} = \sum_{E \in \mathcal{E}} (u - u_h, [\![\vec{p}_h]\!])_E$$

- For each edge  $E$ ,

$$(u - u_h, [\![\vec{p}_h]\!])_E \leq \|u - \mathcal{I}u\|_E \|[\![\vec{p}_h]\!]\|_E \lesssim h_K^{1/2} \|u\|_{H^1(\Omega_E)} \|[\![\vec{p}_h]\!]\|_E$$

where  $\Omega_E$  denotes the patch of elements associated with the edge  $E$ .

- Upper bound follows from manipulating the supremum
- Lower bound follows from Verfürth's bubble function technique

# Sketch of the Proof

$$\sup_{\vec{q} \in H(\text{div}, \Omega)} \frac{\langle \text{tr}_n(\vec{q}), v_h \rangle_h}{\|\vec{q}\|_{H(\text{div}, \Omega)}}$$

- Gauss & Stokes:

$$\langle (\text{curl } \phi) \cdot \vec{n}_K, v \rangle_{\partial K} = (\text{curl } \phi, \text{grad } v)_K + (\text{div curl } \phi, v)_K = \langle \phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K}$$

## Sketch of the Proof

$$\sup_{\vec{q} \in H(\text{div}, \Omega)} \frac{\langle \text{tr}_n(\vec{q}), v_h \rangle_h}{\|\vec{q}\|_{H(\text{div}, \Omega)}}$$

- Gauss & Stokes:

$$\langle (\text{curl } \phi) \cdot \vec{n}_K, v \rangle_{\partial K} = (\text{curl } \phi, \text{grad } v)_K + (\text{div curl } \phi, v)_K = \langle \phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K}$$

- Let  $\sigma_h = \vec{\mathcal{I}} \text{curl } \phi = \text{curl } \mathcal{I}\phi$ . Then

$$\begin{aligned} \langle (\phi - \sigma_h) \cdot \vec{n}_K, v \rangle_{\partial K} &= \langle \phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K} \\ &= \sum_{K \in \mathcal{T}} (\phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K)_{\partial K} \\ &= \sum_{E \in \mathcal{E}} (\phi - \mathcal{I}\phi, [(\text{grad } v) \cdot \vec{t}_K])_E \end{aligned}$$

## Sketch of the Proof

$$\sup_{\vec{q} \in H(\text{div}, \Omega)} \frac{\langle \text{tr}_n(\vec{q}), v_h \rangle_h}{\|\vec{q}\|_{H(\text{div}, \Omega)}}$$

- Gauss & Stokes:

$$\langle (\text{curl } \phi) \cdot \vec{n}_K, v \rangle_{\partial K} = (\text{curl } \phi, \text{grad } v)_K + (\text{div curl } \phi, v)_K = \langle \phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K}$$

- Let  $\sigma_h = \vec{\mathcal{I}} \text{curl } \phi = \text{curl } \mathcal{I}\phi$ . Then

$$\begin{aligned} \langle (\phi - \sigma_h) \cdot \vec{n}_K, v \rangle_{\partial K} &= \langle \phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K} \\ &= \sum_{K \in \mathcal{T}} (\phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K)_{\partial K} \\ &= \sum_{E \in \mathcal{E}} (\phi - \mathcal{I}\phi, [(\text{grad } v) \cdot \vec{t}_K])_E \end{aligned}$$

- Manipulate as before to get upper bound

# Sketch of the Proof

$$\sup_{\vec{q} \in H(\text{div}, \Omega)} \frac{\langle \text{tr}_n(\vec{q}), v_h \rangle_h}{\|\vec{q}\|_{H(\text{div}, \Omega)}}$$

- Gauss & Stokes:

$$\langle (\text{curl } \phi) \cdot \vec{n}_K, v \rangle_{\partial K} = (\text{curl } \phi, \text{grad } v)_K + (\text{div curl } \phi, v)_K = \langle \phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K}$$

- Let  $\sigma_h = \vec{\mathcal{I}} \text{curl } \phi = \text{curl } \mathcal{I}\phi$ . Then

$$\begin{aligned} \langle (\phi - \sigma_h) \cdot \vec{n}_K, v \rangle_{\partial K} &= \langle \phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K \rangle_{\partial K} \\ &= \sum_{K \in \mathcal{T}} (\phi - \mathcal{I}\phi, (\text{grad } v) \cdot \vec{t}_K)_{\partial K} \\ &= \sum_{E \in \mathcal{E}} (\phi - \mathcal{I}\phi, [(\text{grad } v) \cdot \vec{t}_K])_E \end{aligned}$$

- Manipulate as before to get upper bound
- Lower bound from Verfürth

# Summary

- Early work and applications
- DPG/DPG\* paradigm
- A *priori* error analysis
  - DPG error controlled by regularity of solution
  - DPG\* error controlled by regularity of solution **and** Lagrange multiplier
- Duality
- A *posteriori* error control
- Goal-oriented adaptivity

# CSEM objectives

## Area A:

- Analysis of the saddle point paradigm
- *A priori* and *a posteriori* error analysis of DPG\* methods
- Goal-oriented adaptive mesh refinement

## Area B:

- Implement finite element methods and compute *a posteriori* error indicators

## Area C:

- Analysis of viscoelastic fluid flow

# CSEM objectives

## Area A:

- Analysis of the saddle point paradigm
- *A priori* and *a posteriori* error analysis of DPG\* methods
- Goal-oriented adaptive mesh refinement
- Well-posedness of multiple/coupled variational formulations (linear elasticity)

## Area B:

- Implement finite element methods and compute *a posteriori* error indicators
- ESEAS shape functions library
- New assembly algorithm for DPG methods

## Area C:

- Analysis of viscoelastic fluid flow
- Structural mechanics and wave propagation problems

## References

- L. Demkowicz, J. Gopalakrishnan, and B. Keith.  
The DPG-star method.  
*In preparation*, 2018.
- F. Fuentes, B. Keith, L. Demkowicz, and S. Nagaraj.  
Orientation embedded high order shape functions for  
the exact sequence elements of all shapes.  
*Comput. Math. Appl.*, 70(4):353–458, 2015.
- F. Fuentes, B. Keith, L. Demkowicz, and P. L. Tallec.  
Coupled variational formulations of linear elasticity and  
the DPG methodology.  
*J. Comput. Phys.*, 348:715–731, 2017.
- B. Keith, L. Demkowicz, and J. Gopalakrishnan.  
DPG\* method.  
ICES Report 17-25, The University of Texas at Austin, 2017.
- B. Keith, F. Fuentes, and L. Demkowicz.  
The DPG methodology applied to different variational  
formulations of linear elasticity.  
*Comput. Methods Appl. Mech. Engrg.*, 309:579–609,  
2016.
- B. Keith, P. Knechtges, N. V. Roberts, S. Elgeti, M. Behr,  
and L. Demkowicz.  
An ultraweak DPG method for viscoelastic fluids.  
*J. Non-Newton. Fluid Mech.*, 247:107–122, 2017.
- B. Keith, S. Petrides, F. Fuentes, and L. Demkowicz.  
Discrete least-squares finite element methods.  
*Comput. Methods Appl. Mech. Engrg.*, 327:226–255,  
2017.
- B. Keith, N. V. Roberts, and L. Demkowicz.  
Goal-oriented adaptive mesh refinement with a DPG  
method for viscoelastic fluids.  
*In preparation*, 2018.
- B. Keith, A. Vaziri Astaneh, and L. Demkowicz.  
Goal-oriented adaptive mesh refinement for  
non-symmetric functional settings.  
*arXiv:1711.01996 (math.NA)*, 2017.
- A. Vaziri Astaneh, B. Keith, and L. Demkowicz.  
On perfectly matched layers for discontinuous  
Petrov–Galerkin methods.  
*arXiv:1804.04496 (math.NA)*, 2018.

Thank you!

# Extra slides

# The ultraweak form of Poisson's problem

Strong form:

$$\begin{aligned}-\operatorname{div} \boldsymbol{\sigma} &= f, \\ \boldsymbol{\sigma} - \operatorname{grad} u &= 0\end{aligned}$$

# The ultraweak form of Poisson's problem

Strong form:

$$\begin{aligned}-\operatorname{div} \boldsymbol{\sigma} &= f, \\ \boldsymbol{\sigma} - \operatorname{grad} u &= 0\end{aligned}$$

(Ultra)weak form:

$$\begin{aligned}(\boldsymbol{\sigma}, \operatorname{grad} v)_\Omega &= f, \quad \forall v \\ (\boldsymbol{\sigma}, \boldsymbol{\tau})_\Omega + (u, \operatorname{div} \boldsymbol{\tau})_\Omega &= 0, \quad \forall \boldsymbol{\tau}\end{aligned}$$

# The ultraweak form of Poisson's problem

Strong form:

$$\begin{aligned}-\operatorname{div} \boldsymbol{\sigma} &= f, \\ \boldsymbol{\sigma} - \operatorname{grad} u &= 0\end{aligned}$$

(Ultra)weak form:

$$\begin{aligned}(\boldsymbol{\sigma}, \operatorname{grad} v)_\Omega &= f, \quad \forall v \\ (\boldsymbol{\sigma}, \boldsymbol{\tau})_\Omega + (u, \operatorname{div} \boldsymbol{\tau})_\Omega &= 0, \quad \forall \boldsymbol{\tau}\end{aligned}$$

Broken ultraweak form:

$$\begin{aligned}(\boldsymbol{\sigma}, \operatorname{grad} v)_\tau - \langle \hat{\sigma}_n, v \rangle_{\partial\tau} &= f, \quad \forall v \\ (\boldsymbol{\sigma}, \boldsymbol{\tau})_\tau + (u, \operatorname{div} \boldsymbol{\tau})_\tau - \langle \hat{u}, \boldsymbol{\tau} \cdot \hat{n} \rangle_{\partial\tau} &= 0, \quad \forall \boldsymbol{\tau}\end{aligned}$$

# The ultraweak form of Poisson's problem

Strong form:

$$\begin{aligned}-\operatorname{div} \boldsymbol{\sigma} &= f, \\ \boldsymbol{\sigma} - \operatorname{grad} u &= 0\end{aligned}$$

(Ultra)weak form:

$$\begin{aligned}(\boldsymbol{\sigma}, \operatorname{grad} v)_\Omega &= f, \quad \forall v \\ (\boldsymbol{\sigma}, \boldsymbol{\tau})_\Omega + (u, \operatorname{div} \boldsymbol{\tau})_\Omega &= 0, \quad \forall \boldsymbol{\tau}\end{aligned}$$

Broken ultraweak form:

$$\begin{aligned}(\boldsymbol{\sigma}, \operatorname{grad} v)_\tau - \langle \hat{\sigma}_n, v \rangle_{\partial\tau} &= f, \quad \forall v \\ (\boldsymbol{\sigma}, \boldsymbol{\tau})_\tau + (u, \operatorname{div} \boldsymbol{\tau})_\tau - \langle \hat{u}, \boldsymbol{\tau} \cdot \hat{n} \rangle_{\partial\tau} &= 0, \quad \forall \boldsymbol{\tau}\end{aligned}$$

N.B.

- The functions  $u$  and  $\boldsymbol{\sigma}$  are discontinuous

$$u \in L^2(\Omega) \quad \boldsymbol{\sigma} \in (L^2(\Omega))^3 \quad u = (u, \boldsymbol{\sigma}, \hat{u}, \hat{\sigma}_n)$$

- All QoI can be written

$$g(\mathbf{u}) = \int_{\Omega} g_1 \cdot u + \int_{\Omega} \mathbf{g}_2 \cdot \boldsymbol{\sigma} + \int_{\partial\Omega} \hat{g}_1 \cdot \hat{u} + \int_{\partial\Omega} \hat{g}_2 \cdot \hat{\sigma}_n$$

# Goal-oriented error estimation and adaptivity

Quantity of interest

$$g(\mathbf{u}) = \int_{\Omega} u(\mathbf{x}) g_1(\mathbf{x}) d\mathbf{x}$$

where

$$g_1(\mathbf{x}) = \begin{cases} 1 & x \in \Omega_0 \\ 0 & \text{otherwise} \end{cases}$$

Manufactured solution

$$u^{\text{man}}(x, y, z) = f(x/4)f(y)f(z)$$

where


$$f(x) = x(1-x)((x/4) + (1-4x)^2)$$

and

$$\Omega = [0, 4] \times [0, 1] \times [0, 1]$$


1D perspective

Shallow gradient



Steep gradient

Domain



# Goal-oriented error estimation and adaptivity

Quantity of interest

$$g(\mathbf{u}) = \int_{\Omega} u(\mathbf{x}) g_1(\mathbf{x}) \, d\mathbf{x}$$

where

$$g_1(\mathbf{x}) = \begin{cases} 1 & x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Manufactured solution


$$u^{\text{man}}(x, y, z) = f(x/4)f(y)f(z)$$

where

$$f(x) = x(1-x)((x/4) + (1-4x)^2)$$

and

$$\Omega = [0, 4] \times [0, 1] \times [0, 1]$$





# Goal-oriented error estimation and adaptivity

Quantity of interest

$$g(\mathbf{u}) = \int_{\Omega} u(\mathbf{x}) g_1(\mathbf{x}) d\mathbf{x}$$

where

$$g_1(\mathbf{x}) = \begin{cases} 1 & x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$






# Goal-oriented error estimation and adaptivity

Quantity of interest

$$g(\mathbf{u}) = \int_{\Omega} u(\mathbf{x}) g_1(\mathbf{x}) d\mathbf{x}$$


where

$$g_1(\mathbf{x}) = \begin{cases} 1 & x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$




# Goal-oriented error estimation and adaptivity

Energy strategy



Goal-oriented strategy






# Goal-oriented error estimation and adaptivity

Quantity of interest (flux)

$$g(\mathbf{u}) = \int_{\Omega} \boldsymbol{\sigma}(\mathbf{x}) \cdot \mathbf{g}_2(\mathbf{x}) \, d\mathbf{x}$$

where

$$\mathbf{g}_2(\mathbf{x}) = \begin{cases} [1, 0, 0]^T & x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$






# Goal-oriented error estimation and adaptivity

Quantity of interest (boundary)

$$g(\mathbf{u}) = \int_{\partial\Omega} \hat{u}(\mathbf{x}) \hat{g}_1(\mathbf{x}) d\mathbf{x}$$

where

$$\hat{g}_1(\mathbf{x}) = \begin{cases} 1 & x = 0 \\ 0 & \text{otherwise} \end{cases}$$






# Goal-oriented error estimation and adaptivity

Quantity of interest (boundary flux)

$$g(\mathbf{u}) = \int_{\partial\Omega} \hat{\sigma}_n(\mathbf{x}) \hat{g}_2(\mathbf{x}) d\mathbf{x}$$




where

$$\hat{g}_2(\mathbf{x}) \approx \begin{cases} 1 & x = 0 \\ 0 & \text{otherwise} \end{cases}$$



# Goal-oriented error estimation and adaptivity

Pointwise goal



Final mesh

12 refinements